1   Chemical signatures of social information in Barbary macaques

3   Brigitte M. Weiß[1,2], Claudia Birkemeyer[3], Marlen Kücklich[1,2]* & Anja Widdig[1,2]*

4   [1] Behavioural Ecology Group, Leipzig University, Leipzig, Germany

5   [2] Department of Human Behaviour, Ecology and Culture, Max-Planck-Institute for Evolutionary

6   Anthropology, Leipzig, Germany

7   [3] now Claudia Wiesner, Mass Spectrometry Group, Institute of Analytical Chemistry, Leipzig

8   University, Leipzig, Germany

9   *contributed equally

Corresponding author: Brigitte M. Weiß, Institute of Biology, Talstraße 33, 04103 Leipzig, Germany,

phone: ++49-341-9736-754, email: brigitte.schloegl@uni-leipzig.de

Abstract

Primates are well-known for their complex social lives and intricate social relationships, which

requires them to obtain and update social knowledge about conspecifics. The sense of smell may

provide access to social information that is unavailable in other sensory domains or enhance the

precision and reliability of other sensory cues. However, the cognition of social information in

catarrhine primates has been studied primarily in the visual and auditory domain. We assessed the

social information content of body odor in a catarrhine primate, the Barbary macaque (*Macaca*

sylvanus), in three semi-free ranging groups at Affenberg Salem, Germany. In particular, we related

socially relevant attributes (identity, genetic relatedness, rank, sex, age, group membership) to

chemical profiles of body odor. We applied non-invasive active sampling via thermal desorption

tubes, and analyzed samples by gas chromatography–mass spectrometry. We found robust evidence

for olfactory individual signatures, and limited support for kin signatures. Chemical profiles were also

related to group membership, but little to rank, sex or age. The detected associations between

chemical profiles and socially relevant attributes form the theoretical foundations for olfactory

information transfer, highlighting the potential of body odor to provide valuable social information in

this highly visually oriented primate.

Keywords: individual signature, catarrhine primate, relatedness, social information, body odor

  Introduction

The development and maintenance of social relationships requires animals to acquire and update

social information about conspecifics. For instance, animals that form stable relationships need to be

able to recognize the unique features of other individuals, and indeed, individual recognition is

widespread in the animal kingdom (e.g. Rendall et al., 1996; Scheiber et al., 2011; reviewed in

Tibbetts and Dale, 2007). Additionally, social animals may interact differently with each other

depending on attributes such as sex, rank, group membership or genetic relatedness (Ward and

Webster, 2016). Primates are particularly well-known for forming intricate social relationships and

frequently show strong and long-lasting relationships among kin (reviewed in Silk, 2001), but also

with unrelated individuals (e.g. Goffe et al., 2016; Ostner and Schülke, 2014). Representations of

these social relationships are based on the recognition of traits such as identity, rank and kinship

(Seyfarth et al., 2005). For instance, rhesus macaques (*Macaca mulatta)* discriminate individuals, and

familiar as well as unfamiliar kin (Pfefferle et al., 2014a, 2014b; Rendall et al., 1996). Various

macaque species also appear to categorize individuals according to their relationships with others

(see overview in Cheney and Seyfarth, 1999). Similarly, baboons discriminate individuals according to

their identity and rank, and they also appear to recognize third-party relationships (Cheney et al.,

1996; Cheney and Seyfarth, 1999; Seyfarth et al., 2005).

The social information underlying differentiated interactions can be emitted and received via

different sensory modalities (Moreira et al., 2013; Yorzinski, 2017). Olfaction is among the

evolutionarily oldest of the animal senses (Wyatt, 2014), and social communication is considered as

one of its major functions (Stevenson, 2010). Indeed, social interactions are mediated via olfaction in

numerous mammals (e.g. African elephants, *Loxodonta africana*, Bates et al., 2008; black rhinoceros,

Diceros bicornis, Linklater et al., 2013), including primates. For instance, many strepsirrhine (lemurs,

lorises and galagos) and platyrrhine (South American) primates show pronounced scent-marking

behavior (e.g. delBarco-Trillo et al., 2012; Janda et al., 2019; Scordato and Drea, 2007).

Observational, experimental and chemical studies suggest that primate scent marks as well as odors

emanating directly from the body may convey a wealth of information about a sender such as

identity, sex, age, rank or reproductive state (e.g. Drea and Scordato, 2008; Fontani et al., 2022;

Janda et al., 2019; Kücklich et al., 2019), and that scent marks primarily function in intra-sexual

competition and mediating reproductive behavior (Drea and Scordato, 2008; Heymann, 2006a;

Moreira et al., 2013).

Notably, vision has gained more and more importance during primate evolution (see overview in

Veilleux et al., 2022), and catarrhine (African and Asian) primates, in particular, are highly visually

oriented animals (Kawamura, 2016). Scent-marking, on the other hand, is uncommon in catarrhines

(Freeman et al., 2012), and in contrast to evolutionarily older primates, catarrhines lack a functional

accessory olfactory system (the vomeronasal organ, VNO, Smith et al., 2014). Accordingly, catarrhine

primates have long been regarded as little predisposed for olfactory communication (Drea, 2015).

However, also the main olfactory system has an important part in the processing of cues relevant to

social recognition, rendering the lack of a functional VNO of little significance for the communicative

functions of olfaction (see Stevenson, 2010; Tang-Martinez, 2001). Furthermore, catarrhine olfactory

sensitivity may be on par or even exceed those of other mammals (Laska and Seibt, 2002a, 2002b).

Research on catarrhine olfaction has primarily focused on humans (Heymann, 2006b). Similar to

other primates, human body odor has been associated with socially relevant information such as

identity, sex, age, kinship or reproductive status (reviewed in Loos et al., 2023), although the latter is

debated (see Zetzsche et al., 2024). Also in mandrills (*Mandrillus sphinx*), one of the few catarrhine

species known to have scent glands, body odor has been related to sex, age, dominance status,

group membership and, possibly, individual identity (Setchell et al., 2010; Vaglio et al., 2016). Results

of recent studies further show that also the body odor of other catarrhine species contains

information on socially relevant attributes such as sex, rank, female fertility, genetic relatedness or

group membership may be recognized and acted upon by conspecifics (*M. mulatta*: Henkel et al.,

2015; Weiß et al., 2018a; *Pan troglodytes*: Henkel and Setchell, 2018; *Papio anubis*: Vaglio et al.,

2021; *M. fuscata*: Rigaill et al., 2022). Nonetheless, with the exception of a few species such as

humans or mandrills, olfaction remains severely understudied in catarrhine primates (Drea, 2015).

This is particularly true for chemical profiles of body odors (i.e. the presence/absence and relative

abundance of chemical compounds), which restricts our understanding of which olfactory cues may

be available to conspecifics within catarrhines at all. To reduce this gap, we assessed the social

information content of (i.e. potential olfactory cues in) body odor in a catarrhine primate by relating

chemical profiles of body odor to socially relevant attributes of Barbary macaques living in natural

social structures under semi-free conditions at Affenberg Salem, Germany.

Like other macaque species, Barbary macaques live in multi-male, multi-female groups with female

philopatry and male dispersal (Kuester and Paul, 1999; von Segesser et al., 1999). Matrilineal kin

(especially mother-offspring and maternal half-sibling dyads) form strong bonds throughout their

lives and are therefore highly familiar with each other (Roubová et al., 2015; Silk, 2001). Patrilineal

kin (e.g. father-offspring or paternal half-sibling dyads), on the other hand, typically are less closely

associated and may even reside in different groups so that they can be expected to be less familiar

with each other(see Kuester et al., 1994; Widdig, 2007). Females show matrilineal rank acquisition in

which female relatives take similar ranks (Paul and Kuester, 1987; Prud'homme and Chapais, 1993).

Adult males form dominance hierarchies in which older males tend to rank higher, but males beyond

their prime (i.e. older than 14-16 years) are subordinate to prime age males due to being in inferior

physical condition (Berghänel et al., 2011b). Male Barbary macaques also form strong, enduring

affiliative relationships often referred to as "friendships" (Young et al., 2014), whereby triadic

interactions between two males and an infant represent an important form of male affiliative

interaction (Paul et al., 1996).

Various studies suggest that Barbary macaques can discriminate individuals, sexes, age classes and

group members using visual or acoustic cues (Fischer, 2004; Ritter, 2022; Schell et al., 2011). They

further distinguish familiar maternal kin from non-kin (acoustic cues: Hammerschmidt and Fischer,

1998). However, Barbary macaques, which are considered to be particularly promiscuous (Small,

1990), do not seem to differentiate between unfamiliar kin and non-kin either in a mating context

(Kuester et al., 1994) or in infant care (Paul et al., 1996, 1992). It remains unclear whether this is due

to i) a lack of kin-related cues, ii) monkeys not being able to recognize available cues or iii) monkeys

recognizing kin-related cues but not acting upon them. Kin-related cues in body odor composition

have been shown in a closely related species (rhesus macaques, Weiß et al., 2018a), and Barbary

macaques also use olfaction in social contexts (Simon et al., 2024), but systematic studies on

olfactory kin signatures or other socially relevant information in Barbary macaque body odor are, to

date, absent.

In the present study we aimed to i) identify variation in body odor composition (i.e. presence and

abundance of chemicals) that is systematically associated with socially relevant information about

individuals, and ii) to pinpoint potential semiochemicals (i.e. compounds and compound mixtures

involved in chemical interactions between organisms, Wyatt 2014) associated with the respective

traits. Our particular focus was on traits that form the basis for differentiated social relationships, i.e.

individual identity, genetic relatedness between and rank of individuals. We further assessed effects

of sex, age and group membership while statistically controlling for potential seasonal and

methodological variation in the chemical profiles of body odors. We hypothesized that the odorants

emanating from the skin and fur of Barbary macaques contain socially relevant information. In

particular, we expected the similarity between chemical profiles to be greater within than between

individuals, and higher among related than among unrelated individuals. Because maternal kin are in

frequent physical contact and may exchange odors from the environment or share more of the odor-

producing microbiome, we expected maternal kin to exhibit stronger kin signatures than paternal

kin, for whom kin signatures should primarily result from similarities in their genetic architecture. We

also expected individuals of similar rank to have more similar chemical profiles and/or to show rank-

related differences in the relative abundance of particular compounds (as also suggested in Burgener

et al., 2009), because individuals of similar rank may have similar hormonal profiles (e.g. androgens

or glucocorticoids), access to food resources and/or space use (see Majolo et al., 2012; Mehta and

Josephs, 2010; Naud et al., 2016; Schoof et al., 2011). Finally, given the comprehensive evidence for

olfactory differences with respect to sex, age and group membership in other primates, we also

expected similarities between chemical profiles and/or the relative abundance of compounds to be

associated with these traits.

<u>Methods</u>

Study animals

The study was conducted at Affenberg Salem, Germany, home to ~ 200 Barbary macaques in three

social groups roaming year-round within a 20 ha forested enclosure. The monkeys are well-

habituated to humans, as the park is open to visitors from spring to autumn, and to researchers year-

round (see de Turckheim and Merz, 1984 for details on the park). The monkeys' diet consists of

plants and insects found naturally within the park as well as fruit, vegetables and grain that is

provided daily by park staff. Ponds and water troughs are distributed throughout the park and grant

ad libitum access to water. Individuals differ in facial pigmentation patterns, coat color and stature

and have an alphanumeric code tattooed on the inner thigh, which makes them well distinguishable

for experienced observers. Park staff monitor the groups on a near-daily basis and record major life

history events such as births, deaths or group changes. About two thirds of the adult females receive

hormonal implants (Implanon NXT) to control population size.

We collected odor samples from a total of 72 focal males and females from all three groups ("C", "F"

& "H") and across all ages (2 - 27 years, mean ± SD = 10.5 ± 5.4 years) that were most suitable to

address the main study questions. Focal animals included 13 females without and 18 females with

hormonal contraception throughout the study period; two females received contraception for part of

the study period (see supplement 2). To address individual signatures, we included particularly well-

habituated reproductively mature individuals (> 4 years) that could be sampled frequently (10

samples per ID, Pollard et al. 2010). To address kinship, we included mothers/fathers and their

offspring (> 2 years, because younger individuals could not be reliably habituated to sampling yet).

For rank of males and females, we selected individuals from the highest- and lowest-ranking adults in

their respective groups.

Odor sampling

One experimenter (B.M.W.) collected 476 odor samples from the focal animals (mean ± SD = 6.6 ±

2.6 samples per ID) in five sampling periods between October 2020 and April 2022. Sampling took

place in spring, summer and autumn, but, for logistic reasons, not during the peak of the mating

season in the months of November, December and January. As Barbary macaques, like other

macaque species, do not appear to have specialized scent glands, we non-invasively collected the

airborne volatile and semi-volatile compounds emanating from the skin or fur of the animals across

different parts of the body (see below). We employed active sampling with thermal desorption (TD)

tubes (stainless steel, Supelco), which contained two adsorbent polymers (Tenax TA and XAD-2,

Sigma Aldrich) that trap a wide range of chemical compounds (Kücklich et al., 2017). The tubes were

connected to a silently operating air pump (BiVOC2, Holbach) via a silicone hose (Weiß et al., 2018b).

To be able to trap the air in the immediate vicinity of the animals (1 to 15 cm, mean ± SD = 3.5 ± 2.6

cm), monkeys were habituated to retrieve pieces of peanuts from a pipe collar (7.5 cm diameter)

strapped to a horizontal branch or rail while the experimenter stood next to the setup and pointed a

wooden stick with the TD tube mounted at its end towards the monkey. Once a monkey calmly

started to retrieve the peanuts, the experimenter opened the TD tube, positioned it close to the

monkey and started the air flow through the tube with the flow rate set to 1 L/min. Sampling was

resumed until an air volume of 0.5 – 1 L was collected. If the animal moved away before the sample

was completed, the pump was immediately stopped, the tube closed, the volume already sampled

noted and sampling resumed as soon as possible onto the same TD tube (mean ± SD = 2.3 ± 1.5

subsamples). A sample was typically completed with < 5 min between subsamples, except for 10 of

the 476 samples in which the sample was completed on different days (see supplement 3). Samples

were collected from body parts that could be reliably sampled without disturbing the animals, which

primarily constituted the shoulders, flanks or outer thighs (54 % of samples being collected

exclusively from these regions, and 86 % of samples included these regions). Whenever possible we

avoided sampling the hands, feet or facial area, which were the body parts most likely contaminated

with food, feces or other items from the environment. To be able to statistically control for potential

variation arising from the sampled body parts we grouped them into 4 categories: "upper"

constituted samples collected from the upper part of the body (i.e. arms, shoulders and/or the back

of the head), "central" the middle part of the body (i.e. flanks, central back or chest), and "lower" the

lower part of the body (i.e. hips, thighs and legs). Finally, we grouped all samples collected from

multiple regions of the body (i.e. upper, central and/or lower) into the category "mixed" (see

supplement 3).

We avoided sampling when conspecifics were within 3 m of the target animal or higher-ranking

individuals were within sight, when animals had body contact with group members less than 5

minutes before, when the monkeys were wet or when it was very windy. To avoid disturbances

during sampling and contamination with human scents, we did not collect any samples when visitors

or park staff were within 15 m. All equipment was handled wearing disposable gloves. The

experimenter wore a medical mouth-nose cover during sampling and avoided the use of perfumed

products or strongly smelling foods before and on sampling days.

To control for chemical compounds originating from the sampling material and sampling

environment *per se*, we further collected samples of air in the absence of animals (N = 17), TD tubes

mounted for sampling in the absence of animals but without drawing air through the tube (N = 6)

and TD tubes taken to the field but remaining closed at all times (N = 9) as blanks. Before each use,

we cleaned the TD tubes in a thermal conditioner (TD Clean-Cube, Scientific Instruments

Manufacturer) under a constant stream of nitrogen for 120 minutes. When not in use, the tubes

were closed tightly with Swagelok brass caps, wrapped in aluminum foil and stored in airtight bags at

ambient temperature.

Chemical analysis & profiling

Samples were analyzed with gas chromatography – mass spectrometry (GC-MS) on a Shimadzu

TQ8040 GC-MS coupled to a thermal desorption unit (TD-20) with a scan range from 30 to 500 *m/z*

following established GC-MS protocols for the analysis of TD tube samples (Kücklich et al., 2017;

Weiß et al., 2018b). In the process, the trapped organic compounds were entirely desorbed from the

tube, which then was conditioned and re-used for sampling. To remove any potential residues from

previous measurements from the column, we ran three blank samples at the start and end of each

batch as well as after every tenth sample.

GC-MS data were processed following an established, semi-automated procedure (Kücklich et al.,

2017; Weiß et al., 2018b), in which a preliminary compound library was created through automated

peak detection and alignment of peaks by retention time using AMDIS (version 2.71, Stein 1999). We

manually corrected this library by taking into account characteristic mass-to-charge ratios of peaks in

addition to retention times. For the 264 compounds in this library, we extracted the peak areas for

the highest ion not present in adjacent peaks using the Shimadzu GCMS Browser software. In a

subsequent data cleaning step we excluded 81 compounds consistently found in blanks with peak

areas higher than in the animal samples from further analysis, as they are unlikely to have derived

from the target animal (detailed in Weiß et al., 2018b), and performed statistical analyses using the

remaining 183 compounds. Furthermore, we assigned undetected peaks, peaks with a signal to noise

ratio < 1 and with a peak area below the noise threshold (visually determined to be at 1000) a peak

area of 0, because these peaks were not reliably separable from background noise. We tentatively

identified compounds statistically related to particular traits (see below) by comparing their mass

spectra with the best matches of the NIST Mass Spectral Library (NIST14; National Institute of

Standards and Technologies, Gaithersburg, MD, USA).

Parentage analysis

Pregnancies and births have been monitored consistently by Affenberg staff since the 1970s, so that

partial pedigrees could be built tracing back maternal relatedness over 5-6 generations for most

individuals based on these long-term observational records.

For genetic parentage analysis we used DNA extracted from blood, tissue, hair or feces of 144

individuals (including 68 of the 72 odor-sampled animals, see supplement 1 for details on genetic

samples). These included 28 offspring from cohorts 2017-2019 (for 22 of which we also collected

odor samples), their mothers and potential sires. Because Barbary macaques may also reproduce

with extra-group partners (Brauch et al., 2008), we considered all males in the population that were

alive and at least 4 years of age at the onset of the respective mating season as potential sires. For

animals born in cohort 2017 we were able to obtain samples for 96 %, and for cohorts 2018-2019,

97 % of all potential sires.

We genotyped the DNA samples using a panel of up to 23 polymorphic microsatellite markers

established for the study population by (Westphal, 2020), whereby we eventually focused on the 16

markers with the lowest dropout rate, that were most reliable, scorable and combinable in

polymerase chain reactions (PCR, see Table S1). To increase the amount of host DNA in the samples

we used the two-step multiplex approach for PCR amplification (Arandjelovic et al., 2009) by running

a first PCR with all markers simultaneously, followed by a second set of PCRs containing diluted

multiplex products as amplification templates and 4 – 6 fluorescently labelled primers with different

allele ranges. PCRs were conducted in a Mastercycler® vapo.protect thermal cycler (Eppendorf,

Hamburg, Germany) following the protocols of Westphal (2020). We performed fragment analysis on

an *ABI 3730* sequencer. For allele calling we used the Peak Scanner 2 software (Applied Biosystems®).

We typed all individuals at an average of 16.7 ± 2.8 loci. For 27 of the 28 offspring born 2017 – 2019

the observed mothers showed no mismatches to their offspring. For one offspring we could not

confirm the observed mother genetically, probably due to poor sample quality. In this case we

considered the observed mother to be the biological mother, but excluded the offspring from

paternity analysis. We could also genetically confirm the observed mother for 38 females and males

initially sampled as mothers or potential sires.

We used confirmed maternities in the paternity analysis, applying a combination of exclusion and

likelihood methods for mother-offspring-putative father trios with the programs Findsire and Cervus

following established criteria (e.g. Minkner et al., 2018; Ruiz-Lambides et al., 2018) (see supplement

1 for details).

We were able to assign fathers to 23 of the 28 offspring from cohorts 2017-2019 with high certainty

(see supplement 1 for details). Furthermore, we accepted fathers assigned with high confidence for

18 individuals initially sampled as mothers or potential sires in the present study and born between

2011 and 2016. For these cohorts, 58-87 % of potential sires were genotyped. We did not accept

fathers assigned to older individuals due to the increasingly lower proportion of potential sires

sampled. For the 68 odor-sampled and genotyped individuals, we were able to genetically confirm 25

fathers and 41 mothers.

Based on the genetic analysis and maternal pedigree data from long-term records, we calculated

pedigree relatedness (r) per kin line (maternal or paternal/mixed) for each possible dyad in our study

using the software TRACE (version 0.1.0, Westphal et al., 2023). Maternal r reflected genetic

relatedness exclusively via female ancestors, while paternal/mixed r (hereafter just called "paternal")

reflected any connections involving at least one paternal link between two given individuals. As input

we used genetically confirmed mothers whenever available, and observed mothers (based on

observational records from the field) for all remaining subjects, as well as all available paternity data.

Given that mother-offspring dyads are available from long-term observational records, our value of

pedigree relatedness for maternal relatedness can be considered as highly precise, reflecting close

maternal (r > 0.125) but also distant maternal kin (r ≤ 0.125). However, because no genetic samples

could be obtained for potential sires deceased before the onset of the study, paternities could only

be reliably determined for the younger cohorts, and therefore our values of pedigree relatedness for

the paternal line are likely underestimated in this analysis.

Dominance and rank

At the onset of the study, we asked experienced park staff to classify individuals from low to high-

ranking, and to identify the alpha male and female in each group based on observed agonistic

interactions and priority of access to resources. This initial judgement was used to pre-select focal

individuals for habituation to the sampling setup. We further conducted dynamic Elo-rating (Newton-

Fisher, 2017) using *ad libitum* observations of agonistic interactions (as defined in Berghänel et al.,

2011a) throughout all sampling periods. Dynamic Elo-rating allows for incorporating intensities of

interactions as well as prior knowledge and therefore requires fewer observed interactions and

interacting dyads than unmodified Elo-rating (Neumann et al., 2011) or matrix methods to obtain

robust rank estimates (Newton-Fisher, 2017). We used the initial classification by park staff,

supplemented by a classification by B.M.W., to define starting values for Elo-rating, and updated this

information with 462 female-female and 240 male-male interactions of varying intensities observed

between October 2020 and April 2022. We computed ranks separately per sex and group, using the

observations during the first two sampling periods as a "burn-in" phase, and extracted the ranks of

the focal individuals at the ends of the second to fifth sampling period for further analysis.

Ethical note

The study was approved by Affenberg Salem. All samples collected specifically for this study (i.e. odor

and fecal samples) were collected non-invasively and the animals participated in the odor sampling

procedure voluntarily. In line with the animal protection law in Germany, we therefore required no

further animal ethics approval.

Statistical analysis

General procedures

We conducted all analyses in R (version 4.3.3., R Core Team, 2024).

To investigate chemical similarity between samples, we performed permutational multivariate

analyses of variance (perMANOVAs) using the function "adonis2" from the package "vegan" (version

2.6-4, Oksanen et al., 2022). These analyses relate pairwise chemical (dis)similarities between all

samples entered into a given analysis to (dis)similarities between traits using distance matrices

automatically assembled from the sample-specific information provided. As measure of distance (or

dissimilarity) we used Bray-Curtis indices calculated from standardized, log(x+1)-transformed peak

areas (Stoffel et al., 2015; Weiß et al., 2018b). Depending on the investigated trait(s) we used

different data subsets and predictor variables as described for the different traits below. Unless

noted otherwise, we set ID as stratum to account for repeated measures per individual. For

perMANOVAs with multiple test predictors we first conducted a full-null model comparison by

setting the argument "by" to "NULL", and only if this was significant ($p < 0.05$) we investigated the

effects of the single predictor variables by setting the argument "by" to "margin". In case of non-

significant interactions, we re-ran the marginal model without the interaction to be able to assess the

effect of the respective main terms. We ran all perMANOVAs with the default of 999 iterations.

When the analysis of chemical (dis)similarities involved predictor variables describing pairs of

samples (e.g. pedigree relatedness) and/or required the selection of specific pairs of samples (which

is not possible with the perMANOVA), we further conducted multi-membership models. Multi-

membership models extend multilevel models by allowing to fit data with multiple membership

structures that are not purely hierarchical (Chung and Beretvas, 2012), such as terms whose levels

are present in more than one variable (e.g. ID in either sample 1 or sample 2 of a given pair-wise

comparison). For these models we computed Bray-Curtis dissimilarities from standardized, log(x+1)-

transformed peak areas using the package "vegan" and, for easier interpretation, transformed them

into similarities by subtracting the dissimilarity scores from 1. We fitted these models with chemical

similarity as response variable and various test and control predictors as described below. As

similarity scores follow a beta distribution, we computed the models using a Bayesian approach in

the package "brms" (version 2.21.0, Bürkner, 2017), which is the only package we are aware of that

implements multi-membership models with beta distribution. We used the default settings for

priors, i.e. uninformative priors.

We further conducted compositional mixed model analyses, suitable for data whose components are

constrained to sum to a constant value, for a detailed analysis of chemical profiles that considers the

peak areas of all compounds per sample rather than overall (dis)similarities between samples, and

may therefore aid in detecting (suites of) compounds associated with particular traits. For this

purpose, we vectorized the data matrix and included the matrix rows and columns (i.e. the samples

and compounds) as random intercepts to avoid pseudo-replication and heteroscedastic variance

(Jamil et al., 2013; Weiß et al., 2018b). As response variable we used standardized peak areas.

Because standardized areas are bound between 0 and 1 and sum up to 1 per sample, we applied a

centered log-ratio (clr) transformation (Boogaart and Tolosana-Delgado, 2013) and modelled the

transformed values with a Gaussian error distribution. As effects of interest in this analysis we

modelled the interactions between the predictors and compound, i.e. the random slopes & random

interactions of fixed effects test predictors within compound (see Weiß et al., 2018b). To follow a

consistent modelling approach, we also performed this model in the package brms.

In all mixed models we z-transformed continuous predictor variables to a mean of 0 and an SD of 1

prior to fitting the model to facilitate convergence and interpretation of model estimates (Schielzeth,

2010). We assessed collinearity between predictor variables using the function "vif" from the R

package "car" (version 3.1-2 Fox and Weisberg, 2019), whereby we considered values < 4 as

unproblematic. To assess model convergence we inspected rhat values and effective sample sizes

(ESS), with rhat values deviating from 1 by less than 0.01and ESS values > 400 indicating satisfactory

convergence (Vehtari et al., 2021).

We present model results most relevant for the interpretation of the test predictors in the main text,

and full model results, including rhat and ESS values, in supplement 1 (Tables S2-S5).

Individual identity

To investigate individual signatures, we analyzed distance matrices using a perMANOVA with ID as

test predictor while controlling for sex, age (in years), group and season of the year (spring, summer

or autumn). Because ID was the test predictor, we did not set ID as stratum. For this analysis we used

data from 24 individuals (12 males, 12 females) sampled 10 times each. As individual signatures

commonly result from the unique combination and abundance of compounds (i.e. the overall

similarity between chemical profiles) rather than certain compounds that vary consistently between

all individuals (see Smith, 2006; but see Rigaill et al., 2022), we did not conduct a more detailed

analysis of chemical profiles (compositional models) for individual identity.

Genetic relatedness

Similar to ID, kin signatures should result from similarities in the overall composition of compounds

rather than specific compounds. We therefore assessed whether the chemical (dis)similarities

between samples are explained by genetic relatedness. In the first of two sets of analyses, we

computed perMANOVAs using only samples of genetically confirmed units of close kin. Each kin unit

comprised 2 – 4 individuals in up to 3 generations and as such, parent-offspring, grandparent-

grandchild and/or half-sibling dyads, reflecting close kin with relatedness coefficients ranging from

0.25 to 0.5. We ran the perMANOVAs separately per kin line, using the maternal or paternal kin unit

(i.e. belonging to the same or a different kin unit) as test predictor, while controlling for sex, age and

season of the year (spring, summer or autumn). All individuals used in these analyses were sampled

for odor at least 5 times. For maternal kin we used 220 samples from 31 individuals and for paternal

kin 179 samples from 29 individuals in 12 kin units per kin line (see supplement 2). Maternal/paternal

relatedness between individuals from different units ranged from 0 to 0.125, reflecting distant or

non-kin.

While belonging to a certain kin unit was genetically confirmed in this first set of analyses, it treated

relatedness categorically as belonging to a given kin unit or not, but lacked the possibility to

differentiate the pair-wise degree of relatedness. In a second analytical approach we therefore

conducted a multi-membership model to relate pair-wise chemical similarities between samples to

kinship while accounting for a variety of fixed and random predictor variables. For the multi-

membership model we used the data from all genotyped and odor-sampled individuals (N = 68),

irrespective of whether their maternities or paternities were genetically solved. We fitted dyadic

Bray-Curtis similarities as beta-distributed response variable and relatedness between members of a

dyad as two separate test predictors: one reflecting maternal kinship (i.e. individuals linked

exclusively via the maternal line) and one reflecting paternal & mixed kinship (i.e. containing at least

one male linking two individuals, which breaks the familiarity between maternal kin typical for

matrilineally structured primate groups). To control for other sources of biological variation we also

fitted the age and rank difference between the two individuals in a given dyad, the number of days

between the two samples and the difference in the time of day the sample was collected. We further

included whether or not the two individuals were of the same sex, belonged to the same group, were

sampled in the same season of the year and from the same body region. To control for variation in

sampling and measurement we included the difference in sampling distance, sampling volume, the

number of subsamples needed to complete a sample and whether or not samples were analyzed in

the same GC-MS batch. To account for repeated measures we fitted the sample, the individual and

the dyad (i.e. the unique combination of individuals) and random intercepts, whereby sample and

individual were fitted as multi-membership terms because a given sample or individual could occur in

either of the two columns describing the elements of a dyad. For a more accurate estimation of fixed

effects, we further included all technically possible random slopes of fixed effects within the random

intercepts of individual and dyad. Using 9000 post-warmup draws, the model converged with all rhat

values deviating from 1 by < 0.01 and all ESS > 400 (see Table 1).

Dominance and rank

To address potential dominance and rank-related signatures, we first conducted a perMANOVA with

rank as a categorical test predictor (high/low) in interaction with sex while controlling for age, group

and season of the year. For this analysis we used 260 samples from 34 individuals sampled at least 5

times each and classified as particularly high- or low-ranking in their respective groups based on

judgements of staff and confirmed with behavioral data (high-ranking: 9 females & 8 males, low-

ranking: 8 females & 9 males). To keep confounds between dominance class and kinship arising from

matrilineal rank acquisition to a minimum, we chose no more than two females from the same

matriline, so that the 17 females used for this analysis represented 12 different matrilines.

We further conducted a multi-membership model using chemical data and rank as a continuous

measure (i.e. Elo scores) using 476 samples from all 72 sampled individuals. We only considered

same-sex dyads for this analysis, because Barbary macaque males outrank females and Elo scores

were thus computed separately per sex. We fitted dyadic Bray-Curtis similarities as beta-distributed

response variable and the difference in Elo scores and its interaction with sex as test predictors.

Similar to the model on genetic relatedness we also controlled for potential sources of biological and

technical variance by fitting dyad-specific overall relatedness and differences in age, sampling day,

time of day, group, season of the year, body region, sampling distance, sampling volume, number of

subsamples and GC-MS batch. As above we fitted sample and individual as multi-membership terms

as well as dyad as random intercepts, and all technically possible random slopes within individual and

dyad. We subsequently removed the interaction between the difference in Elo scores and sex

because it explained no variance (estimate 0.0004 +- 0.0028, lower/upper CI -0.005 – 0.006) and

fitted both as main terms instead. Using 12000 post-warmup draws, the model without interaction

converged with all rhat values deviating from 1 by < 0.01 and all ESS > 400 (see Table 2).

To consider the possibility that rank information could be encoded by a few specific compounds

(rather than or additional to overall similarities), we further included rank as predictor in a

compositional mixed model as described in the analyses of sex, age and group below.

Sex, age and group

To address potential signatures of sex, age or group membership we conducted a perMANOVA with

these terms as test predictors while controlling for season of the year. For this analysis we used all

476 samples from 72 individuals.

To assess to what extent specific (suites of) compounds may encode information about individual

traits, we further computed a compositional mixed model. As fixed effects predictors we fitted sex,

age, group and rank (Elo scores) along with season of the year, the sampled body part, time of day,

sampling distance, sampling volume and the number of subsamples. As random terms we fitted the

intercepts of sample, compound, individual, date, GC-MS batch and TD tube used for sampling, as

well as all technically possible random slopes, but no random correlations to avoid an overly complex

model. Using 8000 post-warmup draws, the model converged with all rhat values deviating from 1 by

< 0.01 and all ESS > 400 (see Table 3).

<u>Results</u>

Individual identity

Odor samples from the same individuals were significantly more similar than samples from different

individuals, with ID explaining ~ 20 % of the variance in Bray-Curtis indices (perMANOVA, r^2 = 0.203,

df = 21, F = 3.838, p = 0.001, example chromatograms in supplement 1, Fig. S1). An individual

chemical signature was further supported by the posterior mean estimates for ID in the multi-

membership models on genetic relatedness and rank (see Tables 1 and 2).

Genetic relatedness

Odor samples from close maternal kin were significantly more similar than samples from different kin

units (perMANOVA, r^2 = 0.099, df = 11, F = 2.537 , p = 0.001). Furthermore, individuals from the

same paternal kin unit tended to have more similar chemical profiles than individuals from different

kin units (perMANOVA , r^2 = 0.132, df = 11, F = 2.772 , p = 0.067).

Table 1: Estimates for fixed effects and random intercepts from the multi-membership model on
genetic relatedness. Terms of primary interest are indicated in italics. CrI = Credible Interval. ESS =
effective sample size.

fixed terms	Estimate	Est.Error	2.5 % CrI	97.5 % CrI	Rhat	bulk ESS	tail ESS
Intercept	0.1419	0.0264	0.0891	0.1935	1.0017	1592	2884
maternal r	*0.0001*	*0.0010*	*-0.0018*	*0.0020*	*1.0000*	*9718*	*7456*
paternal/mixed r	*-0.0016*	*0.0015*	*-0.0048*	*0.0012*	*1.0003*	*6365*	*5522*
d_elo score	-0.0002	0.0010	-0.0022	0.0018	1.0001	10551	7505
d_sex (same)	-0.0018	0.0018	-0.0053	0.0017	1.0001	9026	7614
d_age	-0.0010	0.0016	-0.0042	0.0022	1.0003	6827	6459
d_group (same)	0.0120	0.0055	0.0009	0.0229	1.0014	2163	3956
d_season (same)	-0.0244	0.0119	-0.0480	-0.0010	1.0032	562	1399
d_date	-0.3204	0.0095	-0.3391	-0.3013	1.0075	528	944
d_time	0.0005	0.0013	-0.0019	0.0030	0.9999	4987	6121
d_bodyregion (same)	0.0026	0.0028	-0.0028	0.0082	1.0005	4418	5675
d_s.distance	0.0043	0.0023	-0.0002	0.0088	1.0006	3783	5261
d_s.volume	0.0065	0.0051	-0.0035	0.0164	1.0032	984	1902
d_subsamples	-0.0072	0.0020	-0.0112	-0.0034	1.0007	4902	6101
d_batch (same)	0.1994	0.0154	0.1690	0.2302	1.0011	1051	1941
random terms							
sample (Intercept)	0.3994	0.0144	0.3724	0.4293	1.0031	1074	2166
ID (Intercept)	0.1490	0.0277	0.0960	0.2037	1.0033	471	973
dyad (Intercept)	0.0189	0.0013	0.0164	0.0214	1.0010	1690	3502

When relatedness was treated as a continuous variable (i.e. using pedigree r) across all genotyped

individuals in the data set, neither maternal nor paternal relatedness affected similarities in chemical

profiles (Table 1).

Dominance and rank

The interaction between dominance classes (high vs. low) and sex (male vs. female) had no

significant effect on the similarity between chemical profiles (perMANOVA, r^2 = 0.003, df= 1, F =

0.932 , p = 0.982), but individuals from the same dominance class had slightly more similar chemical

profiles than individuals from different dominance classes (perMANOVA, r^2 = 0.002, df = 1, F = 0.879,

p = 0.008). When treated as a continuous variable across all individuals in the data set, rank (Elo

score) had no effect on similarities between chemical profiles (Table 2).

Also the specific composition of chemical profiles showed no pronounced variation in relation to rank

(see random slope of Elo score within compound in Table 4).

Table 2: Estimates for fixed effects and random intercepts from the multi-membership model on

rank. Terms of primary interest are indicated in italics. CrI = Credible Interval. ESS = effective sample

size.

fixed term	Estimate	Est.Error	2.5 % CrI	97.5% CrI	Rhat	bulk ESS	tail ESS
Intercept	0.1544	0.0394	0.0756	0.2315	1.0010	2680	5074
d_Elo score	*-0.0001*	*0.0014*	*-0.0027*	*0.0026*	*1.0002*	*19102*	*9841*
total r	-0.0021	0.0014	-0.0049	0.0007	1.0002	18584	8977
sex (male)	-0.0228	0.0526	-0.1265	0.0798	1.0014	2879	5192
d_age	0.0013	0.0021	-0.0030	0.0054	1.0001	14446	9510
d_group (same)	0.0159	0.0070	0.0022	0.0297	1.0001	5497	7723
d_season (same)	-0.0267	0.0108	-0.0484	-0.0059	1.0015	2736	4694
d_date	-0.3250	0.0104	-0.3456	-0.3053	1.0023	1088	2508
d_time	0.0005	0.0012	-0.0019	0.0030	1.0002	15821	9039
d_bodyregion (same)	0.0041	0.0033	-0.0024	0.0107	1.0002	11941	8932
d_s.distance	0.0054	0.0025	0.0006	0.0103	1.0007	11074	9460
d_s.volume	0.0081	0.0054	-0.0026	0.0187	1.0014	2684	3564
d_subsamples	-0.0070	0.0021	-0.0113	-0.0029	0.9999	15094	10231
d_batch (same)	0.1838	0.0182	0.1477	0.2188	1.0011	2652	5513
random term							
sample (Intercept)	0.3952	0.0141	0.3684	0.4239	1.0016	2096	4665
ID (Intercept)	0.1509	0.0282	0.0988	0.2076	1.0058	891	1598
dyad (Intercept)	0.0189	0.0019	0.0152	0.0226	1.0016	2254	4976

Sex, age and group

When testing sex, age and group membership, the set of predictor variables significantly affected

chemical similarities (perMANOVA, r^2 = 0.213, df = 6, F = 21.145, p = 0.001). In particular, sex, age as

well as group explained significant, albeit small proportions of the variance in Bray-Curtis indices (3.5

– 5 %, Table 3). In comparison, season of the year (included as a control predictor) explained more

variance than sex, age and group together (14.5 %, Table 3).

Table 3: Results of the perMANOVA investigating effects of sex, age and group on chemical
similarities.

Term	Df	SumOfSqs	R^2	F	P
sex	1	0.466	0.004	2.332	0.001
age	1	0.425	0.004	2.124	0.001
group	2	6.106	0.051	15.273	0.001
season	2	17.225	0.145	43.083	0.001
Residual	469	93.756	0.787		
Total	475	119.117	1		

This pattern was similar when considering chemical profiles in detail. Slope estimates from the

compositional model suggest that the effects of sex, age and group on relative peak areas differed

between chemical compounds, whereby the variance was close to 0 for age, only slightly higher for

sex and more pronounced for group. Again, variance related to control predictors such as season of

the year was considerably higher than for sex, age and group (see Table 4).

Table 4: Estimates for random slope terms within the random intercept of compound for the
compositional model. Terms of primary interest are indicated in italics. CrI = Credible Interval. ESS =
effective sample size.

slope in compound	Estimate	Est.Error	2.5 % CrI	97.5% CrI	Rhat	bulk ESS	tail ESS
Elo score	*0.0166*	*0.0118*	*0.0007*	*0.0435*	*1.0020*	*3304*	*3855*
sex	*0.1023*	*0.0191*	*0.0626*	*0.1380*	*1.0006*	*2819*	*3082*
age	*0.0524*	*0.0175*	*0.0121*	*0.0831*	*1.0017*	*1709*	*1218*
group	*0.2901*	*0.0157*	*0.2606*	*0.3214*	*1.0012*	*3772*	*4889*
season	1.0605	0.0417	0.9822	1.1461	1.0019	2698	4371
time	0.0397	0.0182	0.0041	0.0724	1.0032	1657	2624
bodyregion	0.1333	0.0176	0.0987	0.1672	1.0009	2690	3865
s.distance	0.2624	0.0177	0.2290	0.2983	1.0003	2989	5028
s.volume	0.8104	0.0420	0.7341	0.8990	1.0015	1140	2388
subsamples	0.1280	0.0147	0.0993	0.1569	0.9999	3441	5449

When investigating the compound-specific estimates per test predictor, the credible intervals for all

compound-specific age and sex estimates overlapped 0. The group estimates for 8 compounds did

not overlap between at least 2 of the 3 groups, suggesting that the peak areas of these compounds

differ distinctly between groups (Fig. 1, supplementary Table S6). We tentatively identified 5 of these

compounds as a ketone, a carboxylic acid, an alcohol, an aromatic hydrocarbon and a catechol

derivate (Table 5).

Figure 1: Group differences (C, F and H) in peak areas (centered log-ratio transformed) of 8

compounds with retention times in [min] as indicated in the upper left corner of each graph.

Compounds are sorted by order of importance as derived from their compound-specific slope

estimates. Vertical lines show mean peak areas per group, points the estimates per group (with all

other predictors at their average) and whiskers the 95 % Credible Intervals.

Table 5: Compounds associated with group differences in chemical profiles of Barbary macaque

samples. RT = retention time (in min.), base peak describes the largest mass fragment of the peak. SI

= similarity index (from 0 - 100) for the match between mass spectra and their respective best hit in

the NIST library.

RT	base peak	best NIST library hit	SI	substance class	found in
2.64	43	2-Pentanone, 3-methyl-	91	ketone	humans [1,2], Japanese macaques[3]
2.68	43	Acetic acid	94	carboxylic acid	various mammals [1,4-9]
5.16	149	unidentified	< 80		
9.08	73	3-Pentanol, 2,4-dimethyl-	81	alcohol	*human excretions [1,2]
10.38	79	unidentified	< 80		
13.23	115	1H-Indene, 3-methyl-	92	aromatic hydrocarbon	unknown
19.50	179	4-tert-Butylcatechol, dimethyl ether	87	catechol derivate	** beaver castoreum 6
28.15	91	unidentified	< 80		

530 [1] Costello et al (2014), [2] Garner et al (2007), [3] Rigaill et al 2022, [4] Bernier et al (2000), [5] Smith et al
531 (2001), [6] Burger (2005), [7] Gallagher et al (2008), [8] Vaglio et al (2016), [9] Marneweck et al (2017)
532

* these studies identified a different derivate (3-Pentanol, 3-methyl)

** this study identified two different catechol derivates

<u>Discussion</u>

In the present study, we investigated to what extent the chemical composition of Barbary macaque

body odor is related to socially relevant traits and might thus have the potential to convey social

information. Our analyses support the existence of olfactory individual signatures, while support for

kin signatures differed between our analytical approaches. Chemical profiles were also related to

group membership, but we found little to no support that rank, sex or age affect chemical similarity

or detailed composition of scents emanating from the skin and fur of Barbary macaques.

Individual identity showed the strongest and most robust signature related to social information in

Barbary macaque body odor. Our results are in line with the extensive evidence for individually

distinct body odor across a range of mammals (e.g. Buesching et al., 2002; Caspers et al., 2008; Kwak

et al., 2008), including primates (e.g. Kulahci et al., 2014; Palagi and Dapporto, 2006; Rigaill et al.,

2022; Spence-Aizenberg et al., 2018; Scordato et al., 2007; reviewed in Smith, 2006). While the

majority of evidence thus far has come from scent-marking species, our results, along with studies in

Japanese macaques (Rigaill et al., 2022) and on human axillary odor (Penn et al., 2007), indicate that

individually distinct odors may also be present in catarrhine primates that do not show apparent

scent-marking. Individual identity explained only slightly less variance (r^2 of 0.2) in the chemical

profiles of Barbary macaques than in Japanese macaque urine (r^2 of 0.25) or vaginal secretions (r^2 of

0.29). Although the comparison between Japanese and Barbary macaques certainly needs to be

viewed in light of different methodologies and a larger and more heterogeneous sample of

individuals in Barbary macaques, it strongly suggests that odor emitted from the body surface may

carry identity information of similar magnitude as potentially more intense odor sources.

Furthermore, the present study shows that individually distinct information can be conveyed by the

chemicals emitted from the body surface, not just by scents deposited in the environment in the

form of scent marks or excretions.

The comparison of close vs. distant or non-kin based on overall chemical similarities suggested

olfactory signatures of both maternal and paternal relatedness, although paternal signatures were

statistically less robust. Units of close kin also showed a more similar chemical composition in owl

monkeys (*Aotus nancymaae* MacDonald et al., 2008; but see Spence-Aizenberg et al., 2018) and

rhesus macaques (Weiß et al., 2018a). Similar to our results in Barbary macaques, kin signatures in

rhesus macaques were only statistically robust among maternal, but not paternal half-siblings (Weiß

et al., 2018a). These patterns suggest that similarities in body odor between maternal relatives may

to some extent be mediated by frequent physical contact and/or a shared environment, which could

result in an exchange of odor-producing microbes, as e.g. suggested for group signatures in hyenas

(Theis et al., 2012). However, when treating relatedness as a continuous variable as well as

accounting for more other potential sources of variation in body odor, we found no support

whatsoever for relatedness predicting chemical similarity between individual odors. Possibly,

chemical profiles of surface body odor allow only a coarse, but no fine-scaled differentiation of

relatedness. Alternatively, what appeared to be kin signatures in the perMANOVAs may actually have

been variation related to other predictors that could not be as accurately controlled for in these

analyses. Furthermore, kin signatures may only be robustly expressed in certain contexts or seasons,

when kin discrimination is particularly important. For instance, kin signatures in glandular secretions

of ring-tailed lemurs (*Lemur catta*) were robustly expressed only in the breeding season, when they

may aid in inbreeding avoidance (Boulet et al., 2009; Charpentier et al., 2008), but we did not collect

odor samples during the mating season for logistic reasons.

The lack of robust chemical information about kinship involving the paternal line (at least outside the

mating season) is especially interesting with respect to the triadic interactions and care that Barbary

macaque males frequently engage in with infants. Particularly, male-infant interactions have been

suggested to be independent of paternity to the respective infant (Paul et al., 1996), which may be

due to low phenotypic similarity between father and offspring when the offspring is still in its infancy.

Studies in rhesus macaques testing exclusively adult subjects suggest that unfamiliar, paternal kin

discriminate relatives based on visual or acoustic phenotypic cues (Pfefferle et al., 2014a, 2014b).

Moreover, humans rating the facial similarity of rhesus macaque parents and offspring performed

above chance level (Kazem and Widdig, 2013) and better if the offspring were older (Kazem et al.,

2018). The paternal kin units sampled in the present study indeed comprised cases in which the

offspring was still rather young (2 years old), but in half of the cases all members of a paternal kin

unit were reproductively mature (i.e. 4 years or older). Hence, the chemical data presented here do

not point towards olfactory signatures alone being robust enough to allow Barbary macaque males

to reliably distinguish between own and unrelated offspring.

Regardless of the analytical approach, we detected no robust support for chemical profiles being

related to dominance classes or rank. Although we found a significant effect of dominance class on

the similarity between chemical profiles, the association was very weak, and ranks across the entire

dominance hierarchy were associated with neither chemical similarity nor the abundance of specific

compounds. Dominance-related odors have been described in various mammals (e.g. Burgener et al.,

2009; reviewed in Scordato et al., 2007), including male mandrills (*Mandrillus sphinx*, Setchell et al.,

2010; Vaglio et al., 2016), but were not detected in ring-tailed lemurs (Scordato et al., 2007) or

female mandrills (Setchell et al., 2010). In female rhesus macaques, individual rank had no effect on

overall chemical similarities, but affected the relative abundance of certain compounds (Weiß et al.,

2018a). As pointed out by Smith (2006), olfactory communication of dominance requires either a

specific dominance-related odor or olfactory individual signatures that can be associated with other

dominance-related information. Evidence for the latter comes from a study in ring-tailed lemurs, in

which males discriminated between the scents of dominant and subordinate individuals, but only if

they were familiar with the odor donors (Scordato and Drea, 2007). Hence, while our results provide

no robust evidence for dominance-related odor in either male or female Barbary macaques, the

detected individual signatures certainly open the possibility for a more indirect olfactory

communication of dominance among familiar individuals.

Based on behavioral and/or chemical evidence, the scents of numerous primates and other

mammals convey sex-specific (sex: e.g. Kean et al., 2011; Leclaire et al., 2017; Linklater et al., 2013;

Spence-Aizenberg et al., 2018; Weiß et al., 2018a) and age-specific information (e.g. Buesching et al.,

2002; Jänig et al., 2019; Kean et al., 2011; Kücklich et al., 2019; MacDonald et al., 2008; Marneweck

et al., 2017), although this need not always be the case (Jänig et al., 2019; also see review in Drea,

2015). Age, for instance, was related to chemical profiles in one subset of individuals in rhesus

macaques, but not in another (Weiß et al., 2018a). In the present study, sex and age, although

significantly associated with chemical similarities, explained very little variation in chemical profiles,

and none of the chemical compounds distinctly differed in abundance between sexes or showed

consistent changes with age. It is possible that sex and age information may actually be more

pronounced in a different context (e.g. mating), in other scent sources (e.g. urine) or other body

parts (e.g. the ano-genital region), all of which, along with differences in sampling and analytical

methods, may affect the expression of chemical signatures or our ability to detect them (see

Charpentier et al., 2008; Drea, 2015; Drea et al., 2013; Rigaill et al., 2022). Furthermore, sex

differences may have been attenuated in the study population due to the use of contraceptives in

60 % of the adult females (see below).

Group membership explained about 5 % of the variance in chemical similarities and affected the

relative abundance of 8 compounds. For most of these compounds, abundances were consistently

higher or lower in group F, but similar in the other two groups. Olfactory group signatures have been

described in a range of group-living mammals, including rhesus macaques (Henkel et al., 2015; Weiß

et al., 2018a), mandrills (Vaglio et al., 2016) and chimpanzees (Henkel and Setchell, 2018), with

explanations for the origins of these differences including the environment, diet and microbial

differences between groups (e.g. Leclaire et al., 2017; Spence-Aizenberg et al., 2018; Theis et al.,

2012). In species living in family groups, group signatures may further reflect relatedness among

individuals (e.g. owl monkeys, MacDonald et al., 2008). Relatedness is an unlikely explanation for the

group differences observed in the present study, because all groups comprised multiple matrilines

and average relatedness among genotyped focal animals was low both within (r = 0.04) and between

groups (r = 0.02, data not shown). The three groups also received the same supplemental food and

had spatially (but not temporally) overlapping home ranges, but there are subtle differences in the

vegetation in different areas of the park that could potentially affect natural food sources as well as

background volatiles or substances the animals get into contact with. Several of the tentatively

identified compounds could fit such an explanation. 3-methyl-2-pentanone and 2,4-dimethyl-3-

pentanol, for instance, could be microbial or plant products metabolized by the mammalian body

(see Charpentier et al., 2012) and similar compounds have been found in human feces before

(Costello et al., 2014). Furthermore, indenes have been described in plants (e.g. Mansur et al., 2018),

although we could not find a source specifically for 3-methyl-1H-indene, which was the best mass

spectral library match for the compound at RT 13.23 min. Alternatively, group differences could be

related to demographical and social differences between groups. The focal animals of group F

comprised a slightly higher proportion of females than groups C and H (57 % vs. 43 % and 32 %) and

were marginally younger (average age in group F 9.2 years vs. 11.7 and 11.3 yrs in groups C and H,

see supplement 2). However, because sex and age effects on body odor were very small (see above)

and statistically accounted for, they are unlikely to contribute much to the observed group

differences. Group F also is the largest and highest-ranking of the three groups, which could affect

stress levels or other physiological parameters (see overview in Dunbar, 2018) and thereby, body

odor (reviewed in Roberts et al., 2022), but we have no physiological data available to further assess

this possibility.

Main assets of our study certainly are the large numbers of samples and focal individuals, the semi-

natural environment and natural social conditions. We also were able to collect all samples non-

invasively, thereby greatly reducing the chance that odors were affected by sampling-related stress.

At the same time, the study faced several limitations, some of which we were able to account for

during data analysis, and others not. For instance, we collected samples from different body parts

that we could approach most easily, i.e. primarily the shoulders, flanks and thighs, and controlled for

the variation in sampled body regions during statistical analysis. However, it is difficult to assess how

relevant these body parts actually are for day-to-day information exchange. Recent observational

data in the study population suggests that sniffs directed at conspecifics frequently target the ano-

genital region of adult females in a mating context (Simon et al., 2024). The ano-genital region might

thus be a more appropriate sampling region at least for certain types of individuals and contexts, but

would have been feasible to sample only in a subset of very well-habituated females at frequent

intervals. As the social information detectable in animal scents may differ between scent sources,

even within species (see Drea, 2015; Rigaill et al., 2022), it is possible that results would have differed

if we had sampled other body regions or scent sources such as urine.

Our data collection periods spanned much of the year, but for logistic reasons we did not collect data

during most of the park's winter break, which corresponds to the peak of the mating season in the

study population. Indeed, the season of the year did affect chemical profiles, even more so than the

socially relevant variables investigated. In particular, seasonal variation may have affected the

volatile background in the environment, which we could at least in part account for by using

environmental control samples. Furthermore, aspects such as reproductive physiology, diet, ambient

temperature or state of the fur (winter vs. summer coat) may additionally affect the chemical

compounds emitted by the body and how readily they become available to recipients (or sampling

tube). Such aspects are likely to be at the base of the detected seasonal changes in chemical profiles,

and we statistically accounted for them in our analyses of socially related olfactory signatures by

using season of the year as a control variable. What remains open is whether any of the investigated

effects might have been more pronounced during the mating season, as shown, e.g., for kin

signatures in ring-tailed lemurs (Boulet et al., 2009; Charpentier et al., 2008).

Another aspect to consider is that the size of the study population is managed via hormonal

contraception, and 60 % of the sampled females received contraception during part or all of the

study period. Studies on the effects of hormonal contraception on primate body odor are scarce and

contradicting: hormonal contraception using medroxyprogesterone acetate has previously been

shown to alter the chemical composition of female scents in ring-tailed lemurs (Crawford et al., 2011)

but not in owl monkeys (Spence-Aizenberg et al., 2018). Male mandrills showed more flehmen

responses towards scent samples from cycling females than from females contracepted with

etonogestrel (Charpentier et al., 2013), which is the active ingredient of the implants used in the

studied Barbary macaques. However, while contraception also tended to modulate age-specific

differences in ano-genital sniffs by male Barbary macaques, it did not affect subsequent sexual

behavior in our studied population (Simon et al., 2024). As we did not assess individual attributes

immediately related to mating and female fertility, contraception might be considered as a source of

noise in the data. Unlike several other sources of noise, however, we could not account for

contraception statistically given that it applied only to a minority of samples and individuals. Overall,

it is thus possible that certain signatures, in particular those of sex, might have been more

pronounced in a sample without females receiving contraception.

Our results also suggest some variation in chemical profiles related to sampling and GC-MS

measurement, whereby different variables affected different aspects of the composition: chemical

similarities between samples were greater if samples were collected closer in time, and to a lesser

extent also when analyzed within the same GC-MS batch. In contrast, variables such as the sampling

distance, sampling volume or the sampled body part had little effect on chemical similarities of

samples, but appeared to have some effect on the relative abundance of certain compounds.

However, while we were unable to eliminate these sources of variation in the study design, we

statistically accounted for them in all mixed model analyses.

Finally, results need to be put into perspective of what we can measure vs. what animals can

perceive. Statistically detectable associations between chemical profiles and socially relevant traits

show the potential for olfactory information transfer, but we do not know if or how animals use this

information. Animal sensory systems have evolved for the detection of relevant information and may

thus be superior to even the most modern of our analytical techniques, at least for certain

compounds (reviewed in Maidodou et al., 2023). Accordingly, it may be possible that animals are

able to pick up upon even very subtle differences in odors, or differences that our analyses are

unable to pick up at all, especially as we are unlikely to sample and detect all chemical substances

emitted by an animal's body across the possible ranges of volatility, polarity etc. (Drea et al., 2013)

despite the constant improvements in sampling and analytical methods. Hence, future studies should

combine chemical data with (experimental) assessments of behavioral and physiological reactions

towards conspecific odors across a range of contexts to obtain a detailed understanding of the

complex role of olfaction in primate social interactions. Particularly under natural and semi-natural

conditions, controlled presentations of conspecific odors are challenging to implement (e.g. Ritter,

2022), as stimuli need to be presented in a manner that is salient and realistic enough to elicit

interest and closer inspection without leading to quick habituation. Promising next steps would be

multi-modal experiments and experimental manipulation of *in vivo* odors to investigate how

olfactory discrimination of social information falls into place between the other senses, i.e. whether

macaques are able to discriminate such information based on odors alone, and/or if olfactory

signatures improve the reliability of other sensory cues.

In conclusion, our results add to the limited evidence that socially relevant information may not only

be encoded in scent secretions and excretions, but also in the volatile and semi-volatile components

emanating from the body surface of primates. Based on chemical profiles alone, i.e. without knowing

what the monkeys can make of it, data suggest that Barbary macaque body odor could directly

contribute to recognizing individuals and groups, while olfactory signatures may not be pronounced

or robust enough to allow differentiating kinship, rank, sex or age of conspecifics from surface body

odor alone. However, given the central role of individual recognition in forming and maintaining

social relationships, surface body odor could potentially still contribute to managing social

relationships, if monkeys associate social knowledge obtained via other sensory modalities or direct

social interactions with olfactory individual signatures.

Acknowledgements

We thank Roland & Mamisolo Hilgartner and the staff of the Affenberg Salem for continuous support

and Ellen Merz for permission to conduct this study. We further thank Roger Mundry for an

introduction to multi-membership models and help with the centred log-ratio transformation.

Hendrikje Westphal established the methods for genetic paternity analyses and helped with faecal

sample collection. Stefanie Bley managed the paternity analyses. Antonia Krüger and Anja Schiffner

conducted DNA extractions and PCRs and helped with GC-MS measurements and TD tube cleaning.

Annika Freudiger helped with the implementation of the TRACE software. Ramona Oehme and Susan

Billig in the mass spectrometry core facility at the Faculty of Chemistry and Mineralogy at Leipzig

University provided technical support in GC-MS maintenance.

Funding

This work was supported by the German Research Foundation, DFG (grant no. SCHL 2011/2-1

awarded to B. M. Weiß). The GC-MS instrument was funded by the European Fund for Regional

Structure Development, EFRE ("Europe funds Saxony", grant #100195810 to A. Widdig).

References

Arandjelovic, M., Guschanski, K., Schubert, G., Harris, T.R., Thalmann, O., Siedel, H., Vigilant, L., 2009.
Two-step multiplex polymerase chain reaction improves the speed and accuracy of genotyping
using DNA from noninvasive and museum samples. Mol. Ecol. Resour. 9, 28–36.
https://doi.org/10.1111/j.1755-0998.2008.02387.x

Bates, L.A., Sayialel, K.N., Njiraini, N.W., Poole, J.H., Moss, C.J., Byrne, R.W., 2008. African elephants
have expectations about the locations of out-of-sight family members. Biol. Lett. 4, 34–36.
https://doi.org/10.1098/rsbl.2007.0529

Berghänel, A., Ostner, J., Schröder, U., Schülke, O., 2011a. Social bonds predict future cooperation in
male Barbary macaques, Macaca sylvanus. Anim. Behav. 81, 1109–1116.
https://doi.org/10.1016/j.anbehav.2011.02.009

Berghänel, A., Ostner, J., Schülke, O., 2011b. Coalitions destabilize dyadic dominance relationships in
male Barbary macaques (Macaca sylvanus). Behaviour 148, 1256–1274.
https://doi.org/10.1163/000579511X600628

Bernier, U.R., Kline, D.L., Barnard, D.R., Schreck, C.E., Yost, R.A., 2000. Analysis of Human Skin
Emanations by Gas Chromatography/Mass Spectrometry. 2. Identification of Volatile
Compounds That Are Candidate Attractants for the Yellow Fever Mosquito (*Aedes aegypti*).
Anal. Chem. 72, 747–756. https://doi.org/10.1021/ac990963k

Boogaart, K.G. van den, Tolosana-Delgado, R., 2013. Analyzing Compositional Data with R. Springer,
Heidelberg.

Boulet, M., Charpentier, M., Drea, C., 2009. Decoding an olfactory mechanism of kin recognition and
inbreeding avoidance in a primate. BMC Evol. Biol. 9, 281. https://doi.org/10.1186/1471-2148-9-
281

Brauch, K., Hodges, K., Engelhardt, A., Fuhrmann, K., Shaw, E., Heistermann, M., 2008. Sex-specific
reproductive behaviours and paternity in free-ranging Barbary macaques (*Macaca sylvanus*).
Behav. Ecol. Sociobiol. 62, 1453–1466. https://doi.org/10.1007/s00265-008-0575-7

Buesching, C.D., Waterhouse, J.S., Macdonald, D.W., 2002. Gas-chromatographic analyses of the
subcaudal gland secretion of the European badger (*Meles meles*) part II: time-related variation in
the individual-specific composition. J. Chem. Ecol. 28, 57–69.

Burgener, N., Dehnhard, M., Hofer, H., East, M.L., 2009. Does anal gland scent signal identity in the
spotted hyaena? Anim. Behav. 77, 707–715. https://doi.org/10.1016/j.anbehav.2008.11.022

Burger, B.V., 2005. Mammalian Semiochemicals. Top. Curr. Chem. 240, 231–278.
https://doi.org/10.1007/b98318

Bürkner, P.-C., 2017. brms: An R Package for Bayesian Multilevel Models Using Stan. J. Stat. Softw.
80, 1–28. https://doi.org/10.18637/jss.v080.i01

Caspers, B., Franke, S., Voigt, C.C., 2008. The Wing-Sac Odour of Male Greater Sac-Winged Bats
Saccopteryx bilineata (Emballonuridae) as a Composite Trait: Seasonal and Individual

Differences, in: Hurst, J.L., Beynon, R.J., Roberts, S.C., Wyatt, T.D. (Eds.), Chemical Signals in Vertebrates 11. Springer New York, New York, pp. 151–160.

Charpentier, M.J.E., Barthes, N., Proffit, M., Bessière, J.-M., Grison, C., 2012. Critical thinking in the chemical ecology of mammalian communication: roadmap for future studies. Funct. Ecol. 26, 769–774. https://doi.org/10.1111/j.1365-2435.2012.01998.x

Charpentier, M.J.E., Boulet, M., Drea, C.M., 2008. Smelling right: the scent of male lemurs advertises genetic quality and relatedness. Mol. Ecol. 17, 3225–3233. https://doi.org/10.1111/j.1365-294X.2008.03831.x

Charpentier, M.J.E., Mboumba, S., Ditsoga, C., Drea, C.M., 2013. Nasopalatine Ducts and Flehmen Behavior in the Mandrill: Reevaluating Olfactory Communication in Old World Primates. Am. J. Primatol. 75, 703–714. https://doi.org/10.1002/ajp.22146

Cheney, D.L., Seyfarth, R.M., 1999. Recognition of other individuals' social relationships by female baboons. Anim. Behav. 58, 67–75. https://doi.org/10.1006/anbe.1999.1131

Cheney, D.L., Seyfarth, R.M., Palombit, R., 1996. The function and mechanisms underlying baboon "contact" barks. Anim. Behav. 52, 507–518.

Chung, H., Beretvas, S.N., 2012. The impact of ignoring multiple membership data structures in multilevel models. Br. J. Math. Stat. Psychol. 65, 185–200. https://doi.org/10.1111/j.2044-8317.2011.02023.x

Costello, B. de L., Amann, A., Al-Kateb, H., Flynn, C., Filipiak, W., Khalid, T., Osborne, D., Ratcliffe, N.M., 2014. A review of the volatiles from the healthy human body. J. Breath Res. 8, 014001. https://doi.org/10.1088/1752-7155/8/1/014001

Crawford, J.C., Boulet, M., Drea, C.M., 2011. Smelling wrong: hormonal contraception in lemurs alters critical female odour cues. Proc. R. Soc. B Biol. Sci. 278, 122–130. https://doi.org/10.1098/rspb.2010.1203

de Turckheim, G., Merz, E., 1984. Breeding Barbary macaques in outdoor open enclosures, in: Fa, J.F. (Ed.), The Barbary Macaque: A Case Study in Conservation. Plenum Press, New York, pp. 241–261.

delBarco-Trillo, J., Sacha, C.R., Dubay, G.R., Drea, C.M., 2012. Eulemur, me lemur: the evolution of scent-signal complexity in a primate clade. Philos. Trans. R. Soc. Lond. B Biol. Sci. 367, 1909–1922. https://doi.org/10.1098/rstb.2011.0225

Drea, C.M., 2015. D'scent of man: A comparative survey of primate chemosignaling in relation to sex. Horm. Behav. 68, 117–133. https://doi.org/10.1016/j.yhbeh.2014.08.001

Drea, C.M., Boulet, M., Delbarco-Trillo, J., Greene, L.K., Sacha, C.R., Goodwin, T.E., Dubay, G.R., 2013. The "Secret" in Secretions: Methodological Considerations in Deciphering Primate Olfactory Communication. Am. J. Primatol. 75, 621–642. https://doi.org/10.1002/ajp.22143

Drea, C.M., Scordato, E.S., 2008. Olfactory Communication in the Ringtailed Lemur (Lemur catta): Form and Function of Multimodal Signals, in: Hurst, J.L., Beynon, R.J., Roberts, S.C., Wyatt, T.D. (Eds.), Chemical Signals in Vertebrates. Springer, New York, NY, pp. 91–102. https://doi.org/10.1007/978-0-387-73945-8_8

Dunbar, R.I.M., 2018. Social structure as a strategy to mitigate the costs of group living: a comparison of gelada and guereza monkeys. Anim. Behav. 136, 53–64. https://doi.org/10.1016/j.anbehav.2017.12.005

Fischer, J., 2004. Emergence of individual recognition in young macaques. Anim. Behav. 67, 655–661.

Fontani, S., Kaburu, S.S.K., Marliani, G., Accorsi, P.A., Vaglio, S., 2022. Anogenital scent-marking signals fertility in a captive female Alaotran gentle lemur. Front. Vet. Sci. 9.

Fox, J., Weisberg, S., 2019. An {R} companion to applied regression. Sage, Thousand Oaks CA.

Freeman, N.J., Pasternak, G.M., Rubi, T.L., Barrett, L., Henzi, S.P., 2012. Evidence for scent marking in vervet monkeys? Primates 53, 311–315.

Gallagher, M., Wysocki, C.J., Leyden, J.J., Spielman, A.I., Sun, X., Preti, G., 2008. Analyses of volatile
organic compounds from human skin. Br. J. Dermatol. 159, 780–791.
https://doi.org/10.1111/j.1365-2133.2008.08748.x

Garner, C.E., Smith, S., de Lacy Costello, B., White, P., Spencer, R., Probert, C.S.J., Ratcliffe, N.M.,
2007. Volatile organic compounds from feces and their potential for diagnosis of gastrointestinal
disease. FASEB J. 21, 1675–1688. https://doi.org/10.1096/fj.06-6927com

Goffe, A.S., Zinner, D., Fischer, J., 2016. Sex and friendship in a multilevel society: behavioural
patterns and associations between female and male Guinea baboons. Behav. Ecol. Sociobiol. 70,
323–336. https://doi.org/10.1007/s00265-015-2050-6

Hammerschmidt, K., Fischer, J., 1998. Maternal discrimination of offspring vocalizations in Barbary
macaques (*Macaca sylvanus*). Primates 39, 231–236.

Henkel, S., Lambides, A., Berger, A., Thomsen, R., Widdig, A., 2015. Rhesus macaques (*Macaca*
mulatta) recognize group membership via olfactory cues alone. Behav. Ecol. Sociobiol. 69, 2019–
2034. https://doi.org/10.1007/s00265-015-2013-y

Henkel, S., Setchell, J.M., 2018. Group and kin recognition via olfactory cues in chimpanzees (Pan
troglodytes). Proc R Soc B 285, 20181527. https://doi.org/10.1098/rspb.2018.1527

Heymann, E.W., 2006a. Scent marking strategies of new world primates. Am. J. Primatol. 68, 650–
661.

Heymann, E.W., 2006b. The neglected sense-olfaction in primate behavior, ecology, and evolution.
Am. J. Primatol. 68, 519–524. https://doi.org/10.1002/ajp.20249

Jamil, T., Ozinga, W.A., Kleyer, M., ter Braak, C.J.F., 2013. Selecting traits that explain species–
environment relationships: a generalized linear mixed model approach. J. Veg. Sci. 24, 988–
1000. https://doi.org/10.1111/j.1654-1103.2012.12036.x

Janda, E.D., Perry, K.L., Hankinson, E., Walker, D., Vaglio, S., 2019. Sex differences in scent-marking in
captive red-ruffed lemurs. Am. J. Primatol. 0, e22951. https://doi.org/10.1002/ajp.22951

Jänig, S., Weiß, B.M., Birkemeyer, C., Widdig, A., 2019. Comparative chemical analysis of body odor in
great apes. Am. J. Primatol. 0, e22976. https://doi.org/10.1002/ajp.22976

Kawamura, S., 2016. Color vision diversity and significance in primates inferred from genetic and field
studies. Genes Genomics 38, 779–791. https://doi.org/10.1007/s13258-016-0448-9

Kazem, A.J.N., Barth, Y., Pfefferle, D., Kulik, L., Widdig, A., 2018. Parent–offspring facial resemblance
increases with age in rhesus macaques. Proc R Soc B 285, 20181208.
https://doi.org/10.1098/rspb.2018.1208

Kazem, A.J.N., Widdig, A., 2013. Visual phenotype matching: cues to paternity are present in rhesus
macaque faces. PLoS ONE 8, e55846. https://doi.org/10.1371/journal.pone.0055846

Kean, E.F., Müller, C.T., Chadwick, E.A., 2011. Otter scent signals age, sex, and reproductive status.
Chem. Senses 36, 555–564. https://doi.org/10.1093/chemse/bjr025

Kücklich, M., Möller, M., Marcillo, A., Einspanier, A., Weiß, B.M., Birkemeyer, C., Widdig, A., 2017.
Different Methods for Volatile Sampling in Mammals. PLOS ONE 12, e0183440.
https://doi.org/10.1371/journal.pone.0183440

Kücklich, M., Weiß, B.M., Birkemeyer, C., Einspanier, A., Widdig, A., 2019. Chemical cues of female
fertility states in a non-human primate. Sci. Rep. 9, 1–12. https://doi.org/10.1038/s41598-019-
50063-w

Kuester, J., Paul, A., 1999. Male migration in Barbary macaques (*Macaca sylvanus*) at Affenberg
Salem. Int. J. Primatol. 20, 85–106.

Kuester, J., Paul, A., Arnemann, J., 1994. Kinship, familiarity and mating avoidance in Barbary
macaques (*Macaca sylvanus*). Anim. Behav. 48, 1183–1194.

Kulahci, I.G., Drea, C.M., Rubenstein, D.I., Ghazanfar, A.A., 2014. Individual recognition through
olfactory-auditory matching in lemurs. Proc. R. Soc. B Biol. Sci. 281, 20140071.
https://doi.org/10.1098/rspb.2014.0071

Kwak, J., Willse, A., Matsumura, K., Curran Opiekun, M., Yi, W., Preti, G., Yamazaki, K., Beauchamp,
G.K., 2008. Genetically-Based Olfactory Signatures Persist Despite Dietary Variation. PLoS ONE 3,
e3591. https://doi.org/10.1371/journal.pone.0003591

Laska, M., Seibt, A., 2002a. Olfactory sensitivity for aliphatic alcohols in squirrel monkeys and pigtail
macaques. J. Exp. Biol. 205, 1633–1643.

Laska, M., Seibt, A., 2002b. Olfactory sensitivity for aliphatic esters in squirrel monkeys and pigtail
macaques. Behav. Brain Res. 134, 165–174.

Leclaire, S., Jacob, S., Greene, L.K., Dubay, G.R., Drea, C.M., 2017. Social odours covary with bacterial
community in the anal secretions of wild meerkats. Sci. Rep. 7, 3240.
https://doi.org/10.1038/s41598-017-03356-x

Linklater, W.L., Mayer, K., Swaisgood, R.R., 2013. Chemical signals of age, sex and identity in black
rhinoceros. Anim. Behav. 85, 671–677. https://doi.org/10.1016/j.anbehav.2012.12.034

Loos, H.M., Schaal, B., Pause, B.M., Smeets, M.A.M., Ferdenzi, C., Roberts, S.C., de Groot, J., Lübke,
K.T., Croy, I., Freiherr, J., Bensafi, M., Hummel, T., Havlíček, J., 2023. Past, Present, and Future of
Human Chemical Communication Research. Perspect. Psychol. Sci. 17456916231188147.
https://doi.org/10.1177/17456916231188147

MacDonald, E.A., Fernandez-Duque, E., Evans, S., Hagey, L.R., 2008. Sex, age, and family differences
in the chemical composition of owl monkey (*Aotus nancymaae*) subcaudal scent secretions. Am.
J. Primatol. 70, 12–18. https://doi.org/10.1002/ajp.20450

Maidodou, L., Clarot, I., Leemans, M., Fromantin, I., Marchioni, E., Steyer, D., 2023. Unraveling the
potential of breath and sweat VOC capture devices for human disease detection: a systematic-
like review of canine olfaction and GC-MS analysis. Front. Chem. 11, 1282450.
https://doi.org/10.3389/fchem.2023.1282450

Majolo, B., Lehmann, J., de Bortoli Vizioli, A., Schino, G., 2012. Fitness-related benefits of dominance
in primates. Am. J. Phys. Anthropol. 147, 652–660. https://doi.org/10.1002/ajpa.22031

Mansur, M.A.A., Siddiqi, M.M.A., Akbor, M.A., Saha, K., 2018. Phytochemical Screening and GC-MS
Chemical Profiling of Ethyl Acetate Extract of Seed and Stem of Anethum sowa Linn. Dhaka Univ.
J. Pharm. Sci. 16, 187–194. https://doi.org/10.3329/dujps.v16i2.35256

Marneweck, C., Jürgens, A., Shrader, A.M., 2017. Dung odours signal sex, age, territorial and oestrous
state in white rhinos. Proc R Soc B 284, 20162376. https://doi.org/10.1098/rspb.2016.2376

Mehta, P.H., Josephs, R.A., 2010. Testosterone and cortisol jointly regulate dominance: Evidence for
a dual-hormone hypothesis. Horm. Behav. 58, 898–906.
https://doi.org/10.1016/j.yhbeh.2010.08.020

Minkner, M.M.I., Young, C., Amici, F., McFarland, R., Barrett, L., Grobler, J.P., Henzi, S.P., Widdig, A.,
2018. Assessment of male reproductive skew via highly polymorphic STR markers in wild vervet
monkeys, *Chlorocebus pygerythrus*. J. Hered. 109, 780–790.
https://doi.org/10.1093/jhered/esy048

Moreira, L.A.A., Pessoa, D.M.A., Sousa, M.B.C., 2013. Socio-sexual communication: A review of the
sensory modalities used by non-human primates. Oecologia Aust. 17, 303–319.

Naud, A., Chailleux, E., Kestens, Y., Bret, C., Desjardins, D., Petit, O., Ngoubangoye, B., Sueur, C.,
2016. Relations between Spatial Distribution, Social Affiliations and Dominance Hierarchy in a
Semi-Free Mandrill Population. Front. Psychol. 7. https://doi.org/10.3389/fpsyg.2016.00612

Neumann, C., Duboscq, J., Dubuc, C., Ginting, A., Irwan, A.M., Agil, M., Widdig, A., Engelhardt, A.,
2011. Assessing dominance hierarchies: Validation and advantages of progressive evaluation
with Elo-rating. Anim. Behav. 82, 911–921. https://doi.org/10.1016/j.anbehav.2011.07.016

Newton-Fisher, N.E., 2017. Modeling Social Dominance: Elo-Ratings, Prior History, and the Intensity of Aggression. Int. J. Primatol. 38, 427–447. https://doi.org/10.1007/s10764-017-9952-2

Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M.D., Durand, S., Evangelista, H.B.A., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M.O., Lahti, L., McGlinn, D., Ouellette, M.-H., Cunha, E.R., Smith, T., Stier, A., Braak, C.J.F.T., Weedon, J., 2022. vegan: Community Ecology Package.

Ostner, J., Schülke, O., 2014. The evolution of social bonds in primate males. Behaviour 151, 871–906. https://doi.org/10.1163/1568539X-00003191

Palagi, E., Dapporto, L., 2006. Beyond Odor Discrimination: Demonstrating Individual Recognition by Scent in *Lemur Catta*. Chem. Senses 31, 437–443. https://doi.org/10.1093/chemse/bjj048

Paul, A., Kuester, J., 1987. Sex ratio adjustment in a seasonally breeding primate species: evidence from Barbary macaque Population at Affenberg Salem. Ethology 74, 117–132.

Paul, A., Kuester, J., Arnemann, J., 1996. The sociobiology of male-infant interactions in Barbary macaques, *Macaca sylvanus*. Anim. Behav. 51, 155–170. https://doi.org/10.1006/anbe.1996.0013

Paul, A., Kuester, J., Arnemann, J., 1992. DNA fingerprinting reveals that infant care by male Barbary macaques (Macaca sylvanus) is not paternal investment. Folia Primatol. (Basel) 58, 93–98.

Penn, D.J., Oberzaucher, E., Grammer, K., Fischer, G., Soini, H.A., Wiesler, D., Novotny, M.V., Dixon, S.J., Xu, Y., Brereton, R.G., 2007. Individual and gender fingerprints in human body odour. J. R. Soc. Interface 4, 331–340. https://doi.org/10.1098/rsif.2006.0182

Pfefferle, D., Kazem, A.J.N., Brockhausen, R.R., Ruiz-Lambides, A.V., Widdig, A., 2014a. Monkeys spontaneously discriminate their unfamiliar paternal kin under natural conditions using facial cues. Curr. Biol. 24, 1806–1810. https://doi.org/10.1016/j.cub.2014.06.058

Pfefferle, D., Ruiz-Lambides, A.V., Widdig, A., 2014b. Female rhesus macaques discriminate unfamiliar paternal sisters in playback experiments – support for acoustic phenotype matching. Proc. R. Soc. B 281, 1–9.

Prud'homme, J., Chapais, B., 1993. Aggressive interventions and matrilineal dominance relations in semifree-ranging Barbary macaques (Macaca sylvanus). Primates 34, 271–283.

R Core Team, 2024. R: A Language and Environment for Statistical Computing.

Rendall, D., Rodman, P.S., Emond, R.E., 1996. Vocal recognition of individuals and kin in free-ranging rhesus monkeys. Anim. Behav. 51, 1007–1015. https://doi.org/10.1006/anbe.1996.0103

Rigaill, L., Vaglio, S., Setchell, J.M., Suda-Hashimoto, N., Furuichi, T., Garcia, C., 2022. Chemical cues of identity and reproductive status in Japanese macaques. Am. J. Primatol. 84, e23411. https://doi.org/10.1002/ajp.23411

Ritter, N., 2022. Multimodal perception of social stimuli in Barbary macaques at the Affenberg Salem (MSc thesis). Leipzig University, Leipzig.

Roberts, S.C., Fialová, J.T., Sorokowska, A., Langford, B., Sorokowski, P., Třebický, V., Havlíček, J., 2022. Emotional expression in human odour. Evol. Hum. Sci. 1–35. https://doi.org/10.1017/ehs.2022.44

Roubová, V., Konečná, M., Šmilauer, P., Wallner, B., 2015. Whom to Groom and for What? Patterns of Grooming in Female Barbary Macaques (*Macaca sylvanus*). PLoS ONE 10, e0117298. https://doi.org/10.1371/journal.pone.0117298

Ruiz-Lambides, A.V., Weiß, B.M., Kulik, L., Widdig, A., 2018. Which male and female characteristics influence the probability of extragroup paternities in rhesus macaques, *Macaca mulatta*? Anim. Behav. 140, 119–127. https://doi.org/10.1016/j.anbehav.2018.04.018

Scheiber, I.B.R., Hohnstein, A., Kotrschal, K., Weiß, B.M., 2011. Juvenile Greylag Geese (Anser anser) Discriminate between Individual Siblings. PLOS ONE 6, e22853. https://doi.org/10.1371/journal.pone.0022853

Schell, A., Rieck, K., Schell, K., Hammerschmidt, K., Fischer, J., 2011. Adult but not juvenile Barbary macaques spontaneously recognize group members from pictures. Anim. Cogn. 14, 503–509. https://doi.org/10.1007/s10071-011-0383-8

Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients. Methods Ecol. Evol. 1, 103–113. https://doi.org/10.1111/j.2041-210X.2010.00012.x

Schoof, V.A.M., Jack, K.M., Carnegie, S.D., 2011. Rise to Power: A Case Study of Male Fecal Androgen and Cortisol Levels before and after a Non-Aggressive Rank Change in a Group of Wild White-Faced Capuchins (Cebus capucinus). Folia Primatol. (Basel) 82, 299–307. https://doi.org/10.1159/000337220

Scordato, E.S., Drea, C.M., 2007. Scents and sensibility: information content of olfactory signals in the ringtailed lemur, Lemur catta. Anim. Behav. 73, 301–314. https://doi.org/10.1016/j.anbehav.2006.08.006

Scordato, E.S., Dubay, G., Drea, C.M., 2007. Chemical composition of scent marks in the ringtailed lemur (*Lemur catta*): glandular differences, seasonal variation, and individual signatures. Chem. Senses 32, 493–504. https://doi.org/10.1093/chemse/bjm018

Setchell, J.M., Vaglio, S., Moggi-Cecchi, J., Boscaro, F., Calamai, L., Knapp, L.A., 2010. Chemical composition of scent-gland secretions in an old world monkey (Mandrillus sphinx): influence of sex, male status, and individual identity. Chem. Senses 35, 205–220. https://doi.org/10.1093/chemse/bjp105

Seyfarth, R.M., Cheney, D.L., Bergman, T.J., 2005. Primate social cognition and the origins of language. Trends Cogn. Sci. 9, 264–266. https://doi.org/10.1016/j.tics.2005.04.001

Silk, J.B., 2001. Ties that bond: the role of kinship in primate societies, in: Stone, L. (Ed.), New Directions in Anthropological Kinship. Rowman & Littlefield Publishers, Inc., Boulder, pp. 71–92.

Simon, M., Widdig, A., Weiß, B.M., 2024. Sniffing behavior of semi free-ranging Barbary macaques (Macaca sylvanus). Am. J. Primatol. 86, e23611. https://doi.org/10.1002/ajp.23611

Small, M., 1990. Promiscuity in Barbary macaques (*Macaca sylvanus*). Am. J. Primatol. 20, 267–282.

Smith, T., 2006. Individual olfactory signatures in common marmosets (Callithrix jacchus). Am. J. Primatol. 68, 585–604. https://doi.org/10.1002/ajp.20254

Smith, T., Tomlinson, A., Mlotkiewicz, J., Abbott, D., 2001. Female marmoset monkeys (*Callithrix jacchus*) can be identified from the chemical composition of their scent marks. Chem. Senses 26, 449–458.

Smith, T.D., Laitman, J.T., Bhatnagar, K.P., 2014. The shrinking anthropoid nose, the human vomeronasal organ, and the language of anatomical reduction. Anat. Rec. 2196–2204. https://doi.org/DOI: 10.1002/ar.23035

Spence-Aizenberg, A., Kimball, B.A., Williams, L.E., Fernandez-Duque, E., 2018. Chemical composition of glandular secretions from a pair-living monogamous primate: Sex, age, and gland differences in captive and wild owl monkeys (Aotus spp.). Am. J. Primatol. 80, e22730. https://doi.org/10.1002/ajp.22730

Stevenson, R.J., 2010. An Initial Evaluation of the Functions of Human Olfaction. Chem. Senses 35, 3–20. https://doi.org/10.1093/chemse/bjp083

Stoffel, M.A., Caspers, B.A., Forcada, J., Giannakara, A., Baier, M., Eberhart-Phillips, L., Müller, C., Hoffman, J.I., 2015. Chemical fingerprints encode mother–offspring similarity, colony membership, relatedness, and genetic quality in fur seals. Proc. Natl. Acad. Sci. 112, E5005–E5012. https://doi.org/10.1073/pnas.1506076112

Tang-Martinez, Z., 2001. The mechanisms of kin discrimination and the evolution of kin recognition in vertebrates: a critical re-evaluation. Behav. Processes 53, 21–40. https://doi.org/10.1016/S0376-6357(00)00148-0

Theis, K.R., Schmidt, T.M., Holekamp, K.E., 2012. Evidence for a bacterial mechanism for group-specific social odors among hyenas. Sci. Rep. 2, 1–8. https://doi.org/10.1038/srep00615

Tibbetts, E.A., Dale, J., 2007. Individual recognition: it is good to be different. Trends Ecol. Evol. 22, 529–537. https://doi.org/10.1016/j.tree.2007.09.001

Vaglio, S., Minicozzi, P., Kessler, S.E., Walker, D., Setchell, J.M., 2021. Olfactory signals and fertility in olive baboons. Sci. Rep. 11, 8506. https://doi.org/10.1038/s41598-021-87893-6

Vaglio, S., Minicozzi, P., Romoli, R., Boscaro, F., Pieraccini, G., Moneti, G., Moggi-Cecchi, J., 2016. Sternal Gland Scent-Marking Signals Sex, Age, Rank, and Group Identity in Captive Mandrills. Chem. Senses 41, 177–186. https://doi.org/10.1093/chemse/bjv077

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B., Bürkner, P.-C., 2021. Rank-Normalization, Folding, and Localization: An Improved R̂ for Assessing Convergence of MCMC (with Discussion). Bayesian Anal. 16, 667–718. https://doi.org/10.1214/20-BA1221

Veilleux, C.C., Dominy, N.J., Melin, A.D., 2022. The sensory ecology of primate food perception, revisited. Evol. Anthropol. Issues News Rev. 31, 281–301. https://doi.org/10.1002/evan.21967

von Segesser, F., Menard, N., Gaci, B., Martin, R.D., 1999. Genetic differentiation within and between isolated Algerian subpopulations of Barbary macaques (Macaca sylvanus): evidence from microsatellites. Mol. Ecol. 8, 433–442.

Ward, A., Webster, M., 2016. Sociality: The Behaviour of Group-Living Animals. Springer.

Weiß, B.M., Kücklich, M., Thomsen, R., Henkel, S., Jänig, S., Kulik, L., Birkemeyer, C., Widdig, A., 2018a. Chemical composition of axillary odorants reflects social and individual attributes in rhesus macaques. Behav. Ecol. Sociobiol. 72, 65. https://doi.org/10.1007/s00265-018-2479-5

Weiß, B.M., Marcillo, A., Manser, M., Holland, R., Birkemeyer, C., Widdig, A., 2018b. A non-invasive method for sampling the body odour of mammals. Methods Ecol. Evol. 9, 420–429. https://doi.org/10.1111/2041-210X.12888

Westphal, H., 2020. Paternity assessment in Barbary macaques (BSc Thesis). Leipzig University, Leipzig.

Westphal, H., Freudiger, A., Gatter, T., Stadler, P.F., Widdig, A., 2023. TRACE - Tool for pedigree Relatedness Analysis and Coefficient Estimation.

Widdig, A., 2007. Paternal kin discrimination: The evidence and likely mechanisms. Biol. Rev. 82, 319–334. https://doi.org/10.1111/j.1469-185X.2007.00011.x

Wyatt, T., 2014. Pheromones and animal behavior: Chemical signals and signatures. Cambridge University Press.

Yorzinski, J.L., 2017. The cognitive basis of individual recognition. Curr. Opin. Behav. Sci., Comparative cognition 16, 53–57. https://doi.org/10.1016/j.cobeha.2017.03.009

Young, C., Majolo, B., Schülke, O., Ostner, J., 2014. Male social bonds and rank predict supporter selection in cooperative aggression in wild Barbary macaques. Anim. Behav. 95, 23–32. https://doi.org/10.1016/j.anbehav.2014.06.007

Zetzsche, M., Weiß, B.M., Kücklich, M., Stern, J., Birkemeyer, C., Widdig, A., Penke, L., 2024. Combined perceptual and chemical analyses show no compelling evidence for ovulatory cycle shifts in women's axillary odour. Proc. R. Soc. B Biol. Sci. 291, 20232712. https://doi.org/10.1098/rspb.2023.2712