Dynamic mean and variance of microparasite load give key insights into population dynamics and underlying mechanisms

<u>Abstract</u>

Heterogeneity among individuals, in number of parasites, body size, etc., can have critical interactions with population dynamics. We tease these out with a relatively simple model of such interactions in a case study of microparasite load with empirically supported assumptions to ask a key question. How does variance in microparasite load interact with population-level dynamics? We show how the mean and variance of infection load vary throughout an epidemic. Further, we show how mean and variance have mutual negative feedbacks on each other mediated by high disease-induced death rates at high loads. Helpfully, we find that mean and variance provide information into underlying processes as well. Population-level trends in the mean and variance reveal underlying trends in within-host processes, e.g., differentiating host evolution of defence that manifests as tolerance, constitutive resistance, inducible resistance, or acquired resistance. Such inference may be critical for managing endangered Mountain Yellow Legged Frog populations recovering from *Batrachochytrium dendrobatidis* epidemics. Lastly, we demonstrate the impact of pathogen load variance on host fitness, pathogen fitness, and host population depression by pathogens. Our results demonstrate the importance of trait heterogeneity and the insights available from relatively simple models, both for microparasite infection load and possibly other traits.

Keywords: Epidemiology, within-host dynamics, trait heterogeneity, parasite aggregation, host regulation

<u>Introduction</u>

Individual organisms differ from each other critically in traits such as body size, the number of parasites they host, and vital rates with implications for population dynamics including

evolution, ecological resilience, and our ability to make ecological inference. Quantitative

genetics models have long considered how the mean and variance (most often small and fixed

but not always) of heritable traits impact population dynamics [1,2]. Non-heritable sources of

trait heterogeneity, our focus here, may also be of critical importance. Individual-level

heterogeneity can interact with population dynamics for traits such as parasite load [3], body size

[4], or vital rates [5]. Further, heterogeneity interacts with evolution as heterogeneity may slow

evolution [6] or selection may favor alleles that alter variance in fitness, as classically expressed

in terms of geometric mean fitness [7]. Further, heterogeneity can interact with ecological

resilience, reflecting population declines or helping drive recoveries [4]. Additionally, trends in

heterogeneity may contain important information about unobserved biological processes [8]. For

these reasons, ecologists have frequently sought to understand and model individual

heterogeneity in populations.

But one barrier to understanding the role of heterogeneity is the complexity of jointly

modelling heterogeneity and population dynamics, particularly in such a way as to gain general

analytical insight. For example, Partial Differential Equations, Integral Projection Models,

Class-structured Matrix Models (for discrete traits) and Individual Based Models all provide

approaches to relatively easily simulate the interaction between individual heterogeneity and

population dynamics, but going beyond simulation-based results often requires biologically

unrealistic assumptions or advanced mathematical techniques. In general, the literature is lacking

somewhat for simpler models that i) allow for the joint modelling of individual heterogeneity and

population dynamics ii) are based on biologically and empirically supported assumptions and iii)

provide novel and generalizable conceptual insight.

Such models are particularly needed in epidemiology where individual heterogeneity

among hosts can have substantial feedbacks on host and parasite population dynamics. For

example, one major aspect of individual heterogeneity in host-parasite systems is that the

minority of the hosts hold the majority of parasites. Such variance (or aggregation) in the number

of parasites on each infected host (infection load) holds implications for host population

dynamics [3,9,10] and variance patterns in wild populations can indicate within-host

mechanisms [8,11]. These observations highlight the need for model-based, conceptual insight to

address our focal question. How does variance in microparasite load interact with population-

level dynamics? Classically, variance in infection load has been considered important for

macroparasites (e.g., arthropods or helminths) but often ignored for microparasites (e.g., fungi,

bacteria, viruses). In fact, microparasitic infections have often been treated as present or absent

[12]. This binary view reflects an assumption that variation in microparasite infection load is

negligible for one of two reasons; either, load variation is negligible because it is small due to

homogeneous hosts and rapid microparasite growth to a steady state or load variation need not be

small but lacks impact on key outcomes.

But many microparasite taxa display significant variance in infection load with important

impacts on population dynamics, evolution, and resilience ecology. For example, HIV patients

display wide variability in viral load, impacting HIV spread and likely also viral evolution [13].

Extreme variation in SARS-CoV-2 viral load may be critical for superspreading [14]. Variation

in bacterial pathogen load can help predict clinical outcomes for patients [15,16] and likely also

impacts pathogen evolution as bacterial genotypes vary in their loads [16]. The fungal agent of

White Nose Syndrome achieves very different pathogen loads in bats, both across hibernation

duration, within species at a given time, and between bat species with conservation implications

[17]. Frogs infected with the fungal agent of chytridiomycosis, *Batrachochytrium dendrobatidis*

(*Bd*), display widely varying infection loads within and between populations, impacting the

survival of individual frogs as well as the persistence of populations [18]. Because heterogeneity

in microparasite load is critical for many taxa, we need new theory for how this heterogeneity

interplays with population dynamics.

While existing theory for macroparasite load provides a critical foundation, we need

further development for microparasites. Microparasite load, as quantified by molecular

techniques (e.g., qPCR), can follow continuous distributions of infection load [14,19–21] rather

than the discrete distributions of macroparasite counts. Further, the availability of laboratory data

allows us to parameterize more biologically realistic, non-linear functions for the relationships

between infection load and key processes such as pathogen growth rate or the mortality of

infected hosts [22] contrasted with the linear functions typically used in classic macroparasite

theory. Seizing these opportunities, we develop novel theory for heterogeneity in microparasite

load focused on dynamic mean and variance in load. With a simple model (relative to models

with full, arbitrary trait distributions like Integral Projection Models), we address our focal

question of how microparasite load variance interacts with population dynamics. We show how

the mean and variance i) are affected by population dynamics throughout epidemic stages ii)

affect population dynamics, especially host suppression and iii) have mutual negative feedbacks

on each other leading to multiple possible qualitative patterns for load distributions. Importantly,

one implication of these cross-scale, bi-directional feedbacks is that the dynamic patterns of

mean and variance in infection intensity contain distinct, qualitative signatures of biological

mechanisms that are impossible to identify from data such as trends in abundance. We illustrate

this phenomenon with a case study of recovering populations of endangered Mountain Yellow

Legged Frogs depressed by *Bd*. Using our model, we demonstrate that the mean and standard

deviation of infection load provide enough information to infer the type of host defence strategy

that hosts are evolving to recover following disease-induced declines, differentiating tolerance,

constitutive resistance, inducible resistance, and acquired resistance. Altogether, we find that a

relatively simple model of heterogeneity gives us new insights into population dynamics,

evolution, and our ability to infer within-individual processes from population-level patterns.

Model structure

Trait heterogeneity within a population interacts with the population-level processes,

birth, death, infection, etc., that determine how many individuals are in the population. We

model these interactions with an Integral Projection Model (IPM) that tracks the population

density of susceptible hosts, infected hosts, and environmental pathogens ("zoospores" per our

focal case study of amphibian chytrid) at each time step (t, $t+1$, etc). We use one day time steps,

but this assumption is easily relaxed [23]. We assume that log load x is normally distributed in

the class of infected hosts, a reasonable assumption for *Bd* [24] and many other microparasites

[13,14,20,21]. This assumption allows us to derive a much more tractable and simpler model (see

Appendix for derivation of our reduced dimension IPM from the full IPM via moment closure),

in which we only need to track the mean and standard deviation of log load rather than the full

distribution. This simpler model allows us to analytically tease out how the mean and variance of

log load impact population dynamics and vice versa (Fig. 1). This is all but impossible to do with

the full IPM directly. Three key processes govern load dynamics. First, newly infected hosts tend

to have low loads. Second, pathogen growth increases loads. Third, infected hosts with higher

loads are more likely to die.

Figure 1. Conceptual diagram of disease model. Hosts begin life susceptible then transmission

moves them into the infected class, which is structured by the distribution of log pathogen load,

x, at a given time, t. New infections predominantly add hosts at low loads, bringing the mean

load down. Pathogen growth increases loads, increasing the mean. Loss of infection, partially

from host recovery but predominantly from host death, disproportionately occurs at high loads

and brings the mean down. Pathogens are shed back into the environment at a rate proportional

to load. $I(x, t)$ dx is the population density of hosts with a load between x and $x +$ dx at time t, and

$I(t)$ is the total population density of infected hosts at time t.

Further, we consider host evolution in this model, for our focal case study of recovering

frog populations, via changes in genotype frequencies. For simplicity, we assume clonal

competition among frog genotypes where those that are more defended in terms of their

epidemiological traits have lower fecundity (r_j); in particular, we assume this defence-fecundity

trade-off has a curvature that favors evolution toward intermediate values of defence in the

presence of pathogens. In each time step, susceptible hosts of genotype j are born when

susceptible or infected hosts $[S_j(t)+I_j(t)=N_j(t)]$ reproduce, with some negative density dependence

$(e^{-qN(t)}$ where $N(t)$ is all hosts and q is the strength of negative density dependence), and some

probability of mutation (m, Eq. 1a). Negative density dependence that regulates host populations

in the absence of disease.

Susceptible hosts may survive (probability s_0) and remain uninfected [depending on

transmission rate β_j and zoospore density $Z(t)$] or become infected while infected hosts that

survive may recover back to the susceptible class (with probability l, Eq. 1b and 1c). The

probability an infected host survives a time step declines with log load x in a non-linear fashion

[survival = $1-\square((x-\mu_{\text{LD50},j})/\sigma_{\text{LD50}})$ where $\square$ is the cumulative distribution function of the standard

normal, an empirically supported functional form [22]]; higher values of σ_{LD50} model a more

gradual, sigmoidal decline of survival probability from 1 to 0 and $\mu_{\text{LD50},j}$ gives the LD50 of load.

The average survival probability of infected hosts of genotype j always declines with their mean

load, $\mu_j(t)$ [average survival: $\hat{s}(\alpha_j(t))$ where $\hat{s}(z) = 1- \square(z)$ and $\alpha_j(t) = (\mu_j(t)-$

$\mu_{\text{LD50},j})/(\sigma_j^2(t)+\sigma_{LD50}^2)^{1/2}$ is a quantity that arises multiple times]. Higher standard deviation of

load, $\sigma_j(t)$, leads to higher death rates, as long as the minority of infected hosts are dying each

time step, because a higher standard deviation increases the minority of hosts with high and

deadly loads $[\partial\hat{s}(\alpha_j(t))/\partial\sigma_j(t)<0$ if and only if $\mu_j(t) < \mu_{\text{LD50}}$, i.e., $\hat{s}(\alpha_j(t)) > 0.5]$.

The dynamics of the mean and variance are captured in the dynamics of the first and

second moment of log load. Mean log load is related to the first moment of log load $[\mu_j(t) =$

$P_j(t)/I_j(t)$; $P_j(t) = \int xI_j(x,t)\mathrm{d}x$ and $I_j(t) = \int I_j(x, t)\ \mathrm{d}x$ over all x where $I_j(x,t)\mathrm{d}x$ is the population

density of hosts of genotype j with a log infection load between x and $x + \mathrm{d}x$]; the first moment

captures all sources that contribute to pathogen load (Eq. 1d) as newly infected hosts have low

log loads (drawn from a normal distribution with mean μ_0 and standard deviation σ_0) and

pathogens grow within hosts with some variability and negative density dependence [$x(t+1)$ is

drawn from a normal distribution with mean $a+bx(t)$ and standard deviation σ_G; $0<b<1$ captures

negative density dependence]; this functional form of microparasite growth is empirically

supported for our focal system [22] and such within-host growth on a log scale up to a carrying

capacity may be appropriate for many host-microparasite systems. The second moment of log

load is related to the standard deviation of log load [$\sigma_j(t) = (Q_j(t)/I_j(t) - \mu_j^2(t))^{1/2}$ where $Q_j(t) =$

$\int x^2 I_j(x,t)dx$ over all x]; this second moment captures all sources that contribute to log load and

variation in log load such as variation in the loads of newly infected hosts and variation in

pathogen growth (e.g., unlike Eq. 1d, σ_0 and σ_G appear in Eq. 1e). Key to both the first and

second moments of log load (and thus mean and standard deviation) are the three processes

controlling pathogen load: new infections, pathogen growth, and death (see Fig. 1). Both the

mean and standard deviation of log load contribute to the mean of load on the natural scale

($e^{\mu+\sigma^{\wedge}2/2}$ is mean load on the natural scale) and thus pathogen shedding from infected hosts (with

λ giving the zoospores shed per within-host pathogen per time unit) and pathogens persist for

some time in the environment (with probability v per time step, Eq. 1f).

<u>Eq. 1a: Birthrate of genotype j</u>

$$f_j\big(N_{1:n}(t)\big) = [\underbrace{r_j N_j(t)(1-m)}_{\text{Birth less mutation}} + \underbrace{\frac{m}{n-1}\sum_{k=1,\ k\neq j}^{n} r_k N_k(t)}_{\text{Gains from mutation}}]\underbrace{e^{-qN(t)}}_{\text{Crowding}}$$

193   <u>Eq. 1b for susceptible individuals of genotype j</u>

$$S_j(t+1) = \underbrace{f_j\big(N_{1:n}(t)\big)}_{\text{Birth}} + \underbrace{S_j(t)s_0 e^{-\beta_j Z(t)}}_{S_j \text{ survive, remain uninfected}} + \underbrace{I_j(t)\hat{s}(\alpha_j(t))l}_{I_j \text{ survive, recover}}$$

194   Eq. 1c for infected individuals of genotype j

$$I_j(t+1) = \underbrace{S_j(t)s_0[1 - e^{-\beta_j Z(t)}]}_{S_j \text{ survive, get infected}} + \underbrace{I_j(t)\hat{s}(\alpha_j(t))(1-l)}_{I_j \text{ surive, do not recover}}$$

195   Eq. 1d for first moment of log load on infected individuals of genotype j ("0[th]" etc. refer to terms

196   of pathogen growth of different order with respect to pathogen density dependence, b_j)

$$P_j(t+1) = \underbrace{S_j(t)s_0[1 - e^{-\beta_j Z(t)}]\mu_0}_{S_j \text{ survive, get infected}} +$$

$$\underbrace{I_j(t)\hat{s}\left(\alpha_j(t)\right)(1-l)}_{I_j \text{ survive, do not recover}} \left\{ \underbrace{a_j}_{0th} + \underbrace{b_j\left[\mu_j(t) - \frac{\sigma_j^2(t)\varphi\left(\alpha_j(t)\right)}{\hat{s}\left(\alpha_j(t)\right)\sqrt{\sigma_j^2(t) + \sigma_{LD50}^2}}\right]}_{1st \ modified \ by \ load-dependent \ survival} \right\}$$

198   Eq. 1e for second moment of log load on infected individuals of genotype j

$$Q_j(t+1)$$

$$= \underbrace{S_j(t)s_0[1 - e^{-\beta_j Z(t)}](\sigma_0^2 + \mu_0^2)}_{S_j \text{ survive, get infected}}$$

$$+ \underbrace{I_j(t)\hat{s}\left(\alpha_j(t)\right)(1-l)}_{I_j \text{ survive, do not recover}} \left\{ \underbrace{\sigma_G^2 + a_j^2}_{0th} + \underbrace{2a_jb_j\left[\mu_j(t) - \frac{\sigma_j^2(t)\varphi\left(\alpha_j(t)\right)}{\hat{s}\left(\alpha_j(t)\right)\sqrt{\sigma_j^2(t) + \sigma_{LD50}^2}}\right]}_{1st \ modified \ by \ load-dependent \ survival} \right.$$

$$\left. + \underbrace{\left[b_j^2(\sigma_j^2(t) + \mu_j^2(t)) - \frac{\sigma_j^2(t)\varphi\left(\alpha_j(t)\right)\left(2\mu_j(t)\sigma_{LD50}^2 + \left(\mu_{LD50,j} + \mu_j(t)\right)\sigma_j^2(t)\right)}{\hat{s}\left(\alpha_j(t)\right)\left(\sigma_j^2(t) + \sigma_{LD50}^2\right)^{\frac{3}{2}}}\right]}_{2nd \ modified \ by \ load-dependent \ survival} \right\}$$

199   Eq. 1f: Environmental pathogens

$$Z(t + 1) = \lambda \underbrace{\sum_{j=1}^{n} I_j(t) e^{\mu_j(t) + \frac{\sigma_j^2(t)}{2}}}_{\text{Pathogen shedding}} + \underbrace{vZ(t)}_{\text{Environmental persistence}}$$

Note that $\varphi(z)$ is the probability density function of the standard normal distribution at z

$[\varphi(z)=(1/\sqrt{(2\pi)}) \, e^{-(z^2)/2}]$. We also derive a very similar version of the model in which recovered

hosts enter a recovered class with acquired immunity (until immunity wanes) such that they have

different traits upon reinfection and the load of reinfected hosts follows its own lognormal

distribution (see Appendix).

<u>Load dynamics from epidemic outbreak to endemic equilibrium</u>

With these equations for the dynamics of the host population, pathogen population, and

the distribution of load, we can gain insight into how population dynamics impact load. We

simulated a small density of hosts who just became infected in a host population of one genotype

at its disease-free equilibrium (using biologically reasonable parameters for Mountain Yellow

Legged Frogs and *Bd*; see Appendix for values and simulation procedure). At different stages of

the epidemic (beginning, log growth, peak, and endemic equilibrium), we see different

distributions of load reflecting how the population-level processes impact the distribution of load

(beginning, log growth, and peak stages are marked by coloured squares in Fig. 2A).

At the beginning of the epidemic, there are very few infected hosts and those who are

infected have only just become infected [i.e., $\mu(t = 0)=\mu_0$ and $\sigma(t = 0) = \sigma_0$], causing low mean

and standard deviation of load (Day 0 in Fig. 2B). Pathogen growth during the log growth of

prevalence stage (Day 49 in Fig. 2B) quickly raises the mean load and there is now separation in

the infected population between newly infected hosts and hosts who have been infected long

enough to experience pathogen growth (Fig. 2C shows how pathogen growth raises the mean

while new infections bring it down) so there is now a wider spread of loads [i.e., $\sigma(t = 49) > \sigma(t =$

0), compare blue and yellow bell curves in Fig. 2D]. As the rate of increase of prevalence slows

and even becomes slightly negative (from around day 200 to orange square in Fig. 2B), mean log

load rises and the standard deviation declines (compare orange curve for peak stage to yellow log

growth curve in Fig. 2D). Mean load rises again at this stage because there are far fewer new,

low load infections entering the infected class and dragging the mean down (see sharp decrease

in New curve in Fig. 2C). The higher mean pushes the right tail of the load distribution up

against the survival constraint (Fig. 2D, orange bell curve's right tail extends far into the

declining portion of the pink survival curve); loss of these high load infections shrinks the

standard deviation of load down [see how $\sigma(t)$ has declined at day 342 in Fig. 2B]. Loads have

become so high that the survival constraint forces mean loads back down to approach their

endemic equilibrium value and the standard deviation is allowed to recover to somewhat higher

values [see rising Loss curve in Fig. 2C around day 342 as well as decreasing $\mu(t)$ and increasing

$\sigma(t)$ in Fig. 2B from day 342 onward]. These general patterns for $\mu(t)$ and $\sigma(t)$ hold across a broad

range of parameters (see code), capturing the impacts of epidemic stage on load.

Figure 2. Epidemic stage shifts the distribution of pathogen load. (A) As a rare pathogen

outbreaks, host density (black curve) declines and pathogen density (gold curve) rises. Coloured

squares denote epidemic stages of beginning (blue), log growth (yellow), and peak (orange). (B)

Prevalence (black) rises over time until it reaches a peak then settles slightly lower toward

equilibrium. Mean (solid gold) and standard deviation (dashed gold) of log load change during

these epidemic stages as well. (C) The change in the mean is determined by a positive

contribution from pathogen growth (black) and negative contributions from new infections (solid

gold) and infection loss (dashed gold), which is predominantly due to high load infections being

lost at a higher rate. (D) The bell curve of log load shifts higher or lower and broader or narrower

246 at different epidemic stages (days 0, 49, and 342 shown). Survival declines with log load,

247 particularly at high loads, constraining the distribution of load (dashed pink).

249    The impact of epidemic stage on the distribution of load also prompts investigation of the

250 long-term, endemic equilibrium mean and standard deviation. We found implicit expressions for

251 the equilibrium mean (μ^*_j) and standard deviation (σ^*_j) of log load for a genotype j (Eq. 2a, b):

252 Eq. 2a: Equilibrium mean of log load

$$\mu^*_j = \underbrace{[1 - \hat{s}(\alpha^*_j)(1 - l)]}_{\text{Prob. inf is lost}} \underbrace{\mu_0}_{\text{Initial}} +$$

$$\underbrace{\hat{s}(\alpha^*_j)(1 - l)}_{\text{Prob. inf remains}} \left\{ a_j + b_j \underbrace{\left[\mu^*_j - \frac{\sigma^{*2}_j \varphi(\alpha^*_j)}{\hat{s}(\alpha^*_j)\sqrt{\sigma^{*2}_j + \sigma^2_{LD50}}} \right]}_{\text{Contribution of within-host growth}} \right\} = F(\mu^*_j, \sigma^*_j, y_1, y_2 \dots y_n)$$

253 Eq. 2b: Equilibrium variance plus mean squared of log load

$$\sigma^{*2}_j + \mu^{*2}_j = \underbrace{[1 - \hat{s}(\alpha^*_j)(1 - l)]}_{\text{Prob. inf is lost}} \underbrace{(\sigma^2_0 + \mu^2_0)}_{\text{Initial}}$$

$$+ \underbrace{\hat{s}(\alpha^*_j)(1 - l)}_{\text{Prob. inf remains}} \left\{ \sigma^2_G + a^2_j + 2a_j b_j \underbrace{\left[\mu^*_j - \frac{\sigma^{*2}_j \varphi(\alpha^*_j)}{\hat{s}(\alpha^*_j)\sqrt{\sigma^{*2}_j + \sigma^2_{LD50}}} \right]}_{\text{Contribution of within-host growth}} \right.$$

$$\left. + \underbrace{\left[b^2_j (\sigma^{*2}_j + \mu^{*2}_j) - \frac{\sigma^{*2}_j \varphi(\alpha^*_j)\left(2\mu^*_j \sigma^2_{LD50} + (\mu_{LD50,j} + \mu^*_j)\sigma^{*2}_j\right)}{\hat{s}(\alpha^*_j)(\sigma^{*2}_j + \sigma^2_{LD50})^{\frac{3}{2}}} \right]}_{\text{Contribution of within-host growth continued}} \right\}$$

$$= G(\mu^*_j, \sigma^*_j, y_1, y_2 \dots y_n)$$

254 Eq. 2c: Pathways affecting equilibrium mean log load

$$\frac{d\mu_j^*}{dy_k} = 1/[\underbrace{\left(1 - \frac{\partial F}{\partial \mu_j^*}\right)\left(1 - \frac{\partial K}{\partial \sigma_j^*}\right) - \frac{\partial F}{\partial \sigma_j^*}\frac{\partial K}{\partial \mu_j^*}}_{\text{Feedback scaling}}] [\underbrace{\left(1 - \frac{\partial K}{\partial \sigma_j^*}\right)}_{\text{Feedback scaling}} \underbrace{\frac{\partial F}{\partial y_k}}_{\text{Direct}} + \underbrace{\frac{\partial F}{\partial \sigma_j^*}\frac{\partial K}{\partial y_k}}_{\text{Indirect}}]$$

255   Eq. 2d: Pathways affecting equilibrium SD of log load

$$\frac{d\sigma_j^*}{dy_k} = 1/[\underbrace{\left(1 - \frac{\partial F}{\partial \mu_j^*}\right)\left(1 - \frac{\partial K}{\partial \sigma_j^*}\right) - \frac{\partial F}{\partial \sigma_j^*}\frac{\partial K}{\partial \mu_j^*}}_{\text{Feedback scaling}}] [\underbrace{\left(1 - \frac{\partial F}{\partial \mu_j^*}\right)}_{\text{Feedback scaling}} \underbrace{\frac{\partial K}{\partial y_k}}_{\text{Direct}} + \underbrace{\frac{\partial K}{\partial \mu_j^*}\frac{\partial F}{\partial y_k}}_{\text{Indirect}}]$$

where K is the positive square root of $G-\mu_j^{*2}$. The expressions for the mean and standard

deviation of log load reflect the balance between the influences of new infection, pathogen

growth on the host, and infection loss (particularly through death because it is load-dependent).

The mean is the sum of initial mean weighted by the probability of infection loss plus the

contribution of within-host growth (see terms in Eq. 2a that mirror terms in Eq. 1d) weighted by

the probability that infection remains (i.e., an infected individual survives and does not recover).

The expression for the equilibrium value of the variance plus mean squared follows the same

structure (Eq. 2b). If we assume no loss of infection, we find explicitly that $\mu_j^* = a/(1-b)$ and σ_j^*

$= \sigma_G/\sqrt{(1-b)}$. Because there is loss of infection, we retain the somewhat more complex expression

in Eqs. 2a and 2b.

What factors influence the mean and standard deviation at equilibrium? We analyse these

expressions to find how any parameter (y_k with k in 1 to n) affects μ_j^* and σ_j^*, considered as

functions of that fixed parameter. As long as the terms we label "feedback scaling" remain

positive (Eqs. 2c, d; they do remain positive across a broad parameter range, see "Quantifying

numerical robustness" in the Appendix), we can interpret these expressions as how y_k affects μ_j^*

directly ("direct" term in Eq. 2c) and indirectly by affecting σ_j^* ("indirect" term in Eq. 2c). The

direct term, $\partial F/\partial y_k$, captures how a change in the parameter y_k directly changes the F function

(where $F = \mu_j^*$). The indirect term comes from how y_k directly alters σ_j^* (i.e., $\partial K/\partial y_k$) multiplied

by how a change in σ^*_j feeds back to change μ^*_j $(\partial F/\partial \sigma^*_j)$. The same logic applies for direct and

indirect effects on σ^*_j in Eq. 2d.

We find that pathogen growth and host death at high loads creates negative feedbacks

between μ^*_j and σ^*_j (i.e., indirect effects, $\partial F/\partial \sigma^*_j$, $\partial K/\partial \mu^*_j$, are negative). These negative

feedbacks hold as long as certain inequalities, such as $\mu^*_j > \mu_0$, $\sigma^{*2}_j + \mu^{*2}_j > \sigma_0^2 + \mu_0^2$, and $\mu^*_j <$

$\mu_{LD50,j}$ are met (see Appendix for one more inequalities and proofs; there is a fourth inequality

that is sufficient but not necessary and less interpretable). Most of these assumptions are

biologically reasonable, interpretable, and likely to hold for many systems. If $\mu^*_j > \mu_0$, then the

loss of infections will bring the mean down as the lost infections are replaced with infections that

have lower loads, on average. Similarly, if $\sigma^{*2}_j + \mu^{*2}_j > \sigma_0^2 + \mu_0^2$, then the loss of infections will

bring the variance plus mean squared down. If $\mu^*_j < \mu_{LD50,j}$ then fewer than 50% of infected

hosts die each time step and increased variance increases death rate instead of decreasing it (see

comment above about the effect of load variance on average survival). These negative feedbacks

held across a broad range of parameter sets we numerically searched to compare to our results

for our focal parameter set (see Quantifying Numerical Robustness section in the Appendix) and

only failed to hold in parameter sets where $\sigma^{*2}_j + \mu^{*2}_j < \sigma_0^2 + \mu_0^2$, $\mu^*_j < \mu_0$, or both (3.7% of

parameter sets considered; these parameter sets seem unlikely to be broadly, biologically

relevant). In summary, an increase in μ^*_j increases the minority of infected hosts with high,

deadly loads and their death squeezes the distribution of load, reducing σ^*_j. An increase in σ^*_j

also increases the minority of infected hosts with high, deadly loads and their death pulls μ^*_j

down. These mutual negative feedbacks arise because of growth of infection and the fact that the

minority of hosts at equilibrium have lethal loads.

Understanding these mutual negative feedbacks allows us to understand how a

perturbation shifts the distribution of load directly vs. indirectly by shifting these feedbacks.

Consider only the *direct effects* of a perturbation that directly increases μ^*_j, shifting the

distribution of load higher and having no direct impact on σ^*_j; biologically, this could be

envisioned as a perturbation that changes all natural loads by the same percentage (because a

fixed, multiplicative change in natural load is a fixed, additive change in log load), e.g., brief,

favorable pathogen growth conditions that suddenly boost all pathogen loads by the same

proportion. Every percentile of the load distribution is now found at higher load values,

increasing disease-induced death rates for higher percentiles in particular. Then we consider the

response in loads as enough time passes for the *indirect effects* of the perturbation to manifest.

These higher percentiles of load experience higher death rates, shrinking the right tail of the

distribution of load, *indirectly* decreasing σ^*_j (see how the orange bell curve in Fig. 2D is

narrower than the yellow bell curve). Similarly, a perturbation that only *directly* increases σ^*_j,

say an increase in environmental stochasticity that increases stochasticity in pathogen growth

(σ_G), moves the left tail of load lower, with negligible impact on survival, and the right tail of

load higher against the survival constraint, *indirectly* decreasing μ^*_j. Thus, a parameter that shifts

both μ^*_j and σ^*_j also acts via their feedbacks; e.g., increased recovery rate (*l*) directly decreases

μ^*_j and σ^*_j because both are constrained to be closer to initial infection levels; but lowering μ^*_j

strongly reduces the survival-based constraints on σ^*_j, creating a net increase in σ^*_j; this example

helps us understand how various parameters affect μ^*_j and σ^*_j directly and via their mutual

negative feedbacks (Table 1).

These direct and indirect effects give us broadly applicable, conceptual insight into how

various processes will impact the distribution of pathogen load. The signs of direct effects of

these various parameters on μ^*_j and σ^*_j can be analytically guaranteed given biologically

meaningful, sufficient conditions (see conditions under Table 1 and proofs in Appendix). Many

of these conditions depend on the equilibrium, average probability of infected hosts surviving for

one day, $\hat{s}(\alpha^*_j)$, not being too low; the most restrictive of these sufficient conditions is $\hat{s}(\alpha^*_j) >$

0.958, which may be biologically reasonable given that we find $\hat{s}(\alpha^*_j) = 0.984$ for our focal

parameter values (see Table A1 for biologically reasonable parameter values and sources) and

even higher for genotypes that evolve defence (also see Quantifying Numerical Robustness for

the percentage of parameter sets in which these conditions held). Thus, we believe the signs of

the direct effects that we found will hold in many biologically meaningful scenarios, giving us

broad, conceptual insight into load dynamics. Additionally, these findings help us see how

various processes would impact load, e.g., lower a_j due to higher constitutive resistance should

strongly decrease mean and have a weak impact on the standard deviation; a population with a

higher variance in initial load (σ_0), e.g., due to some greater host variability, should have a higher

standard deviation of load due to the direct effects of that heterogeneity but lower mean due to

feedback and indirect effects. Further, feedbacks, direct effects, and indirect effects may prove

insightful for the equilibrium distributions of traits other than load.

Table 1. Analytical results for direct and numerical results for total effects of parameters

on μ^*_j and σ^*_j. Superscripts in columns 2 and 3 denote conditions analytically found to be

sufficient (though not always necessary) for that sign to hold (see Appendix for derivations); the

feedback scaling terms are always positive in a broad, biologically relevant parameter range.

Columns 2 and 3 analytically determine the signs of the direct effects of parameter y_k on μ^*_j and

σ^*_j respectively. Columns 4 ($d\mu^*_j/dy_k$) and 5 ($d\sigma^*_j/dy_k$) quantify the total effects (direct

effects+indirect effects, each multiplied by relevant feedback scaling from Eq. 2) found for the

wild-type genotype in our focal parameter set. Mutual negative feedbacks between μ^*_j and σ^*_j

guarantee that indirect effect of any y_k on μ^*_j always has the opposite sign as y_k's direct effect on

σ^*_j, and vice versa, due to mutual negative feedbacks. Parameters not listed do not affect μ^*_j and

σ^*_j (e.g., note that β_j does not appear in Eqs. 2a, b).

Changing parameter y_k	Direct effect on mean: $\partial F/\partial y_k$	Direct effect on SD: $\partial K/\partial y_k$	Total effect on mean $d\mu^*_j/dy_k$	Total effect on SD $d\sigma^*_j/dy_k$
SD of load upon initial infection: σ_0	0	≥ 0	-0.073	0.078
Mean load upon initial infection: μ_0	≥ 0	$\geq 0^1$	0.20	-0.094
a_j contributes to pathogen growth rate at all loads	≥ 0	$\geq 0^{2,\,3}$	5.1	0.80
b_j contributes to pathogen growth rate at high loads	$\geq 0^{2,\,3}$	$\geq 0^{2,\,3,\,4,\,5}$	26	16
σ_{LD50} makes the survival function more gradual	$\leq 0^{2,\,4,\,6}$	$\leq 0^{2,\,5,\,6,\,7}$	-0.64	0.020
$\mu_{LD50,j}$ is the LD50 of the survival function	$\geq 0^{2,\,6}$	$\geq 0^{2,\,4,\,8}$	0.42	0.042

σ_G controls stochasticity in pathogen growth	0	≥ 0	-1.5	1.6
Recovery rate from infection l	≤ 0[6]	≤ 0[8]	-28	6.8

[1] $\mu_0 \geq 0$, i.e., newly infected hosts have non-negative mean log load.

[2] $\mu^*_j < \mu_{\mathrm{LD50},j}$, i.e., infected host survival probability for one day is $> 50\%$.

[3] $\mu^*_j > 0.8\sigma^*_j$, i.e., SD is not too much larger than mean.

[4] $\alpha^*_j < -1$, i.e., infected host survival probability for one day is $> 84.1\%$.

[5] $\mu^*_j > 0.177\mu_{\mathrm{LD50},j}$, i.e., mean is not too much lower than LD50.

[6] $\mu^*_j > \mu_0$, i.e., infection grows so that mean log load is greater than the mean of new infections

[7] $\alpha^*_j < -(3^{1/2})$, i.e., satisfied if infected host survival probability for one day is $> 95.8\%$

[8] $\mu^{*2}_j + \sigma^{*2}_j > \mu^2_0 + \sigma^2_0$. Guaranteed if condition 6 is met and $\sigma^*_j > \sigma_0 + 40$

<u>Load signals in evolutionary recovery</u>

The mutual negative feedbacks between the mean and standard deviation of load greatly

aid our ability to infer differences in underlying mechanisms via differences in load. Due to their

mutual negative feedbacks, the mean and standard deviation of load can shift in opposite

directions in response to a change in a single mechanism (the direct effect columns of Table 1

never have opposite sign but the net effect columns do). This greatly improves our ability to infer

underlying processes from the distribution of load as the mean and standard deviation can each

have three qualitative responses (+, flat, -) and nine theoretical combinations of responses. We

demonstrate the utility of these load signatures with a focal case study of Mountain Yellow

Legged Frog populations recovering from *Bd*.

Population abundance and genetic data indicate Mountain Yellow Legged frog

populations are recovering in the presence of a deadly fungal pathogen (*Bd*) due to some form of

evolution relevant to host defences [25]. However, the precise process of host evolution is

unknown. Frogs may evolve tolerance that improves their survival of high load infections (higher

$\mu_{\mathrm{LD50,\,j}}$ in our model), resistance that prevents infection (lower β_{j}), or resistance that decreases

pathogen growth. Resistance to pathogen growth may be constitutive and apply equally to all

infections (lower a_{j}), be inducible to disproportionately resist growth of high load infections

(lower b_{j}), or be acquired immunity that only protects reinfected frogs (lower a_{j} in reinfected

frogs; see Appendix for the extension of Eq. 1 to the case of acquired immunity). These different

processes correspond to different timing and likely molecular mechanisms of host immunity

evolution. We simulated outbreak of fungal disease in a population of mostly wild-type frogs

with a low starting frequency of nine other genotypes spanning a range of defence in one of these

traits with a fecundity cost of this higher defence (see Appendix). The outbreak of the pathogen

drives host density down (Fig. 3A) as prevalence increases (Fig. 3B) but the more defended

genotypes then increase in frequency (Fig. 3C) and alter the distribution of load (Fig. 3D and E)

with implications for pathogen abundance (Fig. 3F).

Figure 3. Evolutionary recovery via constitutive resistance. Host genotypes vary in the a_j

parameter and more defended genotypes (lower a_j) have lower fecundity. (A) Pathogen invasion

causes host density to decline then recover as hosts evolve. (B) Invasion increases prevalence but

then evolution of constitutive resistance decreases prevalence. (C) Host evolution occurs as the

frequencies of clonal genotypes shift from the least defended, wild-type genotype to more

defended genotypes. (D) More defended genotypes have lower mean log loads (coloured curves)

and the overall mean log load of the entire infected population declines as hosts evolve resistance

(black curve with gray arrow emphasizing change from wild-type stable equilibrium until the

end of the simulation). (E) The standard deviation of load has a weaker, more mixed dependence

on constitutive resistance but weakly increases overall as hosts evolve. Note a transient spike in

the overall standard deviation around year 5 due to a temporary mixture of more and less well

defended genotypes. (F) Host evolution decreases pathogen density in the environment.

Critically, we find that these different defence hypotheses correspond to qualitatively

different patterns in pathogen load. As more defended genotypes increase in frequency, the

overall load pattern in the entire infected host population generally follows that predicted by the

equilibrium analysis for a single genotype in Table 1 (though Table 1, alone, is insufficient to

predict the reinfection case). Increasing tolerance allows a higher and wider spread of loads,

leading to a higher mean and weakly higher standard deviation (see Fig. A1 in Appendix).

Resistance that prevents infection has no direct impact on the mean and standard deviation but

changes the timing of epidemic stage, leading to weak and transient changes in the mean and

standard deviation (see Fig. A3). Different forms of resistance to pathogen growth (depicted

graphically in Fig. 4A) also correspond to qualitatively different trends. Constitutive resistance

decreases the mean and weakly increases the standard deviation (Fig. 4B); this result arises

because constitutive resistance has a weak, negative, direct effect on the standard deviation but

the strong decrease in mean and negative feedbacks of the mean on the standard deviation leads

to a net positive effect on the standard deviation (see the row for a in Table 1). Inducible

resistance has a strong, direct, negative impact on both the mean and standard deviation (Fig.

4C). Acquired resistance to pathogen growth in reinfections drives lower mean and strongly

higher standard deviation (Fig. 4D); this increase arises because the infected population is now a

mixture of hosts infected for the first time, with relatively high loads, and reinfected hosts, with

lower loads (see Appendix for explanation of when and why this increases the standard

deviation). This increases the variance in log load in the total infected host population. These

results were highly robust to variation in the model parameters and arose in the model without

the simplifying assumption of lognormal load (see Quantifying Numerical Robustness section in

the Appendix), though the model without the simplifying assumption provides less conceptual

clarity as to the reason for these results than our focal model here does. Thus, the distribution of

load can qualitatively differentiate among hypotheses for the defence evolution of Mountain

Yellow Legged Frogs, or other hosts.

Figure 4. Load trends qualitatively differ for different forms of resistance. (A) We show

deterministic ln(load+1) on a single host witsh all three types of resistance tracked over 100 days

with infection on day 16, recovery on day 51, and reinfection on day 66. Constitutive resistance

is always on. Inducible resistance acts most strongly at high loads. Acquired resistance only

applies to reinfections. (B) Increasing constitutive resistance decreases mean log load (gold)

while the standard deviation increases weakly (blue). Black arrows emphasize the overall trends.

(C) Inducible resistance decreases the mean and strongly decreases the standard deviation. (D)

Acquired resistance decreases the mean and increases the standard deviation because the entire

infected population is now composed of first-time infected hosts with high mean loads and

reinfections with lower mean loads. These simulations were initialized at the wild-types endemic

equilibrium with low frequencies of the nine, more defended genotypes (see Appendix).

Impacts of the load distribution on population dynamics

Figure 5. Higher standard deviation contributes to host death, pathogen spread, and host

population depression. We show results for single-genotype simulations for a genotype that

either possesses strong constitutive ($a = 0.35$; all other parameters as wild-type) or inducible ($b =$

0.88) resistance. For comparability, we parameterized these to give the same, average infected

host fitness at equilibrium. (A) Constitutive resistance (blue) leads to a lower mean and higher

standard deviation of log load compared to inducible resistance (orange). Constitutive resistance

then has a slightly higher proportion of hosts with very high loads (see magnified inset panel).

(B) More high load infections lead to stronger host population depression for constitutive

resistance than inducible (solid blue lower than solid orange) due to greater pathogen spread

(dashed blue higher than dashed orange).

We exemplify the impact of the distribution of load on population dynamics by

contrasting how different forms of resistance affect host and pathogen fitness as well as endemic

conditions. Consider the dynamics of a single genotype with strong constitutive resistance

experiencing an epidemic in isolation contrasted to those of a single genotype with strong

inducible resistance. Both genotypes keep mean loads low but constitutive resistance has a

higher standard deviation of log load at equilibrium ($\sigma^*_{\text{Constitutive}} > \sigma^*_{\text{Inducible}}$); because mortality

falls disproportionately on high load infections (assuming $\mu^*_{\text{Constitutive}} < \mu_{\text{LD50}}$ and $\mu^*_{\text{Inducible}} <$

μ_{LD50}, i.e., fewer than 50% of infected hosts die each day), the higher standard deviation of

constitutive resistance contributes to higher death rates. For constitutive and inducible resistance

to achieve the same, average fitness while infected [$\hat{s}(\alpha^*_{\text{Constitutive}}) = \hat{s}(\alpha^*_{\text{Inducible}})$], the genotype

with constitutive resistance must reduce mean log load further [we set $a = 0.35$ for the

constitutive genotype and $b = 0.88$ for inducible with all other parameters as wild-type; see Fig.

5A; $\mu^*_{\text{Constitutive}} < \mu^*_{\text{Inducible}}$ is required by $\sigma^*_{\text{Constitutive}} > \sigma^*_{\text{Constitutive}}$, $\mu^*_{\text{Inducible}} < \mu_{\text{LD50}}$, and $\mu^*_{\text{Inducible}}$

$< \mu_{\text{LD50}}$]. Despite lower mean loads, the higher standard deviation of the constitutively resistant

genotype means there are more individuals with rare, very high loads that lead to higher average

pathogen shedding rates per infected hosts [$\mu^*_{\text{Constitutive}} + \sigma^*_{\text{Constitutive}}{}^2/2 > \mu^*_{\text{Inducible}} + \sigma^*_{\text{Inducible}}{}^2/2$; see

Appendix for biologically reasonable conditions under which this inequality will hold]. Higher

pathogen shedding rates contribute directly to pathogen fitness (see R_0 expression in [23]) and

host population depression in the case of constitutive resistance compared to inducible resistance

(Fig. 5B). Therefore, different forms of host defence and thus standard deviations of load can

have strongly different implications for pathogen fitness, even given equal host fitness while

infected (acquired resistance falls on this same continuum, see Fig. A6).

<u>Discussion</u>

How does variance in microparasite load interact with population-level dynamics? We

show how epidemic stage alters the mean and standard deviation of load due to the rate of influx

of new infections, pathogen growth, and high mortality at high loads. Further, we show that the

mean and standard deviation have negative feedbacks with each other due to high mortality at

high loads. These negative feedbacks create qualitatively different trends in the mean and

standard deviation for different forms of host defence evolution, improving our ability to infer

within-host processes from population-level patterns. Finally, we show how different trends in

the variance in load, e.g., due to different forms of host defence evolution, have different impacts

on host fitness, pathogen fitness, and host suppression by pathogens. Thus, we have gained new,

conceptual insights into the interactions among variance in microparasite load and population-

level dynamics that improves our understanding of the focal frog-*Bd* system and likely many

other microparasite systems.

Many of our key results will apply to microparasite load broadly. While other

microparasites may differ structurally in their between-host processes (e.g., frequency-dependent

transmission instead of environmental transmission), many of our key findings depend primarily

on assumptions regarding within-host processes that should be more general. Microparasite

infections grow beyond their initial load, negative density-dependence within hosts constrains

the loads that can be attained, survival probability declines in some non-linear fashion from one

to zero as loads become very high, and the minority of infected hosts dies in a given time step.

Thus, many of our key, conceptual findings should apply to other microparasites despite

differences in parameter values or details of model structure. For example, our finding that mean

loads rise as an epidemic slows, due to a reduced influx of new infections bringing the mean

down, should apply to many microparasites. Negative feedbacks between the mean and standard

deviation, due to a non-linear survival constraint, should also arise frequently as they depend on

conceptually reasonable conditions, such as mean loads being higher than initial loads and less

than half of infected hosts dying every day; these negative feedbacks should allow different

forms of host defence to generate qualitatively different impacts on mean and standard deviation

in other microparasite systems as well. Further, our specific model and findings will be

particularly, directly applicable to microparasites that display lognormal distributions of load like

HIV and SARS-CoV-2 [13,14,20,21]. Because, any of our key results were supported

analytically and those supported numerically had broad numerical support, we conclude that our

key results should apply broadly to many other microparasite systems.

We were able to draw broader conclusions because our model was relatively simple and

tractable for analytical work, showing broad conditions in which our patterns would hold. The

relative simplicity of our model relies on a lognormal distribution of pathogen load, which is

empirically supported for *Bd* [24] and likely many other microparasites [13,14,20,21]. Second,

our model provides insights that are not readily available in simpler models with dynamic mean

and fixed variance [23] or in more complex models that make no simplifying assumptions about

the distribution of load [22]. Without our simplifying assumption of lognormal load, it would

have been much more difficult to derive expressions for the mean and variance of load at

equilibrium and clarify the existence of and reason for various key results. For mutual negative

feedbacks, for example, our simplifications allowed us to find that the decline in host survival at

high loads leads to negative feedbacks between the mean and the standard deviation. Thus, any

change (e.g., a change in a host trait) that directly affects mean and standard deviation also

indirectly affects them by shifting their feedbacks (Table 1). Disentangling these direct and

indirect pathways clarifies how the distribution of log load changes given a parameter

perturbation, e.g., constitutive resistance (lower a) directly lowers the standard deviation but can

actually cause it to weakly increase due to the sum of direct and indirect effects (Figs. 3 and 4).

A notable result from our modelling approach is that different forms of resistance have

qualitatively different impacts on the variance of load (Fig. 4), greatly improving our ability to

infer within-host processes from population-level patterns as well as predict population-level

impacts of epidemics (Fig. 5).

Improved understanding of load dynamics, in general, also helps us recognize how the

stochastic processes governing the standard deviation of load can become a battleground for

evolution by hosts and/or microparasites. Our model confirms the predictions from classic

compartmental models that tolerance (higher μ_{LD50} in our model) increases prevalence and

environmental pathogens (Råberg et al. 2008), confirming the clear intuition that tolerance

should increase mean log loads but illustrates with greater novelty that tolerance also increases

the standard deviation of log load. Resistance mechanisms, on the other hand, should decrease

pathogen fitness [26], as we found. We were able to differentiate resistance mechanisms even

further. We found that different forms of resistance had different impacts on the variance in load

(Fig. 4). Higher standard deviation, e.g., associated with constitutive resistance vs. inducible

resistance, increases the likelihood of exceptionally high load infections, leading to host death.

Even assuming lower mean log load for constitutive resistance to achieve the same average death

rate of infected hosts, we found that higher variance should lead to higher pathogen shedding per

infected host, increasing pathogen fitness, prevalence, and depressing the host population more

for constitutive resistance than inducible. Thus on the spectrum of resistance (increasing host

fitness and decreasing pathogen) to tolerance (increasing host fitness without decreasing

pathogen fitness), constitutive resistance is more "tolerant" than inducible resistance. Previous

compartmental models of constitutive vs. inducible resistance have considered how the two

forms of resistance may differ in whether their costs and benefits are expressed while susceptible

or infected [27,28]; but we assume that all three of our forms of resistance, constitutive,

inducible, and acquired, have constant costs and only provide benefits while infected yet we still

find they have different implications for the standard deviation, host fitness, pathogen fitness,

pathogen prevalence, and host density. Thus, dynamic heterogeneity in pathogen load reveals

important differences among different forms of resistance and points the way toward new

considerations for host-parasite coevolution.

Our work builds toward a greater understanding of how load dynamics shape host and

microparasite fitness (and thus perhaps coevolution) at the within- and between-host scales.

Previously, nested models have demonstrated that within-host dynamics, like pathogen growth,

influence tradeoffs between transmission rate and virulence, impacting pathogen evolution at

different epidemic stages [29]. And a previous moment closure of an IPM with dynamic mean

and constant variance showed that faster within-host growth can increase pathogen fitness

through higher shedding or decrease it if host mortality becomes too high [23]. Our work

emphasizes the importance of a dynamic standard deviation of load for both host and pathogen

fitness for their impact on pathogen spread as well as our ability to infer underlying processes.

These results suggest new avenues for studying host-pathogen coevolution. For example,

evolution by host or microparasite will affect how strongly stochastic mechanisms can generate

heterogeneity in load, altering the standard deviation that controls the probability of very high

load infections that shed at high rates (good for pathogens) and die quickly (bad for hosts).

Further, a model like ours could help investigate more fully the likely critical feedbacks between

the within-host and between-host scales, a criterion that makes such models particularly vital

[30]. For example, high force of infection from the environment may increase the mean load of a

new infection (would create a $\beta_j Z$ dependence of our G_0 function and allow avoidance resistance

to alter equilibrium load dynamics). Further, pathogen growth rates may depend critically on

dose (as dose has been found to influence many within-host dynamics: Kamiya et al. 2020) such

that low dose infections are eliminated by hosts but high dose infections grow out of control (we

could readily model this by modifying our $a+bx$ growth parameters to have $a < 0$, $b > 1$ or using

some non-linear function with a third parameter). Further empirical work could determine the

relevance of such functional forms while further theoretical work could determine their

implications, e.g., possible bistability between endemic and disease-free equilibria.

In addition to multiple conceptual advances, our model also provides a methodological

advance for certain applications to merge theory and data. We demonstrate that investigators

may be able to use the load dynamics of the mean *and* standard deviation to infer underlying

processes, such as which form of defence hosts are evolving. Additionally, the reduced

dimension of our model with the moment closure, compared to without the moment closure, will

improve the efficiency of fitting a model to population-level data (similar to the moment closure

in Bolker and Pacala 1997 for improving fitting efficiency for a different problem) including host

abundance, infection prevalence, and load dynamics. Because our model links within-host and

population-level processes, such a fitting procedure may be very useful for combining multiple

data types. For example, laboratory data may provide informative prior probability distributions

for some parameters, improving the ability of a model fitting procedure to infer other within-host

processes from population-level data. In our focal system, Mountain yellow-legged frogs are

likely evolving some form of resistance to pathogen growth [18,25,33]; ample data are collected

on abundance, infection prevalence, and the distribution of *Bd* load in recovering, wild

populations so that fitting a version of our model to these data may identify whether this

resistance is more likely constitutive, inducible, or acquired helping target future immunological

investigations and conservation actions (e.g., whether or not to immune prime reintroduced

frogs). While we use temporal trends in load patterns to infer temporal differences in within-host

processes, the same approach could be used for other trends (e.g., latitudinal trends in load and

within-host processes). Our model also serves as a resource for investigators interested in

lognormally distributed pathogens and may inspire investigators to use this same moment closure

approach to derive helpful models for other distributions (e.g., gamma distributions).

Further, our findings call for more investigation of how dynamic trait heterogeneity can

inform relatively simple ecological models for new insight. Our approach could be applied

directly to track dynamic mean and variance of other lognormally distributed, ecological traits

outside of disease systems (e.g., body size), and is particularly useful when there are non-linear

transition probabilities (e.g., survival), e.g., to Stubberud et al. 2019 [34]. Some of our insights

would apply most readily to traits that tend to change directionally over time, like body size, so

that the loss of individuals constrains the mean and variance closer to starting values, depending

on mean-variance feedbacks (see Table 1 for how these can work out). Our assumption of a

normal distribution of a non-heritable trait parallels the common simplifying assumption in

quantitative genetics of a normal distribution of a heritable, heterogeneous trait [1]. Additionally,

we found that trends in the variance of load can be very informative regarding underlying

mechanisms, a concept that may apply well to quantitative genetics. For example, [2] used a

quantitative genetics model to show how changes in mean trait can reveal upcoming population

shifts; within their model framework, changes in the trait variance may also reveal mechanisms

by which the mean shifts as we expect a change via selection should decrease trait variance (at

least temporarily) while a shift via adaptive plasticity may not. Given the conceptual insights that

can be obtained with relatively simple models, we recommend further investigations of how

heritable and non-heritable sources of trait variation interact to shape trait variance and

population-level processes.

Overall, our model results emphasize the value of considering load dynamics for critical,

ongoing questions of microparasite diseases. Previous theory and theory-data integration have

effectively utilized the mean of infection load, providing a strong theoretical foundation and

many datasets to leverage when considering a dynamic standard deviation of infection load.

Relatively simple models like ours provide computationally efficient, empirically relevant, and

analytically tractable tools for advancing our understanding of load dynamics for microparasites.

Further, we can address how classic aggregation metrics change with these models to see how

microparasite dynamics follow, or deviate from, patterns found for macroparasites. Improving

our understanding of load dynamics will allow us to infer within-host processes more efficiently,

predict host population resilience more accurately, and build a more complete understanding of

(co-)evolution of host-microparasite systems.

References

1. Abrams PA, Matsuda H, Harada Y. 1993 Evolutionarily unstable fitness maxima and stable fitness
minima of continuous traits. *Evolutionary Ecology* **7**, 465–487.

2. Baruah G, Clements CF, Guillaume F, Ozgul A. 2019 When do shifts in trait dynamics precede
population declines? *The American Naturalist* **193**, 633–644.

3. May RM, Anderson RM. 1978 Regulation and stability of host-parasite population interactions: II.
Destabilizing processes. *The Journal of Animal Ecology* , 249–267.

4. Lachs L, Biondi P, Gouezo M, Nestor V, Olsudong D, Guest J, Golbuu Y. 2024 Demographic recovery
of corals at a wave-exposed reef following catastrophic disturbance. *Coral Reefs* **43**, 193–199.

5. Stover JP, Kendall BE, Fox GA. 2012 Demographic heterogeneity impacts density-dependent
population dynamics. *Theoretical Ecology* **5**, 297–309.

6. Cressler CE, Bengtson S, Nelson WA. 2017 Unexpected nongenetic individual heterogeneity and trait
covariance in Daphnia and its consequences for ecological and evolutionary dynamics. *The American*
Naturalist **190**, E13–E27.

7. Gillespie J. 1973 Polymorphism in random environments. *Theoretical Population Biology* **4**, 193–195.

8. Wilber MQ, Briggs CJ, Johnson PT. 2020 Disease's hidden death toll: Using parasite aggregation
patterns to quantify landscape-level host mortality in a wildlife system. *Journal of Animal Ecology*

9. Anderson RM, May RM. 1978 Regulation and stability of host-parasite population interactions: I.
Regulatory processes. *The journal of animal ecology* , 219–247.

10. Kretzschmar M, Adler F. 1993 Aggregated distributions in models for patchy populations.
Theoretical population biology **43**, 1–30.

11. Wilson K, Bjørnstad O, Dobson A, Merler S, Poglayen G, Randolph S, Read A, Skorping A. 2002
Heterogeneities in macroparasite infections: patterns and processes. In *The ecology of wildlife*
diseases (ed PJR Hudson A Grenfell, BT Heesterbeek, H Dobson, A), pp. 6–44. Oxford University
Press.

12. Anderson RM, May RM. 1981 The population dynamics of microparasites and their invertebrate
hosts. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **291**, 451–
524.

13. Fraser C, Hollingsworth TD, Chapman R, de Wolf F, Hanage WP. 2007 Variation in HIV-1 set-point
viral load: epidemiological analysis and an evolutionary hypothesis. *Proceedings of the National*
Academy of Sciences **104**, 17441–17446.

14. Yang Q *et al.* 2021 Just 2% of SARS-CoV-2– positive individuals carry 90% of the virus circulating in
communities. *Proceedings of the National Academy of Sciences* **118**, e2104547118.

15. Lisboa T, Waterer G, Rello J. 2010 We should be measuring genomic bacterial load and virulence
factors. *Critical care medicine* **38**, S656–S662.

16. Wang G *et al.* 2023 Genomic bacterial load associated with bacterial genotypes and clinical
characteristics in patients with scrub typhus in Hainan Island, Southern China. *PLOS Neglected*
Tropical Diseases **17**, e0011243.

17. Frick WF, Cheng TL, Langwig KE, Hoyt JR, Janicki AF, Parise KL, Foster JT, Kilpatrick AM. 2017
Pathogen dynamics during invasion and establishment of white-nose syndrome explain mechanisms
of host persistence. Wiley Online Library.

18. Briggs CJ, Knapp RA, Vredenburg VT. 2010 Enzootic and epizootic dynamics of the chytrid fungal
pathogen of amphibians. *Proceedings of the National Academy of Sciences* **107**, 9695–9700.

19. Yang J-R, Lo J, Ho Y-L, Wu H-S, Liu M-T. 2011 Pandemic H1N1 and seasonal H3N2 influenza infection
in the human population show different distributions of viral loads, which substantially affect the
performance of rapid influenza tests. *Virus research* **155**, 163–167.

20. Bonhoeffer S, Fraser C, Leventhal GE. 2015 High heritability is compatible with the broad
distribution of set point viral load in HIV carriers. *PLoS pathogens* **11**, e1004634.

21. Euser S *et al.* 2021 SARS-CoV-2 viral-load distribution reveals that viral loads increase with age: a
retrospective cross-sectional cohort study. *International journal of epidemiology* **50**, 1795–1803.

22. Wilber MQ, Langwig KE, Kilpatrick AM, McCallum HI, Briggs CJ. 2016 Integral projection models for
host–parasite systems with an application to amphibian chytrid fungus. *Methods in Ecology and*
Evolution **7**, 1182–1194.

23. Wilber MQ, Pfab F, Ohmer ME, Briggs CJ. 2021 Integrating infection intensity into within-and
between-host pathogen dynamics: Implications for invasion and virulence evolution. *The American*
Naturalist **198**, 661–677.

24. Schrock SA *et al.* 2025 Do fungi look like macroparasites? Quantifying the patterns and mechanisms
of aggregation for host–fungal parasite relationships. *Proceedings B* **292**, 20242013.

25. Knapp RA, Wilber MQ, Joseph MB, Smith TC, Grasso RL. 2024 Reintroduction of resistant frogs
facilitates landscape-scale recovery in the presence of a lethal fungal disease. *Nature*
Communications **15**, 9436.

26. Råberg L, Graham AL, Read AF. 2008 Decomposing health: tolerance and resistance to parasites in
animals. *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**, 37–49.

27. Westra ER *et al.* 2015 Parasite exposure drives selective evolution of constitutive versus inducible
defense. *Current biology* **25**, 1043–1049.

28. Boots M, Best A. 2018 The evolution of constitutive and induced defences to infectious disease.
Proceedings of the Royal Society B-Biological Sciences **285**. (doi:10.1098/rspb.2018.0658)

29. Day T, Alizon S, Mideo N. 2011 Bridging scales in the evolution of infectious disease life histories:
theory. *Evolution: International Journal of Organic Evolution* **65**, 3448–3461.

30. Mideo N, Alizon S, Day T. 2008 Linking within-and between-host dynamics in the evolutionary
epidemiology of infectious diseases. *Trends in Ecology & Evolution* **23**, 511–517.

31. Kamiya T, Greischar MA, Schneider DS, Mideo N. 2020 Uncovering drivers of dose-dependence and
individual variation in malaria infection outcomes. *PLOS Computational Biology* **16**, e1008211.

32. Bolker B, Pacala SW. 1997 Using moment equations to understand stochastically driven spatial
pattern formation in ecological systems. *Theoretical population biology* **52**, 179–197.

33. Knapp RA, Fellers GM, Kleeman PM, Miller DA, Vredenburg VT, Rosenblum EB, Briggs CJ. 2016
Large-scale recovery of an endangered amphibian despite ongoing exposure to multiple stressors.
Proceedings of the National Academy of Sciences **113**, 11889–11894.

34. Stubberud MW, Vindenes Y, Vøllestad LA, Winfield IJ, Stenseth NC, Langangen Ø. 2019 Effects of
size-and sex-selective harvesting: An integral projection model approach. *Ecology and Evolution* **9**,
12556–12570.

708