The wheat VIH2-3B, a functional PPIP5K controls the localization of

fasciclin-like arabinogalactan protein

Author names and affiliations:

Anuj Shukla[1,2,3], Rahul Tanwar[1,4], Kanupriya Agrwal[1], Riya Ghosh[5], Henning Jacob

Jessen[2,3], Debabrata Laha[5] and Ajay K Pandey[1,*]

[1]National Agri-Food Biotechnology Institute (Department of Biotechnology), Sector 81,

Knowledge City, S.A.S. Nagar, Punjab, India.

[2] Institute of Organic Chemistry, University of Freiburg, Albertstraße 21, 79104,

Freiburg, Germany.

[3]CIBSS - Centre for Integrative Biological Signalling Studies, University of Freiburg,

79104 Freiburg, Germany.

[4]Regional Centre for Biotechnology, Faridabad - 121001 Haryana (NCR), Delhi, India

[5]Department of Biochemistry, Indian Institute of Science, Bengaluru-560012,

Karnataka, India

Corresponding author:

Dr. Ajay K Pandey, Scientist-F

Email: pandeyak@nabi.res.in; pandeyak1974@gmail.com

ORCID iD: https://orcid.org/0000-0003-1064-139X

Abstract

Inositol pyrophosphates (PP☐InsPs) are important signalling molecules, that participate in multiple physiological processes across a wide range of eukaryotes. Metabolic pathway kinases (VIP1/VIHs) leading to the production of PP-InsPs are now well characterized in yeast and plants. Previously, the wheat inositol pyrophosphate kinase (*TaVIH2*) was shown to encode a catalytic active kinase domain. Heterologousexpression of *TaVIH2* in *Arabidopsis thaliana* showed enhanced drought tolerance by modulating the cell composition. In this study, we attempted to identify the interacting protein targets of wheat VIH2-3B using a yeast two-hybrid (Y2H) cDNA library screen, identifying 52 putative interactors that are primarily involved in cell wall-related functions. Notably, fasciclin-like arabinogalactan protein (FLA7), a glycosylphosphatidylinositol (GPI)-anchored protein emerged as the most frequently interacting partner in the screen. Further analysis using pull-down assays validated the interaction between TaVIH2-3B and TaFLA7 *in vivo*. Using the reporter fusion studies, we observed the localization of TaFLA7 to be a plasma membrane. We also observed that this localization of the FLA7 was perturbed in the yeast *vip1Δ* strain. The expression of TaVIH2-3B bearing PP-InsP5K enzymatic activity in yeast mutants rescued the localisation of the FLA7 to the membrane. Expression analysis of *TaFLA7* showed differential expression response under drought in wheat shoot tissues. *TaFLA7* was also found to be highly expressed during grain development, particularly in the endosperm and seed coat during grain maturation. Taken together, these findings highlight the potential role of TaVIH2 in cell wall remodelling and stress response pathways, offering new insights into the functional roles of VIH proteins in plants.

Key word: Inositol pyrophosphates, kinases, cell wall maintenance.

Introduction

Inositol pyrophosphates (PP-InsPs) are high-energy molecules that are ubiquitous in various eukaryotic organisms and serve as critical cellular signaling entities(Zsolt et al. 2011). Over the last two decades, inositol pyrophosphate signalling has gained significant attention and proven to regulate different cellular functions such as regulating telomere length, DNA repair, phosphate homeostasis, vesicular trafficking and protein phosphorylation (Saiardi 2004; Fridy et al. 2007; Lee et al. 2007; Azevedo et al. 2009; Thota and Bhandari 2015; Shears 2015; Sanchez et al. 2019; Pipercevic et al. 2023; Chabert et al. 2023). Researchers have identified two distinct classes of inositol pyrophosphate synthases, inositol hexakisphosphate kinases (referred to as IP6Ks or Kcs1-like proteins) and diphosphoinositol pentakisphosphate kinases (known as PP-IP5Ks or Vip proteins) (Saiardi et al. 1999; Mulugu et al. 2007). These enzymes are primarily classified based on their catalytic specificity towards various positions on the inositol ring and their dual-domain protein architecture having a "rim-K" domain at the N-terminal and an ATP-grasp domain at the C-terminal (Lee et al. 2007; Mulugu et al. 2007).

Genes encoding VIH-type PP-InsP synthase could be identified in green algae and embryophytes (Desai et al. 2014; Laha et al. 2015). *Arabidopsis* VIH proteins catalyze the synthesis of the inositol pyrophosphate, $InsP_8$ and VIH-derived $InsP_8$ has been linked with diverse physiological processes such as plant defence and phosphate homeostasis (Laha et al. 2016; Jung et al. 2018; Dong et al. 2019; Zhu et al. 2019; Ried et al. 2021; Riemer et al. 2021) . It was further shown that $InsP_8$ binds to SPX1 protein and controls its interaction with the phosphate starvation response 1 (PHR1), a central regulator of phosphate starvation (Jung et al. 2018; Dong et al. 2019; Ried et al. 2021). It seems logical to conclude that higher levels of InsPx are co-related with the gain of function against different physiological processes.

Although the level of PP-InsPs seems to be important in asserting such function, it remains to be studied if the responsible kinases, especially plant VIH could interact physically with the components of the specific signalling system. One consequence could be local enrichment of PP-InsP species in the vicinity of their target proteins. There are multiple reports that suggests the employment of such gating of PP-InsP to its target proteins through physical interaction of the responsible PP-InsP synthase (Rao et al. 2014; Zhou et al. 2021; Laha et al. 2022; Bhandari et al. 2024). A comprehensive interaction studies for VIH protein is missing in plants that could provide new insights into how their signaling output is regulated.

Multiple VIH proteins were identified in hexaploid wheat with diphosphoinositol

pentakisphosphate kinases (PP-InsP5K) like enzymatic activity (Shukla et al. 2021). Notably,

when wheat VIH homolog (TaVIH2-3B) was expressed heterologously in *Arabidopsis*, they

conferred tolerance to drought conditions (Shukla et al. 2021). Further, it was shown that

VIH2 overexpressed lines show reprogramming of genes involved in cell-wall-related

metabolic pathway genes. At the metabolome level, these transgenic lines showed high

accumulation of cell-wall cementing polysaccharides such as arabinoxylan and

arabinogalactan (Shukla et al. 2021). Although, the role of VIH was established in imparting

drought tolerance in plants; it remains to be dissected if VIH could be involved at the protein

level. Furthermore, it remains to be studied if VIH could function by protein-protein

interactions.

Using a yeast two-hybrid (Y2H) screening system, we identified multiple interactors

of TaVIH2-3B. Further, a detailed characterization was done for one of the interactors

referred to as fasciclin-like arabinogalactan protein (FLA7) which is involved in the cell wall

cementing process. Using biochemical experiments and a yeast genetic screen, we

demonstrated that the presence of VIH-type PP-InsP synthase is required for FLA7

localization. This study expands our knowledge of the functioning of the plant VIHs towards

new roles including cell-wall homeostasis.

Materials and methods

Wheat cDNA library construction and yeast two-hybrid screening

Total RNA extracted from the wheat seedlings was used to purify mRNA. mRNA was

isolated by a plant mRNA isolation kit (Sigma). The mRNA concentration and purity were

checked by nanodrop and 0.8% (w/v) agarose gel, then visualized by ethidium bromide

staining under U.V light. Multiple batches of mRNA were pooled together to prepare the

cDNA library, and single-strand cDNA was constructed as described in the materials and

methods section. Next, the double-strand cDNA was prepared through long-distance PCR,

and aliquots were analyzed in the 1.2% agarose/EtBr gel. Finally, the ds-cDNA was first

purified using CHROMA SPIN+TE-400 columns. The wheat cDNA library was prepared by

the co-transformation of the pGADT7-Rec vector provided in the kit and purified cDNA

prepared (in the above section) in the Y187 yeast strain. The transformation mixture was

diluted (10^{-1} and 10^{-2}), and 100 µl of each dilution was plated on the SD-L agar plates to

select independent clones, indicating the readiness of our library's complexity.

Gene expression analysis by quantitative real-time PCR

The hexaploid wheat cv. C-306 was used for this study. The grains of cv. C306 were

stratified overnight in the dark at 4°C and then germinated for 4 days on Petri dishes lined

with Whatman filter paper. The seven-day-old seedlings were subjected to drought mimic

stress by exposing the roots of seedlings to 30% glycerol for six days along with Hoagland

media. Control treatment was without any presence of glycerol in the media. Root and shoot

tissue were collected post-treatment and Total RNA was extracted from the tissue from the

three biological replicates. Expression in grains was studied as described earlier (Bhati et al.,

2014). To quantify the gene expression in roots, quantitative real-time PCR (qRT-PCR) was

performed using the QuantiTect SYBR Green RT-PCR Kit (Qiagen, Germany). The gene-

specific primers capable of amplifying the genes mentioned in Table S1 were carefully

designed using Oligocalc software. The cDNA dilutions to be used were optimized by

running a semi-quantitative PCR for 22-25 cycles using Thermo Scientific™ DreamTaq™

Green PCR Master Mix (2X). The qRT-PCR reaction was set up in a 96-well plate by mixing

the reaction components including 1X QuantiTect SYBR Green RT-PCR Master Mix,

appropriate cDNA dilution, 900nM of forward and reverse primers and MQ to final volume

of 10µl in each well and the plate was run in 7500 Fast Real-Time PCR System. 4-5 technical

replicates for each set of primers and a minimum of 2 biological replicates were used to

validate the experiment. ADP-ribosylation factor gene (*TaARF*) was used as an internal

control in all the expression studies (Meena et al. 2024). The Ct values obtained after the run

were normalized against the internal control and relative expression was quantified using

$2-\Delta\Delta CT$ method (Livak and Schmittgen 2001). The results for each sample are presented as

the mean ± standard deviation (SD) from three replicates. Student's t-test was performed to

assess the significance of differences between samples. A p-value of less than 0.05 (*) was

considered statistically significant.*P* value < 0.05.

Co-immunoprecipitation of interacting proteins

For pull-down assays, respective yeast cultures were grown in SD media with appropriate

amino acid dropouts. The primary culture was grown overnight at 30 ˚C until an OD_{600} of 1

was achieved. The secondary culture was incubated for 6 hours, cells were washed, and the

lysate was prepared using glass beads in buffer (1% SDS, 100mM NaCl,100mM Tris-Cl,

1mM EDTA, 2% Triton and 1mM protease inhibitor (100X Halt protease inhibitor,

Thermofisher, U.S.A.). The protein concentration in lysate was calculated at two dilutions.

Equal amounts of each lysate were taken and incubated with 100% of Protein-G agarose

beads and 2 ul of anti-c-MYC antibody for overnight at 4 ˚C. The lysate was subsequently

washed three times with lysis buffer and centrifuged for 30 sec at 800 x g. 50 ul of loading

dye was added to the washed beads and heated at 65 C for 10 min. After the SDS-PAGE run,

the proteins were transferred to the PVDF membrane and processed. Blot was separated into

two parts to detect TaVIH2 and TaFLA7 separately. Different primary antibodies were used

for probing (mouse Anti-c-MYC and rabbit anti-HA with 1:2000 dilution). After washing the

blots with TBST, they were treated with the secondary antibody (Goat Anti-Mouse IgG (H +

L); and Goat Anti-Rabbit IgG (H + L) with 1:5000 dilution. After subsequent washing with

TBST, the blot was developed by using BIO-RAD clarity western ECL Substrate.

Localization studies

TaFLA7 was cloned in pGADT7 vector at *EcoR*1 and *BamH*1 sites for the localization

experiments. TaVIH2 cDNA was cloned in the pGBKT7 vector using Sma1 and Not1

restriction sites. The constructs were transformed in the Y2H Gold yeast strain and selected

on SD-Leu or SD-Trp plates. Yeast spheroplasts were prepared for localization as described

earlier (Severance et al., 2004). Mouse monoclonal Anti-HA (H.A.:FLA7-pGADT7) : or

rabbit Anti-c-Myc (cMYC: VIH2-pGBKT7) primary antibody (Invitrogen, USA) was used

for the respective preparations at a ratio of 1:200 followed by 5 washing with blocking buffer.

Yeast cells were incubated with Goat Anti-Mouse IgG (H+L) Alexa Flour Plus 488 or Goat

Anti-Rabbit IgG (H+L) Alexa Flour Plus 647 (Invitrogen, USA) at a ratio of 1:500 for 4hr at

room temperature. Cells were washed with blocking buffer and mounted with Fluor mount

(Sigma, USA). Representative fluorescent images were taken using LEICA DM600CS using

a 64X oil objective.

Luciferase Assay

Coding sequences (CDS) of TaVIH2 and TaFLA7 was amplified using forward primers

containing *BamH*I and reverse primers containing *EcoR*I, ensuring that the open reading

frame (ORF) remained intact. The amplified products were then cloned into pCAMBIA1300-

NLuc and pCAMBIA1300-CLuc vectors. The recombinant plasmids were initially

transformed into *E. coli* DH5α cells for propagation and were confirmed by sequencing.

Subsequently, the verified plasmids were transformed into the *Agrobacterium tumefaciens*

GV3101 strain. Positive clones of GV3101 containing the recombinant plasmids were

selected and cultured overnight with appropriate antibiotics. The cells were pelleted and

resuspended in freshly prepared induction media containing 10 mM MgCl□, 10 mM MES

(pH 5.7), and 150 μM acetosyringone. The optical density (OD) of each culture was adjusted

to 0.5, and 1 mL of the Nluc-expressing cells was mixed with 1 mL of the Cluc-expressing

cells. The mixture was incubated at 28°C without shaking for 5–6 hours before

agroinfiltration into tobacco leaves (10 μL per spot, with three spots per leaf). The infiltrated

plants were maintained in a humidity chamber, kept in the dark for 12 hours, and then

exposed to light for 24 hours. Finally, the leaves were harvested and chemiluminescence was

detected, with an exposure time set at 20 seconds.

Extraction of Inositol Pyrophosphates

S. cerevisiae cultures were grown at 30°C in Synthetic Complete Dextrose (SCD) media until

they reached exponential growth, with an A_{600} between 0.6 and 0.8. Cells equivalent to 10

A_{600} units were collected by centrifugation. After washing with ice-cold water, the cells were

resuspended in 500 μL of 1 M perchloric acid. The samples were quickly frozen using liquid

nitrogen and stored at −80°C. Upon thawing, the samples were briefly vortexed, and cell

debris was removed by centrifugation at 16,200 g for 5 minutes at 4°C. The supernatants

containing acid-extracted inositol polyphosphates were then purified using titanium dioxide

beads (GL Sciences 5020–75000), following the method described by (Wilson and Saiardi

2017). Specifically, the supernatants were combined with 5 mg of TiO2 beads per sample,

pre-washed with water and 1 M perchloric acid, followed by incubation on a nutator at 4°C

for 20 minutes. The beads with bound inositol polyphosphates were collected by

centrifugation at 5,000 g for 1 minute at 4°C, followed by two washes with 1 M perchloric

acid. To elute the inositol polyphosphates, the beads were resuspended twice in 250 μL of 3%

ammonium hydroxide, rotated for 5 minutes at 4°C, and then centrifuged at 5,000 g for 1

minute. The combined eluates (totalling 500 μL per sample) were dried using a vacuum

centrifuge (Eppendorf SpeedVac) at room temp. for approximately 5 hours. The dried

samples were reconstituted in 20 μL of water for capillary electrophoresis electrospray

ionization mass spectrometry (CE-MS) analysis.

Capillary electrophoresis–mass spectrometry (CE-MS) analysis

The CE-MS analyses were conducted as described in (Qiu et al. 2020, 2023; Saiardi et al.

2021). An Agilent CE-QQQ system was used for the analysis, which featured an Agilent

7100 capillary electrophoresis unit, an Agilent 6495C Triple Quadrupole mass spectrometer,

and an Agilent Jet Stream electrospray ionization source, all connected via an Agilent CE-

ESI-MS interface. The sheath liquid, a 50:50 combination of isopropanol and water, was

given at a steady flow rate of 10 µL/min using an isocratic Agilent 1200 LC pump and a

splitter to ensure exact flow. A 100 cm long fused silica capillary with an internal diameter of

50 µm and an outside diameter of 365 µm was employed to separate the samples.

The background electrolyte (BGE) contained 40 mM ammonium acetate that had been

adjusted to pH 9.08 with ammonium hydroxide. Before each sample analysis, the capillary

was flushed with BGE for 400 seconds. Samples were delivered into the system by applying

a pressure of 100 mbar for 15 seconds, resulting in an injection volume of around 30 nL. The

MS source settings included a gas temperature of 150˚C, flow rate of 11 L/min, nebulizer

pressure of 8 psi, and sheath gas temperature of 175˚C. The capillary voltage was set to -

2000V, with a nozzle voltage of 2000V. Furthermore, the negative high-pressure radio

frequency (RF) and negative low-pressure RF were kept at 70 and 40 volts, respectively. The

multiple reaction monitoring (MRM) parameters were set up as shown in Table 1.

The internal standard (IS) stock solution was created using the method described in (Qiu et al.

2020; Saiardi et al. 2021). Which was then added to the samples to help with isomer

assignment and quantification of InsPs and PP-InsPs. 5 µL of IS was added to 5 µL of sample

and carefully mixed. IPs were quantified by adding known amounts of the matching heavy

isotopic references to the samples. After spiking, the final concentrations in the samples were

as follows: 4 µM [$^{13}C_6$] 2-OH InsP$_5$, 20 µM [$^{13}C_6$] InsP$_6$, 1 µM [$^{13}C_6$] 5- InsP$_7$, 1 µM [$^{13}C_6$] 1-

InsP$_7$, and 1 µM [$^{13}C_6$] 1,5- InsP$_8$. Dorothea Fiedler provided the [13C6] inositol STD

(Harmel et al. 2019).

Results

VIH proteins contain variable IDR regions

Studies in the yeast and mammalian systems have shown that different inositol

pyrophosphate kinase function is based on their protein interaction networks (Gokhale 2013;

Rao et al. 2014; Wright and Dyson 2015; Rogers et al. 2021a). The human PPIP5K

(HsPPIP5K) interaction studies show the typical dual domains involvement and requirement

of the C-terminal intrinsically disordered region (IDR) for interaction (Machkalyan et al.

2016). This HsPPIP5K contains a large C-terminal IDR region (~250-550 aa), a missing

characteristic for most plant VIHs (Figure 1& Figure S1). The IDR region was checked in the

plant VIH proteins. Our analysis indicated that plant VIH proteins have a variable and lower

percentage of IDR representation. The plants VIH show 12-22 % of IDR region (Table 2).

Interestingly, wheat TaVIH possesses a higher IDR % when compared to the *Arabidopsis*

(Figure 1).

To get insights into the direct and indirect interactions, we performed a String analysis

of VIH proteins. Human PPIP5K and yeast (VIP1) analysis resulted in the protein-protein

network analysis with nodes pointing towards the PHO-mediated interactors for the VIP and

involvement of HsPPIP5K in other biochemical activities (Figure 2). A similar analysis of

wheat VIH2-3B protein shows multifunctional globular proteins such as actin as probable

interactors. The presence of IDR in the wheat VIH2 and the preliminary String analysis led us

to explore its interacting partners.

Screening of TaVIH2 interacting partners

To get molecular insights into the role of TaVIH2 proteins in stress tolerance, we used yeast

two-hybrid (Y2H) cDNA library screen to identify its possible interacting partners. Protein

expression of bait (VIH2) in the yeast cells was confirmed by immunoblot analysis

(Supplementary Figure S2A). Two pooled wheat cDNA libraries were prepared for Y2H

studies. The resulting smear between 300 bp to 5 kb confirmed the good quality of ds-cDNA

(Supplementary Figure S2B). Y2H was performed using TaVIH2-3B as a bait and this

resulted in the identification of 89 putative yeast colonies with a mating efficiency of 3.8 %.

Subsequent screening of the colonies led to the identification of ~52 putative interactors

(Supplementary Figure S2C&D). Amplicons were sequenced and clones appearing more

than twice were considered for further studies (Supplementary Figure S3). The analysis led to

the shortlisting of ten strong potential interactors with high occurrence in the screening

(Table 3). Amongst these interactors, most of the genes encode for cell-wall-related function,

including fasciclin-like arabinogalactan protein (FLA), glycosyl-transferases, and glycine-

rich structural proteins. The most frequently (nine times) interacting clone was identified as

FLA7 (*TraesCS2D02G172900*) protein, and further detailed characterization was performed

with this protein. Our analysis indicated that TaFLA7 encodes a 367 aa protein containing

FAS-like arabinogalactan protein and the presence of a typical transmembrane domain

(TMD) with glycophosphatidyl-inositol (GPI) domain along with hydrophobic domains

(Figure S4). Additionally, the protein hydropathy plot identified a hydrophobic region near

the GPI region at the C-terminal.

Wheat VIH2 interacts with fasciclin-like arabinogalactan protein (FLA)

To further confirm this interaction, full-length coding sequence of *TaFLA7* (1.2 kb) was
cloned (TaFLA7:AD+TaVIH2:BD) and co-transformed along with TaVIH2 in the yeast
strain. The yeast co-transformed with TaFLA7 and TaVIH2 showed strong reporter activity
(Figure 3A). To further confirm this interaction, yeast colonies were grown on -His/-Leu/-Trp
(Auerobasidin) media (Figure 3B) and the protein extracts were subjected to *in
vivo* pull☐down assay. For this, first we checked the expression of HA-tagged FLA7 and c-
MYC-tagged TaVIH2 in the Y2H Gold yeast strain. Western analysis for both proteins
(TaFLA7 ~55 kDa) indicated optimal expression in the cells (Figure 3B). Furthermore, only
the TaVIH2 was detected in the yeast strain lacking TaFLA7 (Figure 3B) compared to the
control (Figure 3B). However, in the pull-down analysis of TaFLA7 and TaVIH2 co-
transformed cells, both the TaFLA7 and TaVIH2 proteins were detected (Figure 3B; Lane 4).
These findings imply that TaVIH2 and TaFLA7 show protein-protein interaction. To further
validate this interaction, we performed split luciferase complementation assayusing tobacco
leaf (*Nicotiana benthamiana*) with the fused protein NLuc-TaFLA7)/CLuc-TaVIH2.
Luciferase complementation imaging (LCI) pair reconstituted luciferase activity in regions
where both the fused proteins were expressed when compared to controls (Figure 4A). RLU
measurement suggested an increase of 7-8 folds of the luminescence activity compared to the
controls (Figure 4B). Overall, these results confirmed that TaVIH2 nteract with cell-wall
cementing proteins.

VIH2 facilitates the localization of FLA7

Next, we studied the localization of the FLA7 protein in yeast cells. Confocal imaging of
yeast cells expressing TaFLA7:GFP suggests its presence on the plasma membrane (Figure
5A). Since the wheat FLA7-VIH2 interact and FLA7 is localized to the membrane, we
wonder whether VIH2-3B contributes to the localization of FLA7. Previously, we reported
that *TaVIH2-3B* overexpressing lines enhanced the expression of genes involved in cell wall
biosynthesis (Shukla et al. 2021). TaVIH2-3B was shown to utilize IP$_7$ as a substrate likewise
to yeast VIP1. Therefore, we tested the localization of FLA7 in different yeast backgrounds.
As expected, in the WT yeast, FLA7 was localized on the plasma membrane, but
surprisingly this localisation was affected in the VIP1-deficient yeast cells (Figure 5B). This
suggested that localization of the FLA7 is dependent on VIP1 activity. Next, we
complemented the yeast *vip1Δ* mutant and checked for the FLA7 localization. Interestingly,

the presence of TaVIH2-3B restores the localisation of TaFL6 to the plasma membrane in the

the yeast *vip1Δ* strain.

Additionally, we have also profiled the cellular pools of $InsP_6$, $5\text{-}InsP_7$, $1\text{-}InsP_7$, and

$1,5\text{-}InsP_8$ with capillary electrophoresis electrospray ionization mass spectrometry (CE-ESI-

MS), as described in (Qiu et al. 2020, 2023; Saiardi et al. 2021; Liu et al. 2023; Eisenbeis et

al. 2023). This method allows us to investigate the cellular levels of different InsP and PP-

InsP species in the yeast transformants. To account for any differences in sample recovery

during extraction and TiO_2 bead affinity enrichment, the levels of $5\text{-}InsP_7$, $1\text{-}InsP_7$, and $1,5\text{-}$

$InsP_8$ were normalized to $InsP_6$ levels (Figure. 6). The observed changes in the $1,5\text{-}$

$InsP_8$ levels (Figure. 6) between the WT, *vip1Δ* mutant and *vip1Δ*::TaVIH2-3B complimented

extracts confirmed that TaVIH2 is a functional $InsP_8$ synthase in yeast and it is likely that the

localization of TaFLA7 is dependent of the TaVIH2-derived $InsP_8$. Collectively, our results

indicate that TaVIH2-3B interacts with the FLA7 protein and could facilitate its localization

to the plasma membrane. Moreover, the differences in $1,5\text{-}InsP_8$ levels witnessed among the

WT, *vip1Δ* mutant, and *vip1Δ* mutant complemented with TaVIH2-3B suggest that the

appropriate positioning of TaFLA7 is associated with the activity of the PP-InsP5K enzyme.

FLA7 is differentially expressed in wheat tissue and is induced by drought condition

The FLA proteins are known to respond to different stress conditions and are involved in

plant growth and development (Zang et al. 2015; Cagnola et al. 2018; Wu et al. 2020; Ma et

al. 2022; Yao et al. 2023). These observations suggest that VIH2 and its interacting partners

could participate in similar pathways. To further characterize the wheat *FLA7*, we studied its

expression in different tissues and under different stress conditions. Our analysis suggested

that TaFLA7 was expressed in the grain at the Z71 stage (watery ripe with the start of grain

filling) and in the spike at Z65 (50% of the spikes had yellow anthers) of wheat development

(Figure 7A). Expression analysis of *TaFLA7* in different wheat grain tissue using the eFP

wheat browser revealed its high expression in embryo and aleurone layers of the grain

(Figure 7B). The eFP wheat browser analysis indicates the high expression of FLA7 in

grains, shoot axis and spikelet (Figure 7B&C; Figure S5, Figure S6). In grains the expression

was higher in the outer and inner pericarp region. Our qRT-PCR during garin development

suggest increasing expression of TaFLA7 mRNA with its maturation with highest expression

at 28 days after the anthesis DAA (Figure 7D). Based on the previous in-silico data the

transcript expression of *TaFLA7* showed significant upregulation in the shoot tissue when

subjected to drought stress after 1 hr and 6 hrs (Figure 7E). Our quantitative RT-PCR results

also confirmed the significant upregulation of TaFLA7 in the shoot or root tissue subjected to

drought stress (Figure 7F). Altogether the expression data of the *TaFLA7* show transcriptional

changes during desiccation, a prerequisite step in grain maturation.

Discussion

Protein-protein interaction networks provide valuable insights into biological processes and

complex metabolic functions within living cells. The Y2H system leverages eukaryotic

transcriptional activators with distinct functional domains for DNA-binding and

transactivation, making it a powerful and rapid approach to discovering new protein

interactions (He et al. 2019). In this study, a high-quality yeast two-hybrid (Y2H) library

using RNA extracted from wheat seedlings was constructed. Our work identified several

interactors of TaVIH2 proteins, which serve as promising entry points for further exploration

of their regulatory network in the developmental process. Recently, it was demonstrated that

the *Triticum aestivum* (hexaploid wheat) genome encodes functional PPIP5K isoforms,

identified as TaVIH2-3B (Shukla et al. 2021). Moreover, the heterologous expression of

wheat TaVIH2-3B in *Arabidopsis* imparts drought tolerance. These *Arabidopsis* transgenic

lines expressing TaVIH2 also exhibit a reprogramming of gene expression involved in cell-

wall metabolism and stress response pathways (Shukla et al. 2021). In particular differential

expression of genes linked to ABA biosynthesis, cell wall maintenance, and cytochrome P-

450 monooxygenases were highly upregulated. Although the level of InsPs seems to be

important in imparting such a function, it remains to be studied if plant PP-InsP synthase,

such as VIH at the protein level, could be involved directly.

Multi-protein interactions were more accessible by IDRs, which helped them connect

with other proteins in the cell. Abelson (Abl), a protein non-receptor tyrosine kinase, has an

IDR domain (around 900 amino acids) that has been implicated in a variety of tasks in

Drosophila, including the regulation of cell shape and cytoskeleton development (Rogers et

al. 2021b). Similarly, the human PPIP5K domains' association with proteins involved in actin

cytoskeleton structure and phosphatidylinositol biosynthesis was attributed to the IDR's

existence (Machkalyan et al. 2016). The interaction studies were pursued using human

PPIP5K (HsPPIP5K) (Machkalyan et al. 2016). Human PPIP5K, besides containing the

typical dual-domain, also possesses a C-terminal IDR (Machkalyan et al. 2016). This IDR in

HsPPIP5K spans a large portion (~250-550 amino acids), a feature that is not very common

in most plant VIHs (Figure 1). IDRs assist in their interaction with multiple proteins to

facilitate their cellular activities (Ward et al. 2004).

Although VIH proteins have been identified in several plants (Laha et al. 2015; Zhu et

al. 2019), including wheat (Shukla et al. 2021), the IDR domains are relatively smaller,

ranging from approximately 33-56 amino acids, and their functional roles remain to be

elucidated. Our network analysis using the String database can provide clues for the

regulatory network of the VIH proteins (Figure 2). Human PPIP5K and yeast (VIP1) analysis

resulted in the protein-protein network analysis with nodes pointing towards the PHO-

mediated interactors for the VIP and involvement of HsPPIP5K in other biochemical

activities (Figure 2). A similar analysis of wheat VIH shows a network with cytoskeletal actin

as a probable interactor. Actin is involved in forming microfilaments in the cytoskeleton and

can modulate the cellular shape (Pollard and Cooper 2009). This study identified multiple

interactors of VIH2, yet most of them are speculated to be involved in cell wall maintenance.

The speculative parallels of VIP1 in yeast and VIH encouraged us to postulate that plant VIH

is likely involved in cell wall cementing and homeostasis. The conserved role of VIH was

demonstrated in the important cellular function of Pi-homeostasis (Dong et al. 2019; Zhu et

al. 2019; Riemer et al. 2021). Under nutrient-limiting conditions, VIP1 and its ortholog in

Schizosaccharomyces pombe (Asp1) kinase were shown to control physiological changes

such as dimorphic switch, cell-to-cell, cell-substrate adhesion and unipolar budding

(Pöhlmann and Fleig 2010; Pöhlmann et al. 2014). The ability of TaVIH2 to facilitate the

localization of TaFLA7 (Figure 5) provides new insight into the functioning of VIH proteins.

The involvement of these proteins in this process points towards their ability to control the

cellular boundaries and shape (Figure 8). Such phenotypic changes, including the formation

of pseudo-hyphal filaments in yeast (Norman et al. 2018), help sustain themselves under

stress conditions, yet their similar involvement in plants needs more investigation. Previous

biochemical investigations of TaVIH2-3B overexpressing lines in *Arabidopsis* indicated

changes in cell wall cementing metabolites such as cellulose, arabino-xylan, and arabino-

galactan (Shukla et al. 2021). Furthermore, RNAseq analysis of the Col-0 and TaVIH2

overexpressing lines suggested high expression of genes involved in cell wall biosynthesis

(Shukla et al. 2021). Together with these results, our findings suggest that VIH2 can regulate

the composition of cell wall cementing pathways by localizing the involved proteins to their

correct positions. The intriguing role of VIH protein overexpression in modulating cell wall

composition requires further investigation. Specifically, the *in-vivo* levels of $InsP_7/InsP_8$

(Figure 6) suggest that TaVIH2-3B possess PP-IP5K enzymatic activity. Therefore, either the

synthesis of InsP$_8$ or the presence of TaVIH2-3B protein in the yeast cell directly influences
the correct localization of the TaFLA7 to the plasma membrane. Furthermore, it is crucial to
investigate how this localization occurs and to identify which region of TaFLA7 interacts
with TaVIH2-3B and understanding the role in such interaction may also be critical for future
study. In the aftermath of the current study, it will be interesting to check if InsP$_7$/InsP$_8$ could
modulate the protein-protein interactions of transcription factors that are primarily involved
in water-deficient stress conditions. This remains to be investigated and could offer new
mechanistic insights into drought tolerance. Studies using *Arabidopsis* have deciphered
multiple roles of VIHs in different biological processes. However, such efforts should be
expanded to crop plants such as wheat for translational research.
447

Acknowledgements
The authors thank the Executive Director for the facilities and support. This study was
supported by the Department of Biotechnology, Basic Plant Biology Grant to AKP
[BT/PR12432/BPA/118/35/2014]. Part of this work was also supported by the NABI-CORE
grant to A.K.P. and by the Deutsche Forschungsgemeinschaft (DFG) under Germany's
Excellence Strategy CIBSS, EXC-2189, Project ID 390939984 to H.J. DBT-eLibrary
Consortium (DeLCON) is acknowledged for providing timely support and access to e-
resources for this work. A.S. acknowledges Centre For Integrative Biological Signalling
Studies for PostDoctral Fellowship. D. L. acknowledges the Department of Biotechnology
(DBT) for HGK-Innovative Young Biotechnologist Award (BT/13/IYBA/2020/04).

458
459 **Declarations**
460 **Conflict of interest**
461 The authors declare that they have no known competing financial interests or personal
462 relationships that could have appeared to influence the work reported in this manuscript.
463
464 **Authors contribution**
465 AKP and AS conceived the project. AS, AKP and RT designed the experiments. AS, RT, RG
466 and KA performed the experiments. AKP, AS, DL and HJ wrote and revised the draft
467 manuscript. All the authors commented on the manuscript and agreed to the submission of
468 the manuscript.
469
Data Availability

No datasets were generated or analysed during the current study.

Figures for the legends:

Figure 1: *In-silico* calculation of the Intrinsic Disorder Regions (IDR) of VIH and

PPIP5K proteins. IDR was calculated based on the xxx along with the percentage region

score of the mentioned proteins.

Figure 2: Protein association networks were identified for each of the human

(PPIP5K1), yeast (VIP1) and wheat (VIH; Traes_3B_3567813F3.1). The network analysis

for each of the proteins was performed using STRING and was represented as the coloured

lines with the description as given in the figure.

Figure 3: Interaction of TaVIH2-3B with TaFLA7 as identified during Y2H screening.

(A) Y2H assay using full-length TaFLA7 and VIH as represented by GAL plates containing -

L-T and without GAL plates containing -HLT. (B) Pull-down assay and Western analysis of

the wheat FLA7 in the yeast strains.

Figure 4: Luciferase reporter assay to characterization of TaFLA7 interaction with

FLA7. (A) Luciferase assay of TaFLA7 with VIH2 using tobacco leaves. **(B)** RLU

measurement for the strength of interaction. The values shown are the mean $\pm$ SE; n = 4.

Student's *t* test was performed (*$p < 0.001$).

Figure 5: The localization of TaFLA7 and its dependency on the VIH2 activity. (A)

Protein structure arrangements of TaFLA7 protein and localization in yeast cells. (B) In situ

localization of the TaFLA7 to the membrane in the yeast vip1 mutant transformed with

TaVIH2-3B. Localization of FLA7 using HAtagged-TaFLA7 in different yeast strains.

Fluorescence was measured using a Zeiss fluorescence microscope (Axio Imager Z2) with an

Axiocam MRm camera at 63X. Representative images are shown and similar observations

were noted for 3-4 independent transformed colonies of yeast.

Figure 6: Quantification of inositol pyrophosphate levels using CE-ESI-MS. The inositol

pyrophosphate levels in extracts of the WT + pYES2 , *vip1Δ* + pYES2, and *vip1Δ* + TaVIH2-

3B strains that were transformed with TaFLA7 are displayed for 1,5-InsP$_8$, 5-InsP$_7$, and 1-

InsP$_7$, normalized to the levels of InsP$_6$. The values shown are the mean $\pm$ SE; n = 3.

Student's *t* test was performed (*$p < 0.001$).

Figure 7. Expression characterization of TaFLA7 in wheat tissue. (A) Relative gene

expression level during plant development (at Zadosky scale; Exvip database;

https://dubcovskylab.ucdavis.edu/expression_data/). (B) Expression (TPM: transcript per

million) of the FLA7 in developed grains of wheat. Heat map indicates Log$_2$ratio. (C)

Expression (TPM: transcript per million) of TaFLA6 in grain layers; endosperm, inner

pericarp and outer pericarp.   (D) Expression of TaFLA7 during grain development. stages of

embryo development. (E) The expression profiles of the TaFLA7 in roots and shoot tissue of

wheat seedlings collected at 6 days under drought stress. (F) qRT-PCR analysis of TaFLA7

after drought mimic condition (30% glycerol) in shoots and roots. CR: control roots; RT:

Root treatment; CS: Control shoot: and ST: shoot treatment.

Figure 8: Schematic representation drawing parallels for the possible function of VIH

proteins in multiple cellular processes. In yeast, VIP proteins and their products were

shown to be involved in the cell boundary rearrangements which include regulation of

microtubule rearrangement and filamentous growth of the fungi under nutrient-limiting

conditions (Pöhlmann and Fleig 2010; Pöhlmann et al. 2014). Plant VIHs are known to be

involved in phosphate (Pi) homeostasis and plant defence (Laha et al. 2015; Dong et al. 2019;

Riemer et al. 2021). Recently, the role of plant VIH in cell wall-related rearrangements has

been established (Shukla et al. 2021). The participation of genes encoding VIP1/VIH proteins

in cell wall rearrangement in both yeast and plants could reflect the conserved function of the

proteins across species (Pöhlmann et al. 2014; Shukla et al. 2021).

Table 1: MS parameters for MRM transitions

Molecular name	Precursor Ion	Product Ion	Collision Energy (V)	Cell Accelerator Voltage
$[^{13}C_6]$ InsP$_5$	292	504.9	9	3
InsP$_5$	289	498.9	9	3
$[^{13}C_6]$ InsP$_6$	331.9	79.1	53	4
InsP$_6$	328.9	79.1	53	4
$[^{13}C_6]$ InsP$_7$	371.9	322.9	9	3
$[^{18}O_2]$ 4-InsP$_7$	370.9	319.9	9	3
InsP$_7$	368.9	319.9	9	3
$1,5[^{13}C_6]$ InsP$_8$	411.9	362.8	9	1
$1,5$InsP$_8$	408.9	359.8	9	1

543
544
545 Polarity: Negative, Dwell: 80, Fragmentor (V): 166

Table 2: Intrinsic disorder region (IDR) percentage, length and position in the human and plant encoded PPIP5K-like proteins. The efficient disorder prediction was made using ESpritz (ver1.3).

Name	% Disorder	Length of largest segment	Position	Position in AA	Total AA
HsIP6K1	29.58	167	After 8th	1107 to 1274	1433
HsIP6K2	29.92	155	After 9th	873 to 1028	1243
AtVIH1	13.34	65	After 4th	792 to 857	1049
AtVIH2	12.40	69	After 4th	797 to 866	1056
OsVIH1	16.47	81	After 3rd	786 to 867	1043
OsVIH2	10.54	30	After 3rd	648 to 678	758
TaVIH2-3B	12.93	37	After 5th	815 to 837	1036
TaVIH1-4D	13.65	56	After 5th	797 to 853	1047

553 **Table 3: Interacting partners of wheat VIH2 proteins.** The tables indicate the gene IDs
554 with their annotated function based on homology. The numbers indicate the frequency of the
555 occurrence of the clone as an interacting partner from the library.
556

S.No	Gene Function	Known functional in Planta	New TRIAE_ ID	Number of Occurrences
1.	Fasciclin-like protein FLA7	Cell wall biosynthesis, stress response,	TraesCS2D02G172900.1	9
2.	Glycosyl-transferases (GT1)	Cell wall biosynthesis, stress response, cell wall cementing, transfer sugar moieties	TraesCS6B02G339100.1	6
3.	Xylan arabinosyl transferase (Xat1)	Cell wall biosynthesis, stress response, cell wall cementing	TraesCS6A02G309400.1	5
4.	Glycine-rich cell wall structural protein-like	Cell wall biosynthesis, stress response	TraesCS2B02G541900	5
5.	NADH-ubiquinone reductase complex 1 MLRQ subunit;	Uncharacterized protein	TraesCS7A02G238600	3
6.	Hypothetical protein TRIUR3_12806	Auxin efflux transporter that acts as a negative regulator of light signalling to promote hypocotyl elongation	TraesCS4A02G016700	3
7.	ABC	Metal transporter involved	TraesCS5A02G392600	4

	transporter B family member	in the sequestration of aluminium into vacuoles		
8.	**Short-chain dehydrogenase reductase**	Glucose signalling and abscisic acid biosynthesis and functions	TraesCS2B02G116700	3
9.	*Ethylene-responsive element binding protein 1 (EREB1)*	Increases tolerance to abiotic stress	TraesCS7B02G062200	4
10	**W.W. domain-binding protein (containing PPG PPP motif)**	Modules mediate protein-protein interactions by recognizing proline-rich peptide motifs and phosphorylated serine/threonine-proline sites.	TraesCS7B02G062200	3

557
558
559 **Reference:**
560

Azevedo C, Burton A, Ruiz-Mateos E, et al (2009) Inositol pyrophosphate mediated

pyrophosphorylation of AP3B1 regulates HIV-1 Gag release. Proc Natl Acad Sci

106:21161–21166. https://doi.org/10.1073/pnas.0909176106

Bhandari R, Hamid A, Ladke JS, et al (2024) Interaction with IP6K1 supports

pyrophosphorylation of substrate proteins by the inositol pyrophosphate 5-InsP7. Biosci

Rep BSR20240792. https://doi.org/10.1042/BSR20240792

Bhati K, Aggarwal S, Sharma S, et al (2014) Differential expression of structural genes for

the late phase of phytic acid biosynthesis in developing seeds of wheat (*Triticum*

aestivum L.). Plant Science 224: 74-85

Cagnola JI, Dumont de Chassart GJ, Ibarra SE, et al (2018) Reduced expression of selected

FASCICLIN-LIKE ARABINOGALACTAN PROTEIN genes associates with the

abortion of kernels in field crops of Zea mays (maize) and of Arabidopsis seeds. Plant

Cell Environ 41:661–674. https://doi.org/https://doi.org/10.1111/pce.13136

Chabert V, Kim G-D, Qiu D, et al (2023) Inositol pyrophosphate dynamics reveals control of

575   the yeast phosphate starvation program through 1,5-IP(8) and the SPX domain of

576   Pho81. Elife 12:. https://doi.org/10.7554/eLife.87956

577   Desai M, Rangarajan P, Donahue JL, et al (2014) Two inositol hexakisphosphate kinases

578   drive inositol pyrophosphate synthesis in plants. Plant J 80:642–653.

579   https://doi.org/10.1111/tpj.12669

580   Dong J, Ma G, Sui L, et al (2019) Inositol Pyrophosphate InsP8 Acts as an Intracellular

581   Phosphate Signal in Arabidopsis. Mol Plant 12:1463–1473.

582   https://doi.org/10.1016/j.molp.2019.08.002

583   Eisenbeis VB, Qiu D, Gorka O, et al (2023) β-lapachone regulates mammalian inositol

584   pyrophosphate levels in an NQO1- and oxygen-dependent manner. Proc Natl Acad Sci

585   120:e2306868120. https://doi.org/10.1073/pnas.2306868120

586   Fridy PC, Otto JC, Dollins DE, York JD (2007) Cloning and Characterization of Two Human

587   VIP1-like Inositol Hexakisphosphate and Diphosphoinositol Pentakisphosphate Kinases.

588   J Biol Chem 282:30754–30762. https://doi.org/10.1074/jbc.m704656200

589   Gokhale NA (2013) Membrane phosphoinositides and protein–membrane interactions.

590   Amino Acids 45:751–754. https://doi.org/10.1007/s00726-013-1512-2

591   Harmel RK, Puschmann R, Nguyen Trung M, et al (2019) Harnessing (13)C-labeled myo-

592   inositol to interrogate inositol phosphate messengers by NMR. Chem Sci 10:5267–5274.

593   https://doi.org/10.1039/c9sc00151d

594   He Y, Wu L, Liu X, et al (2019) Yeast two-hybrid screening for proteins that interact with

595   PFT in wheat. Sci Rep 9:15521. https://doi.org/10.1038/s41598-019-52030-x

596   Jung J-Y, Ried MK, Hothorn M, Poirier Y (2018) Control of plant phosphate homeostasis by

597   inositol pyrophosphates and the SPX domain. Curr Opin Biotechnol 49:156–162.

598   https://doi.org/https://doi.org/10.1016/j.copbio.2017.08.012

599   Laha D, Johnen P, Azevedo C, et al (2015) VIH2 Regulates the Synthesis of Inositol

600   Pyrophosphate InsP8 and Jasmonate-Dependent Defenses in Arabidopsis. Plant Cell

601   27:1082–1097. https://doi.org/10.1105/tpc.114.135160

602   Laha D, Parvin N, Dynowski M, et al (2016) Inositol Polyphosphate Binding Specificity of

603   the Jasmonate Receptor Complex . Plant Physiol 171:2364–2370.

604   https://doi.org/10.1104/pp.16.00694

605   Laha NP, Giehl RFH, Riemer E, et al (2022) INOSITOL (1,3,4) TRIPHOSPHATE 5/6

606   KINASE1-dependent inositol polyphosphates regulate auxin responses in Arabidopsis.

607   Plant Physiol 190:2722–2738. https://doi.org/10.1093/plphys/kiac425

Lee Y-S, Mulugu S, York JD, O'Shea EK (2007) Regulation of a cyclin-CDK-CDK inhibitor complex by inositol pyrophosphates. Science 316:109–112. https://doi.org/10.1126/science.1139080

Liu G, Riemer E, Schneider R, et al (2023) The phytase RipBL1 enables the assignment of a specific inositol phosphate isomer as a structural component of human kidney stones. RSC Chem Biol 4:300–309. https://doi.org/10.1039/d2cb00235c

Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. Methods 25:402–408. https://doi.org/10.1006/meth.2001.1262

Ma Y, MacMillan CP, de Vries L, et al (2022) FLA11 and FLA12 glycoproteins fine-tune stem secondary wall properties in response to mechanical stresses. New Phytol 233:1750–1767. https://doi.org/10.1111/nph.17898

Machkalyan G, Trieu P, Pétrin D, et al (2016) PPIP5K1 interacts with the exocyst complex through a C-terminal intrinsically disordered domain and regulates cell motility. Cell Signal 28:. https://doi.org/10.1016/j.cellsig.2016.02.002

Meena V, Kaur G, Joon R, et al (2024) Transcriptome and biochemical analysis in hexaploid wheat with contrasting tolerance to iron deficiency pinpoints multi-layered molecular process. Plant Physiol Biochem 207:108336. https://doi.org/https://doi.org/10.1016/j.plaphy.2024.108336

Mulugu S, Bai W, Fridy P, et al (2007) A Conserved Family of Enzymes That Phosphorylate Inositol Hexakisphosphate. Science 316:106–109. https://doi.org/10.1126/science.1139099

Norman KL, Shively CA, De La Rocha AJ, et al (2018) Inositol polyphosphates regulate and predict yeast pseudohyphal growth phenotypes. PLoS Genet 14:e1007493–e1007493. https://doi.org/10.1371/journal.pgen.1007493

Pipercevic J, Kohl B, Gerasimaite R, et al (2023) Inositol pyrophosphates activate the vacuolar transport chaperone complex in yeast by disrupting a homotypic SPX domain interaction. Nat Commun 14:2645. https://doi.org/10.1038/s41467-023-38315-w

Pöhlmann J, Fleig U (2010) Asp1, a Conserved 1/3 Inositol Polyphosphate Kinase, Regulates the Dimorphic Switch in Schizosaccharomyces pombe. Mol Cell Biol 30:4535. https://doi.org/10.1128/MCB.00472-10

Pöhlmann J, Risse C, Seidel C, et al (2014) The Vip1 inositol polyphosphate kinase family regulates polarized growth and modulates the microtubule cytoskeleton in fungi. PLoS Genet 10:e1004586–e1004586. https://doi.org/10.1371/journal.pgen.1004586

Pollard TD, Cooper JA (2009) Actin, a central player in cell shape and movement. Science 326:1208–1212. https://doi.org/10.1126/science.1175862

Qiu D, Gu C, Liu G, et al (2023) Capillary electrophoresis mass spectrometry identifies new isomers of inositol pyrophosphates in mammalian tissues. Chem Sci 14:658–667. https://doi.org/10.1039/d2sc05147h

Qiu D, Wilson MS, Eisenbeis VB, et al (2020) Analysis of inositol phosphate metabolism by capillary electrophoresis electrospray ionization mass spectrometry. Nat Commun 11:6035. https://doi.org/10.1038/s41467-020-19928-x

Rao F, Cha J, Xu J, et al (2014) Inositol pyrophosphates mediate the DNA-PK/ATM-p53 cell death pathway by regulating CK2 phosphorylation of Tti1/Tel2. Mol Cell 54:119–132. https://doi.org/10.1016/j.molcel.2014.02.020

Ried MK, Wild R, Zhu J, et al (2021) Inositol pyrophosphates promote the interaction of SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant phosphate homeostasis. Nat Commun 12:384. https://doi.org/10.1038/s41467-020-20681-4

Riemer E, Qiu D, Laha D, et al (2021) ITPK1 is an InsP6/ADP phosphotransferase that controls phosphate signaling in Arabidopsis. Mol Plant 14:1864–1880. https://doi.org/https://doi.org/10.1016/j.molp.2021.07.011

Rogers EM, Allred SC, Peifer M (2021a) Abelson kinase's intrinsically disordered region plays essential roles in protein function and protein stability. Cell Commun Signal 19:27. https://doi.org/10.1186/s12964-020-00703-w

Rogers EM, Allred SC, Peifer M (2021b) Abelson kinase's intrinsically disordered region plays essential roles in protein function and protein stability. Cell Commun Signal 19:27. https://doi.org/10.1186/s12964-020-00703-w

Saiardi A (2004) Phosphorylation of Proteins by Inositol Pyrophosphates. Science (80-) 306:2101–2105. https://doi.org/10.1126/science.1103344

Saiardi A, Erdjument-Bromage H, Snowman AM, et al (1999) Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of higher inositol polyphosphate kinases. Curr Biol 9:1323–1326. https://doi.org/https://doi.org/10.1016/S0960-9822(00)80055-X

Saiardi A, Qiu D, Eisenbeis VB, Jessen HJ (2021) Absolute Quantitation of Inositol Pyrophosphates by Capillary Electrophoresis Electrospray Ionization Mass Spectrometry. JoVE e62847. https://doi.org/doi:10.3791/62847

Sanchez AM, Garg A, Shuman S, Schwer B (2019) Inositol pyrophosphates impact

676 phosphate homeostasis via modulation of RNA 3' processing and transcription

677 termination. Nucleic Acids Res 47:8452–8469. https://doi.org/10.1093/nar/gkz567

678 Shears SB (2015) Inositol pyrophosphates: why so many phosphates? Adv Biol Regul

679 57:203–216. https://doi.org/10.1016/j.jbior.2014.09.015

680 Shukla A, Kaur M, Kanwar S, et al (2021) Wheat inositol pyrophosphate kinase TaVIH2-3B

681 modulates cell-wall composition and drought tolerance in Arabidopsis. BMC Biol

682 19:261. https://doi.org/10.1186/s12915-021-01198-8

683 Thota SG, Bhandari R (2015) The emerging roles of inositol pyrophosphates in eukaryotic

684 cell physiology. J Biosci 40:593–605. https://doi.org/10.1007/s12038-015-9549-x

685 Ward JJ, Sodhi JS, McGuffin LJ, et al (2004) Prediction and Functional Analysis of Native

686 Disorder in Proteins from the Three Kingdoms of Life. J Mol Biol 337:635–645.

687 https://doi.org/https://doi.org/10.1016/j.jmb.2004.02.002

688 Wilson MSC, Saiardi A (2017) Importance of Radioactive Labelling to Elucidate Inositol

689 Polyphosphate Signalling. Top Curr Chem 375:14. https://doi.org/10.1007/s41061-016-

690 0099-y

691 Wright PE, Dyson HJ (2015) Intrinsically disordered proteins in cellular signalling and

692 regulation. Nat Rev Mol Cell Biol 16:18–29. https://doi.org/10.1038/nrm3920

693 Wu X, Lai Y, Lv L, et al (2020) Fasciclin-like arabinogalactan gene family in Nicotiana

694 benthamiana: genome-wide identification, classification and expression in response to

695 pathogens. BMC Plant Biol 20:305. https://doi.org/10.1186/s12870-020-02501-5

696 Yao K, Yao Y, Ding Z, et al (2023) Characterization of the FLA Gene Family in Tomato

697 (Solanum lycopersicum L.) and the Expression Analysis of SlFLAs in Response to

698 Hormone and Abiotic Stresses. Int J Mol Sci 24:. https://doi.org/10.3390/ijms242216063

699 Zang L, Zheng T, Chu Y, et al (2015) Genome-Wide Analysis of the Fasciclin-Like

700 Arabinogalactan Protein Gene Family Reveals Differential Expression Patterns,

701 Localization, and Salt Stress Response in Populus   . Front. Plant Sci.   6

702 Zhou J, Hu Q, Xiao X, et al (2021) Mechanism of phosphate sensing and signaling revealed

703 by rice SPX1-PHR2 complex structure. Nat Commun 12:7040.

704 https://doi.org/10.1038/s41467-021-27391-5

705 Zhu J, Lau K, Puschmann R, et al (2019) Two bifunctional inositol pyrophosphate

706 kinases/phosphatases control plant phosphate homeostasis. Elife 8:e43582.

707 https://doi.org/10.7554/eLife.43582

708 Zsolt S, Assegid G, Cristina A, Adolfo S (2011) Influence of Inositol Pyrophosphates on

709 Cellular Energy Dynamics. Science (80-) 334:802–805.

710   https://doi.org/10.1126/science.1211908

Figure 1

Figure 2

(A)

-ive control (Lam+T)
+ive control (p53+SV40)
TaVIH1+FLA7
TaVIH2+FLA7

SD(-LT) (X-alpha-Gal)

SD(-HLT) (+AbA)

(B)

HA-FLA7

cMYC-TaVIH2
HA-FLA7
Y187
cMYC-TaVIH2Y187: HA-FLA7

IP with c-MYC

IB::Anti-MYC

IB::Anti-HA

kDa
70
55
35

kDa
250
130
100
70

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8