**Oxytocin receptor controls distinct components of pair bonding and development in prairie voles**

Ruchira Sharma[1,2,3,4†], Kristen M. Berendzen[1,2,3,4†], Amanda Everitt[1,3,10,11], Belinda Wang[1,3], Gina Williams[1,2,3,4,5], Shuyu Wang[1,2,3,4], Kara Quine[1,2,3,4], Rose D. Larios[1,2,3,4,5], Kimberly L. P. Long[1,2,3,4], Nerissa Hoglen[1,2,3,4,5], Bibi Alika Sulaman[1,2,3,4], Marie C. Heath[1,2,3,4], Michael Sherman[1,2,3,4], Matthew Kinkel[1,2,3,4,], Angela Cai[1,2,3,4,], Denis Galo[1,2,3,4,], Lizandro Chan Caamal[1,2,3,4,], Nastacia L. Goodwin[1,2,3,4], Annaliese Beery[6,7], Karen L. Bales[8,9], Katherine S. Pollard[10,11,12], Arthur Jeremy Willsey[1,13], Devanand S. Manoli[1,2,3,4,*]

[1]Department of Psychiatry and Behavioral Sciences

[2]Center for Integrative Neuroscience

[3]Weill Institute for Neurosciences

[4]Kavli Institute for Fundamental Neuroscience

[5]Neurosciences Graduate Program, University of California, San Francisco; San Francisco, CA 95158, USA

[6]Department of Integrative Biology

[7]Department of Neuroscience, UC Berkeley, Berkeley, CA, United States

[8]Department of Psychology

[9]Department of Neurobiology, Physiology, and Behavior, University of California, Davis; Davis, CA 95616, USA

10Gladstone Institutes, San Francisco, CA, USA

11Department of Epidemiology & Biostatistics, University of California, San Francisco,

San Francisco, CA, USA

12Chan Zuckerberg Biohub, San Francisco, CA, USA

13Calico Labs, San Francisco, CA, USA 94080

* Corresponding authors. Devanand.manoli@ucsf.edu

 Lead Contact: Devanand.manoli@ucsf.edu

†These authors contributed equally to this work

**Summary:**

Oxytocin receptor (Oxtr) signaling influences complex social behaviors in diverse species, including social monogamy in prairie voles. How Oxtr regulates specific components of social attachment behaviors and the neural mechanisms mediating them remains unknown. Here, we examine prairie voles lacking Oxtr and demonstrate that pair bonding comprises distinct behavioral modules: the preference for a bonded partner, and the rejection of novel potential mates. Our longitudinal study of social attachment shows that Oxtr sex-specifically influences early interactions between novel partners facilitating the formation of partner preference. Additionally, Oxtr suppresses promiscuity towards novel potential mates following pair bonding, contributing to rejection. Oxtr function regulates coordinated patterns of gene expression in regions implicated in attachment behaviors and regulates the expression of oxytocin in the paraventricular nucleus of the hypothalamus, a principal source of oxytocin. Thus, Oxtr controls genetically separable components of pair bonding behaviors and coordinates development of the neural substrates of attachment.

**Keywords:**

Attachment behaviors, prairie vole, pair-bonding, oxytocin receptor, partner preference, monogamy, promiscuity, oxytocin signaling, paraventricular nucleus of the hypothalamus, nucleus accumbens

## Introduction

Social attachments, such as the bonds between parents and children, family and kin, friends, and the long-term relationships between mates and romantic partners form the basis of interpersonal and group dynamics within societies[1–7]. Various selective pressures have given rise to diverse patterns of complex social organization involving long-term relationships between members of a species, across mammals and other taxa, one of the most robust and intriguing of which is the enduring bond between mates[2,4,8–13]. In mammals, ~4% of species demonstrate social monogamy, forming persistent pair bonds between mating partners[9,14–16]. Comparative studies between closely related species that display distinct patterns of these social behaviors, including social monogamy, reveal correlated species-specific differences in the nonapeptide hormones oxytocin (Oxt) and arginine vasopressin (Avp), or their homologs, suggesting that they serve as critical modulators of affiliative behaviors and pair bonding[9–11,16–20]. Despite the central role of social attachment in human behavior and a rich history of investigation into the cognitive processes underlying social cognition and affiliative behaviors, it has been difficult to probe the neural mechanisms underlying these fascinating behaviors at the cellular, molecular, and genetic level, as most species that serve as model organisms for such studies in the laboratory do not display long-term social attachment as adults [6,21–27].

Prairie voles (*Microtus ochrogaster*) display long-term social attachment such that mating partners show an enduring pair bond and social monogamy[13,28–31]. Like many complex innate behaviors, pair bonding consists of distinct modules: the development of

increased prosocial behaviors with partners over strangers (partner preference), and

increased agonistic behaviors towards novel opposite sex potential mates (stranger

rejection)[13,30,32–37]. As in humans and other species that show affiliative social behaviors,

disruption of pair bonds results in increased anxiety-like behaviors and markers of

stress, supporting integrated neural and physiologic mechanisms that facilitate the

preservation of such attachments[38–40]. Pioneering work in prairie voles identified Oxt

and Avp as critical mediators of pair bonding[41,42]. Interspecific variations in the patterns

of expression of the oxytocin receptor (Oxtr) and the vasopressin 1a receptor (V1ar)

correlate with the potential for pair bonding within closely related vole species[43–49].

Consistently, pharmacologic inhibition of Oxt and Avp signaling via their respective

cognate receptors disrupts pair bonding, while exogenous administration of these

hormones promotes pair bonding without mating[28,50,51].

Given the central role of Oxt signaling in affiliative behaviors, we tested the genetic

requirement for signaling via Oxtr for pair bonding and social behaviors in prairie voles.

We generated prairie voles lacking Oxtr and unexpectedly observed that these animals

continue to display a strong preference for partners following mating and cohabitation[52].

To gain deeper insight into the modulation of attachment and social behaviors by Oxtr

function, we developed a series of behavioral assays to probe distinct aspects of pair

bonding and associated social behaviors. To interrogate the molecular mechanisms

mediating social attachment, we profiled patterns of gene expression in the nucleus

accumbens (NAc), a key site of species differences associated with social monogamy

that is dramatically enriched for Oxt binding in prairie voles[45,53,54]. We find that Oxtr

controls the timing of pair bonding in prairie voles and promiscuity independent of

partner preference formation. Oxtr sex-specifically controls reciprocal social behaviors,

such that naïve wildtype (WT) males preferentially interact with wildtype females over

those lacking Oxtr, while wildtype females show no such bias towards WT males. Oxtr

function regulates patterns of gene expression in the NAc, as well as other regions of

the vole brain implicated in social and attachment behaviors, in distinct ways in each

sex. Finally, we find that Oxtr function regulates the development of Oxt and Avp

expressing neurons in the paraventricular nucleus, a principal source of these

neuropeptides. Our studies reveal genetically separable components of pair bonding

and suggest that Oxtr signaling sex-specifically participates in the development of the

circuits underlying attachment behaviors.

**Results**

**Oxtr function facilitates the formation of partner preference**

We previously found that Oxtr is not required for the display of partner preference

following 1 week of cohabitation[52]. However, acute modulation of Oxtr signaling can

alter patterns of social behaviors between individuals[29,50,55]. These observations

suggest that changes in Oxtr function may influence early social interactions that

contribute to the trajectory of pair bond formation and/or specific aspects of attachment

behaviors following the formation of partner preference. We first wished to determine

whether Oxtr function facilitated pair bond formation by examining the display of partner

preference following shorter periods of cohabitation. Consistent with previous studies

showing that the cohabitation time required for the display of partner preference differs

between males and females, we find that WT females demonstrate partner preference

after only 6 hours of cohousing with a WT male partner, while WT males require 5 days

following introduction to a potential WT mate (Fig. S1A-C)[56]. We therefore cohoused WT

or Oxtr$^{1-/-}$ females with WT males for 6 hours, and WT or Oxtr$^{1-/-}$ males with WT females

for 5 days following introduction (short cohabitation), or in these combinations for 1 or 7

days (long cohabitation) for females or males, respectively (Fig. 1A, Fig. S1A).

Both males and females lacking Oxtr show an absence of partner preference following

short cohabitation with WT partners (Fig. 1B). Oxtr$^{1-/-}$ females, but not males, showed a

decrease in the fraction of huddling time with their partner and an increase in the

fraction of huddling time with a stranger compared to WT controls (Fig. 1C-D). Oxtr$^{1-/-}$

males present an absence of partner preference (Fig. 1B), displayed due to both

decreased time with their cohoused partner and increased time with strangers (Fig. 1C-

D). Consistent with our previous studies, we find no difference between WT and Oxtr$^{1-/-}$

females when the pairs are cohoused for the longer duration (Fig. S1B-F).

To further understand the dynamics of social interactions that contribute to the

difference in huddling behavior between WT animals vs their siblings lacking Oxtr

following short cohabitation, we quantified the cumulative time subjects spent huddling

with their cohoused partners or strangers over the course of the assay. Oxtr$^{1-/-}$ females

show a significant decrease in the amount of time spent huddling with partners in the

last 30 mins of the assay (150-180mins) (Fig. 1E). We also observe a significant

increase in huddling time of Oxtr$^{1-/-}$ males with strangers during the middle periods (75 –

130mins) (Fig. 1F), suggesting that they engage in more early prosocial behavior with

strangers than WT males. Consistent with our previous work, we find no difference

between cumulative huddling times with either partner or stranger between WTs and

Oxtr$^{1-/-}$ animals of both sexes after long cohabitation periods (Fig. 1G-H)[45]. These

observations demonstrate that Oxtr function changes the temporal dynamics of social

interactions with both familiar and unfamiliar animal in a sex-specific manner. Taken

together, our findings suggest that Oxtr function acts early in the process of pair bonding

to facilitate the formation of the partner preference.

**Oxtr suppresses promiscuous behaviors in a state dependent manner in females**

Oxtr influences the dynamics of interactions with partners and strangers differently

between the sexes. We therefore wished to examine the detailed patterns of these

interactions separately in females and males (Fig. 2A). We examined the distribution of

huddles by WT vs Oxtr$^{1-/-}$ females following short cohabitation as a measure of prosocial

behavior displayed by these animals. We find that the top quartile of huddle durations

(long huddles) from each subject animal make up an average of ~90% of the total

huddle duration, functionally separating these interactions from incidental side-by-side

contact that may not constitute affiliative huddling (Fig. S2A Left). We classified long

huddles as partner huddles or stranger huddles and discovered that, consistent with the

overall time spent, the median huddle duration carried out by Oxtr$^{1-/-}$ females is

significantly shorter with partners and significantly longer with strangers, compared to

WT females (Fig. 2B). The longest huddle displayed with partner is significantly longer

for WT females compared to Oxtr$^{1-/-}$ females though we find no difference in the length

of the overall longest huddle. This is due to some Oxtr$^{1-/-}$ females conducting their

longest huddle with the stranger instead (Fig. S2B). Oxtr[1-/-] females additionally

executed a larger number of huddles only with the stranger, but not the partner,

compared to WT controls (Fig. 2C). Taken together, these results suggest that Oxtr

function in females both facilitates prosocial behavior with partners and suppresses

such behavior with strangers during the early stages of pair bond formation.

We next wished to determine whether the loss of Oxtr function influences patterns of

prosocial behaviors after the display of partner preference. We therefore examined

patterns of huddling, in particular the duration and frequency of these interactions,

following long cohabitation, a time point at which both WT and Oxtr[1-/-] animals display

robust preference for partners over strangers[52] (Fig. S1D). As with short cohabitation,

the top quartile of huddles (by duration) for each subject animal following long

cohabitation, described an average of 90% of total huddle time (Fig. S2A right).

Following long cohabitation, preference and huddling behavior in Oxtr[1-/-] females is

indistinguishable from their WT siblings (Fig. 2D-E). Comparing changes in the

frequency of top quartile huddles displayed by Oxtr[1-/-] females after short vs long

cohabitation, we find that both WT and Oxtr[1-/-] females show decreased huddle

frequency with stranger following long cohabitation (Fig. S2C-D). On comparing

durations of long huddles, we find that Oxtr[1-/-] females significantly increase their

median huddle duration with partner post long cohousing (Fig. 2F-G). Thus, loss of Oxtr

function in females reduces the duration of early prosocial interactions (short

cohabitation) with partners and strangers while Oxtr-independent mechanisms

eventually promote WT patterns of affiliative huddling and partner preference following

longer periods of cohabitation.

Our results suggest that Oxtr function influences specific aspects of early pair bonding

and attachment behaviors. We next wished to dissociate the influence of Oxtr on the

formation and display of partner preference from the display of agonistic rejection

behaviors towards novel strangers following cohabitation. We therefore developed a

battery of assays to examine initial social interactions between mating partners at early

and later stages of pair bonding, as well as stranger rejection following cohabitation

(Fig.2A). We examined the behavior of subject females from each genotype with WT

males when naive animals were first introduced to each other (Introduction), mating

behaviors following estrus induction (Timed Mating), prosocial interactions following

reunification with their bonded mate (Partner Reunification), and agonistic rejection of

novel, opposite sex strangers (Stranger Rejection) (Fig. 2A see Methods). We find no

differences in social behaviors during Introduction, Timed Mating and Partner

Reunification (Fig. 2H-J). In contrast, during Stranger Rejection, Oxtr$^{1-/-}$ females show

more social behaviors than WT females towards novel naive males in the absence of

their partner (Fig. 2K). Taken together, these studies demonstrate that Oxtr function

regulates specific components of pair bonding in females in a state-dependent manner,

suppressing promiscuous social interactions with stranger males after cohabitation

sufficient for the display of partner preference.

**Oxtr function influences different prosocial behaviors across the stages of pair bonding in males**

Given the Oxtr-dependent difference in huddling behavior between males and females after short cohabitation (Fig. 1C-D), we wished to assess sex differences in the patterns of behaviors displayed by Oxtr[1-/-] animals and to further investigate Oxtr function in males (Fig. 3A). Examining long huddles (as above, Fig. 2), we find that such huddles by Oxtr[1-/-] males with their partners following short cohabitation are significantly shorter when compared to WT siblings, contributing to diminished partner preference early in pair bonding (Fig. 3B Left). Interestingly, while Oxtr[1-/-] males show no difference in the total amount of time spent huddling with strangers when compared to WT siblings (Fig. 1D), they display shorter and more frequent huddles towards them at this time (Fig. 3B, 3C Right). In males, Oxtr thus influences the pattern as well as amount of early affiliative behavior to distinct behavioral stimuli in different ways.

Oxtr[1-/-] males show no difference in time spent with either partners or strangers after long cohabitation (Fig. S1E-F). However, these males show a persistent reduction in the duration of their long huddles with strangers at this time point (Fig. 3D). Following long cohabitation, WT and Oxtr[1-/-] males show no difference in the duration of long huddles with partners and a reduction in the duration of long huddles with strangers (Fig. S3A-B). We find no difference in the frequency of long huddles between Oxtr[1-/-] and WT males after long cohabitation (Fig. 3E). Consistently, comparing Oxtr[1-/-] males' behavior following short vs long cohabitation reveals that the frequency with which Oxtr[1-/-] males huddle with strangers decreases following long cohabitation (Fig. 3F-G). Thus,

specifically in males, loss of Oxtr function does not change the overall time spent

huddling with strangers after short or long cohabitation but does cause a continuing

disruption in the patterns of affiliative behavior that comprises huddling with strangers.

Our observations suggest that the loss of Oxtr disrupts certain characteristics of

prosocial behaviors in males regardless of the time spent bonding with their partner. We

next wished to dissect whether such loss of function resulted in atypical affiliative or

agonistic behaviors across stages of bonding. We examined the behavior of Oxtr$^{1-/-}$

males and their WT siblings in the same series of paradigms described previously to

characterize the formation and maintenance of their pair bonds (Fig. 3A). Oxtr$^{1-/-}$ males

spend similar amounts of time as WT males engaging in prosocial behaviors with

partners regardless of bonding state (Fig. 3H-J, S3E). Similar to our observations in

Oxtr$^{1-/-}$ females, Oxtr$^{1-/-}$ males spend more time engaging in prosocial behaviors with

strangers in the absence of their partners (Fig. 3K).

In contrast to time spent conducting mating behaviors with WT female partners (Fig.

S3E), Oxtr$^{1-/-}$ males demonstrate increased frequency of mating attempts when these

females are in estrus (Fig. 3I). Examining all instances of mounting, we find that Oxtr$^{1-/-}$

males carry out shorter mounting bouts more frequently than WT males (Fig. S3F). To

determine whether the shorter bouts of mounting in Oxtr$^{1-/-}$ males influence the

receptivity of their partners, we compared the mount-to-intromission ratio for each

animal[57] and find that females are equally receptive to Oxtr$^{1-/-}$ males as their WT siblings

(Fig. S3G). Thus, consistent with our previous observations, Oxtr influences patterns,

but not the amount or success, of male mating behaviors, suggesting that such changes

in males' behaviors are equally able to elicit receptivity in females.

Compared to their WT siblings, naïve Oxtr[1-/-] males show decreased agonistic behaviors

towards an unfamiliar WT female upon first encountering her but display similar levels of

rejection of novel (unfamiliar) strangers upon bonding (Fig. S3C-D). Thus, Oxtr function

influences distinct components of males' attachment-related social behaviors towards

unfamiliar females at different points across pair-bonding, promoting agonistic displays

and lengthening mating bout durations when naïve, and suppressing promiscuous

prosocial behaviors upon bonding. Taken together our results suggest that Oxtr

influences patterns of social behaviors with future partners during the early phases of

pair-bonding differently in females and males, but upon bond formation, functions in

both sexes to suppress promiscuous affiliative behaviors towards strangers (Fig. S3H)

without affecting agonistic behaviors.

**Oxtr sex-specifically influences social interactions between potential mates**

Oxtr function influences patterns of social interactions between mates early during the

formation of a pair-bond (Short Cohabitation in both sexes, Introduction and Timed

Mating in males). We therefore investigated the effect of Oxtr loss on the choice made

by a potential mate. We developed a variation of the PPT three-chamber assay to

present a naive WT "chooser" with a choice between a naive opposite sex WT or naive

opposite sex Oxtr[1-/-] animal (Fig. 4A, Fig. S4A, see Methods). Strikingly, we find that

Oxtr influences the choice by a WT mate in a sex-specific manner. WT females choose

WT or Oxtr$^{1-/-}$ males with equal frequency (Fig. 4B). In contrast, WT male choosers

show significant preference for WT females compared to Oxtr$^{1-/-}$ females, particularly

during the middle of the assay (150-280 mins) (Fig. 4B-C, Fig. S4B). These

observations suggest that disruption of Oxtr function specifically in females may alter

their behavior such that WT males prefer WT future partners over such mutants.

While naïve WT females show no difference in overall social choice between naive WT

and Oxtr$^{1-/-}$ males (WT-Oxtr$^{1-/-}$, Fig. 4B), they may show distinct behavioral dynamics in

the presence of an Oxtr$^{1-/-}$ male. To assess this, we compared the WT-Oxtr$^{1-/-}$ condition

with a control group in which the naive female chooser is given access to two naive WT

males (WT-WT) (Fig. 4D). We designated the tethered animal with whom the subject

spent <u>M</u>ore <u>o</u>f <u>A</u>ssay <u>T</u>ime ("chosen") as the MOAT. Intriguingly, we found that the

presence of an Oxtr$^{1-/-}$ male (WT-Oxtr$^{1-/-}$) influenced the behavior of the WT female

chooser, such that they spent significantly more time with WT or Oxtr$^{1-/-}$ MOAT males

compared to the time spent with WT MOAT males from the WT-WT condition (Fig. 4E).

Consistent with this finding, naive WT female choosers break contact more frequently in

WT-WT conditions when compared to WT-Oxtr$^{1-/-}$ (Fig. 4F). Similarly, the divergence in

the amount of cumulative time spent with the MOAT for WT female choosers occurs

after 5 hours have elapsed (320-360 mins) suggesting that complex interactions with

males lacking Oxtr influence how females "choose" a male partner regardless of

whether they end up choosing a WT or Oxtr$^{1-/-}$ MOAT (Fig. 4G). In contrast, we find no

difference in the contact time or number of such transitions when comparing naive WT

male choosers with MOAT females of a WT-WT vs WT-Oxtr$^{1-/-}$ condition (Fig. 4E-F).

We next examined the pattern of a WT chooser's interactions with both chosen (MOAT)

and unchosen animals (Less of Assay Time, LOAT) over time using a sliding preference

index (see Methods). We observed periods of convergence when choosers from a

condition (WT-WT or WT-Oxtr[1-/-]) displayed significantly concurrent interactions with

their MOAT, indicated by the black bar above the trace. WT female choosers show a

longer and more consistent preference for their MOAT (regardless of genotype) in WT-

Oxtr[1-/-] vs WT-WT conditions, whereas WT male choosers show an earlier onset but

equivalent length of preference for MOATs in the WT-Oxtr[1-/-] condition compared with

the WT-WT condition (Fig. 4H). These observations suggest that interactions with males

lacking Oxtr influence the behavior of the WT females making a social choice, even

when these females interact more with and eventually choose a WT male. Taken

together, loss of Oxtr leads to differences in prosocial behaviors during initial

interactions with WT naive animals of the opposite sex that may later impact aspects of

pair bonding.

Finally, to determine if naive prairie voles lacking Oxtr display generalized changes to

their behaviors independent of attachment-associated social behavior, we examined

general pro-social, anxiety-related, and locomotor behaviors. With the exception of an

increase in locomotion displayed by Oxtr[1-/-] females when compared to WT females in

an open field paradigm, we found no differences between Oxtr[1-/-] and WT voles of either

sex (Fig. S4C).

We have examined social interactions during the initial contact between mates, the

early stages of pair bonding, and during social attachment behaviors following partner

preference formation. Consistent with our findings and previous work, Oxtr function is

not required for the eventual display of partner preference. Here, we demonstrate that

Oxtr function sex-specifically facilitates early social interactions that contribute to the

formation of partner preference, and controls promiscuity.

**Oxtr controls the molecular signature of pair bonding in the nucleus accumbens**

We previously observed that Oxtr function is not required for the formation of partner

preference and here demonstrate that it suppresses prosocial behaviors towards novel

strangers following pair bonding. These results suggest that distinct components of pair

bonding are differentially influenced by Oxtr function or compensated upon its loss. We

next characterized the molecular signature associated with pair bonding to determine

whether changes in gene expression following pair bond formation are controlled by

Oxtr. Comparative studies demonstrate that Oxt binding in multiple brain regions differs

significantly between promiscuous and monogamous rodent species. Oxt binding is

particularly enriched in the NAc in prairie voles, while vasopressin binding is largely

absent in this area[45,47]. Thus, analyzing gene expression in the NAc allows us to identify

bonding-related expression that is primarily due to Oxtr function and largely

independent of the direct consequences of AVP signaling. Additionally, given the well-

established role of the NAc in the association of reward or aversion with specific

sensory and environmental cues[58–60] this region is well poised to play a role in the

development of partner preference, even without Oxtr signaling, as well as stranger

rejection, which appears to be dependent upon Oxtr function.

We performed bulk RNA-sequencing on NAc tissue from WT and Oxtr$^{1-/-}$ animals of

both sexes that were either: group housed with same-sex peers and sexually naïve

(pre-pairing); or paired with WT opposite-sex partners for six days, using our timed

mating paradigm (post-pairing) (Fig. 5A). PCA showed robust separation of WT and

Oxtr$^{1-/-}$ samples (Fig. S5A). Differential expression (DE) analysis of pairing condition,

regardless of genotype or sex, identified 13 genes that show significant changes in

expression post-pairing (Fig. S5B), all of which showed decreased expression (Fig. 5B).

These genes include *Agt, Sgk1, Ptgds, Fosb, Pdk4*, and *Acsm5* (Fig. 5C) and, while

these genes have not previously been linked to shared gene regulatory or functional

pathways, the predicted protein-protein interaction network for the 13 DE genes shows

more connections than would be expected by chance ($p=2.47 \ e^{-05}$) (Fig. S5C).

Intriguingly, even in the limited set of genes with altered expression in the context of pair

bonding, changes are driven primarily by expression in WT females (Fig. 5C, S5D).

Subgroup analyses by genetic background and sex identified additional genes including

*Adora2a, Fkbp5*, and *Spred3* in WT females and collagen-related genes, including

*Col4a1* and *Col4a2*, in Oxtr$^{1-/-}$ females (Fig. 5D, S5D). Only one gene, *Sema3b,* is

significantly different in Oxtr$^{1-/-}$ males following pairing (Fig. 5D).

To further examine changes in patterns of gene expression in the NAc associated with

pair bonding, we performed weighted gene coexpression network analysis (WGCNA)[61]

and calculated three sets of modules (M) from different sample subsets (modules$^{WT}$,

modules$^{Mut}$, and modules$^{All}$) (Fig. S5F, S6D). Using the modules$^{WT}$ subset, we identified

15 coexpression modules (Fig. S5F, Table 2) representing genes that share highly

similar expression patterns within the NAc. We identified one module (Mpink$^{WT}$) that

significantly and specifically correlated with pre-pairing status in WT animals (Pearson R

correlation of 0.7, Bonferroni adjusted p-value ($p_{adj}$)=0.018) (Fig. 5E, S5E-F).  Genes

with high module membership for the Mpink$^{WT}$ are also those with high gene

significance for pairing condition supporting the module's association with behavior

condition (Fig. 5SG). Further, genes significantly differentially expressed with bonding

are significantly enriched in Mpink$^{WT}$ (Bonferroni $p_{adj}$=7.44 e$^{-08}$) and serve as hub genes

within this module (Fig. S5E, G). These results suggest that the Mpink$^{WT}$ module might

be driving, or under the influence of pathways driving, the pre-pairing to post-pairing

state change.

We asked whether any of the modules$^{WT}$ identified were regulated by Oxtr and thus lost

in Oxtr null animals. To address this, we assessed how well modules$^{WT}$ are preserved in

mutant samples. We calculated a $Z_{summary}$ measure that combines module density and

intramodular connectivity metrics into a composite statistic. We find that of the

modules$^{WT}$, Mpink$^{WT}$ had the lowest $Z_{summary}$ ($Z_{summary}$ = 2) in Oxtr$^{1-/-}$ samples (Fig. 5F),

indicating that dynamics of gene expression that correlate with bonding states are lost in

the absence of Oxtr. Finally, the module eigengene (ME) for Mpink$^{WT}$ when calculated

for all 30 samples and compared between pre- and post-pairing, was significant only for

WT females (Wilcoxon rank sum, Benjamini-Hochberg $p_{adj}$=0.03) and not for WT males

or Oxtr[1-/-] animals of either sex (Fig. 5G). These observations suggest there is sex-

specific, coordinated regulation of gene expression in the NAc following pairing of mates

that may be associated with pair bond associated behaviors.

**Loss of Oxtr sex-specifically disrupts patterns of gene expression associated**

**with neurodevelopmental disorders**

We next wished to determine how Oxtr signaling contributes to changes in gene

expression in the NAc associated with the naive state or the process of pair bonding

(Fig. 6A) by comparing the transcriptional profile of WT and Oxtr[1-/-] animals of each sex.

We observed significant genotype specific differences in gene expression in animals

from both sexes, with 466 genes showing differences between WT and Oxtr[1-/-] females,

270 between WT and Oxtr[1-/-] males, and 1,014 genes when samples from both sexes

were combined (using a $p_{adj}$<0.05 and absolute ($log_2$(FC)) > 0.25 as criteria). Of the

combined DE gene set, 370 genes are upregulated and 644 are downregulated (Fig.

6B, S6A-B). Genes with the greatest fold change across both sexes include *Slc2a3,*

*Kcna1, Rgs8, Cnksr2, Acvr*, and *Cnot6* (Fig. 6B). Genes specifically downregulated in

Oxtr[1-/-] females with $log_2$(FC) less than 1 include *Abcb11* (a gene recently associated

with treatment resistant schizophrenia[62]), *Ngfr, Pon3,* and *Shh*. While there was no

significant difference in Oxtr transcript levels by genotype ($p_{adj}$ = 0.753)—consistent with

minimal nonsense-mediated decay despite a premature stop codon—we have

previously shown that no functional protein is produced in Oxtr[1-/-] animals[52]. V1aR

showed a modest decrease in expression in Oxtr[1-/-] animals ($p_{adj}$=0.03) (Fig. S6C),

though this is likely influenced by variability in WT samples. WGCNA of all samples

identified two modules (Mbrown$^{All}$ and Mturquoise$^{All}$) that were significantly positively

correlated with the WT genotype compared to mutant (Pearson R = 0.55, $p_{adj}$ = 0.011 for

brown and Pearson R = 0.8, $p_{adj}$ = 6.6e-7 for turquoise) (Fig. S6D).  None of the

identified modules showed correlation with pairing condition (Fig. S6D). Our results thus

reveal a dramatic change in patterns of gene expression in the NAc with global loss of

Oxtr signaling, independent of bonding state.

Examination of the sets of genes whose expression is altered by the loss of Oxtr,

regardless of bonding state or sex, revealed several distinct cellular processes

regulated by Oxtr signaling. These processes comprise larger domains of cell function

including extracellular channel binding activity, axoneme assembly complex, axon glial

ensheathment development, and metal cation transport (Fig. 6C), as well as pathways

for G protein signaling and dopaminergic neurogenesis (Fig. S6E). When analyzed by

sex, more of the identified processes were enriched in female Oxtr gene sets than in

male ones (female: 19/62, male: 6/62). To further link differentially expressed genes to

behavioral traits, we asked whether genes altered by loss of Oxtr are selectively

enriched for high-confidence associations with autism spectrum disorders (ASD) and

other neurodevelopmental diseases (NDD) affecting social attachment in humans as

compared to epilepsy, a disorder that does not disrupt social attachment[63–66].

Consistently, we find that the genes upregulated with loss of Oxtr ($p_{adj}$<0.01, log$_2$(FC)>0)

are specifically enriched for ASD- and NDD-associations but not epilepsy associations

(Fig. 6D "Joint"). This enrichment is again primarily driven by genes upregulated in

females and not males (Fig. 6D). Thus, similar to the factors that influence gene

expression changes associated with pair bonding, many of the changes we identify in

the NAc resulting from loss of Oxtr are primarily driven by female-specific effects that

may play a conserved role in mediating social attachment in humans.

**Oxtr regulates coordinated gene expression across multiple neural populations**

**implicated in social attachment behaviors**

Given the expression of Oxtr in regions of the brain known to mediate social behavior in

voles and other species[67,68], we wished to determine if genes whose expression is

altered in the NAc by the loss of Oxtr were expressed not only in specific populations of

cells within this region, but in other regions of the prairie vole brain implicated in social

and attachment behaviors. We therefore examined the expression of 15 genes that

showed significant changes in expression in WT females when compared to those

lacking Oxtr using *in situ* hybridization (ISH) to both validate our RNA-seq findings as

well as examine the expression patterns of these genes. Several of these genes, such

as phosphodiesterase 10A (*Pde10A)* and diacylglycerol kinase beta (*Dgkb)*, show

robust expression throughout the striatum (data not shown), while genes such as the

calcitonin receptor (*Calcr*) and delta like non-canonical notch ligand 1 (*Dlk-1*) are

expressed in restricted populations of cells in the NAc (Fig. 6F-H, top panels). The

expression of these genes is significantly reduced in Oxtr$^{1-/-}$ females (Fig. 6E-H),

supporting the specificity and resolution of our RNA-seq findings.

In parallel to our examination of gene expression in the NAc, our ISH revealed that loss

of Oxtr in females altered the expression of the genes we examined in other regions of

the prairie vole brain. We examined the expression of these genes in the lateral septum

(LS), a region that has been implicated in agonistic and attachment behaviors in prairie

voles and other rodents and shows species dimorphic patterns of Oxt and Avp binding.

We observe that both *Calcr* and *Dlk-1* are decreased in the absence of Oxtr (Fig. 6G, H

middle panel) in the LS. We also find that expression of *Calcr* and *Dlk-1* is decreased in

the paraventricular nucleus of the hypothalamus (PVN), a principal source of Oxt and

Avp in the rodent brain which sends projections to the NAc (Fig. 6G-H bottom panels)[69–

71]. These findings suggest that Oxtr function regulates patterns of gene expression in

multiple regions of the brain implicated in social and attachment behaviors in a

coordinated and region-specific manner.

**Loss of Oxtr results in the decrease in density of Oxt and Avp expressing**

**neurons in the PVN**

We find here that loss of Oxtr results in coordinated changes in gene expression in

multiple neural populations, including the PVN. We therefore wished to determine if the

signatures of gene expression regulated by O̲xtr in the N̲Ac in p̲rairie voles (ONP set),

regardless of sex and bonding state, were enriched for patterns of gene expression

associated with distinct neural populations in other regions of the brain, particularly

those implicated in social and attachment behaviors. We compared the ONP gene set to

the molecular signatures of 33 hypothalamic subtypes previously identified by single-cell

RNA sequencing in mice[72]. Of the 33 clusters examined, the four with highest overlap

with the ONP gene set represent subsets of PVN neurons, in particular those that

express Oxt and AVP (Fig. 7A, clusters 43 and 26), as well as PVN neurons that

express *Dlk1* and *Calcr* (cluster 29), both of which show reduced expression in the NAc

and PVN in females lacking Oxtr (Fig. 6G-H bottom panel).

We next sought to determine if the overlap between Oxtr regulated genes and gene

expression patterns of neural populations in the PVN correlated with changes in the

expression of neuropeptides within PVN itself. We examined patterns of Oxt and Avp

expression using immunohistochemistry in naive adults and compared patterns of

expression between WT animals of both sexes and those lacking Oxtr (Fig. 7B, Fig.

S7A-B). We find that the density of both Oxt and Avp producing neurons is lower in

Oxtr$^{1-/-}$ animals (Fig. 7C-E). To determine if the loss of Oxt or Avp expression was

restricted to specific populations of PVN neurons, we examined the expression of these

peptides across the anterior-posterior axis of the PVN in both WT and Oxtr$^{1-/-}$ males and

females. We find that animals of both sexes show a preferential loss of Oxt and Avp

expressing neurons in the anterior PVN (Fig. 7F-G). Finally, we examined patterns of

Oxt and Avp over the course of development to determine when loss of Oxtr function

influences the development of their expression. We find that the density of Oxt neurons

at P0 (Fig. S7C-F) and both Oxt and Avp neurons at P21 is lower in Oxtr$^{1-/-}$ females

compared to age matched WT females (Fig. S7G-J). In contrast, we find that Oxt and

Avp neuron density loss occurs only in adult Oxtr$^{1-/-}$ males (Fig. 7D-G, Fig. S7B). These

observations suggest the coordinated, region-, and sex-specific influence of Oxtr

function on the development of neurons expressing Oxt (and Avp) within the PVN,

consistent with a trophic role for this pathway identified in mice[73–76].

**Discussion**

Robust reproductive behaviors are essential for survival. Accordingly, species have

evolved diverse strategies, including social monogamy, which mediate locating,

identifying, and attracting mates, facilitate reproduction, and promote the survival of

progeny[16,77]. Signaling via Oxtr influences a wide range of physiology and behavior

associated with social interactions and attachment between conspecifics, including pair

bonding between prairie vole mates[41,78]. Our recent findings that Oxtr is not genetically

required for the formation of partner preference in prairie voles suggests that, like other

genes and neuromodulatory pathways across diverse species, Oxtr likely functions in

parallel to multiple genetically specified pathways to facilitate pair bonding by

influencing specific behavioral modules that contribute to the complex behaviors that

comprise social attachment[33,35,79–85].

Here we demonstrate that loss of Oxtr in prairie voles influences the dynamics of social

interactions between sexually naïve animals and WT potential mates in distinct social

contexts in sex-specific ways. When given a choice, naive WT males display a

preference for WT females rather than for those lacking Oxtr, supporting the role of Oxtr

in controlling behavior during an early, sensitive period in bond formation[30]. The

presence of an Oxtr[1-/-] male alters levels and patterns of early prosocial behavior

exhibited by WT females, without ultimately affecting their choice of potential partner.

Taken together, these observations suggest that Oxtr may modulate social behaviors

that act early in a sensitive pair bond formation period, and via these behaviors or yet-

to-be-determined sensory cues such as pheromone profiles or patterns of vocal

communication, influences the reciprocal behavior of partners.

Animals lacking Oxtr take longer to display partner preference, and despite strong

partner preference following longer cohabitation, demonstrate promiscuous prosocial

behaviors towards novel potential mates, including huddling with strangers. These

observations are consistent with a state-dependent role of Oxtr function, similar to

findings in multiple species, including humans[86–88]. Oxtr function may thus promote

prosocial behavior in the naïve state to facilitate attachment with an unfamiliar potential

mate, but following pair bond formation, suppresses further prosocial interactions to new

individuals, or even promotes further separation from and rejection of unfamiliar

individuals or groups. Across species and outside of the context of pair bonding,

oxytocin signaling may play a conserved role in mediating other state-specific aversive

behavior, for example, defeat-induced social avoidance learning[89].

Our findings reveal that patterns of social behaviors contributing to 1) pair bond

formation, 2) partner preference, and 3) rejection of novel mates are genetically

separable components of pair bonding representing distinct behavioral modules

regulated by mechanisms that require Oxtr function, i.e. reciprocal behaviors and

stranger rejection, or those that can be displayed in its absence, i.e. partner preference.

Further work will allow us to determine the Oxtr-independent mechanisms that mediate

the closure of an early sensitive period and facilitate the display of partner preference[52].

Where Oxtr functions to control the switch from prosocial behavior towards a potential

mate in naïve animals to the suppression of such behaviors towards, and active

rejection of, opposite-sex conspecific strangers following bonding remains to be

determined.

Examination of the molecular signatures of pair bonding and the role of Oxtr in

regulating these changes in gene expression in the NAc reveals modest, Oxtr-

dependent changes in the expression of a specific network of genes in WT animals

following pair bonding[45,90]. Intriguingly, loss of Oxtr preferentially impacts patterns of

gene expression in the NAc in females. The NAc demonstrates little endogenous sex

steroid hormone receptor expression difference in gene expression in prairie voles and

other species[67,91–93]. Sex differences in the effects of Oxtr on gene expression this

region may therefore suggest that there are significant sex-differences in the inputs into

this region that act through Oxtr during development[94–99]. The larger impact of loss of

Oxtr in females is consistent with its more general conservation in mammals to regulate

aspects of female physiology such as lactation and prosocial behaviors or sensory cues

that promote interactions with males[100].

Genes with altered expression in the absence of Oxtr are enriched for genes associated

with neuropsychiatric disorders, in particular neurodevelopmental disorders and ASD.

Given the sex-biased enrichment of these genes, such developmental differences in the

impacts of Oxtr function may contribute to sex differences in the pathophysiology of

these disorders[101–103]. Accordingly, the molecular signatures of attachment in the NAc,

both dependent upon and independent of Oxtr function, are subtle relative to these sex-

biased differences in the role of Oxtr and possibly reside in other neural populations that

function in attachment.

Validation of changes in gene expression in the NAc resulting from loss of Oxtr

suggests coordinated regulation of these genes by Oxtr across multiple brain regions

implicated in social and attachment behaviors. Oxtr may be one of multiple factors that

organizes specific components of the circuits underlying these displays to control

discreet modules of behavior. Consistent with a trophic role for Oxtr function during

development, we find that the genes regulated by Oxtr within the NAc are enriched for

those that identify cell types in the PVN, a principal source of neuropeptides implicated

in pair bonding, which sends oxytocinergic projections to the NAc[68,70,104–106]. Loss of

Oxtr preferentially impacts Oxt and Avp expression in the anterior PVN, suggesting that

specific populations of these cells may be more dependent upon such trophic function

for their development, survival, or expression of these hormones. Whole brain analysis

will likely reveal if such neurons have distinct projections when compared to those that

remain in the absence of Oxtr, contributing to our understanding of how these circuits

and pathways regulate distinct aspects of social and attachment behaviors and

physiology[4,78,107].

Taken together, our behavioral and molecular studies reveal previously unappreciated

nuances in the control of social behavior and pair bonding by Oxtr function. They

uncover intriguing sex differences in the patterns of behavior influenced by loss of Oxtr

function. These observations suggest that circuits with sex-specific patterns of

development, or even function in ancestral species, may mechanistically converge in

the context of Oxtr function to generate displays of reciprocal interactions, pair bonding

and attachment behaviors in WT prairie voles that are similar between males and

females[108,109]. Manipulation of Oxtr expression at specific developmental timepoints and

in specific neural populations will help to elucidate how this pathway influences social

and attachment behaviors in the context of normal development to regulate distinct

components of pair bonding. Understanding the specific neural populations that express

Oxtr and other molecular pathways that contribute to pair bonding will allow us to

understand the mechanisms controlling the early experience-dependent, sensitive

period of bond formation. In parallel, examination of the activity in these populations

following manipulations of Oxtr function will allow us to determine the circuit

mechanisms by which similar but distinct sensory cues contribute to the identification of

partners or novel conspecifics and consequently, the display of dramatically different

patterns of social behaviors. The sum of these mechanisms represents the preference

for partners over strangers and reflects the formation and maintenance of pair bonds

between mates, affiliation between individuals, and the behavioral consequences of

identifying others as unfamiliar or outside of these bonds.

**Acknowledgments**

The authors would like to thank members of the Manoli lab for assistance, advice,

helpful discussion and comments on the manuscript. Additionally, we would like to thank

Sudhanshu Garg for his assistance in data analysis and help designing the analytic

pipeline. The authors received funding from National Institutes of Health grant

R01MH123513 (D.S.M.), National Science Foundation grant 1556974 (D.S.M.),

Burroughs Wellcome Fund 1015667 (D.S.M.), Whitehall Foundation grant 2018-08-83

(D.S.M.), One Mind Foundation A137726 (D.S.M.), National Institutes of Health grant

R25MH060482 (K.M.B.), AP Giannini Foundation Fellowship P0534952 (K.M.B.), Larry

L. Hillblom Foundation Fellowship 2020-A-023-FEL (K.M.B.), and National Institute of

Mental Health grant R01MH123178 (K.S.P.)

**Author contributions**

Conceptualization, R.S., K.M.B., K.L., and D.S.M.; methodology, A.B., A.J.W., D.S.M.,

K.L.B., K.M.B., K.S.P., and R.S.; software, K.Q., and S.W.; formal analysis, A.E., R.S.,

and K.M.B.; investigation, R.S., K.M.B., A.E., B.W., G.W., S.W., K.Q., R.D.L., K.L., N.H.,

B.A.S., M.C.H., M.S., M.K., A.C., D.G., L.C.C., N.L.G.; visualization, R.S., K.M.B., B.W.,

and A.E.; funding acquisition, D.S.M., K.S.P., and A.J.W; writing– original draft, R.S.,

and K.M.B.; writing-review & editing, R.S., K.M.B., D.S.M., N.H., K.S.P., A.E., and A.B.

**Declaration of interests**

The authors declare no competing interests.

**Star Methods**

**Key resources table**

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|
| **Chemicals, peptides, and recombinant proteins** | | |

| | | |
|---|---|---|
| DAPI | Sigma-Aldrich | Cat #D9542; CAS #28718-90-3 |
| **Experimental models: Organisms/strains** | | |
| Prairie vole: OxtR$^{1-/-}$ | Berendzen et al. | |
| **Oligonucleotides** | | |
| ISH probes | This paper | N/A |
| Oxtr$^{1-/-}$ genotyping oligos | This paper | N/A |
| **Antibodies** | | |
| Oxytocin Polyclonal | Invitrogen | Cat#PA5-26701; RRID |
| Anti-Neurophysin 2/NP-AVP | Millipore | Cat#MABN845; RRID |
| Cy™3 AffiniPure Donkey Anti-Rabbit IgG | Jackson ImmunoResearch | Cat #711-165-152; RRID |
| Alexa Flour 488 Donkey Anti Mouse IgG | Thermo Fisher Scientific | Cat #711-165-152; RRID |
| **Software and algorithms** | | |
| ImageJ | NIH | ; RRID: SCR_003070 |
| R (version 1.1.423) | R Core | https://www.r-project.org/ |
| GraphPad Prism (version | GraphPad | |

| 7.03) | Software | |
|---|---|---|
| Boris | Boris | www.boris.unito.it |
| MATLAB | MathWorks | https://www.mathworks.com/products.html<br><br>RRID: SCR_001622 |

**Resource availability**

**Lead contact**

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Devanand Manoli (devanand.manoli@ucsf.edu).

**Materials availability**

All unique reagents generated in this study are available from the lead contact upon

request.

**Data**

Transcriptomics data discussed in this publication are accessible through GEO Series

accession number GSE:TBD.

**Code**

All code used for RNA-seq data processing and differential expression analyses, as well

as post-processed expression data, can be found at

https://github.com/aseveritt/OXTR_prairie_voles

## EXPERIMENTAL MODEL AND SUBJECT DETAILS

### Animals

Subjects were laboratory-bred prairie voles (*Microtus ochrogaster*) which originated through systematic outbreeding of a wild stock captured near Champaign, Illinois. CRISPR/Cas9 editing was used to generate 2 independent null alleles, $Oxtr^{1-/-}$ and $Oxtr^{2-/-}$, both of which were validated using autoradiography to show loss of receptor function[52] (data not shown). $Oxtr^{1-/-}$ was backcrossed for >7 generations to mitigate any possible off-target effects and this line was used for all subsequent experiments. Sexually naïve male and female animals were group weaned at 21 ± 1 days and separated to group housing with same-sex siblings and age-matched same-sex non-siblings. Voles were maintained under a 14:10 h light-dark cycle in clear plastic cages (45 × 25 × 15 cm) with bedding, nesting material (nestlet), and a PVC hiding tube. Rooms were maintained at approximately 20°C, and food and water were available *ad libitum*.

Breeding pairs were established between two heterozygotes or a homozygous $Oxtr^{1-/-}$ mutant male and heterozygous female partner from a breeding line maintained in the lab. Voles were randomly assigned into experimental groups when they reached 7-9 weeks of age at the start of testing. This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals published by the National Research Council. The protocol was approved by the Institutional Animal Care and Use Committee at each respective institution.

**Genotyping**

The following primers were used to amplify a region where XcmI restriction for Oxtr[1] or

Bpu10I restriction for Oxtr[2] determined the genotype of all animals studied. XcmI cuts

the mutant allele in the case of Oxtr[1] and Bpu10I restricts the WT sequence for Oxtr[2].

(Forward: ACTGGAGCTTCGAGTTGGAC, Reverse: ATGCCCACCACTTGCAAGTA)

**METHOD DETAILS**

**Behavior**

***Triadic naive choice assay***

The linear three chamber partner preference apparatus with open top and 10 x 32 inch

walls[52] was modified to capture the interactions of the naive, freely moving WT subject

and the naive, tethered, opposite sex potential partner (either 2 WTs or a WT and an

Oxtr[1-/-]) at either end of the chamber. Behaviors were recorded for 6 hours using a top

view camera and the videos generated were analyzed using a contour detection

algorithm. Specifically, contours were detected on individual video frames using a

threshold-based strategy that took advantage of the contrast between dark-colored vole

fur coats and light-colored bedding and chamber apparatus. A standalone vole was

detected as a single contour, and two voles in physical contact were likewise detected

as a single contour. Consistent vole identities across video frames were determined

using the Hungarian algorithm. Since the assay began with one vole in each of the three

chambers, and since the subject vole remained freely mobile throughout, it was possible

to unambiguously assign each contour to a single vole identity (e.g., subject, right

chamber, or left chamber) or to that of a pair of voles (e.g., subject and right chamber

voles or subject and left chamber vole), resulting in continuous trajectories for each of

the three animals. Time in chamber was determined by the chamber location of the

subject vole in each frame, since the chamber location of the right chamber and left

chamber animals was fixed throughout the assay by physical tethering and any time the

contour of the subject animal merged with that of the tethered animal (right or left), the

subject was termed in contact with it. The algorithm also determined the boundaries of

the chambers and calculated chamber times. We used the chamber time data from

when both ends of the chamber contained tethered WTs (WT-WT condition) to ensure

that there was no bias in which side the choosers preferred.

***Bond formation assays***

Initial introduction of $Oxtr^{1-/-}$ and WT animals of either sex with an opposite sex WT

animal was filmed for 30 minutes. Observed behaviors included anogenital sniffing,

investigation (subject animal's nose contacts any non-genital region), duration of

stationary huddling or >50% side-by-side contact between animals, rearing, frequency

of aggressive behavior including lunges, receipt of aggression, and tussles, as well as

mating behavior including mounting, intromission, and ejaculation. These animals

remained in their home cage for 18 hours following pairing, after which, they were

separated by a clear, plastic barrier. The barrier was left in place for 24 hours and then

removed allowing the subject free access to the partner animal thereafter. A 30-minute

period directly following barrier removal was also filmed to assess mating behavior,

termed the timed mating assay.

Subjects remained cohoused with the opposite-sex, wildtype partner for durations as

indicated in the text. The partner preference test (PPT), as previously published[52], was

used to assess pair bond formation. Subject, partner, and an opposite sex stranger are

placed in a three-chamber apparatus with open top and 10 x 32-inch walls. The partner

and stranger animals are tethered on either end of the three-chamber arena and the

subject allowed free access over a period of 3 hours. Behaviors are recorded from a top

view camera capturing the entire apparatus. Videos were scored using BORIS[110] post-

test by validated scorers blind to condition. Observed behaviors included location (i.e.,

duration of time in partner, stranger, and neutral chambers), duration of stationary

huddling or >50% side-by-side contact between the partner and stranger animals, and

frequency of aggressive behavior (i.e., lunges). Of note, analysis of partner preference

between wild type siblings from the $Oxtr^{1-/-}$ background and wild type animals from the

breeding colony showed no difference in time spent huddling with partner vs stranger

animals.

***Bond maintenance assays***

Four days after the timed mating assay, female animals underwent a separation-

reunification assay. Females remained in their home cage while their male partners

were placed in a separate cage in another room for one hour. The males were then

reintroduced into the home cage and behavior was video recorded for 30 minutes. Male

subject animals underwent the same procedure five days following timed mating. Two

days following the separation reunification test, animals underwent a selective

aggression assay. Partners were again removed from the home cage and separated to

a separate room for an hour. After one hour, a novel sexually naïve animal of the

opposite sex was introduced into the home cage with the subject animal and behavior

observed for 20 minutes. If significant aggression occurred, defined as >3 tussles during

the assay, the animals were separated and the assay terminated. Overall, two WT

female assays and one Oxtr[1-/-] were terminated for aggression. The same behaviors

listed above for the introduction were observed and recorded in the reunification and

aggression assays.

**RNA Sequencing**

*Sample collection*

Adult 7-9 week old WT and Oxtr[1-/-] voles were either cohoused with same sex-siblings

or paired with an animal of the opposite sex for four days following the timed mating

protocol described above. These animals were euthanized by $CO_2$ inhalation and

decapitated. The brain was quickly dissected out and sectioned into 500μm coronal

slices using a brain matrix mold (BrainTree Scientific) chilled on ice. Slices were floated

in ice-cold phosphate buffered saline (PBS), the NAc was identified using anatomical

landmarks, and dissected using a Zeiss microscope. A total of 45 WT females, 39 WT

males, 21 Oxtr[1-/-] females, and 18 Oxtr[1-/-] males were split between paired and unpaired

conditions and tissue from three animals per condition was pooled prior to RNA

extraction. Tissue was flash-frozen in liquid nitrogen and stored at -80°C until further

processing. Total RNA was extracted using TRIzol according to manufacturer's

instructions and quantified using NanoDrop (Thermo Scientific). Library preparation for

next generation sequencing, comprising 35 samples and 4 technical replicates, was

done using TruSeq Stranded mRNA Kit according to manufacturer recommendations

and sequenced on a NovaSeq 6000 to an average depth of 2.9e07 reads per sample.

Sequencing was performed across three runs, but all data was processed together.

***Read processing***

Adapter and low-quality sequence trimming was performed with Trimmomatic v0.39

using parameters: 'TruSeq3-PE.fa:2:30:10:2:keepBothReads LEADING:3 TRAILING:3

SLIDINGWINDOW:4:20 MINLEN:25' (Bolger et al., 2014). Trimmed RNA-Seq reads

were aligned to a custom MicOch1.0 prairie vole genome, which incorporated a known

missing gene V1A, using STAR v2.7.3a in gene annotation mode (Dobin et al., 2013).

Alignment, RNA-Seq, and Insert Size quality control metrics were generated using

Picard v 2.10.10 (http://broadinstitute.github.io/picard). Homologous genes between

vole, mouse, and human references were identified using biomart v2.56.1. When

multiple mouse or human genes mapped to the same vole gene, the homologous gene

with the highest percent identity to vole was selected.

***Differential expression analysis***

To remove non- and lowly-expressed genes, only genes with more than 2 counts per

million (cpm) in at least 3 samples were retained. Principal component analysis of the

cpm revealed that technical replicates all cluster tightly but showed a slight batch effect

due to sequencing run (Fig. S5A). To remove this effect, we used RUVSeq v1.34.0 to

estimate one factor of unwanted variation, $W\_1$, which we later used to normalize the

data prior to differential expression[111]. The factor analysis was performed on the

deviance residuals from an initial generalized linear model regression of the upper

quartile normalized counts on our covariates of interest (pair bonding status, genotype,

sex). Expressed genes were tested for differential expression (DE), after combining

technical replicates, with DESeq2 v1.40.2 using a design matrix that included the

covariates and $W\_1^{112}$. The adaptive shrinkage estimator ashr v2.2-63 was used in

order to be more robust to genes with low cpm values[113]. For DE analyses in general,

while most changes were driven by female samples, the effect was maintained in the

males at a smaller magnitude; thus, analyses were primarily performed for both sexes

jointly to maximize accuracy and power.

*Bonding effect*

To compare pre- to post-pairing individuals, we performed the DE analysis in two ways.

First, we compared pairing status regardless of genotype or sex to increase our

statistical power and to identify a reproducible gene set of thirteen genes (Fig. 5B, 5C,

S5B). While the relationship between the genes is unknown, there is support that the

proteins are functionally related. Using the STRING protein database

(STRINGdb_2.12.1), the human-homologs of the DE genes share more connections

than we would expect by chance (Fig. S5C, p-value= 3.08E-0.7). Next, we compared

pairing status using all three covariates, to identify subgroup-specific genes of interest

and effect sizes (Fig. 5D, S5D). In both analyses, genes were considered significantly

DE with a BH adjusted p-value < 0.05.

*Genotype effect*

To examine the effect of genotype, we first performed DE analyses between WT and

Oxtr$^{-/-}$ individuals, not including sex as a covariate to identify a reproducible and

stringent gene set. We considered genes DE with an adjusted p-value < 0.05 and

absolute log2 fold-change > 0.25 (Fig. 6B, S6A, S6B). To examine the functional

relationship of the DE genes, we performed a Gene Ontology enrichment analysis using

goseq v1.52.0 which controls for gene length bias[114]. We considered all expressed vole

genes with a mouse homolog as our background and all DE genes with a mouse

homolog as our foreground. GO categories with an adjusted p-value < 0.05 were

considered significantly enriched. To help visualize the results, enriched GO categories

were clustered according to their semantic similarity calculated with the R package

GOSemSim v2.26.1 using method "Wang"[115]. Large, general, GO categories with over

1000 genes were excluded in visualizations (Fig. 6C). WikiPathways[116] over-

representation analysis was performed with R package clusterProfiler v4.8.3[117] (Fig.

S6D).

For disorder and disease set overrepresentation analysis, Chi-squared tests were used

to assess whether Oxtr$^{1-/-}$ disrupted genes occur in disease-associated gene lists at a

frequency higher than we would expect by chance.  Genes lists for autism spectrum

disorder[63], schizophrenia[64], neurodevelopmental disorders[66], and epilepsy[65] were

sourced and all alternate or outdated gene symbols were updated manually. For this

test, we were interested in sex- and direction-specific enrichments, so we performed a

DE analysis separately for males and females. Across all DE analyses, we defined DE

genes more stringently (adjusted p-value < 0.01). For enrichment tests, our universe

was all expressed vole genes with a human homolog. We split this analysis by fold-

change direction to highlight the distinct enrichment in upregulated genes. P-values

underwent BH correction across all 24 comparisons.

To compare to mouse neuronal populations, we leveraged single-cell data on mouse

hypothalamus development[72]. The released Seurat object

GSE132730_subset_cca_to_velo.rds.gz was accessed at GSE132730. Using the

authors' defined clusters (accessed at "prim_walktrap"), we assessed how similar each

clusters' gene expression is to the genes disrupted by the $Oxtr^{1-/-}$ mutation. This was

calculated as the Jaccard similarity coefficient using R package clustifyR v1.12.0[118].

***Weighted Gene Coexpression Network Analysis***

<u>Data</u>

Starting with the RNAseq dataset of n=30 samples (one WT sample and 4 replicate

samples were removed) and n=13357 genes, we removed genes with low expression

(n=1738 genes with expression level in lowest quantile for all 30 samples) and/or low

variability (n=8230 genes with coefficient of variation < 0.05), resulting in a trimmed

dataset of n=3389 genes.

<u>Identifying co-expression modules</u>

We used the R package WGCNA (v1.70-3) to conduct WGCNA analyses[61]. Briefly, we

calculated Pearson correlations for all gene pairs. Next, we constructed an unsigned

weighted correlation network to identify co-expression modules comprised of highly

correlated genes (positive or negative correlation) with high topological overlap[119].

Modules were defined as branches of a hierarchical cluster tree using the top-down

dynamic tree cut method[61]. For each module, the expression pattern was summarized

by the module eigengene (ME), which is defined as the right singular vector of the

expression patterns. Pairs of modules with high module eigengene correlations (r >

0.75) were merged to maintain a level of decorrelation among MEs.

In more detail, a weighted unsigned network was computed based on a fit to scale-free

topology, and the lowest thresholding power that resulted in a scale-free $R^2$ fit of 0.9

was selected, and the pairwise topological overlap (TO) between genes was

calculated[119] . We constructed a TO dendrogram using hierarchical clustering (R::hclust

function, method= "average"), and defined modules using the WGCNA::cutreeHybrid

function with minimum module size set to 30 genes. We then merged modules with high

module eigengene correlations as describe above.

We calculated 3 sets of modules from different sample subsets:

1. Modules$^{WT}$: n=15 modules constructed from n=18 WT samples with

thresholding power $\beta = 5$.

2. Modules$^{Mut}$: n= 43 modules constructed from n=12 Oxtr$^1$-mutant samples

with thresholding power $\beta = 6$.

3. Modules$^{All}$: n=7 modules constructed from all n=30 samples with

thresholding power $\beta = 3$

## Relating modules to sample traits

In order to correlate modules with experimental conditions, we calculated the Pearson

correlations between module eigengenes and sample metrics, including species

(Mut=0, WT=1), condition (Pre=1, Post=0), sex (F=1, M=0), and quality control (QC)

metrics (RIN, yield, sequencing run). We identified three modules that significantly

correlated with a sample metric of interest but not with any of the QC metrics.

Significant correlation was defined to be $p_{adj} <= 0.05$, where $p_{adj}$ = BH correction by

number of modules. These modules (M) were:

1. Mpink$^{WT}$: positive correlation with pre-bonding status (Pearson R = 0.7, $p_{adj}$ =

0.018)

2. Mbrown$^{All}$: positive correlation with WT status (Pearson R = 0.55 , $p_{adj}$ = 0.011)

3. Mturquoise$^{All}$: positive correlation with WT status (Pearson R = 0.8, $p_{adj}$ = 6.6e-7)

We calculated the module eigengenes (ME) for Mpink$^{WT}$, Mbrown$^{All}$, and Mturquoise$^{All}$

for all n=30 samples using the WGCNA::moduleEigengenes function. We then

conducted Wilcoxon rank sum tests (ggpubr::stat_compare_means function) to assess

whether MEs were significantly different between the following groups:

1. MEpink$^{WT}$: pre- versus post-bonded animals, separated by sex and species

2. Mbrown$^{All}$: WT versus mutant animals, separated by sex and condition

3. Mturquoise$^{All}$: WT versus mutant animals, separated by sex and condition

## Comparing gene significance with module membership:

We assessed whether genes whose expression highly correlate with module

eigengenes of modules of interest (Mpink$^{WT}$, Mbrown$^{All}$, and Mturquoise$^{All}$) are

themselves highly correlated with traits of interest (bonding status or mutation status).

Specifically, for the genes in Mpink$^{WT}$, Mbrown$^{All}$, and Mturquoise$^{All}$, we calculated 1)

gene significance (GS), defined as the absolute Pearson's correlation between

expression of a given gene and a trait of interest; and 2) module membership (MM),

which is defined as the absolute Pearson's correlation between the expression profile of

a gene with the ME of a module to quantify the relationship between a gene and a given

module[61]. We then calculated the Pearson correlations between:

1.  GS$^{condition}$ and MMpink$^{WT}$ for Mpink$^{WT}$ genes (R = 0.69, p = 5.5e-18)

2.  GS$^{species}$ and MMbrown$^{All}$ for Mbrown$^{All}$ genes (R = 0.63, p = 2.2e-32)

3.  GS$^{species}$ and MMturquoise$^{All}$ for MMturquoise$^{All}$ genes (R= 0.82, p<1e-200)

Hub genes were defined as those genes in the top 25% for both the absolute value of

pink module membership and the absolute value of correlation with bonding status.

<u>Assessing whether modules are enriched for DE genes</u>

We were interested in assessing whether Mpink$^{WT}$ genes are significantly enriched for

any the DE pre vs post pairing gene set described above.  We assessed for enrichment

of DE gene sets using a hypergeometric test.

<u>WT module preservation analysis (Zsummary)</u>

We assessed how well modules$^{WT}$ are preserved in mutant samples.  We calculated a

Z$_{summary}$ statistic for each module$^{WT}$ [120] using the WGCNA::modulePreservation function

with 200 permutations. The $Z_{summary}$ measure combines module density and

intramodular connectivity metrics to a composite statistic where Z>2 suggests moderate

preservation and Z>10 suggests high preservation. We found of all the n=15

modules$^{WT}$, Mpink$^{WT}$ had the lowest $Z_{summary}$ ($Z_{summary}$=2) in mutant samples.

**Histology**

***Immunohistochemistry (IHC)***

Sexually naïve voles group housed by sex after weaning (P0, P21, adults:10 - 14 weeks

old), were perfused with 25mL ice cold phosphate buffer (PBS) and fixed in 25mL of 4%

paraformaldehyde in PBS (PFA). Whole brain was dissected and post fixed overnight at

4°C in 4% PFA in the dark. The tissue was cryoprotected for 24 hours in 30% sucrose in

PBS and then embedded in OCT over dry ice to prevent cracking. 60μm serial sections

from the entire PVN were collected in PBS and processed as free-floating sections.

Sections were blocked using 10% donkey serum in 0.1% Triton X 100 in PBS and then

stained with primary antibodies (Oxytocin Polyclonal Antibody, Invitrogen, Rabbit

Polyclonal, Cat #PA5-26701, 1:1000 and Anti-Neurophysin 2/NP-AVP Antibody, clone

PS 41, Millipore Sigma, Mouse Monoclonal, Cat#MABN845, 1:10,000) in 1% donkey

serum with 0.1% Triton X 100 (staining buffer) in PBS at 4°C with shaking. The sections

were rinsed and soaked in secondary antibodies (Cy™3 AffiniPure Donkey Anti-Rabbit

IgG, Jackson ImmunoResearch Laboratory, Cat #711-165-152, 1:500 and Alexa Flour

488 Donkey Anti Mouse IgG, Thermo Fisher Scientific, Cat #A21202, 1:500) in staining

buffer for 2 hours at room temperature. Sections were DAPI stained and transferred to

glass slides in sequence and mounted in Aquamount. Slides were imaged using a

confocal microscope (Zeiss LSM 700) and the zstacks were quantified (cell count by

hand and area hand selected and measured) in ImageJ. Comparisons of cell density

(count/area) were made in age and sex matched samples.

***In situ hybridization (ISH)***

ISH was performed using RNA probes prepared as described previously[121], to detect

expression of *Calcr, Dlk1*, and *Dgkb* mRNA in the adult vole brain. We prepared RNA

sense and anti-sense probes corresponding to 706bp (CalcR), 773bp (Dgkb), and

796bp (Dlk1) for genes of interest identified in the WT vs Oxtr DEX list from our RNAseq

data. For ISH on adult animals, 7-9 week old WT and Oxtr[1-/-] voles of both sexes were

either group housed with same sex conspecifics or paired for 4 days with a WT animal

of the opposite sex following the timed mating protocol described above. The brains

were processed as described for IHC, then embedded in OCT and sectioned at 50µm,

mounted immediately after sectioning onto SuperFrost Plus glass microscope slides

and stored at -80°C until ISH staining. Slides were fixed in 4% PFA for 20 minutes,

rinsed in PBS, treated with proteinase K (10µg/mL, Roche) for 20 minutes, rinsed in

PBS, and fixed again in 4% PFA for 5 minutes at room temperature. Slides were

acylated for 10 mins then rinsed in 1% Triton100X in PBS followed by PBS. The slides

were equilibrated in warm hybridization solution for 1-2 hours at 65°C and subsequently

incubated with a temporary paraffin cover for 14 - 18 hours at 65°C in fresh

hybridization buffer containing RNA probe. After incubation, the slides were dipped in

5X SSC warmed to 72°C to remove the parafilm coverslips and then washed in 0.2X

SSC warmed to 72°C. The slides were blocked with 10% heat inactivated sheep serum

(HISS) solution then stained for 12 - 18 hours at 4°C in buffer containing 10% HISS and

alkaline phosphatase-conjugated sheep anti- digoxigenin antibody (1:2000, Roche).

After extensive washing, the slides were incubated for ~72 hours at 37°C in staining

solution containing nitro blue tetrazolium and 5- bromo-4-chloro-3-indolyl-phosphate

(Roche). The slides were finally washed, fixed in 4% PFA, and coverslipped. Slides

were imaged using a Keyence fluorescence microscope and the z-stacks were

quantified (cell count and area) in ImageJ. For signal quantitation, positive puncta were

counted in every third coronal section encompassing the entire anterior to posterior

gene expression domain. Cell counts were normalized by total area for a gene

expression domain in each section. Positive signal was quantified by calculating total

area of signal multiplied by the average intensity of signal (total area x mean pixel

intensity) in gene expression domains of a section; every third section was quantified

and added. Comparisons of cell counts were made in age and sex matched samples.

**Quantification and Behavioral Analysis**

For detailed statistical methods please see Table S1. The number of animals used was

based on previous studies in the field by our group and others, combined with a power

analysis. Assumptions of independence and normality were considered with multiple

tests. Preference index (PI) was calculated as (partner huddle duration – stranger

huddle duration)/(partner huddle duration + stranger huddle duration). Sliding partner

preference was calculated by calculating PI for a single animal in 20-minute intervals

where each interval overlapped the next by 15 minutes (e.g. 0-20mins, 5-25mins…). To

look at convergence in sliding PI the area under the curve (Area above 0 – Area below

0, AUC) for each hour of the assay was determined for each animal and then pooled for

each choosing animal by sex and then condition. Preference was determined as

significantly convergent if the AUC from the hour of assay for all animals within the

group significantly differed from 0 using a One sample t Test. The level of statistical

significance for each test was set at p =< 0.05. Outliers were detected using a

combination of z-score (+/- 3) and plots of original and log-transformed data. Analyses

were completed in RStudio (version 2024.04.2+764) and GraphPad Prism (version

10.3.1)

1010

**FIGURE LEGENDS**

**Figure1: Oxtr reduces the amount of time naive animals take to show partner**

**preference**

A. Schematic of the partner preference tests (PPTs) and the day on which they were

conducted to study pair bonding behaviors. The prosocial, agonistic and mating

behaviors scored are shown below. During the bond formation phase, the ratio of

partner-directed vs stranger-directed affiliative behaviors increases in wild types

and following the bond formation in the bond maintenance phase, wild types

show decreased prosocial behaviors and increased agonistic behaviors towards

strangers.

B. A preference for partner is induced in the WTs but not Oxtr$^{1-/-}$ animals following

short cohabitation.

C. WT females spend more time huddling with their partners.

D. Oxtr$^{1-/-}$ females spend more time huddling with strangers.

E. WT females show a significant increase in partner huddling time towards the end

of a 3-hour assay.

F. Oxtr$^{1-/-}$ males show a significant increase in stranger huddling time in the middle

of a 3-hour assay.

G. No difference in partner huddling time between WT and Oxtr$^{1-/-}$ animals following

long cohabitation.

H. No difference in stranger huddling time between WT and Oxtr$^{1-/-}$ animals

following long cohabitation.

See Supplementary Figure 1 and Supplementary Table 1.

Mean +/- SD, n =>8, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT =

male WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$.

**Figure2: Oxtr suppresses promiscuous behaviors in a state dependent manner in**

**females**

A. Timeline of the behavioral battery carried out on females. Bottom row shows the

behaviors scored.

B. Oxtr$^{1-/-}$ female long huddle duration is shorter with partners and longer with

strangers after a short cohabitation.

C. Oxtr$^{1-/-}$ female long huddle frequency is greater with strangers, but not partners

after a short cohabitation.

D. No difference in median long huddle durations between WT and Oxtr$^{1-/-}$ females

post long cohabitation.

E. No difference in long huddle frequencies between WT and Oxtr$^{1-/-}$ females post

long cohabitation.

F. Increasing cohabitation time increases the median long huddle duration an Oxtr$^{1-}$

$^{/-}$ female executes with the partner.

G. There is no significant change in the median long huddle duration conducted by

females with strangers when cohabitation time is increased.

H. Naive Oxtr$^{1-/-}$ and WT females are equally prosocial towards a naive WT male.

I. There is no difference in the counts of mating behaviors between Oxtr$^{1-/-}$ and WT

females.

J. Oxtr$^{1-/-}$ and WT females are equally prosocial towards their partners after a brief

separation.

K. Oxtr$^{1-/-}$ females are significantly more prosocial towards a naive WT stranger

male compared to WT females.

See Supplementary Figure 2 and Supplementary Table 1.

Mean ± SD, n =>8, *p < 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT = male

WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$.

**Figure3: Oxtr function influences different prosocial behaviors across pair**

**bonding in males**

A. Timeline of the behavioral battery carried out on males. Bottom row shows the

behaviors scored.

B. Oxtr$^{1-/-}$ male long huddle duration is shorter with partners and strangers after a

short cohabitation.

C. $Oxtr^{1-/-}$ male long huddle frequency is greater with strangers, but not partners after a short cohabitation.

D. The median long huddle executed by $Oxtr^{1-/-}$ males with the stranger after a long cohabitation, is shorter than their WT siblings.

E. No difference in the long huddle frequencies between WT and $Oxtr^{1-/-}$ males post long cohabitation.

F. There is no change in the frequency of long huddles with partner between $Oxtr^{1-/-}$ and WT males.

G. Increasing cohabitation time decreases the frequency of long huddles with strangers in $Oxtr^{1-/-}$ males.

H. Naive $Oxtr^{1-/-}$ and WT males are equally prosocial towards naive WT females.

I. $Oxtr^{1-/-}$ males execute higher counts of mating behaviors compared to WT.

J. Prosocial behaviors following a brief separation with their partner are not different between $Oxtr^{1-/-}$ and WT males.

K. $Oxtr^{1-/-}$ males are significantly more prosocial towards a naive WT stranger female compared to WT males.

See Supplementary Figure 3 and Supplementary Table 1.

Mean +/- SD, n =>8, *p =< 0.05, FWT = female WT, $FOxtr^{1-/-}$ = female $Oxtr^{1-/-}$, MWT = male WT, $MOxtr^{1-/-}$ = male $Oxtr^{1-/-}$

**Figure4: Oxtr sex-specifically influences social interactions between potential mates**

A. Schematic of naive social choice with a naive WT animal choosing between an opposite sex WT and opposite sex $Oxtr^{1-/-}$ animal (red triangles).

B. Male WT choosers prefer WT females.

C. Preference for WT females over Oxtr$^{1-/-}$ females by WT males manifests during the middle of a 6-hour assay.

D. Schematic of comparison of time in contact with the MOAT from WT-Oxtr$^{1-/-}$ condition to contact with the MOAT from WT-WT condition.

E. WT female chooser spends more time in contact with a MOAT from a WT-Oxtr$^{1-/-}$ condition than the WT-WT. The color of the dots represents the genotype of the MOAT.

F. WT female choosers transition between contacts more frequently in the WT-WT condition compared to the WT-Oxtr$^{1-/-}$ condition.

G. WT female chooser spend significantly more time in contact with the MOAT from the WT-Oxtr$^{1-/-}$ condition in the last 40 mins of the assay.

H. WT female choosers prefer their MOAT over longer periods of time in the WT-Oxtr$^{1-/-}$ condition. WT male choosers prefer their MOAT earlier in the WT-Oxtr$^{1-/-}$ condition, but then show preference for the same amount of time as the WT-WT.

See Supplementary Figure 4 and Supplementary Table 1.

Mean +/- SD, n = 8, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT = male WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$

**Figure 5: Post pair bonding gene expression is driven by female specific changes dependent on Oxtr signaling**

A. Schematic of experimental groups used for sequencing highlighting pre- vs post-pairing comparisons between groups. Animals with symbols represent those used for tissue collection.

B. Scatterplot illustrating the relationship between fold change and mean

expression, emphasizing the 13 differentially expressed (DE) genes in post-

pairing individuals compared to pre-pairing. Gray dots indicate non-significant

genes ($p_{adj} > 0.05$); Blue dots indicate downregulated genes ($p_{adj} < 0.05$, $\log_2FC$

<0) and upregulated genes in red ($p_{adj} < 0.05$, $\log_2FC>0$). The right panel shows

box plots of normalized gene expression for the top six DE genes by fold-change

between pre- and post-pairing conditions.

C. Bar graph emphasizing how each combination of genotype and sex (denoted at

bottom as 'subgroup') influence the 13 overall pairing-DE genes identified. The

top graph depicts the absolute fold change and standard error of the differential

expression analysis including all pre- and post-bonding individuals in which the

13 genes of the x axis were identified. The bottom graph depicts the absolute fold

change and standard error in analyses when only individuals of that subgroup

were considered. (* denotes $p_{adj} < 0.05$ for the pre- vs post-pairing analysis).

D. Box plots of select genes that show sex- and genotype-specific changes in

expression with pairing condition. *Adora2A* and *Fkbp5* are DE when considering

WT females, *Spred3* is DE when considering WT and Oxtr[1-/-] females, and

*Sema3b* is DE when considering Oxtr[1-/-] males. Note: Sema3b is the mouse

ortholog gene symbol for vole gene "ENSMOCG00000019704" which does not

have an assigned gene symbol.

E. Heatmap showing correlations between module eigengenes (MEs) for modules[WT]

and pairing condition (Pre=1, Post=0). Heatmap color and value reflect Pearson

correlation coefficient; adjusted p values shown in parentheses ($p_{adj}$ = p * number

of modules).

F. Preservation of modules$^{WT}$ in OXTR-mutant samples. The $Z_{summary}$ measure

combines module density and intramodular connectivity metrics to a composite

statistic where Z>2 suggests moderate preservation and Z>10 suggests high

preservation[119]. Of all modules$^{WT}$(n=15), Mpink$^{WT}$ had the lowest $Z_{summary}$

($Z_{summary}$=2) in mutant samples.

G. Box plots showing comparison of MEpink$^{WT}$ for pre- vs post-pairing condition

separated by sex and genotype. (Wilcoxon rank sum test, Bonferroni adjusted for

4 comparisons).

For all box plots center = median, box boundaries = 1$^{st}$ and 3$^{rd}$ quartile, whiskers =

1.5*IQR from boundaries. (*n* = 16 pre-pairing and 15 post-pairing)

See Supplementary Figure 5.

**Figure 6: Oxtr regulated genes are common to social behavior relevant processes**

**across species and are similarly regulated across brain regions**

A. Schematic of experimental groups for sequencing, highlighting genotype

comparisons between groups. Animals with symbols represent those used for

tissue collection.

B. Scatterplot illustrating the relationship between fold change and mean

expression, emphasizing differentially expressed (DE) genes in Oxtr$^{1-/-}$

individuals compared to WT. Significantly downregulated genes are shown in

blue ($log_2FC$<-0.25, $p_{adj}$<0.05) and upregulated genes in red ($log_2FC$>0.25, $p_{adj}$ <

0.05). Box plots of normalized gene expression for the top six DE genes by fold

change between genotypes.

C.  Gene Ontology enrichment analysis for all genotype-DE genes (n=1014)

highlights processes related to extracellular channel binding activity, axonal

processes, behavior related gene expression, and cation transport. The leaves

represent significant GO categories (BH adjusted p value < 0.05) which have

been hierarchically clustered by their semantic similarity. Branches are colored

by cluster and high-frequency words are displayed to the right. Leaf size and

color correspond to the number and median fold change of DE genes in each

category.

D.  Enrichment analysis shows that genes which are DE in $Oxtr^{1-/-}$ samples are over-

represented in some neuropsychiatric disease-associated gene sets. The first DE

gene set evaluated ('Joint') represents genes which are DE in $Oxtr^{1-/-}$ samples

when considering both sexes in the differential expression analysis ($p_{adj}$ < 0.01).

The second gene set are DE genes when only considering females, and the third

when only considering males. Gene sets are further separated by direction of

effect across the x-axis to highlight the enrichment in upregulated DE genes.

Colors correspond to the Benjamini-Hochberg corrected Chi-square p values;

asterisks denote $p_{adj}$ < 0.05. Y-axis abbreviations are as follows:

SCZ=schizophrenia, NDD=neurodevelopmental disorder, EPI=epilepsy, and

ASD=autism spectrum disorder.

E. Box plots of *Calcr* and *dlk1* normalized gene counts by genotype. Log$_2$FC:-0.962

and p$_{adj}$: 1.75e-06 for *Calcr* and log$_2$FC: -0.276 and p$_{adj}$: 0.00334 for *dlk1* from

comparison considering all WT and Oxtr$^{1-/-}$ samples.

F. Brain regions selected for in situ analysis. (NAc=nucleus accumbens,

PVN=paraventricular nucleus of the hypothalamus, LS=lateral septum)

G. ISH images and quantification for *Calcr* expression in NAc, PVN, and LS from

WT and Oxtr$^{1-/-}$ animals.

H. ISH images and quantification for *Dlk-1* expression in NAc, PVN, and LS from

WT and Oxtr$^{1-/-}$ animals.

Box plots in G. and H. show comparison of the area-normalized cell number by

genotype. (*n*=3 per condition, Wilcoxon sign rank, *=p<0.05, ***=p<0.001)

See Supplementary Figure 6 and Supplementary Table 1.

**Figure7: Loss of Oxtr results in the decrease in density of Oxt and Avp**

**expressing neurons in the PVN**

A. UMAP visualization of mouse hypothalamus single-cell RNAseq dataset[66] with

proposed PVN clusters labeled. The authors' original numeric cluster labels and

embedding are shown. Each cluster is colored by the Jaccard similarity

coefficient between the mouse marker gene sets and the vole Oxtr$^{1-/-}$ DE genes

(ONP set). This is also represented in a heat plot to the right. Some of the

highest similarity is seen in Cluster 43, which represents Oxt+ cells, and Cluster

26, which represents Avp+ cells in the PVN.

B. Representative images of IHC staining against OT and AVP neuropeptides from

FWT and FOxtr$^{1-/-}$ PVN showing the loss of structural integrity and OT and AVP

positive cells from the knock-out. (White dotted line denotes the PVN area, scale

bar represents 100µm)

C. Mean of positive cells per unit area across all sections is lower for Oxtr$^{1-/-}$s vs WT

for both OT (left) and AVP (right). Each dot represents 1 animal.

D. The number of sections with a high density of OT positive cells is greater in both

female (left) and male (right) WT animals.

E. Similarly, WT animals of both sexes show an increase in the number of sections

containing AVP positive neurons in high densities.

F. The average OT cell density from matched PVN sections arranged from anterior

to posterior shows that the decrease in cell density in Oxtr$^{1-/-}$ animals is biased

towards the anterior. Each dot represents the average cell density, the lines

represent the sixth order non-linear regression model for cell density by sex and

by genotype (n=3) and the ribbons represent the standard deviation.

G. Oxtr$^{1-/-}$ animals show a significant decrease in AVP cell density in the anterior

PVN in both sexes.

See Supplementary Figure 7 and Supplementary Table 1.

Mean +/- SD, n = 3, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT =

male WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$.

1220

1221

1222  **References**

1223

1. Wilson, R.S., Krueger, K.R., Arnold, S.E., Schneider, J.A., Kelly, J.F., Barnes, L.L.,

Tang, Y., and Bennett, D.A. (2007). Loneliness and Risk of Alzheimer Disease. Arch

Gen Psychiatry *64*, 234. https://doi.org/10.1001/archpsyc.64.2.234.

2. Bowlby, J., and Bowlby, J. (1982). Attachment 2. ed. (Basic Books).

3. Farinelli, M., Cevolani, D., Gestieri, L., Romaniello, C., Maffei, M., Agati, R., Leo,

M.R., Huang, Z., Pedone, V., and Northoff, G. (2020). Brain and behaviour in post-

acute stroke: Reduction in seeking and posterior cingulate neuronal variability.

Journal of Clinical and Experimental Neuropsychology *42*, 584–601.

https://doi.org/10.1080/13803395.2020.1780417.

4. Feldman, R., Monakhov, M., Pratt, M., and Ebstein, R.P. (2016). Oxytocin Pathway

Genes: Evolutionary Ancient System Impacting on Human Affiliation, Sociality, and

Psychopathology. Biological Psychiatry *79*, 174–184.

https://doi.org/10.1016/j.biopsych.2015.08.008.

5. Ebstein, R.P., Israel, S., Chew, S.H., Zhong, S., and Knafo, A. (2010). Genetics of

human social behavior. Neuron *65*, 831–844.

https://doi.org/10.1016/j.neuron.2010.02.020.

6. Winslow, J.T., Shapiro, L., Carter, C.S., and Insel, T.R. (1993). Oxytocin and

complex social behavior: species comparisons. Psychopharmacol Bull *29*, 409–414.

7. Goodson, J.L. (2013). Deconstructing sociality, social evolution and relevant

nonapeptide functions. Psychoneuroendocrinology *38*, 465–478.

https://doi.org/10.1016/j.psyneuen.2012.12.005.

8. Harlow, H.F., and Harlow, M.K. (1965). THE EFFECT OF REARING CONDITIONS

ON BEHAVIOR. Int J Psychiatry *1*, 43–51.

9. Shultz, S., Opie, C., and Atkinson, Q.D. (2011). Stepwise evolution of stable sociality

in primates. Nature *479*, 219–222.

10. Insel, T.R., Winslow, J.T., Wang, Z., and Young, L.J. (1998). Oxytocin, vasopressin,

and the neuroendocrine basis of pair bond formation. Adv. Exp. Med. Biol *449*, 215–

224.

11. Klatt, J.D., and Goodson, J.L. (2013). Oxytocin-like receptors mediate pair bonding

in a socially monogamous songbird. Proceedings of the Royal Society B: Biological

Sciences *280*, 20122396. https://doi.org/10.1098/rspb.2012.2396.

12. Ainsworth, M.S. (1979). Infant–mother attachment. American Psychologist *34*, 932–

937. https://doi.org/10.1037/0003-066X.34.10.932.

13. Getz, L.L., Carter, C.S., and Gavish, L. (1981). The mating system of the prairie

vole, Microtus ochrogaster: Field and laboratory evidence for pair-bonding. Behav

Ecol Sociobiol *8*, 189–194. https://doi.org/10.1007/BF00299829.

14. Donaldson, Z.R. (2010). We're the Same... but Different: Addressing Academic

Divides in the Study of Brain and Behavior. Front Behav Neurosci *4*.

https://doi.org/10.3389/fnbeh.2010.00041.

15. Insel, T.R., Winslow, J.T., Wang, Z.X., Young, L., and Hulihan, T.J. (1995). Oxytocin

and the molecular basis of monogamy. Adv. Exp. Med. Biol *395*, 227–234.

16. Lukas, D., and Clutton-Brock, T.H. (2013). The evolution of social monogamy in

mammals. Science *341*, 526–530.

17. Donaldson, Z.R., and Young, L.J. (2008). Oxytocin, vasopressin, and the

neurogenetics of sociality. Science *322*, 900–904.

https://doi.org/10.1126/science.1158668.

18. Bielsky, I.F., and Young, L.J. (2004). Oxytocin, vasopressin, and social recognition in

mammals. Peptides *25*, 1565–1574. https://doi.org/10.1016/j.peptides.2004.05.019.

19. Nowicki, J.P., Pratchett, M.S., Walker, S.P.W., Coker, D.J., and O'Connell, L.A.

(2020). Gene expression correlates of social evolution in coral reef butterflyfishes.

Proceedings of the Royal Society B: Biological Sciences *287*, 20200239.

https://doi.org/10.1098/rspb.2020.0239.

20. O'Connor, C.M., Marsh-Rollo, S.E., Aubin-Horth, N., and Balshine, S. (2016).

Species-specific patterns of nonapeptide brain gene expression relative to pair-

bonding behavior in grouping and non-grouping cichlids. Hormones and Behavior

*80*, 30–38. https://doi.org/10.1016/j.yhbeh.2015.10.015.

21. Stanley, D.A., and Adolphs, R. (2013). Toward a neural basis for social behavior.

Neuron *80*, 816–826.

22. Insel, T.R. (1992). Oxytocin--a neuropeptide for affiliation: evidence from behavioral,

receptor autoradiographic, and comparative studies. Psychoneuroendocrinology *17*,

3–35.

23. Lim, M.M., and Young, L.J. (2006). Neuropeptidergic regulation of affiliative behavior

and social bonding in animals. Horm Behav *50*, 506–517.

https://doi.org/10.1016/j.yhbeh.2006.06.028.

24. Carter, C.S., and Getz, L.L. (1993). Monogamy and the prairie vole. Sci. Am *268*,

100–106.

25. Beery, A.K., and Shambaugh, K.L. (2021). Comparative assessment of

familiarity/novelty preferences in rodents. Frontiers in Behavioral Neuroscience *15*,

648830.

26. Beery, A.K., Christensen, J.D., Lee, N.S., and Blandino, K.L. (2018). Specificity in

Sociality: Mice and Prairie Voles Exhibit Different Patterns of Peer Affiliation.

Frontiers in Behavioral Neuroscience *12*, 50.

https://doi.org/10.3389/fnbeh.2018.00050.

27. Schweinfurth, M.K., Neuenschwander, J., Engqvist, L., Schneeberger, K., Rentsch,

A.K., Gygax, M., and Taborsky, M. (2017). Do female Norway rats form social

bonds? Behavioral Ecology and Sociobiology *71*, 1–9.

28. Cho, M.M., DeVries, A.C., Williams, J.R., and Carter, C.S. (1999). The effects of

oxytocin and vasopressin on partner preferences in male and female prairie voles

(Microtus ochrogaster). Behav. Neurosci *113*, 1071–1079.

29. Insel, T.R., and Hulihan, T.J. (1995). A gender-specific mechanism for pair bonding:

oxytocin and partner preference formation in monogamous voles. Behav. Neurosci

*109*, 782–789.

30. Williams, J.R., Catania, K.C., and Carter, C.S. (1992). Development of partner

preferences in female prairie voles (Microtus ochrogaster): the role of social and

sexual experience. Hormones and behavior *26*, 339–349.

31. Wang, Z., Hulihan, T.J., and Insel, T.R. (1997). Sexual and social experience is

associated with different patterns of behavior and neural activation in male prairie

voles. Brain Res *767*, 321–332.

32. Xu, X., Coats, J.K., Yang, C.F., Wang, A., Ahmed, O.M., Alvarado, M., Izumi, T., and

Shah, N.M. (2012). Modular genetic control of sexually dimorphic behaviors. Cell

*148*, 596–607. https://doi.org/10.1016/j.cell.2011.12.018.

33. Villella, A., and Hall, J.C. (2008). Chapter 3 Neurogenetics of Courtship and Mating

in Drosophila. In Advances in Genetics (Elsevier), pp. 67–184.

https://doi.org/10.1016/S0065-2660(08)00603-2.

34. Konopka, R.J., and Benzer, S. (1971). Clock Mutants of *Drosophila melanogaster*.

Proc Natl Acad Sci USA *68*, 2112–2116. https://doi.org/10.1073/pnas.68.9.2112.

35. Takahashi, J.S., Shimomura, K., and Kumar, V. (2008). Searching for genes

underlying behavior: lessons from circadian rhythms. Science *322*, 909–912.

https://doi.org/10.1126/science.1158822.

36. Baker, B.S., Taylor, B.J., and Hall, J.C. (2001). Are complex behaviors specified by

dedicated regulatory genes? Reasoning from Drosophila. Cell *105*, 13–24.

37. Manoli, D.S., and Baker, B.S. (2004). Median bundle neurons coordinate behaviours

during Drosophila male courtship. Nature *430*, 564–569.

https://doi.org/10.1038/nature02713.

38. Martin, L.B., Glasper, E.R., Nelson, R.J., and DeVries, A.C. (2006). Prolonged

separation delays wound healing in monogamous California mice, Peromyscus

californicus, but not in polygynous white-footed mice, P. leucopus. Physiology &

Behavior *87*, 837–841. https://doi.org/10.1016/j.physbeh.2006.01.035.

39. Ferland, C.L., and Schrader, L.A. (2011). Cage mate separation in pair-housed male

rats evokes an acute stress corticosterone response. Neuroscience Letters *489*,

154–158. https://doi.org/10.1016/j.neulet.2010.12.006.

40. Okuda, T., Osako, Y., Hidaka, C., Nishihara, M., Young, L.J., Mitsui, S., and Yuri, K.

(2022). Separation from a bonded partner alters neural response to inflammatory

pain in monogamous rodents. Behavioural Brain Research *418*, 113650.

https://doi.org/10.1016/j.bbr.2021.113650.

41. Carter, C.S., Williams, J.R., Witt, D.M., and Insel, T.R. (1992). Oxytocin and social

bonding. Ann. N. Y. Acad. Sci *652*, 204–211.

42. Winslow, J.T., Hastings, N., Carter, C.S., Harbaugh, C.R., and Insel, T.R. (1993). A

role for central vasopressin in pair bonding in monogamous prairie voles. Nature

*365*, 545–548. https://doi.org/10.1038/365545a0.

43. Carter, C.S., Grippo, A.J., Pournajafi-Nazarloo, H., Ruscio, M.G., and Porges, S.W.

(2008). Oxytocin, vasopressin and sociality. Prog. Brain Res *170*, 331–336.

https://doi.org/10.1016/S0079-6123(08)00427-5.

44. Ophir, A.G., Gessel, A., Zheng, D.-J., and Phelps, S.M. (2012). Oxytocin receptor

density is associated with male mating tactics and social monogamy. Hormones and

behavior *61*, 445–453.

45. Shapiro, L.E., and Insel, T.R. (1992). Oxytocin receptor distribution reflects social

organization in monogamous and polygamous voles. Ann. N. Y. Acad. Sci *652*, 448–

451.

46. Heinrichs, M., von Dawans, B., and Domes, G. (2009). Oxytocin, vasopressin, and

human social behavior. Front Neuroendocrinol *30*, 548–557.

https://doi.org/10.1016/j.yfrne.2009.05.005.

47. Insel, T.R., Wang, Z.X., and Ferris, C.F. (1994). Patterns of brain vasopressin

receptor distribution associated with social organization in microtine rodents. J.

Neurosci *14*, 5381–5392.

48. King, L.B., Walum, H., Inoue, K., Eyrich, N.W., and Young, L.J. (2016). Variation in

the oxytocin receptor gene predicts brain region–specific expression and social

attachment. Biological psychiatry *80*, 160–169.

49. Walum, H., and Young, L.J. (2018). The neural mechanisms and circuitry of the pair

bond. Nat Rev Neurosci *19*, 643–654. https://doi.org/10.1038/s41583-018-0072-6.

50. Williams, J.R., Insel, T.R., Harbaugh, C.R., and Carter, C.S. (1994). Oxytocin

administered centrally facilitates formation of a partner preference in female prairie

voles (Microtus ochrogaster). J. Neuroendocrinol *6*, 247–250.

51. Bales, K.L., and Carter, C.S. (2003). Developmental exposure to oxytocin facilitates

partner preferences in male prairie voles (Microtus ochrogaster). Behavioral

neuroscience *117*, 854.

52. Berendzen, K.M., Sharma, R., Mandujano, M.A., Wei, Y., Rogers, F.D., Simmons,

T.C., Seelke, A.M., Bond, J.M., Larios, R., Goodwin, N.L., et al. (2023). Oxytocin

receptor is not required for social attachment in prairie voles. Neuron *111*, 787–796.

53. Young, L.J., Lim, M.M., Gingrich, B., and Insel, T.R. (2001). Cellular mechanisms of

social attachment. Hormones and behavior *40*, 133–138.

54. Tripp, J.A., Berrio, A., McGraw, L.A., Matz, M.V., Davis, J.K., Inoue, K., Thomas,

J.W., Young, L.J., and Phelps, S.M. (2021). Comparative neurotranscriptomics

reveal widespread species differences associated with bonding. BMC Genomics *22*,

399. https://doi.org/10.1186/s12864-021-07720-0.

55. Young, L.J., and Wang, Z. (2004). The neurobiology of pair bonding. Nat. Neurosci

*7*, 1048–1054. https://doi.org/10.1038/nn1327.

56. DeVries, A.C., and Carter, C.S. (1999). Sex differences in temporal parameters of

partner preference in prairie voles (Microtus ochrogaster). Can. J. Zool. *77*, 885–

889. https://doi.org/10.1139/z99-054.

57. Fraser, E.J., and Shah, N.M. (2014). Complex Chemosensory Control of Female

Reproductive Behaviors. PLOS ONE *9*, e90368.

https://doi.org/10.1371/journal.pone.0090368.

58. Aragona, B.J., Liu, Y., Curtis, J.T., Stephan, F.K., and Wang, Z. (2003). A Critical

Role for Nucleus Accumbens Dopamine in Partner-Preference Formation in Male

Prairie Voles. J. Neurosci. *23*, 3483–3490. https://doi.org/10.1523/JNEUROSCI.23-

08-03483.2003.

59. Dölen, G., Darvishzadeh, A., Huang, K.W., and Malenka, R.C. (2013). Social reward

requires coordinated activity of nucleus accumbens oxytocin and serotonin. Nature

*501*, 179–184. https://doi.org/10.1038/nature12518.

60. Ross, H.E., Freeman, S.M., Spiegel, L.L., Ren, X., Terwilliger, E.F., and Young, L.J.

(2009). Variation in Oxytocin Receptor Density in the Nucleus Accumbens Has

Differential Effects on Affiliative Behaviors in Monogamous and Polygamous Voles.

J. Neurosci. *29*, 1312–1318. https://doi.org/10.1523/JNEUROSCI.5039-08.2009.

61. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted

correlation network analysis. BMC Bioinformatics *9*, 559.

https://doi.org/10.1186/1471-2105-9-559.

62. Maury, E.A., Sherman, M.A., Genovese, G., Gilgenast, T.G., Kamath, T., Burris, S.J.,

Rajarajan, P., Flaherty, E., Akbarian, S., Chess, A., et al. (2023). Schizophrenia-

associated somatic copy-number variants from 12,834 cases reveal recurrent

NRXN1 and ABCB11 disruptions. Cell Genom *3*, 100356.

https://doi.org/10.1016/j.xgen.2023.100356.

63. Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.-Y.,

Peng, M., Collins, R., Grove, J., Klei, L., et al. (2020). Large-Scale Exome

Sequencing Study Implicates Both Developmental and Functional Changes in the

Neurobiology of Autism. Cell *180*, 568-584.e23.

https://doi.org/10.1016/j.cell.2019.12.036.

64. Fromer, M., Pocklington, A.J., Kavanagh, D.H., Williams, H.J., Dwyer, S., Gormley,

P., Georgieva, L., Rees, E., Palta, P., Ruderfer, D.M., et al. (2014). De novo

mutations in schizophrenia implicate synaptic networks. Nature *506*, 179–184.

https://doi.org/10.1038/nature12929.

65. EuroEPINOMICS-RES Consortium, Epilepsy Phenome/Genome Project, and Epi4K

Consortium (2014). De novo mutations in synaptic transmission genes including

DNM1 cause epileptic encephalopathies. Am J Hum Genet *95*, 360–370.

https://doi.org/10.1016/j.ajhg.2014.08.013.

66. Kaplanis, J., Samocha, K.E., Wiel, L., Zhang, Z., Arvai, K.J., Eberhardt, R.Y.,

Gallone, G., Lelieveld, S.H., Martin, H.C., McRae, J.F., et al. (2020). Evidence for 28

genetic disorders discovered by combining healthcare and research data. Nature

*586*, 757–762. https://doi.org/10.1038/s41586-020-2832-5.

67. Inoue, K., Ford, C.L., Horie, K., and Young, L.J. (2022). Oxytocin receptors are

widely distributed in the prairie vole (Microtus ochrogaster) brain: Relation to social

behavior, genetic polymorphisms, and the dopamine system. J Comp Neurol *530*,

2881–2900. https://doi.org/10.1002/cne.25382.

68. Lefevre, A., Benusiglio, D., Tang, Y., Krabichler, Q., Charlet, A., and Grinevich, V.

(2021). Oxytocinergic Feedback Circuitries: An Anatomical Basis for

Neuromodulation of Social Behaviors. Front. Neural Circuits *15*, 688234.

https://doi.org/10.3389/fncir.2021.688234.

69. Harris, G.W. (1955). PITUITARY-HYPOTHALAMIC MECHANISMS. Arch NeurPsych

*73*, 124. https://doi.org/10.1001/archneurpsyc.1955.02330080002002.

70. Ye, Q., Nunez, J., and Zhang, X. (2022). Oxytocin Receptor-Expressing Neurons in

the Paraventricular Thalamus Regulate Feeding Motivation through Excitatory

Projections to the Nucleus Accumbens Core. J. Neurosci. *42*, 3949–3964.

https://doi.org/10.1523/JNEUROSCI.2042-21.2022.

71. Zhang, B., Qiu, L., Xiao, W., Ni, H., Chen, L., Wang, F., Mai, W., Wu, J., Bao, A., Hu,

H., et al. (2021). Reconstruction of the Hypothalamo-Neurohypophysial System and

Functional Dissection of Magnocellular Oxytocin Neurons in the Brain. Neuron *109*,

331-346.e7. https://doi.org/10.1016/j.neuron.2020.10.032.

72. Romanov, R.A., Tretiakov, E.O., Kastriti, M.E., Zupancic, M., Häring, M.,

Korchynska, S., Popadin, K., Benevento, M., Rebernik, P., Lallemend, F., et al.

(2020). Molecular design of hypothalamus development. Nature *582*, 246–252.

https://doi.org/10.1038/s41586-020-2266-0.

73. Mitre, M., Saadipour, K., Williams, K., Khatri, L., Froemke, R.C., and Chao, M.V.

(2022). Transactivation of TrkB Receptors by Oxytocin and Its G Protein-Coupled

Receptor. Front Mol Neurosci *15*, 891537.

https://doi.org/10.3389/fnmol.2022.891537.

74. Seoyoung Son, Steffy B. Manjila, Kyra T. Newmaster, Yuan-ting Wu, Daniel J.

Vanselow, Matt Ciarletta, Todd E. Anthony, Keith C. Cheng, and Yongsoo Kim

(2022). Whole-Brain Wiring Diagram of Oxytocin System in Adult Mice. J. Neurosci.

*42*, 5021. https://doi.org/10.1523/JNEUROSCI.0307-22.2022.

75. Vaidyanathan, R., and Hammock, E.A.D. (2020). Oxytocin receptor gene loss

influences expression of the oxytocin gene in C57BL/6J mice in a sex- and age-

dependent manner. Journal of Neuroendocrinology *32*, e12821.

https://doi.org/10.1111/jne.12821.

76. Newmaster, K.T., Nolan, Z.T., Chon, U., Vanselow, D.J., Weit, A.R., Tabbaa, M.,

Hidema, S., Nishimori, K., Hammock, E.A.D., and Kim, Y. (2020). Quantitative

cellular-resolution map of the oxytocin receptor in postnatally developing mouse

brains. Nature Communications *11*, 1885. https://doi.org/10.1038/s41467-020-

15659-1.

77. Kleiman, D.G. (1977). Monogamy in Mammals. The Quarterly Review of Biology *52*,

39–69. https://doi.org/10.1086/409721.

78. Feldman, R. (2017). The Neurobiology of Human Attachments. Trends in Cognitive

Sciences *21*, 80–99. https://doi.org/10.1016/j.tics.2016.11.007.

79. Yang, C.F., and Shah, N.M. (2014). Representing sex in the brain, one module at a

time. Neuron *82*, 261–278. https://doi.org/10.1016/j.neuron.2014.03.029.

80. Mong, J.A., and Pfaff, D.W. (2004). Hormonal symphony: steroid orchestration of

gene modules for sociosexual behaviors. Mol Psychiatry *9*, 550–556.

https://doi.org/10.1038/sj.mp.4001493.

81. Mong, J., Easton, A., Kow, L.M., and Pfaff, D. (2003). Neural, hormonal and genetic

mechanisms for the activation of brain and behavior. Eur J Pharmacol *480*, 229–

231. https://doi.org/10.1016/j.ejphar.2003.08.109.

82. Barron, A.B., and Robinson, G.E. (2008). The utility of behavioral models and

modules in molecular analyses of social behavior. Genes Brain and Behavior *7*,

257–265. https://doi.org/10.1111/j.1601-183X.2007.00344.x.

83. Kyriacou, C.P., and Hall, J.C. (1980). Circadian rhythm mutations in Drosophila

melanogaster affect short-term fluctuations in the male's courtship song. Proc Natl

Acad Sci U S A *77*, 6729–6733. https://doi.org/10.1073/pnas.77.11.6729.

84. Konopka, R.J., and Benzer, S. (1971). Clock Mutants of *Drosophila melanogaster*.

Proc Natl Acad Sci USA *68*, 2112–2116. https://doi.org/10.1073/pnas.68.9.2112.

85. Bendesky, A., Kwon, Y.-M., Lassance, J.-M., Lewarch, C.L., Yao, S., Peterson, B.K.,

He, M.X., Dulac, C., and Hoekstra, H.E. (2017). The genetic basis of parental care

evolution in monogamous mice. Nature *544*, 434–439.

https://doi.org/10.1038/nature22074.

86. Scheele, D., Striepens, N., Güntürkün, O., Deutschländer, S., Maier, W., Kendrick,

K.M., and Hurlemann, R. (2012). Oxytocin modulates social distance between males

and females. J Neurosci *32*, 16074–16079.

https://doi.org/10.1523/JNEUROSCI.2755-12.2012.

87. Ditzen, B., Schaer, M., Gabriel, B., Bodenmann, G., Ehlert, U., and Heinrichs, M.

(2009). Intranasal Oxytocin Increases Positive Communication and Reduces

Cortisol Levels During Couple Conflict. Biological Psychiatry *65*, 728–731.

https://doi.org/10.1016/j.biopsych.2008.10.011.

88. De Dreu, C.K.W., Greer, L.L., Handgraaf, M.J.J., Shalvi, S., Van Kleef, G.A., Baas,

M., Ten Velden, F.S., Van Dijk, E., and Feith, S.W.W. (2010). The Neuropeptide

Oxytocin Regulates Parochial Altruism in Intergroup Conflict Among Humans.

Science *328*, 1408–1411. https://doi.org/10.1126/science.1189047.

89. Osakada, T., Yan, R., Jiang, Y., Wei, D., Tabuchi, R., Dai, B., Wang, X., Zhao, G.,

Wang, C.X., Liu, J.-J., et al. (2024). A dedicated hypothalamic oxytocin circuit

controls aversive social learning. Nature *626*, 347–356.

https://doi.org/10.1038/s41586-023-06958-w.

90. Wang, Z., and Young, L.J. (1997). Ontogeny of oxytocin and vasopressin receptor

binding in the lateral septum in prairie and montane voles. Brain Res. Dev. Brain

Res *104*, 191–195.

91. Smith, M.L., Asada, N., and Malenka, R.C. (2021). Anterior cingulate inputs to

nucleus accumbens control the social transfer of pain and analgesia. Science *371*,

153–159. https://doi.org/10.1126/science.abe3040.

92. Gegenhuber, B., Wu, M.V., Bronstein, R., and Tollkuhn, J. (2020). Regulation of

neural gene expression by estrogen receptor alpha (Neuroscience)

https://doi.org/10.1101/2020.10.21.349290.

93. Cao, J., and Patisaul, H.B. (2013). Sex-specific expression of estrogen receptors α

and β and Kiss1 in the postnatal rat amygdala. J of Comparative Neurology *521*,

465–478. https://doi.org/10.1002/cne.23185.

94. Mitsushima, D., Yamada, K., Takase, K., Funabashi, T., and Kimura, F. (2006). Sex

differences in the basolateral amygdala: the extracellular levels of serotonin and

dopamine, and their responses to restraint stress in rats. Eur J Neurosci *24*, 3245–

3254. https://doi.org/10.1111/j.1460-9568.2006.05214.x.

95. McCann, K.E., Sinkiewicz, D.M., Rosenhauer, A.M., Beach, L.Q., and Huhman, K.L.

(2019). Transcriptomic Analysis Reveals Sex-Dependent Expression Patterns in the

Basolateral Amygdala of Dominant and Subordinate Animals After Acute Social

Conflict. Mol Neurobiol *56*, 3768–3779. https://doi.org/10.1007/s12035-018-1339-7.

96. Trainor, B.C. (2011). Stress responses and the mesolimbic dopamine system: social

contexts and sex differences. Horm Behav *60*, 457–469.

https://doi.org/10.1016/j.yhbeh.2011.08.013.

97. Wright, E.C., Luo, P.X., Zakharenkov, H.C., Serna Godoy, A., Lake, A.A., Prince,

Z.D., Sekar, S., Culkin, H.I., Ramirez, A.V., Dwyer, T., et al. (2023). Sexual

differentiation of neural mechanisms of stress sensitivity during puberty. Proc. Natl.

Acad. Sci. U.S.A. *120*, e2306475120. https://doi.org/10.1073/pnas.2306475120.

98. Dölen, G. (2015). Oxytocin: parallel processing in the social brain? J.

Neuroendocrinol. *27*, 516–535. https://doi.org/10.1111/jne.12284.

99. Choi, T.-Y., Jeon, H., Jeong, S., Kim, E.J., Kim, J., Jeong, Y.H., Kang, B., Choi, M.,

and Koo, J.W. (2023). Distinct prefrontal projection activity and transcriptional state

conversely orchestrate social competition and hierarchy. Neuron, S0896-

6273(23)00886-3. https://doi.org/10.1016/j.neuron.2023.11.012.

100.   Yaguchi, K., Hagihara, M., Konno, A., Hirai, H., Yukinaga, H., and Miyamichi, K.

(2023). Dynamic modulation of pulsatile activities of oxytocin neurons in lactating

wild-type mice. PLoS One *18*, e0285589.

https://doi.org/10.1371/journal.pone.0285589.

101.   Manoli, D.S., and State, M.W. (2021). Autism Spectrum Disorder Genetics and

the Search for Pathological Mechanisms. AJP *178*, 30–38.

https://doi.org/10.1176/appi.ajp.2020.20111608.

102.   Werling, D.M. (2016). The role of sex-differential biology in risk for autism

spectrum disorder. Biology of Sex Differences *7*, 58. https://doi.org/10.1186/s13293-

016-0112-8.

103. Werling, D.M., and Geschwind, D.H. (2013). Sex differences in autism spectrum

disorders. Curr. Opin. Neurol. *26*, 146–153.

https://doi.org/10.1097/WCO.0b013e32835ee548.

104. Perissi, V., and Rosenfeld, M.G. (2005). Controlling nuclear receptors: the

circular logic of cofactor cycles. Nature Reviews: Molecular Cell Biology *6*, 542–554.

https://doi.org/10.1038/nrm1682.

105. Horie, K., Inoue, K., Nishimori, K., and Young, L.J. (2020). Investigation of Oxtr-

expressing Neurons Projecting to Nucleus Accumbens using Oxtr-ires-Cre Knock-in

prairie Voles (Microtus ochrogaster). Neuroscience *448*, 312–324.

https://doi.org/10.1016/j.neuroscience.2020.08.023.

106. Villanueva, C., Jacquier, S., and de Roux, N. (2012). DLK1 Is a Somato-Dendritic

Protein Expressed in Hypothalamic Arginine-Vasopressin and Oxytocin Neurons.

PLoS One *7*, e36134. https://doi.org/10.1371/journal.pone.0036134.

107. Insel, T.R. (1997). A neurobiological basis of social attachment. Am J Psychiatry

*154*, 726–735.

108. Carter, C.S., and Perkeybile, A.M. (2018). The Monogamy Paradox: What Do

Love and Sex Have to Do With It? Front. Ecol. Evol. *6*, 202.

https://doi.org/10.3389/fevo.2018.00202.

109. De Vries, G.J. (2004). Minireview: Sex Differences in Adult and Developing

Brains: Compensation, Compensation, Compensation. Endocrinology *145*, 1063–

1068. https://doi.org/10.1210/en.2003-1504.

110.   Friard, O., and Gamba, M. (2016). BORIS: a free, versatile open-source event-

logging software for video/audio coding and live observations. Methods in ecology

and evolution *7*, 1325–1330.

111.   Risso, D., Ngai, J., Speed, T.P., and Dudoit, S. (2014). Normalization of RNA-seq

data using factor analysis of control genes or samples. Nat Biotechnol *32*, 896–902.

https://doi.org/10.1038/nbt.2931.

112.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold

change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

https://doi.org/10.1186/s13059-014-0550-8.

113.   Stephens, M., Carbonetto, P., Gerard, D., Lu, M., Sun, L., Willwerscheid, J., and

Xiao, N. (2020). Methods for Adaptive Shrinkage, using Empirical Bayes [R package

ashr version 2.2-47]. In.

114.   Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene

ontology analysis for RNA-seq: accounting for selection bias. Genome Biol *11*, R14.

https://doi.org/10.1186/gb-2010-11-2-r14.

115.   Yu, G. (2020). Gene Ontology Semantic Similarity Analysis Using GOSemSim. In

Stem Cell Transcriptional Networks Methods in Molecular Biology., B. L. Kidder, ed.

(Springer US), pp. 207–215. https://doi.org/10.1007/978-1-0716-0301-7_11.

116.   Pico, A.R., Kelder, T., Van Iersel, M.P., Hanspers, K., Conklin, B.R., and Evelo, C.

(2008). WikiPathways: Pathway Editing for the People. PLoS Biol *6*, e184.

https://doi.org/10.1371/journal.pbio.0060184.

117.  Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W.,

Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting

omics data. The Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

118.  Fu, R., Gillen, A.E., Sheridan, R.M., Tian, C., Daya, M., Hao, Y., Hesselberth,

J.R., and Riemondy, K.A. (2020). clustifyr: an R package for automated single-cell

RNA sequencing cluster classification. F1000Res *9*, 223.

https://doi.org/10.12688/f1000research.22969.2.

119.  Zhang, B., and Horvath, S. (2005). A general framework for weighted gene co-

expression network analysis. Stat Appl Genet Mol Biol *4*, Article17.

https://doi.org/10.2202/1544-6115.1128.

120.  Langfelder, P., Luo, R., Oldham, M.C., and Horvath, S. (2011). Is my network

module preserved and reproducible? PLoS Comput Biol *7*, e1001057.

https://doi.org/10.1371/journal.pcbi.1001057.

121.  Gandrillon, O., Solari, F., Legrand, C., Jurdic, P., and Samarut, J. (1996). A rapid

and convenient method to prepare DIG-labelled RNA probes for use in non-

radioactive in situ hybridization. Molecular and cellular probes *10*, 51–56.

1603

# Figure 1: Oxtr reduces the amount of time naïve animals take to show partner preference

**Figure 2: Oxtr suppresses promiscuous behaviors in a state dependent manner in females**

# Figure 3: Oxtr function influences different prosocial behaviors across pair bonding in males

# Figure 4: Oxtr sex-specifically influences social interactions between potential mates

# Figure 5: Post pair bonding gene expression is driven by female specific changes dependent on Oxtr signaling

# Figure 6: Oxtr regulated genes are common to social behavior relevant processes across species and are similarly regulated across brain regions

# Figure 7: Loss of Oxtr results in the decrease in density of Oxt and Avp expressing neurons in the PVN