Oxytocin receptor controls promiscuity and development in prairie voles

Ruchira Sharma[1,2,3,4†], Kristen M. Berendzen[1,2,3,4†], Amanda Everitt[1,3,10,11], Belinda Wang[1,3], Gina Williams[1,2,3,4,5], Shuyu Wang[1,2,3,4], Kara Quine[1,2,3,4], Rose D. Larios[1,2,3,4,5,14], Kimberly L. P. Long[1,2,3,4], Nerissa Hoglen[1,2,3,4,5], Bibi Alika Sulaman[1,2,3,4,15], Marie C. Heath[1,2,3,4,16], Michael Sherman[1,2,3,4], Robert Klinkel[1,2,3,4,17], Angela Cai[1,2,3,4,20], Denis Galo[1,2,3,4,], Lizandro Chan Caamal[1,2,3,4,18], Nastacia L. Goodwin[1,2,3,4,19], Annaliese Beery[6,7], Karen L. Bales[8,9], Katherine S. Pollard[10,11,12], Arthur Jeremy Willsey[1,13], Devanand S. Manoli[1,2,3,4,*]

[1]Department of Psychiatry and Behavioral Sciences

[2]Center for Integrative Neuroscience

[3]Weill Institute for Neurosciences

[4]Kavli Institute for Fundamental Neuroscience

[5]Neurosciences Graduate Program, University of California, San Francisco; San Francisco, CA 95158, USA

[6]Department of Integrative Biology

[7]Department of Neuroscience, UC Berkeley, Berkeley, CA, United States

[8]Department of Psychology

[9]Department of Neurobiology, Physiology, and Behavior, University of California, Davis; Davis, CA 95616, USA

[10]Gladstone Institutes, San Francisco, CA, USA

[11]Department of Epidemiology & Biostatistics, University of California, San Francisco,

San Francisco, CA, USA

[12]Chan Zuckerberg Biohub, San Francisco, CA, USA

[13]Calico Labs, San Francisco, CA 94080, USA

[14]Neurona Therapeutics, South San Francisco, CA 94080, USA

[15]Department of Neuroscience, University of Michigan, Ann Arbor, MI 48109, USA

[16]Department of Neurobiology and Behavior, University of California, Irvine; Irvine, CA

92697, USA

[17]Branch Out School, Los Angeles, CA 90026, USA

[18]University of California, Los Angeles; Los Angeles, CA 90095, USA

[19]Department of Psychology, University of Washington; Seattle, WA 98195, USA

[20]Department of Rehabilitation and Regenerative Medicine, Columbia University Irving

Medical Center; New York, NY 10032, USA

* Corresponding authors. Devanand.manoli@ucsf.edu

Lead Contact: Devanand.manoli@ucsf.edu

†These authors contributed equally to this work

Summary:

Oxytocin receptor (Oxtr) signaling influences complex social behaviors in diverse species,

including social monogamy in prairie voles. How Oxtr regulates specific components of

social attachment behaviors and the neural mechanisms mediating them remains

unknown. Here, we examine prairie voles lacking Oxtr and demonstrate that pair bonding

comprises distinct behavioral modules: the preference for a bonded partner, and the

rejection of novel potential mates. Our longitudinal study of social attachment shows that

Oxtr sex-specifically influences early interactions between novel partners facilitating the

formation of partner preference. Additionally, Oxtr suppresses promiscuity towards novel

potential mates following pair bonding, contributing to rejection. Oxtr function regulates

coordinated patterns of gene expression in regions implicated in attachment behaviors

and regulates the expression of oxytocin in the paraventricular nucleus of the

hypothalamus, a principal source of oxytocin. Thus, Oxtr controls genetically separable

components of pair bonding behaviors and coordinates development of the neural

substrates of attachment.

Keywords:

Attachment behaviors, prairie vole, pair-bonding, oxytocin receptor, partner preference,

monogamy, promiscuity, oxytocin signaling, paraventricular nucleus of the

hypothalamus, nucleus accumbens

Introduction

Social attachments, such as the bonds between parents and children, family and kin, friends, and the long-term relationships between mates and romantic partners form the basis of interpersonal and group dynamics within societies[1–7]. Across mammals and other taxa, various selective pressures have given rise to diverse patterns of complex social organization involving long-term relationships between members of a species, one of the most robust and intriguing of which is the enduring bond between mates[2,4,8–13]. In mammals, ~4% of species demonstrate social monogamy, forming persistent pair bonds between mating partners[9,14–16]. Comparative studies of closely related species that display distinct patterns of social behaviors, including social monogamy, reveal correlated species-specific differences in the nonapeptide hormones oxytocin (Oxt) and arginine vasopressin (Avp), or their homologs, suggesting that these hormones serve as critical modulators of affiliative behaviors and pair bonding[9–11,16–20]. Despite the central role of social attachment in human behavior and a rich history of investigation into the cognitive processes underlying social cognition and affiliative behaviors, it has been difficult to probe the neural mechanisms underlying these fascinating behaviors at the cellular, molecular, and genetic level, as most species that serve as model organisms for such studies in the laboratory do not display long-term social attachment as adults [6,21–27].

Prairie voles (*Microtus ochrogaster*) display long-term social attachment such that mating partners show an enduring pair bond and social monogamy[13,28–31]. Like many complex innate behaviors, pair bonding consists of distinct modules: the development of increased prosocial behaviors with partners over strangers (partner preference), and increased

agonistic behaviors towards novel opposite sex potential mates (stranger rejection)[13,30,32–

37]. As in humans and other species that show affiliative social behaviors, disruption of pair

bonds results in increased anxiety-like behaviors and markers of stress, supporting

integrated neural and physiologic mechanisms that facilitate the preservation of such

attachments[38–40]. Pioneering work in prairie voles identified Oxt and Avp as critical

mediators of pair bonding[41,42]. Interspecific variations in the patterns of expression of the

oxytocin receptor (Oxtr) and the vasopressin 1a receptor (V1ar) correlate with the

potential for pair bonding within closely related vole species[43–49].

Across phyla, manipulations of oxytocin or its species-specific homolog alters social

behaviors. In prairie voles, pharmacologic inhibition of Oxt and Avp signaling via their

respective cognate receptors disrupts pair bonding, while exogenous administration of

these hormones promotes pair bonding without mating[28,50,51]. Transient blockade of Oxtr

only during the first few hours of cohabitation delays the formation of a partner

preference[28,29] and artificial overexpression of Oxtr accelerates the formation of a pair

bond[52]. Exogenous oxytocin administration leads to to increased prosocial behaviors

towards conspecifics as well as humans in domestic dogs, initiation of huddling towards

a potential partner in marmosets (*Callithrix penicillata*), and food sharing with conspecifics

in pinyon jays (*Gymnorhinus cyanocephalus*, a highly social corvid species)[53–56].

Conversely, OT receptor antagonists administered during early pair-bond formation to

monogamous cichlid fish (*Amatitlania nigrofasciata*), socially monogamous zebra finch

(*Taeniopygia guttata*) and marmosets (*Callithrix penicillata*) lead to reduced affiliative

behaviors towards potential mates[55,57,58]. Taken together, these studies underscore the

broad conservation of this neuropeptide in the modulation of social behaviors, as well an

early temporal window during which oxytocin-dependent behaviors mediate partner

preference.

Given the central role of Oxt signaling in affiliative behaviors, we tested the genetic

requirement for signaling via Oxtr for pair bonding and social behaviors in prairie voles.

We generated prairie voles lacking Oxtr and unexpectedly observed that these animals

continue to display a strong preference for partners following mating and cohabitation[59].

To gain deeper insight into the modulation of attachment and social behaviors by Oxtr

function, we developed a series of behavioral assays to probe distinct aspects of pair

bonding and associated social behaviors. To interrogate the molecular mechanisms

mediating social attachment, we profiled patterns of gene expression in the nucleus

accumbens (NAc), a key site of species differences associated with social monogamy

that is dramatically enriched for Oxt binding in prairie voles[45,60,61]. We find that Oxtr

controls the timing of pair bonding in prairie voles and promiscuity independent of partner

preference formation. Oxtr sex-specifically influences the behavior of potential mates,

increasing agonistic displays in naive wildtype (WT) females, evidenced by decreased

aggression and increased prosocial interactions shown by potential female WT mates in

the presence of a male lacking Oxtr. Oxtr function regulates patterns of gene expression

in the NAc, as well as other regions of the vole brain implicated in social and attachment

behaviors, in distinct ways in each sex. Furthermore, we see changes in expression of

genes regulating neurodevelopmental processes and disease, supporting a role for Oxtr

function early in life[62–64]. Finally, consistent with potential trophic effects of Oxtr function

during development, we find that loss of Oxtr alters patterns of Oxt and Avp expression

in the paraventricular nucleus, a principal source of these neuropeptides[63,65]. Our studies

reveal genetically separable components of pair bonding and suggest that Oxtr signaling

sex-specifically participates in the development of the circuits underlying attachment

behaviors.

Results

Oxtr function facilitates the formation of partner preference

We previously found that partner preference formation, following 1 week of cohabitation,

can occur in the absence of Oxtr function[59]. However, acute modulation of Oxtr signaling

can alter patterns of social behaviors between individuals[29,50,66]. These observations

suggest that changes in Oxtr function may influence early social interactions that

contribute to the trajectory of pair bond formation and/or specific aspects of attachment

behaviors following the formation of partner preference. We first wished to determine

whether Oxtr function facilitated pair bond formation by examining the display of partner

preference following shorter periods of cohabitation. Consistent with previous studies

showing that the cohabitation time required for the display of partner preference differs

between males and females, we find that WT females demonstrate partner preference

after only 6 hours of cohousing with a WT male partner, while WT males require 5 days

following introduction to a potential WT mate (Fig. S1A-C)[67]. We therefore cohoused WT

or Oxtr$^{1-/-}$ females with WT males for 6 hours, and WT or Oxtr$^{1-/-}$ males with WT females

for 5 days following introduction (short cohabitation), or in these combinations for 1 or 7

days (long cohabitation) for females or males, respectively (Fig. 1A, Fig. S1A).

Both males and females lacking Oxtr show an absence of partner preference following

short cohabitation with WT partners (Fig. 1B). Oxtr$^{1-/-}$ females, but not males, showed a

decrease in the fraction of huddling time with their partner and an increase in the fraction

of huddling time with a stranger compared to WT controls (Fig. 1C-D). Oxtr$^{1-/-}$ males

present an absence of partner preference (Fig. 1B), resulting from a trend towards

decreased time with their cohoused partner and increased time with strangers (Fig. 1C-

D). Consistent with our previous studies, we find no difference between WT and Oxtr$^{1-/-}$

females when the pairs are cohoused for the longer duration (Fig. S1B-F).

To better understand the social interaction dynamics underlying differences in huddling

behavior between WT animals and their siblings lacking Oxtr after short-term

cohabitation, we quantified the cumulative time subjects spent huddling with either their

cohoused partners or strangers across the assay.Oxtr$^{1-/-}$ females show a significant

decrease in the amount of time spent huddling with partners in the last 30 mins of the

assay (150-180mins) (Fig. 1E). We also observe a significant increase in huddling time

of Oxtr$^{1-/-}$ males with strangers during the middle periods (75 – 130mins) (Fig. 1F),

suggesting that they engage in more early prosocial behavior with strangers than WT

males. Consistent with our previous work, we find no difference between cumulative

huddling times with either partner or stranger between WTs and Oxtr$^{1-/-}$ animals of both

sexes after long cohabitation periods (Fig. 1G-H)[45]. These observations demonstrate that

Oxtr function changes the temporal dynamics of social interactions with both familiar and

unfamiliar animal in a sex-specific manner. Taken together, our findings suggest that Oxtr

function acts early in the process of pair bonding to facilitate the formation of the partner

preference.

Oxtr suppresses promiscuous behaviors in a state dependent manner in females

Oxtr influences the dynamics of interactions with partners and strangers differently

between the sexes. We therefore wished to examine the detailed patterns of these

interactions separately in females and males (Fig. 2A). We examined the distribution of

huddles by WT vs Oxtr[1-/-] females following short cohabitation as a measure of prosocial

behavior displayed by these animals. We find that the top quartile of huddle durations

(long huddles) from each subject animal make up an average of ~90% of the total huddle

duration, functionally separating these interactions from incidental side-by-side contact

that may not constitute affiliative huddling (Fig. S2A Left). We classified long huddles as

partner huddles or stranger huddles and discovered that, consistent with the overall time

spent, the median huddle duration carried out by Oxtr[1-/-] females is significantly shorter

with partners and significantly longer with strangers, compared to WT females (Fig. 2B).

The longest huddle displayed with partner is significantly longer for WT females compared

to Oxtr[1-/-] females though we find no difference in the length of the overall longest huddle.

This is due to some Oxtr[1-/-] females conducting their longest huddle with the stranger

instead (Fig. S2B). Oxtr[1-/-] females additionally executed a larger number of huddles only

with the stranger, but not the partner, compared to WT controls (Fig. 2C). Taken together,

these results suggest that Oxtr function in females both facilitates prosocial behavior with

partners and suppresses such behavior with strangers during the early stages of pair

bond formation.

We next wished to determine whether the loss of Oxtr function influences patterns of prosocial behaviors after the display of partner preference. We therefore examined patterns of huddling, in particular the duration and frequency of these interactions, following long cohabitation, a time point at which both WT and Oxtr[1-/-] animals display robust preference for partners over strangers[59] (Fig. S1D). As with short cohabitation, the top quartile of huddles (by duration) for each subject animal following long cohabitation, described an average of 90% of total huddle time (Fig. S2A right). Following long cohabitation, preference and huddling behavior in Oxtr[1-/-] females is indistinguishable from their WT siblings (Fig. 2D-E). Comparing changes in the frequency of top quartile huddles displayed by Oxtr[1-/-] females after short vs long cohabitation, we find that both WT and Oxtr[1-/-] females show decreased huddle frequency with stranger following long cohabitation (Fig. S2C-D). On comparing durations of long huddles, we find that Oxtr[1-/-] females significantly increase their median huddle duration with partner post long cohousing (Fig. 2F-G). Thus, loss of Oxtr function in females reduces the duration of early prosocial interactions (short cohabitation) with partners and strangers while Oxtr-independent mechanisms eventually promote WT patterns of affiliative huddling and partner preference following longer periods of cohabitation.

Our results suggest that Oxtr function influences specific aspects of early pair bonding and attachment behaviors. We next wished to dissociate the influence of Oxtr on the formation and display of partner preference from the display of agonistic rejection behaviors towards novel strangers following cohabitation. We therefore developed a

battery of assays to examine initial social interactions between mating partners at early

and later stages of pair bonding, as well as stranger rejection following cohabitation

(Fig.2A). We examined the behavior of subject females from each genotype with WT

males when naive animals were first introduced to each other (Introduction), mating

behaviors following estrus induction (Timed Mating), prosocial interactions following

reunification with their bonded mate (Partner Reunification), and agonistic rejection of

novel, opposite sex strangers (Stranger Rejection) (Fig. 2A see Methods). We find no

differences in social behaviors during Introduction, Timed Mating and Partner

Reunification (Fig. 2H-J). In contrast, during Stranger Rejection, $Oxtr^{1-/-}$ females show

more social behaviors towards novel naive males than WT females do in the absence of

their partner (Fig. 2K). Taken together, these studies demonstrate that Oxtr function

regulates specific components of pair bonding in females in a state-dependent manner,

suppressing promiscuous social interactions with stranger males after cohabitation

sufficient for the display of partner preference.

Oxtr function influences different prosocial behaviors across the stages of pair

bonding in males

Given the Oxtr-dependent difference in huddling behavior between males and females

after short cohabitation (Fig. 1C-D), we wished to assess sex differences in the patterns

of behaviors displayed by $Oxtr^{1-/-}$ animals and to further investigate Oxtr function in males

(Fig. 3A). Examining long huddles (as above, Fig. 2), we find that such huddles by $Oxtr^{1-/-}$

males with their partners following short cohabitation are significantly shorter when

compared to WT siblings, contributing to diminished partner preference early in pair

bonding (Fig. 3B Left). Interestingly, while Oxtr[1-/-] males show no difference in the total

amount of time spent huddling with strangers when compared to WT siblings (Fig. 1D),

they display shorter and more frequent huddles towards them at this time (Fig. 3B, 3C

Right). In males, Oxtr thus influences the pattern as well as amount of early affiliative

behavior to distinct behavioral stimuli in different ways.

Oxtr[1-/-] males show no difference in time spent with either partners or strangers after long

cohabitation (Fig. S1E-F). However, these males show a persistent reduction in the

duration of their long huddles with strangers at this time point (Fig. 3D). Following long

cohabitation, WT and Oxtr[1-/-] males show no difference in the duration of long huddles

with partners and a reduction in the duration of long huddles with strangers (Fig. S3A-B).

We find no difference in the frequency of long huddles between Oxtr[1-/-] and WT males

after long cohabitation (Fig. 3E). Consistently, comparing Oxtr[1-/-] males' behavior following

short vs long cohabitation reveals that the frequency with which Oxtr[1-/-] males huddle with

strangers decreases following long cohabitation (Fig. 3F-G). Thus, specifically in males,

loss of Oxtr function does not change the overall time spent huddling with strangers after

short or long cohabitation but does cause a continuing disruption in the patterns of

affiliative behavior that comprises huddling with strangers.

Our observations suggest that the loss of Oxtr disrupts certain characteristics of prosocial

behaviors in males regardless of the time spent bonding with their partner. We next

wished to dissect whether such loss of function resulted in atypical affiliative or agonistic

behaviors across stages of bonding. We examined the behavior of Oxtr[1-/-] males and their

WT siblings in the same series of paradigms described previously to characterize the

formation and maintenance of their pair bonds (Fig. 3A). Oxtr$^{1-/-}$ males spend similar

amounts of time as WT males engaging in prosocial behaviors with partners regardless

of bonding state (Fig. 3H-J, S3E). Similar to our observations in Oxtr$^{1-/-}$ females, Oxtr$^{1-/-}$

males spend more time engaging in prosocial behaviors with strangers in the absence of

their partners (Fig. 3K).

In contrast to time spent conducting mating behaviors with WT female partners (Fig. S3E),

Oxtr$^{1-/-}$ males demonstrate increased frequency of mating attempts when these females

are in estrus (Fig. 3I). Examining all instances of mounting, we find that Oxtr$^{1-/-}$ males

carry out shorter mounting bouts more frequently than WT males (Fig. S3F). To determine

whether the shorter bouts of mounting in Oxtr$^{1-/-}$ males influence the receptivity of their

partners, we compared the mount-to-intromission ratio for each animal[68] and find that

females are equally receptive to Oxtr$^{1-/-}$ males as their WT siblings (Fig. S3G). Thus,

consistent with our previous observations, Oxtr influences patterns, but not the amount

or success, of male mating behaviors, suggesting that such changes in males' behaviors

are equally able to elicit receptivity in females.

Oxtr function influences distinct components of males' attachment-related social

behaviors towards unfamiliar females at different stages of pair-bonding, lengthening

mating bout durations when naïve and suppressing promiscuous prosocial behaviors

upon bonding. Taken together our results suggest that Oxtr influences patterns of social

behaviors with future partners during the early phases of pair-bonding differently in

females and males, but upon bond formation, functions in both sexes to suppress

promiscuous affiliative behaviors towards strangers (Fig. S3H).

Oxtr sex-specifically influences social interactions between potential mates

Oxtr function influences patterns of social interactions between mates early during the

formation of a pair-bond (Short Cohabitation in both sexes, Introduction and Timed Mating

in males). We therefore investigated the effect of Oxtr loss on the choice made by a

potential mate. We developed a variation of the PPT three-chamber assay to present a

naive WT "chooser" with a choice between a naive opposite sex WT or naive opposite

sex $Oxtr^{1-/-}$ animal (Fig. 4A, Fig. S4A, see Methods). Using the "naïve-choice" paradigm

we find that, over the course of 6 hours, WT choosers spend significantly more time with

one of the two stimulus animals (winner) (Fig. 4B) and that WT and $Oxtr^{1-/-}$ animals are

chosen as the winner with equal frequency(Fig. 4C).

To examine whether Oxtr influences the dynamics of the choice made by a potential WT

mate, we compared the time spent with the winner from a lane where the chooser had a

choice between a naïve WT or $Oxtr^{1-/-}$ animal (WTOxtr lanes) versus that of a chooser

choosing between 2 naive WT animals (WTWT lanes) of the opposite sex (Fig. 4D). While

naïve WT choosers show no difference in overall social choice between naive WT and

$Oxtr^{1-/-}$ opposite sex animals (Fig. 4C), WT female choosers show distinct behavioral

dynamics in the presence of an $Oxtr^{1-/-}$ male (WTOxtr lane). In the presence of an Oxtr

mutant (WTOxtr lane), WT females spend significantly more time in contact with the

winner, regardless of genotype, when compared to contact with the winner between two

WT males (WTWT lane) (Fig. 4E). In contrast, we observe no difference in contact time

with a winner when a naive WT male is choosing in WTOxtr versus WTWT lane (Fig. 4E).

We therefore manually examined the specific behaviors shown by female WT choosers

during their interactions with both the winner and loser in WTWT and WTOxtr lanes.

Preference indices using these metrics were highly concordant with automated scoring

(Fig. S4A). The time spent by the female chooser displaying prosocial behaviors towards

the winner and the time spent huddling with the winner in WTOxtr lanes was significantly

greater than WTWT lanes (Fig. S4B) (Fig. 4E). The divergence between the amount of

cumulative time spent huddling with the winner from WTOxtr versus WTWT lanes occurs

after 5 hours have elapsed (313-360 mins, Fig. 4F), suggesting that complex interactions

in the presence of a male lacking Oxtr influence how females "choose" a male partner

regardless of whether they end up choosing a WT or Oxtr$^{1-/-}$ winner.

We evaluated the duration of the longest huddles conducted by all WT female choosers,

and find that the longest huddle is always conducted with the winner. The longest huddles

conducted in the WTOxtr lanes, regardless of the genotype of the male winner, are greater

than those from WTWT lanes (Fig. 4G). There is no difference in the latency to start the

longest huddle between WTOxtr and WTWT lanes (Fig. S4C). Thus, the presence of the

Oxtr$^{1-/-}$ male in the lane leads a WT chooser female to execute the longest huddle for a

greater amount of time compared to the longest huddle by the WT female chooser from

a WTWT lane (Fig. 4G). These observations suggests either that naïve Oxtr$^{1-/-}$ males

elicit longer longest-huddle bouts from WT females, even when this huddle occurs with a

WT male winner, or that the behavior of naïve WT males reduces the duration of WT

females' longest huddle - a reduction that is absent in the presence of a male lacking Oxtr.

To determine if specific patterns of behavior preceded the longest huddle with a winner,

we aligned the onset of the longest huddle for all the WTOxtr and WTWT lanes and

evaluated the behaviors performed 15 mins prior to its initiation. We find that the duration

of sniffs (anogenital investigation, see methods) by WT female choosers in WTWT lanes

is greater than those by choosers in WTOxtr lanes (Fig. 4H). We did not find any

difference in the duration of sniffs specifically with the winner or the loser when comparing

WTWT and WTOxtr lanes, or in the total time spent sniffing over the course of the assay

(Fig S4D, E). Thus, the interactions with both winner and loser contribute to the increase

in duration spent sniffing 15 minutes prior to the longest huddle in WTWT lanes. These

observations may suggest that a female chooser from the WTWT lane continues to seek

chemosensory input before the longest huddle and these huddles are shorter due to

ambivalence in a choice between two WT males. In contrast, a WT female chooser in a

WTOxtr lane may encounter larger differences between a WT versus Oxtr$^{1-/-}$ male, leading

to a display of more prosocial behavior towards the chosen male independent of the

winner's Oxtr function.

We next examined antagonistic behaviors displayed during the naive choice assay as

such behaviors may reduce prosocial behaviors displayed by chooser females. We found

no difference in the overall counts of aggressive behaviors between WTWT and WTOxtr

lanes (Fig. S4F). However, WT female choosers attack the loser male in the WTWT lanes

more compared to WTOxtr lanes (Fig. 4I, Fig. S4G). Thus, in WTWT lanes, winner

selection may be influenced by the level of aggression elicited by naïve WT males, a

factor that is lower in WTOxtr lanes. We find a correlation between the amount of time

spent socializing (with both winner and loser) and the count of all the aggression during

the assay (Fig. 4J). Female choosers from WTOxtr lanes tend to spend more time

engaged in prosocial behaviors with the winner (Fig. 4E-F, S4B) and display inceased

prosocial and reduced aggressive behaviors. Female choosers choosing between two

WT males, however, display either high prosocial and low aggressive behaviors or low

prosocial behavior and high aggression. These observations suggest that the presence

of a second male may influence the behavioral state of a WT female. The presence of

only WT males may induce aggression by females against eventual losers, while such

displays are absent in the presence of a male lacking Oxtr regardless of a female's choice.

Given the differences in the influence of the presence Oxtr$^{1-/-}$ males on displays of

aggression by WT females when choosing between males, we examined antagonistic

behaviors exhibited during interactions between naive (Introduction) or bonded (Stranger

Rejection) females with WT or Oxtr$^{1-/-}$ males in the absence of a third animal. Strikingly,

we find that counts of aggressive behaviors were elevated only when naive WT females

encountered WT males compared to Oxtr$^{1-/-}$ males, while this difference is lost following

pair bonding (Fig. 4K). In contrast, we find no differences in aggressive behaviors when

comparing naïve or bonded WT and Oxtr$^{1-/-}$ females (Fig. S4H). Taken together, these

data suggest sex- and state-specific roles for Oxtr function. Specifically, in naive males,

Oxtr function induces displays of aggression by naive females, and the presence of a

male lacking Oxtr diminishes a naïve WT female's propensity to display agonistic

behaviors even in the presence of a WT male. Following bonding, however, the absence

of Oxtr function in males does not appear to influence the behavior of WT females despite

the promiscuous prosocial displays such mutant males.

Finally, to determine if naive prairie voles lacking Oxtr display generalized changes to

their behaviors independent of attachment-associated social behavior, we examined

general pro-social, anxiety-related, and locomotor behaviors. With the exception of an

increase in locomotion displayed by Oxtr$^{1-/-}$ females when compared to WT females in

an open field paradigm, we found no differences between Oxtr$^{1-/-}$ and WT voles of either

sex (Fig. S4I).

We have examined social interactions during the initial contact between mates, the

early stages of pair bonding, and during social attachment behaviors following partner

preference formation. Consistent with our findings and previous work, Oxtr function is

not required for the eventual display of partner preference. Here, we demonstrate that

Oxtr function sex-specifically facilitates early social interactions that contribute to the

formation of partner preference, and controls promiscuity.

Oxtr controls the molecular signature of pair bonding in the nucleus accumbens

While partner preference formation can occur in the absence of Oxtr function, we find that

it suppresses prosocial behaviors towards novel strangers following pair bonding. These

results suggest that distinct components of pair bonding are differentially influenced by

Oxtr function or compensated upon its loss. We next characterized the molecular

signature associated with pair bonding to determine whether changes in gene expression

following pair bond formation are controlled by Oxtr. Comparative studies demonstrate

that Oxt binding in multiple brain regions differs significantly between promiscuous and

monogamous rodent species. Oxt binding is particularly enriched in the NAc in prairie

voles, while vasopressin binding is largely absent in this area[45,47]. Thus, analyzing gene

expression in the NAc allows us to identify bonding-related expression that is primarily

due to Oxtr function and largely independent of the direct consequences of AVP signaling.

Additionally, given the well-established role of the NAc in the association of reward or

aversion with specific sensory and environmental cues[69–71] this region is well poised to

play a role in the development of partner preference, even without Oxtr signaling, as well

as stranger rejection, which appears to be dependent upon Oxtr function.

We performed bulk RNA-sequencing on NAc tissue from WT and Oxtr[1-/-] animals of both

sexes that were either: group housed with same-sex peers and sexually naïve (pre-

pairing); or paired with WT opposite-sex partners for six days, using our timed mating

paradigm (post-pairing) (Fig. 5A). PCA showed robust separation of WT and Oxtr[1-/-]

samples (Fig. S5A). Differential expression (DE) analysis of pairing condition, regardless

of genotype or sex, identified 13 genes that show significant changes in expression post-

pairing (Fig. S5B), all of which showed decreased expression (Fig. 5B). These genes

include *Agt, Sgk1, Ptgds, Fosb, Pdk4*, and *Acsm5* (Fig. 5C) and, while these genes have

not previously been linked to shared gene regulatory or functional pathways, the predicted

protein-protein interaction network for the 13 DE genes shows more connections than

would be expected by chance (p=2.47 e^{-05}) (Fig. S5C). Intriguingly, even in the limited set

of genes with altered expression in the context of pair bonding, changes are driven

primarily by expression in WT females (Fig. 5C, S5D). Subgroup analyses by genetic

background and sex identified additional genes including *Adora2a, Fkbp5*, and *Spred3* in

WT females and collagen-related genes, including *Col4a1* and *Col4a2*, in Oxtr$^{1-/-}$ females

(Fig. 5D, S5D). Only one gene, *Sema3b,* is significantly different in Oxtr$^{1-/-}$ males following

pairing (Fig. 5D).

To further examine changes in patterns of gene expression in the NAc associated with

pair bonding, we performed weighted gene coexpression network analysis (WGCNA)[72]

and calculated three sets of modules (M) from different sample subsets (modules^{WT} ,

modules^{Mut}, and modules^{All}) (Fig. S5F, S6D). Using the modules^{WT} subset, we identified

15 coexpression modules (Fig. S5F, Table 2) representing genes that share highly similar

expression patterns within the NAc. We identified one module (Mpink^{WT}) that significantly

and specifically correlated with pre-pairing status in WT animals (Pearson R correlation

of 0.7, Bonferroni adjusted p-value (p$_{adj}$)=0.018) (Fig. 5E, S5E-F). Genes with high

module membership for the Mpink^{WT} are also those with high gene significance for pairing

condition supporting the module's association with behavior condition (Fig. 5SG). Further,

genes significantly differentially expressed with bonding are significantly enriched in

Mpink^{WT} (Bonferroni p$_{adj}$=7.44 e^{-08}) and serve as hub genes within this module (Fig. S5E,

G). These results suggest that the Mpink^{WT} module might be driving, or under the

influence of pathways driving, the pre-pairing to post-pairing state change.

We asked whether any of the modules[WT] identified were regulated by Oxtr and thus lost

in Oxtr null animals. To address this, we assessed how well modules[WT] are preserved in

mutant samples. We calculated a $Z_{summary}$ measure that combines module density and

intramodular connectivity metrics into a composite statistic. We find that of the modules[WT],

Mpink[WT] had the lowest $Z_{summary}$ ($Z_{summary}$ = 2) in Oxtr[1-/-] samples (Fig. 5F), indicating that

dynamics of gene expression that correlate with bonding states are lost in the absence of

Oxtr. Finally, the module eigengene (ME) for Mpink[WT] when calculated for all 30 samples

and compared between pre- and post-pairing, was significant only for WT females

(Wilcoxon rank sum, Benjamini-Hochberg p_{adj}=0.03) and not for WT males or Oxtr[1-/-]

animals of either sex (Fig. 5G). These observations suggest there is sex-specific,

coordinated regulation of gene expression in the NAc following pairing of mates that may

be associated with pair bond associated behaviors.

Loss of Oxtr sex-specifically disrupts patterns of gene expression associated

with neurodevelopmental disorders

We next wished to determine how Oxtr signaling contributes to changes in gene

expression in the NAc associated with the naive state or the process of pair bonding (Fig.

6A) by comparing the transcriptional profile of WT and Oxtr[1-/-] animals of each sex. Given

that our lines are maintained on an outbred background, in order to minimize sample

variability, tissue from three animals was pooled for each sample. We observed significant

genotype specific differences in gene expression in animals from both sexes, with 466

genes showing differences between WT and Oxtr[1-/-] females, 270 between WT and Oxtr[1-]

[/-] males, and 1,014 genes when samples from both sexes were combined (using a

p_{adj}<0.05 and absolute ($\log_2$(FC)) > 0.25 as criteria). Of the combined DE gene set, 370

genes are upregulated and 644 are downregulated (Fig. 6B, S6A-B). Genes with the

greatest fold change across both sexes include *Slc2a3, Kcna1, Rgs8, Cnksr2, Acvr*, and

Cnot6 (Fig. 6B). Genes specifically downregulated in Oxtr[1-/-] females with $\log_2$(FC) less

than 1 include *Abcb11* (a gene recently associated with treatment resistant

schizophrenia[73]), *Ngfr, Pon3,* and *Shh*. While there was no significant difference in Oxtr

transcript levels by genotype (p_{adj} = 0.753)—consistent with minimal nonsense-mediated

decay despite a premature stop codon—we have previously shown that no functional

protein is produced in Oxtr[1-/-] animals via autoradiography[59]. V1aR showed a modest

decrease in expression in Oxtr[1-/-] animals (p_{adj}=0.03) (Fig. S6C), though this is likely

influenced by variability in WT samples. WGCNA of all samples identified two modules

(Mbrown[All] and Mturquoise[All]) that were significantly positively correlated with the WT

genotype compared to mutant (Pearson R = 0.55, p_{adj} = 0.011 for brown and Pearson R

= 0.8, p_{adj} = 6.6e-7 for turquoise) (Fig. S6D). None of the identified modules showed

correlation with pairing condition (Fig. S6D). Our results thus reveal a dramatic change in

patterns of gene expression in the NAc with global loss of Oxtr signaling, independent of

bonding state.

Examination of the sets of genes whose expression is altered by the loss of Oxtr,

regardless of bonding state or sex, revealed several distinct cellular processes regulated

by Oxtr signaling. These processes comprise larger domains of cell function including

extracellular channel binding activity, axoneme assembly complex, axon glial

ensheathment development, and metal cation transport (Fig. 6C), as well as pathways

for G protein signaling and dopaminergic neurogenesis (Fig. S6E). When analyzed by

sex, more of the identified processes were enriched in female Oxtr gene sets than in male

ones (female: 19/62, male: 6/62). To further link differentially expressed genes to

behavioral traits, we asked whether genes altered by loss of Oxtr are selectively enriched

for high-confidence associations with autism spectrum disorders (ASD) and other

neurodevelopmental diseases (NDD) affecting social attachment in humans as compared

to epilepsy, a disorder that does not disrupt social attachment[74–77]. Consistently, we find

that the genes upregulated with loss of Oxtr ($p_{adj}<0.01$, $\log_2(FC)>0$) are specifically

enriched for ASD- and NDD-associations but not epilepsy associations (Fig. 6D "Joint").

This enrichment is again primarily driven by genes upregulated in females and not males

(Fig. 6D). Thus, similar to the factors that influence gene expression changes associated

with pair bonding, many of the changes we identify in the NAc resulting from loss of Oxtr

are primarily driven by female-specific effects that may play a conserved role in mediating

social attachment in humans.

Oxtr regulates coordinated gene expression across multiple neural populations

implicated in social attachment behaviors

Given the expression of Oxtr in regions of the brain known to mediate social behavior in

voles and other species[78,79], we wished to determine if genes whose expression is altered

in the NAc by the loss of Oxtr were expressed not only in specific populations of cells

within this region, but in other regions of the prairie vole brain implicated in social and

attachment behaviors. We therefore examined the expression of 15 genes that showed

significant changes in expression in WT females when compared to those lacking Oxtr

using *in situ* hybridization (ISH) to both validate our RNA-seq findings as well as examine

the expression patterns of these genes. Several of these genes, such as

phosphodiesterase 10A (*Pde10A)* and diacylglycerol kinase beta (*Dgkb)*, show robust

expression throughout the striatum (data not shown), while genes such as the calcitonin

receptor (*Calcr*) and delta like non-canonical notch ligand 1 (*Dlk-1*) are expressed in

restricted populations of cells in the NAc (Fig. 6F-H, top panels). The expression of these

genes is significantly reduced in Oxtr[1-/-] females (Fig. 6E-H), supporting the specificity and

resolution of our RNA-seq findings.

In parallel to our examination of gene expression in the NAc, our ISH revealed that loss

of Oxtr in females altered the expression of the genes we examined in other regions of

the prairie vole brain. We examined the expression of these genes in the lateral septum

(LS), a region that has been implicated in agonistic and attachment behaviors in prairie

voles and other rodents and shows species dimorphic patterns of Oxt and Avp binding.

We observe that both *Calcr* and *Dlk-1* are decreased in the absence of Oxtr (Fig. 6G, H

middle panel) in the LS. We also find that expression of *Calcr* and *Dlk-1* is decreased in

the paraventricular nucleus of the hypothalamus (PVN), a principal source of Oxt and Avp

in the rodent brain which sends projections to the NAc (Fig. 6G-H bottom panels)[80–82].

These findings suggest that Oxtr function regulates patterns of gene expression in

multiple regions of the brain implicated in social and attachment behaviors in a

coordinated and region-specific manner.

Loss of Oxtr results in the decrease in density of Oxt and Avp expressing

neurons in the PVN

We find here that loss of Oxtr results in coordinated changes in gene expression in multiple neural populations, including the PVN. We therefore wished to determine if the signatures of gene expression regulated by Oxtr in the NAc in prairie voles (ONP set), regardless of sex and bonding state, were enriched for patterns of gene expression associated with distinct neural populations in other regions of the brain, particularly those implicated in social and attachment behaviors. We compared the ONP gene set to the molecular signatures of 33 hypothalamic subtypes previously identified by single-cell RNA sequencing in mice[83]. Of the 33 clusters examined, the four with highest overlap with the ONP gene set represent subsets of PVN neurons, in particular those that express Oxt and AVP (Fig. 7A, clusters 43 and 26), as well as PVN neurons that express *Dlk1* and *Calcr* (cluster 29), both of which show reduced expression in the NAc and PVN in females lacking Oxtr (Fig. 6G-H bottom panel).

We next sought to determine if the overlap between Oxtr regulated genes and gene expression patterns of neural populations in the PVN correlated with changes in the expression of neuropeptides within PVN itself. We examined patterns of Oxt and Avp expression using immunohistochemistry in naive adults and compared patterns of expression between WT animals of both sexes and those lacking Oxtr (Fig. 7B, Fig. S7A-B). We find that the density of both Oxt and Avp producing neurons is lower in Oxtr$^{1-/-}$ animals (Fig. 7C-E). To determine if the loss of Oxt or Avp expression was restricted to specific populations of PVN neurons, we examined the expression of these

peptides across the anterior-posterior axis of the PVN in both WT and Oxtr[1-/-] males and

females. We find that animals of both sexes show a preferential loss of Oxt and Avp

expressing neurons in the anterior PVN (Fig. 7F-G). Finally, we examined patterns of

Oxt and Avp over the course of development to determine when loss of Oxtr function

influences the development of their expression. We find that the density of Oxt neurons

at P0 (Fig. S7C-F) and both Oxt and Avp neurons at P21 is lower in Oxtr[1-/-] females

compared to age matched WT females (Fig. S7G-J). In contrast, we find that Oxt and

Avp neuron density loss occurs only in adult Oxtr[1-/-] males (Fig. 7D-G, Fig. S7B). These

observations suggest the coordinated, region-, and sex-specific influence of Oxtr

function on the development of neurons expressing Oxt (and Avp) within the PVN,

consistent with a trophic role for this pathway identified in mice[63,65,84,85].

Discussion

Robust reproductive behaviors are essential for survival. Accordingly, species have

evolved diverse strategies, including social monogamy, which mediate locating,

identifying, and attracting mates, facilitate reproduction, and promote the survival of

progeny[16,86]. Signaling via Oxtr influences a wide range of physiology and behavior

associated with social interactions and attachment between conspecifics, including pair

bonding between prairie vole mates[41,87]. Acute blockade of Oxtr signaling via central or

nucleus accumbens infusions of Oxtr antagonists prevents the formation of partner

preference[28,50,51,60,88]. In contrast, constitutive loss of Oxtr in prairie voles disrupts specific

components of pair bonding, while the capacity for partner preference ultimately remains

intact[59]. Similarly, Oxtr null mice show select behavioral deficits, including impaired

sociability, reduced preference for social novelty, and increased aggression towards a

novel intruder[89,90].Thus, like other circuit and neuromodulatory pathways across diverse

species, Oxtr likely functions in parallel to multiple genetically specified pathways to

influence specific behavioral modules that comprise complex behavioral states, including

social attachment.[33,35,91–97]. These actions are likely dependent on the developmental

context and engagement of potentially compensatory parallel pathways, leading to

divergent phenotypes depending on the timing and nature of Oxtr disruption.

Here we demonstrate that loss of Oxtr in prairie voles influences the dynamics of social

interactions between sexually naïve animals and WT potential mates in distinct social

contexts in sex-specific ways. The presence of a naive $Oxtr^{1-/-}$ male changes levels and

patterns of both early prosocial and antagonistic behaviors displayed by naive WT

females, supporting the role of Oxtr in controlling individuals' and reciprocal behavior [30].

Thus, Oxtr function orchestrates male social behaviors operating early in the sensitive

period of pair-bond formation and, through these behaviors or through additional sensory

cues yet to be identified (pheromone signatures or vocal communication), shapes

bonding.

Animals lacking Oxtr take longer to display partner preference, and despite strong partner

preference following longer cohabitation, demonstrate promiscuous prosocial behaviors

towards novel potential mates, including huddling with strangers. These observations are

consistent with a state-dependent role of Oxtr function, similar to findings in multiple

species, including humans[98–100]. Oxtr function may thus promote prosocial behavior in the

naïve state to facilitate attachment with an unfamiliar potential mate, but following pair

bond formation, suppresses further prosocial interactions to new individuals, or even

promotes further separation from and rejection of unfamiliar individuals or groups. Across

species and outside of the context of pair bonding, oxytocin signaling may play a

conserved role in mediating other state-specific aversive behavior, for example, defeat-

induced social avoidance learning[101].

Our findings reveal that patterns of social behaviors contributing to 1) pair bond formation,

2) partner preference, and 3) rejection of novel mates are genetically separable

components of pair bonding representing distinct behavioral modules regulated by

mechanisms that require Oxtr function, i.e. reciprocal behaviors and stranger rejection, or

those that can be displayed in its absence, i.e. partner preference. Further work will allow

us to determine the Oxtr-independent mechanisms that mediate the closure of an early

sensitive period and facilitate the display of partner preference[59]. Where Oxtr functions

to control the switch from prosocial behavior towards a potential mate in naïve animals to

the suppression of such behaviors towards, and active rejection of, opposite-sex

conspecific strangers following bonding remains to be determined.

Examination of the molecular signatures of pair bonding and the role of Oxtr in regulating

these changes in gene expression in the NAc reveals modest, Oxtr-dependent changes

in the expression of a specific network of genes in WT animals following pair bonding[45,102].

Intriguingly, loss of Oxtr preferentially impacts patterns of gene expression in the NAc in

females. The NAc demonstrates little endogenous sex steroid hormone receptor

expression difference in gene expression in prairie voles and other species[78,103–105]. Sex

differences in the effects of Oxtr on gene expression this region may therefore suggest

that there are significant sex-differences in the inputs into this region that act through Oxtr

during development[106–111]. The larger impact of loss of Oxtr in females is consistent with

its more general conservation in mammals to regulate aspects of female physiology such

as lactation and prosocial behaviors or sensory cues that promote interactions with

males[112].

Oxt and its receptor show species-specific changes in expression through embryonic

development and early postnatal life, impacting the development of cortical and

subcortical circuits[62,64,113,114]. Consistent with this, genes with altered expression in the

absence of Oxtr are enriched for genes associated with neuropsychiatric disorders, in

particular neurodevelopmental disorders and ASD. Given the sex-biased enrichment of

these genes, such developmental differences in the impacts of Oxtr function may

contribute to sex differences in the pathophysiology of these disorders[115–117]. Accordingly,

the molecular signatures of attachment in the NAc, both dependent upon and

independent of Oxtr function, are subtle relative to these sex-biased differences in the

role of Oxtr and possibly reside in other neural populations that function in attachment.

Validation of changes in gene expression in the NAc resulting from loss of Oxtr suggests

coordinated regulation of these genes by Oxtr across multiple brain regions implicated in

social and attachment behaviors. Oxtr may be one of multiple factors that organizes

specific components of the circuits underlying these displays to control discreet modules

of behavior. Consistent with a trophic role for Oxtr function during development, we find

that the genes regulated by Oxtr within the NAc are enriched for those that identify cell

types in the PVN, a principal source of neuropeptides implicated in pair bonding, which

sends oxytocinergic projections to the NAc[79,81,118–120]. Neonatal OT manipulations in mice

sex-specifically modulate levels of neuropeptides in PVN neurons, increasing Oxt

expression in females and reducing Avp expression in males[64]. We demonstrate that, in

prairie voles, loss of Oxtr preferentially impacts Oxt and Avp expression in the anterior

PVN, suggesting that specific populations of these cells may be more dependent upon

such trophic function for their development, survival, or expression of these hormones.

Whole brain analysis will likely reveal if such neurons have distinct projections when

compared to those that remain in the absence of Oxtr, contributing to our understanding

of how these circuits and pathways regulate distinct aspects of social and attachment

behaviors and physiology[4,87,121].

Taken together, our behavioral and molecular studies reveal previously unappreciated

nuances in the control of social behavior and pair bonding by Oxtr function. They uncover

intriguing sex differences in the patterns of behavior influenced by loss of Oxtr function.

These observations suggest that circuits with sex-specific patterns of development, or

even function in ancestral species, may mechanistically converge in the context of Oxtr

function to generate displays of reciprocal interactions, pair bonding and attachment

behaviors in WT prairie voles that are similar between males and females[122,123].

Manipulation of Oxtr expression at specific developmental timepoints and in specific

neural populations will help to elucidate how this pathway influences social and

attachment behaviors in the context of normal development to regulate distinct

components of pair bonding. Understanding the specific neural populations that express

Oxtr and other molecular pathways that contribute to pair bonding will allow us to

understand the mechanisms controlling the early experience-dependent, sensitive period

of bond formation. In parallel, examination of the activity in these populations following

manipulations of Oxtr function will allow us to determine the circuit mechanisms by which

similar but distinct sensory cues contribute to the identification of partners or novel

conspecifics and consequently, the display of dramatically different patterns of social

behaviors. The sum of these mechanisms represents the preference for partners over

strangers and reflects the formation and maintenance of pair bonds between mates,

affiliation between individuals, and the behavioral consequences of identifying others as

unfamiliar or outside of these bonds.

Acknowledgments

The authors would like to thank members of the Manoli lab for assistance, advice,

helpful discussion and comments on the manuscript. J. Tollkuhn and M. Brainard

provided valuable insights to the structure, content and design of the manuscript.

Additionally, we would like to thank S. Garg for his assistance in data analysis and help

designing the analytic pipeline. The authors received funding from National Institutes of

Health grant R01MH123513 (D.S.M.), National Science Foundation grant 1556974

(D.S.M.), Burroughs Wellcome Fund 1015667 (D.S.M.), Whitehall Foundation grant

2018-08-83 (D.S.M.), One Mind Foundation A137726 (D.S.M.), National Institutes of

Health grant R25MH060482 (K.M.B.), AP Giannini Foundation Fellowship P0534952

(K.M.B.), Larry L. Hillblom Foundation Fellowship 2020-A-023-FEL (K.M.B.), and

National Institute of Mental Health grant R01MH123178 (K.S.P.)

Author contributions

Conceptualization, R.S., K.M.B., K.L., and D.S.M.; methodology, A.B., A.J.W., D.S.M.,

K.L.B., K.M.B., K.S.P., and R.S.; software, K.Q., and S.W.; formal analysis, A.E., R.S.,

and K.M.B.; investigation, R.S., K.M.B., A.E., B.W., G.W., S.W., K.Q., R.D.L., K.L., N.H.,

B.A.S., M.C.H., M.S., R.K., A.C., D.G., L.C.C., N.L.G.; visualization, R.S., K.M.B., B.W.,

and A.E.; funding acquisition, D.S.M., K.S.P., and A.J.W; writing– original draft, R.S., and

K.M.B.; writing-review & editing, R.S., K.M.B., D.S.M., N.H., K.S.P., A.E., and A.B.

Declaration of interests

724   The authors declare no competing interests.

Star Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
DAPI	Sigma-Aldrich	Cat #D9542; CAS #28718-90-3

Experimental models: Organisms/strains		
Prairie vole: OxtR[1-/-]	Berendzen et al.	
Oligonucleotides		
ISH probes	This paper	N/A
Oxtr[1-/-] genotyping oligos	This paper	N/A
Antibodies		
Oxytocin Polyclonal	Invitrogen	Cat#PA5-26701; RRID
Anti-Neurophysin 2/NP-AVP	Millipore	Cat#MABN845; RRID
Cy™3 AffiniPure Donkey Anti-Rabbit IgG	Jackson ImmunoResearch	Cat #711-165-152; RRID
Alexa Flour 488 Donkey Anti Mouse IgG	Thermo Fisher Scientific	Cat #711-165-152; RRID
Software and algorithms		
ImageJ	NIH	; RRID: SCR_003070
R (version 1.1.423)	R Core	https://www.r-project.org/
GraphPad Prism (version 7.03)	GraphPad Software	
Boris	Boris	www.boris.unito.it

MATLAB	MathWorks	https://www.mathworks.com/products.html RRID: SCR_001622

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Devanand Manoli (devanand.manoli@ucsf.edu).

Materials availability

All unique reagents generated in this study are available from the lead contact upon

request.

Data

Transcriptomics data discussed in this publication are accessible through GEO Series

accession number GSE279248.

Code

All code used for RNA-seq data processing and differential expression analyses, as well

as post-processed expression data, can be found at

https://github.com/aseveritt/OXTR_prairie_voles

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals

Subjects were laboratory-bred prairie voles (*Microtus ochrogaster*) which originated through systematic outbreeding of a wild stock captured near Champaign, Illinois. CRISPR/Cas9 editing was used to generate 2 independent null alleles, Oxtr$^{1-/-}$ and Oxtr$^{2-/-}$, both of which were validated using autoradiography to show loss of receptor function[59] (data not shown). Oxtr$^{1-/-}$ was backcrossed for >7 generations to mitigate any possible off-target effects and this line was used for all subsequent experiments. Sexually naïve male and female animals were group weaned at 21 ± 1 days and separated to group housing with same-sex siblings and age-matched same-sex non-siblings. Voles were maintained under a 14:10 h light-dark cycle in clear plastic cages (45 × 25 × 15 cm) with bedding, nesting material (nestlet), and a PVC hiding tube. Rooms were maintained at approximately 20°C, and food and water were available *ad libitum*.

Breeding pairs were established between two heterozygotes or a homozygous Oxtr$^{1-/-}$ mutant male and heterozygous female partner from a breeding line maintained in the lab. Voles were randomly assigned into experimental groups when they reached 7-9 weeks of age at the start of testing. This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals published by the National Research Council. The protocol was approved by the Institutional Animal Care and Use Committee at each respective institution.

Genotyping

The following primers were used to amplify a region where XcmI restriction for Oxtr[1] or

Bpu10I restriction for Oxtr[2] determined the genotype of all animals studied. XcmI cuts

the mutant allele in the case of Oxtr[1] and Bpu10I restricts the WT sequence for Oxtr[2].

(Forward: ACTGGAGCTTCGAGTTGGAC, Reverse: ATGCCCACCACTTGCAAGTA)

METHOD DETAILS

Behavior

Triadic naive choice assay

The linear three chamber partner preference apparatus with open top and 10 x 32 inch

walls[59] was modified to capture the interactions of the naive, freely moving WT subject

and the naive, tethered, opposite sex potential partner (either 2 WTs or a WT and an

Oxtr[1-/-]) at either end of the chamber. Behaviors were recorded for 6 hours using a top

view camera and the videos generated were analyzed using a contour detection

algorithm. Specifically, contours were detected on individual video frames using a

threshold-based strategy that took advantage of the contrast between dark-colored vole

fur coats and light-colored bedding and chamber apparatus. A standalone vole was

detected as a single contour, and two voles in physical contact were likewise detected

as a single contour. Consistent vole identities across video frames were determined

using the Hungarian algorithm. Since the assay began with one vole in each of the three

chambers, and since the subject vole remained freely mobile throughout, it was possible

to unambiguously assign each contour to a single vole identity (e.g., subject, right

chamber, or left chamber) or to that of a pair of voles (e.g., subject and right chamber

voles or subject and left chamber vole), resulting in continuous trajectories for each of

the three animals. Time in chamber was determined by the chamber location of the

subject vole in each frame, since the chamber location of the right chamber and left

chamber animals was fixed throughout the assay by physical tethering and any time the

contour of the subject animal merged with that of the tethered animal (right or left), the

subject was termed in contact with it. The algorithm also determined the boundaries of

the chambers and calculated chamber times. We used the chamber time data from

when both ends of the chamber contained tethered WTs (WT-WT condition) to ensure

that there was no bias in which side the choosers preferred. This footage was then

scored manually to produce greater resolution of behaviors.

Bond formation assays

Initial introduction of $Oxtr^{1-/-}$ and WT animals of either sex with an opposite sex WT animal

was filmed for 30 minutes. Observed behaviors included anogenital sniffing, investigation

(subject animal's nose contacts any non-genital region), duration of stationary huddling

or >50% side-by-side contact between animals, rearing, frequency of aggressive behavior

including lunges, receipt of aggression, and tussles, as well as mating behavior including

mounting, intromission, and ejaculation. These animals remained in their home cage for

18 hours following pairing, after which, they were separated by a clear, plastic barrier. The

barrier was left in place for 24 hours and then removed allowing the subject free access

to the partner animal thereafter. A 30-minute period directly following barrier removal was

also filmed to assess mating behavior, termed the timed mating assay.

Subjects remained cohoused with the opposite-sex, wildtype partner for durations as

indicated in the text. The partner preference test (PPT), as previously published[59], was

used to assess pair bond formation. Subject, partner, and an opposite sex stranger are

placed in a three-chamber apparatus with open top and 10 x 32-inch walls. The partner

and stranger animals are tethered on either end of the three-chamber arena and the

subject allowed free access over a period of 3 hours. Behaviors are recorded from a top

view camera capturing the entire apparatus. Videos were scored using BORIS[124] post-

test by validated scorers blind to condition. Observed behaviors included location (i.e.,

duration of time in partner, stranger, and neutral chambers), duration of stationary

huddling or >50% side-by-side contact between the partner and stranger animals, and

frequency of aggressive behavior (i.e., lunges). Of note, analysis of partner preference

between wild type siblings from the $Oxtr^{1-/-}$ background and wild type animals from the

breeding colony showed no difference in time spent huddling with partner vs stranger

animals.

Bond maintenance assays

Four days after the timed mating assay, female animals underwent a separation-

reunification assay. Females remained in their home cage while their male partners

were placed in a separate cage in another room for one hour. The males were then

reintroduced into the home cage and behavior was video recorded for 30 minutes. Male

subject animals underwent the same procedure five days following timed mating. Two

days following the separation reunification test, animals underwent a selective

aggression assay. Partners were again removed from the home cage and separated to

a separate room for an hour. After one hour, a novel sexually naïve animal of the

opposite sex was introduced into the home cage with the subject animal and behavior

observed for 20 minutes. If significant aggression occurred, defined as >3 tussles during

the assay, the animals were separated and the assay terminated. Overall, two WT

female assays and one Oxtr[1-/-] were terminated for aggression. The same behaviors

listed above for the introduction were observed and recorded in the reunification and

aggression assays.

RNA Sequencing

Sample collection

Adult 7-9 week old WT and Oxtr[1-/-] voles were either cohoused with same sex-siblings or

paired with an animal of the opposite sex for four days following the timed mating protocol

described above. These animals were euthanized by CO_2 inhalation and decapitated. The

brain was quickly dissected out and sectioned into 500µm coronal slices using a brain

matrix mold (BrainTree Scientific) chilled on ice. Slices were floated in ice-cold phosphate

buffered saline (PBS), the NAc was identified using anatomical landmarks, and dissected

using a Zeiss microscope. A total of 45 WT females, 39 WT males, 21 Oxtr[1-/-] females,

and 18 Oxtr[1-/-] males were split between paired and unpaired conditions and tissue from

three animals per condition was pooled prior to RNA extraction. Tissue was flash-frozen

in liquid nitrogen and stored at -80°C until further processing. Total RNA was extracted

using TRIzol according to manufacturer's instructions and quantified using NanoDrop

(Thermo Scientific). Library preparation for next generation sequencing, comprising 35

samples and 4 technical replicates, was done using TruSeq Stranded mRNA Kit

according to manufacturer recommendations and sequenced on a NovaSeq 6000 to an

average depth of 2.9e07 reads per sample. Sequencing was performed across three

runs, but all data was processed together.

Read processing

Adapter and low-quality sequence trimming was performed with Trimmomatic v0.39

using parameters: 'TruSeq3-PE.fa:2:30:10:2:keepBothReads LEADING:3 TRAILING:3

SLIDINGWINDOW:4:20 MINLEN:25' (Bolger et al., 2014). Trimmed RNA-Seq reads

were aligned to a custom MicOch1.0 prairie vole genome, which incorporated a known

missing gene V1A, using STAR v2.7.3a in gene annotation mode (Dobin et al., 2013).

Alignment, RNA-Seq, and Insert Size quality control metrics were generated using

Picard v 2.10.10 (http://broadinstitute.github.io/picard). Homologous genes between

vole, mouse, and human references were identified using biomart v2.56.1. When

multiple mouse or human genes mapped to the same vole gene, the homologous gene

with the highest percent identity to vole was selected.

Differential expression analysis

To remove non- and lowly-expressed genes, only genes with more than 2 counts per

million (cpm) in at least 3 samples were retained. Principal component analysis of the

cpm revealed that technical replicates all cluster tightly but showed a slight batch effect

due to sequencing run (Fig. S5A). To remove this effect, we used RUVSeq v1.34.0 to

estimate one factor of unwanted variation, W_1, which we later used to normalize the

data prior to differential expression[125]. The factor analysis was performed on the

deviance residuals from an initial generalized linear model regression of the upper

quartile normalized counts on our covariates of interest (pair bonding status, genotype,

sex). Expressed genes were tested for differential expression (DE), after combining

technical replicates, with DESeq2 v1.40.2 using a design matrix that included the

covariates and W_1[126]. The adaptive shrinkage estimator ashr v2.2-63 was used in

order to be more robust to genes with low cpm values[127]. For DE analyses in general,

while most changes were driven by female samples, the effect was maintained in the

males at a smaller magnitude; thus, analyses were primarily performed for both sexes

jointly to maximize accuracy and power.

Bonding effect

To compare pre- to post-pairing individuals, we performed the DE analysis in two ways.

First, we compared pairing status regardless of genotype or sex to increase our

statistical power and to identify a reproducible gene set of thirteen genes (Fig. 5B, 5C,

S5B). While the relationship between the genes is unknown, there is support that the

proteins are functionally related. Using the STRING protein database

(STRINGdb_2.12.1), the human-homologs of the DE genes share more connections

than we would expect by chance (Fig. S5C, p-value= 3.08E-0.7). Next, we compared

pairing status using all three covariates, to identify subgroup-specific genes of interest

and effect sizes (Fig. 5D, S5D). In both analyses, genes were considered significantly

DE with a BH adjusted p-value < 0.05.

 Genotype effect

To examine the effect of genotype, we first performed DE analyses between WT and

Oxtr$^{-/-}$ individuals, not including sex as a covariate to identify a reproducible and

stringent gene set. We considered genes DE with an adjusted p-value < 0.05 and

absolute log2 fold-change > 0.25 (Fig. 6B, S6A, S6B). To examine the functional

relationship of the DE genes, we performed a Gene Ontology enrichment analysis using

goseq v1.52.0 which controls for gene length bias[128]. We considered all expressed vole

genes with a mouse homolog as our background and all DE genes with a mouse

homolog as our foreground. GO categories with an adjusted p-value < 0.05 were

considered significantly enriched. To help visualize the results, enriched GO categories

were clustered according to their semantic similarity calculated with the R package

GOSemSim v2.26.1 using method "Wang"[129]. Large, general, GO categories with over

1000 genes were excluded in visualizations (Fig. 6C). WikiPathways[130] over-

representation analysis was performed with R package clusterProfiler v4.8.3[131] (Fig.

S6D).

For disorder and disease set overrepresentation analysis, Chi-squared tests were used

to assess whether Oxtr$^{1-/-}$ disrupted genes occur in disease-associated gene lists at a

frequency higher than we would expect by chance. Genes lists for autism spectrum

disorder[74], schizophrenia[75], neurodevelopmental disorders[77], and epilepsy[76] were

sourced and all alternate or outdated gene symbols were updated manually. For this

test, we were interested in sex- and direction-specific enrichments, so we performed a

DE analysis separately for males and females. Across all DE analyses, we defined DE

genes more stringently (adjusted p-value < 0.01). For enrichment tests, our universe

was all expressed vole genes with a human homolog. We split this analysis by fold-

change direction to highlight the distinct enrichment in upregulated genes. P-values

underwent BH correction across all 24 comparisons.

To compare to mouse neuronal populations, we leveraged single-cell data on mouse

hypothalamus development[83]. The released Seurat object

GSE132730_subset_cca_to_velo.rds.gz was accessed at GSE132730. Using the

authors' defined clusters (accessed at "prim_walktrap"), we assessed how similar each

clusters' gene expression is to the genes disrupted by the $Oxtr^{1-/-}$ mutation. This was

calculated as the Jaccard similarity coefficient using R package clustifyR v1.12.0[132].

Weighted Gene Coexpression Network Analysis

Data

Starting with the RNAseq dataset of n=30 samples (one WT sample and 4 replicate

samples were removed) and n=13357 genes, we removed genes with low expression

(n=1738 genes with expression level in lowest quantile for all 30 samples) and/or low

variability (n=8230 genes with coefficient of variation < 0.05), resulting in a trimmed

dataset of n=3389 genes.

Identifying co-expression modules

We used the R package WGCNA (v1.70-3) to conduct WGCNA analyses[72]. Briefly, we

calculated Pearson correlations for all gene pairs. Next, we constructed an unsigned

weighted correlation network to identify co-expression modules comprised of highly

correlated genes (positive or negative correlation) with high topological overlap[133].

Modules were defined as branches of a hierarchical cluster tree using the top-down

dynamic tree cut method[72]. For each module, the expression pattern was summarized

by the module eigengene (ME), which is defined as the right singular vector of the

expression patterns. Pairs of modules with high module eigengene correlations (r >

0.75) were merged to maintain a level of decorrelation among MEs.

In more detail, a weighted unsigned network was computed based on a fit to scale-free

topology, and the lowest thresholding power that resulted in a scale-free R^2 fit of 0.9

was selected, and the pairwise topological overlap (TO) between genes was

calculated[133] . We constructed a TO dendrogram using hierarchical clustering (R::hclust

function, method= "average"), and defined modules using the WGCNA::cutreeHybrid

function with minimum module size set to 30 genes. We then merged modules with high

module eigengene correlations as describe above.

We calculated 3 sets of modules from different sample subsets:

965   1. Modules^{WT}: n=15 modules constructed from n=18 WT samples with

966   thresholding power β = 5.

967   2. Modules^{Mut}: n= 43 modules constructed from n=12 Oxtr^1-mutant samples

968   with thresholding power β = 6.

969   3. Modules^{All}: n=7 modules constructed from all n=30 samples with

970   thresholding power β = 3

<u>Relating modules to sample traits</u>

In order to correlate modules with experimental conditions, we calculated the Pearson

correlations between module eigengenes and sample metrics, including species

(Mut=0, WT=1), condition (Pre=1, Post=0), sex (F=1, M=0), and quality control (QC)

metrics (RIN, yield, sequencing run). We identified three modules that significantly

correlated with a sample metric of interest but not with any of the QC metrics.

Significant correlation was defined to be $p_{adj} <= 0.05$, where p_{adj} = BH correction by

number of modules. These modules (M) were:

1. Mpink[WT]: positive correlation with pre-bonding status (Pearson R = 0.7, p_{adj} =

   0.018)

2. Mbrown[All]: positive correlation with WT status (Pearson R = 0.55 , p_{adj} = 0.011)

3. Mturquoise[All]: positive correlation with WT status (Pearson R = 0.8, p_{adj} = 6.6e-7)

We calculated the module eigengenes (ME) for Mpink[WT], Mbrown[All], and Mturquoise[All]

for all n=30 samples using the WGCNA::moduleEigengenes function. We then

conducted Wilcoxon rank sum tests (ggpubr::stat_compare_means function) to assess

whether MEs were significantly different between the following groups:

1. MEpink[WT]: pre- versus post-bonded animals, separated by sex and species

2. Mbrown[All]: WT versus mutant animals, separated by sex and condition

3. Mturquoise[All]: WT versus mutant animals, separated by sex and condition

<u>Comparing gene significance with module membership:</u>

We assessed whether genes whose expression highly correlate with module

eigengenes of modules of interest (Mpink^{WT}, Mbrown^{All}, and Mturquoise^{All}) are

themselves highly correlated with traits of interest (bonding status or mutation status).

Specifically, for the genes in Mpink^{WT}, Mbrown^{All}, and Mturquoise^{All}, we calculated 1)

gene significance (GS), defined as the absolute Pearson's correlation between

expression of a given gene and a trait of interest; and 2) module membership (MM),

which is defined as the absolute Pearson's correlation between the expression profile of

a gene with the ME of a module to quantify the relationship between a gene and a given

module[72]. We then calculated the Pearson correlations between:

1. GS^{condition} and MMpink^{WT} for Mpink^{WT} genes ($R = 0.69$, $p = 5.5e-18$)

2. GS^{species} and MMbrown^{All} for Mbrown^{All} genes ($R = 0.63$, $p = 2.2e-32$)

3. GS^{species} and MMturquoise^{All} for MMturquoise^{All} genes ($R = 0.82$, $p<1e-200$)

Hub genes were defined as those genes in the top 25% for both the absolute value of

pink module membership and the absolute value of correlation with bonding status.

1007

1008 <u>Assessing whether modules are enriched for DE genes</u>

1009 We were interested in assessing whether Mpink^{WT} genes are significantly enriched for

1010 any the DE pre vs post pairing gene set described above. We assessed for enrichment

1011 of DE gene sets using a hypergeometric test.

1012

<u>WT module preservation analysis (Zsummary)</u>

We assessed how well modules^{WT} are preserved in mutant samples. We calculated a

Z$_{summary}$ statistic for each module^{WT} [134] using the WGCNA::modulePreservation function

with 200 permutations. The $Z_{summary}$ measure combines module density and

intramodular connectivity metrics to a composite statistic where Z>2 suggests moderate

preservation and Z>10 suggests high preservation. We found of all the n=15

modules^{WT}, Mpink^{WT} had the lowest $Z_{summary}$ ($Z_{summary}$=2) in mutant samples.

1020

Histology

Immunohistochemistry (IHC)

Sexually naïve voles group housed by sex after weaning (P0, P21, adults:10 - 14 weeks

old), were perfused with 25mL ice cold phosphate buffer (PBS) and fixed in 25mL of 4%

paraformaldehyde in PBS (PFA). Whole brain was dissected and post fixed overnight at

4°C in 4% PFA in the dark. The tissue was cryoprotected for 24 hours in 30% sucrose in

PBS and then embedded in OCT over dry ice to prevent cracking. 60μm serial sections

from the entire PVN were collected in PBS and processed as free-floating sections.

Sections were blocked using 10% donkey serum in 0.1% Triton X 100 in PBS and then

stained with primary antibodies (Oxytocin Polyclonal Antibody, Invitrogen, Rabbit

Polyclonal, Cat #PA5-26701, 1:1000 and Anti-Neurophysin 2/NP-AVP Antibody, clone

PS 41, Millipore Sigma, Mouse Monoclonal, Cat#MABN845, 1:10,000) in 1% donkey

serum with 0.1% Triton X 100 (staining buffer) in PBS at 4°C with shaking. The sections

were rinsed and soaked in secondary antibodies (Cy™3 AffiniPure Donkey Anti-Rabbit

IgG, Jackson ImmunoResearch Laboratory, Cat #711-165-152, 1:500 and Alexa Flour

488 Donkey Anti Mouse IgG, Thermo Fisher Scientific, Cat #A21202, 1:500) in staining

buffer for 2 hours at room temperature. Sections were DAPI stained and transferred to

glass slides in sequence and mounted in Aquamount. Slides were imaged using a

confocal microscope (Zeiss LSM 700) and the zstacks were quantified (cell count by

hand and area hand selected and measured) in ImageJ. Comparisons of cell density

(count/area) were made in age and sex matched samples.

1042

In situ hybridization (ISH)

ISH was performed using RNA probes prepared as described previously[135], to detect

expression of *Calcr, Dlk1*, and *Dgkb* mRNA in the adult vole brain. We prepared RNA

sense and anti-sense probes corresponding to 706bp (CalcR), 773bp (Dgkb), and

796bp (Dlk1) for genes of interest identified in the WT vs Oxtr DEX list from our RNAseq

data. For ISH on adult animals, 7-9 week old WT and Oxtr$^{1-/-}$ voles of both sexes were

either group housed with same sex conspecifics or paired for 4 days with a WT animal

of the opposite sex following the timed mating protocol described above. The brains

were processed as described for IHC, then embedded in OCT and sectioned at 50µm,

mounted immediately after sectioning onto SuperFrost Plus glass microscope slides

and stored at -80°C until ISH staining. Slides were fixed in 4% PFA for 20 minutes,

rinsed in PBS, treated with proteinase K (10µg/mL, Roche) for 20 minutes, rinsed in

PBS, and fixed again in 4% PFA for 5 minutes at room temperature. Slides were

acylated for 10 mins then rinsed in 1% Triton100X in PBS followed by PBS. The slides

were equilibrated in warm hybridization solution for 1-2 hours at 65°C and subsequently

incubated with a temporary paraffin cover for 14 - 18 hours at 65°C in fresh

hybridization buffer containing RNA probe. After incubation, the slides were dipped in

5X SSC warmed to 72°C to remove the parafilm coverslips and then washed in 0.2X

SSC warmed to 72°C. The slides were blocked with 10% heat inactivated sheep serum

(HISS) solution then stained for 12 - 18 hours at 4°C in buffer containing 10% HISS and

alkaline phosphatase-conjugated sheep anti- digoxigenin antibody (1:2000, Roche).

After extensive washing, the slides were incubated for ~72 hours at 37°C in staining

solution containing nitro blue tetrazolium and 5- bromo-4-chloro-3-indolyl-phosphate

(Roche). The slides were finally washed, fixed in 4% PFA, and coverslipped. Slides

were imaged using a Keyence fluorescence microscope and the z-stacks were

quantified (cell count and area) in ImageJ. For signal quantitation, positive puncta were

counted in every third coronal section encompassing the entire anterior to posterior

gene expression domain. Cell counts were normalized by total area for a gene

expression domain in each section. Positive signal was quantified by calculating total

area of signal multiplied by the average intensity of signal (total area x mean pixel

intensity) in gene expression domains of a section; every third section was quantified

and added. Comparisons of cell counts were made in age and sex matched samples.

Quantification and Behavioral Analysis

For detailed statistical methods please see Table S1. The number of animals used was

based on previous studies in the field by our group and others, combined with a power

analysis. Assumptions of independence and normality were considered with multiple

tests. Preference index (PI) was calculated as (partner huddle duration – stranger huddle

duration)/(partner huddle duration + stranger huddle duration). Sliding partner preference

was calculated by calculating PI for a single animal in 20-minute intervals where each

interval overlapped the next by 15 minutes (e.g. 0-20mins, 5-25mins…). To look at

convergence in sliding PI the area under the curve (Area above 0 – Area below 0, AUC)

for each hour of the assay was determined for each animal and then pooled for each

choosing animal by sex and then condition. Preference was determined as significantly

convergent if the AUC from the hour of assay for all animals within the group significantly

differed from 0 using a One sample t Test. The level of statistical significance for each test

was set at $p =< 0.05$. Outliers were detected using a combination of z-score (+/- 3) and

plots of original and log-transformed data. Analyses were completed in RStudio (version

2024.04.2+764) and GraphPad Prism (version 10.3.1)

 FIGURE LEGENDS

Figure 1: Oxtr reduces the amount of time naïve animals take to show partner preference

Figure1: Oxtr reduces the amount of time naive animals take to show partner

preference

A. Schematic of the partner preference tests (PPTs) and the day on which they were

conducted to study pair bonding behaviors. The prosocial, agonistic and mating

behaviors scored are shown below. During the bond formation phase, the ratio of

partner-directed vs stranger-directed affiliative behaviors increases in wild types

and following the bond formation in the bond maintenance phase, wild types show

decreased prosocial behaviors and increased agonistic behaviors towards

strangers.

B. A preference for partner is induced in the WTs but not Oxtr[1-/-] animals following

short cohabitation.

C. WT females spend more time huddling with their partners.

D. Oxtr[1-/-] females spend more time huddling with strangers.

E. WT females show a significant increase in partner huddling time towards the end

of a 3-hour assay.

F. Oxtr[1-/-] males show a significant increase in stranger huddling time in the middle of

a 3-hour assay.

G. No difference in partner huddling time between WT and Oxtr[1-/-] animals following

long cohabitation.

H. No difference in stranger huddling time between WT and Oxtr[1-/-] animals following

long cohabitation.

See Supplementary Figure 1 and Supplementary Table 1.

Mean +/- SD, n =>8, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT =

male WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$.

Figure 2: Oxtr suppresses promiscuous behaviors in a state dependent manner in females

Figure2: Oxtr suppresses promiscuous behaviors in a state dependent manner in females

A. Timeline of the behavioral battery carried out on females. Bottom row shows the behaviors scored.

B. Oxtr[1-/-] female long huddle duration is shorter with partners and longer with strangers after a short cohabitation.

C. Oxtr[1-/-] female long huddle frequency is greater with strangers, but not partners after a short cohabitation.

D. No difference in median long huddle durations between WT and Oxtr[1-/-] females post long cohabitation.

E. No difference in long huddle frequencies between WT and Oxtr[1-/-] females post long cohabitation.

F. Increasing cohabitation time increases the median long huddle duration an Oxtr[1-/-] female executes with the partner.

G. There is no significant change in the median long huddle duration conducted by females with strangers when cohabitation time is increased.

H. Naive Oxtr[1-/-] and WT females are equally prosocial towards a naive WT male.

I. There is no difference in the counts of mating behaviors between Oxtr[1-/-] and WT females.

J. Oxtr[1-/-] and WT females are equally prosocial towards their partners after a brief separation.

K. Oxtr[1-/-] females are significantly more prosocial towards a naive WT stranger male compared to WT females.

1142　See Supplementary Figure 2 and Supplementary Table 1.

1143　Mean ± SD, n =>8, *p < 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT = male

1144　WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$.

Figure 3: Oxtr function influences different prosocial behaviors across pair bonding in males

1145
Figure3: Oxtr function influences different prosocial behaviors across pair

bonding in males

A. Timeline of the behavioral battery carried out on males. Bottom row shows the

behaviors scored.

B. Oxtr[1-/-] male long huddle duration is shorter with partners and strangers after a

short cohabitation.

C. Oxtr[1-/-] male long huddle frequency is greater with strangers, but not partners after

a short cohabitation.

D. The median long huddle executed by Oxtr[1-/-] males with the stranger after a long

cohabitation, is shorter than their WT siblings.

E. No difference in the long huddle frequencies between WT and Oxtr[1-/-] males post

long cohabitation.

F. There is no change in the frequency of long huddles with partner between Oxtr[1-/-]

and WT males.

G. Increasing cohabitation time decreases the frequency of long huddles with

strangers in Oxtr[1-/-] males.

H. Naive Oxtr[1-/-] and WT males are equally prosocial towards naive WT females.

I. Oxtr[1-/-] males execute higher counts of mating behaviors compared to WT.

J. Prosocial behaviors following a brief separation with their partner are not different

between Oxtr[1-/-] and WT males.

K. Oxtr[1-/-] males are significantly more prosocial towards a naive WT stranger female

compared to WT males.

See Supplementary Figure 3 and Supplementary Table 1.

Mean +/- SD, n =>8, *p =< 0.05, FWT = female WT, FOxtr[1-/-] = female Oxtr[1-/-], MWT =

male WT, MOxtr[1-/-] = male Oxtr[1-/-]

Figure 4: Oxtr sex-specifically influences social interactions between potential mates

Figure4: Oxtr sex-specifically influences social interactions between potential

mates

A. Schematic of naive social choice with a naive WT animal choosing between an

opposite sex WT and opposite sex $Oxtr^{1-/-}$ animal (red triangles).

B. WT choosers of both sexes spend significantly more time with one stimulus

animal (winner) over the other (loser). The color of each dot denotes the

genotype of the winner and loser.

C. WT and $Oxtr^{1-/-}$ animals are chosen as winners with equal frequency by WT

choosers of both sexes.

D. Schematic of comparison of contact with winner from WTOxtr condition to that

with winner from WTWT condition.

E. Female WT choosers spends more time in contact with a winner from a WTOxtr

condition than the WTWT. The color of the dots represents the genotype of the

winner.

F. Female WT choosers spend significantly more time huddling with the winner from

the WTOxtr condition in the last 47 mins of the assay. Solid lines depict the

average cumulative huddle time, the solid black bar denotes the time at which

the lines significantly diverge.

G. Longest huddle from WTOxtr lanes is significantly higher compared to those from

WTWT lanes. The color of each dot denotes the genotype/position of the animal

with whom the longest huddle took place.

H. Female WT choosers from WTWT lanes spend more time sniffing both stimulus

animals 15 minutes before the longest huddle compared to female choosers from

WTOxtr lanes

I. Female WT choosers attack the loser in WTWT lanes more frequently than

choosers from WTOxtr lanes (right). There is no difference in the number of

attacks on the winners by female choosers from WTWT lanes compared to those

from WTOxtr lanes (left).

J. Total prosocial behavior time shows significant negative correlation with the total

counts of aggressive behaviors displayed by female WT choosers.

K. Naive male WTs attack a potential female WT mate more frequently than male

Oxtr$^{1-/-}$ animals (left). There is no difference in the number of attacks against an

unfamiliar naive WT female stranger by bonded WT and Oxtr$^{1-/-}$ males (right).

See Supplementary Figure 4 and Supplementary Table 1.

Mean +/- SD, n = 8, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT = male

WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$, WTWT = Lane with 2 naive WT animals, WTOxtr = Lane with

a WT and a Oxtr$^{1-/-}$ animal.

Figure 5: Post pair bonding gene expression is driven by female specific changes dependent on Oxtr signaling

Figure 5: Post pair bonding gene expression is driven by female specific changes

dependent on Oxtr signaling

A. Schematic of experimental groups used for sequencing highlighting pre- vs post-

pairing comparisons between groups. Animals with symbols represent those used

for tissue collection.

B. Scatterplot illustrating the relationship between fold change and mean expression,

emphasizing the 13 differentially expressed (DE) genes in post-pairing individuals

compared to pre-pairing. Gray dots indicate non-significant genes ($p_{adj} > 0.05$);

Blue dots indicate downregulated genes ($p_{adj} < 0.05$, $\log_2FC < 0$) and upregulated

genes in red ($p_{adj} < 0.05$, $\log_2FC > 0$). The right panel shows box plots of normalized

gene expression for the top six DE genes by fold-change between pre- and post-

pairing conditions.

C. Bar graph emphasizing how each combination of genotype and sex (denoted at

bottom as 'subgroup') influence the 13 overall pairing-DE genes identified. The top

graph depicts the absolute fold change and standard error of the differential

expression analysis including all pre- and post-bonding individuals in which the 13

genes of the x axis were identified. The bottom graph depicts the absolute fold

change and standard error in analyses when only individuals of that subgroup were

considered. (* denotes $p_{adj} < 0.05$ for the pre- vs post-pairing analysis).

D. Box plots of select genes that show sex- and genotype-specific changes in

expression with pairing condition. *Adora2A* and *Fkbp5* are DE when considering

WT females, *Spred3* is DE when considering WT and Oxtr[1-/-] females, and *Sema3b*

is DE when considering Oxtr[1-/-] males. Note: Sema3b is the mouse ortholog gene

symbol for vole gene "ENSMOCG00000019704" which does not have an assigned

gene symbol.

E. Heatmap showing correlations between module eigengenes (MEs) for modules[WT]

and pairing condition (Pre=1, Post=0). Heatmap color and value reflect Pearson

correlation coefficient; adjusted p values shown in parentheses (p_{adj} = p * number

of modules).

F. Preservation of modules[WT] in OXTR-mutant samples. The $Z_{summary}$ measure

combines module density and intramodular connectivity metrics to a composite

statistic where Z>2 suggests moderate preservation and Z>10 suggests high

preservation[134]. Of all modules[WT](n=15), Mpink[WT] had the lowest $Z_{summary}$

($Z_{summary}$=2) in mutant samples.

G. Box plots showing comparison of MEpink[WT] for pre- vs post-pairing condition

separated by sex and genotype. (Wilcoxon rank sum test, Bonferroni adjusted for

4 comparisons).

For all box plots center = median, box boundaries = 1st and 3rd quartile, whiskers =

1.5*IQR from boundaries. (*n* = 16 pre-pairing and 15 post-pairing)

See Supplementary Figure 5.

Figure 6: Oxtr regulated genes are common to social behavior relevant processes across species and are similarly regulated across brain regions

Figure 6: Oxtr regulated genes are common to social behavior relevant processes across species and are similarly regulated across brain regions

A. Schematic of experimental groups for sequencing, highlighting genotype comparisons between groups. Animals with symbols represent those used for tissue collection.

B. Scatterplot illustrating the relationship between fold change and mean expression, emphasizing differentially expressed (DE) genes in Oxtr[1-/-] individuals compared to WT. Significantly downregulated genes are shown in blue ($log_2FC<-0.25$, $p_{adj}<0.05$) and upregulated genes in red ($log_2FC>0.25$, $p_{adj} < 0.05$). Box plots of normalized gene expression for the top six DE genes by fold change between genotypes.

C. Gene Ontology enrichment analysis for all genotype-DE genes (n=1014) highlights processes related to extracellular channel binding activity, axonal processes, behavior related gene expression, and cation transport. The leaves represent significant GO categories (BH adjusted p value < 0.05) which have been hierarchically clustered by their semantic similarity. Branches are colored by cluster and high-frequency words are displayed to the right. Leaf size and color correspond to the number and median fold change of DE genes in each category.

D. Enrichment analysis shows that genes which are DE in Oxtr[1-/-] samples are over-represented in some neuropsychiatric disease-associated gene sets. The first DE gene set evaluated ('Joint') represents genes which are DE in Oxtr[1-/-] samples when considering both sexes in the differential expression analysis ($p_{adj} < 0.01$). The second gene set are DE genes when only considering females, and the third

when only considering males. Gene sets are further separated by direction of effect

across the x-axis to highlight the enrichment in upregulated DE genes. Colors

correspond to the Benjamini-Hochberg corrected Chi-square p values; asterisks

denote $p_{adj} < 0.05$. Y-axis abbreviations are as follows: SCZ=schizophrenia,

NDD=neurodevelopmental disorder, EPI=epilepsy, and ASD=autism spectrum

disorder.

E. Box plots of *Calcr* and *dlk1* normalized gene counts by genotype. Log_2FC:-0.962

and p_{adj}: 1.75e-06 for *Calcr* and log_2FC: -0.276 and p_{adj}: 0.00334 for *dlk1* from

comparison considering all WT and Oxtr$^{1-/-}$ samples.

F. Brain regions selected for in situ analysis. (NAc=nucleus accumbens,

PVN=paraventricular nucleus of the hypothalamus, LS=lateral septum)

G. ISH images and quantification for *Calcr* expression in NAc, PVN, and LS from WT

and Oxtr$^{1-/-}$ animals.

H. ISH images and quantification for *Dlk-1* expression in NAc, PVN, and LS from WT

and Oxtr$^{1-/-}$ animals.

Box plots in G. and H. show comparison of the area-normalized cell number by

genotype. (*n*=3 per condition, Wilcoxon sign rank, *=p<0.05, ***=p<0.001)

See Supplementary Figure 6 and Supplementary Table 1.

Figure 7: Loss of Oxtr results in the decrease in density of Oxt and Avp expressing neurons in the PVN

1292
1293

Figure7: Loss of Oxtr results in the decrease in density of Oxt and Avp

expressing neurons in the PVN

A. UMAP visualization of mouse hypothalamus single-cell RNAseq dataset[83] with proposed PVN clusters labeled. The authors' original numeric cluster labels and embedding are shown. Each cluster is colored by the Jaccard similarity coefficient between the mouse marker gene sets and the vole $Oxtr^{1-/-}$ DE genes (ONP set). This is also represented in a heat plot to the right. Some of the highest similarity is seen in Cluster 43, which represents Oxt+ cells, and Cluster 26, which represents Avp+ cells in the PVN.

B. Representative images of IHC staining against OT and AVP neuropeptides from FWT and $FOxtr^{1-/-}$ PVN showing the loss of structural integrity and OT and AVP positive cells from the knock-out. (White dotted line denotes the PVN area, scale bar represents 100μm)

C. Mean of positive cells per unit area across all sections is lower for $Oxtr^{1-/-}$s vs WT for both OT (left) and AVP (right). Each dot represents 1 animal.

D. The number of sections with a high density of OT positive cells is greater in both female (left) and male (right) WT animals.

E. Similarly, WT animals of both sexes show an increase in the number of sections containing AVP positive neurons in high densities.

F. The average OT cell density from matched PVN sections arranged from anterior to posterior shows that the decrease in cell density in $Oxtr^{1-/-}$ animals is biased towards the anterior. Each dot represents the average cell density, the lines

represent the sixth order non-linear regression model for cell density by sex and

by genotype (n=3) and the ribbons represent the standard deviation.

G. $Oxtr^{1-/-}$ animals show a significant decrease in AVP cell density in the anterior PVN

in both sexes.

See Supplementary Figure 7 and Supplementary Table 1.

Mean +/- SD, n = 3, *p =< 0.05, FWT = female WT, $FOxtr^{1-/-}$ = female $Oxtr^{1-/-}$, MWT = male

WT, $MOxtr^{1-/-}$ = male $Oxtr^{1-/-}$.

1323

1324

1325   **References**

1326

1. Wilson, R.S., Krueger, K.R., Arnold, S.E., Schneider, J.A., Kelly, J.F., Barnes, L.L., Tang, Y.,
and Bennett, D.A. (2007). Loneliness and Risk of Alzheimer Disease. Arch Gen Psychiatry
64, 234. https://doi.org/10.1001/archpsyc.64.2.234.

2. Bowlby, J., and Bowlby, J. (1982). Attachment 2. ed. (Basic Books).

3. Farinelli, M., Cevolani, D., Gestieri, L., Romaniello, C., Maffei, M., Agati, R., Leo, M.R.,
Huang, Z., Pedone, V., and Northoff, G. (2020). Brain and behaviour in post-acute stroke:
Reduction in seeking and posterior cingulate neuronal variability. Journal of Clinical and
Experimental Neuropsychology *42*, 584–601.
https://doi.org/10.1080/13803395.2020.1780417.

4. Feldman, R., Monakhov, M., Pratt, M., and Ebstein, R.P. (2016). Oxytocin Pathway Genes:
Evolutionary Ancient System Impacting on Human Affiliation, Sociality, and
Psychopathology. Biological Psychiatry *79*, 174–184.
https://doi.org/https://doi.org/10.1016/j.biopsych.2015.08.008.

5. Ebstein, R.P., Israel, S., Chew, S.H., Zhong, S., and Knafo, A. (2010). Genetics of human
social behavior. Neuron *65*, 831–844. https://doi.org/10.1016/j.neuron.2010.02.020.

6. Winslow, J.T., Shapiro, L., Carter, C.S., and Insel, T.R. (1993). Oxytocin and complex social
behavior: species comparisons. Psychopharmacol Bull *29*, 409–414.

7. Goodson, J.L. (2013). Deconstructing sociality, social evolution and relevant nonapeptide
functions. Psychoneuroendocrinology *38*, 465–478.
https://doi.org/10.1016/j.psyneuen.2012.12.005.

8. Harlow, H.F., and Harlow, M.K. (1965). THE EFFECT OF REARING CONDITIONS ON
BEHAVIOR. Int J Psychiatry *1*, 43–51.

9. Shultz, S., Opie, C., and Atkinson, Q.D. (2011). Stepwise evolution of stable sociality in
primates. Nature *479*, 219–222.

10. Insel, T.R., Winslow, J.T., Wang, Z., and Young, L.J. (1998). Oxytocin, vasopressin, and the
neuroendocrine basis of pair bond formation. Adv. Exp. Med. Biol *449*, 215–224.

11. Klatt, J.D., and Goodson, J.L. (2013). Oxytocin-like receptors mediate pair bonding in a
socially monogamous songbird. Proceedings of the Royal Society B: Biological Sciences
280, 20122396. https://doi.org/10.1098/rspb.2012.2396.

12. Ainsworth, M.S. (1979). Infant–mother attachment. American Psychologist *34*, 932–937.
https://doi.org/10.1037/0003-066X.34.10.932.

13. Getz, L.L., Carter, C.S., and Gavish, L. (1981). The mating system of the prairie vole,
Microtus ochrogaster: Field and laboratory evidence for pair-bonding. Behav Ecol Sociobiol
8, 189–194. https://doi.org/10.1007/BF00299829.

14. Donaldson, Z.R. (2010). We're the Same... but Different: Addressing Academic Divides in
the Study of Brain and Behavior. Front Behav Neurosci *4*.
https://doi.org/10.3389/fnbeh.2010.00041.

15. Insel, T.R., Winslow, J.T., Wang, Z.X., Young, L., and Hulihan, T.J. (1995). Oxytocin and the
molecular basis of monogamy. Adv. Exp. Med. Biol *395*, 227–234.

16. Lukas, D., and Clutton-Brock, T.H. (2013). The evolution of social monogamy in mammals.
Science *341*, 526–530.

17. Donaldson, Z.R., and Young, L.J. (2008). Oxytocin, vasopressin, and the neurogenetics of
sociality. Science *322*, 900–904. https://doi.org/10.1126/science.1158668.

18. Bielsky, I.F., and Young, L.J. (2004). Oxytocin, vasopressin, and social recognition in
mammals. Peptides *25*, 1565–1574. https://doi.org/10.1016/j.peptides.2004.05.019.

19. Nowicki, J.P., Pratchett, M.S., Walker, S.P.W., Coker, D.J., and O'Connell, L.A. (2020).
Gene expression correlates of social evolution in coral reef butterflyfishes. Proceedings of
the Royal Society B: Biological Sciences *287*, 20200239.
https://doi.org/10.1098/rspb.2020.0239.

20. O'Connor, C.M., Marsh-Rollo, S.E., Aubin-Horth, N., and Balshine, S. (2016). Species-
specific patterns of nonapeptide brain gene expression relative to pair-bonding behavior in

grouping and non-grouping cichlids. Hormones and Behavior *80*, 30–38.
https://doi.org/10.1016/j.yhbeh.2015.10.015.

21. Stanley, D.A., and Adolphs, R. (2013). Toward a neural basis for social behavior. Neuron *80*,
816–826.

22. Insel, T.R. (1992). Oxytocin--a neuropeptide for affiliation: evidence from behavioral,
receptor autoradiographic, and comparative studies. Psychoneuroendocrinology *17*, 3–35.

23. Lim, M.M., and Young, L.J. (2006). Neuropeptidergic regulation of affiliative behavior and
social bonding in animals. Horm Behav *50*, 506–517.
https://doi.org/10.1016/j.yhbeh.2006.06.028.

24. Carter, C.S., and Getz, L.L. (1993). Monogamy and the prairie vole. Sci. Am *268*, 100–106.

25. Beery, A.K., and Shambaugh, K.L. (2021). Comparative assessment of familiarity/novelty
preferences in rodents. Frontiers in Behavioral Neuroscience *15*, 648830.

26. Beery, A.K., Christensen, J.D., Lee, N.S., and Blandino, K.L. (2018). Specificity in Sociality:
Mice and Prairie Voles Exhibit Different Patterns of Peer Affiliation. Frontiers in Behavioral
Neuroscience *12*, 50. https://doi.org/10.3389/fnbeh.2018.00050.

27. Schweinfurth, M.K., Neuenschwander, J., Engqvist, L., Schneeberger, K., Rentsch, A.K.,
Gygax, M., and Taborsky, M. (2017). Do female Norway rats form social bonds? Behavioral
Ecology and Sociobiology *71*, 1–9.

28. Cho, M.M., DeVries, A.C., Williams, J.R., and Carter, C.S. (1999). The effects of oxytocin
and vasopressin on partner preferences in male and female prairie voles (Microtus
ochrogaster). Behav. Neurosci *113*, 1071–1079.

29. Insel, T.R., and Hulihan, T.J. (1995). A gender-specific mechanism for pair bonding:
oxytocin and partner preference formation in monogamous voles. Behav. Neurosci *109*, 782–
789.

30. Williams, J.R., Catania, K.C., and Carter, C.S. (1992). Development of partner preferences in
female prairie voles (Microtus ochrogaster): the role of social and sexual experience.
Hormones and behavior *26*, 339–349.

31. Wang, Z., Hulihan, T.J., and Insel, T.R. (1997). Sexual and social experience is associated
with different patterns of behavior and neural activation in male prairie voles. Brain Res *767*,
321–332.

32. Xu, X., Coats, J.K., Yang, C.F., Wang, A., Ahmed, O.M., Alvarado, M., Izumi, T., and Shah,
N.M. (2012). Modular genetic control of sexually dimorphic behaviors. Cell *148*, 596–607.
https://doi.org/10.1016/j.cell.2011.12.018.

33. Villella, A., and Hall, J.C. (2008). Chapter 3 Neurogenetics of Courtship and Mating in
Drosophila. In Advances in Genetics (Elsevier), pp. 67–184. https://doi.org/10.1016/S0065-
2660(08)00603-2.

34. Konopka, R.J., and Benzer, S. (1971). Clock Mutants of *Drosophila melanogaster*. Proc Natl
Acad Sci USA *68*, 2112–2116. https://doi.org/10.1073/pnas.68.9.2112.

35. Takahashi, J.S., Shimomura, K., and Kumar, V. (2008). Searching for genes underlying
behavior: lessons from circadian rhythms. Science *322*, 909–912.
https://doi.org/10.1126/science.1158822.

36. Baker, B.S., Taylor, B.J., and Hall, J.C. (2001). Are complex behaviors specified by
dedicated regulatory genes? Reasoning from Drosophila. Cell *105*, 13–24.

37. Manoli, D.S., and Baker, B.S. (2004). Median bundle neurons coordinate behaviours during
Drosophila male courtship. Nature *430*, 564–569. https://doi.org/10.1038/nature02713.

38. Martin, L.B., Glasper, E.R., Nelson, R.J., and DeVries, A.C. (2006). Prolonged separation
delays wound healing in monogamous California mice, Peromyscus californicus, but not in
polygynous white-footed mice, P. leucopus. Physiology & Behavior *87*, 837–841.
https://doi.org/10.1016/j.physbeh.2006.01.035.

39. Ferland, C.L., and Schrader, L.A. (2011). Cage mate separation in pair-housed male rats
evokes an acute stress corticosterone response. Neuroscience Letters *489*, 154–158.
https://doi.org/10.1016/j.neulet.2010.12.006.

40. Okuda, T., Osako, Y., Hidaka, C., Nishihara, M., Young, L.J., Mitsui, S., and Yuri, K. (2022).
Separation from a bonded partner alters neural response to inflammatory pain in
monogamous rodents. Behavioural Brain Research *418*, 113650.
https://doi.org/10.1016/j.bbr.2021.113650.

41. Carter, C.S., Williams, J.R., Witt, D.M., and Insel, T.R. (1992). Oxytocin and social bonding.
Ann. N. Y. Acad. Sci *652*, 204–211.

42. Winslow, J.T., Hastings, N., Carter, C.S., Harbaugh, C.R., and Insel, T.R. (1993). A role for
central vasopressin in pair bonding in monogamous prairie voles. Nature *365*, 545–548.
https://doi.org/10.1038/365545a0.

43. Carter, C.S., Grippo, A.J., Pournajafi-Nazarloo, H., Ruscio, M.G., and Porges, S.W. (2008).
Oxytocin, vasopressin and sociality. Prog. Brain Res *170*, 331–336.
https://doi.org/10.1016/S0079-6123(08)00427-5.

44. Ophir, A.G., Gessel, A., Zheng, D.-J., and Phelps, S.M. (2012). Oxytocin receptor density is
associated with male mating tactics and social monogamy. Hormones and behavior *61*, 445–
453.

45. Shapiro, L.E., and Insel, T.R. (1992). Oxytocin receptor distribution reflects social
organization in monogamous and polygamous voles. Ann. N. Y. Acad. Sci *652*, 448–451.

46. Heinrichs, M., von Dawans, B., and Domes, G. (2009). Oxytocin, vasopressin, and human
social behavior. Front Neuroendocrinol *30*, 548–557.
https://doi.org/10.1016/j.yfrne.2009.05.005.

47. Insel, T.R., Wang, Z.X., and Ferris, C.F. (1994). Patterns of brain vasopressin receptor
distribution associated with social organization in microtine rodents. J. Neurosci *14*, 5381–
5392.

48. King, L.B., Walum, H., Inoue, K., Eyrich, N.W., and Young, L.J. (2016). Variation in the
oxytocin receptor gene predicts brain region–specific expression and social attachment.
Biological psychiatry *80*, 160–169.

49. Walum, H., and Young, L.J. (2018). The neural mechanisms and circuitry of the pair bond.
Nat Rev Neurosci *19*, 643–654. https://doi.org/10.1038/s41583-018-0072-6.

50. Williams, J.R., Insel, T.R., Harbaugh, C.R., and Carter, C.S. (1994). Oxytocin administered
centrally facilitates formation of a partner preference in female prairie voles (Microtus
ochrogaster). J. Neuroendocrinol *6*, 247–250.

51. Bales, K.L., and Carter, C.S. (2003). Developmental exposure to oxytocin facilitates partner
preferences in male prairie voles (Microtus ochrogaster). Behavioral neuroscience *117*, 854.

52. Keebaugh, A.C., and Young, L.J. (2011). Increasing oxytocin receptor expression in the
nucleus accumbens of pre-pubertal female prairie voles enhances alloparental responsiveness
and partner preference formation as adults. Hormones and Behavior *60*, 498–504.
https://doi.org/https://doi.org/10.1016/j.yhbeh.2011.07.018.

53. Romero, T., Nagasawa, M., Mogi, K., Hasegawa, T., and Kikusui, T. (2014). Oxytocin
promotes social bonding in dogs. Proc Natl Acad Sci U S A *111*, 9085–9090.
https://doi.org/10.1073/pnas.1322868111.

54. Nagasawa, M., Mitsui, S., En, S., Ohtani, N., Ohta, M., Sakuma, Y., Onaka, T., Mogi, K., and
Kikusui, T. (2015). Social evolution. Oxytocin-gaze positive loop and the coevolution of
human-dog bonds. Science *348*, 333–336. https://doi.org/10.1126/science.1261022.

55. Smith, A.S., Agmo, A., Birnie, A.K., and French, J.A. (2010). Manipulation of the oxytocin
system alters social behavior and attraction in pair-bonding primates, Callithrix penicillata.
Horm Behav *57*, 255–262. https://doi.org/10.1016/j.yhbeh.2009.12.004.

56. Duque, J.F., Leichner, W., Ahmann, H., and Stevens, J.R. (2018). Mesotocin influences
pinyon jay prosociality. Biol Lett *14*. https://doi.org/10.1098/rsbl.2018.0105.

57. Oldfield, R.G., and Hofmann, H.A. (2011). Neuropeptide regulation of social behavior in a
monogamous cichlid fish. Physiology & Behavior *102*, 296–303.
https://doi.org/https://doi.org/10.1016/j.physbeh.2010.11.022.

58. Klatt, J.D., and Goodson, J.L. (2013). Oxytocin-like receptors mediate pair bonding in a
socially monogamous songbird. Proc Biol Sci *280*, 20122396.
https://doi.org/10.1098/rspb.2012.2396.

59. Berendzen, K.M., Sharma, R., Mandujano, M.A., Wei, Y., Rogers, F.D., Simmons, T.C.,
Seelke, A.M., Bond, J.M., Larios, R., Goodwin, N.L., et al. (2023). Oxytocin receptor is not
required for social attachment in prairie voles. Neuron *111*, 787–796.

60. Young, L.J., Lim, M.M., Gingrich, B., and Insel, T.R. (2001). Cellular mechanisms of social
attachment. Hormones and behavior *40*, 133–138.

61. Tripp, J.A., Berrio, A., McGraw, L.A., Matz, M.V., Davis, J.K., Inoue, K., Thomas, J.W.,
Young, L.J., and Phelps, S.M. (2021). Comparative neurotranscriptomics reveal widespread
species differences associated with bonding. BMC Genomics *22*, 399.
https://doi.org/10.1186/s12864-021-07720-0.

62. Grinevich, V., Desarménien, M.G., Chini, B., Tauber, M., and Muscatelli, F. (2015).
Ontogenesis of oxytocin pathways in the mammalian brain: late maturation and psychosocial
disorders. Front. Neuroanat. *8*. https://doi.org/10.3389/fnana.2014.00164.

63. Newmaster, K.T., Nolan, Z.T., Chon, U., Vanselow, D.J., Weit, A.R., Tabbaa, M., Hidema, S.,
Nishimori, K., Hammock, E.A.D., and Kim, Y. (2020). Quantitative cellular-resolution map
of the oxytocin receptor in postnatally developing mouse brains. Nature Communications *11*,
1885. https://doi.org/10.1038/s41467-020-15659-1.

64. Yamamoto, Y., Cushing, B.S., Kramer, K.M., Epperson, P.D., Hoffman, G.E., and Carter,
C.S. (2004). Neonatal manipulations of oxytocin alter expression of oxytocin and
vasopressin immunoreactive cells in the paraventricular nucleus of the hypothalamus in a
gender-specific manner. Neuroscience *125*, 947–955.
https://doi.org/10.1016/j.neuroscience.2004.02.028.

65. Mitre, M., Saadipour, K., Williams, K., Khatri, L., Froemke, R.C., and Chao, M.V. (2022).
Transactivation of TrkB Receptors by Oxytocin and Its G Protein-Coupled Receptor. Front
Mol Neurosci *15*, 891537. https://doi.org/10.3389/fnmol.2022.891537.

66. Young, L.J., and Wang, Z. (2004). The neurobiology of pair bonding. Nat. Neurosci *7*, 1048–
1054. https://doi.org/10.1038/nn1327.

67. DeVries, A.C., and Carter, C.S. (1999). Sex differences in temporal parameters of partner
preference in prairie voles (Microtus ochrogaster). Can. J. Zool. *77*, 885–889.
https://doi.org/10.1139/z99-054.

68. Fraser, E.J., and Shah, N.M. (2014). Complex Chemosensory Control of Female
Reproductive Behaviors. PLOS ONE *9*, e90368.
https://doi.org/10.1371/journal.pone.0090368.

69. Aragona, B.J., Liu, Y., Curtis, J.T., Stephan, F.K., and Wang, Z. (2003). A Critical Role for
Nucleus Accumbens Dopamine in Partner-Preference Formation in Male Prairie Voles. J.
Neurosci. *23*, 3483–3490. https://doi.org/10.1523/JNEUROSCI.23-08-03483.2003.

70. Dölen, G., Darvishzadeh, A., Huang, K.W., and Malenka, R.C. (2013). Social reward
requires coordinated activity of nucleus accumbens oxytocin and serotonin. Nature *501*,
179–184. https://doi.org/10.1038/nature12518.

71. Ross, H.E., Freeman, S.M., Spiegel, L.L., Ren, X., Terwilliger, E.F., and Young, L.J. (2009).
Variation in Oxytocin Receptor Density in the Nucleus Accumbens Has Differential Effects
on Affiliative Behaviors in Monogamous and Polygamous Voles. J. Neurosci. *29*, 1312–
1318. https://doi.org/10.1523/JNEUROSCI.5039-08.2009.

72. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation
network analysis. BMC Bioinformatics *9*, 559. https://doi.org/10.1186/1471-2105-9-559.

73. Maury, E.A., Sherman, M.A., Genovese, G., Gilgenast, T.G., Kamath, T., Burris, S.J.,
Rajarajan, P., Flaherty, E., Akbarian, S., Chess, A., et al. (2023). Schizophrenia-associated
somatic copy-number variants from 12,834 cases reveal recurrent NRXN1 and ABCB11
disruptions. Cell Genom *3*, 100356. https://doi.org/10.1016/j.xgen.2023.100356.

74. Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.-Y., Peng, M.,
Collins, R., Grove, J., Klei, L., et al. (2020). Large-Scale Exome Sequencing Study
Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. Cell
180, 568-584.e23. https://doi.org/10.1016/j.cell.2019.12.036.

75. Fromer, M., Pocklington, A.J., Kavanagh, D.H., Williams, H.J., Dwyer, S., Gormley, P.,
Georgieva, L., Rees, E., Palta, P., Ruderfer, D.M., et al. (2014). De novo mutations in
schizophrenia implicate synaptic networks. Nature *506*, 179–184.
https://doi.org/10.1038/nature12929.

76. EuroEPINOMICS-RES Consortium, Epilepsy Phenome/Genome Project, and Epi4K
Consortium (2014). De novo mutations in synaptic transmission genes including DNM1
cause epileptic encephalopathies. Am J Hum Genet *95*, 360–370.
https://doi.org/10.1016/j.ajhg.2014.08.013.

77. Kaplanis, J., Samocha, K.E., Wiel, L., Zhang, Z., Arvai, K.J., Eberhardt, R.Y., Gallone, G.,
Lelieveld, S.H., Martin, H.C., McRae, J.F., et al. (2020). Evidence for 28 genetic disorders
discovered by combining healthcare and research data. Nature *586*, 757–762.
https://doi.org/10.1038/s41586-020-2832-5.

78. Inoue, K., Ford, C.L., Horie, K., and Young, L.J. (2022). Oxytocin receptors are widely
distributed in the prairie vole (Microtus ochrogaster) brain: Relation to social behavior,
genetic polymorphisms, and the dopamine system. J Comp Neurol *530*, 2881–2900.
https://doi.org/10.1002/cne.25382.

79. Lefevre, A., Benusiglio, D., Tang, Y., Krabichler, Q., Charlet, A., and Grinevich, V. (2021).
Oxytocinergic Feedback Circuitries: An Anatomical Basis for Neuromodulation of Social
Behaviors. Front. Neural Circuits *15*, 688234. https://doi.org/10.3389/fncir.2021.688234.

80. Harris, G.W. (1955). PITUITARY-HYPOTHALAMIC MECHANISMS. Arch NeurPsych *73*,
124. https://doi.org/10.1001/archneurpsyc.1955.02330080002002.

81. Ye, Q., Nunez, J., and Zhang, X. (2022). Oxytocin Receptor-Expressing Neurons in the
Paraventricular Thalamus Regulate Feeding Motivation through Excitatory Projections to the
Nucleus Accumbens Core. J. Neurosci. *42*, 3949–3964.
https://doi.org/10.1523/JNEUROSCI.2042-21.2022.

82. Zhang, B., Qiu, L., Xiao, W., Ni, H., Chen, L., Wang, F., Mai, W., Wu, J., Bao, A., Hu, H., et
al. (2021). Reconstruction of the Hypothalamo-Neurohypophysial System and Functional
Dissection of Magnocellular Oxytocin Neurons in the Brain. Neuron *109*, 331-346.e7.
https://doi.org/10.1016/j.neuron.2020.10.032.

83. Romanov, R.A., Tretiakov, E.O., Kastriti, M.E., Zupancic, M., Häring, M., Korchynska, S.,
Popadin, K., Benevento, M., Rebernik, P., Lallemend, F., et al. (2020). Molecular design of
hypothalamus development. Nature *582*, 246–252. https://doi.org/10.1038/s41586-020-2266-
0.

84. Seoyoung Son, Steffy B. Manjila, Kyra T. Newmaster, Yuan-ting Wu, Daniel J. Vanselow,
Matt Ciarletta, Todd E. Anthony, Keith C. Cheng, and Yongsoo Kim (2022). Whole-Brain
Wiring Diagram of Oxytocin System in Adult Mice. J. Neurosci. *42*, 5021.
https://doi.org/10.1523/JNEUROSCI.0307-22.2022.

85. Vaidyanathan, R., and Hammock, E.A.D. (2020). Oxytocin receptor gene loss influences
expression of the oxytocin gene in C57BL/6J mice in a sex- and age-dependent manner.
Journal of Neuroendocrinology *32*, e12821. https://doi.org/10.1111/jne.12821.

86. Kleiman, D.G. (1977). Monogamy in Mammals. The Quarterly Review of Biology *52*, 39–
69. https://doi.org/10.1086/409721.

87. Feldman, R. (2017). The Neurobiology of Human Attachments. Trends in Cognitive Sciences
21, 80–99. https://doi.org/10.1016/j.tics.2016.11.007.

88. Johnson, Z.V., Walum, H., Jamal, Y.A., Xiao, Y., Keebaugh, A.C., Inoue, K., and Young, L.J.
(2016). Central oxytocin receptors mediate mating-induced partner preferences and enhance
correlated activation across forebrain nuclei in male prairie voles. Horm Behav *79*, 8–17.
https://doi.org/10.1016/j.yhbeh.2015.11.011.

89. Takayanagi, Y., Yoshida, M., Bielsky, I.F., Ross, H.E., Kawamata, M., Onaka, T.,
Yanagisawa, T., Kimura, T., Matzuk, M.M., Young, L.J., et al. (2005). Pervasive social
deficits, but normal parturition, in oxytocin receptor-deficient mice. PNAS *102*, 16096–
16101.

90. Pobbe, R.L.H., Pearson, B.L., Defensor, E.B., Bolivar, V.J., Young, W.S., Lee, H.-J.,
Blanchard, D.C., and Blanchard, R.J. (2012). Oxytocin receptor knockout mice display
deficits in the expression of autism-related behaviors. Hormones and Behavior *61*, 436–444.
https://doi.org/https://doi.org/10.1016/j.yhbeh.2011.10.010.

91. Yang, C.F., and Shah, N.M. (2014). Representing sex in the brain, one module at a time.
Neuron *82*, 261–278. https://doi.org/10.1016/j.neuron.2014.03.029.

92. Mong, J.A., and Pfaff, D.W. (2004). Hormonal symphony: steroid orchestration of gene
modules for sociosexual behaviors. Mol Psychiatry *9*, 550–556.
https://doi.org/10.1038/sj.mp.4001493.

93. Mong, J., Easton, A., Kow, L.M., and Pfaff, D. (2003). Neural, hormonal and genetic
mechanisms for the activation of brain and behavior. Eur J Pharmacol *480*, 229–231.
https://doi.org/10.1016/j.ejphar.2003.08.109.

94. Barron, A.B., and Robinson, G.E. (2008). The utility of behavioral models and modules in
molecular analyses of social behavior. Genes Brain and Behavior *7*, 257–265.
https://doi.org/10.1111/j.1601-183X.2007.00344.x.

95. Kyriacou, C.P., and Hall, J.C. (1980). Circadian rhythm mutations in Drosophila
melanogaster affect short-term fluctuations in the male's courtship song. Proc Natl Acad Sci
U S A *77*, 6729–6733. https://doi.org/10.1073/pnas.77.11.6729.

96. Konopka, R.J., and Benzer, S. (1971). Clock Mutants of *Drosophila melanogaster*. Proc Natl
Acad Sci USA *68*, 2112–2116. https://doi.org/10.1073/pnas.68.9.2112.

97. Bendesky, A., Kwon, Y.-M., Lassance, J.-M., Lewarch, C.L., Yao, S., Peterson, B.K., He,
M.X., Dulac, C., and Hoekstra, H.E. (2017). The genetic basis of parental care evolution in
monogamous mice. Nature *544*, 434–439. https://doi.org/10.1038/nature22074.

98. Scheele, D., Striepens, N., Güntürkün, O., Deutschländer, S., Maier, W., Kendrick, K.M., and
Hurlemann, R. (2012). Oxytocin modulates social distance between males and females. J
Neurosci *32*, 16074–16079. https://doi.org/10.1523/JNEUROSCI.2755-12.2012.

99. Ditzen, B., Schaer, M., Gabriel, B., Bodenmann, G., Ehlert, U., and Heinrichs, M. (2009).
Intranasal Oxytocin Increases Positive Communication and Reduces Cortisol Levels During
Couple Conflict. Biological Psychiatry *65*, 728–731.
https://doi.org/10.1016/j.biopsych.2008.10.011.

100.   De Dreu, C.K.W., Greer, L.L., Handgraaf, M.J.J., Shalvi, S., Van Kleef, G.A., Baas, M.,
Ten Velden, F.S., Van Dijk, E., and Feith, S.W.W. (2010). The Neuropeptide Oxytocin
Regulates Parochial Altruism in Intergroup Conflict Among Humans. Science *328*, 1408–
1411. https://doi.org/10.1126/science.1189047.

101.   Osakada, T., Yan, R., Jiang, Y., Wei, D., Tabuchi, R., Dai, B., Wang, X., Zhao, G., Wang,
C.X., Liu, J.-J., et al. (2024). A dedicated hypothalamic oxytocin circuit controls aversive
social learning. Nature *626*, 347–356. https://doi.org/10.1038/s41586-023-06958-w.

102. Wang, Z., and Young, L.J. (1997). Ontogeny of oxytocin and vasopressin receptor
binding in the lateral septum in prairie and montane voles. Brain Res. Dev. Brain Res *104*,
191–195.

103. Smith, M.L., Asada, N., and Malenka, R.C. (2021). Anterior cingulate inputs to nucleus
accumbens control the social transfer of pain and analgesia. Science *371*, 153–159.
https://doi.org/10.1126/science.abe3040.

104. Gegenhuber, B., Wu, M.V., Bronstein, R., and Tollkuhn, J. (2020). Regulation of neural
gene expression by estrogen receptor alpha (Neuroscience)
https://doi.org/10.1101/2020.10.21.349290.

105. Cao, J., and Patisaul, H.B. (2013). Sex-specific expression of estrogen receptors α and β
and Kiss1 in the postnatal rat amygdala. J of Comparative Neurology *521*, 465–478.
https://doi.org/10.1002/cne.23185.

106. Mitsushima, D., Yamada, K., Takase, K., Funabashi, T., and Kimura, F. (2006). Sex
differences in the basolateral amygdala: the extracellular levels of serotonin and dopamine,
and their responses to restraint stress in rats. Eur J Neurosci *24*, 3245–3254.
https://doi.org/10.1111/j.1460-9568.2006.05214.x.

107. McCann, K.E., Sinkiewicz, D.M., Rosenhauer, A.M., Beach, L.Q., and Huhman, K.L.
(2019). Transcriptomic Analysis Reveals Sex-Dependent Expression Patterns in the
Basolateral Amygdala of Dominant and Subordinate Animals After Acute Social Conflict.
Mol Neurobiol *56*, 3768–3779. https://doi.org/10.1007/s12035-018-1339-7.

108. Trainor, B.C. (2011). Stress responses and the mesolimbic dopamine system: social
contexts and sex differences. Horm Behav *60*, 457–469.
https://doi.org/10.1016/j.yhbeh.2011.08.013.

109. Wright, E.C., Luo, P.X., Zakharenkov, H.C., Serna Godoy, A., Lake, A.A., Prince, Z.D.,
Sekar, S., Culkin, H.I., Ramirez, A.V., Dwyer, T., et al. (2023). Sexual differentiation of
neural mechanisms of stress sensitivity during puberty. Proc. Natl. Acad. Sci. U.S.A. *120*,
e2306475120. https://doi.org/10.1073/pnas.2306475120.

110. Dölen, G. (2015). Oxytocin: parallel processing in the social brain? J. Neuroendocrinol.
27, 516–535. https://doi.org/10.1111/jne.12284.

111. Choi, T.-Y., Jeon, H., Jeong, S., Kim, E.J., Kim, J., Jeong, Y.H., Kang, B., Choi, M., and
Koo, J.W. (2023). Distinct prefrontal projection activity and transcriptional state conversely
orchestrate social competition and hierarchy. Neuron, S0896-6273(23)00886-3.
https://doi.org/10.1016/j.neuron.2023.11.012.

112. Yaguchi, K., Hagihara, M., Konno, A., Hirai, H., Yukinaga, H., and Miyamichi, K.
(2023). Dynamic modulation of pulsatile activities of oxytocin neurons in lactating wild-type
mice. PLoS One *18*, e0285589. https://doi.org/10.1371/journal.pone.0285589.

113. Newmaster, K.T., Nolan, Z.T., Chon, U., Vanselow, D.J., Weit, A.R., Tabbaa, M.,
Hidema, S., Nishimori, K., Hammock, E.A.D., and Kim, Y. (2020). Quantitative cellular-
resolution map of the oxytocin receptor in postnatally developing mouse brains. Nat
Commun *11*, 1885. https://doi.org/10.1038/s41467-020-15659-1.

114. Vaidyanathan, R., and Hammock, E.A.D. (2017). Oxytocin receptor dynamics in the
brain across development and species. Developmental Neurobiology *77*, 143–157.
https://doi.org/10.1002/dneu.22403.

115. Manoli, D.S., and State, M.W. (2021). Autism Spectrum Disorder Genetics and the
Search for Pathological Mechanisms. AJP *178*, 30–38.
https://doi.org/10.1176/appi.ajp.2020.20111608.

116. Werling, D.M. (2016). The role of sex-differential biology in risk for autism spectrum
disorder. Biology of Sex Differences *7*, 58. https://doi.org/10.1186/s13293-016-0112-8.

117. Werling, D.M., and Geschwind, D.H. (2013). Sex differences in autism spectrum
disorders. Curr. Opin. Neurol. *26*, 146–153.
https://doi.org/10.1097/WCO.0b013e32835ee548.

118. Perissi, V., and Rosenfeld, M.G. (2005). Controlling nuclear receptors: the circular logic
of cofactor cycles. Nature Reviews: Molecular Cell Biology *6*, 542–554.
https://doi.org/10.1038/nrm1682.

119. Horie, K., Inoue, K., Nishimori, K., and Young, L.J. (2020). Investigation of Oxtr-
expressing Neurons Projecting to Nucleus Accumbens using Oxtr-ires-Cre Knock-in prairie
Voles (Microtus ochrogaster). Neuroscience *448*, 312–324.
https://doi.org/10.1016/j.neuroscience.2020.08.023.

120. Villanueva, C., Jacquier, S., and de Roux, N. (2012). DLK1 Is a Somato-Dendritic
Protein Expressed in Hypothalamic Arginine-Vasopressin and Oxytocin Neurons. PLoS One
7, e36134. https://doi.org/10.1371/journal.pone.0036134.

121. Insel, T.R. (1997). A neurobiological basis of social attachment. Am J Psychiatry *154*,
726–735.

122. Carter, C.S., and Perkeybile, A.M. (2018). The Monogamy Paradox: What Do Love and
Sex Have to Do With It? Front. Ecol. Evol. *6*, 202. https://doi.org/10.3389/fevo.2018.00202.

123. De Vries, G.J. (2004). Minireview: Sex Differences in Adult and Developing Brains:
Compensation, Compensation, Compensation. Endocrinology *145*, 1063–1068.
https://doi.org/10.1210/en.2003-1504.

124. Friard, O., and Gamba, M. (2016). BORIS: a free, versatile open-source event-logging
software for video/audio coding and live observations. Methods in ecology and evolution *7*,
1325–1330.

125. Risso, D., Ngai, J., Speed, T.P., and Dudoit, S. (2014). Normalization of RNA-seq data
using factor analysis of control genes or samples. Nat Biotechnol *32*, 896–902.
https://doi.org/10.1038/nbt.2931.

126. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

127. Stephens, M., Carbonetto, P., Gerard, D., Lu, M., Sun, L., Willwerscheid, J., and Xiao, N.
(2020). Methods for Adaptive Shrinkage, using Empirical Bayes [R package ashr version
2.2-47]. In.

128. Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology
analysis for RNA-seq: accounting for selection bias. Genome Biol *11*, R14.
https://doi.org/10.1186/gb-2010-11-2-r14.

129. Yu, G. (2020). Gene Ontology Semantic Similarity Analysis Using GOSemSim. In Stem
Cell Transcriptional Networks Methods in Molecular Biology., B. L. Kidder, ed. (Springer
US), pp. 207–215. https://doi.org/10.1007/978-1-0716-0301-7_11.

130. Pico, A.R., Kelder, T., Van Iersel, M.P., Hanspers, K., Conklin, B.R., and Evelo, C.
(2008). WikiPathways: Pathway Editing for the People. PLoS Biol *6*, e184.
https://doi.org/10.1371/journal.pbio.0060184.

131. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.,
et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The
Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

132. Fu, R., Gillen, A.E., Sheridan, R.M., Tian, C., Daya, M., Hao, Y., Hesselberth, J.R., and
Riemondy, K.A. (2020). clustifyr: an R package for automated single-cell RNA sequencing
cluster classification. F1000Res *9*, 223. https://doi.org/10.12688/f1000research.22969.2.

133. Zhang, B., and Horvath, S. (2005). A general framework for weighted gene co-expression
network analysis. Stat Appl Genet Mol Biol *4*, Article17. https://doi.org/10.2202/1544-
6115.1128.

134. Langfelder, P., Luo, R., Oldham, M.C., and Horvath, S. (2011). Is my network module
preserved and reproducible? PLoS Comput Biol *7*, e1001057.
https://doi.org/10.1371/journal.pcbi.1001057.

135. Gandrillon, O., Solari, F., Legrand, C., Jurdic, P., and Samarut, J. (1996). A rapid and
convenient method to prepare DIG-labelled RNA probes for use in non-radioactive in situ
hybridization. Molecular and cellular probes *10*, 51–56.

1729