Oxytocin receptor controls promiscuity and development in prairie voles

Ruchira Sharma[1,2,3,4†], Kristen M. Berendzen[1,2,3,4†], Amanda Everitt[1,3,10,11], Belinda

Wang[1,3], Gina Williams[1,2,3,4,5], Shuyu Wang[1,2,3,4], Kara Quine[1,2,3,4], Rose D.

Larios[1,2,3,4,5,14], Kimberly L. P. Long[1,2,3,4], Nerissa Hoglen[1,2,3,4,5], Bibi Alika

Sulaman[1,2,3,4,15], Marie C. Heath[1,2,3,4,16], Michael Sherman[1,2,3,4], Robert Klinkel[1,2,3,4,17],

Angela Cai[1,2,3,4,20], Denis Galo[1,2,3,4,], Lizandro Chan Caamal[1,2,3,4,18], Nastacia L.

Goodwin[1,2,3,4,19], Annaliese Beery[6,7], Karen L. Bales[8,9], Katherine S. Pollard[10,11,12], Arthur

Jeremy Willsey[1,13], Devanand S. Manoli[1,2,3,4,*]

[1]Department of Psychiatry and Behavioral Sciences

[2]Center for Integrative Neuroscience

[3]Weill Institute for Neurosciences

[4]Kavli Institute for Fundamental Neuroscience

[5]Neurosciences Graduate Program, University of California, San Francisco; San

Francisco, CA 95158, USA

[6]Department of Integrative Biology

[7]Department of Neuroscience, UC Berkeley, Berkeley, CA, United States

[8]Department of Psychology

[9]Department of Neurobiology, Physiology, and Behavior, University of California, Davis;

Davis, CA 95616, USA

[10]Gladstone Institutes, San Francisco, CA, USA

[11]Department of Epidemiology & Biostatistics, University of California, San Francisco,

San Francisco, CA, USA

[12]Chan Zuckerberg Biohub, San Francisco, CA, USA

[13]Calico Labs, San Francisco, CA 94080, USA

[14]Neurona Therapeutics, South San Francisco, CA 94080, USA

[15]Department of Neuroscience, University of Michigan, Ann Arbor, MI 48109, USA

[16]Department of Neurobiology and Behavior, University of California, Irvine; Irvine, CA

92697, USA

[17]Branch Out School, Los Angeles, CA 90026, USA

[18]University of California, Los Angeles; Los Angeles, CA 90095, USA

[19]Department of Psychology, University of Washington; Seattle, WA 98195, USA

[20]Department of Rehabilitation and Regenerative Medicine, Columbia University Irving

Medical Center; New York, NY 10032, USA

* Corresponding authors. Devanand.manoli@ucsf.edu

Lead Contact: Devanand.manoli@ucsf.edu

†These authors contributed equally to this work

Summary:

Oxytocin receptor (Oxtr) signaling influences complex social behaviors in diverse species, including social monogamy in prairie voles. How Oxtr regulates specific components of social attachment behaviors and the neural mechanisms mediating them remains unknown. Here, we examine prairie voles lacking Oxtr and demonstrate that pair bonding comprises distinct behavioral modules: the preference for a bonded partner, and the rejection of novel potential mates. Our longitudinal study of social attachment shows that Oxtr sex-specifically influences early interactions between novel partners facilitating the formation of partner preference. Additionally, Oxtr suppresses promiscuity towards novel potential mates following pair bonding, contributing to rejection. Oxtr function regulates coordinated patterns of gene expression in regions implicated in attachment behaviors and regulates the expression of oxytocin in the paraventricular nucleus of the hypothalamus, a principal source of oxytocin. Thus, Oxtr controls genetically separable components of pair bonding behaviors and coordinates development of the neural substrates of attachment.

Keywords:

Attachment behaviors, prairie vole, pair-bonding, oxytocin receptor, partner preference, monogamy, promiscuity, oxytocin signaling, paraventricular nucleus of the hypothalamus, nucleus accumbens

Introduction

Social attachments, such as the bonds between parents and children, family and kin, friends, and the long-term relationships between mates and romantic partners form the basis of interpersonal and group dynamics within societies[1–7]. Various selective pressures have given rise to diverse patterns of complex social organization involving long-term relationships between members of a species, across mammals and other taxa, one of the most robust and intriguing of which is the enduring bond between mates[2,4,8–13]. In mammals, ~4% of species demonstrate social monogamy, forming persistent pair bonds between mating partners[9,14–16]. Comparative studies between closely related species that display distinct patterns of these social behaviors, including social monogamy, reveal correlated species-specific differences in the nonapeptide hormones oxytocin (Oxt) and arginine vasopressin (Avp), or their homologs, suggesting that they serve as critical modulators of affiliative behaviors and pair bonding[9–11,16–20]. Despite the central role of social attachment in human behavior and a rich history of investigation into the cognitive processes underlying social cognition and affiliative behaviors, it has been difficult to probe the neural mechanisms underlying these fascinating behaviors at the cellular, molecular, and genetic level, as most species that serve as model organisms for such studies in the laboratory do not display long-term social attachment as adults [6,21–27].

Prairie voles (*Microtus ochrogaster*) display long-term social attachment such that mating partners show an enduring pair bond and social monogamy[13,28–31]. Like many complex innate behaviors, pair bonding consists of distinct modules: the development of increased prosocial behaviors with partners over strangers (partner preference), and increased

agonistic behaviors towards novel opposite sex potential mates (stranger rejection)[13,30,32–

37]. As in humans and other species that show affiliative social behaviors, disruption of pair

bonds results in increased anxiety-like behaviors and markers of stress, supporting

integrated neural and physiologic mechanisms that facilitate the preservation of such

attachments[38–40]. Pioneering work in prairie voles identified Oxt and Avp as critical

mediators of pair bonding[41,42]. Interspecific variations in the patterns of expression of the

oxytocin receptor (Oxtr) and the vasopressin 1a receptor (V1ar) correlate with the

potential for pair bonding within closely related vole species[43–49]. Consistently,

pharmacologic inhibition of Oxt and Avp signaling via their respective cognate receptors

disrupts pair bonding, while exogenous administration of these hormones promotes pair

bonding without mating[28,50,51].

Given the central role of Oxt signaling in affiliative behaviors, we tested the genetic

requirement for signaling via Oxtr for pair bonding and social behaviors in prairie voles.

We generated prairie voles lacking Oxtr and unexpectedly observed that these animals

continue to display a strong preference for partners following mating and cohabitation[52].

To gain deeper insight into the modulation of attachment and social behaviors by Oxtr

function, we developed a series of behavioral assays to probe distinct aspects of pair

bonding and associated social behaviors. To interrogate the molecular mechanisms

mediating social attachment, we profiled patterns of gene expression in the nucleus

accumbens (NAc), a key site of species differences associated with social monogamy

that is dramatically enriched for Oxt binding in prairie voles[45,53,54]. We find that Oxtr

controls the timing of pair bonding in prairie voles and promiscuity independent of partner

preference formation. Oxtr sex-specifically controls reciprocal social behaviors, such that

naïve wildtype (WT) males preferentially interact with wildtype females over those lacking

Oxtr, while wildtype females show no such bias towards WT males. Oxtr function

regulates patterns of gene expression in the NAc, as well as other regions of the vole

brain implicated in social and attachment behaviors, in distinct ways in each sex. Finally,

we find that Oxtr function regulates the development of Oxt and Avp expressing neurons

in the paraventricular nucleus, a principal source of these neuropeptides. Our studies

reveal genetically separable components of pair bonding and suggest that Oxtr signaling

sex-specifically participates in the development of the circuits underlying attachment

behaviors.

Results

Oxtr function facilitates the formation of partner preference

We previously found that Oxtr is not required for the display of partner preference

following 1 week of cohabitation[52]. However, acute modulation of Oxtr signaling can alter

patterns of social behaviors between individuals[29,50,55]. These observations suggest that

changes in Oxtr function may influence early social interactions that contribute to the

trajectory of pair bond formation and/or specific aspects of attachment behaviors following

the formation of partner preference. We first wished to determine whether Oxtr function

facilitated pair bond formation by examining the display of partner preference following

shorter periods of cohabitation. Consistent with previous studies showing that the

cohabitation time required for the display of partner preference differs between males and

females, we find that WT females demonstrate partner preference after only 6 hours of

cohousing with a WT male partner, while WT males require 5 days following introduction

to a potential WT mate (Fig. S1A-C)[56]. We therefore cohoused WT or Oxtr$^{1-/-}$ females with

WT males for 6 hours, and WT or Oxtr$^{1-/-}$ males with WT females for 5 days following

introduction (short cohabitation), or in these combinations for 1 or 7 days (long

cohabitation) for females or males, respectively (Fig. 1A, Fig. S1A).

Both males and females lacking Oxtr show an absence of partner preference following

short cohabitation with WT partners (Fig. 1B). Oxtr$^{1-/-}$ females, but not males, showed a

decrease in the fraction of huddling time with their partner and an increase in the fraction

of huddling time with a stranger compared to WT controls (Fig. 1C-D). Oxtr$^{1-/-}$ males

present an absence of partner preference (Fig. 1B), displayed due to both decreased time

with their cohoused partner and increased time with strangers (Fig. 1C-D). Consistent

with our previous studies, we find no difference between WT and Oxtr$^{1-/-}$ females when

the pairs are cohoused for the longer duration (Fig. S1B-F).

To further understand the dynamics of social interactions that contribute to the difference

in huddling behavior between WT animals vs their siblings lacking Oxtr following short

cohabitation, we quantified the cumulative time subjects spent huddling with their

cohoused partners or strangers over the course of the assay. Oxtr$^{1-/-}$ females show a

significant decrease in the amount of time spent huddling with partners in the last 30 mins

of the assay (150-180mins) (Fig. 1E). We also observe a significant increase in huddling

time of Oxtr$^{1-/-}$ males with strangers during the middle periods (75 – 130mins) (Fig. 1F),

suggesting that they engage in more early prosocial behavior with strangers than WT

males. Consistent with our previous work, we find no difference between cumulative

huddling times with either partner or stranger between WTs and Oxtr[1-/-] animals of both

sexes after long cohabitation periods (Fig. 1G-H)[45]. These observations demonstrate that

Oxtr function changes the temporal dynamics of social interactions with both familiar and

unfamiliar animal in a sex-specific manner. Taken together, our findings suggest that Oxtr

function acts early in the process of pair bonding to facilitate the formation of the partner

preference.

Oxtr suppresses promiscuous behaviors in a state dependent manner in females

Oxtr influences the dynamics of interactions with partners and strangers differently

between the sexes. We therefore wished to examine the detailed patterns of these

interactions separately in females and males (Fig. 2A). We examined the distribution of

huddles by WT vs Oxtr[1-/-] females following short cohabitation as a measure of prosocial

behavior displayed by these animals. We find that the top quartile of huddle durations

(long huddles) from each subject animal make up an average of ~90% of the total huddle

duration, functionally separating these interactions from incidental side-by-side contact

that may not constitute affiliative huddling (Fig. S2A Left). We classified long huddles as

partner huddles or stranger huddles and discovered that, consistent with the overall time

spent, the median huddle duration carried out by Oxtr[1-/-] females is significantly shorter

with partners and significantly longer with strangers, compared to WT females (Fig. 2B).

The longest huddle displayed with partner is significantly longer for WT females compared

to Oxtr[1-/-] females though we find no difference in the length of the overall longest huddle.

This is due to some Oxtr[1-/-] females conducting their longest huddle with the stranger

instead (Fig. S2B). Oxtr[1-/-] females additionally executed a larger number of huddles only

with the stranger, but not the partner, compared to WT controls (Fig. 2C). Taken together,

these results suggest that Oxtr function in females both facilitates prosocial behavior with

partners and suppresses such behavior with strangers during the early stages of pair

bond formation.

We next wished to determine whether the loss of Oxtr function influences patterns of

prosocial behaviors after the display of partner preference. We therefore examined

patterns of huddling, in particular the duration and frequency of these interactions,

following long cohabitation, a time point at which both WT and $Oxtr^{1-/-}$ animals display

robust preference for partners over strangers[52] (Fig. S1D). As with short cohabitation, the

top quartile of huddles (by duration) for each subject animal following long cohabitation,

described an average of 90% of total huddle time (Fig. S2A right). Following long

cohabitation, preference and huddling behavior in $Oxtr^{1-/-}$ females is indistinguishable

from their WT siblings (Fig. 2D-E). Comparing changes in the frequency of top quartile

huddles displayed by $Oxtr^{1-/-}$ females after short vs long cohabitation, we find that both

WT and $Oxtr^{1-/-}$ females show decreased huddle frequency with stranger following long

cohabitation (Fig. S2C-D). On comparing durations of long huddles, we find that $Oxtr^{1-/-}$

females significantly increase their median huddle duration with partner post long

cohousing (Fig. 2F-G). Thus, loss of Oxtr function in females reduces the duration of early

prosocial interactions (short cohabitation) with partners and strangers while Oxtr-

independent mechanisms eventually promote WT patterns of affiliative huddling and

partner preference following longer periods of cohabitation.

Our results suggest that Oxtr function influences specific aspects of early pair bonding

and attachment behaviors. We next wished to dissociate the influence of Oxtr on the

formation and display of partner preference from the display of agonistic rejection

behaviors towards novel strangers following cohabitation. We therefore developed a

battery of assays to examine initial social interactions between mating partners at early

and later stages of pair bonding, as well as stranger rejection following cohabitation

(Fig.2A). We examined the behavior of subject females from each genotype with WT

males when naive animals were first introduced to each other (Introduction), mating

behaviors following estrus induction (Timed Mating), prosocial interactions following

reunification with their bonded mate (Partner Reunification), and agonistic rejection of

novel, opposite sex strangers (Stranger Rejection) (Fig. 2A see Methods). We find no

differences in social behaviors during Introduction, Timed Mating and Partner

Reunification (Fig. 2H-J). In contrast, during Stranger Rejection, $Oxtr^{1-/-}$ females show

more social behaviors than WT females towards novel naive males in the absence of their

partner (Fig. 2K). Taken together, these studies demonstrate that Oxtr function regulates

specific components of pair bonding in females in a state-dependent manner, suppressing

promiscuous social interactions with stranger males after cohabitation sufficient for the

display of partner preference.

Oxtr function influences different prosocial behaviors across the stages of pair

bonding in males

Given the Oxtr-dependent difference in huddling behavior between males and females

after short cohabitation (Fig. 1C-D), we wished to assess sex differences in the patterns

of behaviors displayed by Oxtr[1-/-] animals and to further investigate Oxtr function in males

(Fig. 3A). Examining long huddles (as above, Fig. 2), we find that such huddles by Oxtr[1-/-]

males with their partners following short cohabitation are significantly shorter when

compared to WT siblings, contributing to diminished partner preference early in pair

bonding (Fig. 3B Left). Interestingly, while Oxtr[1-/-] males show no difference in the total

amount of time spent huddling with strangers when compared to WT siblings (Fig. 1D),

they display shorter and more frequent huddles towards them at this time (Fig. 3B, 3C

Right). In males, Oxtr thus influences the pattern as well as amount of early affiliative

behavior to distinct behavioral stimuli in different ways.

Oxtr[1-/-] males show no difference in time spent with either partners or strangers after long

cohabitation (Fig. S1E-F). However, these males show a persistent reduction in the

duration of their long huddles with strangers at this time point (Fig. 3D). Following long

cohabitation, WT and Oxtr[1-/-] males show no difference in the duration of long huddles

with partners and a reduction in the duration of long huddles with strangers (Fig. S3A-B).

We find no difference in the frequency of long huddles between Oxtr[1-/-] and WT males

after long cohabitation (Fig. 3E). Consistently, comparing Oxtr[1-/-] males' behavior following

short vs long cohabitation reveals that the frequency with which Oxtr[1-/-] males huddle with

strangers decreases following long cohabitation (Fig. 3F-G). Thus, specifically in males,

loss of Oxtr function does not change the overall time spent huddling with strangers after

short or long cohabitation but does cause a continuing disruption in the patterns of

affiliative behavior that comprises huddling with strangers.

Our observations suggest that the loss of Oxtr disrupts certain characteristics of prosocial

behaviors in males regardless of the time spent bonding with their partner. We next

wished to dissect whether such loss of function resulted in atypical affiliative or agonistic

behaviors across stages of bonding. We examined the behavior of Oxtr[1-/-] males and their

WT siblings in the same series of paradigms described previously to characterize the

formation and maintenance of their pair bonds (Fig. 3A). Oxtr[1-/-] males spend similar

amounts of time as WT males engaging in prosocial behaviors with partners regardless

of bonding state (Fig. 3H-J, S3E). Similar to our observations in Oxtr[1-/-] females, Oxtr[1-/-]

males spend more time engaging in prosocial behaviors with strangers in the absence of

their partners (Fig. 3K).

In contrast to time spent conducting mating behaviors with WT female partners (Fig. S3E),

Oxtr[1-/-] males demonstrate increased frequency of mating attempts when these females

are in estrus (Fig. 3I). Examining all instances of mounting, we find that Oxtr[1-/-] males

carry out shorter mounting bouts more frequently than WT males (Fig. S3F). To determine

whether the shorter bouts of mounting in Oxtr[1-/-] males influence the receptivity of their

partners, we compared the mount-to-intromission ratio for each animal[57] and find that

females are equally receptive to Oxtr[1-/-] males as their WT siblings (Fig. S3G). Thus,

consistent with our previous observations, Oxtr influences patterns, but not the amount

or success, of male mating behaviors, suggesting that such changes in males' behaviors

are equally able to elicit receptivity in females.

Compared to their WT siblings, naïve Oxtr$^{1-/-}$ males show decreased agonistic behaviors

towards an unfamiliar WT female upon first encountering her but display similar levels of

rejection of novel (unfamiliar) strangers upon bonding (Fig. S3C-D). Thus, Oxtr function

influences distinct components of males' attachment-related social behaviors towards

unfamiliar females at different points across pair-bonding, promoting agonistic displays

and lengthening mating bout durations when naïve, and suppressing promiscuous

prosocial behaviors upon bonding. Taken together our results suggest that Oxtr influences

patterns of social behaviors with future partners during the early phases of pair-bonding

differently in females and males, but upon bond formation, functions in both sexes to

suppress promiscuous affiliative behaviors towards strangers (Fig. S3H) without affecting

agonistic behaviors.

Oxtr sex-specifically influences social interactions between potential mates

Oxtr function influences patterns of social interactions between mates early during the

formation of a pair-bond (Short Cohabitation in both sexes, Introduction and Timed Mating

in males). We therefore investigated the effect of Oxtr loss on the choice made by a

potential mate. We developed a variation of the PPT three-chamber assay to present a

naive WT "chooser" with a choice between a naive opposite sex WT or naive opposite

sex Oxtr$^{1-/-}$ animal (Fig. 4A, Fig. S4A, see Methods). Strikingly, we find that Oxtr influences

the choice by a WT mate in a sex-specific manner. WT females choose WT or Oxtr$^{1-/-}$

males with equal frequency (Fig. 4B). In contrast, WT male choosers show significant

preference for WT females compared to Oxtr$^{1-/-}$ females, particularly during the middle of

the assay (150-280 mins) (Fig. 4B-C, Fig. S4B). These observations suggest that

disruption of Oxtr function specifically in females may alter their behavior such that WT

males prefer WT future partners over such mutants.

While naïve WT females show no difference in overall social choice between naive WT

and Oxtr$^{1-/-}$ males (WT-Oxtr$^{1-/-}$, Fig. 4B), they may show distinct behavioral dynamics in

the presence of an Oxtr$^{1-/-}$ male. To assess this, we compared the WT-Oxtr$^{1-/-}$ condition

with a control group in which the naive female chooser is given access to two naive WT

males (WT-WT) (Fig. 4D). We designated the tethered animal with whom the subject

spent <u>M</u>ore <u>of</u> <u>A</u>ssay <u>T</u>ime ("chosen") as the MOAT. Intriguingly, we found that the

presence of an Oxtr$^{1-/-}$ male (WT-Oxtr$^{1-/-}$) influenced the behavior of the WT female

chooser, such that they spent significantly more time with WT or Oxtr$^{1-/-}$ MOAT males

compared to the time spent with WT MOAT males from the WT-WT condition (Fig. 4E).

Consistent with this finding, naive WT female choosers break contact more frequently in

WT-WT conditions when compared to WT-Oxtr$^{1-/-}$ (Fig. 4F). Similarly, the divergence in

the amount of cumulative time spent with the MOAT for WT female choosers occurs after

5 hours have elapsed (320-360 mins) suggesting that complex interactions with males

lacking Oxtr influence how females "choose" a male partner regardless of whether they

end up choosing a WT or Oxtr$^{1-/-}$ MOAT (Fig. 4G). In contrast, we find no difference in the

contact time or number of such transitions when comparing naive WT male choosers with

MOAT females of a WT-WT vs WT-Oxtr$^{1-/-}$ condition (Fig. 4E-F).

We next examined the pattern of a WT chooser's interactions with both chosen (MOAT)

and unchosen animals (<u>L</u>ess <u>of</u> <u>A</u>ssay <u>T</u>ime, LOAT) over time using a sliding preference

index (see Methods). We observed periods of convergence when choosers from a

condition (WT-WT or WT-Oxtr$^{1-/-}$) displayed significantly concurrent interactions with their

MOAT, indicated by the black bar above the trace. WT female choosers show a longer

and more consistent preference for their MOAT (regardless of genotype) in WT-Oxtr$^{1-/-}$ vs

WT-WT conditions, whereas WT male choosers show an earlier onset but equivalent

length of preference for MOATs in the WT-Oxtr$^{1-/-}$ condition compared with the WT-WT

condition (Fig. 4H). These observations suggest that interactions with males lacking Oxtr

influence the behavior of the WT females making a social choice, even when these

females interact more with and eventually choose a WT male. Taken together, loss of Oxtr

leads to differences in prosocial behaviors during initial interactions with WT naive

animals of the opposite sex that may later impact aspects of pair bonding.

Finally, to determine if naive prairie voles lacking Oxtr display generalized changes to

their behaviors independent of attachment-associated social behavior, we examined

general pro-social, anxiety-related, and locomotor behaviors. With the exception of an

increase in locomotion displayed by Oxtr$^{1-/-}$ females when compared to WT females in

an open field paradigm, we found no differences between Oxtr$^{1-/-}$ and WT voles of either

sex (Fig. S4C).

We have examined social interactions during the initial contact between mates, the

early stages of pair bonding, and during social attachment behaviors following partner

preference formation. Consistent with our findings and previous work, Oxtr function is

not required for the eventual display of partner preference. Here, we demonstrate that

Oxtr function sex-specifically facilitates early social interactions that contribute to the

formation of partner preference, and controls promiscuity.

Oxtr controls the molecular signature of pair bonding in the nucleus accumbens

We previously observed that Oxtr function is not required for the formation of partner

preference and here demonstrate that it suppresses prosocial behaviors towards novel

strangers following pair bonding. These results suggest that distinct components of pair

bonding are differentially influenced by Oxtr function or compensated upon its loss. We

next characterized the molecular signature associated with pair bonding to determine

whether changes in gene expression following pair bond formation are controlled by Oxtr.

Comparative studies demonstrate that Oxt binding in multiple brain regions differs

significantly between promiscuous and monogamous rodent species. Oxt binding is

particularly enriched in the NAc in prairie voles, while vasopressin binding is largely

absent in this area[45,47]. Thus, analyzing gene expression in the NAc allows us to identify

bonding-related expression that is primarily due to Oxtr function and largely independent

of the direct consequences of AVP signaling. Additionally, given the well-established role

of the NAc in the association of reward or aversion with specific sensory and

environmental cues[58–60] this region is well poised to play a role in the development of

partner preference, even without Oxtr signaling, as well as stranger rejection, which

appears to be dependent upon Oxtr function.

We performed bulk RNA-sequencing on NAc tissue from WT and Oxtr[1-/-] animals of both

sexes that were either: group housed with same-sex peers and sexually naïve (pre-

pairing); or paired with WT opposite-sex partners for six days, using our timed mating

paradigm (post-pairing) (Fig. 5A). PCA showed robust separation of WT and Oxtr[1-/-]

samples (Fig. S5A). Differential expression (DE) analysis of pairing condition, regardless

of genotype or sex, identified 13 genes that show significant changes in expression post-

pairing (Fig. S5B), all of which showed decreased expression (Fig. 5B). These genes

include *Agt, Sgk1, Ptgds, Fosb, Pdk4*, and *Acsm5* (Fig. 5C) and, while these genes have

not previously been linked to shared gene regulatory or functional pathways, the predicted

protein-protein interaction network for the 13 DE genes shows more connections than

would be expected by chance (p=2.47 e^{-05}) (Fig. S5C). Intriguingly, even in the limited set

of genes with altered expression in the context of pair bonding, changes are driven

primarily by expression in WT females (Fig. 5C, S5D). Subgroup analyses by genetic

background and sex identified additional genes including *Adora2a, Fkbp5*, and *Spred3* in

WT females and collagen-related genes, including *Col4a1* and *Col4a2*, in Oxtr[1-/-] females

(Fig. 5D, S5D). Only one gene, *Sema3b,* is significantly different in Oxtr[1-/-] males following

pairing (Fig. 5D).

To further examine changes in patterns of gene expression in the NAc associated with

pair bonding, we performed weighted gene coexpression network analysis (WGCNA)[61]

and calculated three sets of modules (M) from different sample subsets (modules[WT] ,

modules[Mut], and modules[All]) (Fig. S5F, S6D). Using the modules[WT] subset, we identified

15 coexpression modules (Fig. S5F, Table 2) representing genes that share highly similar

expression patterns within the NAc. We identified one module (Mpink[WT]) that significantly

and specifically correlated with pre-pairing status in WT animals (Pearson R correlation

of 0.7, Bonferroni adjusted p-value (p_{adj})=0.018) (Fig. 5E, S5E-F). Genes with high

module membership for the Mpink[WT] are also those with high gene significance for pairing

condition supporting the module's association with behavior condition (Fig. 5SG). Further,

genes significantly differentially expressed with bonding are significantly enriched in

Mpink[WT] (Bonferroni p_{adj}=7.44 e^{-08}) and serve as hub genes within this module (Fig. S5E,

G). These results suggest that the Mpink[WT] module might be driving, or under the

influence of pathways driving, the pre-pairing to post-pairing state change.

We asked whether any of the modules[WT] identified were regulated by Oxtr and thus lost

in Oxtr null animals. To address this, we assessed how well modules[WT] are preserved in

mutant samples. We calculated a $Z_{summary}$ measure that combines module density and

intramodular connectivity metrics into a composite statistic. We find that of the modules[WT],

Mpink[WT] had the lowest $Z_{summary}$ ($Z_{summary}$ = 2) in Oxtr[1-/-] samples (Fig. 5F), indicating that

dynamics of gene expression that correlate with bonding states are lost in the absence of

Oxtr. Finally, the module eigengene (ME) for Mpink[WT] when calculated for all 30 samples

and compared between pre- and post-pairing, was significant only for WT females

(Wilcoxon rank sum, Benjamini-Hochberg p_{adj}=0.03) and not for WT males or Oxtr[1-/-]

animals of either sex (Fig. 5G). These observations suggest there is sex-specific,

coordinated regulation of gene expression in the NAc following pairing of mates that may

be associated with pair bond associated behaviors.

Loss of Oxtr sex-specifically disrupts patterns of gene expression associated

with neurodevelopmental disorders

We next wished to determine how Oxtr signaling contributes to changes in gene expression in the NAc associated with the naive state or the process of pair bonding (Fig. 6A) by comparing the transcriptional profile of WT and Oxtr[1-/-] animals of each sex. We observed significant genotype specific differences in gene expression in animals from both sexes, with 466 genes showing differences between WT and Oxtr[1-/-] females, 270 between WT and Oxtr[1-/-] males, and 1,014 genes when samples from both sexes were combined (using a p_{adj}<0.05 and absolute ($\log_2$(FC)) > 0.25 as criteria). Of the combined DE gene set, 370 genes are upregulated and 644 are downregulated (Fig. 6B, S6A-B). Genes with the greatest fold change across both sexes include *Slc2a3, Kcna1, Rgs8, Cnksr2, Acvr*, and *Cnot6* (Fig. 6B). Genes specifically downregulated in Oxtr[1-/-] females with $\log_2$(FC) less than 1 include *Abcb11* (a gene recently associated with treatment resistant schizophrenia[62]), *Ngfr, Pon3,* and *Shh*. While there was no significant difference in Oxtr transcript levels by genotype (p_{adj} = 0.753)—consistent with minimal nonsense-mediated decay despite a premature stop codon—we have previously shown that no functional protein is produced in Oxtr[1-/-] animals[52]. V1aR showed a modest decrease in expression in Oxtr[1-/-] animals (p_{adj}=0.03) (Fig. S6C), though this is likely influenced by variability in WT samples. WGCNA of all samples identified two modules (Mbrown[All] and Mturquoise[All]) that were significantly positively correlated with the WT genotype compared to mutant (Pearson R = 0.55, p_{adj} = 0.011 for brown and Pearson R = 0.8, p_{adj} = 6.6e-7 for turquoise) (Fig. S6D). None of the identified modules showed correlation with pairing

condition (Fig. S6D). Our results thus reveal a dramatic change in patterns of gene

expression in the NAc with global loss of Oxtr signaling, independent of bonding state.

Examination of the sets of genes whose expression is altered by the loss of Oxtr,

regardless of bonding state or sex, revealed several distinct cellular processes regulated

by Oxtr signaling. These processes comprise larger domains of cell function including

extracellular channel binding activity, axoneme assembly complex, axon glial

ensheathment development, and metal cation transport (Fig. 6C), as well as pathways

for G protein signaling and dopaminergic neurogenesis (Fig. S6E). When analyzed by

sex, more of the identified processes were enriched in female Oxtr gene sets than in male

ones (female: 19/62, male: 6/62). To further link differentially expressed genes to

behavioral traits, we asked whether genes altered by loss of Oxtr are selectively enriched

for high-confidence associations with autism spectrum disorders (ASD) and other

neurodevelopmental diseases (NDD) affecting social attachment in humans as compared

to epilepsy, a disorder that does not disrupt social attachment[63–66]. Consistently, we find

that the genes upregulated with loss of Oxtr ($p_{adj}<0.01$, $\log_2(FC)>0$) are specifically

enriched for ASD- and NDD-associations but not epilepsy associations (Fig. 6D "Joint").

This enrichment is again primarily driven by genes upregulated in females and not males

(Fig. 6D). Thus, similar to the factors that influence gene expression changes associated

with pair bonding, many of the changes we identify in the NAc resulting from loss of Oxtr

are primarily driven by female-specific effects that may play a conserved role in mediating

social attachment in humans.

Oxtr regulates coordinated gene expression across multiple neural populations

implicated in social attachment behaviors

Given the expression of Oxtr in regions of the brain known to mediate social behavior in voles and other species[67,68], we wished to determine if genes whose expression is altered in the NAc by the loss of Oxtr were expressed not only in specific populations of cells within this region, but in other regions of the prairie vole brain implicated in social and attachment behaviors. We therefore examined the expression of 15 genes that showed significant changes in expression in WT females when compared to those lacking Oxtr using *in situ* hybridization (ISH) to both validate our RNA-seq findings as well as examine the expression patterns of these genes. Several of these genes, such as phosphodiesterase 10A (*Pde10A)* and diacylglycerol kinase beta (*Dgkb)*, show robust expression throughout the striatum (data not shown), while genes such as the calcitonin receptor (*Calcr*) and delta like non-canonical notch ligand 1 (*Dlk-1*) are expressed in restricted populations of cells in the NAc (Fig. 6F-H, top panels). The expression of these genes is significantly reduced in Oxtr[1-/-] females (Fig. 6E-H), supporting the specificity and resolution of our RNA-seq findings.

In parallel to our examination of gene expression in the NAc, our ISH revealed that loss of Oxtr in females altered the expression of the genes we examined in other regions of the prairie vole brain. We examined the expression of these genes in the lateral septum (LS), a region that has been implicated in agonistic and attachment behaviors in prairie voles and other rodents and shows species dimorphic patterns of Oxt and Avp binding. We observe that both *Calcr* and *Dlk-1* are decreased in the absence of Oxtr (Fig. 6G, H

middle panel) in the LS. We also find that expression of *Calcr* and *Dlk-1* is decreased in

the paraventricular nucleus of the hypothalamus (PVN), a principal source of Oxt and Avp

in the rodent brain which sends projections to the NAc (Fig. 6G-H bottom panels)[69–71].

These findings suggest that Oxtr function regulates patterns of gene expression in

multiple regions of the brain implicated in social and attachment behaviors in a

coordinated and region-specific manner.

Loss of Oxtr results in the decrease in density of Oxt and Avp expressing

neurons in the PVN

We find here that loss of Oxtr results in coordinated changes in gene expression in

multiple neural populations, including the PVN. We therefore wished to determine if the

signatures of gene expression regulated by Oxtr in the NAc in prairie voles (ONP set),

regardless of sex and bonding state, were enriched for patterns of gene expression

associated with distinct neural populations in other regions of the brain, particularly those

implicated in social and attachment behaviors. We compared the ONP gene set to the

molecular signatures of 33 hypothalamic subtypes previously identified by single-cell RNA

sequencing in mice[72]. Of the 33 clusters examined, the four with highest overlap with the

ONP gene set represent subsets of PVN neurons, in particular those that express Oxt

and AVP (Fig. 7A, clusters 43 and 26), as well as PVN neurons that express *Dlk1* and

Calcr (cluster 29), both of which show reduced expression in the NAc and PVN in females

lacking Oxtr (Fig. 6G-H bottom panel).

We next sought to determine if the overlap between Oxtr regulated genes and gene

expression patterns of neural populations in the PVN correlated with changes in the

expression of neuropeptides within PVN itself. We examined patterns of Oxt and Avp

expression using immunohistochemistry in naive adults and compared patterns of

expression between WT animals of both sexes and those lacking Oxtr (Fig. 7B, Fig.

S7A-B). We find that the density of both Oxt and Avp producing neurons is lower in

Oxtr$^{1-/-}$ animals (Fig. 7C-E). To determine if the loss of Oxt or Avp expression was

restricted to specific populations of PVN neurons, we examined the expression of these

peptides across the anterior-posterior axis of the PVN in both WT and Oxtr$^{1-/-}$ males and

females. We find that animals of both sexes show a preferential loss of Oxt and Avp

expressing neurons in the anterior PVN (Fig. 7F-G). Finally, we examined patterns of

Oxt and Avp over the course of development to determine when loss of Oxtr function

influences the development of their expression. We find that the density of Oxt neurons

at P0 (Fig. S7C-F) and both Oxt and Avp neurons at P21 is lower in Oxtr$^{1-/-}$ females

compared to age matched WT females (Fig. S7G-J). In contrast, we find that Oxt and

Avp neuron density loss occurs only in adult Oxtr$^{1-/-}$ males (Fig. 7D-G, Fig. S7B). These

observations suggest the coordinated, region-, and sex-specific influence of Oxtr

function on the development of neurons expressing Oxt (and Avp) within the PVN,

consistent with a trophic role for this pathway identified in mice[73–76].

Discussion

Robust reproductive behaviors are essential for survival. Accordingly, species have

evolved diverse strategies, including social monogamy, which mediate locating,

identifying, and attracting mates, facilitate reproduction, and promote the survival of

progeny[16,77]. Signaling via Oxtr influences a wide range of physiology and behavior

associated with social interactions and attachment between conspecifics, including pair

bonding between prairie vole mates[41,78]. Our recent findings that Oxtr is not genetically

required for the formation of partner preference in prairie voles suggests that, like other

genes and neuromodulatory pathways across diverse species, Oxtr likely functions in

parallel to multiple genetically specified pathways to facilitate pair bonding by influencing

specific behavioral modules that contribute to the complex behaviors that comprise social

attachment[33,35,79–85].

Here we demonstrate that loss of Oxtr in prairie voles influences the dynamics of social

interactions between sexually naïve animals and WT potential mates in distinct social

contexts in sex-specific ways. When given a choice, naive WT males display a preference

for WT females rather than for those lacking Oxtr, supporting the role of Oxtr in controlling

behavior during an early, sensitive period in bond formation[30]. The presence of an Oxtr[1-

/-] male alters levels and patterns of early prosocial behavior exhibited by WT females,

without ultimately affecting their choice of potential partner. Taken together, these

observations suggest that Oxtr may modulate social behaviors that act early in a sensitive

pair bond formation period, and via these behaviors or yet-to-be-determined sensory cues

such as pheromone profiles or patterns of vocal communication, influences the reciprocal

behavior of partners.

Animals lacking Oxtr take longer to display partner preference, and despite strong partner

preference following longer cohabitation, demonstrate promiscuous prosocial behaviors

towards novel potential mates, including huddling with strangers. These observations are

consistent with a state-dependent role of Oxtr function, similar to findings in multiple

species, including humans[86-88]. Oxtr function may thus promote prosocial behavior in the

naïve state to facilitate attachment with an unfamiliar potential mate, but following pair

bond formation, suppresses further prosocial interactions to new individuals, or even

promotes further separation from and rejection of unfamiliar individuals or groups. Across

species and outside of the context of pair bonding, oxytocin signaling may play a

conserved role in mediating other state-specific aversive behavior, for example, defeat-

induced social avoidance learning[89].

Our findings reveal that patterns of social behaviors contributing to 1) pair bond formation,

2) partner preference, and 3) rejection of novel mates are genetically separable

components of pair bonding representing distinct behavioral modules regulated by

mechanisms that require Oxtr function, i.e. reciprocal behaviors and stranger rejection, or

those that can be displayed in its absence, i.e. partner preference. Further work will allow

us to determine the Oxtr-independent mechanisms that mediate the closure of an early

sensitive period and facilitate the display of partner preference[52]. Where Oxtr functions

to control the switch from prosocial behavior towards a potential mate in naïve animals to

the suppression of such behaviors towards, and active rejection of, opposite-sex

conspecific strangers following bonding remains to be determined.

571 Examination of the molecular signatures of pair bonding and the role of Oxtr in regulating

572 these changes in gene expression in the NAc reveals modest, Oxtr-dependent changes

573 in the expression of a specific network of genes in WT animals following pair bonding[45,90].

574 Intriguingly, loss of Oxtr preferentially impacts patterns of gene expression in the NAc in

575 females. The NAc demonstrates little endogenous sex steroid hormone receptor

576 expression difference in gene expression in prairie voles and other species[67,91–93]. Sex

577 differences in the effects of Oxtr on gene expression this region may therefore suggest

578 that there are significant sex-differences in the inputs into this region that act through Oxtr

579 during development[94–99]. The larger impact of loss of Oxtr in females is consistent with its

580 more general conservation in mammals to regulate aspects of female physiology such as

581 lactation and prosocial behaviors or sensory cues that promote interactions with males[100].

583 Genes with altered expression in the absence of Oxtr are enriched for genes associated

584 with neuropsychiatric disorders, in particular neurodevelopmental disorders and ASD.

585 Given the sex-biased enrichment of these genes, such developmental differences in the

586 impacts of Oxtr function may contribute to sex differences in the pathophysiology of these

587 disorders[101–103]. Accordingly, the molecular signatures of attachment in the NAc, both

588 dependent upon and independent of Oxtr function, are subtle relative to these sex-biased

589 differences in the role of Oxtr and possibly reside in other neural populations that function

590 in attachment.

592 Validation of changes in gene expression in the NAc resulting from loss of Oxtr suggests

593 coordinated regulation of these genes by Oxtr across multiple brain regions implicated in

social and attachment behaviors. Oxtr may be one of multiple factors that organizes

specific components of the circuits underlying these displays to control discreet modules

of behavior. Consistent with a trophic role for Oxtr function during development, we find

that the genes regulated by Oxtr within the NAc are enriched for those that identify cell

types in the PVN, a principal source of neuropeptides implicated in pair bonding, which

sends oxytocinergic projections to the NAc[68,70,104–106]. Loss of Oxtr preferentially impacts

Oxt and Avp expression in the anterior PVN, suggesting that specific populations of these

cells may be more dependent upon such trophic function for their development, survival,

or expression of these hormones. Whole brain analysis will likely reveal if such neurons

have distinct projections when compared to those that remain in the absence of Oxtr,

contributing to our understanding of how these circuits and pathways regulate distinct

aspects of social and attachment behaviors and physiology[4,78,107].

Taken together, our behavioral and molecular studies reveal previously unappreciated

nuances in the control of social behavior and pair bonding by Oxtr function. They uncover

intriguing sex differences in the patterns of behavior influenced by loss of Oxtr function.

These observations suggest that circuits with sex-specific patterns of development, or

even function in ancestral species, may mechanistically converge in the context of Oxtr

function to generate displays of reciprocal interactions, pair bonding and attachment

behaviors in WT prairie voles that are similar between males and females[108,109].

Manipulation of Oxtr expression at specific developmental timepoints and in specific

neural populations will help to elucidate how this pathway influences social and

attachment behaviors in the context of normal development to regulate distinct

components of pair bonding. Understanding the specific neural populations that express

Oxtr and other molecular pathways that contribute to pair bonding will allow us to

understand the mechanisms controlling the early experience-dependent, sensitive period

of bond formation. In parallel, examination of the activity in these populations following

manipulations of Oxtr function will allow us to determine the circuit mechanisms by which

similar but distinct sensory cues contribute to the identification of partners or novel

conspecifics and consequently, the display of dramatically different patterns of social

behaviors. The sum of these mechanisms represents the preference for partners over

strangers and reflects the formation and maintenance of pair bonds between mates,

affiliation between individuals, and the behavioral consequences of identifying others as

unfamiliar or outside of these bonds.

Acknowledgments

The authors would like to thank members of the Manoli lab for assistance, advice,

helpful discussion and comments on the manuscript. J. Tollkuhn and M. Brainard

provided valuable insights to the structure, content and design of the manuscript.

Additionally, we would like to thank S. Garg for his assistance in data analysis and help

designing the analytic pipeline. The authors received funding from National Institutes of

Health grant R01MH123513 (D.S.M.), National Science Foundation grant 1556974

(D.S.M.), Burroughs Wellcome Fund 1015667 (D.S.M.), Whitehall Foundation grant

2018-08-83 (D.S.M.), One Mind Foundation A137726 (D.S.M.), National Institutes of

Health grant R25MH060482 (K.M.B.), AP Giannini Foundation Fellowship P0534952

(K.M.B.), Larry L. Hillblom Foundation Fellowship 2020-A-023-FEL (K.M.B.), and

National Institute of Mental Health grant R01MH123178 (K.S.P.)

Author contributions

Conceptualization, R.S., K.M.B., K.L., and D.S.M.; methodology, A.B., A.J.W., D.S.M.,

K.L.B., K.M.B., K.S.P., and R.S.; software, K.Q., and S.W.; formal analysis, A.E., R.S.,

and K.M.B.; investigation, R.S., K.M.B., A.E., B.W., G.W., S.W., K.Q., R.D.L., K.L., N.H.,

B.A.S., M.C.H., M.S., R.K., A.C., D.G., L.C.C., N.L.G.; visualization, R.S., K.M.B., B.W.,

and A.E.; funding acquisition, D.S.M., K.S.P., and A.J.W; writing– original draft, R.S., and

K.M.B.; writing-review & editing, R.S., K.M.B., D.S.M., N.H., K.S.P., A.E., and A.B.

Declaration of interests

650 The authors declare no competing interests.

Star Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
DAPI	Sigma-Aldrich	Cat #D9542; CAS #28718-90-3

Experimental models: Organisms/strains		
Prairie vole: OxtR[1-/-]	Berendzen et al.	
Oligonucleotides		
ISH probes	This paper	N/A
Oxtr[1-/-] genotyping oligos	This paper	N/A
Antibodies		
Oxytocin Polyclonal	Invitrogen	Cat#PA5-26701; RRID
Anti-Neurophysin 2/NP-AVP	Millipore	Cat#MABN845; RRID
Cy™3 AffiniPure Donkey Anti-Rabbit IgG	Jackson ImmunoResearch	Cat #711-165-152; RRID
Alexa Flour 488 Donkey Anti Mouse IgG	Thermo Fisher Scientific	Cat #711-165-152; RRID
Software and algorithms		
ImageJ	NIH	; RRID: SCR_003070
R (version 1.1.423)	R Core	https://www.r-project.org/
GraphPad Prism (version 7.03)	GraphPad Software	
Boris	Boris	www.boris.unito.it

		https://www.mathworks.com/products.ht
MATLAB	MathWorks	ml
		RRID: SCR_001622

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Devanand Manoli (devanand.manoli@ucsf.edu).

Materials availability

All unique reagents generated in this study are available from the lead contact upon

request.

Data

Transcriptomics data discussed in this publication are accessible through GEO Series

accession number GSE279248.

Code

All code used for RNA-seq data processing and differential expression analyses, as well

as post-processed expression data, can be found at

https://github.com/aseveritt/OXTR_prairie_voles

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals

Subjects were laboratory-bred prairie voles (*Microtus ochrogaster*) which originated through systematic outbreeding of a wild stock captured near Champaign, Illinois. CRISPR/Cas9 editing was used to generate 2 independent null alleles, Oxtr[1-/-] and Oxtr[2-/-], both of which were validated using autoradiography to show loss of receptor function[52] (data not shown). Oxtr[1-/-] was backcrossed for >7 generations to mitigate any possible off-target effects and this line was used for all subsequent experiments. Sexually naïve male and female animals were group weaned at 21 ± 1 days and separated to group housing with same-sex siblings and age-matched same-sex non-siblings. Voles were maintained under a 14:10 h light-dark cycle in clear plastic cages (45 × 25 × 15 cm) with bedding, nesting material (nestlet), and a PVC hiding tube. Rooms were maintained at approximately 20°C, and food and water were available *ad libitum*.

Breeding pairs were established between two heterozygotes or a homozygous Oxtr[1-/-] mutant male and heterozygous female partner from a breeding line maintained in the lab. Voles were randomly assigned into experimental groups when they reached 7-9 weeks of age at the start of testing. This study was carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals published by the National Research Council. The protocol was approved by the Institutional Animal Care and Use Committee at each respective institution.

Genotyping

The following primers were used to amplify a region where XcmI restriction for Oxtr[1] or

Bpu10I restriction for Oxtr[2] determined the genotype of all animals studied. XcmI cuts

the mutant allele in the case of Oxtr[1] and Bpu10I restricts the WT sequence for Oxtr[2].

(Forward: ACTGGAGCTTCGAGTTGGAC, Reverse: ATGCCCACCACTTGCAAGTA)

METHOD DETAILS

Behavior

Triadic naive choice assay

The linear three chamber partner preference apparatus with open top and 10 x 32 inch

walls[52] was modified to capture the interactions of the naive, freely moving WT subject

and the naive, tethered, opposite sex potential partner (either 2 WTs or a WT and an

Oxtr[1-/-]) at either end of the chamber. Behaviors were recorded for 6 hours using a top

view camera and the videos generated were analyzed using a contour detection

algorithm. Specifically, contours were detected on individual video frames using a

threshold-based strategy that took advantage of the contrast between dark-colored vole

fur coats and light-colored bedding and chamber apparatus. A standalone vole was

detected as a single contour, and two voles in physical contact were likewise detected

as a single contour. Consistent vole identities across video frames were determined

using the Hungarian algorithm. Since the assay began with one vole in each of the three

chambers, and since the subject vole remained freely mobile throughout, it was possible

to unambiguously assign each contour to a single vole identity (e.g., subject, right

chamber, or left chamber) or to that of a pair of voles (e.g., subject and right chamber

voles or subject and left chamber vole), resulting in continuous trajectories for each of

the three animals. Time in chamber was determined by the chamber location of the

subject vole in each frame, since the chamber location of the right chamber and left

chamber animals was fixed throughout the assay by physical tethering and any time the

contour of the subject animal merged with that of the tethered animal (right or left), the

subject was termed in contact with it. The algorithm also determined the boundaries of

the chambers and calculated chamber times. We used the chamber time data from

when both ends of the chamber contained tethered WTs (WT-WT condition) to ensure

that there was no bias in which side the choosers preferred.

Bond formation assays

Initial introduction of $Oxtr^{1-/-}$ and WT animals of either sex with an opposite sex WT animal

was filmed for 30 minutes. Observed behaviors included anogenital sniffing, investigation

(subject animal's nose contacts any non-genital region), duration of stationary huddling

or >50% side-by-side contact between animals, rearing, frequency of aggressive behavior

including lunges, receipt of aggression, and tussles, as well as mating behavior including

mounting, intromission, and ejaculation. These animals remained in their home cage for

18 hours following pairing, after which, they were separated by a clear, plastic barrier. The

barrier was left in place for 24 hours and then removed allowing the subject free access

to the partner animal thereafter. A 30-minute period directly following barrier removal was

also filmed to assess mating behavior, termed the timed mating assay.

Subjects remained cohoused with the opposite-sex, wildtype partner for durations as

indicated in the text. The partner preference test (PPT), as previously published[52], was

used to assess pair bond formation. Subject, partner, and an opposite sex stranger are

placed in a three-chamber apparatus with open top and 10 x 32-inch walls. The partner

and stranger animals are tethered on either end of the three-chamber arena and the

subject allowed free access over a period of 3 hours. Behaviors are recorded from a top

view camera capturing the entire apparatus. Videos were scored using BORIS[110] post-

test by validated scorers blind to condition. Observed behaviors included location (i.e.,

duration of time in partner, stranger, and neutral chambers), duration of stationary

huddling or >50% side-by-side contact between the partner and stranger animals, and

frequency of aggressive behavior (i.e., lunges). Of note, analysis of partner preference

between wild type siblings from the $Oxtr^{1-/-}$ background and wild type animals from the

breeding colony showed no difference in time spent huddling with partner vs stranger

animals.

Bond maintenance assays

Four days after the timed mating assay, female animals underwent a separation-

reunification assay. Females remained in their home cage while their male partners

were placed in a separate cage in another room for one hour. The males were then

reintroduced into the home cage and behavior was video recorded for 30 minutes. Male

subject animals underwent the same procedure five days following timed mating. Two

days following the separation reunification test, animals underwent a selective

aggression assay. Partners were again removed from the home cage and separated to

a separate room for an hour. After one hour, a novel sexually naïve animal of the

opposite sex was introduced into the home cage with the subject animal and behavior

observed for 20 minutes. If significant aggression occurred, defined as >3 tussles during

the assay, the animals were separated and the assay terminated. Overall, two WT

female assays and one Oxtr[1-/-] were terminated for aggression. The same behaviors

listed above for the introduction were observed and recorded in the reunification and

aggression assays.

RNA Sequencing

Sample collection

Adult 7-9 week old WT and Oxtr[1-/-] voles were either cohoused with same sex-siblings or

paired with an animal of the opposite sex for four days following the timed mating protocol

described above. These animals were euthanized by CO_2 inhalation and decapitated. The

brain was quickly dissected out and sectioned into 500μm coronal slices using a brain

matrix mold (BrainTree Scientific) chilled on ice. Slices were floated in ice-cold phosphate

buffered saline (PBS), the NAc was identified using anatomical landmarks, and dissected

using a Zeiss microscope. A total of 45 WT females, 39 WT males, 21 Oxtr[1-/-] females,

and 18 Oxtr[1-/-] males were split between paired and unpaired conditions and tissue from

three animals per condition was pooled prior to RNA extraction. Tissue was flash-frozen

in liquid nitrogen and stored at -80°C until further processing. Total RNA was extracted

using TRIzol according to manufacturer's instructions and quantified using NanoDrop

(Thermo Scientific). Library preparation for next generation sequencing, comprising 35

samples and 4 technical replicates, was done using TruSeq Stranded mRNA Kit

according to manufacturer recommendations and sequenced on a NovaSeq 6000 to an

average depth of 2.9e07 reads per sample. Sequencing was performed across three

runs, but all data was processed together.

Read processing

Adapter and low-quality sequence trimming was performed with Trimmomatic v0.39

using parameters: 'TruSeq3-PE.fa:2:30:10:2:keepBothReads LEADING:3 TRAILING:3

SLIDINGWINDOW:4:20 MINLEN:25' (Bolger et al., 2014). Trimmed RNA-Seq reads

were aligned to a custom MicOch1.0 prairie vole genome, which incorporated a known

missing gene V1A, using STAR v2.7.3a in gene annotation mode (Dobin et al., 2013).

Alignment, RNA-Seq, and Insert Size quality control metrics were generated using

Picard v 2.10.10 (http://broadinstitute.github.io/picard). Homologous genes between

vole, mouse, and human references were identified using biomart v2.56.1. When

multiple mouse or human genes mapped to the same vole gene, the homologous gene

with the highest percent identity to vole was selected.

Differential expression analysis

To remove non- and lowly-expressed genes, only genes with more than 2 counts per

million (cpm) in at least 3 samples were retained. Principal component analysis of the

cpm revealed that technical replicates all cluster tightly but showed a slight batch effect

due to sequencing run (Fig. S5A). To remove this effect, we used RUVSeq v1.34.0 to

estimate one factor of unwanted variation, W_1, which we later used to normalize the

data prior to differential expression[111]. The factor analysis was performed on the

deviance residuals from an initial generalized linear model regression of the upper

quartile normalized counts on our covariates of interest (pair bonding status, genotype,

sex). Expressed genes were tested for differential expression (DE), after combining

technical replicates, with DESeq2 v1.40.2 using a design matrix that included the

covariates and W_1[112]. The adaptive shrinkage estimator ashr v2.2-63 was used in

order to be more robust to genes with low cpm values[113]. For DE analyses in general,

while most changes were driven by female samples, the effect was maintained in the

males at a smaller magnitude; thus, analyses were primarily performed for both sexes

jointly to maximize accuracy and power.

Bonding effect

To compare pre- to post-pairing individuals, we performed the DE analysis in two ways.

First, we compared pairing status regardless of genotype or sex to increase our

statistical power and to identify a reproducible gene set of thirteen genes (Fig. 5B, 5C,

S5B). While the relationship between the genes is unknown, there is support that the

proteins are functionally related. Using the STRING protein database

(STRINGdb_2.12.1), the human-homologs of the DE genes share more connections

than we would expect by chance (Fig. S5C, p-value= 3.08E-0.7). Next, we compared

pairing status using all three covariates, to identify subgroup-specific genes of interest

and effect sizes (Fig. 5D, S5D). In both analyses, genes were considered significantly

DE with a BH adjusted p-value < 0.05.

Genotype effect

To examine the effect of genotype, we first performed DE analyses between WT and

Oxtr$^{-/-}$ individuals, not including sex as a covariate to identify a reproducible and

stringent gene set. We considered genes DE with an adjusted p-value < 0.05 and

absolute log2 fold-change > 0.25 (Fig. 6B, S6A, S6B). To examine the functional

relationship of the DE genes, we performed a Gene Ontology enrichment analysis using

goseq v1.52.0 which controls for gene length bias[114]. We considered all expressed vole

genes with a mouse homolog as our background and all DE genes with a mouse

homolog as our foreground. GO categories with an adjusted p-value < 0.05 were

considered significantly enriched. To help visualize the results, enriched GO categories

were clustered according to their semantic similarity calculated with the R package

GOSemSim v2.26.1 using method "Wang"[115]. Large, general, GO categories with over

1000 genes were excluded in visualizations (Fig. 6C). WikiPathways[116] over-

representation analysis was performed with R package clusterProfiler v4.8.3[117] (Fig.

S6D).

For disorder and disease set overrepresentation analysis, Chi-squared tests were used

to assess whether Oxtr$^{1-/-}$ disrupted genes occur in disease-associated gene lists at a

frequency higher than we would expect by chance. Genes lists for autism spectrum

disorder[63], schizophrenia[64], neurodevelopmental disorders[66], and epilepsy[65] were

sourced and all alternate or outdated gene symbols were updated manually. For this

test, we were interested in sex- and direction-specific enrichments, so we performed a

DE analysis separately for males and females. Across all DE analyses, we defined DE

genes more stringently (adjusted p-value < 0.01). For enrichment tests, our universe

was all expressed vole genes with a human homolog. We split this analysis by fold-

change direction to highlight the distinct enrichment in upregulated genes. P-values

underwent BH correction across all 24 comparisons.

To compare to mouse neuronal populations, we leveraged single-cell data on mouse

hypothalamus development[72]. The released Seurat object

GSE132730_subset_cca_to_velo.rds.gz was accessed at GSE132730. Using the

authors' defined clusters (accessed at "prim_walktrap"), we assessed how similar each

clusters' gene expression is to the genes disrupted by the $Oxtr^{1-/-}$ mutation. This was

calculated as the Jaccard similarity coefficient using R package clustifyR v1.12.0[118].

Weighted Gene Coexpression Network Analysis

<u>Data</u>

Starting with the RNAseq dataset of n=30 samples (one WT sample and 4 replicate

samples were removed) and n=13357 genes, we removed genes with low expression

(n=1738 genes with expression level in lowest quantile for all 30 samples) and/or low

variability (n=8230 genes with coefficient of variation < 0.05), resulting in a trimmed

dataset of n=3389 genes.

<u>Identifying co-expression modules</u>

We used the R package WGCNA (v1.70-3) to conduct WGCNA analyses[61]. Briefly, we

calculated Pearson correlations for all gene pairs. Next, we constructed an unsigned

weighted correlation network to identify co-expression modules comprised of highly

correlated genes (positive or negative correlation) with high topological overlap[119].

Modules were defined as branches of a hierarchical cluster tree using the top-down

dynamic tree cut method[61]. For each module, the expression pattern was summarized

by the module eigengene (ME), which is defined as the right singular vector of the

expression patterns. Pairs of modules with high module eigengene correlations (r >

0.75) were merged to maintain a level of decorrelation among MEs.

In more detail, a weighted unsigned network was computed based on a fit to scale-free

topology, and the lowest thresholding power that resulted in a scale-free R^2 fit of 0.9

was selected, and the pairwise topological overlap (TO) between genes was

calculated[119] . We constructed a TO dendrogram using hierarchical clustering (R::hclust

function, method= "average"), and defined modules using the WGCNA::cutreeHybrid

function with minimum module size set to 30 genes. We then merged modules with high

module eigengene correlations as describe above.

We calculated 3 sets of modules from different sample subsets:

1. Modules[WT]: n=15 modules constructed from n=18 WT samples with

891   thresholding power β = 5.

892   2. Modules[Mut]: n= 43 modules constructed from n=12 Oxtr[1]-mutant samples

893   with thresholding power β = 6.

894   3. Modules[All]: n=7 modules constructed from all n=30 samples with

895   thresholding power β = 3

<u>Relating modules to sample traits</u>

In order to correlate modules with experimental conditions, we calculated the Pearson

correlations between module eigengenes and sample metrics, including species

(Mut=0, WT=1), condition (Pre=1, Post=0), sex (F=1, M=0), and quality control (QC)

metrics (RIN, yield, sequencing run). We identified three modules that significantly

correlated with a sample metric of interest but not with any of the QC metrics.

Significant correlation was defined to be p_{adj} <= 0.05, where p_{adj} = BH correction by

number of modules. These modules (M) were:

1. Mpink^{WT}: positive correlation with pre-bonding status (Pearson R = 0.7, p_{adj} =
   0.018)

2. Mbrown^{All}: positive correlation with WT status (Pearson R = 0.55 , p_{adj} = 0.011)

3. Mturquoise^{All}: positive correlation with WT status (Pearson R = 0.8, p_{adj} = 6.6e-7)

We calculated the module eigengenes (ME) for Mpink^{WT}, Mbrown^{All}, and Mturquoise^{All}

for all n=30 samples using the WGCNA::moduleEigengenes function. We then

conducted Wilcoxon rank sum tests (ggpubr::stat_compare_means function) to assess

whether MEs were significantly different between the following groups:

1. MEpink^{WT}: pre- versus post-bonded animals, separated by sex and species

2. Mbrown^{All}: WT versus mutant animals, separated by sex and condition

3. Mturquoise^{All}: WT versus mutant animals, separated by sex and condition

<u>Comparing gene significance with module membership:</u>

We assessed whether genes whose expression highly correlate with module

eigengenes of modules of interest (Mpink^{WT}, Mbrown^{All}, and Mturquoise^{All}) are

themselves highly correlated with traits of interest (bonding status or mutation status).

Specifically, for the genes in Mpink^{WT}, Mbrown^{All}, and Mturquoise^{All}, we calculated 1)

gene significance (GS), defined as the absolute Pearson's correlation between

expression of a given gene and a trait of interest; and 2) module membership (MM),

which is defined as the absolute Pearson's correlation between the expression profile of

a gene with the ME of a module to quantify the relationship between a gene and a given

module[61]. We then calculated the Pearson correlations between:

1. GS^{condition} and MMpink^{WT} for Mpink^{WT} genes (R = 0.69, p = 5.5e-18)

2. GS^{species} and MMbrown^{All} for Mbrown^{All} genes (R = 0.63, p = 2.2e-32)

3. GS^{species} and MMturquoise^{All} for MMturquoise^{All} genes (R= 0.82, p<1e-200)

Hub genes were defined as those genes in the top 25% for both the absolute value of

pink module membership and the absolute value of correlation with bonding status.

<u>Assessing whether modules are enriched for DE genes</u>

We were interested in assessing whether Mpink^{WT} genes are significantly enriched for

any the DE pre vs post pairing gene set described above. We assessed for enrichment

of DE gene sets using a hypergeometric test.

<u>WT module preservation analysis (Zsummary)</u>

We assessed how well modules^{WT} are preserved in mutant samples. We calculated a

Z$_{summary}$ statistic for each module^{WT} [120] using the WGCNA::modulePreservation function

with 200 permutations. The $Z_{summary}$ measure combines module density and

intramodular connectivity metrics to a composite statistic where Z>2 suggests moderate

preservation and Z>10 suggests high preservation. We found of all the n=15

modules^{WT}, Mpink^{WT} had the lowest $Z_{summary}$ ($Z_{summary}$=2) in mutant samples.

Histology

Immunohistochemistry (IHC)

Sexually naïve voles group housed by sex after weaning (P0, P21, adults:10 - 14 weeks

old), were perfused with 25mL ice cold phosphate buffer (PBS) and fixed in 25mL of 4%

paraformaldehyde in PBS (PFA). Whole brain was dissected and post fixed overnight at

4°C in 4% PFA in the dark. The tissue was cryoprotected for 24 hours in 30% sucrose in

PBS and then embedded in OCT over dry ice to prevent cracking. 60μm serial sections

from the entire PVN were collected in PBS and processed as free-floating sections.

Sections were blocked using 10% donkey serum in 0.1% Triton X 100 in PBS and then

stained with primary antibodies (Oxytocin Polyclonal Antibody, Invitrogen, Rabbit

Polyclonal, Cat #PA5-26701, 1:1000 and Anti-Neurophysin 2/NP-AVP Antibody, clone

PS 41, Millipore Sigma, Mouse Monoclonal, Cat#MABN845, 1:10,000) in 1% donkey

serum with 0.1% Triton X 100 (staining buffer) in PBS at 4°C with shaking. The sections

were rinsed and soaked in secondary antibodies (Cy™3 AffiniPure Donkey Anti-Rabbit

IgG, Jackson ImmunoResearch Laboratory, Cat #711-165-152, 1:500 and Alexa Flour

488 Donkey Anti Mouse IgG, Thermo Fisher Scientific, Cat #A21202, 1:500) in staining

buffer for 2 hours at room temperature. Sections were DAPI stained and transferred to

glass slides in sequence and mounted in Aquamount. Slides were imaged using a

confocal microscope (Zeiss LSM 700) and the zstacks were quantified (cell count by

hand and area hand selected and measured) in ImageJ. Comparisons of cell density

(count/area) were made in age and sex matched samples.

In situ hybridization (ISH)

ISH was performed using RNA probes prepared as described previously[121], to detect

expression of *Calcr, Dlk1*, and *Dgkb* mRNA in the adult vole brain. We prepared RNA

sense and anti-sense probes corresponding to 706bp (CalcR), 773bp (Dgkb), and

796bp (Dlk1) for genes of interest identified in the WT vs Oxtr DEX list from our RNAseq

data. For ISH on adult animals, 7-9 week old WT and Oxtr$^{1-/-}$ voles of both sexes were

either group housed with same sex conspecifics or paired for 4 days with a WT animal

of the opposite sex following the timed mating protocol described above. The brains

were processed as described for IHC, then embedded in OCT and sectioned at 50μm,

mounted immediately after sectioning onto SuperFrost Plus glass microscope slides

and stored at -80°C until ISH staining. Slides were fixed in 4% PFA for 20 minutes,

rinsed in PBS, treated with proteinase K (10μg/mL, Roche) for 20 minutes, rinsed in

PBS, and fixed again in 4% PFA for 5 minutes at room temperature. Slides were

acylated for 10 mins then rinsed in 1% Triton100X in PBS followed by PBS. The slides

were equilibrated in warm hybridization solution for 1-2 hours at 65°C and subsequently

incubated with a temporary paraffin cover for 14 - 18 hours at 65°C in fresh

hybridization buffer containing RNA probe. After incubation, the slides were dipped in

5X SSC warmed to 72°C to remove the parafilm coverslips and then washed in 0.2X

SSC warmed to 72°C. The slides were blocked with 10% heat inactivated sheep serum

(HISS) solution then stained for 12 - 18 hours at 4°C in buffer containing 10% HISS and

alkaline phosphatase-conjugated sheep anti- digoxigenin antibody (1:2000, Roche).

After extensive washing, the slides were incubated for ~72 hours at 37°C in staining

solution containing nitro blue tetrazolium and 5- bromo-4-chloro-3-indolyl-phosphate

(Roche). The slides were finally washed, fixed in 4% PFA, and coverslipped. Slides

were imaged using a Keyence fluorescence microscope and the z-stacks were

quantified (cell count and area) in ImageJ. For signal quantitation, positive puncta were

counted in every third coronal section encompassing the entire anterior to posterior

gene expression domain. Cell counts were normalized by total area for a gene

expression domain in each section. Positive signal was quantified by calculating total

area of signal multiplied by the average intensity of signal (total area x mean pixel

intensity) in gene expression domains of a section; every third section was quantified

and added. Comparisons of cell counts were made in age and sex matched samples.

Quantification and Behavioral Analysis

For detailed statistical methods please see Table S1. The number of animals used was

based on previous studies in the field by our group and others, combined with a power

analysis. Assumptions of independence and normality were considered with multiple

tests. Preference index (PI) was calculated as (partner huddle duration – stranger huddle

duration)/(partner huddle duration + stranger huddle duration). Sliding partner preference

was calculated by calculating PI for a single animal in 20-minute intervals where each

interval overlapped the next by 15 minutes (e.g. 0-20mins, 5-25mins…). To look at

convergence in sliding PI the area under the curve (Area above 0 – Area below 0, AUC)

for each hour of the assay was determined for each animal and then pooled for each

choosing animal by sex and then condition. Preference was determined as significantly

convergent if the AUC from the hour of assay for all animals within the group significantly

differed from 0 using a One sample t Test. The level of statistical significance for each test

was set at $p =< 0.05$. Outliers were detected using a combination of z-score (+/- 3) and

plots of original and log-transformed data. Analyses were completed in RStudio (version

2024.04.2+764) and GraphPad Prism (version 10.3.1)

1016

FIGURE LEGENDS

Figure1: Oxtr reduces the amount of time naive animals take to show partner

preference

A. Schematic of the partner preference tests (PPTs) and the day on which they were

conducted to study pair bonding behaviors. The prosocial, agonistic and mating

behaviors scored are shown below. During the bond formation phase, the ratio of

partner-directed vs stranger-directed affiliative behaviors increases in wild types

and following the bond formation in the bond maintenance phase, wild types show

decreased prosocial behaviors and increased agonistic behaviors towards

strangers.

B. A preference for partner is induced in the WTs but not Oxtr[1-/-] animals following

short cohabitation.

C. WT females spend more time huddling with their partners.

D. Oxtr[1-/-] females spend more time huddling with strangers.

E. WT females show a significant increase in partner huddling time towards the end

of a 3-hour assay.

F. $Oxtr^{1-/-}$ males show a significant increase in stranger huddling time in the middle of

a 3-hour assay.

G. No difference in partner huddling time between WT and $Oxtr^{1-/-}$ animals following

long cohabitation.

H. No difference in stranger huddling time between WT and $Oxtr^{1-/-}$ animals following

long cohabitation.

See Supplementary Figure 1 and Supplementary Table 1.

Mean +/- SD, n =>8, *p =< 0.05, FWT = female WT, $FOxtr^{1-/-}$ = female $Oxtr^{1-/-}$, MWT =

male WT, $MOxtr^{1-/-}$ = male $Oxtr^{1-/-}$.

Figure2: Oxtr suppresses promiscuous behaviors in a state dependent manner in

females

A. Timeline of the behavioral battery carried out on females. Bottom row shows the

behaviors scored.

B. $Oxtr^{1-/-}$ female long huddle duration is shorter with partners and longer with

strangers after a short cohabitation.

C. $Oxtr^{1-/-}$ female long huddle frequency is greater with strangers, but not partners

after a short cohabitation.

D. No difference in median long huddle durations between WT and $Oxtr^{1-/-}$ females

post long cohabitation.

E. No difference in long huddle frequencies between WT and $Oxtr^{1-/-}$ females post

long cohabitation.

F. Increasing cohabitation time increases the median long huddle duration an $Oxtr^{1-/-}$ female executes with the partner.

G. There is no significant change in the median long huddle duration conducted by females with strangers when cohabitation time is increased.

H. Naive $Oxtr^{1-/-}$ and WT females are equally prosocial towards a naive WT male.

I. There is no difference in the counts of mating behaviors between $Oxtr^{1-/-}$ and WT females.

J. $Oxtr^{1-/-}$ and WT females are equally prosocial towards their partners after a brief separation.

K. $Oxtr^{1-/-}$ females are significantly more prosocial towards a naive WT stranger male compared to WT females.

See Supplementary Figure 2 and Supplementary Table 1.

Mean ± SD, n =>8, *p < 0.05, FWT = female WT, $FOxtr^{1-/-}$ = female $Oxtr^{1-/-}$, MWT = male WT, $MOxtr^{1-/-}$ = male $Oxtr^{1-/-}$.

Figure3: Oxtr function influences different prosocial behaviors across pair bonding in males

A. Timeline of the behavioral battery carried out on males. Bottom row shows the behaviors scored.

B. $Oxtr^{1-/-}$ male long huddle duration is shorter with partners and strangers after a short cohabitation.

C. $Oxtr^{1-/-}$ male long huddle frequency is greater with strangers, but not partners after a short cohabitation.

D. The median long huddle executed by Oxtr$^{1-/-}$ males with the stranger after a long

cohabitation, is shorter than their WT siblings.

E. No difference in the long huddle frequencies between WT and Oxtr$^{1-/-}$ males post

long cohabitation.

F. There is no change in the frequency of long huddles with partner between Oxtr$^{1-/-}$

and WT males.

G. Increasing cohabitation time decreases the frequency of long huddles with

strangers in Oxtr$^{1-/-}$ males.

H. Naive Oxtr$^{1-/-}$ and WT males are equally prosocial towards naive WT females.

I. Oxtr$^{1-/-}$ males execute higher counts of mating behaviors compared to WT.

J. Prosocial behaviors following a brief separation with their partner are not different

between Oxtr$^{1-/-}$ and WT males.

K. Oxtr$^{1-/-}$ males are significantly more prosocial towards a naive WT stranger female

compared to WT males.

See Supplementary Figure 3 and Supplementary Table 1.

Mean +/- SD, n =>8, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT =

male WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$

Figure4: Oxtr sex-specifically influences social interactions between potential

mates

A. Schematic of naive social choice with a naive WT animal choosing between an

opposite sex WT and opposite sex Oxtr$^{1-/-}$ animal (red triangles).

B. Male WT choosers prefer WT females.

C. Preference for WT females over Oxtr$^{1-/-}$ females by WT males manifests during

the middle of a 6-hour assay.

D. Schematic of comparison of time in contact with the MOAT from WT-Oxtr$^{1-/-}$

condition to contact with the MOAT from WT-WT condition.

E. WT female chooser spends more time in contact with a MOAT from a WT-Oxtr$^{1-/-}$

condition than the WT-WT. The color of the dots represents the genotype of the

MOAT.

F. WT female choosers transition between contacts more frequently in the WT-WT

condition compared to the WT-Oxtr$^{1-/-}$ condition.

G. WT female chooser spend significantly more time in contact with the MOAT from

the WT-Oxtr$^{1-/-}$ condition in the last 40 mins of the assay.

H. WT female choosers prefer their MOAT over longer periods of time in the WT-

Oxtr$^{1-/-}$ condition. WT male choosers prefer their MOAT earlier in the WT-Oxtr$^{1-/-}$

condition, but then show preference for the same amount of time as the WT-WT.

See Supplementary Figure 4 and Supplementary Table 1.

Mean +/- SD, n = 8, *p =< 0.05, FWT = female WT, FOxtr$^{1-/-}$ = female Oxtr$^{1-/-}$, MWT = male

WT, MOxtr$^{1-/-}$ = male Oxtr$^{1-/-}$

Figure 5: Post pair bonding gene expression is driven by female specific changes

dependent on Oxtr signaling

A. Schematic of experimental groups used for sequencing highlighting pre- vs post-

pairing comparisons between groups. Animals with symbols represent those used

for tissue collection.

B. Scatterplot illustrating the relationship between fold change and mean expression, emphasizing the 13 differentially expressed (DE) genes in post-pairing individuals compared to pre-pairing. Gray dots indicate non-significant genes ($p_{adj} > 0.05$); Blue dots indicate downregulated genes ($p_{adj} < 0.05$, $\log_2FC < 0$) and upregulated genes in red ($p_{adj} < 0.05$, $\log_2FC > 0$). The right panel shows box plots of normalized gene expression for the top six DE genes by fold-change between pre- and post-pairing conditions.

C. Bar graph emphasizing how each combination of genotype and sex (denoted at bottom as 'subgroup') influence the 13 overall pairing-DE genes identified. The top graph depicts the absolute fold change and standard error of the differential expression analysis including all pre- and post-bonding individuals in which the 13 genes of the x axis were identified. The bottom graph depicts the absolute fold change and standard error in analyses when only individuals of that subgroup were considered. (* denotes $p_{adj} < 0.05$ for the pre- vs post-pairing analysis).

D. Box plots of select genes that show sex- and genotype-specific changes in expression with pairing condition. *Adora2A* and *Fkbp5* are DE when considering WT females, *Spred3* is DE when considering WT and Oxtr[1-/-] females, and *Sema3b* is DE when considering Oxtr[1-/-] males. Note: Sema3b is the mouse ortholog gene symbol for vole gene "ENSMOCG00000019704" which does not have an assigned gene symbol.

E. Heatmap showing correlations between module eigengenes (MEs) for modules[WT] and pairing condition (Pre=1, Post=0). Heatmap color and value reflect Pearson

correlation coefficient; adjusted p values shown in parentheses ($p_{adj} = p *$ number

of modules).

F. Preservation of modules^{WT} in OXTR-mutant samples. The $Z_{summary}$ measure

combines module density and intramodular connectivity metrics to a composite

statistic where Z>2 suggests moderate preservation and Z>10 suggests high

preservation[119]. Of all modules^{WT}(n=15), Mpink^{WT} had the lowest $Z_{summary}$

($Z_{summary}$=2) in mutant samples.

G. Box plots showing comparison of MEpink^{WT} for pre- vs post-pairing condition

separated by sex and genotype. (Wilcoxon rank sum test, Bonferroni adjusted for

4 comparisons).

For all box plots center = median, box boundaries = 1^{st} and 3^{rd} quartile, whiskers =

1.5*IQR from boundaries. (n = 16 pre-pairing and 15 post-pairing)

See Supplementary Figure 5.

Figure 6: Oxtr regulated genes are common to social behavior relevant processes

across species and are similarly regulated across brain regions

A. Schematic of experimental groups for sequencing, highlighting genotype

comparisons between groups. Animals with symbols represent those used for

tissue collection.

B. Scatterplot illustrating the relationship between fold change and mean expression,

emphasizing differentially expressed (DE) genes in Oxtr$^{1-/-}$ individuals compared

to WT. Significantly downregulated genes are shown in blue ($\log_2$FC<-0.25,

p_{adj}<0.05) and upregulated genes in red ($\log_2$FC>0.25, p_{adj} < 0.05). Box plots of

normalized gene expression for the top six DE genes by fold change between

genotypes.

C. Gene Ontology enrichment analysis for all genotype-DE genes (n=1014) highlights

processes related to extracellular channel binding activity, axonal processes,

behavior related gene expression, and cation transport. The leaves represent

significant GO categories (BH adjusted p value < 0.05) which have been

hierarchically clustered by their semantic similarity. Branches are colored by

cluster and high-frequency words are displayed to the right. Leaf size and color

correspond to the number and median fold change of DE genes in each category.

D. Enrichment analysis shows that genes which are DE in $Oxtr^{1-/-}$ samples are over-

represented in some neuropsychiatric disease-associated gene sets. The first DE

gene set evaluated ('Joint') represents genes which are DE in $Oxtr^{1-/-}$ samples

when considering both sexes in the differential expression analysis ($p_{adj} < 0.01$).

The second gene set are DE genes when only considering females, and the third

when only considering males. Gene sets are further separated by direction of effect

across the x-axis to highlight the enrichment in upregulated DE genes. Colors

correspond to the Benjamini-Hochberg corrected Chi-square p values; asterisks

denote $p_{adj} < 0.05$. Y-axis abbreviations are as follows: SCZ=schizophrenia,

NDD=neurodevelopmental disorder, EPI=epilepsy, and ASD=autism spectrum

disorder.

E. Box plots of *Calcr* and *dlk1* normalized gene counts by genotype. Log_2FC:-0.962

and p_{adj}: 1.75e-06 for *Calcr* and log_2FC: -0.276 and p_{adj}: 0.00334 for *dlk1* from

comparison considering all WT and $Oxtr^{1-/-}$ samples.

F. Brain regions selected for in situ analysis. (NAc=nucleus accumbens,

PVN=paraventricular nucleus of the hypothalamus, LS=lateral septum)

G. ISH images and quantification for *Calcr* expression in NAc, PVN, and LS from WT

and Oxtr[1-/-] animals.

H. ISH images and quantification for *Dlk-1* expression in NAc, PVN, and LS from WT

and Oxtr[1-/-] animals.

Box plots in G. and H. show comparison of the area-normalized cell number by

genotype. (*n*=3 per condition, Wilcoxon sign rank, *=$p<0.05$, ***=$p<0.001$)

See Supplementary Figure 6 and Supplementary Table 1.

Figure7: Loss of Oxtr results in the decrease in density of Oxt and Avp

expressing neurons in the PVN

A. UMAP visualization of mouse hypothalamus single-cell RNAseq dataset[66] with

proposed PVN clusters labeled. The authors' original numeric cluster labels and

embedding are shown. Each cluster is colored by the Jaccard similarity coefficient

between the mouse marker gene sets and the vole Oxtr[1-/-] DE genes (ONP set).

This is also represented in a heat plot to the right. Some of the highest similarity is

seen in Cluster 43, which represents Oxt+ cells, and Cluster 26, which represents

Avp+ cells in the PVN.

B. Representative images of IHC staining against OT and AVP neuropeptides from

FWT and FOxtr[1-/-] PVN showing the loss of structural integrity and OT and AVP

positive cells from the knock-out. (White dotted line denotes the PVN area, scale

bar represents 100µm)

C. Mean of positive cells per unit area across all sections is lower for $Oxtr^{1-/-}$s vs WT for both OT (left) and AVP (right). Each dot represents 1 animal.

D. The number of sections with a high density of OT positive cells is greater in both female (left) and male (right) WT animals.

E. Similarly, WT animals of both sexes show an increase in the number of sections containing AVP positive neurons in high densities.

F. The average OT cell density from matched PVN sections arranged from anterior to posterior shows that the decrease in cell density in $Oxtr^{1-/-}$ animals is biased towards the anterior. Each dot represents the average cell density, the lines represent the sixth order non-linear regression model for cell density by sex and by genotype (n=3) and the ribbons represent the standard deviation.

G. $Oxtr^{1-/-}$ animals show a significant decrease in AVP cell density in the anterior PVN in both sexes.

See Supplementary Figure 7 and Supplementary Table 1.

Mean +/- SD, n = 3, *p =< 0.05, FWT = female WT, $FOxtr^{1-/-}$ = female $Oxtr^{1-/-}$, MWT = male WT, $MOxtr^{1-/-}$ = male $Oxtr^{1-/-}$.

References

1. Wilson, R.S., Krueger, K.R., Arnold, S.E., Schneider, J.A., Kelly, J.F., Barnes, L.L., Tang, Y., and Bennett, D.A. (2007). Loneliness and Risk of Alzheimer Disease. Arch Gen Psychiatry *64*, 234. https://doi.org/10.1001/archpsyc.64.2.234.

2. Bowlby, J., and Bowlby, J. (1982). Attachment 2. ed. (Basic Books).

3. Farinelli, M., Cevolani, D., Gestieri, L., Romaniello, C., Maffei, M., Agati, R., Leo,

M.R., Huang, Z., Pedone, V., and Northoff, G. (2020). Brain and behaviour in post-

acute stroke: Reduction in seeking and posterior cingulate neuronal variability.

Journal of Clinical and Experimental Neuropsychology *42*, 584–601.

https://doi.org/10.1080/13803395.2020.1780417.

4. Feldman, R., Monakhov, M., Pratt, M., and Ebstein, R.P. (2016). Oxytocin Pathway

Genes: Evolutionary Ancient System Impacting on Human Affiliation, Sociality, and

Psychopathology. Biological Psychiatry *79*, 174–184.

https://doi.org/10.1016/j.biopsych.2015.08.008.

5. Ebstein, R.P., Israel, S., Chew, S.H., Zhong, S., and Knafo, A. (2010). Genetics of

human social behavior. Neuron *65*, 831–844.

https://doi.org/10.1016/j.neuron.2010.02.020.

6. Winslow, J.T., Shapiro, L., Carter, C.S., and Insel, T.R. (1993). Oxytocin and

complex social behavior: species comparisons. Psychopharmacol Bull *29*, 409–414.

7. Goodson, J.L. (2013). Deconstructing sociality, social evolution and relevant

nonapeptide functions. Psychoneuroendocrinology *38*, 465–478.

https://doi.org/10.1016/j.psyneuen.2012.12.005.

8. Harlow, H.F., and Harlow, M.K. (1965). THE EFFECT OF REARING CONDITIONS

ON BEHAVIOR. Int J Psychiatry *1*, 43–51.

9. Shultz, S., Opie, C., and Atkinson, Q.D. (2011). Stepwise evolution of stable sociality

in primates. Nature *479*, 219–222.

10. Insel, T.R., Winslow, J.T., Wang, Z., and Young, L.J. (1998). Oxytocin, vasopressin,

and the neuroendocrine basis of pair bond formation. Adv. Exp. Med. Biol *449*, 215–

224.

11. Klatt, J.D., and Goodson, J.L. (2013). Oxytocin-like receptors mediate pair bonding

in a socially monogamous songbird. Proceedings of the Royal Society B: Biological

Sciences *280*, 20122396. https://doi.org/10.1098/rspb.2012.2396.

12. Ainsworth, M.S. (1979). Infant–mother attachment. American Psychologist *34*, 932–

937. https://doi.org/10.1037/0003-066X.34.10.932.

13. Getz, L.L., Carter, C.S., and Gavish, L. (1981). The mating system of the prairie

vole, Microtus ochrogaster: Field and laboratory evidence for pair-bonding. Behav

Ecol Sociobiol *8*, 189–194. https://doi.org/10.1007/BF00299829.

14. Donaldson, Z.R. (2010). We're the Same... but Different: Addressing Academic

Divides in the Study of Brain and Behavior. Front Behav Neurosci *4*.

https://doi.org/10.3389/fnbeh.2010.00041.

15. Insel, T.R., Winslow, J.T., Wang, Z.X., Young, L., and Hulihan, T.J. (1995). Oxytocin

and the molecular basis of monogamy. Adv. Exp. Med. Biol *395*, 227–234.

16. Lukas, D., and Clutton-Brock, T.H. (2013). The evolution of social monogamy in

mammals. Science *341*, 526–530.

17. Donaldson, Z.R., and Young, L.J. (2008). Oxytocin, vasopressin, and the

neurogenetics of sociality. Science *322*, 900–904.

https://doi.org/10.1126/science.1158668.

18. Bielsky, I.F., and Young, L.J. (2004). Oxytocin, vasopressin, and social recognition in

mammals. Peptides *25*, 1565–1574. https://doi.org/10.1016/j.peptides.2004.05.019.

19. Nowicki, J.P., Pratchett, M.S., Walker, S.P.W., Coker, D.J., and O'Connell, L.A.

(2020). Gene expression correlates of social evolution in coral reef butterflyfishes.

Proceedings of the Royal Society B: Biological Sciences *287*, 20200239.

https://doi.org/10.1098/rspb.2020.0239.

20. O'Connor, C.M., Marsh-Rollo, S.E., Aubin-Horth, N., and Balshine, S. (2016).

Species-specific patterns of nonapeptide brain gene expression relative to pair-

bonding behavior in grouping and non-grouping cichlids. Hormones and Behavior

80, 30–38. https://doi.org/10.1016/j.yhbeh.2015.10.015.

21. Stanley, D.A., and Adolphs, R. (2013). Toward a neural basis for social behavior.

Neuron *80*, 816–826.

22. Insel, T.R. (1992). Oxytocin--a neuropeptide for affiliation: evidence from behavioral,

receptor autoradiographic, and comparative studies. Psychoneuroendocrinology *17*,

3–35.

23. Lim, M.M., and Young, L.J. (2006). Neuropeptidergic regulation of affiliative behavior

and social bonding in animals. Horm Behav *50*, 506–517.

https://doi.org/10.1016/j.yhbeh.2006.06.028.

24. Carter, C.S., and Getz, L.L. (1993). Monogamy and the prairie vole. Sci. Am *268*,

100–106.

25. Beery, A.K., and Shambaugh, K.L. (2021). Comparative assessment of

familiarity/novelty preferences in rodents. Frontiers in Behavioral Neuroscience *15*,

648830.

26. Beery, A.K., Christensen, J.D., Lee, N.S., and Blandino, K.L. (2018). Specificity in

Sociality: Mice and Prairie Voles Exhibit Different Patterns of Peer Affiliation.

Frontiers in Behavioral Neuroscience *12*, 50.

https://doi.org/10.3389/fnbeh.2018.00050.

27. Schweinfurth, M.K., Neuenschwander, J., Engqvist, L., Schneeberger, K., Rentsch,

A.K., Gygax, M., and Taborsky, M. (2017). Do female Norway rats form social

bonds? Behavioral Ecology and Sociobiology *71*, 1–9.

28. Cho, M.M., DeVries, A.C., Williams, J.R., and Carter, C.S. (1999). The effects of

oxytocin and vasopressin on partner preferences in male and female prairie voles

(Microtus ochrogaster). Behav. Neurosci *113*, 1071–1079.

29. Insel, T.R., and Hulihan, T.J. (1995). A gender-specific mechanism for pair bonding:

oxytocin and partner preference formation in monogamous voles. Behav. Neurosci

109, 782–789.

30. Williams, J.R., Catania, K.C., and Carter, C.S. (1992). Development of partner

preferences in female prairie voles (Microtus ochrogaster): the role of social and

sexual experience. Hormones and behavior *26*, 339–349.

31. Wang, Z., Hulihan, T.J., and Insel, T.R. (1997). Sexual and social experience is

associated with different patterns of behavior and neural activation in male prairie

voles. Brain Res *767*, 321–332.

32. Xu, X., Coats, J.K., Yang, C.F., Wang, A., Ahmed, O.M., Alvarado, M., Izumi, T., and

Shah, N.M. (2012). Modular genetic control of sexually dimorphic behaviors. Cell

148, 596–607. https://doi.org/10.1016/j.cell.2011.12.018.

33. Villella, A., and Hall, J.C. (2008). Chapter 3 Neurogenetics of Courtship and Mating

in Drosophila. In Advances in Genetics (Elsevier), pp. 67–184.

https://doi.org/10.1016/S0065-2660(08)00603-2.

34. Konopka, R.J., and Benzer, S. (1971). Clock Mutants of *Drosophila melanogaster*.

Proc Natl Acad Sci USA *68*, 2112–2116. https://doi.org/10.1073/pnas.68.9.2112.

35. Takahashi, J.S., Shimomura, K., and Kumar, V. (2008). Searching for genes

underlying behavior: lessons from circadian rhythms. Science *322*, 909–912.

https://doi.org/10.1126/science.1158822.

36. Baker, B.S., Taylor, B.J., and Hall, J.C. (2001). Are complex behaviors specified by

dedicated regulatory genes? Reasoning from Drosophila. Cell *105*, 13–24.

37. Manoli, D.S., and Baker, B.S. (2004). Median bundle neurons coordinate behaviours

during Drosophila male courtship. Nature *430*, 564–569.

https://doi.org/10.1038/nature02713.

38. Martin, L.B., Glasper, E.R., Nelson, R.J., and DeVries, A.C. (2006). Prolonged

separation delays wound healing in monogamous California mice, Peromyscus

californicus, but not in polygynous white-footed mice, P. leucopus. Physiology &

Behavior *87*, 837–841. https://doi.org/10.1016/j.physbeh.2006.01.035.

39. Ferland, C.L., and Schrader, L.A. (2011). Cage mate separation in pair-housed male

rats evokes an acute stress corticosterone response. Neuroscience Letters *489*,

154–158. https://doi.org/10.1016/j.neulet.2010.12.006.

40. Okuda, T., Osako, Y., Hidaka, C., Nishihara, M., Young, L.J., Mitsui, S., and Yuri, K.

(2022). Separation from a bonded partner alters neural response to inflammatory

pain in monogamous rodents. Behavioural Brain Research *418*, 113650.

https://doi.org/10.1016/j.bbr.2021.113650.

41. Carter, C.S., Williams, J.R., Witt, D.M., and Insel, T.R. (1992). Oxytocin and social

bonding. Ann. N. Y. Acad. Sci *652*, 204–211.

42. Winslow, J.T., Hastings, N., Carter, C.S., Harbaugh, C.R., and Insel, T.R. (1993). A

role for central vasopressin in pair bonding in monogamous prairie voles. Nature

365, 545–548. https://doi.org/10.1038/365545a0.

43. Carter, C.S., Grippo, A.J., Pournajafi-Nazarloo, H., Ruscio, M.G., and Porges, S.W.

(2008). Oxytocin, vasopressin and sociality. Prog. Brain Res *170*, 331–336.

https://doi.org/10.1016/S0079-6123(08)00427-5.

44. Ophir, A.G., Gessel, A., Zheng, D.-J., and Phelps, S.M. (2012). Oxytocin receptor

density is associated with male mating tactics and social monogamy. Hormones and

behavior *61*, 445–453.

45. Shapiro, L.E., and Insel, T.R. (1992). Oxytocin receptor distribution reflects social

organization in monogamous and polygamous voles. Ann. N. Y. Acad. Sci *652*, 448–

451.

46. Heinrichs, M., von Dawans, B., and Domes, G. (2009). Oxytocin, vasopressin, and

human social behavior. Front Neuroendocrinol *30*, 548–557.

https://doi.org/10.1016/j.yfrne.2009.05.005.

47. Insel, T.R., Wang, Z.X., and Ferris, C.F. (1994). Patterns of brain vasopressin

receptor distribution associated with social organization in microtine rodents. J.

Neurosci *14*, 5381–5392.

48. King, L.B., Walum, H., Inoue, K., Eyrich, N.W., and Young, L.J. (2016). Variation in

the oxytocin receptor gene predicts brain region–specific expression and social

attachment. Biological psychiatry *80*, 160–169.

49. Walum, H., and Young, L.J. (2018). The neural mechanisms and circuitry of the pair

bond. Nat Rev Neurosci *19*, 643–654. https://doi.org/10.1038/s41583-018-0072-6.

50. Williams, J.R., Insel, T.R., Harbaugh, C.R., and Carter, C.S. (1994). Oxytocin

administered centrally facilitates formation of a partner preference in female prairie

voles (Microtus ochrogaster). J. Neuroendocrinol *6*, 247–250.

51. Bales, K.L., and Carter, C.S. (2003). Developmental exposure to oxytocin facilitates

partner preferences in male prairie voles (Microtus ochrogaster). Behavioral

neuroscience *117*, 854.

52. Berendzen, K.M., Sharma, R., Mandujano, M.A., Wei, Y., Rogers, F.D., Simmons,

T.C., Seelke, A.M., Bond, J.M., Larios, R., Goodwin, N.L., et al. (2023). Oxytocin

receptor is not required for social attachment in prairie voles. Neuron *111*, 787–796.

53. Young, L.J., Lim, M.M., Gingrich, B., and Insel, T.R. (2001). Cellular mechanisms of

social attachment. Hormones and behavior *40*, 133–138.

54. Tripp, J.A., Berrio, A., McGraw, L.A., Matz, M.V., Davis, J.K., Inoue, K., Thomas,

J.W., Young, L.J., and Phelps, S.M. (2021). Comparative neurotranscriptomics

reveal widespread species differences associated with bonding. BMC Genomics *22*,

399. https://doi.org/10.1186/s12864-021-07720-0.

55. Young, L.J., and Wang, Z. (2004). The neurobiology of pair bonding. Nat. Neurosci

7, 1048–1054. https://doi.org/10.1038/nn1327.

56. DeVries, A.C., and Carter, C.S. (1999). Sex differences in temporal parameters of

partner preference in prairie voles (Microtus ochrogaster). Can. J. Zool. *77*, 885–

889. https://doi.org/10.1139/z99-054.

57. Fraser, E.J., and Shah, N.M. (2014). Complex Chemosensory Control of Female

Reproductive Behaviors. PLOS ONE *9*, e90368.

https://doi.org/10.1371/journal.pone.0090368.

58. Aragona, B.J., Liu, Y., Curtis, J.T., Stephan, F.K., and Wang, Z. (2003). A Critical

Role for Nucleus Accumbens Dopamine in Partner-Preference Formation in Male

Prairie Voles. J. Neurosci. *23*, 3483–3490. https://doi.org/10.1523/JNEUROSCI.23-

08-03483.2003.

59. Dölen, G., Darvishzadeh, A., Huang, K.W., and Malenka, R.C. (2013). Social reward

requires coordinated activity of nucleus accumbens oxytocin and serotonin. Nature

501, 179–184. https://doi.org/10.1038/nature12518.

60. Ross, H.E., Freeman, S.M., Spiegel, L.L., Ren, X., Terwilliger, E.F., and Young, L.J.

(2009). Variation in Oxytocin Receptor Density in the Nucleus Accumbens Has

Differential Effects on Affiliative Behaviors in Monogamous and Polygamous Voles.

J. Neurosci. *29*, 1312–1318. https://doi.org/10.1523/JNEUROSCI.5039-08.2009.

61. Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted

correlation network analysis. BMC Bioinformatics *9*, 559.

https://doi.org/10.1186/1471-2105-9-559.

62. Maury, E.A., Sherman, M.A., Genovese, G., Gilgenast, T.G., Kamath, T., Burris, S.J.,

Rajarajan, P., Flaherty, E., Akbarian, S., Chess, A., et al. (2023). Schizophrenia-

associated somatic copy-number variants from 12,834 cases reveal recurrent

NRXN1 and ABCB11 disruptions. Cell Genom *3*, 100356.

https://doi.org/10.1016/j.xgen.2023.100356.

63. Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.-Y.,

Peng, M., Collins, R., Grove, J., Klei, L., et al. (2020). Large-Scale Exome

Sequencing Study Implicates Both Developmental and Functional Changes in the

Neurobiology of Autism. Cell *180*, 568-584.e23.

https://doi.org/10.1016/j.cell.2019.12.036.

64. Fromer, M., Pocklington, A.J., Kavanagh, D.H., Williams, H.J., Dwyer, S., Gormley,

P., Georgieva, L., Rees, E., Palta, P., Ruderfer, D.M., et al. (2014). De novo

mutations in schizophrenia implicate synaptic networks. Nature *506*, 179–184.

https://doi.org/10.1038/nature12929.

65. EuroEPINOMICS-RES Consortium, Epilepsy Phenome/Genome Project, and Epi4K

Consortium (2014). De novo mutations in synaptic transmission genes including

DNM1 cause epileptic encephalopathies. Am J Hum Genet *95*, 360–370.

https://doi.org/10.1016/j.ajhg.2014.08.013.

66. Kaplanis, J., Samocha, K.E., Wiel, L., Zhang, Z., Arvai, K.J., Eberhardt, R.Y.,

Gallone, G., Lelieveld, S.H., Martin, H.C., McRae, J.F., et al. (2020). Evidence for 28

genetic disorders discovered by combining healthcare and research data. Nature

586, 757–762. https://doi.org/10.1038/s41586-020-2832-5.

67. Inoue, K., Ford, C.L., Horie, K., and Young, L.J. (2022). Oxytocin receptors are

widely distributed in the prairie vole (Microtus ochrogaster) brain: Relation to social

behavior, genetic polymorphisms, and the dopamine system. J Comp Neurol *530*,

2881–2900. https://doi.org/10.1002/cne.25382.

68. Lefevre, A., Benusiglio, D., Tang, Y., Krabichler, Q., Charlet, A., and Grinevich, V.

(2021). Oxytocinergic Feedback Circuitries: An Anatomical Basis for

Neuromodulation of Social Behaviors. Front. Neural Circuits *15*, 688234.

https://doi.org/10.3389/fncir.2021.688234.

69. Harris, G.W. (1955). PITUITARY-HYPOTHALAMIC MECHANISMS. Arch NeurPsych

73, 124. https://doi.org/10.1001/archneurpsyc.1955.02330080002002.

70. Ye, Q., Nunez, J., and Zhang, X. (2022). Oxytocin Receptor-Expressing Neurons in

the Paraventricular Thalamus Regulate Feeding Motivation through Excitatory

Projections to the Nucleus Accumbens Core. J. Neurosci. *42*, 3949–3964.

https://doi.org/10.1523/JNEUROSCI.2042-21.2022.

71. Zhang, B., Qiu, L., Xiao, W., Ni, H., Chen, L., Wang, F., Mai, W., Wu, J., Bao, A., Hu,

H., et al. (2021). Reconstruction of the Hypothalamo-Neurohypophysial System and

Functional Dissection of Magnocellular Oxytocin Neurons in the Brain. Neuron *109*,

331-346.e7. https://doi.org/10.1016/j.neuron.2020.10.032.

72. Romanov, R.A., Tretiakov, E.O., Kastriti, M.E., Zupancic, M., Häring, M.,

Korchynska, S., Popadin, K., Benevento, M., Rebernik, P., Lallemend, F., et al.

(2020). Molecular design of hypothalamus development. Nature *582*, 246–252.

https://doi.org/10.1038/s41586-020-2266-0.

73. Mitre, M., Saadipour, K., Williams, K., Khatri, L., Froemke, R.C., and Chao, M.V.

(2022). Transactivation of TrkB Receptors by Oxytocin and Its G Protein-Coupled

Receptor. Front Mol Neurosci *15*, 891537.

https://doi.org/10.3389/fnmol.2022.891537.

74. Seoyoung Son, Steffy B. Manjila, Kyra T. Newmaster, Yuan-ting Wu, Daniel J.

Vanselow, Matt Ciarletta, Todd E. Anthony, Keith C. Cheng, and Yongsoo Kim

(2022). Whole-Brain Wiring Diagram of Oxytocin System in Adult Mice. J. Neurosci.

42, 5021. https://doi.org/10.1523/JNEUROSCI.0307-22.2022.

75. Vaidyanathan, R., and Hammock, E.A.D. (2020). Oxytocin receptor gene loss

influences expression of the oxytocin gene in C57BL/6J mice in a sex- and age-

dependent manner. Journal of Neuroendocrinology *32*, e12821.

https://doi.org/10.1111/jne.12821.

76. Newmaster, K.T., Nolan, Z.T., Chon, U., Vanselow, D.J., Weit, A.R., Tabbaa, M.,

Hidema, S., Nishimori, K., Hammock, E.A.D., and Kim, Y. (2020). Quantitative

cellular-resolution map of the oxytocin receptor in postnatally developing mouse

brains. Nature Communications *11*, 1885. https://doi.org/10.1038/s41467-020-

15659-1.

77. Kleiman, D.G. (1977). Monogamy in Mammals. The Quarterly Review of Biology *52*,

39–69. https://doi.org/10.1086/409721.

78. Feldman, R. (2017). The Neurobiology of Human Attachments. Trends in Cognitive

Sciences *21*, 80–99. https://doi.org/10.1016/j.tics.2016.11.007.

79. Yang, C.F., and Shah, N.M. (2014). Representing sex in the brain, one module at a

time. Neuron *82*, 261–278. https://doi.org/10.1016/j.neuron.2014.03.029.

80. Mong, J.A., and Pfaff, D.W. (2004). Hormonal symphony: steroid orchestration of

gene modules for sociosexual behaviors. Mol Psychiatry *9*, 550–556.

https://doi.org/10.1038/sj.mp.4001493.

81. Mong, J., Easton, A., Kow, L.M., and Pfaff, D. (2003). Neural, hormonal and genetic

mechanisms for the activation of brain and behavior. Eur J Pharmacol *480*, 229–

231. https://doi.org/10.1016/j.ejphar.2003.08.109.

82. Barron, A.B., and Robinson, G.E. (2008). The utility of behavioral models and

modules in molecular analyses of social behavior. Genes Brain and Behavior *7*,

257–265. https://doi.org/10.1111/j.1601-183X.2007.00344.x.

83. Kyriacou, C.P., and Hall, J.C. (1980). Circadian rhythm mutations in Drosophila

melanogaster affect short-term fluctuations in the male's courtship song. Proc Natl

Acad Sci U S A *77*, 6729–6733. https://doi.org/10.1073/pnas.77.11.6729.

84. Konopka, R.J., and Benzer, S. (1971). Clock Mutants of *Drosophila melanogaster*.

Proc Natl Acad Sci USA *68*, 2112–2116. https://doi.org/10.1073/pnas.68.9.2112.

85. Bendesky, A., Kwon, Y.-M., Lassance, J.-M., Lewarch, C.L., Yao, S., Peterson, B.K.,

He, M.X., Dulac, C., and Hoekstra, H.E. (2017). The genetic basis of parental care

evolution in monogamous mice. Nature *544*, 434–439.

https://doi.org/10.1038/nature22074.

86. Scheele, D., Striepens, N., Güntürkün, O., Deutschländer, S., Maier, W., Kendrick,

K.M., and Hurlemann, R. (2012). Oxytocin modulates social distance between males

and females. J Neurosci *32*, 16074–16079.

https://doi.org/10.1523/JNEUROSCI.2755-12.2012.

87. Ditzen, B., Schaer, M., Gabriel, B., Bodenmann, G., Ehlert, U., and Heinrichs, M.

(2009). Intranasal Oxytocin Increases Positive Communication and Reduces

Cortisol Levels During Couple Conflict. Biological Psychiatry *65*, 728–731.

https://doi.org/10.1016/j.biopsych.2008.10.011.

88. De Dreu, C.K.W., Greer, L.L., Handgraaf, M.J.J., Shalvi, S., Van Kleef, G.A., Baas,

M., Ten Velden, F.S., Van Dijk, E., and Feith, S.W.W. (2010). The Neuropeptide

Oxytocin Regulates Parochial Altruism in Intergroup Conflict Among Humans.

Science *328*, 1408–1411. https://doi.org/10.1126/science.1189047.

89. Osakada, T., Yan, R., Jiang, Y., Wei, D., Tabuchi, R., Dai, B., Wang, X., Zhao, G.,

Wang, C.X., Liu, J.-J., et al. (2024). A dedicated hypothalamic oxytocin circuit

controls aversive social learning. Nature *626*, 347–356.

https://doi.org/10.1038/s41586-023-06958-w.

90. Wang, Z., and Young, L.J. (1997). Ontogeny of oxytocin and vasopressin receptor

binding in the lateral septum in prairie and montane voles. Brain Res. Dev. Brain

Res *104*, 191–195.

91. Smith, M.L., Asada, N., and Malenka, R.C. (2021). Anterior cingulate inputs to

nucleus accumbens control the social transfer of pain and analgesia. Science *371*,

153–159. https://doi.org/10.1126/science.abe3040.

92. Gegenhuber, B., Wu, M.V., Bronstein, R., and Tollkuhn, J. (2020). Regulation of

neural gene expression by estrogen receptor alpha (Neuroscience)

https://doi.org/10.1101/2020.10.21.349290.

93. Cao, J., and Patisaul, H.B. (2013). Sex-specific expression of estrogen receptors α

and β and Kiss1 in the postnatal rat amygdala. J of Comparative Neurology *521*,

465–478. https://doi.org/10.1002/cne.23185.

94. Mitsushima, D., Yamada, K., Takase, K., Funabashi, T., and Kimura, F. (2006). Sex

differences in the basolateral amygdala: the extracellular levels of serotonin and

dopamine, and their responses to restraint stress in rats. Eur J Neurosci *24*, 3245–

3254. https://doi.org/10.1111/j.1460-9568.2006.05214.x.

95. McCann, K.E., Sinkiewicz, D.M., Rosenhauer, A.M., Beach, L.Q., and Huhman, K.L.

(2019). Transcriptomic Analysis Reveals Sex-Dependent Expression Patterns in the

Basolateral Amygdala of Dominant and Subordinate Animals After Acute Social

Conflict. Mol Neurobiol *56*, 3768–3779. https://doi.org/10.1007/s12035-018-1339-7.

96. Trainor, B.C. (2011). Stress responses and the mesolimbic dopamine system: social

contexts and sex differences. Horm Behav *60*, 457–469.

https://doi.org/10.1016/j.yhbeh.2011.08.013.

97. Wright, E.C., Luo, P.X., Zakharenkov, H.C., Serna Godoy, A., Lake, A.A., Prince,

Z.D., Sekar, S., Culkin, H.I., Ramirez, A.V., Dwyer, T., et al. (2023). Sexual

differentiation of neural mechanisms of stress sensitivity during puberty. Proc. Natl.

Acad. Sci. U.S.A. *120*, e2306475120. https://doi.org/10.1073/pnas.2306475120.

98. Dölen, G. (2015). Oxytocin: parallel processing in the social brain? J.

Neuroendocrinol. *27*, 516–535. https://doi.org/10.1111/jne.12284.

99. Choi, T.-Y., Jeon, H., Jeong, S., Kim, E.J., Kim, J., Jeong, Y.H., Kang, B., Choi, M.,

and Koo, J.W. (2023). Distinct prefrontal projection activity and transcriptional state

conversely orchestrate social competition and hierarchy. Neuron, S0896-

6273(23)00886-3. https://doi.org/10.1016/j.neuron.2023.11.012.

100.   Yaguchi, K., Hagihara, M., Konno, A., Hirai, H., Yukinaga, H., and Miyamichi, K.

(2023). Dynamic modulation of pulsatile activities of oxytocin neurons in lactating

wild-type mice. PLoS One *18*, e0285589.

https://doi.org/10.1371/journal.pone.0285589.

101.   Manoli, D.S., and State, M.W. (2021). Autism Spectrum Disorder Genetics and

the Search for Pathological Mechanisms. AJP *178*, 30–38.

https://doi.org/10.1176/appi.ajp.2020.20111608.

102.   Werling, D.M. (2016). The role of sex-differential biology in risk for autism

spectrum disorder. Biology of Sex Differences *7*, 58. https://doi.org/10.1186/s13293-

016-0112-8.

103.   Werling, D.M., and Geschwind, D.H. (2013). Sex differences in autism spectrum

disorders. Curr. Opin. Neurol. *26*, 146–153.

https://doi.org/10.1097/WCO.0b013e32835ee548.

104. Perissi, V., and Rosenfeld, M.G. (2005). Controlling nuclear receptors: the

circular logic of cofactor cycles. Nature Reviews: Molecular Cell Biology *6*, 542–554.

https://doi.org/10.1038/nrm1682.

105. Horie, K., Inoue, K., Nishimori, K., and Young, L.J. (2020). Investigation of Oxtr-

expressing Neurons Projecting to Nucleus Accumbens using Oxtr-ires-Cre Knock-in

prairie Voles (Microtus ochrogaster). Neuroscience *448*, 312–324.

https://doi.org/10.1016/j.neuroscience.2020.08.023.

106. Villanueva, C., Jacquier, S., and de Roux, N. (2012). DLK1 Is a Somato-Dendritic

Protein Expressed in Hypothalamic Arginine-Vasopressin and Oxytocin Neurons.

PLoS One *7*, e36134. https://doi.org/10.1371/journal.pone.0036134.

107. Insel, T.R. (1997). A neurobiological basis of social attachment. Am J Psychiatry

154, 726–735.

108. Carter, C.S., and Perkeybile, A.M. (2018). The Monogamy Paradox: What Do

Love and Sex Have to Do With It? Front. Ecol. Evol. *6*, 202.

https://doi.org/10.3389/fevo.2018.00202.

109. De Vries, G.J. (2004). Minireview: Sex Differences in Adult and Developing

Brains: Compensation, Compensation, Compensation. Endocrinology *145*, 1063–

1068. https://doi.org/10.1210/en.2003-1504.

110. Friard, O., and Gamba, M. (2016). BORIS: a free, versatile open-source event-

logging software for video/audio coding and live observations. Methods in ecology

and evolution *7*, 1325–1330.

111.   Risso, D., Ngai, J., Speed, T.P., and Dudoit, S. (2014). Normalization of RNA-seq

1575   data using factor analysis of control genes or samples. Nat Biotechnol *32*, 896–902.

1576   https://doi.org/10.1038/nbt.2931.

1577   112.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold

1578   change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

1579   https://doi.org/10.1186/s13059-014-0550-8.

1580   113.   Stephens, M., Carbonetto, P., Gerard, D., Lu, M., Sun, L., Willwerscheid, J., and

1581   Xiao, N. (2020). Methods for Adaptive Shrinkage, using Empirical Bayes [R package

1582   ashr version 2.2-47]. In.

1583   114.   Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene

1584   ontology analysis for RNA-seq: accounting for selection bias. Genome Biol *11*, R14.

1585   https://doi.org/10.1186/gb-2010-11-2-r14.

1586   115.   Yu, G. (2020). Gene Ontology Semantic Similarity Analysis Using GOSemSim. In

1587   Stem Cell Transcriptional Networks Methods in Molecular Biology., B. L. Kidder, ed.

1588   (Springer US), pp. 207–215. https://doi.org/10.1007/978-1-0716-0301-7_11.

1589   116.   Pico, A.R., Kelder, T., Van Iersel, M.P., Hanspers, K., Conklin, B.R., and Evelo, C.

1590   (2008). WikiPathways: Pathway Editing for the People. PLoS Biol *6*, e184.

1591   https://doi.org/10.1371/journal.pbio.0060184.

1592   117.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W.,

1593   Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting

1594   omics data. The Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

118.   Fu, R., Gillen, A.E., Sheridan, R.M., Tian, C., Daya, M., Hao, Y., Hesselberth,

J.R., and Riemondy, K.A. (2020). clustifyr: an R package for automated single-cell

RNA sequencing cluster classification. F1000Res *9*, 223.

https://doi.org/10.12688/f1000research.22969.2.

119.   Zhang, B., and Horvath, S. (2005). A general framework for weighted gene co-

expression network analysis. Stat Appl Genet Mol Biol *4*, Article17.

https://doi.org/10.2202/1544-6115.1128.

120.   Langfelder, P., Luo, R., Oldham, M.C., and Horvath, S. (2011). Is my network

module preserved and reproducible? PLoS Comput Biol *7*, e1001057.

https://doi.org/10.1371/journal.pcbi.1001057.

121.   Gandrillon, O., Solari, F., Legrand, C., Jurdic, P., and Samarut, J. (1996). A rapid

and convenient method to prepare DIG-labelled RNA probes for use in non-

radioactive in situ hybridization. Molecular and cellular probes *10*, 51–56.

1608

Figure 1: Oxtr reduces the amount of time naïve animals take to show partner preference

Figure 2: Oxtr suppresses promiscuous behaviors in a state dependent manner in females

Figure 3: Oxtr function influences different prosocial behaviors across pair bonding in males

Figure 4: Oxtr sex-specifically influences social interactions between potential mates

Figure 5: Post pair bonding gene expression is driven by female specific changes dependent on Oxtr signaling

Figure 6: Oxtr regulated genes are common to social behavior relevant processes across species and are similarly regulated across brain regions

Figure 7: Loss of Oxtr results in the decrease in density of Oxt and Avp expressing neurons in the PVN