Why are telomeres the length that they are? Insight from a phylogenetic comparative analysis

Derek M. Benson[a,*], Dylan J. Padilla Perez[b], and Dale F. DeNardo[c]

School of Life Sciences, Arizona State University, Tempe, Arizona 85287, USA.

**\*Corresponding author**

*a:* dmbenso4@asu.edu, *b:* dpadil10@asu.edu, *c:* denardo@asu.edu

**Running title:** Phylogenetic comparison of vertebrate telomeres

**Data Accessibility Statement**

A fully reproducible workflow of the data analyses, including R scripts and additional supporting
material, is available in the following repositories: Github https://dylan-
padilla.github.io/telomere_evo/, a Dryad link will be available upon acceptance.

**Conflict of interest**

The authors declare no conflict of interest.

**Author Contributions**

Derek M. Benson conceptualized the study, collected and curated the data, assisted with the
methodology and data analysis, and wrote the manuscript. Dylan J. Padilla-Perez was responsible
for the majority of the methodology and data analysis, created the figures, and assisted in writing
the manuscript. Dale F. DeNardo oversaw the project and contributed to conceptualization and
writing.
26

**Abstract**

Telomeres are short repeating nucleotide sequences at the ends of chromosomes that shorten with

every cellular replication. Despite the importance of keeping telomere length within a critical

homeostatic range, adult telomere length can differ by two orders of magnitude across vertebrate

species. Why telomere length varies so widely remains unknown, though popular hypotheses

suggest that body size, lifespan, and endothermy are key variables that have coevolved with

telomere length. To test the relationship among telomere length, telomerase activity (which extends

telomeres), and these variables, we modeled the evolution of telomere length across 122 vertebrate

species. We failed to find an influence of body mass, lifespan, or baseline metabolism on telomere

length. However, we found a significant interactive effect between baseline metabolism and body

mass. The presence of telomerase activity was positively correlated with telomere length across the 58

species where data for both existed. Taken together, our findings suggest that body mass may have

differentially influenced the evolution of telomere length in endotherms and ectotherms and indicate that

telomerase activity and telomere length may have coevolved.

**Keywords:** aging, cellular senescence, molecular evolution, oxidative stress.

## Introduction

Telomeres are DNA sequences that are critical for protecting the ends of chromosomes. Accordingly, all vertebrate chromosomes end in telomeres. Yet, despite the crucial role of telomeres and their ubiquity across vertebrates, the existing literature on absolute telomere length has not previously been comparatively analyzed to better understand the variation in telomere biology across major vertebrate clades.

Structurally, vertebrate telomeres are comprised of repeating TTAGGG nucleotide sequences and a protein complex known as shelterin (Rhodes and Giraldo, 1995; De Lange, 2005). The 3' end of telomeres ends in a G-strand overhang, which often forms a T-loop structure by invading and binding to double-stranded telomeric DNA (Cong et al., 2002; Haussmann and Marchetto, 2010). The functional limitations of DNA polymerase and the creation of Okazaki fragments lead to the progressive loss of telomeric DNA with each cellular replication (Gilson and Géli, 2007). Many species, including humans, experience replicative aging where cells enter senescence and stop replicating when telomere length hits a critical minimum (Saretzki and Von Zglinicki, 2002; Shay and Wright, 2005). In addition to the impacts of cellular replication, oxidative stress can also cause significant telomere shortening (Von Zglinicki, 2002). The telomeric 5'-GGG-3' sequence is particularly susceptible to damage by reactive oxygen species and conversion of one or more guanines into 8-oxoG (Kawanishi and Oikawa, 2004). Furthermore, the DNA damage response is repressed at telomeric DNA, and telomeric DNA within the single-stranded T-loop is not protected by hydrogen bonding and, due to the absence of a template strand, not able to be repaired by base excision repair (Ahmed and Lingner, 2018). In contrast to the cellular senescence associated with short telomeres, long telomeres are associated with increased cancer risk in some species, so it is crucial for an organism to maintain its telomere lengths within a critical range (Rode et al., 2016; Aviv et al., 2017; McNally et al., 2019).

To counter the progressive shortening of telomeres resulting from cellular replication,

organisms can prevent telomere shortening through antioxidant production and, in some cases,

elongate telomeres via telomerase activity and alternative telomere lengthening (Neumann et al. 2013;

Armstrong and Boonekamp 2023). Telomerase is a ribonucleoprotein reverse transcriptase comprised

of catalytic (TERT) and RNA (TERC) components (Cong et al., 2002). Telomerase extends the G

strand of telomeric DNA, after which the C strand is filled in by DNA polymerase $\alpha - primase$

(Bonetti et al., 2014). Despite its obvious benefits, in humans and many endotherms telomerase

activity is suppressed in most somatic tissues outside of embryonic stages (Wright et al., 1996). It has

been hypothesized that this inactivity of telomerase is an anti-cancer protection mechanism, as

telomerase activity enables continuous cellular replication and is almost always necessary for

cellular immortalization (Shay and Wright, 2011). In fact, telomerase activity is present in 85-90%

of all human tumors (Shay and Bacchetti, 1997).

Since their discovery, telomeres, telomere length, and telomerase activity, collectively

referred to as "telomere dynamics," have been studied extensively for their potential involvement in

both aging and cancer (Aubert and Lansdorp, 2008; Xu et al., 2013). Additionally, many studies have

investigated telomere dynamics in non-human vertebrates, revealing that telomere length, replicative

senescence, and repressed telomerase activity are not conserved traits but rather vary among species

(Gomes et al., 2010). In fact, telomere length ranges across species from one to hundreds of kb (Rocco

et al., 2001; Zhdanova et al., 2005). Why telomere length varies so widely across vertebrates remains

to be understood, though popular hypotheses suggest that body size, lifespan, and endothermy are key

variables that have coevolved with telomere length (Heidinger et al., 2012; Olsson et al., 2018; Pepke

and Eisenberg, 2022). Larger body size increases the number of cells an organism has which

increases the number of potentially cancerous cells, increased lifespan enables more mutations to

accumulate over time, and endothermy increases the rate of metabolism and cellular turnover, thus

increasing the mutation rate of cells (Peto et al., 1975; Seluanov et al., 2008; Hayes, 2010).

While the importance of telomere dynamics has resulted in a considerable number of

publications regarding the subject across numerous species, no literature has taken a

comprehensive, wide-scale comparative approach to review and analyze the relationships among

telomere dynamics, body size, lifespan, and endothermy across major vertebrate clades. The goals

of this review are to 1. Determine whether lifespan, body mass, and endothermy have affected the

evolution of telomere length; 2. Elucidate whether telomere length is differentially affected by these

variables among major vertebrate clades; and 3. Investigate whether telomere length and

telomerase expression have coevolved.

**Materials and Methods**

*Literature search and data collection*

To search for existing literature on telomere length and telomerase activity, key terms including

"telomere length", "telomerase activity", and specific species or taxonomic groups were input into

both Google Scholar and Web of Science (Figure S1). Given the discrepancies in telomere length

measurements across methodology, we only collected data from studies using telomere restriction

fragment analysis (TRF) as it is the most widely used quantitative measurement of telomere length. As

TRF can be performed with or without completely denaturing the DNA and measuring interstitial

telomere sequences (ITSs), we tested for an effect of denaturing on telomere length across all the

species in our analyses. As we did not detect any significant effect, we compared telomere length

across the methodologies, though we cannot completely discount the fact that methodology may

influence our findings. When available and for consistency, we utilized average adult telomere length

as reported in previous analyses that had collected data across multiple studies (see, Gomes et al., 2011;

Criscuolo et al., 2021; Pepke and Eisenberg, 2022). For species not included in these analyses, we

used the average telomere length value reported in individual studies. For studies that did not report

an average telomere length, we calculated average telomere length as the mean telomere length

across all individuals in a study. We did not include studies or datapoints on telomere length in

extremely young or old individuals (see the supporting information for more information on the

collection of data). Given that telomere length is generally strongly correlated but not constant across

somatic tissues within individuals (McLester-Davis et al., 2023), we tested for an effect of tissue type

on telomere length. As we did not detect an effect of tissue type on telomere length, we compared

telomere length across tissue types (see supporting information for more detail). Additionally, as

previous studies have shown no evidence for sex-based differences in adult telomere length across

vertebrates, we included data for both sexes in our analyses (Remot et al., 2020). Telomerase activity

in the analyzed cells (hereafter referred to as telomerase activity) was recorded as a binary variable,

being either absent or present in the somatic tissue of the study species. Telomerase activity data from

gonadal tissue were not utilized as many species exhibit telomerase activity in gonadal tissue but not

in other somatic tissues (Wright et al., 1996; Sun et al., 2021). For studies that reported telomere

length but not telomerase activity, we recorded telomerase activity as not available (n/a) for the

species. As telomere length data from Seluanov et al. (2007) was found to be highly correlated with

but longer than that of overlapping species analyzed by Gomes et al. (2011), we applied the

same correction factor (0.6617 x TL – 2.4454 kb) that was utilized by Pepke and Eisenberg

(2022). Once telomere length and telomerase activity data were collected, we obtained species-

specific maximum lifespan in years (hereafter referred to as lifespan) and adult mass in grams

primarily from the AnAge database (de Magalhães et al., 2024;

https://genomics.senescence.info/species/index.html). For species that did not have an entry in the

AnAge database, information on adult mass and lifespan was collected from various literature

sources (see supplemental table 1 for more information). If no data on lifespan or adult mass could

be found, the species was not used for any analysis. Additionally, as domestication may have

influenced the evolution of telomere length, animals with a long history of domestication were not

included in our analyses. We utilized telomere dynamics data for 122 vertebrate species across 42

orders and 88 families.

**Figure 1.**

*Vertebrate phylogeny*

To build a vertebrate phylogeny for this review, we used the Build a Timetree function available at

The TimeTree of Life database ([https://timetree.org/](https://timetree.org/)), which outputs a timetree of taxa of interest

extracted from the global timetree connecting species and publication-specific timetrees in the

database. For species that did not have phylogenetic data readily available, we utilized closely related

species or congeners when constructing the phylogeny (see the supporting information for more

detail). Although the placement of taxa in our species-level tree of vertebrates was generally well

supported, phylogenetic comparative methods appear to be robust to perturbations such as tree

misspecification or soft polytomies (Stone, 2011).

*Statistical Analysis*

To better understand the evolution of telomere length, we estimated its ancestral state across

vertebrates. We used the function *fastAnc* in the "phytools" package of R v. 4.2.2 (Revell, 2022),

which performs a reasonably fast estimation of the most likely ancestral states for a continuous

trait by taking advantage of the fact that the state computed for the root node of the tree during

Felsenstein (1985) contrasts algorithm is also the most likely estimate of the root node. Thus, the

function re-roots the tree at all internal nodes and computes the contrasts state at the root each

time. The function also computes variances or 95% confidence intervals on the estimates.

Subsequently, we used Phylogenetic Path Analysis (van der Bijl, 2018) to test our

hypotheses on the relationships among body mass, baseline metabolism (i.e., the dichotomous

distinction between ectotherms and endotherms), lifespan, and telomere length across species.

Phylogenetic path analysis uses Phylogenetic Generalized Least-squares (PGLS) to perform

confirmatory path analysis based on the directed separation method–"d-separation"–as described in

Shipley (2013). We examined the explanatory power of 8 competing models involving different

relationships among variables, which enabled us to identify the most plausible hypothesis given the

available data, and thus infer the relative importance of mechanisms described by optimality models.

To select the most likely path model, we relied on the $C$ statistic of the model, which is a maximum

likelihood estimate that follows a $x^2$ distribution with degrees of freedom $df = 2k$. Therefore, it

provides a convenient statistic for testing the goodness of fit of the whole path model (Shipley, 2013).

If the path model did not provide a good fit to the data because the p-value of the $C$ statistic was below

the alpha value (i.e., $< 0.05$), we proceed to examine the conditional independencies tested in the path

model. A conditional independency specifies the list of pairs of variables that are statistically

independent conditioning on a set of other variables in the causal path model. A conditional

independency is only supported by the data if its p-value is greater than the alpha value (i.e., p-value $>$

$0.05$). The most likely conditional independency suggested by path analysis was then refitted

assuming that the species trait values evolved via a Brownian motion model, an Ornstein–Uhlenbeck

motion model, and a Pagel's lambda model using the R package 'ape' v. 5.6.2. To evaluate the models'

goodness of fit, we used information-theoretic approach such as AICc. We ranked the candidate

models accordingly and selected the most likely one for inferences (lowest value of AICc). To

investigate the effects of sample size on the exploratory power of our models, we conducted a

sensitivity analysis across the entire dataset manually utilizing the previously determined most likely

model. In brief, we ran 1000 simulations for each of five bins randomly removing 10, 20, 30, 40,

and 50% of species, for a total of 5000 simulations. For each bin, we calculated the mean parameter

estimate for the model intercept, body mass, lifespan, baseline metabolism, and the baseline

metabolism x body mass interaction. Additionally, we calculated the standard deviation for each

parameter, and the 95% CI for each parameter. Significant parameter estimates were those where the

95% CI did not include 0.

We also estimated the degree of phylogenetic signal in our data based on the most likely

model. The "ape" package of R enabled us to modify the correlation structure of the residuals in the

function gls. Thus, we assigned different fixed values to the parameter $\lambda$, ranging from zero (no

phylogenetic signal, equivalent to a "star" phylogeny) to one (consistent with Brownian motion).

Intermediate values of $\lambda$ imply that the data support a model that is somewhere between a "star"

phylogeny and Brownian motion. $\lambda$ represents the amount of phylogenetic "signal" in the data,

conditional on the tree. In a regression context, we used maximum likelihood to simultaneously fit

the regression model and estimate $\lambda$. Because confidence intervals for $\lambda$ were provided by the models,

we tested the hypotheses that $\lambda = 0$ (i.e., independence) and $\lambda = 1$ (Brownian Motion).

Lastly, we investigated whether telomerase activity changes as a function of telomere length.

To do that, we modeled the correlated evolution of telomere length and telomerase activity under

the threshold model (Felsenstein, 2012). This model creates a natural framework by which one can

evaluate the evolutionary correlation between a continuous variable (e.g., telomere length) and a

discrete variable (e.g., telomerase activity). In R, the correlational threshold model can be performed

through the function threshBayes available in the package "phytools" (Revell, 2022). It also enabled

us to generate a three-panel plot showing the likelihood profile, the mean acceptance rates (using a

default sliding window), and a profile plot for the correlation coefficient (Figure S2). In addition, we

used the density function to later plot a posterior density of the correlation coefficient and added a

95% high probability density interval (HPD) around the value of $r$. We did this by extracting a post-

burn-in sample of values of $r$ (assuming a 20% burn-in) and using the function HPD interval from

the package "coda". To produce good visualization of our results and ensure that they are reproducible,

we carried out all the analyses in the free software for statistical computing R v.4.2.2 (R Core Team,

2022).

**Results**

Our results indicated no effect of lifespan on telomere length. However, the evolution of telomere

length across vertebrates appeared to be influenced by the interaction between baseline

metabolism and body mass of species (Figure S3-4). The information-theoretic approach revealed

that a model including the main effects of these variables was strongly supported (Table 1).

Generally, telomere length increased with body mass in ectotherms and decreased with body mass

in endotherms (Figure 2). Finally, our analyses indicate that there may have been many transitions

in telomere length across vertebrates (Figure 1).

Figure 1

Table 1.

Figure 2.

As the scatter in the data indicates substantial noise, we examined the impact that different sample

sizes could have on our results. Randomly removing 10, 20, 30, 40, and 50% of species in the

analyses did not affect the likelihood of finding a significant effect of the interaction between

baseline metabolism and body mass on telomere length, though the independent effects of

lifespan, baseline metabolism, and body mass were significantly affected by sample size (Figure

S5).

Figure 3.

Based on a likelihood ratio test (X2), models with $\lambda$ fixed at 0 and 1 were significantly

different from one that had $\lambda$ fixed at 0.5. Thus, we rejected the hypotheses of independent

residuals and Brownian motion. The most likely model suggested an intermediate degree of

phylogenetic signal in the data.

Finally, the threshold model of correlated evolution indicated that the evolution of

telomere length is correlated with the presence of telomerase activity. According to the posterior

probability distribution of the correlation coefficient ($r = 0.51$), the presence of telomerase activity

significantly increased with telomere length. This conclusion was based on the 95% high

probability density ($HPD = 0.18 - 0.79$) interval around $r$, which did not include 0 (Figure 3 and

S2).

**Discussion**

To our knowledge, ours is the first study to test for correlates between telomere length and

various variables across vertebrate clades, though previous studies have investigated within clades

(Gomes et al., 2011; Criscuolo et al., 2021; Pepke and Eisenberg, 2022). Across vertebrates, we

found evidence that the interaction between baseline metabolism and body mass is correlated with

telomere length. However, we failed to find evidence that body mass, lifespan, baseline

metabolism, or any other interactions are correlated with telomere length across major vertebrate

clades. It should be noted that while we did not detect an effect of tissue type or methodology for our

analyses, we cannot discount the possibility that the heterogenous nature of the data analyzed may

influence our results and represent a major limitation of our study. Still, our study lends valuable

insight into the evolution of telomere length and suggests that the evolution of telomere length has

been influenced by baseline metabolism and that the effect of body mass on telomere length may

differ between endotherms and ectotherms.

The relationship between body mass and telomere length remains unclear, as previous

research on birds and mammals has yielded contrasting results (Gomes et al. 2011; Criscuolo et al.

2021; Pepke and Eisenberg 2022). As we included fishes and ectothermic diapsids, our results are

not directly comparable, though they suggest that the correlation between body mass and telomere

length may be dependent on baseline metabolism. One possible explanation for the differential

effect of body mass based on baseline metabolism is that high stable body temperatures may be

associated with an increased risk of malignant tumors (Olsson et al. 2018). While plausible, many

ectotherms prefer and maintain body temperatures above that of endotherms, and previous

research has failed to find a relationship between high stable body temperature and an increase in

malignant tumors (Stevenson 1985; Olsson et al. 2018). Alternatively, it may be that the difference

in metabolic rate between endotherms and ectotherms has significantly affected the evolution of

telomere length, as metabolic rate has been shown to be positively correlated with DNA damage

(Moretton and Loizou 2020). Thus, the selective pressure that body mass imposes on telomere

length may differ between endotherms and ectotherms, as is evident by the difference in slopes

relating telomere length to body mass. Regardless of the reason behind the relationship between

baseline metabolism and telomere length, it is important to consider the disparity in sample size

between endotherms and ectotherms in our analyses. Vertebrate ectotherms are understudied in

terms of telomere biology when compared to endotherms, and information on major clades

including Anurans and Agnatha is nearly or completely absent. Given the importance of these

groups for understanding vertebrate evolution, it is likely that quantifying telomere length in

species from these clades will further elucidate how baseline metabolism is related to telomere

length. Nevertheless, our results indicate that baseline metabolism and body mass may have

played a significant role in the evolution of telomere length.

In contrast to previous research in mammals (Pepke and Eisenberg 2022), we failed to find

evidence that telomere length is correlated with lifespan. Conclusions on this relationship remain

divided, with studies both supporting and refuting the idea that a correlation exists (Gomes et al.

2011; Tricola et al., 2018; Criscuolo et al., 2021; Pepke et al., 2023b). This discrepancy may in

part be explained by the use of either early-life (within the first few weeks of life) or adult

telomere length. We used adult telomere length for several reasons: Firstly, early-life telomere

length changes rapidly for many though not all species (Watson et al., 2015; Pepke et al., 2023a).

Thus, early-life telomere length may not matter as much as preserved telomere length. Secondly,

early-life telomere length fails to account for telomere length modulation later in life. Several

long-lived species have been shown to increase telomere length in adulthood (Haussmann et al.,

2003). Furthermore, telomere rate of change has been repeatedly shown to be correlated with

lifespan (Tricola et al., 2018; Criscuolo et al., 2021; Whittemore et al., 2019). While adult

telomere length does not measure this, adult telomere length is affected by shortening and

lengthening and may serve as a more effective measurement of telomere length for a species.

However, among adults, age can vary considerably in longer lived species and such variation in

age among adults could confound results.

Despite variation in telomere length within and among clades, our analyses indicate that

long telomeres are correlated with the presence of adult telomerase activity. Long telomere length

and the presence of telomerase activity might be thought to increase cancer risk as previous

literature suggests that selection favors short telomeres and the repression of telomerase as a

mechanism to prevent cancer (Risques and Promislow, 2018). However, while cancer is

detrimental to organismal health, the vast majority of wild animals do not live to the age where

cancer becomes a significant risk factor (Rozhok and DeGregori, 2016). Thus, other variables

such as pace of life and reproductive investment, rather than cancer risk, may have exerted

stronger selective pressures on the evolution of telomere length and telomerase activity.

Additionally, given that our and previous analyses are strongly biased towards mammal and bird,

it is possible that the correlations between telomerase activity, long telomeres, and cancer risk are

not reflective of vertebrates in general (Gomes et al., 2011; Pepke and Eisenberg, 2022). For

example, the limited data in fishes demonstrate that most species in our analysis display

telomerase activity with telomeres <15 kb in length, a phenomenon rarely seen among mammals

(Zhdanova et al., 2005; Elmore et al., 2008; Gomes et al., 2011).

Making comparisons across and among subclades of a larger tree can prove valuable,

particularly where large timescales are involved and thus where evolutionary patterns and

processes are unlikely to be constant. However, there are important considerations when choosing

which clades to study, including sample size and whether there is sufficient within-clade diversity

for statistical analysis (Baker et al., 2021). Given our findings and the indications that baseline

metabolism may play a role in shaping the evolution of telomere length as well as the current

literature's bias towards mammals and birds, future studies need to include a wider range of

vertebrates, preferably with a similar degree of representation across the major taxa, to gain a

more thorough understanding of (1) how telomere length varies within and among major clades,

and (2) how the effects of variables such as lifespan and body mass on telomere length may differ

among clades.

Our data may suffer from both an extremely variable number of species per clade and a

large variation in telomere size among species, preventing us from making a general conclusion

about the evolution of telomere size across vertebrates. While sample size may have influenced

our results, our findings appear to be fairly robust given that the random removal of up to 50% of

the species in our study did not affect the chance of finding a significant correlation between

telomere length and the interaction between body mass and baseline metabolism. In support of the

idea that sample size may be impacting our findings, previous studies have found differing results

depending on the clade and range of taxa used when investigating the relationship between

telomere length and lifespan (Seluanov et al., 2007; Gomes et al., 2011; Criscuolo et al., 2021;

Pepke and Eisenberg, 2022). For instance, work in rodents found no evidence that lifespan was

correlated with telomere length, while several analyses that include a wider range of mammals

have concluded that telomere length is strongly correlated with lifespan (Seluanov et al., 2007;

Gomes et al., 2011; Pepke and Eisenberg, 2022).

While our greater overall sample size may explain the differences between our results and

those of previous studies, many variables, such as diet and habitat, may have also significantly

influenced the evolution of telomere length. While diet has been shown to significantly affect

telomere length in mice and humans, it is not known whether carnivory or herbivory differentially

impact telomere length in mammals (Paul, 2011; Gokarn et al., 2018). Furthermore, research on a

wide range of neotropical bat species found that frugivorous, omnivorous, and carnivorous species

differ in oxidative stress and antioxidant capacity (Schneeberger et al., 2014). Given that telomere

length is associated with oxidative stress, it may be that diet has had a significant impact on the

evolution of telomere length (Von Zglinicki, 2002). In addition, habitat and environment have

been shown to impact telomere length. For example, moose (*Alces alces*) populations in boreal

regions have significantly longer telomeres than do those in sarmatic mixed forest and montane

regions (Fohringer et al., 2022). Furthermore, tropical and temperate birds have differentially

evolved responses to oxidative damage, and telomere length in young birds of the genus *Saxicola*

differs between species in tropical and temperate regions (Jimenez et al., 2013; Apfelbeck et al.,

2019). In addition to needing greater and more balanced phylogenic coverage, future studies

investigating the evolution of telomere length should also investigate a wider range of life history

and ecological traits.

In conclusion, despite the possible complications including a mammal-biased sample size,

our study provides evidence that telomerase activity and telomere length in adult life have

coevolved, with longer telomeres being associated with the presence of telomerase activity.

Additionally, we show that the interaction between body mass and baseline metabolism is related

to telomere length. Finally, we failed to find any relationship between lifespan and telomere

length. Future studies should incorporate a wider range of non-mammalian species in their

analyses to further our understanding of the variables affecting telomere length while also

attempting to account for other potentially confounding variables such as unbalanced sample

sizes, chromosome size, diet, habitat, pace of life, and additional life history traits.

**Acknowledgements**

We thank the anonymous reviewers for feedback on earlier drafts.

**References**

Ahmed, W., & Lingner, J. (2018). Impact of oxidative stress on telomere biology. *Differentiation*, *99*, 21–27. https://doi.org/10.1016/j.diff.2017.12.002

Apfelbeck, B., Haussmann, M. F., Boner, W., Flinks, H., Griffiths, K., Illera, J. C., Mortega, K. G., Sisson, Z., Smiddy, P., & Helm, B. (2019). Divergent patterns of telomere shortening in tropical compared to temperate stonechats. *Ecology and Evolution*, *9*(1), 511–521. https://doi.org/10.1002/ece3.4769

Armstrong, E., & Boonekamp, J. (2023). Does oxidative stress shorten telomeres *in vivo*? A meta-analysis. *Ageing Research Reviews*, *85*, 101854. https://doi.org/10.1016/j.arr.2023.101854

Aviv, A., Anderson, J. J., & Shay, J. W. (2017). Mutations, cancer and the telomere length paradox. *Trends in Cancer*, *3*(4), 253–258. https://doi.org/10.1016/j.trecan.2017.02.005

Aubert, G., & Lansdorp, P. M. (2008). Telomeres and aging. *Physiological Reviews*, *88*(2), 557–579. https://doi.org/10.1152/physrev.00026.2007

Baker, J., Meade, A., Pagel, M., & Venditti, C. (2021). Nothing wrong with the analysis of clades in comparative evolutionary studies: A reply to Poe et al. *Systematic Biology*, *70*(1), 197–201. https://doi.org/10.1093/sysbio/syaa067

Bijl, W. van der. (2018). phylopath: Easy phylogenetic path analysis in R. *PeerJ*, *6*, e4718. https://doi.org/10.7717/peerj.4718

Bonetti, D., Martina, M., Falcettoni, M., & Longhese, M. P. (2014). Telomere-end processing: Mechanisms and regulation. *Chromosoma*, *123*(1), 57–66. https://doi.org/10.1007/s00412-013-0440-y

Cong, Y.-S., Wright, W. E., & Shay, J. W. (2002). Human telomerase and its regulation. *Microbiology and Molecular Biology Reviews*, *66*(3), 407–425. https://doi.org/10.1128/mmbr.66.3.407-425.2002

Criscuolo, F., Dobson, F. S., & Schull, Q. (2021). The influence of phylogeny and life history on telomere lengths and telomere rate of change among bird species: A meta-analysis. *Ecology and Evolution*, *11*(19), 12908–12922. https://doi.org/10.1002/ece3.7931

de Magalhães, J. P., Abidi, Z., dos Santos, G. A., Avelar, R. A., Barardo, D., Chatsirisupachai, K., Clark, P., De-Souza, E. A., Johnson, E. J., Lopes, I., Novoa, G., Senez, L., Talay, A., Thornton, D., & To, P. K. P. (2024). Human Ageing Genomic Resources: Updates on key databases in ageing research. *Nucleic Acids Research*, *52*(D1), D900–D908. https://doi.org/10.1093/nar/gkad927

Elmore, L. W., Norris, M. W., Sircar, S., Bright, A. T., McChesney, P. A., Winn, R. N., & Holt, S. E. (2008). Upregulation of telomerase function during tissue regeneration. *Experimental Biology and Medicine (Maywood, N.J.)*, *233*(8), 958–967. https://doi.org/10.3181/0712-RM-345

Felsenstein, J. (2012). A comparative method for both discrete and continuous characters using the threshold model. *The American Naturalist*, *179*(2), 145–156. https://doi.org/10.1086/663681

Fohringer, C., Hoelzl, F., Allen, A. M., Cayol, C., Ericsson, G., Spong, G., Smith, S., & Singh, N. J. (2022). Large mammal telomere length variation across ecoregions. *BMC Ecology and Evolution*, *22*(1), 105. https://doi.org/10.1186/s12862-022-02050-5

Gilson, E., & Géli, V. (2007). How telomeres are replicated. *Nature Reviews Molecular Cell Biology*, *8*(10), Article 10. https://doi.org/10.1038/nrm2259

Gokarn, R., Solon-Biet, S., Youngson, N. A., Wahl, D., Cogger, V. C., McMahon, A. C., Cooney, G. J., Ballard, J. W. O., Raubenheimer, D., Morris, M. J., Simpson, S. J., & Le Couteur, D. G. (2018). The relationship between dietary macronutrients and hepatic telomere length in aging mice. *The Journals of Gerontology: Series A*, *73*(4), 446–449. https://doi.org/10.1093/gerona/glx186

Gomes, N. M. V., Ryder, O. A., Houck, M. L., Charter, S. J., Walker, W., Forsyth, N. R., Austad, S. N., Venditti, C., Pagel, M., Shay, J. W., & Wright, W. E. (2011). Comparative biology of mammalian telomeres: Hypotheses on ancestral states and the roles of telomeres in longevity determination. *Aging Cell*, *10*(5), 761–768. https://doi.org/10.1111/j.1474-9726.2011.00718.x

Gomes, N. M. V., Shay, J. W., & Wright, W. E. (2010). Telomere biology in Metazoa. *FEBS Letters*, *584*(17), 3741–3751. https://doi.org/10.1016/j.febslet.2010.07.031

Haussmann, M. F., & Marchetto, N. M. (2010). Telomeres: Linking stress and survival, ecology and evolution. *Current Zoology*, *56*(6), 714–727. https://doi.org/10.1093/czoolo/56.6.714

Haussmann, M. F., Winkler, D. W., O'Reilly, K. M., Huntington, C. E., Nisbet, I. C. T., & Vleck, C. M. (2003). Telomeres shorten more slowly in long-lived birds and mammals than in short–lived ones. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *270*(1522), 1387–1392. https://doi.org/10.1098/rspb.2003.2385

Hayes, J. P. (2010). Metabolic rates, genetic constraints, and the evolution of endothermy. *Journal of Evolutionary Biology*, *23*(9), 1868–1877. https://doi.org/10.1111/j.1420-9101.2010.02053.x

Heidinger, B. J., Blount, J. D., Boner, W., Griffiths, K., Metcalfe, N. B., & Monaghan, P. (2012). Telomere length in early life predicts lifespan. *Proceedings of the National Academy of Sciences*, *109*(5), 1743–1748. https://doi.org/10.1073/pnas.1113306109

Jimenez, A. G., Harper, J. M., Queenborough, S. A., & Williams, J. B. (2013). Linkages between the life-history evolution of tropical and temperate birds and the resistance of cultured skin fibroblasts to oxidative and non-oxidative chemical injury. *Journal of Experimental Biology*, *216*(8), 1373–1380. https://doi.org/10.1242/jeb.079889

Kawanishi, S., & Oikawa, S. (2004). Mechanism of telomere shortening by oxidative stress.

*Annals of the New York Academy of Sciences*, *1019*(1), 278–284.

https://doi.org/10.1196/annals.1297.047

Lange, T. de. (2005). Shelterin: The protein complex that shapes and safeguards human telomeres.

*Genes & Development*, *19*(18), 2100–2110. https://doi.org/10.1101/gad.1346005

McLester-Davis, L. W. Y., Estrada, P., Hastings, W. J., Kataria, L. A., Martin, N. A.,

Siebeneicher, J. T., Tristano, R. I., Mayne, C. V., Horlick, R. P., O'Connell, S. S., &

Drury, S. S. (2023). A review and meta-analysis: Cross-tissue telomere length correlations

in healthy humans. *Ageing Research Reviews*, *88*, 101942.

https://doi.org/10.1016/j.arr.2023.101942

McNally, E. J., Luncsford, P. J., & Armanios, M. (2019). Long telomeres and cancer risk: The

price of cellular immortality. *The Journal of Clinical Investigation*, *129*(9), 3474–3481.

https://doi.org/10.1172/JCI120851

Moretton, A., & Loizou, J. I. (2020). Interplay between Cellular Metabolism and the DNA

Damage Response in Cancer. *Cancers*, *12*(8), Article 8.

https://doi.org/10.3390/cancers12082051

Neumann, A. A., Watson, C. M., Noble, J. R., Pickett, H. A., Tam, P. P. L., & Reddel, R. R.

(2013). Alternative lengthening of telomeres in normal mammalian somatic cells. *Genes &*

*Development*, *27*(1), 18–23. https://doi.org/10.1101/gad.205062.112

Olsson, M., Wapstra, E., & Friesen, C. (2018). Ectothermic telomeres: It's time they came in from

the cold. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

*373*(1741), 20160449. https://doi.org/10.1098/rstb.2016.0449

Paul, L. (2011). Diet, nutrition and telomere length. *The Journal of Nutritional Biochemistry*,

*22*(10), 895–901. https://doi.org/10.1016/j.jnutbio.2010.12.001

Pepke, M. L., & Eisenberg, D. T. A. (2022). On the comparative biology of mammalian

telomeres: Telomere length co-evolves with body mass, lifespan and cancer risk.

*Molecular Ecology*, *31*(23), 6286–6296. https://doi.org/10.1111/mec.15870

Pepke, M. L., Kvalnes, T., Wright, J., Araya-Ajoy, Y. G., Ranke, P. S., Boner, W., Monaghan, P.,

Sæther, B.-E., Jensen, H., & Ringsby, T. H. (2023b). Longitudinal telomere dynamics

within natural lifespans of a wild bird. *Scientific Reports*, *13*(1), Article 1.

https://doi.org/10.1038/s41598-023-31435-9

Pepke, M. L., Ringsby, T. H., & Eisenberg, D. T. A. (2023a). The evolution of early-life telomere

length, pace-of-life and telomere-chromosome length dynamics in birds. *Molecular

Ecology*, *32*(11), 2898–2912. https://doi.org/10.1111/mec.16907

Peto, R., Roe, F. J., Lee, P. N., Levy, L., & Clack, J. (1975). Cancer and ageing in mice and men.

*British Journal of Cancer*, *32*(4), Article 4. https://doi.org/10.1038/bjc.1975.242

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for

Statistical Computing. https://www.R-project.org

Revell, L. J., & Harmon, L. J. (2022). Phylogenetic comparative methods in R. Princeton

University Press.

Rhodes, D., & Giraldo, R. (1995). Telomere structure and function. *Current Opinion in Structural

Biology*, *5*(3), 311–322. https://doi.org/10.1016/0959-440X(95)80092-1

Risques, R. A., & Promislow, D. E. L. (2018). All's well that ends well: Why large species have

short telomeres. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

*373*(1741), 20160448. https://doi.org/10.1098/rstb.2016.0448

Rocco, L., Costagliola, D., & Stingo, V. (2001). (TTAGGG)n telomeric sequence in selachian

chromosomes. *Heredity*, *87*(5), Article 5. https://doi.org/10.1046/j.1365-

2540.2001.00945.x

Rode, L., Nordestgaard, B. G., & Bojesen, S. E. (2016). Long telomeres and cancer risk among

   95 568 individuals from the general population. *International Journal of Epidemiology*,

   *45*(5), 1634–1643. https://doi.org/10.1093/ije/dyw179

Rozhok, A. I., & DeGregori, J. (2016). The evolution of lifespan and age-dependent cancer risk.

   *Trends in Cancer*, *2*(10), 552–560. https://doi.org/10.1016/j.trecan.2016.09.004

Saretzki, G., & Von Zglinicki, T. (2002). Replicative aging, telomeres, and oxidative stress.

   *Annals of the New York Academy of Sciences*, *959*(1), 24–29.

   https://doi.org/10.1111/j.1749-6632.2002.tb02079.x

Schneeberger, K., Czirják, G. Á., & Voigt, C. C. (2014). Frugivory is associated with low

   measures of plasma oxidative stress and high antioxidant concentration in free-ranging

   bats. *Die Naturwissenschaften*, *101*(4), 285–290. https://doi.org/10.1007/s00114-014-

   1155-5

Seluanov, A., Chen, Z., Hine, C., Sasahara, T. H. C., Ribeiro, A. A. C. M., Catania, K. C.,

   Presgraves, D. C., & Gorbunova, V. (2007). Telomerase activity coevolves with body

   mass not lifespan. *Aging Cell*, *6*(1), 45–52. https://doi.org/10.1111/j.1474-

   9726.2006.00262.x

Seluanov, A., Hine, C., Bozzella, M., Hall, A., Sasahara, T. H. C., Ribeiro, A. A. C. M., Catania,

   K. C., Presgraves, D. C., & Gorbunova, V. (2008). Distinct tumor suppressor mechanisms

   evolve in rodent species that differ in size and lifespan. *Aging Cell*, *7*(6), 813–823.

   https://doi.org/10.1111/j.1474-9726.2008.00431.x

Shay, J. W., & S. A. Bacchetti. (1997). A survey of telomerase activity in human cancer.

   *European Journal of Cancer*, *33*, 787–791. https://doi.org/10.1016/S0959-8049(97)00062-

Shay, J. W., & Wright, W. E. (2005). Senescence and immortalization: Role of telomeres and

   telomerase. *Carcinogenesis*, *26*(5), 867–874. https://doi.org/10.1093/carcin/bgh296

Shay, J. W., & Wright, W. E. (2011). Role of telomeres and telomerase in cancer. *Seminars in Cancer Biology*, *21*(6), 349–353. https://doi.org/10.1016/j.semcancer.2011.10.001

Shipley, B. (2013). The AIC model selection method applied to path analytic models compared using a d-separation test. *Ecology*, *94*(3), 560–564. https://doi.org/10.1890/12-0976.1

Stone, E. A. (2011). Why the phylogenetic regression appears robust to tree misspecification. *Systematic Biology*, *60*(3), 245–260. https://doi.org/10.1093/sysbio/syq098

Sun, J., Liu, W., Guo, Y., Zhang, H., Jiang, D., Luo, Y., Liu, R., & Chen, C. (2021). Characterization of tree shrew telomeres and telomerase. *Journal of Genetics and Genomics*, *48*(7), 631–639. https://doi.org/10.1016/j.jgg.2021.06.004

Tricola, G. M., Simons, M. J. P., Atema, E., Boughton, R. K., Brown, J. L., Dearborn, D. C., Divoky, G., Eimes, J. A., Huntington, C. E., Kitaysky, A. S., Juola, F. A., Lank, D. B., Litwa, H. P., Mulder, E. G. A., Nisbet, I. C. T., Okanoya, K., Safran, R. J., Schoech, S. J., Schreiber, E. A., … Haussmann, M. F. (2018). The rate of telomere loss is related to maximum lifespan in birds. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *373*(1741), 20160445. https://doi.org/10.1098/rstb.2016.0445

Watson, H., Bolton, M., & Monaghan, P. (2015). Variation in early-life telomere dynamics in a long-lived bird: Links to environmental conditions and survival. *Journal of Experimental Biology*, *218*(5), 668–674. https://doi.org/10.1242/jeb.104265

Whittemore, K., Vera, E., Martínez-Nevado, E., Sanpera, C., & Blasco, M. A. (2019). Telomere shortening rate predicts species life span. *Proceedings of the National Academy of Sciences*, *116*(30), 15122–15127. https://doi.org/10.1073/pnas.1902452116

Wright, W. E., Piatyszek, M. A., Rainey, W. E., Byrd, W., & Shay, J. W. (1996). Telomerase activity in human germline and embryonic tissues and cells. *Developmental Genetics*, *18*(2), 173–179. https://doi.org/10.1002/(SICI)1520-6408(1996)18:2<173::AID-DVG10>3.0.CO;2-3

Xu, L., Li, S., & Stohr, B. A. (2013). The role of telomere biology in cancer. *Annual Review of Pathology: Mechanisms of Disease*, *8*(1), 49–78. https://doi.org/10.1146/annurev-pathol-020712-164030

Zglinicki, T. von. (2002). Oxidative stress shortens telomeres. *Trends in Biochemical Sciences*, *27*(7), 339–344. https://doi.org/10.1016/S0968-0004(02)02110-2

Zhdanova, N. S., Karamisheva, T. V., Minina, J., Astakhova, N. M., Lansdorp, P., Kammori, M., Rubtsov, N. B., & Searle, J. B. (2005). Unusual distribution pattern of telomeric repeats in the shrews *Sorex araneus* and *Sorex granarius*. *Chromosome Research*, *13*(6), 617–625. https://doi.org/10.1007/s10577-005-0988-3

**Table with captions**

Table 1: Contrast of parameters estimated by the most likely model according to the information-theoretic criteria of selection.

|  | t-value | p-value |
| --- | --- | --- |
| (Intercept) | 6.84 | <0.001 |
| Baseline metabolism | 0.81 | 0.42 |
| Log mass | 1.11 | 0.27 |
| Log lifespan | 1.37 | 0.18 |
| Log mass x Baseline metabolism | 2.41 | 0.018 |

**Figures with captions**

Figure 1: Telomere length, body mass, baseline metabolism, and telomerase activity across 122 vertebrate species. Warmer colors represent shorter telomeres.

Figure 2: Evolution of telomere length as a function of the interaction between body mase and baseline metabolism across vertebrates. Shading around the regression lines represents the standard error.

Figure 3: Posterior density of the correlation between telomere length and telomerase activity in vertebrates. The dashed line represents the estimated correlation coefficient (r = 0.51), representing the correlation between telomere length and the presence or absence of telomerase activity. Additionally, the 95% high probability density (HPD = 0.18 – 0.79) is represented by the solid black bar below the distribution.

**Supplementary Material**

Figure S1. PRISMA flow diagram depicting the data collection for this study.

Figure S2: Profile plots from a Bayesian MCMC analysis of the threshold model for telomere length and telomerase activity in vertebrates. (a) The likelihood profile and (b) Profile of the correlation coefficient, r.

Figure S3: Path models depicting relationships among telomere length and predictor variables derived from optimality models (see text for details). Abbreviation in the path diagrams are as follows: TL, telomere length; BM, body mass; LS, lifespan; M, baseline metabolism.

Figure S4: Relative importance of the causal models describing the evolution of telomere length across vertebrates. Bar labels are p-values; significance indicates rejection.

Figure S5: Sensitivity analysis investigating the effect of sample size on the parameter estimate of the intercept (red), body mass (purple), lifespan (blue), baseline metabolism (tan), and the baseline metabolism body mass interaction (green) on telomere length. Error bars represent the 95% CI of each parameter. 95% CIs not including zero indicates significance.