Multi-behavioral phenotyping in zebrafish identifies a novel disruptor of non-associative learning with environmental and human relevance

David Leuthold[1*], Nadia K. Herold[1], Jana Nerlich[2], Kristina Bartmann[3,4], Ilka Scharkin[3], Stefan J. Hallermann[2], Nicole Schweiger[1], Ellen Fritsche[3,4,5], Tamara Tal[1,2*]

[1]Department of Ecotoxicology, Helmholtz-Centre for Environmental Research – UFZ, Leipzig, Germany; [2]Medical Faculty, Carl Ludwig Institute of Physiology, University of Leipzig, Leipzig, Germany; [3]IUF – Leibniz Research Institute for Environmental Medicine, Düsseldorf, Germany; [4]DNTOX GmbH, Düsseldorf, Germany; [5]Swiss Centre for Applied Human Toxicology, University of Basel, Basel, Switzerland.

*Corresponding authors: David Leuthold; david.leuthold@ufz.de; Helmholtz-Centre for Environmental Research – UFZ; Permoser Straße 15; 04318 Leipzig; Germany, Tamara Tal; tamara.tal@ufz.de; Helmholtz-Centre for Environmental Research – UFZ; Permoser Straße 15; 04318 Leipzig; Germany.

Declaration of conflicts of interest

K.B. and E.F. are shareholders of the company DNTOX which provides DNT *in vitro* battery assay service. They declare no potential conflict of interest concerning the data included in this article. All other authors declare no competing interests.

Abstract

The vertebrate nervous system is vulnerable to chemical toxicity and the widespread release of chemicals into the environment outstrips the capacity to assess their safety. We devised a battery of automated behavior assays in larval zebrafish (*Danio rerio*), a 3R-compliant model amenable to higher-throughput chemical screens. The battery captures stereotypical visual and acoustic behaviors including habituation, a form of non-associative learning. Known pharmacological modulators of habituation in zebrafish evoked distinct behavioral patterns. By screening chemicals positive for *ex vivo* N-methyl-D-aspartate receptor (NMDAR) modulation, we identified chlorophene, a biocide that caused sedation, paradoxical excitation, and reduced habituation in zebrafish. Using *in silico* target predictions and pharmacological interventions, we discovered that chlorophene acts via gamma-aminobutyric acid A receptors (GABA$_A$Rs), a previously unknown target site. Orthogonal validation in cultured mouse cortical neurons and human stem cell-derived BrainSpheres confirmed chlorophene's interaction with GABA$_A$Rs. Together, multi-behavioral phenotyping in zebrafish can accelerate the identification of neurotoxicants and their underlying mode of action.

Main

The planetary boundary for chemical pollution has been exceeded based on the concept that the production and release of chemicals outstrips the capacity to monitor human and environmental exposure levels and to perform adequate safety assessments[1]. The global inventory of registered chemicals and mixtures used in commerce totals more than 350,000 substances[2]. The widespread release of anthropogenic chemicals into air, land, and water represents an existential threat to environmental and human health as chemical pollution represents a significant environmental risk factor for disease and premature death[3,4]. The nervous system is one of the most vulnerable organ systems to toxic injury due to the interdependence of neurodevelopmental processes and where perturbations can result in profound structural and/or functional consequences[5]. Epidemiological studies indicate that exposure to pesticides (e.g. rotenone, paraquat, chlorpyrifos), metals (e.g. lead, mercury), neuroactive drugs (e.g. benzodiazepines, anticholinergics), or industrial chemicals (e.g. polychlorinated biphenyls, brominated flame retardants, phthalates) contributes to neurobehavioral pathologies such as Parkinson's disease[6,7], Alzheimer's disease[8–10], attention deficit hyperactivity disorder[11,12], and other neurodevelopmental deficits[13]. These associations are supported by experimental findings in mice[14–18], rats[19,20], non-human primates[21,22], and other, non-mammalian models[23,24]. Despite these associations, most neurotoxicants remain unknown[13]. This is highlighted by the fact that, to date, fewer than 200 chemicals[25] have been tested according to guideline neurotoxicity studies including acute (28 d), sub-chronic (90 d), and chronic (≥1 year) adult neurotoxicity[26] and developmental neurotoxicity (DNT) studies[27,28]. This lack of information exists because DNT *in vivo* studies are typically conducted in rats[27,28], costly (~$1,000,000 per chemical[29]), laborious (~1 year per chemical[29,30]), and raise ethical concerns[29,31].

As untested chemicals should not be presumed safe, a new testing paradigm is needed to assess the ever-expanding universe of potentially neurotoxic chemicals[13,29]. To fill this gap, an integrated testing strategy that combines *in silico* models, *in vitro* techniques, and alternative animal models, collectively referred to as New Approach Methods (NAMs), has been proposed[29,32,33]. *In silico* methods (e.g. grouping and read-across, quantitative structure-activity relationships) can be used to predict ADME (Absorption, Distribution, Metabolism, Excretion) processes (e.g. blood-brain-barrier penetration), to model physiologically-based toxicokinetics, and to predict mechanisms based on relationships between chemical structure/properties and biological activity[34]. *In silico* approaches rely on experimental data. International efforts to enhance neurotoxicity testing involve the establishment of a DNT *in vitro* testing battery that depicts key neurodevelopmental processes including proliferation, migration, differentiation, apoptosis, neurite outgrowth,

synaptogenesis, and functional network formation in a suite of rodent- and human-based cell
models[35,36]. The underlying assumption of this approach is that the alteration of any of these
processes is indicative of DNT[35]. Despite advances in depicting the processes that orchestrate
nervous system development and function *in vitro*, more complex functional endpoints at the
organismal level, including complex behaviors, remain elusive. In particular, *in vitro* assays fail to
capture behavior endpoints assessed in guideline DNT studies including motor activity, motor and
sensory function, learning, and memory[27].

The zebrafish is a powerful vertebrate model that can bridge the gap between *in vitro* and
mammalian-based *in vivo* studies. Although zebrafish are phylogenetically more distant from
humans than rodents, 70% of human genes and 82% of disease-associated human genes have
a zebrafish orthologue[37]. Embryo-larval stages of zebrafish up to 5 d post fertilization (dpf) are
non-protected[38,39] and therefore considered to be an alternative to mammalian testing[40]. The high
reproduction rate, rapid *ex utero* development, and small size of embryo-larval stages make
zebrafish amenable to medium-to-high-throughput screens. Early life-stage zebrafish provide a
diverse repertoire of behaviors such as spontaneous tail contractions[41], touch-evoked
responses[42,43], visual and acoustic startle responses[44], phototaxis[45], learning and memory[46,47].
While multi-behavioral phenotyping in zebrafish has been widely applied in drug discovery[48,49],
molecular target identification[50,51], and to study neurological disorders[52–54], the behavior-rich
repertoire of zebrafish has yet to be fully deployed to identify and characterize chemicals that
cause neurotoxicity.

The aim of this study is to identify and mechanistically evaluate chemical compounds that cause
deficits in learning and memory. First, we developed a multi-behavioral phenotyping battery in
early-life stage zebrafish to quantify various visual and acoustic behaviors including non-
associative learning. Second, we validated the assay battery using known pharmacological
modulators of habituation learning. Third, we performed a targeted chemical screen to identify
chemical compounds that affect learning and memory in zebrafish. We then combined
pharmacological interventions and electrophysiology in mouse- and human-based models to
identify mechanisms underlying phenotypic changes and to demonstrate cross-species relevance.
Overall, the zebrafish-based multi-behavioral phenotyping approach represents an efficient tool
for the identification and mechanistic elucidation of neurotoxic environmental chemicals with
relevance across taxa, including humans.

Results

A multi-behavioral phenotyping approach to quantify visual and acoustic behaviors in larval zebrafish

Figure 1 | A battery of visual and acoustic assays provokes stereotyped behaviors in larval zebrafish. (**a**) Motor activity (y-axis) of zebrafish larvae over time and across various stimulus modalities (x-axis). Data are median ± 95% CI for n = 440 larvae obtained from 28 x 96-well plates, each containing 12-16 control individuals. (**b**) A rapid dark-to-light transition caused a visual startle response (VSR1). (**c**) Subsequent exposure to visual light caused freezing behavior in zebrafish larvae (VMR1). (**d**) Light-to-dark transition caused an additional visual startle response (VSR2). (**e**) Motor activity upon light-to-dark transition reached a maximum after approximately 4 min (VMR2) followed by an asymptotic decrease to baseline (VMR3). (**f**) A series of five low- and (**g**) high-intensity acoustic stimuli (1 s) provokes acoustic startle responses with lower or higher amplitude, respectively. (**h**) A sequence of 30 high-intensity acoustic stimuli (inter stimulus interval, ISI = 1 s) induced an exponential response decrement, indicating habituation. (**i**) Representative plot of data used for **h**, excluding ISIs. (**j**) Decrease in responsiveness during first (ASH1), relative to fifth habituation bout (ASH5), indicates potentiation of habituation. (**k**) Responsiveness to non-habituating high-intensity acoustic stimuli is significantly reduced comparing pre- (ASR2) and post-habituation (ASR3) assays, indicating memory retention. Numbers above the split rainclouds in **j** and **k** represent p-values (one-sided Wilcoxon signed rank exact test). ASH, acoustic startle habituation; ASR, acoustic startle response; BSL, baseline motor activity; VMR, visual motor response; VSR, visual startle response. N=439-440.

To identify chemicals that modulate learning and memory in larval zebrafish, we devised a battery of 11 automated, sequential behavioral assays called the Visual and Acoustic Motor Response (VAMR) assay battery. The battery included assays which quantified baseline (BSL) activity, visual

startle responses (VSR1,2), visual motor responses (VMR1-3), acoustic startle responses (ASR1,2), acoustic startle habituation (ASH1), potentiation of habituation (ASH1/ASH5), and memory retention (ASR3/ASR2) (**Fig. 1a**). Overall, four stimulus modalities were used. This included visual light at intensities of 0 lux (dark condition) and 7.5 klux (light condition). In addition, in 5 dpf larvae responding to a 300 Hz frequency, low- (65 dB) and high-volume (75 dB) acoustic stimuli were used (**Supplementary Fig. 1**) These stimuli provoked stereotyped motor activity patterns in larval zebrafish that were quantified as pixel change per second for each individual per well (**Fig. 1a**). First, in the BSL assay, zebrafish showed low average baseline activity per second. In the VSR assays, transitions from dark-to-light (**Fig. 1b**) or light-to-dark (**Fig. 1d**) conditions elicited rapid motor responses lasting 1 s. During the VMR assays, visual light caused low basal activity (**Fig. 1c**), whereas a lack of visual light gradually increased motor activity to a maximum after approximately 4 min which was followed by an asymptotic decrease to baseline (**Fig. 1e**). In the ASR1 and ASR2 assays, five low- or high-volume acoustic stimuli, with an inter-stimulus interval of 1 min, evoked weak or strong motor responses, respectively (**Fig. 1f,g**). Next, a series of 30 high-intensity acoustic stimuli with an inter-stimulus interval of 1 s induced robust habituation (**Fig. 1 h,i**). Repetitive application of habituation trials with an inter-bout interval of 1 min gradually decreased motor activity (**Fig. 1j**), indicating potentiation of habituation. Following a 3 min rest period, five high-volume acoustic stimuli elicited motor responses significantly lower than in the ASR2 assay (**Fig. 1k**), indicating memory retention. Additional data (**Supplementary Fig. 2)** indicate that the phenotyping battery fulfills hallmark criteria of non-associative habituation learning[55]. Thus, we hypothesized that the VAMR battery has the potential to identify compounds that modulate visual and acoustic behaviors in larval zebrafish, including learning and memory.

Known modulators of habituation learning alter multiple zebrafish behaviors

Figure 2 | Modulators of habituation learning cause distinct behavioral phenotypes. (**a-d**) Structures, modes of action, and habituation scores of reference modulators of habituation. Raincloud plots quantify acoustic startle habituation as habituation scores (y-axis) across indicated compounds and concentrations (x-axis). Horizontal dotted lines indicate the median habituation score of untreated larvae. Numbers above the rainclouds represent adjusted p-values (grey: $p \geq 0.05$, black: $p < 0.05$, two-sample bootstrapping test, $n_{control}$ = 24-64, $n_{treatment}$ = 24-32). Habituation scores significantly greater than negative control indicate reduced habituation whereas reduced values reflect enhanced habituation. (**e**) Hierarchical clustering of compound- and concentration-specific profiles (y-axis), across the entire assay battery (**Fig. S4**), based on effect sizes (SSMD, strictly standardized median difference) across behavioral endpoints (x-axis). Heatmap colors indicate deviation from the median control phenotype where red and blue reflect increased and reduced activity, respectively. (**f**) Multidimensional scaling (MDS) representation of pairwise distances between behavioral profiles based on data shown in **e**. Contour labels (grey) and numbers indicate clusters shown in **e**.

To evaluate the assay battery, we first tested a reference set of compounds that act through mechanisms previously reported to modulate habituation learning in zebrafish[46,47] (**Supplementary Table 3**). Selected compounds included those reported to reduce short-term habituation including two NMDAR antagonists (APV, MK-801) and donepezil (acetylcholinesterase inhibitor) and compounds that reportedly enhance short-term habituation including citalopram (selective serotonin reuptake inhibitor), haloperidol (dopamine D2 receptor antagonist), picrotoxin (GABA$_A$R antagonist), and yohimbine (α-2 adrenoreceptor antagonist).

Larval zebrafish were exposed to a series of 7-10 concentrations per compound for 1 h before quantification of motor activity. To quantify habituation behavior, the initial habituation bout (ASH1) was used to calculate a 'habituation score,' achieved by dividing the motor activity during the final ten taps by the cumulative responses during the initial and final ten acoustic stimuli. On average, the untreated vehicle control exhibited a habituation score of 0.32. Compounds that reduced short-term habituation therefore generated values greater than the control group. In contrast, compounds that accelerated the habituation response yielded values lower than the control. In line with previous reports, the prototypical NMDAR antagonists APV and MK-801 significantly reduced non-associative habituation learning in a concentration-dependent manner (**Fig. 2a,d**), whereas citalopram (21.5, 100 µM) and haloperidol (1, 4.6 µM) significantly enhanced habituation (**Fig. 2b,c**). In addition, MK-801 significantly enhanced habituation at the highest test concentration (**Fig. 2d**). Non-significant alterations in habituation were determined in zebrafish exposed to picrotoxin, donepezil, or yohimbine (**Supplementary Fig. 3**). Together, these data suggest that the VAMR assay battery can be used to identify chemicals that modulate habituation behavior in zebrafish.

To determine whether the reference compounds selectively modulated short-term habituation learning, we simplified motor activity patterns by condensing compound- and concentration-dependent responses into behavioral profiles reflecting effect size relative to vehicle control (**Fig. 2e, Supplementary Fig. 4**). Except for habituation (described above), habituation potentiation, and memory retention endpoints, the cumulative motor activity per assay for each individual was calculated. The 'potentiation score' (ASH1/5) reflects the ratio of the scaled sum of motor responses during taps in the final habituation bout (ASH5) to the sum of responses in the initial (ASH1) and final (ASH5) habituation bouts. Motor activity in the ASR3 assay, relative to an individual's overall responses summed across ASR2 and ASR3 assays, was used to quantify memory retention (ASR2/3). Subsequently, we determined effect sizes relative to vehicle control for each of the eleven endpoints, yielding compound- and concentration-specific behavioral profiles. To determine phenotypic similarities between behavioral profiles, we used two complementary approaches, hierarchical clustering (**Fig. 2e**) and multidimensional scaling (**Fig. 2f**). While the majority of compounds caused slight behavioral changes in the lower concentration range (cluster 1), exposure to higher concentrations provoked diverse multi-behavioral phenotypes (clusters 2-5) (**Fig. 2e**). Within these clusters, chemical exposure caused different combinations of behavioral alterations, such as enhanced visual and acoustic startle responses (cluster 5), hyper- and hypolocomotion in response to light or dark conditions (cluster 4), or reduced visual and acoustic startle activity (clusters 2 and 3) (**Fig. 2e**). Behavioral profiles of the

NMDAR antagonist APV (cluster 2) were segregated from all other profiles including
concentrations that significantly reduced habituation (**Fig. 2e,f**). While both NMDAR antagonists
(APV and MK-801) blocked habituation learning and reduced dark-to-light visual startle responses
in zebrafish, disparate phenotypes were also observed. Specific to APV, exposure also reduced
all other visual and acoustic startle responses (**Fig. 2e, Supplementary Fig. 4**). In contrast, MK-
801 exposure enhanced acoustic startle responses, caused light-sensitive hyperlocomotion, and
decreased responsiveness under dark conditions (**Fig. 2e, Supplementary Fig. 4**). At
concentrations that decreased habituation, behavioral profiles of MK-801 differed from APV and
clustered with profiles of picrotoxin (cluster 3), donepezil (clusters 3 and 4), and yohimbine (cluster
4) (**Fig. 2e,f**). Together with *in silico* target predictions (**Supplementary Table 4**), these data
suggest that potential off-target site(s) of MK-801 in zebrafish might involve $GABA_A$ and/or
acetylcholine receptors. In sum, while the assay battery illustrates a lack of selectivity for known
habituation modulators, it highlights the ability to identify and discriminate chemical mode of action
information.

Targeted screen for chemical compounds that affect learning and memory identifies

chlorophene

Figure 3 | Prediction and evaluation of chemical compounds that interfere with NMDAR-dependent signaling.
(**a**) Data on *ex vivo* rat glutamate receptor radioligand binding (AC$_{50}$: blue, pink), *in vitro* cytotoxicity limit (lower bound on the estimate of the cytotoxicity threshold; black), and predicted baseline toxicity in fish (LC$_{50}$; grey) were gathered from the US EPA CompTox Chemical Dashboard and ECOSAR. (**b**) Cytotoxicity- and (**c**) baseline toxicity-based rankings of 26 *ex vivo* positive chemicals (y-axis) predict 11 unique candidate compounds (black) that alter NMDAR agonist activity with higher potency relative to cytotoxicity and predicted baseline toxicity levels. (**d**) Hierarchical clustering of behavioral profiles for 10 predicted candidate compounds and two negative reference chemicals (ascorbate, cypermethrin) (y-axis) across behavioral endpoints (x-axis). Heatmap colors based on effect sizes (SSMD, strictly standardized median difference) indicate deviation from the median control phenotype. Red and blue reflect increased and reduced activity, respectively (*n* = 12-24 larvae per condition). (**e**) MDS representation of pairwise distances between behavioral profiles based on data shown in **d**. Fill color indicates deviation from the median control phenotype in the acoustic startle habituation (ASH1) assay where red and blue reflect reduced and enhanced habituation, respectively. Contour labels (grey) and numbers indicate clusters shown in **d**. (**f**) Structure of the positive hit chlorophene. (**g**) Median motor activity (y-axis) of vehicle control (0.01% DMSO, grey) and chlorophene-exposed (10 μM, red) zebrafish larvae over 30 sequential (ISI = 1 s) high-intensity acoustic stimuli (x-axis) (shaded envelope

Based on these results, we hypothesized that the assay battery could be used to identify chemical
compounds that affect learning and memory in zebrafish. Given the well-characterized
involvement of NMDARs in synaptic plasticity and memory function[56–58], we conducted a targeted
selection of candidate compounds. The EPA CompTox Chemicals Dashboard, which contains
bioactivity data on thousands of chemicals tested across numerous high-throughput screening
assays[59], was mined for relevant assays. We identified two tissue-based, cell-free radioligand
binding assays that used extracted proteins from rat forebrain membranes to determine changes
in NMDAR agonist activity (**Fig. 3a**, **Supplementary Table 5**). In total, these datasets contained
81 unique chemicals of which 26 were annotated as hits (**Supplementary Table 6**). To enrich the
set of hits for compounds that exhibited higher specificity towards NMDARs relative to non-specific
general toxicity, we used *in vitro* cytotoxicity and baseline toxicity data (**Supplementary Table 7**).
These data were used to rank order compounds according to their specificity for NMDAR
interaction relative to non-specific toxicity effects (**Fig. 3a-c**). Rankings predicted 11 compounds
with higher potency for NMDAR modulation relative to cytotoxicity and baseline toxicity levels (**Fig.**
3b,c). For example, besonprodil and MK-968, two preclinical drugs that act as NMDAR
antagonists, and methadone, a µ-opioid receptor antagonist used as a painkiller, were among the
predicted hits that appeared in both rankings (**Fig. 3b,c**). Ten compounds were purchased and
screened in our behavior assay battery. In addition, we selected two chemicals (cypermethrin,
ascorbic acid) as negative reference compounds for NMDAR modulation. Cypermethrin is a
pyrethroid insecticide that prolongs the opening of sodium channels[60]. From a structural
perspective, it is closely related to deltamethrin, a developmental neurotoxicant[61]. In contrast,
ascorbic acid was proposed as a negative chemical for developmental neurotoxicity[25].

To determine whether exposure to the selected compounds affected behavior, we treated
zebrafish for 1 h with multiple concentrations of each test chemical and translated the 11
quantitative features into behavioral profiles (**Supplementary Figure 5**). Hierarchical clustering
segregated compound- and concentration-dependent profiles into five main clusters (**Fig. 3d**).
Cluster 1 was characterized by decreased activity in multiple visual and acoustic startle response
assays and contained compounds such as chlorpromazine (neuroepileptic) and spironolactone
(diuretic) (**Fig. 3d**). A second cluster (cluster 2), characterized by enhanced responsiveness to
high-volume acoustic stimuli, included diphenhydramine (antihistamine) and methadone (**Fig. 3d**).
Cluster 3 exclusively contained cypermethrin which provoked light-sensitive hyperlocomotion in a

concentration-dependent manner and decreased motor activity across multiple visual- and

acoustic-based endpoints (**Fig. 3d**). To visualize chemical modulators of habituation learning

among the variety of phenotypes, we projected all multi-behavioral profiles into two dimensions

using multidimensional scaling and shaded each data point with the effect size determined in the

ASH1 assay (**Fig. 3e**). We found that behavioral profiles with enhanced or reduced startle

habituation spread across the two-dimensional space, reflecting diverse multi-behavioral

phenotypes (**Fig. 3e**). Diphenhydramine (1, 4.6-100 μM), the structurally related fungicides

mancozeb (0.1-4.6 μM) and maneb (0.1-21.5 μM), and spironolactone (46.4 μM) significantly

enhanced acoustic startle habituation (**Fig. 3e**, **Supplementary Fig. 6**, $p < 0.05$, two-sample

bootstrapping test). Three compounds including besonprodil (100 μM, cluster 2), cypermethrin

(10 μM, cluster 3), and chlorophene (10 μM, cluster 4), reduced acoustic startle habituation (**Fig.**

3d), although only the habituation decrease caused by chlorophene was statistically significant

(**Fig. 3f-h**, **Supplementary Fig. 6**, $p = 0.028$, two-sample bootstrapping test). Overall, half (5/10)

of the predicted hits were active in our ASH1 assay. Surprisingly, while one of the negative

reference chemicals, cypermethrin, was indeed inactive, ascorbate (10-21.5 μM, 100 μM)

significantly enhanced acoustic startle habituation (**Supplementary Fig. 6**). Together, these data

suggest that the VAMR behavior assay battery is useful for identifying compounds that also

modulate NMDAR-mediated signaling in rat *ex vivo* preparations. In addition, relative to *in vitro* or

ex vivo biochemical assays, and based on the diversity of potential phenotypic outcomes, the

zebrafish assay has the potential to elucidate additional, previously unknown phenotypes that can

be mechanistically evaluated.

Chlorophene does not directly act on NMDARs and alters neuronal firing

Figure 4 | Chlorophene does not act on NMDARs in cultured mouse cortical neurons but reduces the weighted mean firing rate of glutamatergic units in human stem cell-derived 3D BrainSpheres. (**a**) Representative infrared differential interference contrast image of a patch-clamped cultured mouse cortical neuron. (**b**) Representative current response to NMDA (1 µM) remained unchanged during chlorophene (10 µM) application, but was inhibited by the NMDAR antagonist APV (50 µM). (**c**) Individual holding currents (I_{hold}) before (ctr) and during NMDA administration were not affected by chlorophene (+chloro, $n = 8$), but inhibited by APV ($n = 4$). (**d**) Relative I_{hold} during chlorophene (n=8) and APV (n=4) application compared to I_{hold} evoked by respective NMDA exposure. (**e**) Weighted mean firing rate (y-axis) of human stem cell-derived 3D-differentiated BrainSpheres exposed to increasing concentrations of chlorophene (x-axis). After the first baseline recording, the neurotransmitter glutamate, followed by the antagonists APV/NBQX, were applied to identify glutamatergic units (two-sample bootstrapping test, $n = 9$).

Given its wide use as a biocide, preservative, and disinfectant[62,63], we investigated chlorophene in more detail. Because chlorophene exposure impaired habituation in zebrafish (**Fig. 3g,h**), altered NMDAR agonist activity in rat *ex vivo* preparations (**Fig. 3b**), and given the crucial role of NMDARs in neuroplasticity, we hypothesized that chlorophene reduced habituation learning by targeting NMDARs. We further hypothesized that the underlying molecular mechanisms may also be conserved across taxa, including mice and humans. To evaluate the translational relevance of these findings, we used whole-cell patch-clamp recordings in cultured mouse cortical neurons (**Fig. 4a**). We found that, while the NMDAR antagonist APV effectively inhibited the current response provoked by NMDA application, chlorophene application failed to do so (**Fig. 4b-d**). As a second approach to determine whether chlorophene targeted glutamatergic signaling, the compound was evaluated in a microelectrode array (MEA) assay using human-induced pluripotent stem cell-derived (hiPSC) 3D BrainSpheres. In this *in vitro* model, chlorophene exposure reduced

307 the weighted mean firing rate of glutamatergic units in a concentration-dependent manner (**Fig.**
308 **4e**). Together, although chlorophene may interact with glutamatergic signaling, these data do not
309 support the hypothesis that chlorophene directly antagonizes NMDARs leading to reduced
310 habituation in zebrafish.

Chlorophene acts on GABAARs in zebrafish, mouse, and human models

Figure 5 | Chlorophene affects GABAergic signaling in zebrafish, cultured mouse cortical neurons, and human
stem cell-derived 3D BrainSpheres. (**a**) Motor activity (y-axis) of zebrafish treated with chlorophene (10 µM, red) or
0.01% DMSO (grey) over time and across various stimulus modalities (x-axis). Data represents median ± 95% CI. (**b**)
Split raincloud plots show the effect of chlorophene exposure on zebrafish motor activity (y-axis) across 11 assay
endpoints (x-axis). Motor activity is expressed as effect size (SSMD, strictly standardized median difference) to
represent different assay endpoints on a common scale. Data points are equally shifted to set the median control activity
per assay endpoint to zero. Numbers above the rainclouds represent adjusted p-values (grey: p ≥ 0.05, black: p < 0.05,
two-sample bootstrapping test). (**c**) Representative raincloud plot showing the cumulative motor activity (y-axis) of
zebrafish exposed to the indicated compounds (x-axis) in response to dark conditions (VMR3) (Kruskal-Wallis test, n =
94-96 larvae). (**d**) Habituation scores (y-axis) of zebrafish treated with the indicated compounds (x-axis). A sequence of
30 high-intensity acoustic stimuli with 1 s ISI was used to induce acoustic startle habituation (ASH1) (Kruskal-Wallis
test, n = 94-96 larvae). (**e**) Representative response of a cultured mouse cortical neuron to depolarizing current
injections under the control condition (black) and (**f**) during chlorophene application (10 µM, red). (**g-j**) Passive

membrane and intrinsic properties of cultured mouse cortical neurons without and during application of chlorophene
including (**g**) rheobase, (**h**) action potential (AP) firing frequency at 2x rheobase, (**i**) input resistance (R_{input}), and (**j**)
resting membrane potential (V_{rest}) (paired t-tests, n=7). (**k**) Representative current response of a cultured mouse cortical
neuron to chlorophene (10 µM) was inhibited by co-exposure to the $GABA_A$ receptor antagonists picrotoxin (Ptx, 50 µM)
and SR95531 (SR, 10 µM). (**l**) Individual and median holding current (I_{hold}) under control condition, during chlorophene
application and $GABA_A$ receptor blockade by picrotoxin and SR95531 (repeated measures ANOVA, Bonferroni t-test, *n*
= 6). (**m**) Weighted mean firing rate (y-axis) of 3D BrainSpheres treated with increasing concentrations of chlorophene
(x-axis). After the first baseline recording, the neurotransmitter GABA followed by the antagonist bicuculline were applied
to identify GABAergic units (two-sample bootstrapping test, *n* = 46).

To determine whether chlorophene acted on targets other than NMDARs, we used the Similarity Ensemble Approach[64] to predict protein targets based on structural similarity to their ligands. Among 107 predicted targets were three $GABA_A$R subunits including α2, α3, and α5 (**Supplementary Table 8**). Certain agonists and positive allosteric modulators of $GABA_A$Rs have been reported to cause sedation and paradoxical excitation in zebrafish[51], two phenotypes observed in chlorophene-exposed larvae (**Fig. 3g,h and Fig. 5a,b**). We therefore hypothesized that sedation in visually-mediated responses and increased responsiveness to repeated acoustic stimuli, were related to a GABAergic mode of action in chlorophene-exposed zebrafish. To determine whether chlorophene exposure caused these phenomena via $GABA_A$R agonism, we pretreated zebrafish with the $GABA_A$R antagonist picrotoxin (**Fig. 5c**). Pharmacological intervention rescued chlorophene-induced sedation, restoring visual motor responses (**Fig. 5c**, **Supplementary Fig. 7**) whereas picrotoxin pretreatment did not counteract the habituation deficit caused by chlorophene exposure (**Fig. 5d**). Together, these observations suggest that exposure to chlorophene caused sedation and paradoxical excitation, but not defective habituation, through $GABA_A$Rs in zebrafish.

We next sought to evaluate whether chlorophene acts as a $GABA_A$R agonist in mouse- and human-based models. To determine whether chlorophene acts on $GABA_A$ receptors we first measured currents in cultured mouse cortical neurons. Chlorophene exposure induced a slow tonic chloride current that was blocked by the $GABA_A$R antagonists picrotoxin and SR95531 (**Fig. 5k,l**). This suggests that chlorophene activates rodent $GABA_A$Rs. To investigate the effect of $GABA_A$R activation by chlorophene on neuronal action potential firing, we measured the membrane potential during current injections while blocking glutamatergic receptors with APV and NBQX (**Fig. 5e-f**). Chlorophene exposure increased the minimum current needed to generate an action potential (i.e. the rheobase; **Fig. 5g**), decreased the action potential firing frequency and the input resistance (**Fig. 5h and i**), and did not change the resting membrane potential (**Fig. 5j**). These data suggest that chlorophene-induced $GABA_A$R-mediated chloride conductance shunts

neuronal action potential spiking by lowering the input resistance and increasing the current

threshold required for neuronal spiking. To determine whether chlorophene acts as a GABA$_A$R

modulator in humans, we used the 3D BrainSphere MEA assay where the weighted mean firing

rate of GABAergic units was significantly decreased in chlorophene-treated (6.7-20 µM) 3D

BrainSpheres (**Fig. 5m**). In summary, these data indicate that chlorophene likely acts through

similar mechanisms in zebrafish, mouse, and human-based models to agonize GABA$_A$Rs.

Discussion

Given the ubiquitous exposure to diverse environmental chemicals across lifespan and the vast dearth of information on the hazards presented by these chemical compounds, there is an unmet need to efficiently assess their potential adverse effects on neurodevelopment and function. Here we describe a multi-behavioral phenotyping approach in early life-stage zebrafish that enables (i) the rapid identification of neuroactive compounds and (ii) the elucidation of mechanisms underlying alterations in neurobehavior. While current behavior-based toxicity tests in larval zebrafish, such as the light-dark transition test[65–68], largely focus on a small number of visual motor behaviors, we merged this widely used assay with additional, well-described paradigms, broadly applied in large-scale behavioral drug screens, including acoustic startle responses[48,51], habituation learning[46,47,49], and other memory-related endpoints that have yet to be applied to toxicity testing.

This work focused on the identification of chemical modulators of habituation, which is a form of non-associative learning[55]. Defective habituation is associated with decrements in learning and memory[69,70] and is prevalent in neuropsychiatric disorders such as autism spectrum disorder, schizophrenia, and attention deficit hyperactivity disorder[71]. Here, we demonstrated that the phenotyping battery fulfills hallmark criteria of non-associative learning[55], a phenomenon conserved across phyla including zebrafish[47], rodents[72], and humans[73]. Confirmation of habituation characteristics (**Supplementary Fig. 2**) and the reproducibility of phenotypes observed with known habituation modulators demonstrate that the VAMR battery quantifies non-associative learning. This includes a quantitative demonstration of dishabituation, which is defined as an increased response to a habituated stimulus following a second, brief stimulus using a different stimulus modality[55]. Dishabituation to acoustic stimuli has been quantitatively demonstrated 7 dpf zebrafish using a second, light-based stimulus[47] while in 6 dpf larvae, dishabituation following a user-based tactile stimulus has been reported[46,74]. This study demonstrates dishabituation in 5 dpf zebrafish, using a light-based secondary stimulus and thereby avoiding the use of a tactile stimulus which could cause confounding effects on swimming behavior (**Supplementary Fig. 2**).

In line with previous studies, we found that acute administration of the prototypical NMDAR antagonists APV or MK-801 reduced acoustic startle habituation in zebrafish[46,75]. This observation is consistent with deficits in learning and memory reported in rodent-based test paradigms[76,77]. We found a lack of selectivity for habituation with different chemical modulators. This may result from off-target activities or systemic compound interactions across multiple molecular targets that orchestrate behavior. The mode(s) of action underlying behavioral alterations in zebrafish

exposed to MK-801 for example, are not completely understood. *grin1* encodes the obligatory

GluN1 subunit of NMDARs[78]. While behavioral defects in *grin1* double mutants are generally

phenocopied by short-term MK-801 treatment[79], off-target activities on acetylcholine receptors[80]

and voltage-gated potassium channels[81] may explain the difference in phenotypes observed in

zebrafish exposed to MK-801 and APV. Overall, our data indicate that zebrafish multi-behavioral

phenotyping strategies can be used to identify complex, system-modulating compounds.

We found that exposure to chlorophene, a chlorinated phenolic compound, caused reduced

habituation learning, sedation, and paradoxical excitation in zebrafish. Sedatives and general

anesthetics (e.g. benzodiazepines, propofol, etomidate) can cause paradoxical excitation where

an opposite, paradoxical reaction induces both sedation and increased brain and/or motor

activity[82–84]. It is widely accepted that anesthetics induce neuronal suppression via GABAR

agonism, NMDAR antagonism, or by membrane polarity modulation[85–89]. Several lines of evidence

indicate that chlorophene also acts on $GABA_ARs$. First, among other proteins, SEA algorithms[64]

predicted multiple $GABA_AR$ subunits to be targeted chlorophene, including subunits α2, α3, and

α5. Second, chemical intervention using the $GABA_AR$ antagonist picrotoxin counteracted sedation

of visual motor behaviors in zebrafish. Third, tonic inhibition against neuronal excitability in

cultured mouse cortical neurons was blocked by the $GABA_AR$ antagonists picrotoxin and

SR95531. Fourth, chlorophene exposure reduced the weighted mean firing rate of GABAergic

units in hiPSC-derived 3D BrainSpheres.

Most $GABA_AR$ α subunits are expressed during zebrafish embryogenesis[90]. The amino acid

sequences and electrophysiological properties of $GABA_AR$ subunits are largely conserved among

vertebrates including zebrafish, mice, and humans[91]. Several studies suggest that α5-$GABA_AR$-

mediated tonic inhibition reduces learning and memory[92–98]. Under physiological conditions, tonic

$GABA_AR$ currents cause hyperpolarization and decrease neuronal excitability. Synaptic input,

therefore, becomes subthreshold and interferes with coordinated pre- and post-synaptic spiking

activity, which is required for NMDA-dependent long-term plasticity[99]. Antagonizing GABAergic

tone facilitates neuroplasticity in genetic mouse models for cognitive disabilities[100,101]. In contrast,

the deficits in learning and memory in zebrafish described here were not rescued by the $GABA_AR$

antagonist picrotoxin. Our results indicate that reduced habituation may result from a mechanism

distinct from functional $GABA_AR$ agonism. Alternatively, co-occurring phenotypes including

sedation, paradoxical excitation, and defective learning, may causally relate to a common,

unknown upstream regulator.

NMDARs are essential to mediate experience-dependent neuroplasticity underlying learning and memory[58,78]. In teleost fish, including zebrafish, the telencephalon, and in particular the medial and lateral pallium, are analogous to amygdala and hippocampus in mammals[102–104] and represent the neuroanatomical structures responsible for learning and memory involving NMDAR activation[105–107]. However, our electrophysiological data indicates that chlorophene does not directly interfere with NMDAR-dependent signaling in mouse cortical neurons. This suggests that chlorophene indirectly affects NMDARs, phenocopying the decreased habituation observed in zebrafish exposed to NAMDAR antagonists. A large-scale screen for chemical modulators of non-associative learning identified additional mechanisms reducing short-term habituation in larval zebrafish including potassium channel antagonism[46]. Interestingly, another zebrafish-based small molecule screen implicated that the M-current, a voltage-gated potassium channel that regulates neuronal excitability[108], is related to sedation and paradoxical excitation[51]. Our screen confirmed that exposure to diphenhydramine, an antihistamine used in some hypnotic preparations, caused sedation and paradoxical excitation in early life stage zebrafish[109]. Surprisingly, as compared to chlorophene, diphenhydramine exposure had the opposite effect on non-associative learning, enhancing short-term habituation in zebrafish. Future studies should investigate whether M-current modulators counteract the sedative, paradoxical excitatory, and habituative learning effects of compounds such as chlorophene and diphenhydramine.

Chlorophene and its structural analogues, known as chlorophenes, represent a group of halogenated phenolic compounds that are globally used as antimicrobial agents and preservatives in personal care products and cosmetics[63,110]. Their environmental occurrence in surface waters and soils exclusively results from anthropogenic activities through emissions from sewage treatment plants and application of sewage sludge[110–112]. Based on a predicted environmental concentration (PEC) of 0.2 µg/L (0.9 nM) and a predicted no effect concentration (PNEC) of 0.058 µg/L (0.3 nM), a risk quotient (RQ=PEC/PNEC) of 3.4 has been calculated for the emission of chlorophene into surface waters, indicating that chlorophene may pose an unreasonable risk to the aquatic environment[110,113]. Consequently, chlorophene is included on international lists of environmental chemicals of emerging concern[110]. This, paired with the toxicity data reported here, suggests that (bio)monitoring in human and environmental matrices is justified.

Chlorophenes are hydrophobic (log K_{OW} 4.3-7.5), persistent (ultimate biodegradation half-life 1.9-4.9 years), low-to-moderately bioaccumulative in fish (BCF 55-400 L/kg), and toxic to aquatic organisms including fish, invertebrates, and algae[110,113]. In particular, chlorophene's 30-day no observed effect concentration (NOEC$_{chronic}$) for survival in zebrafish (Fish Early Life Stage Test, OECD TG 210[114]) of 0.58 mg/L (2.7 nM) indicates high chronic toxicity (acute-to-chronic ratio;

ACR=LC50$_{acute}$/NOEC$_{chronic}$=569) and is consistent with a specific mode of action[62,110,113,115]. Our
study suggests that chlorophene acts as a functional agonist at GABA$_A$Rs, indicating a neurotoxic
mode of action. Currently, adult or developmental neurotoxicity assessment is addressed under
the Classification, Labeling and Packaging (CLP) Regulation[116,117], REACH (Registration,
Evaluation, Authorisation and Restriction of Chemicals), and the Biocidal Products Regulation
(BPR). Under CLP, developmental neurotoxicity studies are required if there is evidence that the
nervous system is a target organ including any effect interfering with normal development until
sexual maturation, including death, structural abnormality, altered growth, and functional
deficits[118], whereas both under REACH and BPR, adult neurotoxicity (functional or morphological
alterations in the mature nervous system) triggers tests on developmental neurotoxicity[119,120].
Interestingly, an ECHA (European Chemicals Agency) assessment reports state that chlorophene
has "…no structural similarity to […] known inducers of delayed neurotoxicity" and that "[…] studies
in several species did not reveal the potential for neurotoxic effects […]"[62]. This discrepancy
underscores limitations in current neurotoxicity assessment practices as this conclusion was
reached without conducting regulatory adult or developmental neurotoxicity tests.

In addition to neurotoxicity, there is increasing evidence that chlorophene and related chemical
compounds act as endocrine-disrupting chemicals (EDCs) that interfere with hormonal
homeostasis. In cell-based luciferase reporter assays *in vitro*, chlorophene modulated androgen,
estrogen, and thyroid receptor activity, and meets the criteria for an EDC[121]. A structurally related
compound, bromochlorophene, was shown to act as an antagonist on androgen, estrogen,
glucocorticoid, and thyroid receptors *in vitro*[121]. Given that GABA is widely distributed in endocrine
tissues and serves as a hormone or trophic factor in neuroendocrine systems[122,123], and
considering that changes in hypothalamic GABA levels are involved in endocrine disorders[124,125],
a potential link between altered GABAergic signaling and endocrine disruption should be
addressed in EDC-related studies.

In the future, zebrafish multi-behavioral assays might be utilized for the regulatory assessment of
chemical-induced neurotoxicity. The European Union's chemicals strategy (European Green
Deal) aims to enhance innovation for safe and sustainable chemicals to better protect human
health and the environment[126]. To improve public health and environmental protection under
legislative instruments such as REACH and CLP, current risk assessment should prevent the use
of chemicals that affect neurological systems[126]. Because current regulatory DNT testing is
exclusively conducted in rodents (e.g., OECD Test No. 426: Developmental Neurotoxicity
Study[27]), less than 0.06% of an estimated 350,000 commercially available chemicals[2] have been
evaluated for potential effects on the developing nervous system[25]. As a result, there is an urgent

need for alternative, rapid, inexpensive, and 3R-compliant[38] approaches to evaluate potential DNT effects. Compared to cell-based *in vitro* neurotoxicity methods[127,128], multi-behavioral phenotyping in zebrafish represents a more holistic and untargeted approach to identify neurotoxicants and to infer their potential chemical mode(s) of action. In addition, unlike *in vitro*-based approaches, early life-stage zebrafish capture the functional complexity of the vertebrate nervous system[129], including its conserved developmental biology that is relevant to higher vertebrates including humans[130]. In addition, zebrafish are metabolically competent, enabling the identification and characterization of adverse effects derived from biotransformation products[131,132]. While the zebrafish phenotyping battery contains multiple learning- and memory-related readouts including habituation, potentiation of habituation[46], and memory retention[46,75] that serve as potential correlates to behavior-based endpoints in OECD rodent neurotoxicity test guidelines[27], more work is needed to understand the direct relationship between rodent- and zebrafish-based behavior readouts.

In summary, we combined a series of behavior assays in a non-mammalian vertebrate model to screen chemicals for neurotoxicity in a phenotypically rich, rapid, and inexpensive test system. Together with *in silico* target predictions, and mouse- and human-based models, our findings establish multi-behavioral phenotyping as a powerful toolkit for neurotoxicity testing and mechanism identification, with cross-species relevance. More broadly, multi-behavioral phenotyping in zebrafish can accelerate the identification of neuroactive chemicals that should be prioritized for additional testing to better protect human health and the environment from chemicals that disrupt nervous system function.

Methods

Zebrafish maintenance and embryo collection

All protocols and procedures involving zebrafish were conducted according to German and European animal protection standards and were approved by the Government of Saxony (Landesdirektion Leipzig, file number 75-9185.64). Adult zebrafish (*Danio rerio*), derived from an outcross of WIK and an in-house 'UFZ-OBI' strain, were maintained on a 14-h light, 10-h dark cycle at 28 °C and fed twice a day during the week and once daily on weekends with live brine shrimp and/or SDS 400 dry feed (Special Diets Services). Fertilized eggs were collected from group breeds containing 120-140 fish/120 L water and were bleached for 5 min with 0.05% NaOCl at 3-4 h post-fertilization (hpf). Embryos were raised in crystallization dishes at a density of 50 embryos per 100 mL standard dilution water (ISO 7346-3; 80 mM $CaCl_2 \cdot 2H_2O$, 20 mM $MgSO_4 \cdot 7H_2O$, 31 mM $NaHCO_3$, 3.1 mM KCl, pH 7.4) until 4 dpf. Viable, hatched, and

morphologically normal larvae with an inflated posterior swim bladder were individually distributed
into single wells of clear polystyrene flat-bottom 96-square-well plates (Cytiva, Marlborough, MA,
USA) filled with 200 μL standard dilution water. Plates were covered with microseal 'A' film (Bio-
Rad Laboratories, Hercules, CA, USA) and microplate lids (Cytiva, Marlborough, MA, USA). Plates
were stored at 28 °C on a 14-h light to 10-h dark cycle prior chemical exposure at 5 dpf.

Zebrafish chemical exposure
Chemical compounds are listed in **Supplementary Table 1**. Pharmacological modulators of
habituation learning, selected based on previous studies in larval zebrafish (**Supplementary**
Table 3), were dissolved in standard dilution water. Environmentally-relevant chemical
compounds, selected from *ex vivo* studies, were dissolved in dimethyl sulfoxide (DMSO, Merck,
1.02952.1000, ≥99.9% purity). 100 mM chemical stock aliquots, prepared in DMSO, were stored
at -80 °C. Serial dilutions were prepared from 200 μM chemical stock solutions adjusted to pH 7.4.
At 5 dpf, larvae were exposed to 200 μL of 2✕ exposure solutions for a final volume of 400 μL per
well. Each compound was tested at 7-10 concentrations that ranged from 0.1-100 μM, unless
otherwise indicated. When DMSO was used, the control group was exposed to an equal volume
for a final concentration of 0.01% DMSO. Compounds were tested in 8-24 replicate wells across
2-4 replicate plates using larvae from two independent spawning events. To detect acute,
receptor-mediated effects, larvae were exposed for 40 min (28 °C, light on) prior to behavior
analysis. For behavior phenotype rescue experiments, larvae in 200 μL standard dilution water
were pre-treated with 100 μL 3✕ picrotoxin solution at a concentration of 64.6 μM (21.5 μM final
concentration) or vehicle for 15 min, followed by addition of either 100 μL 4✕ chlorophene solution
at 27.3 μM (6.8 μM final concentration), 21.5 μM picrotoxin, a mixture of chlorophene and
picrotoxin (6.8 μM chlorophene, 21.5 μM picrotoxin final concentration), or vehicle. The final
concentration of DMSO was 0.01% for all groups. Plates were incubated for an additional 25 min
before behavior acquisition.

Zebrafish automated behavior assay battery and motor activity quantification
96-square well plates containing a single 5 dpf larva per well were loaded into a ZebraBox
behavior apparatus (ViewPoint) and the measurement was initiated. The assay commenced with
an initial 21 min acclimation period (1 min 100% light, 20 min 0% light). The temperature was
maintained at 28°C for the duration of the experiment. Videos were recorded at 25 frames per
second and ZebraLab video tracking software (ViewPoint) in 'quantization mode' (sensitivity
threshold = 15) to quantify locomotor activity using pixel intensity changes per second for each of
the 96 evenly-spaced, well-specific regions. In a battery of 11 behavioral assays, sequential

variations in stimulus modalities (**Supplementary Table 2**) were applied to stimulate changes in
locomotor activity, that were optimized considering visual (light- and dark-cycle durations) and
acoustic stimulus parameters (sound frequency, volume, inter-stimulus interval (ISI), inter-bout
interval), and developmental stage (**Supplementary Fig. 1** and **2**). Infrared light was used to
continuously illuminate the measuring chamber and visual light at 100% nominal intensity was
determined to be 7.5 klux (LI-250 light meter, LI-COR). Acoustic stimuli with a frequency of 300 Hz
were applied at nominal volumes of 75% (65 dB for ASR1) and 100% (75 dB for ASR2, ASR3,
ASH1-5).

Motor activity data (Δpixel/second/well) was analyzed using custom KNIME (version 4.1.4, KNIME
AG, www.knime.com) workflows implemented with R scripts[133] for statistical analysis and data
visualization using the add-on packages ggplot2[134], RColorBrewer[135], and ComplexHeatmap[136].
To quantify motor activity, cumulative pixel intensity changes per second were calculated per well
for each assay. The average motor activity per second during the final minute of the acclimation
period was used to quantify baseline activity (assay BSL). Motor activity following dark-to-light
transitions was divided into two periods reflecting rapid visual startle responses lasting 1 s (assay
VSR1) and low basal activity in the following 9:59 min (assay VMR1). Subsequent changes in
motor activity following light-to-dark transitions were binned into three periods including a
one-second visual startle response (assay VSR2), a gradual increase in motor activity in the
following 3:59 min (assay VMR2), and a transient decrease in motor activity to baseline level within
the subsequent 16 min of the dark period (assay VMR3). Then, acoustic startle responses to five
low-intensity (assay ASR1) and high-intensity (assay ASR2) 1-s acoustic stimuli, separated by 1-
min ISIs, were averaged across five trials. Motor responses to 30 high-intensity acoustic stimuli
(1-s with 1-s ISI) were condensed into a single metric reflecting acoustic startle habituation (assay
ASH1) comprising the cumulative motor activity during the final ten taps scaled to the overall
activity during the initial and final ten taps to account for individual differences in baseline startle
activity[137]. Out of five repetitive habituation bouts with an inter-bout interval of 1 min, the ratio of
motor activity during taps in the final bout (ASH5) relative to the sum of activities in the initial
(ASH1) and final bout (ASH5) was calculated to quantify potentiation of habituation (ASH1/5).
Habituation trials were followed by a three-minute rest period. Then, five high-intensity acoustic
stimuli (1-s, 1-min ISI; ASR3) that were averaged across trials and scaled to the overall responses
summed across ASR2 and ASR3 assays to quantify memory retention (ASR2/3). Following each
experiment, dead larvae, identified upon visual inspection, were excluded from behavior data
analyses.

Statistics, behavioral profiling, hierarchical clustering, and multidimensional scaling for zebrafish behavior data

Statistical analyses were performed on 11 behavior metrics including baseline activity (BSL), visual startle responses (VSR1 and VSR2), visual motor responses (VMR1-VMR3), acoustic startle responses (ASR1 and ASR2), acoustic startle habituation (ASH1), potentiation of habituation (ASH1/5), and memory retention (ASR2/3). Given the diverse, mostly nonparametric probability distributions of data across different behavior metrics, we chose a universally applicable, nonparametric bootstrapping method that is not limited by a model assumption[138]. First, the absolute value (two-sided test) of the difference in medians between untreated control and each treatment was calculated as a test statistic. Then, a *N* (number of observations to sample) times *B* (number of bootstrap samples; set to 10,000) matrix was created, where each column represented a bootstrap resample (with replacement) of the observed data. Third, the bootstrap test statistic (absolute value of the difference in medians) was calculated for each bootstrap resample per column, considering the number of observations per group. Finally, the p-value was calculated reflecting the proportion of bootstrap test statistics that were equal or greater than the observed test statistic[139,140]. For multiple comparisons, the Kruskal-Wallis rank sum test was used and the R package "agricolae"[141] was used to report results of all pairwise comparisons using compact letter displays. A Benjamini-Hochberg correction was used to adjust for multiple comparisons in order to control for family-wise Type I error rates. Results were considered significant if $p_{adjusted}<0.05$. To calculate 95% confidence intervals of the median for each one-second time bin, 1,000 bootstraps (with replacement) were sampled from the observed data and the percentile method was used to determine the lower (2.5th percentile) and upper (97.5th percentile) confidence interval bounds[142]. As a measure of effect size, strictly standardized median difference (SSMD) values were calculated as follows: $SSMD=\tilde{x}_{treatment}-\tilde{x}_{control}/(MAD^2_{treatment} + MAD^2_{control})^{1/2}$, where $\tilde{x}$ and MAD represent median and median absolute deviation[143]. SSMD calculations were performed on each of the 11 behavior metrics, collectively representing a compound's behavioral profile at a given concentration. Hierarchical clustering was performed on behavioral profiles using k-means clustering in the R package ComplexHeatmap[136]. For multidimensional scaling (MDS), two-dimensional coordinates for each behavioral profile were calculated using the KNIME nodes 'Numeric Distances' (Euclidean distance configuration), 'Distance Matrix Calculate', and 'MDS (DistMatrix)'.

Identification of candidate compounds for zebrafish behavior-based screening

The U.S. Environmental Protection Agency (US EPA) CompTox Chemicals Dashboard (https://comptox.epa.gov/dashboard/) was searched for assay endpoint names and genes using

the search term "NMDA". Two assays were identified, including "NVS_LGIC_rGluNMDA_Agonist"
and "NVS_LGIC_rGluNMDA_MK801_Agonist", both representing tissue-based, cell-free
radioligand binding assays on extracted gene-proteins from rat forebrain membranes relating to
the gene *grin1*. Detailed assay information can be found in **Supplementary Table 5**. Among 81
unique chemical compounds present in at least one of the two datasets, 26 compounds were
annotated as hits (**Supplementary Table 6**). Positive hits were included in the zebrafish behavior
screen if their AC_{50} (active concentration causing 50% loss in scintillation count) values were
below the lower bound of cytotoxicity[144] or the predicted 96-h LC_{50} in fish (ECOSAR v2.0, U.S.
EPA) (**Supplementary Table 7**). Additionally, two compounds annotated as inactive, including
cypermethrin and ascorbate, were selected as negative reference compounds. Chemical
exposures were performed as described above.

Molecular target prediction

The Similarity Ensemble Approach (SEA[64]) was used to predict molecular target sites for MK-801
(**Supplementary Table 4**) and chlorophene (**Supplementary Table 8**).

Mouse cortical cultures

Neocortical neuronal cultures from P0 to P1 mice were prepared as previously described[145].
Briefly, mice were decapitated and cerebral cortices were removed, dissected and enzymatically
digested with papain (Sigma, CAS 9001-73-4) or trypsin (Sigma, CAS 9007-07-7) in the presence
of DNAse (Sigma, CAS 9003-98-9), followed by mechanical dissociation and centrifugation
through a cushion of 4% bovine serum albumin (Sigma, CAS 9048-46-8). These steps were
completed using Hibernate medium (ThermoFisher). Cells were then plated onto Poly-L-Lysine
(Sigma, CAS 9001-73-4) coated coverslips in 24 well plates. For each coverslip, 25-30 k cells
were allowed to settle in a 40 µL drop for approximately 30 min and then each well was filled with
500 µL of NeurobasalA/B27 growth medium (Invitrogen) supplemented with GlutaMax (0.25%,
Invitrogen), glutamine (0.25-0.5 mM, Sigma), penicillin/streptomycin (1:100, ThermoFisher), and
heat-inactivated fetal calf serum (10%, Sigma, CAS 1943609-65-1). Media was partially
exchanged on day 3 (800 µL) and day 14 (500 µL) with fresh maintenance medium comprised of
BrainPhys (StemCell), B27 (2%, Invitrogen), GlutaMax (0.25%, Invitrogen), and
penicillin/streptomycin (1%, ThermoFisher). Cultures were maintained for up to 2-3 weeks at 37°C
and 5% CO_2 until use.

Electrophysiology

Postsynaptic patch-clamp recordings were performed at mouse cortical neurons [days *in vitro*
(DIV) 15-16] using a HEKA EPC10 amplifier (HEKA Elektronik). Pipette solution for voltage-clamp

recordings contained 120 mM CsCl, 20 mM TEA-Cl, 10 mM HEPES, 5 mM EGTA, 3 mM Mg-ATP,

0.3 mM Na-GTP, 5 mM Na-Phosphocreatin, and 3 mM QX314-Cl. The pH was adjusted with

CsOH to 7.31 and osmolarity was adjusted with sucrose to 298 mOsm. Series resistance (Rs)

was, on average, 9.5 ± 2.3 MΩ and was compensated to remaining Rs of 5.1 ± 0.8 MΩ. Pipettes

were pulled from borosilicate glass (Science Products) with a DMZ Universal Electrode (Zeitz

Instruments) with a resistance of 3-4 MΩ. Recordings were performed at room temperature.

Postsynaptic currents and potentials were analyzed with the NeuroMatic plug-in (Version 3) for

Igor Pro (WaveMetrics, Lake Oswego, OR, USA; Version 9). Obtained parameters were presented

and statistically compared with Sigma Plot 11 (Systat Software).

Recordings of the holding current (I_{hold}) and spontaneous inhibitory postsynaptic currents (sIPSCs)

were performed at a holding potential of -60 mV in extracellular Tyrode's salt solution containing

145 mM NaCl, 2.5 mM KCl, 1.2 mM $MgCl_2$, 2 mM $CaCl_2$, 10 mM HEPES, and 10 mM glucose.

The pH was adjusted by NaOH to 7.4. To isolate $GABA_A$R-mediated currents, the extracellular

solution was supplemented by 10 µM NBQX (Biotechne, CAS 479347-86-9) to block AMPA

receptors and 50 µM APV (Tocris, CAS 79055-68-8) to block NMDA receptors. Voltages were not

corrected for a calculated liquid junction potential of 2 mV. After establishing a baseline recording

period (before condition ~3 min), 10 µM chlorophene was washed in through the perfusion system

and I_{hold} was monitored. After four min, 50 µM picrotoxin (Sigma, CAS 124-87-8) and 20 µM

SR95531 (Sigma, CAS 104104-50-9) were bath-applied to block $GABA_A$R-mediated chloride

conductance. To compare I_{hold} before and during the application of chlorophene and after

application of the $GABA_A$R blockers, the mean I_{hold} was calculated for (i) the before period (I_{hold}

ctr), (ii) 2 min after application of chlorophene (I_{hold} +chloro), and (iii) 1 min after application of

picrotoxin and SR95531 (I_{hold} +PTX/SR) for each cell. The main effect of drug perfusion was tested

with repeated measures ANOVA followed by pairwise comparisons (Bonferroni t-test, n=6

neurons).

To evaluate the effect of chlorophene on NMDA-evoked currents, NMDA currents were

pharmacologically isolated by supplementing the extracelluar solution with 50 µM picrotoxin to

block $GABA_A$ receptors and 10 µM NBQX to block AMPA receptors. After establishing a stable

baseline recording period, NMDA (1 µM, 10 µM or 20 µM, Merck, CAS 6384-92-5) was washed in

through the perfusion system and the change of I_{hold}, representing a NMDA evoked current, was

monitored. After reaching a stable NMDA evoked current (~3 min) chlorophene was bath applied.

In a subset of cells, 50 µM APV was applied at the end of the experiment to block NMDA evoked

currents. To compare the effect of chlorophene and APV exposure on NMDA currents, I_{hold} was

calculated (i) 3 min after NMDA application (I_{hold} +NMDA), (ii) 2 min after chlorophene application

(I_{hold} +chloro) and (iii) 1 min after APV application (I_{hold} +APV). The effect of chlorophene on NMDA

currents was compared by a paired t-test (I_{hold} +NMDA vs. I_{hold} +chloro) and for the subset of cells

with APV application by repeated measures ANOVA followed by pairwise comparisons

(Bonferroni t-test).

The pipette solution for current clamp recordings contained 115 mM KMeSO$_3$, 5 mM NaCl, 10 mM

HEPES, 0.1 mM EGTA, 4 mM Mg-ATP, 0.3 mM Na-GTP, 20 mM K$_2$-Phosphocreatin. The pH was

adjusted with KOH to 7.33 and osmolarity was adjusted with sucrose to 298 mOsm. Voltages were

not corrected for a calculated liquid junction potential of 9.6 mV. To test the effect of chlorophene

on basal neuronal properties and neuronal firing the internal chloride concentration was set to 5

mM to mimic physiological chloride gradients and hyperpolarizing GABA$_A$ receptor-mediated

conductance. To isolate the effect of chlorophene via GABA$_A$ receptors, the extracelluar solution

was supplemented by 10 µM NBQX and 50 µM APV to block AMPA and NMDA receptors. The

built-in bridge compensation was adjusted to 100% of the series resistance determined in voltage-

clamp. Passive and active neuronal properties of single neurons were measured before (control)

and after chlorophene application, respectively. The input resistance (R_{in}) was calculated from

alternating subthreshold current injections from −50 to 20 pA (10 pA steps, 200 ms duration).

Action potential firing was evoked by suprathreshold depolarizing current injections. The current

threshold required for eliciting an action potential was denoted as rheobase. The action potential

frequency was calculated from the number of action potentials in response to a 200 ms current

injection with an amplitude twice as large as the rheobase. The resting membrane potential (V_{rest})

was measured ~1 min after obtaining whole-cell access as the membrane voltage with zero

current in current clamp mode. For statistical comparison of neuronal properties under control and

chlorophene condition, p values were calculated by paired t-tests.

Human multi-neurotransmitter receptor (hMNR) assay

The human multi-neurotransmitter (hMNR) assay used 3D BrainSpheres derived from human-

induced pluripotent stem cell (hiPSC)-based neural progenitor cells (hiNPCs). The cultivation of

the hiPSC line IMR90 (clone 4, WiCell, Madison, WI, USA), including neural induction, was carried

out according to a previously described 2D-NIM protocol[146]. Briefly, hiPSC colonies were

singularized and cultured in defined medium, including dual SMAD inhibitors SB-431542 and LDN-

193189 for 21 d in 2D. On day 21 of neural induction, hiNPCs were cryopreserved in single-cell

suspension. For sphere generation, hiNPCs were thawed, transferred to 6-well plates, and

cultivated in a gyrical shaking incubator (140 rpm, 12.5 mm diameter) in a proliferation medium[146].

To start the 3D neural differentiation, hiNPC spheres (0.3 mm in diameter) were cultivated in

CINDA+ differentiation medium[146,147] for 3 weeks, and half of the medium was replaced twice

weekly. After 3D differentiation, BrainSpheres were plated on pre-coated (Poly-L-Ornithine, Laminin LN521) 96-well microelectrode arrays (MEA, #M768-tMEA-96B, Axion Biosystems) in CINDA+ differentiation medium (1 sphere per well) and fed twice per week by replacing half of the medium. After 4 weeks of differentiation on the MEAs, spontaneous electrical baseline activity was recorded with the Axion Maestro Pro MEA system. For initial unit identification, BrainSpheres were acutely exposed to the respective neurotransmitter agonists including glutamate (50 µM), and γ-aminobutyric acid (GABA; 10 µM). After electrical activity was measured, neurotransmitter antagonists were added to each well (10 µM bicuculline, 50 µM/50 µM D-AP5/NBQX), network activity was recorded and substances were removed with a complete exchange of the medium. After a 3 h washout, the second baseline was recorded and chlorophene was gradually added at sequentially increasing concentrations (0.25, 0.74, 2.22, 6.67, 20 µM). Extracellular recording of electrical activity was performed for 15 min at 37°C and 5% CO_2 in the Axion Maestro Pro System (Axion Biosystems), in line with previous work[146]. For further analysis of detected spikes, recorded .spk files were concatenated in the order of measurement and converted into a single .nex file with a MATLAB (R2021b, R2022b, MathWorks, Natick, MA, USA) script. Spike sorting of the .nex file was performed using the Offline Sorter (OFS, version 4.4, Plexon, Dallas, TX, USA) software, applying the automatic clustering T-Distribution EM method with 10 degrees of freedom (DOF) and an initial number of units of 20. The sorted units were exported as per-unit and per-waveform data. BrainSphere spiking data was analyzed using a custom KNIME (version 4.1.4, KNIME AG, www.knime.com) workflow implemented with R scripts[133] for statistical analysis and data visualization using the add-on package ggplot2[134]. For determination of neuronal subtype-specific responses, only units responding with a minimum fold change of 0.25 (increase or decrease) to the respective neurotransmitter and neurotransmitter receptor antagonist treatment compared to the baseline recording were analyzed for a specific response to chlorophene. Units were classified as GABAergic if, relative to the first baseline recording, the fold change in spiking following bicuculline (10 µM) and GABA (10 µM) treatments were ≤0.75 and ≥1.25, respectively Similarly, units were defined as glutamatergic if glutamate (50 µM) increased baseline spiking activity by factor ≥1.25, and if APV/NBQX (50 µM/50 µM) reduced spiking relative to prior glutamate treatment by factor ≤0.75. To account for differences in baseline spiking frequency between individual electrodes, the weighted mean firing rate (wMFR), representing the number of recorded spikes per second and recording, was scaled to the sum of wMFRs across all recordings including both the second baseline recording and subsequent recordings under chlorophene exposure. To ease visual data accessibility, scaled wMFRs were shifted by 1-median$_{control}$ to set the control median to one. As described above, a two-sample bootstrapping test and Benjamini-Hochberg

769   correction were used to calculate p-values and to determine significant differences between

770   control and treatment groups.

References

1.   Persson, L. *et al.* Outside the Safe Operating Space of the Planetary Boundary for Novel
Entities. *Environ. Sci. Technol.* **56**, 1510–1521 (2022).

2.   Wang, Z., Walker, G. W., Muir, D. C. G. & Nagatani-Yoshida, K. Toward a Global
Understanding of Chemical Pollution: A First Comprehensive Analysis of National and Regional
Chemical Inventories. *Environ. Sci. Technol.* **54**, 2575–2584 (2020).

3.   Landrigan, P. J. *et al.* The Lancet Commission on pollution and health. *The Lancet* **391**,
462–512 (2018).

4.   Fuller, R. *et al.* Pollution and health: a progress update. *The Lancet Planetary Health* **6**,
e535–e547 (2022).

5.   Harris, J. B. Neurotoxicology: what the neurologist needs to know. *Journal of Neurology,*
Neurosurgery & Psychiatry **75**, iii29–iii34 (2004).

6.   Spivey, A. Rotenone and Paraquat Linked to Parkinson's Disease: Human Exposure
Study Supports Years of Animal Studies. *Environ Health Perspect* **119**, a (2011).

7.   Weisskopf, M. G. *et al.* Association of Cumulative Lead Exposure with Parkinson's
Disease. *Environ Health Perspect* **118**, 1609–1613 (2010).

8.   Elonheimo, H. M., Andersen, H. R., Katsonouri, A. & Tolonen, H. Environmental
Substances Associated with Alzheimer's Disease—A Scoping Review. *IJERPH* **18**, 11839
(2021).

9.   Billioti De Gage, S. *et al.* Benzodiazepine use and risk of Alzheimer's disease: case-
control study. *BMJ* **349**, g5205–g5205 (2014).

10.   Lu, C. & Tune, L. E. Chronic Exposure to Anticholinergic Medications Adversely Affects
the Course of Alzheimer Disease. *The American Journal of Geriatric Psychiatry* **11**, 458–461
(2003).

11.   Yousef, S. *et al.* Attention Deficit Hyperactivity Disorder and Environmental Toxic Metal
Exposure in the United Arab Emirates. *Journal of Tropical Pediatrics* **57**, 457–460 (2011).

12.   Moore, S. *et al.* The Association between ADHD and Environmental Chemicals—A
Scoping Review. *IJERPH* **19**, 2849 (2022).

13.   Grandjean, P. & Landrigan, P. J. Neurobehavioural effects of developmental toxicity. *The*
Lancet Neurology **13**, 330–338 (2014).

14.   McCormack, A. L. *et al.* Environmental Risk Factors and Parkinson's Disease: Selective
Degeneration of Nigral Dopaminergic Neurons Caused by the Herbicide Paraquat. *Neurobiology*
of Disease **10**, 119–127 (2002).

15.   Moreno, J. A. *et al.* Age-Dependent Susceptibility to Manganese-Induced Neurological
Dysfunction. *Toxicological Sciences* **112**, 394–404 (2009).

16.   Eriksson, P., Jakobsson, E. & Fredriksson, A. Brominated flame retardants: a novel class
of developmental neurotoxicants in our environment? *Environ Health Perspect* **109**, 903–908
(2001).

17.   Branchi, I., Alleva, E. & Costa, L. G. Effects of Perinatal Exposure to a Polybrominated
Diphenyl Ether (PBDE 99) on Mouse Neurobehavioural Development. *NeuroToxicology* **23**,
375–384 (2002).

18.   Eriksson, P., Viberg, H., Jakobsson, E., Orn, U. & Fredriksson, A. A Brominated Flame
Retardant, 2,2`,4,4`,5-Pentabromodiphenyl Ether: Uptake, Retention, and Induction of
Neurobehavioral Alterations in Mice during a Critical Phase of Neonatal Brain Development.
Toxicological Sciences **67**, 98–103 (2002).

19.   Zhou, C.-C. *et al.* Lead exposure induces Alzheimers's disease (AD)-like pathology and
disturbes cholesterol metabolism in the young rat brain. *Toxicology Letters* **296**, 173–183 (2018).

20.   Ossowska, K. *et al.* A slowly developing dysfunction of dopaminergic nigrostriatal
neurons induced by long-term paraquat administration in rats: an animal model of preclinical
stages of Parkinson's disease? *Eur J of Neuroscience* **22**, 1294–1304 (2005).

21.   Wu, J. *et al.* Alzheimer's Disease (AD)-Like Pathology in Aged Monkeys after Infantile
Exposure to Environmental Metal Lead (Pb): Evidence for a Developmental Origin and
Environmental Link for AD. *J. Neurosci.* **28**, 3–9 (2008).

22.   Masilamoni, G. J. & Smith, Y. Chronic MPTP administration regimen in monkeys: a
model of dopaminergic and non-dopaminergic cell loss in Parkinson's disease. *J Neural Transm*
125, 337–363 (2018).

23.   Lee, J. & Freeman, J. L. Zebrafish as a model for investigating developmental lead (Pb)
neurotoxicity as a risk factor in adult neurodegenerative disease: A mini-review.
NeuroToxicology **43**, 57–64 (2014).

24.   Yen, P.-L., How, C. M. & Hsiu-Chuan Liao, V. Early-life and chronic exposure to di(2-
ethylhexyl) phthalate enhances amyloid-β toxicity associated with an autophagy-related gene in
Caenorhabditis elegans Alzheimer's disease models. *Chemosphere* **273**, 128594 (2021).

25.   Martin, M. M. *et al.* An expert-driven literature review of "negative" chemicals for
developmental neurotoxicity (DNT) in vitro assay evaluation. *Neurotoxicology and Teratology* **93**,
107117 (2022).

26.   OECD. *Test No. 424: Neurotoxicity Study in Rodents*. (OECD, 1997).
doi:10.1787/9789264071025-en.

27.   OECD. *Test No. 426: Developmental Neurotoxicity Study*. (OECD, 2007).
doi:10.1787/9789264067394-en.

28.   U.S. Environmental Protection Agency. Health Effects Guidelines OPPTS 870.6300
Developmental Neurotoxicity Study.
https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=P100IRWO.txt (1998).

29.   Crofton, K. M., Mundy, W. R. & Shafer, T. J. Developmental neurotoxicity testing: A path
forward. *Congenital Anomalies* **52**, 140–146 (2012).

30.   Tohyama, C. Developmental neurotoxicity test guidelines: problems and perspectives. *J.
Toxicol. Sci.* **41**, SP69–SP79 (2016).

31.   Smirnova, L. Developmental neurotoxicity – Challenges in the 21st Century and In Vitro
Opportunities. *ALTEX* (2014) doi:10.14573/altex.1403271.

32. *Toxicity Testing in the 21st Century: A Vision and a Strategy*. (The National Academies
Press, Washington, DC, 2007).

33. Tal, T. *et al.* New approach methods to assess developmental and adult neurotoxicity for
regulatory use: a PARC work package 5 project. *Front. Toxicol.* **6**, 1359507 (2024).

34. Crofton, K. M. *et al.* Current status and future directions for a neurotoxicity hazard
assessment framework that integrates in silico approaches. *Computational Toxicology* **22**,
100223 (2022).

35. Sachana, M. *et al.* International Regulatory and Scientific Effort for Improved
Developmental Neurotoxicity Testing. *Toxicological Sciences* **167**, 45–57 (2019).

36. Blum, J. *et al.* Establishment of a human cell-based in vitro battery to assess
developmental neurotoxicity hazard of chemicals. *Chemosphere* **311**, 137035 (2023).

37. Howe, K. *et al.* The zebrafish reference genome sequence and its relationship to the
human genome. *Nature* **496**, 498–503 (2013).

38. Strähle, U. *et al.* Zebrafish embryos as an alternative to animal experiments—A
commentary on the definition of the onset of protected life stages in animal welfare regulations.
Reproductive Toxicology **33**, 128–132 (2012).

39. European Parliament, Council of the European Union. Directive 2010/63/EU of the
European Parliament and of the Council of 22 September 2010 on the protection of animals
used for scientific purposes Text with EEA relevance. (2010).

40. Cassar, S. *et al.* Use of Zebrafish in Drug Discovery Toxicology. *Chem. Res. Toxicol.* **33**,
95–118 (2020).

41. Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. & Schilling, T. F. Stages of
embryonic development of the zebrafish. *Developmental Dynamics* **203**, 253–310 (1995).

42. Brustein, E., Marandi, N., Kovalchuk, Y., Drapeau, P. & Konnerth, A. 'In vivo' monitoring
of neuronal network activity in zebrafish by two-photon Ca2+ imaging. *Pflugers Arch - Eur J*
Physiol **446**, 766–773 (2003).

43. Fetcho, J. R. & O'Malley, D. M. Visualization of active neural circuitry in the spinal cord of
intact zebrafish. *Journal of Neurophysiology* **73**, 399–406 (1995).

44. Burgess, H. A. & Granato, M. Sensorimotor Gating in Larval Zebrafish. *J. Neurosci.* **27**,
4984–4994 (2007).

45. Burgess, H. A. & Granato, M. Modulation of locomotor activity in larval zebrafish during
light adaptation. *Journal of Experimental Biology* **210**, 2526–2539 (2007).

46. Wolman, M. A., Jain, R. A., Liss, L. & Granato, M. Chemical modulation of memory
formation in larval zebrafish. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 15468–15473 (2011).

47. Best, J. D. *et al.* Non-Associative Learning in Larval Zebrafish. *Neuropsychopharmacol*
33, 1206–1215 (2008).

48. Bruni, G. *et al.* Zebrafish behavioral profiling identifies multitarget antipsychotic-like
compounds. *Nat Chem Biol* **12**, 559–566 (2016).

49.   Jordi, J. *et al.* High-throughput screening for selective appetite modulators: A
multibehavioral and translational drug discovery strategy. *Sci. Adv.* **4**, eaav1966 (2018).

50.   Kokel, D. *et al.* Rapid behavior-based identification of neuroactive small molecules in the
zebrafish. *Nat Chem Biol* **6**, 231–237 (2010).

51.   McCarroll, M. N. *et al.* Zebrafish behavioural profiling identifies GABA and serotonin
receptor ligands related to sedation and paradoxical excitation. *Nat Commun* **10**, 4078 (2019).

52.   Babin, P. J., Goizet, C. & Raldúa, D. Zebrafish models of human motor neuron diseases:
Advantages and limitations. *Progress in Neurobiology* **118**, 36–58 (2014).

53.   Bandmann, O. & Burton, E. A. Genetic zebrafish models of neurodegenerative diseases.
Neurobiology of Disease **40**, 58–65 (2010).

54.   Lee, J. & Freeman, J. Zebrafish as a Model for Developmental Neurotoxicity
Assessment: The Application of the Zebrafish in Defining the Effects of Arsenic, Methylmercury,
or Lead on Early Neurodevelopment. *Toxics* **2**, 464–495 (2014).

55.   Rankin, C. H. *et al.* Habituation revisited: An updated and revised description of the
behavioral characteristics of habituation. *Neurobiology of Learning and Memory* **92**, 135–138
(2009).

56.   Riedel, G. Glutamate receptor function in learning and memory. *Behavioural Brain
Research* **140**, 1–47 (2003).

57.   Li, F. & Tsien, J. Z. Memory and the NMDA Receptors. *N Engl J Med* **361**, 302–303
(2009).

58.   Hunt, D. L. & Castillo, P. E. Synaptic plasticity of NMDA receptors: mechanisms and
functional implications. *Current Opinion in Neurobiology* **22**, 496–508 (2012).

59.   Williams, A. J. *et al.* The CompTox Chemistry Dashboard: a community data resource for
environmental chemistry. *J Cheminform* **9**, 61 (2017).

60.   Kumar Singh, A., Nath Tiwari, M., Prakash, O. & Pratap Singh, M. A Current Review of
Cypermethrin-Induced Neurotoxicity and Nigrostriatal Dopaminergic Neurodegeneration. *CN* **10**,
64–71 (2012).

61.   EFSA Panel on Plant Protection Products and their Residues (EFSA PPR Panel) *et al.*
Development of Integrated Approaches to Testing and Assessment (IATA) case studies on
developmental neurotoxicity (DNT) risk assessment. *EFS2* **19**, (2021).

62.   European Chemicals Agency. Product-type 3 Assessment report: Chlorophene. (2017).

63.   Safety Assessment of Dichlorophene and Chlorophene1. *Int J Toxicol* **23**, 1–27 (2004).

64.   Keiser, M. J. *et al.* Relating protein pharmacology by ligand chemistry. *Nat Biotechnol* **25**,
197–206 (2007).

65.   Emran, F., Rihel, J. & Dowling, J. E. A Behavioral Assay to Measure Responsiveness of
Zebrafish to Changes in Light Intensities. *JoVE* 923 (2008) doi:10.3791/923.

66.   Jarema, K. A. *et al.* Developmental Neurotoxicity and Behavioral Screening in Larval
Zebrafish with a Comparison to Other Published Results. *Toxics* **10**, 256 (2022).

67.   Gaballah, S. *et al.* Evaluation of Developmental Toxicity, Developmental Neurotoxicity,
and Tissue Dose in Zebrafish Exposed to GenX and Other PFAS. *Environ Health Perspect* **128**,
047005 (2020).

68.   Leuthold, D., Klüver, N., Altenburger, R. & Busch, W. Can Environmentally Relevant
Neuroactive Chemicals Specifically Be Detected with the Locomotor Response Test in Zebrafish
Embryos? *Environ. Sci. Technol.* **53**, 482–493 (2019).

69.   Kavšek, M. Predicting later IQ from infant visual habituation and dishabituation: A meta-
analysis. *Journal of Applied Developmental Psychology* **25**, 369–393 (2004).

70.   McCall, R. B. & Carriger, M. S. A Meta-Analysis of Infant Habituation and Recognition
Memory Performance as Predictors of Later IQ. *Child Development* **64**, 57 (1993).

71.   McDiarmid, T. A., Bernardos, A. C. & Rankin, C. H. Habituation is altered in
neuropsychiatric disorders—A comprehensive review with recommendations for experimental
design and analysis. *Neuroscience & Biobehavioral Reviews* **80**, 286–305 (2017).

72.   Valsamis, B. & Schmid, S. Habituation and Prepulse Inhibition of Acoustic Startle in
Rodents. *JoVE* 3446 (2011) doi:10.3791/3446.

73.   Ludewig, K. *et al.* The acoustic startle reflex and its modulation: effects of age and
gender in humans. *Biological Psychology* **63**, 311–323 (2003).

74.   Roberts, A. C. *et al.* Long-term habituation of the C-start escape response in zebrafish
larvae. *Neurobiology of Learning and Memory* **134**, 360–368 (2016).

75.   Roberts, A. C. *et al.* Habituation of the C-Start Response in Larval Zebrafish Exhibits
Several Distinct Phases and Sensitivity to NMDA Receptor Blockade. *PLoS ONE* **6**, e29132
(2011).

76.   Morris, R. Synaptic plasticity and learning: selective impairment of learning rats and
blockade of long-term potentiation in vivo by the N-methyl-D- aspartate receptor antagonist AP5.
J. Neurosci. **9**, 3040–3057 (1989).

77.   Wozniak, D. F., Olney, J. W., Kettinger, L., Price, M. & Miller, J. P. Behavioral effects of
MK-801 in the rat. *Psychopharmacology* **101**, 47–56 (1990).

78.   Paoletti, P., Bellone, C. & Zhou, Q. NMDA receptor subunit diversity: impact on receptor
properties, synaptic plasticity and disease. *Nat Rev Neurosci* **14**, 383–400 (2013).

79.   Zoodsma, J. D. *et al.* A Model to Study NMDA Receptors in Early Nervous System
Development. *J. Neurosci.* **40**, 3631–3645 (2020).

80.   Galligan, J. J. & North, R. A. MK-801 blocks nicotinic depolarizations of guinea pig
myenteric neurons. *Neuroscience Letters* **108**, 105–109 (1990).

81.   Rothman, S. Noncompetitive N-methyl-D-aspartate antagonists affect multiple ionic
currents. *J Pharmacol Exp Ther* **246**, 137–142 (1988).

82.   Mancuso, C. E., Tanzi, M. G. & Gabay, M. Paradoxical Reactions to Benzodiazepines:
Literature Review and Treatment Options. *Pharmacotherapy* **24**, 1177–1185 (2004).

83.   McCarthy, M. M., Brown, E. N. & Kopell, N. Potential Network Mechanisms Mediating
Electroencephalographic Beta Rhythm Changes during Propofol-Induced Paradoxical Excitation.
J. Neurosci. **28**, 13488–13504 (2008).

84.   Reddy, R. V., Moorthy, S. S., Dierdorf, S. F., Deitch, R. D. & Link, L. Excitatory Effects
and Electroencephalographic Correlation of Etomidate, Thiopental, Methohexital, and Propofol:
Anesthesia & Analgesia **77**, 1008???1011 (1993).

85.   Iqbal, F. *et al.* Anesthetics: from modes of action to unconsciousness and neurotoxicity.
Journal of Neurophysiology **122**, 760–787 (2019).

86.   Århem, P., Klement, G. & Nilsson, J. Mechanisms of Anesthesia: Towards Integrating
Network, Cellular, and Molecular Level Modeling. *Neuropsychopharmacol* **28**, S40–S47 (2003).

87.   Armstrong, R. *et al.* Mechanisms of Anesthetic Action and Neurotoxicity: Lessons from
Molluscs. *Front. Physiol.* **8**, 1138 (2018).

88.   Franks, N. P. & Lieb, W. R. Molecular and cellular mechanisms of general anaesthesia.
Nature **367**, 607–614 (1994).

89.   Sun, L. Early childhood general anaesthesia exposure and neurocognitive development.
British Journal of Anaesthesia **105**, i61–i68 (2010).

90.   Monesson-Olson, B. *et al.* Expression of the eight GABAA receptor α subunits in the
developing zebrafish central nervous system. *PLoS ONE* **13**, e0196083 (2018).

91.   Sadamitsu, K. *et al.* Characterization of zebrafish GABAA receptor subunits. *Sci Rep* **11**,
6242 (2021).

92.   Collinson, N. *et al.* Enhanced Learning and Memory and Altered GABAergic Synaptic
Transmission in Mice Lacking the α5 Subunit of the GABA $_A$ Receptor. *J. Neurosci.* **22**, 5572–
5580 (2002).

93.   Crestani, F. *et al.* Trace fear conditioning involves hippocampal α $_5$ GABA $_A$ receptors.
Proc. Natl. Acad. Sci. U.S.A. **99**, 8980–8985 (2002).

94.   Yee, B. K. *et al.* GABA $_A$ receptors containing the α5 subunit mediate the trace effect in
aversive and appetitive conditioning and extinction of conditioned fear. *Eur J of Neuroscience*
20, 1928–1936 (2004).

95.   Navarro, J. F., Burón, E. & Martín-López, M. Anxiogenic-like activity of L-655,708, a
selective ligand for the benzodiazepine site of GABAA receptors which contain the alpha-5
subunit, in the elevated plus-maze test. *Progress in Neuro-Psychopharmacology and Biological*
Psychiatry **26**, 1389–1392 (2002).

96.   Atack, J. R. *et al.* L-655,708 enhances cognition in rats but is not proconvulsant at a dose
selective for α5-containing GABAA receptors. *Neuropharmacology* **51**, 1023–1029 (2006).

97.   Dawson, G. R. *et al.* An Inverse Agonist Selective for α5 Subunit-Containing GABA $_A$
Receptors Enhances Cognition. *J Pharmacol Exp Ther* **316**, 1335–1345 (2006).

98.   Ballard, T. M. *et al.* RO4938581, a novel cognitive enhancer acting at GABAA α5
subunit-containing receptors. *Psychopharmacology* **202**, 207–223 (2009).

99. Herbison, A. E. & Moenter, S. M. Depolarising and Hyperpolarising Actions of GABAA
Receptor Activation on Gonadotrophin-Releasing Hormone Neurones: Towards an Emerging
Consensus: GABAA modulation of GnRH neurones. *Journal of Neuroendocrinology* **23**, 557–
569 (2011).

100. Cui, Y. *et al.* Neurofibromin Regulation of ERK Signaling Modulates GABA Release and
Learning. *Cell* **135**, 549–560 (2008).

101. Yoshiike, Y. *et al.* GABAA Receptor-Mediated Acceleration of Aging-Associated Memory
Decline in APP/PS1 Mice and Its Pharmacological Treatment by Picrotoxin. *PLoS ONE* **3**, e3029
(2008).

102. Portavella, M., Vargas, J. P., Torres, B. & Salas, C. The effects of telencephalic pallial
lesions on spatial, temporal, and emotional learning in goldfish. *Brain Research Bulletin* **57**, 397–
399 (2002).

103. Rodríguez, F. *et al.* Spatial memory and hippocampal pallium through vertebrate
evolution: insights from reptiles and teleost fish. *Brain Research Bulletin* **57**, 499–503 (2002).

104. Saleem, S. & Kannan, R. R. Zebrafish: an emerging real-time model system to study
Alzheimer's disease and neurospecific drug discovery. *Cell Death Discov.* **4**, 45 (2018).

105. Nam, R.-H., Kim, W. & Lee, C.-J. NMDA receptor-dependent long-term potentiation in
the telencephalon of the zebrafish. *Neuroscience Letters* **370**, 248–251 (2004).

106. Tsien, J. Z. Building a Brainier Mouse. *Sci Am* **282**, 62–68 (2000).

107. Newcomer, J. W. & Krystal, J. H. NMDA receptor regulation of memory and behavior in
humans. *Hippocampus* **11**, 529–542 (2001).

108. Brown, D. A. & Passmore, G. M. Neural *KCNQ* (Kv7) channels. *British J Pharmacology*
156, 1185–1195 (2009).

109. De Leon, J. & Nikoloff, D. M. Paradoxical Excitation on Diphenhydramine May Be
Associated with Being a CYP2D6 Ultrarapid Metabolizer: *Three Case Reports. CNS spectr.* **13**,
133–135 (2008).

110. National Industrial Chemicals Notification and Assessment Scheme. Chlorophene
preservatives: Environment tier II assessment. (2019).

111. Yu, J. T., Bouwer, E. J. & Coelhan, M. Occurrence and biodegradability studies of
selected pharmaceuticals and personal care products in sewage effluent. *Agricultural Water
Management* **86**, 72–80 (2006).

112. Kasprzyk-Hordern, B., Dinsdale, R. M. & Guwy, A. J. The removal of pharmaceuticals,
personal care products, endocrine disruptors and illicit drugs during wastewater treatment and
its impact on the quality of receiving waters. *Water Research* **43**, 363–380 (2009).

113. REACH. REACH Dossier for Chlorofene. (2019).

114. OECD. *Test No. 210: Fish, Early-Life Stage Toxicity Test.* (OECD, 2013).
doi:10.1787/9789264203785-en.

115. Kleinstreuer, N. C. *et al.* Development and Validation of a Computational Model for
Androgen Receptor Activity. *Chem. Res. Toxicol.* **30**, 946–964 (2017).

116.   The European Parliament and the Council of the European Union. Regulation (EC) No
1272/2008 of the European Parliament and of the Council of 16 December 2008 on
classification, labelling and packaging of substances and mixtures, amending and repealing
Directives 67/548/EEC and 1999/45/EC, and amending Regulation (EC) No 1907/2006. (2008).

117.   European Chemicals Agency. Minutes of the 62nd Meeting of the Committee for Risk
Assessment (RAC-62). (2022).

118.   European Chemicals Agency. *Guidance on Information Requirements and Chemical*
Safety Assessment: Chapter R.7a : Endpoint Specific Guidance. (Publications Office, LU, 2017).

119.   European Chemicals Agency. *Non-Animal Approaches: Current Status of Regulatory*
Applicability under the REACH, CLP and Biocidal Products Regulations. (Publications Office,
LU, 2017).

120.   European Chemicals Agency. *Guidance on the Biocidal Products Regulation: Volume III :*
Human Health, Assessment and Evaluation (Parts B+C). (Publications Office, LU, 2017).

121.   Kenda, M., Karas Kuželički, N., Iida, M., Kojima, H. & Sollner Dolenc, M. Triclocarban,
Triclosan, Bromochlorophene, Chlorophene, and Climbazole Effects on Nuclear Receptors: An
in Silico and *in Vitro* Study. *Environ Health Perspect* **128**, 107005 (2020).

122.   Gladkevich, A., Korf, J., Hakobyan, V. P. & Melkonyan, K. V. The peripheral GABAergic
system as a target in endocrine disorders. *Autonomic Neuroscience* **124**, 1–8 (2006).

123.   Casoni, F. *et al.* SDF and GABA interact to regulate axophilic migration of GnRH
neurons. *Journal of Cell Science* jcs.101675 (2012) doi:10.1242/jcs.101675.

124.   Eepho, O.-I. *et al.* Modulation of GABA by sodium butyrate ameliorates hypothalamic
inflammation in experimental model of PCOS. *BMC Neurosci* **24**, 62 (2023).

125.   Ingram, R. J., Leverton, L. K., Daniels, V. C., Li, J. & Christian-Hinman, C. A. Increased
GABA transmission to GnRH neurons after intrahippocampal kainic acid injection in mice is sex-
specific and associated with estrous cycle disruption. *Neurobiology of Disease* **172**, 105822
(2022).

126.   European Commission. Communication from the Commission: The European Green
Deal. (2019).

127.   Fritsche, E. *et al.* Current Availability of Stem Cell-Based In Vitro Methods for
Developmental Neurotoxicity (DNT) Testing. *Toxicological Sciences* **165**, 21–30 (2018).

128.   Bal-Price, A. K., Hogberg, H. T., Buzanska, L. & Coecke, S. Relevance of in vitro
neurotoxicity testing for regulatory requirements: Challenges to be considered. *Neurotoxicology*
and Teratology **32**, 36–41 (2010).

129.   Bally-Cuif, L. & Vernier, P. Organization and physiology of the zebrafish nervous system.
in *Fish Physiology* vol. 29 25–80 (Elsevier, 2010).

130.   Kaslin, J. & Ganz, J. Zebrafish Nervous Systems. in *The Zebrafish in Biomedical*
Research 181–189 (Elsevier, 2020). doi:10.1016/B978-0-12-812431-4.00018-X.

131. Brox, S., Seiwert, B., Küster, E. & Reemtsma, T. Toxicokinetics of Polar Chemicals in
Zebrafish Embryo (*Danio rerio*): Influence of Physicochemical Properties and of Biological
Processes. *Environ. Sci. Technol.* **50**, 10264–10272 (2016).

132. Loerracher, A.-K. & Braunbeck, T. Cytochrome P450-dependent biotransformation
capacities in embryonic, juvenile and adult stages of zebrafish (Danio rerio)—a state-of-the-art
review. *Arch Toxicol* **95**, 2299–2334 (2021).

133. R Core Team. R: A language and environment for statistical computing. (2021).

134. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer, Switzerland, 2016).
doi:10.1007/978-3-319-24277-4.

135. Neuwirth, E. RColorBrewer. (2022).

136. Gu, Z. Complex heatmap visualization. *iMeta* **1**, e43 (2022).

137. Lloyd, D. R., Medina, D. J., Hawk, L. W., Fosco, W. D. & Richards, J. B. Habituation of
reinforcer effectiveness. *Front. Integr. Neurosci.* **7**, (2014).

138. Kulesa, A., Krzywinski, M., Blainey, P. & Altman, N. Sampling distributions and the
bootstrap. *Nat Methods* **12**, 477–478 (2015).

139. Harpaz, R. *et al.* Collective behavior emerges from genetically controlled simple
behavioral motifs in zebrafish. *Sci. Adv.* **7**, eabi7460 (2021).

140. Efron, B. Bootstrap Methods: Another Look at the Jackknife. *Ann. Statist.* **7**, (1979).

141. de Mendiburu, F. & Yaseen, M. agricolae: Statistical Procedures for Agricultural
Research. (2020).

142. Cameron, L. P. *et al.* A non-hallucinogenic psychedelic analogue with therapeutic
potential. *Nature* **589**, 474–479 (2021).

143. Zhang, X. D. A pair of new statistical parameters for quality control in RNA interference
high-throughput screening assays. *Genomics* **89**, 552–561 (2007).

144. Judson, R. *et al.* Editor's Highlight: Analysis of the Effects of Cell Stress and Cytotoxicity
on *In Vitro* Assay Activity Across a Diverse Chemical and Assay Space. *Toxicol. Sci.* **152**, 323–
339 (2016).

145. Moutin, E. *et al.* Procedures for Culturing and Genetically Manipulating Murine
Hippocampal Postnatal Neurons. *Front. Synaptic Neurosci.* **12**, 19 (2020).

146. Hartmann, J. *et al.* Molecular and Functional Characterization of Different BrainSphere
Models for Use in Neurotoxicity Testing on Microelectrode Arrays. *Cells* **12**, 1270 (2023).

147. Nimtz, L. *et al.* Characterization and application of electrically active neuronal networks
established from human induced pluripotent stem cell-derived neural progenitor cells for
neurotoxicity evaluation. *Stem Cell Research* **45**, 101761 (2020).

Acknowledgements

We thank Gi Mick Wu (UFZ) for statistical consultation. This work was supported by a W2 Helmholtz Association Grant to T.T. and the Horizon Europe project PARC (Grant Agreement No 101057014) to T.T. and E.F.

Author contributions

D.L. designed and performed the research, analyzed the data, and wrote the manuscript. N.K.H., J.N., K.B., and I.S. performed experiments and analyzed data. T.T. designed the study and wrote the manuscript. All authors contributed to data interpretation and commented on the manuscript.

Corresponding authors

Correspondence should be addressed to T.T. and D.L.

Availability of data and code

All data (DOI: 10.6084/m9.figshare.26819890) and code (DOI: 10.6084/m9.figshare.26819914) were uploaded to FigShare and will be made public upon paper acceptance.