Particular sequence characteristics induce bias in the detection of polymorphic transposable element insertions

Marie Verneret[1,2], Van Anthony Le[1], Thomas Faraut[3], Jocelyn Turpin[2], Emmanuelle Lerat[1*]

1 Universite Claude Bernard Lyon 1, LBBE, UMR5558, CNRS, VAS, Villeurbanne, F-69622, France.

2 IVPC UMR754, INRAE, Universite Claude Bernard Lyon 1, EPHE, Université PSL, Lyon, F-69007, France

3 GenPhySE, Universite de Toulouse, INRAE, INPT, ENVT, 31326, Castanet Tolosan, France

*corresponding author

Abstract

Transposable elements (TEs) have an important role in genome evolution but are challenging for bioinformatics detection due to their repetitive nature and ability to move and replicate within genomes. New sequencing technologies now enable the characterization of nucleotide and structural variations within species. Among them, TE polymorphism is critical to identify as it may influence species adaptation or trigger diseases. Despite the development of numerous bioinformatic programs, identifying the most effective tool is challenging due to non-overlapping results and varying efficiency across studies. Benchmarking efforts have highlighted some of the limitations of these tools, often evaluated on either real or simulated data. However, real data may be incomplete or contain unannotated TEs, while simulated data may not accurately reflect real genomes. This study introduces a simulation method generating data based on real genomes to control all genomic parameters. Evaluating several TE polymorphic detection tools using data from *Drosophila melanogaster* and *Arabidopsis thaliana,* our study investigates factors like copy size, sequence divergence, and GC content that influence detection efficiency. Our results indicate that only a few programs perform satisfactorily and that all are sensitive to TE and genomic characteristics that may differ according to the species considered. Using *Bos taurus* population data as a case study to identify polymorphic LTR-retrotransposon insertions, we found low-frequency insertions particularly challenging to detect due to a high number of false positives. Increased sequencing coverage improved sensitivity but reduced precision. Our work underscores the importance of selecting appropriate tools and thresholds according to the specific research questions.

Introduction

Recognized as being among the most important players in the evolution of genomes, transposable elements (TEs) represent a real challenge for bioinformatics approaches to detect them. TEs are repeated sequences present in almost all eukaryotic genomes. They have the ability to move and replicate, forming different families of similar but not always identical sequences. Several types have been described, depending on their

structure and their mode of transposition, varying both in genomic distribution and in sequence length

(Wicker et al. 2007). For example, LTR-retrotransposons represent sequences of approximately 10 kb but

DNA transposons such as MITEs (Miniature Inverted Repeats Transposable Elements) span only a few

hundred base pairs. Moreover, TEs are not randomly distributed in the genome since their insertion patterns

reflect a balance between selection pressure against their deleterious effects and genetic drift (Bourque et al.

2018). As a consequence, TEs are likely to be found inserted into each other constituting nested insertions,

which are particularly difficult to automatically identify (Bergman and Quesneville 2007). In addition, their

proportion in genomes can vary greatly, ranging from a few percent as for example in the honeybee

(Weinstock et al. 2006) to the major part of the genome as in maize (Schnable et al. 2009). Over the past

twenty years, different bioinformatic tools have been developed allowing their annotation in assembled

genomes (Lerat 2019). However, the rapid development of new sequencing technologies has made it

possible to access numerous data from different individuals or populations in order to characterize the

nucleotide and structural variations within a given species. Indeed, a reference genome for a given species is

not sufficient to reflect the overall diversity of individuals. In particular, although TEs are generally

regulated in a genome to prevent their activity, certain TE families can nevertheless continue to transpose

throughout the life of an individual or may be reactivated due to some stress (Di Stefano 2022). It has been

proposed that in Drosophila, the transposition rate is comparable to that of the nucleotide mutation rate

(Adrion et al. 2017). More recently, according to the TE family, the transposition rate has been shown to be

higher with an average of 4.93×10^{-9} insertions per site per generation corresponding to a new insertion in

each new embryo (Wang et al. 2023). In humans, the most active TEs have a transposition rate of one

insertion every 20 births (Cordaux and Batzer 2009). We can thus expect to find variations in the TE

insertion pattern between individuals, which constitutes the TE polymorphism. Polymorphic TEs are

particularly important to identify since they represent insertions that may be at the basis of

species/population adaptation or triggering diseases. For example, numerous polymorphic TEs have been

detected in sub-populations of the Chinese white poplar (*Populus tomentosa)* some of them being under

positive selection while inserted in genes involved in stress, defense and immune responses (Zhao et al.

2022). In humans, a specific polymorphic TE insertion is associated with the development of the Fukuyama

type congenital muscular distrophy (Kobayashi et al. 1998).

In order to search for polymorphic insertions, bioinformatics tools have been developed to answer

specific questions and on particular organisms such as Drosophila, human or some plants (Lerat 2019). All

these methods follow similar principles in their functioning which consist first in mapping sequenced reads

to a reference genome and a set of reference TE sequences. Then two approaches, that can be combined,

have been proposed to detect the presence/absence of TEs. The first is to consider discordant read pairs with

one read mapping uniquely on a genomic location and the other mapping on different sequences of the same

TE family. The second approach considers split reads, *i.e.*, reads overlapping a junction between the genome

and a TE insertion, with a part of the read mapping uniquely on the genome while the other part maps on

several TE sequences. More than twenty programs have been developed during the past ten years (for an

exhaustive list, see https://tehub.org/), which makes it difficult for users to determine which program is the

most appropriate or the most efficient. In particular, the results of these programs are often not entirely

overlapping (Ewing 2015; Lerat et al. 2019). This makes it more difficult to identify true positives, especially

in the case of non-reference insertions, which correspond to insertions not present in the reference genome

but present in the analyzed read samples. Several attempts have been previously made to benchmark all these

programs (Nelson et al. 2017; Rishishwar et al. 2017; Vendrell-Mir et al. 2019; Chen et al. 2023). These

works showed that all these programs generally are not as efficient as indicated in their original publication.

However, these evaluations were made either on partial real data or on simulated data without controlling all

parameters, or were targeting only particular TE types like for example the approach by Vendrell-Mir

(Vendrell-Mir et al. 2019). A problem with real data is that they may be only partial or may contain

unannotated TE insertions that can blur the results. However, using partially simulated data is also

problematic since it usually does not reflect in a realistic manner a real genome and does not allow to control

all parameters. For example, the approach used by Rishiwar et al. (Rishishwar et al. 2017) consisted in the

random insertions of consensus sequences from three human TE families into human reference

chromosomes. In the work by Nelson et al. and Chen et al. (Nelson et al. 2017; Chen et al. 2023), they

inserted one single TE from one of the four active families of the yeast at positions that are supposed to be

biologically sound. These approaches are thus very biased toward the particularities of a single species.

Then, there are still several unanswered questions regarding the underperformance of certain tools,

particularly in relation to specific characteristics of the studied genome and the TE sequences themselves that

cannot be achieved using real data or simulated approaches used until now.

In this study, we have developed a simulation approach to produce data based on real genomes to

allow the complete control of all genomic parameters. Using data generated for *Drosophila melanogaster*

and *Arabidopsis thaliana*, we evaluated several TE polymorphic detection tools and investigated different

characteristics like the copy size, the sequence divergence, the distance between copies, the GC content of

the surrounding genomic regions, the Target Site Duplicate (TSD) size or the TE family that could explain

why some insertions are better detected than others. Our results show that only very few of the different

tested programs give satisfactory results and that all programs are sensitive to TE and genomic sequence

characteristics that slightly differ according to the species considered. As an application case, we used *Bos*

taurus real population data to identify polymorphic LTR-retrotransposon insertions. Low-frequency

insertions appeared to be more challenging to detect due to a high proportion of false positives. Increasing

sequencing coverage improved the sensitivity but at the expense of precision. Our study emphasizes the

importance of selecting appropriate tools and thresholds depending on the scientific questions asked.

Material and Methods

Genomic data used for simulation

110 The sequence of the *Drosophila melanogaster* 2L chromosome version 6.18 in GenBank format was

obtained from the NCBI GenBank database (accession number: NT_033779). The chromosome sequence is 23,513,712 bp long in which 3,519 genes and 919 TEs are annotated. For *Arabidopsis thaliana*, a GenBank file of the chromosome 1 was generated using the TAIR10 version of the gene and transposable element (TE) annotation in gff format available from the Arabidopsis Information Resource website (https://www.arabidopsis.org/). The chromosome sequence is 30,427,671 bp long in which 7,509 genes and 7,135 TEs are annotated. The sequence of the chromosome 25 from *Bos taurus* was obtained from the GenBank database (version ARS-UCD1.3, accession number: GCF_002263795.2). The chromosome, that is 42,350,435 bp long, contains 1,006 genes but no TEs have been previously annotated. We thus determined the position of endogenous retroviruses (ERV) using RepeatMasker version 2.0.3 with cattle ERV consensus sequences from Repbase version 29.03 (https://www.girinst.org/). ERV insertions from four ERV families were used for the simulations: two class I ERV families (ERV1-1_BT and BtERVF2) and two class II ERV families (ERV2-2_BT and ERV2-3_BT).

Simulation tool *replicaTE*

We have developed a simulation tool based on real data. This tool is implemented as several python3 scripts that need to be run successively, using as a starting point a GenBank file (Figure 1). In summary, we consider three types of sequences (genes, TEs and intergenic regions). The genes are cleaned up from any TE insertions. Intergenic regions are simulated to remove any misannotated TEs and based on the real intergenic regions with respect to their GC content and length. The real characteristics of TE insertions in the genome (number of copies, size of copies, %divergence, etc.) are used to simulate new TE sequences. These TE sequences are randomly assigned to the intergenic regions. Finally, all three parts are reassembled to create a complete simulated genome and a deleted simulated genome in which half of the TE insertions are removed. The tool is available as a git repository (https://github.com/e-lerat/replicaTE). For the simulation of the three species, default parameters for each module were used.

Figure 1: Workflow of the simulation tool *ReplicaTE*. Each step corresponds to the different scripts. In gray are indicated the genes, in blue and red are indicated the TE insertions, a given color corresponding to a given TE family. The simulated TE copies are represented in orange and light blue.

deleTE.py

This script allows us to get the characteristics of each element (genes, intergenic regions, and TEs) for the next steps and to generate a simulated genome without TEs. It takes as an input a GenBank file from which it will extract the annotations. It outputs multiple files which can then be used as input by the other codes. The size of the simulated intergenic regions are drawn from an exponentiated Weibull distribution constructed from the computed gene density (number of genes per Mb) with a minimal size of 200 pb. The GC content of the simulated intergenic regions are drawn from a truncated normal law fitted on the observed %GC of the chromosome sequence, with values between 0 and 100%.

generaTE.py

This script generates TE copies based on different characteristics (copy number, length, Target Site Duplication (TSD) length, strand). It attributes an intergenic region to each copy to be inserted into with the possibility to have nested insertions. For each family, a pool of copy sizes is drawn in a truncated exponential law, with values between 80 bp and 102.5% of the largest sequence of the family to take into account potential small insertions, called the "ancestral" sequence. The sequence divergences of the copies compared to the "ancestral" sequence are drawn from a truncated normal law distribution, with values between 0 and 20% (mean = 10 and standard deviation = 4). By default, the copy number corresponds to the observed copy number in the real chromosome. It is also possible to simulate the copy number. In that case, it is randomly drawn from an exponentiated Weilbull distribution fitted on the data. The TSDs have a length between 0 and 8 bp and are attributed for a given family when the option is specified.

inseraTE.py

This script associates the cleaned genes, the simulated TEs and the simulated intergenic regions to produce a

genomic sequence. The TE copies are randomly inserted into their attributed intergenic region. The insertion

can be 'normal' or 'nested' (inserted into a previous TE) and multiple nested events can arise. The complete

simulated chromosome is provided in fasta format. A "deleted" version is also generated, in which half of the

TE copies are not present.

Short-read simulation

The different tested tools all use short-read sequences as an input. We thus have generated short reads based

on either the "complete" or the "deleted" simulated chromosomes using the program ART Version 2.1.8

(Huang et al. 2012). This program produces theoretical reads expected by an NGS technique on a given

genome. For this analysis, we generated paired-end Illumina reads of 150 bp (with a fragment size of 300 bp)

using three different coverages (10X, 50X and 100X). Only 15X short reads were produced for *B. taurus* to

reflect the landscape of the real cattle data coverage in the public databases.

Polymorphic TE detection tools

Reference and non-reference insertions were detected in the simulated short reads using the either the

"complete" or the "deleted" simulated genomes as a reference with the 12 programs included in

McClintock2 (Nelson et al. 2017, Chen et al. 2023) in addition to TEPID (Stuart et al. 2016) and Jitterbug

(Hénaff et al. 2015) programs. All the programs were run with default parameters. The number of True

Positives and False Negatives were computed from the results of the different programs using two

homemade perl scripts (test_position_ref.pl and test_position_nonref.pl) available in the git repository (see

below).

Statistical analyses

All statistical tests were performed using the R software version 3.6.3 (2020-02-29) (R Core Team 2017).

The programs were evaluated according to different metrics described below.

Recall (sensitivity): it corresponds to the proportion of True Positives (TP) among all the TE insertions

present in the reference genome. It is computed as: $\dfrac{TP}{TP+FN}$

Precision: it corresponds to the proportion of good answers among the predicted TE insertions. It is

computed as: $\dfrac{TP}{TP+FP}$

F-score: it corresponds to the harmonic mean of the recall and the precision. It is computed as:

$2\dfrac{recall \, . \, precision}{recall + precision}$

To compute these different metrics, it is necessary to assess the number of TPs among the identified TE

insertions proposed by each program, using two homemade perl scripts "test_position_ref.pl" and

"test_position_nonref.pl" (available in the git repository). We considered an insertion to be a TP when the

program associates the same TE family name and a position that is close to the real position, with a certain

margin of error. More specifically, we considered four different margins of error to determine whether the

position was correct or not which are 5 bp, 20 bp, 100 bp and 150 bp. The False Negatives (FN) correspond

to insertions present in the reference dataset that were not detected by the program and the False Positives

(FP) correspond to predicted insertions that do not correspond to insertions present in the reference dataset.

False positive rate estimation in real data of *Bos taurus*

Endogenous retroviruses (ERV) insertion detection was performed using TEFLoN (Adrion et al. 2017) with

default parameters on 10 WGS short-read data samples from various individuals of *Bos taurus* (accession

numbers from SRA database are provided in Supplementary Table S1). A homemade python script

(FP_TP_teflon_insertion.py) available in the git repository, was applied to compute the proportion of TPs,

FPs and FPs among the identified insertions. Insertions found in common with the reference were considered

as TPs or FPs compared to the ERV annotation of the *B. taurus* ARS-UCD1.3 assembly. Insertions found in

the samples but not in the reference genome were considered as TPs if they were also present in the variant

output file obtain from a variant calling analysis on long-read data from the same samples using the *call*

function of pbsv version 2.6.2 with default parameters (https://github.com/PacificBiosciences/pbsv). In both

cases, we considered insertions as TPs if the program also associated the correct ERV family name and with

a correct position within 20 bp of error margin.

Results

Chromosome simulation and evaluation approach

The simulation tool *replicaTE* was used on the chromosome 2L of *D. melanogaster* and on the chromosome

1 of *A. thaliana* (all generated files are available as supplementary data). The first script, *deleTE.py*, produces

different output files. Among them, the "gene_clean_tab.csv" file contains the real genes without any

annotated internal TE insertions. The "intergenic_sim_tab.csv" file contains the simulated intergenic regions

with their length and %GC. The "stat_TEs_tab.csv" contains a sequence corresponding to the longest real TE

sequence (that will be considered as the "ancestral" TE sequence) of a given family that will be used to

generate all simulated TE copies and the number of copies for each family, that corresponds to the real

number of annotated copies in the considered chromosome. These two last files are used in the second script,

generaTE.py, to simulate the TE copies. It produces a fasta file containing the simulated sequences

("simulated_TEs.fas") and a text file ("param_TEs_tab.csv") containing all the information regarding each

TE family (length of each copy, sequence divergence of each copy compared to the "ancestral" TE sequence,

the associated intergenic region, the strand and the TSD size). These two files, in addition to the

"gene_clean_tab.csv" and the "stat_TEs_tab.csv" files, are used in the third script *inseraTE.py*. It produces,

among other files, the two simulated genomes in fasta format and the files "annot_TEs.tsv" and

"annot_TEs_del_1" containing all information regarding each TE copy (positions, length, divergence,

insertion type (nested or not), strand, TSD size, distances to the closest TE insertions, and the GC content of

the flanking genomic regions).

For each chromosome, we thus have all information about the inserted copies in addition to their

precise positions. These different parameters will be used to evaluate the tested programs. In particular, we

will be able to determine if particular factors relative to the TE sequences (size, distance to other copies,

divergence, TSD size) or to their genomic region of insertion (%GC) may have an influence on whether they

are correctly detected or not by the tested programs.

In our evaluation approach, the "complete" simulated chromosome and the "deleted" simulated

chromosome can be used alternatively as reference genome or as sample genome in order to evaluate the

possibility to identify reference / absent insertions or non-reference insertions. Indeed, as described on Figure

2, when using the "complete" simulated genome as a reference, the short reads will be generated using the

"deleted" simulated genome, in which half of the TE insertions are missing. This will allow us to evaluate

the capacity of the programs to detect both reference and absent insertions. Alternatively, if the "deleted"

simulated genome is used as a reference and the "complete" simulated genome is used to generate short

reads, then it will allow us to evaluate the capacity of the programs to detect TE insertions not present in the

reference.

Figure 2: Evaluation approach. The black rectangles correspond to genes. The orange and ligth blue circles

correspond to TE copies from two different families. The reads simulated on the "deleted" simulated genome will be

mapped to the "complete" simulated genome, which will allow to identify reference and/or absent TE insertions. The

reads simulated on the "complete" simulated genome will be mapped to the "deleted" simulated genome, which will

allow us to identify non-reference TE insertions.

Using the *D. melanogaster* and *A. thaliana* data, we have generated two simulated chromosomes for

each species. For Drosophila, the "complete" simulated chromosome, which is 23,298,325 bp long, contains

790 TEs whereas the "deleted" simulated chromosome contains 400 TEs. In the case of *A. thaliana*, the

"complete" simulated chromosome, which is 37,444,832 bp long, contains 6,324 TEs whereas the "deleted"

simulated chromosome contains 3,132 TEs. Knowing exactly the positions and name of each insertion, it is

thus possible to compute the number of True Positives (TP), False Positives (FP) and False Negatives (FN)

for each program allowing to determine their efficiency. Additionally, since we have all information about

the different insertions for which we can control all associated parameters (size, distance, %GC etc.), it will

be possible to compare the characteristics of the TP to those of the FN that could indicate any detection bias

in the tested programs.

Tests of the polymorphic TE detection programs

More than 20 programs have been proposed during the last 10 years to identify polymorphic TE insertions.

However, many of them were not possible to evaluate in this analysis. Some programs were no longer

available to be retrieved. Other programs were not flexible about the reference genome that can be used,

unless modifying significantly the source code. We also did not test T-lex3 (Bogaerts-Márquez et al. 2020)

since it cannot detect TE insertions present in the sample but not the reference, but only presence/absence of

annotated TE insertions in a reference genome.

We have finally tested 14 programs for which it was possible to use customized reference genomes

(TEMP (Zhuang et al. 2014), TEMP2 (Yu et al. 2021), ngs_te_mapper (Linheiro and Bergman 2012),

ngs_te_mapper2 (Han et al. 2021), PoPoolationTE (Kofler et al. 2012), PoPoolationTE2 (Kofler et al. 2016),

RetroSeq (Keane et al. 2013), RelocaTE (Robb et al. 2013), RelocaTE2 (Chen et al. 2017), TEBreak

(Schauer et al. 2018), TEFLoN (Adrion et al. 2017), TE-locate (Platzer et al. 2012), TEPID (Stuart et al.

2016), and Jitterbug (Hénaff et al. 2015)). The programs are designed to find non-reference insertions (when

compared to a reference genome) and, except Jitterbug, TEBreak and RetroSeq, to find also shared insertions

(between a reference genome and a genome under investigation). The programs have been developed on

particular organisms but sometimes tested on several of them (human, Drosophila, Arabidopsis, rice, mouse

and Daphnia).

Detection of reference insertions

We have first evaluated the capacities of the programs to identify reference insertions, that is to say,

insertions present in the reference genome and in the genome from which the reads are produced. For the *D.*

melanogaster simulated chromosome, it represents 400 insertions and in *A. thaliana*, it represents 3,133

insertions. On Figure 3, the total number of reference insertions found by each program is represented for

each species, independently of the identification of true positives (TP). For TEPID, this number has been

estimated by subtraction since the program provides information about the absence of reference insertions.

As we can see, globally, the increase of coverage has little influence on the total number of reference

insertions detected, except for PopoolationTE and PopoolationTE2, which do not find many insertions at

10X. For both species, the TEMP and TEMP2 programs, which have the same results, find far more

reference insertions than expected. Other programs find more than expected reference insertions but to a

lesser extent in *A. thaliana* (TEPID) and in *D. melanogaster* (TEFLoN for all coverage and PoPoolationTE2

for coverage 50X and 100X). PopoolationTE2, TEFLoN and ngs_te_mapper2 (for *A. thaliana*) and

PopoolationTE, ngs_te_mapper2 and TEPID (for *D. melanogaster)* find a number of reference insertions

close to what is expected. All the other programs find no or few reference insertions.

301

Figure 3: Number of reference insertions detected by each program. On the left panel is represented the number of

reference insertions found for *D. melanogaster* and on the right panel for *A. thaliana*, for the three different read

coverages. The dashed lines correspond to the expected number of reference insertions in each species; * indicate

programs that are not designed to identify reference insertions.

We have then determined among all these insertions the number of False Positives (FP), False

Negatives (FN) and True Positives (TP) in order to compute various metrics to evaluate the programs. TPs

have been identified according to both the capacity of the program to identify the right TE family and

according to the localization prediction with several margin of errors (see Material and Methods). Figure 4

represents the different metrics for both species using a localization prediction with a margin of error of 20

bp (see supplementary figures S1, S2 and S3 for all cutoffs).

Figure 4: Evaluation metrics for the two species for the three read coverages for the reference insertions with a precision localization of 20 bp; * indicate programs that are not designed to identified reference insertions.

We have retained this particular margin of error since at 5 bp all programs do not perform well whereas at 100 bp and 150 bp the efficiency of the programs is not improved. The *recall* metrics indicate the number of good answers among all the possible answers. In our case, it indicates for each program the number of TPs

among all the TE insertions that should be detected. For both species, five programs give the best results for
these metrics: TEMP, TEMP2, PoPoolationTE2 (starting at 50X), ngs_te_mapper2 and TEFLoN, with *recall*
values of more than 0.5. The other programs find few or no TP among all the TE insertions that can be found
given a localization window of 20 bp. The *precision* gives the number of good answers among all the results
proposed by the programs. According to the species, the tools do not have the same results. For *D.*
melanogaster, ngs_te_mapper2 has the best results for these metrics, whereas it is ngs_te_mapper for *A.*
thaliana. In order to take into account both metrics, we have computed the *Fscore*. For both species, five
programs give the best results: ngs_te_mapper2, PoPoolationTE2, TEMP, TEMP2, and TEFLoN. However,
according to the species, the best program is not the same: ngs_te_mapper2 performs better for *D.*
melanogaster when it is PoPoolationTE2 for *A. thaliana*.
We have observed the overlap of TPs between the top four programs for each species (Figure 5). The
results show that 66.8% for *D. melanogaster* and 61.9% for *A. thaliana* of the TPs are found by the four
programs. Among the remaining TPs, a majority is found in common by at least three programs. Only
TEMP/TEMP2 find a significant proportion of unique TPs (3.5% for *D. melanogaster* and 6.1% for *A.*
thaliana).

Figure 5: Percentage overlap of TP reference insertions found by the programs having the best *Fscores*.

<u>*Detection of non-reference insertions*</u>
We have then evaluated the capacity of the programs to find insertions not present in the reference genome.
They correspond to 390 insertions in the simulated *D. melanogaster* chromosome and 3,192 insertions in the
simulated *A. thaliana* chromosome. All programs find a total number of non-reference insertions inferior to
what is expected (Figure 6). The sequence coverage has an impact on the total number of non-reference
insertions found for the majority of the programs. In particular, a coverage of 10X seems to be insufficient
for most programs. Only ngs-te-mapper2 and TEBreak are not very impacted.

Figure 6: Number of non-reference insertions detected by each program. On the left panel is represented the number of non-reference insertions found for *D. melanogaster* and on the right panel for *A. thaliana,* for the three different read coverages. The dashed lines correspond to the expected number of non-reference insertions in each species.

We have then determined among all the non-reference insertions that are detected which ones are TP according to the same rationale presented above and in the material and methods section. Figure 7 represents the different metrics for both species using a correct localization prediction at 20 bp (see supplementary figures S4, S5, and S6 for all cutoffs). Globally, the recall for each program, and for both species, is not very high, meaning that many TPs are missed by the programs. Three of the programs give the best results considering 50X of coverage (TEBreak, PoPoolationTE2 and ngs_te_mapper2). The *precision* metric on the contrary shows that for most programs, TPs are numerous among all the results produced, especially for *D. melanogaster*. The *Fscore* shows similar results between the two species with four programs having the best results: TEBreak, ngs_te_mapper2, PopoolationTE2 (starting at 50X) and RetroSeq (starting at 50X).

Figure 7: Evaluation metrics for the two species for the three read coverages for the non-reference insertions with a precision localization of 20 bp.

The overlapping of the TP detected by the six best programs accounts for only 10.9% of the TPs for *D. melanogaster* and 10.5% of the TP for *A. thaliana* (Figure 8). PopoolationTE2 and ngs_te_mapper2 each identify 6.5% and 4.4% unique TPs in *D. melanogaster*, and 2.1% and 2.9% respectively in *A. thaliana*. It

should be noted that 11.4% of TPs are found by all the programs except TEFLoN in *A. thaliana*. For *D. melanogaster*, 10.6% of TPs are found in common for all programs except TEMP/TEMP2.

Figure 8: Percentage overlap of TP non-reference insertions found by the programs having the best *Fscores*.

Comparison of the characteristics between True Positives and False Negatives

Since we know with accuracy the characteristics of all insertions present in the two reconstructed chromosomes for both species, it is possible to determine whether some of them may have an impact on the fact that an insertion is detected or not by the programs. We have considered the results of the programs having the best *Fscores* for a coverage of 50X and with enough identified TPs considering a localization precision of 20bp to allow statistical analyses without bias.

First, we have considered the reference insertions in both species (Table 1 and Table 2, Wilcoxon tests). The results show that the TPs have significantly smaller sizes than FNs for all programs (expected for ngs_te_mapper2 with *D. melanogaster*). Moreover, the distance to the closest TE insertions is also important since it is significantly larger for TPs when compared to FNs, for all programs and for the two species. Additionally, in *A. thaliana*, the %GC of the flanking regions of TPs are significantly more GC rich than those around FNs. To summarize, the programs better detect reference insertions that are small and largely distant from other TE insertions.

Table 1: characteristics of TPs vs FNs for reference insertions for *D. melanogaster* (400 reference insertions in total)

	ngs_te_mapper2 (279 TPs)		PoPoolationTE2 (307 TPs)		TEFLoN (274 TPs)		TEMP /TEMP2 (339 TPs)	
	TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs
Mean insertion size (bp)	1,126.5	1,800.4	1,096.5	2,105.7	990.6	2,072	1,086.2	2,702.3
	NS		*0.011116*		*1.696e-05*		*8.091e06*	
%divergence to the reference	9.593	11.092	10.070	9.958	10.054	10.02	10.05	10.013
	0.002068		*NS*		*NS*		*NS*	

Mean distance to the nearest insertions (bp)		31,427	21,329	32,127.3	15,918	30,561.1	23,630	30,332.1	17,416
		1.627e-06		*2.254e-10*		*0.0008189*		*5.21e-05*	
%GC of flanking regions	5'	40.29	40.81	40.39	40.57	40.40	40.54	40.57	40.03
		NS		*NS*		*NS*		*NS*	
	3'	40.57	40.45	40.68	39.82	40.61	39.65	40.81	39.76
		NS		*0.02122*		*NS*		*NS*	
Mean TSD size (bp)		3.62	3.967	3.713	3.761	3.752	3.664	3.735	3.667
		NS		*NS*		*NS*		*NS*	

Table 2: characteristics of TPs vs FNs for reference insertions for *A. thaliana* (3,133 reference insertions in total)

		ngs_te_mapper2 (2,019 TPs)		PoPoolationTE2 (2,191 TPs)		TEFLoN (1,799 TPs)		TEMP /TEMP2 (2,506 TPs)	
		TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs
Mean insertion size (bp)		978	1894.7	1043	1917	950.7	1781.8	1013	2465
		<2.2e-16		*<2.2e-16*		*<2.2e-16*		*<2.2e-16*	
%divergence to the reference		9.70	10.33	9.91	9.93	9.81	10.1	9.88	10.1
		1.435e-05		*NS*		*NS*		*NS*	
Mean distance to the nearest insertions (bp)		5,684	2,455.5	5,494	2,296.3	5,525	3,202.5	5,156	2,059
		<2.2e-16		*<2.2e-16*		*<2.2e-16*		*<2.2e-16*	
%GC of flanking regions	5'	34.91	34.15	34.97	33.96	34.93	34.35	34.74	34.58
		0.04939		*0.006899*		*0.005606*		*NS*	
	3'	35.14	33.89	35.11	33.59	35.09	34.28	34.90	33.85
		5.771e-06		*2.559e-06*		*0.002174*		*0.03816*	
Mean TSD size (bp)		4.02	4.17	4.08	4.08	4.24	3.86	4.08	4.05
		NS		*NS*		*0.0003743*		*NS*	

In the case of the non-reference insertions (Table 3 and Table 4, Wilcoxon tests), the results show slightly different characteristics. For all species and almost all programs, the percentage of divergence of TPs compared to its ancestral sequence is significantly lower than for the FNs. Again, the distance of TPs to the closest TEs is larger than for the FNs, especially for *A. thaliana* but also for *D. melanogaster* for three programs (TEFLoN, RetroSeq, and TEBreak). Also, the size of TSD is significantly larger for TPs than for FNs for both species and for most of the programs. Finally, in *A. thaliana,* the %GC of the flanking regions of TPs are significantly more GC rich than those around FNs. To summarize, the programs better detect non-reference insertions that are not too divergent from the consensus TE used to identify them (so likely to be recent insertions), largely distant from other TE insertions and with specific TSD size.

Table 3: characteristics of TPs vs FNs for non-reference insertions for *D. melanogaster* (390 non-reference insertions in total)

		ngs_te_mapper2 (224 TPs)		PoPoolationTE2 (257 TPs)		TEFLoN (135 TPs)		TEMP2 (119 TPs)		RetroSeq (186 TPs)		TEBreak (248 TPs)	
		TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs
Mean insertion size (bp)		1,257.8	1,518	1,439	1,230.5	636.9	1,756.6	1,503	1,308.5	1,061.1	1,649	1,277.4	1,527
		NS		*NS*		*3.875e-08*		*NS*		*0.01601*		*NS*	
%divergence to		9.22	11.12	9.84	10.387	10.12	9.975	7.11	11.31	9.47	10.53	9.740	10.53

the reference			1				2	1		7		
	4.227e-07		NS		NS		< 2.2e-16		0.003103		0.02933	
Mean distance to the nearest insertions (bp)	27,946	26,612	27,967	26,238	30,095	25,937	28,538	26,869	30,054	24,930	30,147	22,513
	NS		NS		0.03794		NS		0.0136		0.001241	
%GC of flanking regions — 5'	40.59	40.50	40.67	40.29	40.70	40.48	40.74	40.47	40.72	40.39	40.71	40.23
	NS		NS		NS		NS		NS		NS	
%GC of flanking regions — 3'	40.15	40.80	40.38	40.50	40.21	40.53	40.19	40.52	40.25	40.58	40.37	40.50
	NS		NS		NS		NS		NS		NS	
Mean TSD size (bp)	4.045	3.03	3.642	3.561	3.956	3.433	3.899	3.489	3.774	3.468	3.903	3.106
	1.63e-06		NS		0.002307		NS		0.04023		5.961e-05	

Table 4: characteristics of TPs vs FNs for non-reference insertions for *A. thaliana* (3,192 non-reference insertions in total)

	ngs_te_mapper2 (1,564 TP)		PoPoolationTE2 (1,558 TP)		TEFLoN (740 TP)		TEMP2 (826 TP)		RetroSeq (1,323 TP)		TEBreak (1,684 TP)	
	TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs	TPs	FNs
Mean insertion size (bp)	1,358.1	1,326.8	1,351.1	1,334	1,187	1,388.9	1,272.4	1,366.5	1,313.0	1,363	1,370.4	1,310.6
	NS		NS		NS		NS		NS		NS	
%divergence to the reference	9.589	10.43	9.668	10.355	10.101	9.995	7.149	11.022	9.53	10.37	9.765	10.304
	5.138e-10		2.619e-07		NS		< 2.2e-16		6.297e-10		7.778e-05	
Mean distance to the nearest insertions (bp)	5,498	3,853.5	5,682	3,684.3	5,402	4,435	5,280	4,442	5,345	4,173	5,366	3,870
	< 2.2e-16		< 2.2e-16		8.975e-09		3.313e-09		< 2.2e-16		< 2.2e-16	
%GC of flanking regions — 5'	35.43	34.03	35.44	34.04	35.43	34.61	35.36	34.61	35.61	34.12	35.47	33.84
	3.734e-12		5.535e-10		0.0001875		0.004747		4.639e-14		1.087e-15	
%GC of flanking regions — 3'	35.23	34.24	35.14	34.32	35.18	34.60	35.27	34.54	35.20	34.41	35.19	34.23
	1.352e-07		6.72e-06		0.002374		0.0003581		1.399e-06		1.81e-07	
Mean TSD size (bp)	4.503	3.568	4.15	3.907	4.789	3.795	4.063	4.013	4.044	4.013	4.404	3.603
	< 2.2e-16		0.01874		< 2.2e-16		NS		NS		7.709e-15	

Application case: detection of endogenous retroviruses polymorphic insertions in real cattle population data

Although a comprehensive understanding of TEs could have an agricultural interest in improving animal breeding, few TE studies have been conducted on livestock species and more particularly on cattle. We have decided to use cattle as a mammalian genome example to study a subpart of the TEs, the endogenous retroviruses (ERV) insertions. We propose hereafter a workflow to perform such an analysis.

 Find the best configuration using simulated data

The study of simulated _D. melanogaster_ and _A. thaliana_ chromosomes has shown that the performance of

the programs to detect polymorphic TE insertions are different depending on the studied species. In order to

choose the best tool to use, the same pipeline as before has been applied to _Bos taurus_ to further detect

polymorphic insertions in short-read data. Two simulated chromosomes were generated using _ReplicaTE_

from chromosome 25. In this chromosome, 899 random CDS sequences were extracted and 900 intergenic

regions were generated. The obtained "complete" simulated chromosome is 25,638,271 bp long including

936 ERVs whereas the "deleted" simulated chromosome contains 474 ERVs. The "deleted" simulated

chromosome has been used as a reference and the "complete" simulated chromosome has been used to

generate simulated short reads in order to evaluate the capacity of the programs to detect the 462 reference

insertions and 474 non-reference insertions. We have determined among the detected insertions the number

of False Positives (FPs), False Negatives (FNs) and True positives (TPs) to compute the same metrics as for

D. melanogaster and _A. thaliana_ (see material and methods section).

Figure 9 represents the _Fscore_ metric for the detection of ERV reference and non-reference

insertions using each of the tested programs (see supplementary figure S7 for _recall_ and _precision_ metrics).

Similar results are found in cattle compared to the other species but the best programs slightly differ. For the

reference insertions, TEMP2 and TEFLoN give the best results with a _Fscore_ higher than 0.80. For the non-

reference insertions, TEFLoN and TEBreak are the two programs giving the best results with respectively a

Fscore of 0.82 and 0.66. In conclusion TEFLoN appears to be the best performing tool to use on _B. taurus_

data.

Figure 9: **Performance evaluation of McClintock programs on *Bos taurus* simulated data**. The detection of the reference and non-reference insertions is represented in the upper and lower panels respectively. INT and LTR consensus were provided separately. The performance of the programs has been evaluated with the *Fscore* metric.

Repbase consensus sequences are largely used for TE annotation using RepeatMasker. For LTR-
retrotransposons, the LTR sequences and the internal part are often split into two separate sequences.
Different types of input sequences have been evaluated to detect ERV insertions in *B. taurus* simulated data
using TEFLoN (Figure 10A): i) only the LTR sequences, ii) only the internal sequences, iii) the LTR and
internal sequences separately, iv) the LTR and internal sequences concatenated for each ERV family
sequence, and v) the LTR, the internal and the concatenated family sequences together to test redundancy.
Figure 10B represents the *Fscore* metric for each input sequence (see supplementary figure S8 for *recall* and
precision metrics). The use of the internal part alone is not working well contrary to other configurations
involving both LTR and internal parts. The use of internal and LTR parts separately gives satisfying results
for reference insertions but is less efficient for non-reference insertions detection. The input giving the best
results is the one with the concatenated family sequences.
We have used four ERV families to generate the simulated chromosomes from ERV class I and II
clades. The figure 10C shows how the different ERV families have been detected by TEFLoN. ~~Reference~~
~~ERVs insertions from class I families are better recognized than class II but it is not the case for new~~

~~insertions.~~ Each family seems to have its own detection characteristics that might correspond to sequence

characteristics identified for *D. melanogaster* and *A. thaliana*.

Figure 10: **Impact of ERV input consensus sequences and ERV families on insertion detection with TEFLoN on**
Bos taurus **simulated data.** A) ERV copy genomic structure and the different input consensus sequences tested on
cattle simulated data to detect ERV copies, B) *Fscores* (Performance) of TEFLoN with the different input consensus
sequences, C) *Fscores* (Performance) of TEFLoN using the detail of each ERV family when using the consensus
sequences labeled "INT LTR fam". The detection of reference insertions is represented with circles and the detection of
non-reference insertions with triangles.

Detection of insertion polymorphism in real population data

We have used the previously selected tool TEFLoN to analyze 10 cattle WGS short-read dataset. The

detected insertions have been compared to the ERV annotation of the reference assembly and to the output of

a variant calling analysis performed on long-read data from the same samples. Figure 11 represents the

Fscores obtained for these samples and the correlation between the tool performance and the sample short-

read depth sequencing. More than 80% of the expected insertions are detected, on average, in the 10 samples.

ERV insertions also present in the reference genome are significantly better recognized than the non-

reference insertions (Wilcoxon test, $p = 1.1e\text{-}05$). Among the insertions common with the reference, almost

no FP are identified. For insertions not present in the reference, almost a hundred of FP are detected

representing from 30 to 40% of the non-reference insertions detected in each sample. The tool performances

are also more homogeneous between the samples for the detection of reference insertions than for the non-

reference ones mainly due the short-read coverage differences across samples. A higher coverage improves

the detection of insertions but also increases the detection of FPs (see supplementary figure S9).

Furthermore, samples with coverage lower than 10X have a drop in detection rates compared to the others. It

appears that 10X is the minimum coverage to reliably detect a sufficient number of ERV insertions. Finally,

the comparison between the analysis on simulated and real data shows better results in detecting reference

insertions in real data compared to simulated data, with median *Fscores* of 0.97 and 0.81 respectively. On the

contrary, TEFLoN is less effective in identifying non-reference insertions in real data compared to simulated

data with median *Fscores* of 0.75 and 0.82 respectively (Figure 11).

542

Figure 11: **Detection of ERV insertions in 10 *Bos taurus* samples with TEFLoN.** A) *Fscores* for the detection of

ERV insertions in the 10 samples. The red dots indicate the *Fscores* obtained with TEFLoN on cattle simulated data, B)

Impact of the short-read depth sequencing on the number of detected insertions. Depth is computed on trimmed reads

mapped on the cattle reference genome.

Discussion

In this work, we have developed an approach to simulate TE insertions from a known biological context. The

data obtained made it possible to test in a reliable and controlled manner 14 programs for the detection of

polymorphic TEs. For the first time in the benchmarking of these approaches, it is possible to show why

certain insertions are better detected than others by the different programs. Especially, reference and non-

reference insertions show different biases. Reference insertions are more correctly detected if they are small

and largely distant from other TE insertions. In the case of non-reference insertions, they need to be not too

divergent from the consensus or reference TE used to identify them, very distant from other TE insertions

and with specific TSD size.

Generally, full data simulation approaches are often used to test polymorphic TE detection programs.

They make it possible to perfectly control all the information. The major problem is that often these

simulated data do not completely reflect the biological reality. In order to overcome this problem, we have

proposed here an approach that uses real data as a starting point and simulates sequences using the biological

information of the organism of interest. We thus made the choice to completely simulate the intergenic

regions in order to free ourselves from possible bad TE annotations. However, these intergenic regions are

not completely randomly generated sequences. In particular, the %GC of these sequences must correspond to

what is observed in the analyzed genome. The GC content may be an important factor influencing the

detection since it may be a caveat in steps of mapping (Donato et al. 2021). Similarly, the reinserted

sequences of the TEs are not the true sequences but come from a real insertion representative of each family

contained in the analyzed genome. This allows us to control not only the position of the insertions but also to

know with accuracy other information that may play a role on whether an insertion is detected or not. Thus,

among the parameters which are controlled, the size of the insertions, the sequence divergence with respect

to the reference element, the distance to the closest TEs and the size of TSD are perfectly known for each

insertion. It thus allows us to shed light on precise sequence characteristics rather than limiting tests on

specific types of TEs, which is sometimes an approach used to benchmark TE polymorphic tools (Nelson et

al. 2017; Vendrell-Mir et al. 2019; Chen et al. 2023). Our approach seems to be a good compromise between

the use of complete simulation and real but partial biological data with either consensus TE sequences or

using only a small set of TE families ~~representing a small portion of a chromosome~~. However, a number of

improvements can be considered with our approach. Currently, only one chromosome is simulated. It could

be interesting to simulate several chromosomes and in particular, to generate populations of chromosomes in

order to mimic what can be observed in a natural population. Additionally, the tool is currently limited

regarding the number of TE insertions that can be inserted. Thus, for a human chromosome for example, the

tool works only with a limited number of TE families. The input format of the reference chromosome could

be modified to support bed annotation files along with a fasta file containing the chromosome sequence,

rather than one genbank file. However, in any case, TE annotations for the reference species are mandatory

to allow the different programs to be used to identify reference insertions.

Our results show that all the programs tested here are far from obtaining results as good as

announced in their original publication. For some of them, read coverage strongly impacts the ability to find

non-reference insertions, as has already been shown (Rishishwar et al. 2017; Vendrell-Mir et al. 2019) but

this is only true up to 50X coverage from which a plateau is reached. Moreover, the results are not as good

whether we are

interested in the reference insertions (present in the reference genome) or the non-reference insertions

(present only in the read samples). Indeed, non-reference insertions are less correctly detected than the

reference insertions, an observation that was also made by the only other benchmark evaluating non-

reference insertion detection (Vendrell-Mir et al. 2019). Globally, the *Fscores* are better in the first case.

However, the values obtained for the best programs do not indicate exceptional performance. Indeed, for

reference insertions, the best programs ngs-te mapper2 and popoolationTE2 have *Fscores* below 0.8. The

other programs (PopoolationTE2, TEFLoN, TEMP, and TEMP2) show values around 0.6. For non-reference

insertions, the best programs (TEBreak, ngs_te_mapper2, popoolationTE2 and RetroSeq) have values
hovering around 0.6. It is important to note that some programs are more successful in finding reference
insertions than non-reference insertions, and *vice versa*. TEFLoN, TEMP and TEMP2 show poorer
performance in finding non-reference insertions compared to reference insertions. Overall ngs_te_mapper2
and PoPoolationTE2 give consistent results for the two types of insertions. If we compare the results for the
two species, there are some notable differences for the detection of reference insertions. Ngs-te-mapper2
gives better results with *D. melanogaster* while the best program is PoPoolationTE2 (at 50X and 100X) for
A. thaliana. In the case of non-reference insertions, all programs give comparable results for the two species,
although working a little less well in the case of *A. thaliana*. It is to note that TEBreak was proposed as the
best performing tool to identify non-reference insertions in yeast genomes (Chen et al. 2023), which
indicates that the choice of the best performing tool needs to be assessed according to the species under
study.

Given that the programs produce many false positives (FPs), an approach allowing to optimize the
identification of the true positives (TPs), in the absence of comparison, is to use several tools at the same
time to retain only the insertions detected in common. This approach has been used for the analysis of many
natural populations of *D. melanogaster* (Lerat et al. 2019). However, the two tools used showed little overlap
in their results. We observed the overlap between the TPs for the best programs identified in this work for
reference insertions and non-reference insertions. The proportion of common insertions correctly found by
all the programs is quite high in the case of reference insertions since it is almost 70% considering four
programs. This proportion is much lower in the case of non-reference insertions with less than 11% for six
programs. However, the proportion reach 48.1% for *D. melanogaster* and 58,5% for *A. thaliana* when
considering only the results common to TEBreak, ngs_te_mapper2 and PoPoolationTE2, the three programs
giving the best results in our benchmark. This remains lower than for the reference insertions. This lack of
overlap among the tools has already been observed in other benchmarks (Nelson et al. 2017; Vendrell-Mir et
al. 2019). Thus, as proposed by Vendrell-Mir et al. (Vendrell-Mir et al. 2019), an approach consisting of
using several tools at the same time to optimize the number of TPs must be limited to a few tools at a time.
Even with this method, it is important to take into account that some information will be inevitably lost and
that the number of polymorphic TE insertions will be underestimated.

With our approach, it was possible to compare the characteristics of the True Positives (TPs)
compared to those of the False Negatives (FNs), *i.e.* the insertions which are missed by the programs, a point
that has never been assessed previously by the other benchmark analyses. The goal was to determine if there
are biases inherent in the sequences preventing their detection. Between reference and non-reference
insertions, some differences appeared. In particular, reference insertions correctly detected tend to be smaller
than those not detected by the programs. That would indicate that degraded or small size types of TEs will be
better detected as reference insertions. This observation is consistent with the fact that MITE reference
elements were better identified than LTR-retrotransposon reference elements (Vendrell-Mir et al. 2019) since
MITE elements are shorter than LTR-retrotransposons. On the contrary, the non-reference insertions are

better detected when their divergence compared to a reference element is low. Then, recent insertions will be

better detected. Although this could be enough to identify recent events, it remains that some of the non-

reference insertions may be ancient. These particular insertions would be missed by the different programs.

In their original manuscript, almost all programs acknowledge the fact that they cannot detect nested

insertions. This is confirmed by our analysis for both types of insertions since TPs present significantly

larger distances to the nearest TEs than FNs. Globally, the same bias appears between the two explored

species. However, for *A. thaliana*, we observed that the GC content of genomic regions surrounding the

insertions also play a role in whether they are detected or not by the program. This species has globally AT

rich intergenic regions (DeRose-Wilson and Gaut 2007). We observed that the insertions are better detected

when the genomic regions are less AT rich. Since TEs are known to be also AT rich (Lerat et al. 2002;

Boissinot 2022), they may be better identified when their base composition is more different from the

surrounding genomic regions. We also observed for the detection of non-reference insertions that the size of

the TSD is important. Since these sequences may not be well conserved, it may prevent the detection of

many insertions.

The case study provided here, focusing on *B. taurus,* allowed us to identify important criteria that

should be considered before performing studies on polymorphic TEs in real population data. The choice of

the program is crucial and depends on the analyzed species. Indeed, the best identified tool to use on this

species is not the same as for *D. melanogaster* and *A. thaliana*. Therefore, it is essential to first perform tests

on simulated data built with specific elements from the species of interest to identify the most suitable

tool(s) to use. The different programs were all used through the McClintock pipeline (Nelson et al. 2017;

Chen et al. 2023) which has a significant advantage to allow the use of multiple tools simultaneously,

prevents difficulties in program installation and ensures standardized results. It is also important to carefully

select the type of consensus sequences, especially for LTR-retrotransposons. For these elements, usually the

LTR and the internal parts are separated in distinct consensus sequences. The re-association of the LTR

sequences and the internal parts of a given family is thus necessary and require an in-depth annotation of the

reference genome.

Here, we demonstrated the importance of testing a tool also on real data before launching a large-

scale population analysis. Even though our study was limited to 10 samples, the genomic characteristics and

TE content reflected the reality. The results obtained on real data were different compared to the simulated

data, with a better detection of the reference insertions but a less effective identification of the non-reference

insertions. This difference is mainly due to the total number of ERV insertions. In the simulated data, half of

the total insertions were insertions not present in the reference, whereas they constituted approximately 2%

of the insertions in the real data. It appears that detecting non-reference insertions is easier when they

represent a larger fraction of the genome of interest.

We showed that non-reference insertions were overall more challenging to detect than the reference

ones. Moreover, assessing insertions absent from the reference genome in real samples is challenging

because we do not know what to expect, making it difficult to determine whether an insertion is a true or

false positive. In our analysis, we used variant calling results obtained from long-reads sequencing data.

However, this approach might also miss some insertions, raising questions about its reliability as a reference.

Nevertheless, it provides results from two distinct methodologies, ensuring the identification of TPs, even if

some are missed.

In conclusion, most of the tested tools do not achieve extraordinary results. There are several biases

that prevent them from detecting certain insertions. In addition, the FP rate is particularly high for some

tools. Therefore, it is advisable to use a small number of programs simultaneously to optimize the detection

of real insertions while keeping a critical perspective on the results.

Acknowledgments

This work was performed using the computing facilities of the CC LBBE/PRABI and of the IFB-cloud. We

thank the SeqOccin project and the Get-Plage platform (https://get.genotoul.fr/) for sharing the bovine data

set, and Mekki Boussaha (G2B team, INRAE Jouy-en-Josas) for sharing the associated alignments computed

by his team. We thank Carole Lampietro, Claire Kuchly and Caroline Vernette (Get-Plage) for their help

with the submission of the sequencing data to SRA. We thank Caroline Leroux (IVPC) and Vincent Navratil

(PRABI-Doua) for useful discussions about this work. JT and TF were supported by the INRAE

GoatRetrovirome grant for this project. MV PhD fellowship was funded by ANR, grant ANR-22-CE35-

0002-01.

References

Adrion JR, Song MJ, Schrider DR, et al (2017) Genome-Wide Estimates of Transposable Element Insertion

and Deletion Rates in Drosophila Melanogaster. Genome Biology and Evolution 9:1329–1340.

https://doi.org/10.1093/gbe/evx050

Bergman CM, Quesneville H (2007) Discovering and detecting transposable elements in genome sequences.

Briefings in Bioinformatics 8:382–392. https://doi.org/10.1093/bib/bbm048

Bogaerts-Márquez M, Barrón MG, Fiston-Lavier A-S, et al (2020) T-lex3: an accurate tool to genotype and

estimate population frequencies of transposable elements using the latest short-read whole genome

sequencing data. Bioinformatics 36:1191–1197. https://doi.org/10.1093/bioinformatics/btz727

Boissinot S (2022) On the Base Composition of Transposable Elements. Int J Mol Sci 23:4755.

https://doi.org/10.3390/ijms23094755

Bourque G, Burns KH, Gehring M, et al (2018) Ten things you should know about transposable elements.

Genome Biology 19:199. https://doi.org/10.1186/s13059-018-1577-z

Chen J, Basting PJ, Han S, et al (2023) Reproducible evaluation of transposable element detectors with

McClintock 2 guides accurate inference of Ty insertion patterns in yeast. Mobile DNA 14:8.

https://doi-org/10.1186/s13100-023-00296-4

Chen J, Wrightsman TR, Wessler SR, Stajich JE (2017) RelocaTE2: a high resolution transposable element

insertion site mapping tool for population resequencing. PeerJ 5:e2942.

https://doi.org/10.7717/peerj.2942

Cordaux R, Batzer MA (2009) The impact of retrotransposons on human genome evolution. Nature reviews Genetics 10:691–703. https://doi.org/10.1038/nrg2640

DeRose-Wilson LJ, Gaut BS (2007) Transcription-related mutations and GC content drive variation in nucleotide substitution rates across the genomes of Arabidopsis thaliana and Arabidopsis lyrata. BMC Evol Biol 7:1–12. https://doi.org/10.1186/1471-2148-7-66

Di Stefano L (2022) All Quiet on the TE Front? The Role of Chromatin in Transposable Element Silencing. Cells 11:2501. https://doi.org/10.3390/cells11162501

Donato L, Scimone C, Rinaldi C, et al (2021) New evaluation methods of read mapping by 17 aligners on simulated and empirical NGS data: an updated comparison of DNA- and RNA-Seq data from Illumina and Ion Torrent technologies. Neural Comput & Applic 33:15669–15692. https://doi.org/10.1007/s00521-021-06188-z

Ewing AD (2015) Transposable element detection from whole genome sequence data. Mobile DNA 6:24. https://doi.org/10.1186/s13100-015-0055-3

Han S, Basting PJ, Dias GB, et al (2021) Transposable element profiles reveal cell line identity and loss of heterozygosity in Drosophila cell culture. Genetics 219:iyab113. https://doi.org/10.1093/genetics/iyab113

Hénaff E, Zapata L, Casacuberta JM, Ossowski S (2015) Jitterbug: somatic and germline transposon insertion detection at single-nucleotide resolution. BMC Genomics 16:768. https://doi.org/10.1186/s12864-015-1975-5

Huang W, Li L, Myers JR, Marth GT (2012) ART: a next-generation sequencing read simulator. Bioinformatics 28:593–594. https://doi.org/10.1093/bioinformatics/btr708

Keane TM, Wong K, Adams DJ (2013) RetroSeq: Transposable element discovery from next-generation sequencing data. Bioinformatics 29:389–390. https://doi.org/10.1093/bioinformatics/bts697

Kobayashi K, Nakahori Y, Miyake M, et al (1998) An ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy. Nature 394:388–392. https://doi.org/10.1038/28653

Kofler R, Betancourt AJ, Schlötterer C (2012) Sequencing of pooled DNA samples (Pool-Seq) uncovers complex dynamics of transposable element insertions in Drosophila melanogaster. PLoS Genetics 8:. https://doi.org/10.1371/journal.pgen.1002487

Kofler R, Gómez-Sánchez D, Schlötterer C (2016) PoPoolationTE2: Comparative Population Genomics of Transposable Elements Using Pool-Seq. Molecular Biology and Evolution 33:2759–2764. https://doi.org/10.1093/molbev/msw137

Lerat E (2019) Repeat in Genomes: How and Why You Should Consider Them in Genome Analyses? In: Ranganathan S, Nakai K, Schönbach C, Gribskov M (eds) Encyclopedia of Bioinformatics and Computational Biology. Elsevier Inc., pp 210–220

Lerat E, Capy P, Biémont C (2002) Codon usage by transposable elements and their host genes in five species. Journal of Molecular Evolution 54:625–637. https://doi.org/10.1007/s00239-001-0059-0

Lerat E, Goubert C, Guirao-Rico S, et al (2019) Population-specific dynamics and selection patterns of

transposable element insertions in European natural populations. Mol Ecol 28:1506–1522.

https://doi.org/10.1111/mec.14963

Linheiro RS, Bergman CM (2012) Whole genome resequencing reveals natural target site preferences of

transposable elements in Drosophila melanogaster. PLoS ONE 7:.

https://doi.org/10.1371/journal.pone.0030008

Nelson MG, Linheiro RS, Bergman CM (2017) McClintock: An Integrated Pipeline for Detecting

Transposable Element Insertions in Whole-Genome Shotgun Sequencing Data. G3 (Bethesda)

7:2763–2778. https://doi.org/10.1534/g3.117.043893

Platzer A, Nizhynska V, Long Q (2012) TE-Locate: A Tool to Locate and Group Transposable Element

Occurrences Using Paired-End Next-Generation Sequencing Data. Biology 1:395–410.

https://doi.org/10.3390/biology1020395

R Core Team (2017) R: A Language and Environment for Statistical Computing. https://www.r-project.org

Rishishwar L, Mariño-Ramírez L, Jordan IK (2017) Benchmarking computational tools for polymorphic

transposable element detection. Brief Bioinform 18:908–918. https://doi.org/10.1093/bib/bbw072

Robb SMC, Lu L, Valencia E, et al (2013) The use of RelocaTE and unassembled short reads to produce

high-resolution snapshots of transposable element generated diversity in rice. G3 (Bethesda, Md)

3:949–57. https://doi.org/10.1534/g3.112.005348

Schauer SN, Carreira PE, Shukla R, et al (2018) L1 retrotransposition is a common feature of mammalian

hepatocarcinogenesis. Genome Res 28:639–653. https://doi.org/10.1101/gr.226993.117

Schnable PS, Ware D, Fulton RS, et al (2009) The B73 maize genome: complexity, diversity, and dynamics.

Science (New York, NY) 326:1112–5. https://doi.org/10.1126/science.1178534

Stuart T, Eichten SR, Cahn J, et al (2016) Population scale mapping of transposable element diversity reveals

links to gene regulation and epigenomic variation. eLife 5:. https://doi.org/10.7554/eLife.20777

Vendrell-Mir P, Barteri F, Merenciano M, et al (2019) A benchmark of transposon insertion detection tools

using real data. Mob DNA 10:53. https://doi.org/10.1186/s13100-019-0197-9

Wang Y, McNeil P, Abdulazeez R, et al (2023) Variation in mutation, recombination, and transposition rates

in Drosophila melanogaster and Drosophila simulans. Genome Res gr.277383.122.

https://doi.org/10.1101/gr.277383.122

Weinstock GM, Robinson GE, Gibbs RA, et al (2006) Insights into social insects from the genome of the

honeybee Apis mellifera. Nature 443:931–949. https://doi.org/10.1038/nature05260

Wicker T, Sabot F, Hua-Van A, et al (2007) A unified classification system for eukaryotic transposable

elements. Nature reviews Genetics 8:973–982. https://doi.org/10.1038/nrg2165-c4

Yu T, Huang X, Dou S, et al (2021) A benchmark and an algorithm for detecting germline transposon

insertions and measuring de novo transposon insertion frequencies. Nucleic Acids Res 49:e44.

https://doi.org/10.1093/nar/gkab010

Zhao Y, Li X, Xie J, et al (2022) Transposable Elements: Distribution, Polymorphism, and Climate

Adaptation in Populus. Front Plant Sci 13:814718. https://doi.org/10.3389/fpls.2022.814718

Zhuang J, Wang J, Theurkauf W, Weng Z (2014) TEMP: A computational method for analyzing

transposable element polymorphism in populations. Nucleic Acids Research 42:6826–6838.

https://doi.org/10.1093/nar/gku323

Supplementary data

Supplementary files: Output files produced by *ReplicaTE* on the two species *D. melanogaster* and *A.*
thaliana.

Figure S1: Recall metrics for *D. melanogaster* and *A. thaliana* for the three read coverages for the reference
insertions according to the different precision localization tested.

Figure S2: Precision metrics for *D. melanogaster* and *A. thaliana* for the three read coverages for the
reference insertions according to the different precision localization tested.

Figure S3: *Fscore* metrics for *D. melanogaster* and *A. thaliana* for the three read coverages for the reference
insertions according to the different precision localization tested.

Figure S4: Recall metrics for *D. melanogaster* and *A. thaliana* for the three read coverages for the non-
reference insertions according to the different precision localization tested.

Figure S5: Precision metrics for *D. melanogaster* and *A. thaliana* for the three read coverages for the non-
reference insertions according to the different precision localization tested.

Figure S6: *Fscore* metrics for *D. melanogaster* and *A. thaliana* for the three read coverages for the non-
reference insertions according to the different precision localization tested.

Figure S7: Recall and precision metrics of the different tested programs on *Bos taurus* simulated data.
803

804 **Figure S8:** Impact of the structure of the ERV input consensus sequences (panel A) and ERV families (panel
805 B) on recall and precision metrics using TEFLoN on *Bos taurus* simulated data.
806 The detection of reference insertions is represented with circles and the detection of non-reference insertions
807 with triangles.
808

809 **Figure S9**: Performance of TEFLoN and impact of read coverage in the detection of ERV insertions in 10
810 *Bos taurus* samples. A) Recall and precision metrics, B) Number of TP and FP according to the short-read
811 depth sequencing. Depth was computed on trimmed reads mapped on the cattle reference genome.
812

813 **Supplementary Table S1**: Accession numbers of 10 WGS short-read data samples of *Bos taurus*.

Figure 1: Workflow of the simulation tool *ReplicaTE*. Each step corresponds to the different scripts. In gray are indicated the genes, in blue and red are indicated the TE insertions, a given color corresponding to a given TE family. The simulated TE copies are represented in orange and light blue.

Figure 2: Evaluation approach. The black rectangles correspond to genes. The red and blue circles correspond to TE copies from two different families. The reads simulated on the "deleted" simulated genome will be mapped to the "complete" simulated genome, which will allow to identify reference and/or absent TE insertions. The reads simulated on the "complete" simulated genome will be mapped to the "deleted" simulated genome, which will allow to identify non-reference TE insertions.

Figure 3: Number of reference insertions detected by each program. On the left panel is represented the number of reference insertions found for *D. melanogaster* and on the right panel for *A. thaliana*, for the three different read coverages. The dashed lines correspond to the expected number of reference insertions in each species; * indicate programs that are not designed to identify reference insertions.

Figure 4: Evaluation metrics for the two species for the three read coverages for the reference insertions with a precision localization of 20 bp; * indicate programs that are not designed to identified reference insertions.

Figure 5: Percentage overlap of TP reference insertions found by the programs having the best Fscores.

Figure 6: Number of non-reference insertions detected by each program. On the left panel is represented the number of non-reference insertions found for *D. melanogaster* and on the right panel for *A. thaliana,* for the three different read coverages. The dashed lines correspond to the expected number of non-reference insertions in each species.

Figure 7: Evaluation metrics for the two species for the three read coverages for the non-reference insertions with a precision localization of 20 bp.

Figure 8: Percentage overlap of TP non-reference insertions found by the programs having the best *Fscores*.

Figure 9: Performance evaluation of McClintock programs on *Bos taurus* simulated data. The detection of the reference and non-reference insertions is represented in the upper and lower panels respectively. The performance of the programs was evaluated with the Fscore metric.

Figure 10: Impact of ERV input consensus sequences and ERV families on insertion detection with TEFLoN on *Bos taurus* simulated data. A) ERV copy genomic structure and the different input consensus sequences tested on cattle simulated data to detect ERV copies, B) *Fscores* (Performance) of TEFLoN with the different input consensus sequences, C) *Fscores* (Performance) of TEFLoN using the detail of each ERV family when using the consensus sequences labeled "INT LTR fam". The detection of reference insertions is represented with circles and the detection of non-reference insertions with triangles.

Figure 11: Detection of ERV insertions in 10 *Bos taurus* samples with TEFLoN. A) Fscore for the detection of ERV insertions in the 10 samples. The red dots indicate the *Fscore* obtained with TEFLoN on cattle simulated data, B) Impact of the short-read depth sequencing on the number of detected insertions. Depth was computed on trimmed reads mapped on the cattle reference genome.

real genome

Step 1: genomic feature retrieval and intergenic region simulation

Cleaned genes

Simulated intergenic regions

Size and %GC from statistical distributions fitted on the data

TE families

Step 2: simulation of TE families based on real sequences

Step 3: assembly of the simulated genome

simulated TE families

simulated genome

Simulation based on *D. melanogaster*

Simulation based on *A. thaliana*

Left panel:

Y-axis: Number of reference insertions

X-axis: read coverage (10X, 50X, 100X)

Legend:
- ngs_te_mapper
- ngs_te_mapper2
- RelocaTE
- RelocaTE2
- TEMP
- TEMP2
- RetroSeq*
- PoPoolationTE
- PoPoolationTE2
- TE-locate
- TEFLoN
- TEBreak*
- TEPID

Right panel:

Y-axis: Number of reference insertions

X-axis: read coverage (10X, 50X, 100X)

Legend:
- ngs_te_mapper
- ngs_te_mapper2
- RelocaTE
- RelocaTE2
- TEMP
- TEMP
- RetroSeq*
- PoPoolationTE
- PoPoolationTE2
- TE-locate
- TEFLoN
- TEBreak*
- TEPID

Simulation based on *D. melanogaster* — **Simulation based on *A. thaliana***

Exact position at 20 bp (Recall, Precision, Fscore panels)

Legend: ngs_te_mapper, ngs_te_mapper2, RelocaTE, RelocaTE2, TEMP, TEMP2, RetroSeq*, PoPoolationTE, PoPoolationTE2, TE-locate, TEFLoN, TEBreak*

Simulation based on *D. melanogaster*   **Simulation based on *A. thaliana***

Legend (both panels):
- ngs_te_mapper
- ngs_te_mapper2
- RelocaTE
- RelocaTE2
- TEMP
- TEMP2
- RetroSeq
- PoPoolationTE
- PoPoolationTE2
- TE-locate
- TEFLoN
- TEBreak
- TEPID
- Jitterbug

Simulation based on *D. melanogaster*

Exact position at 20 bp

Exact position at 20 bp

Exact position at 20 bp

Simulation based on *A. thaliana*

Exact position at 20 bp

Exact position at 20 bp

Exact position at 20 bp