1 **Title:** CCR5 antagonists as cerebroprotective and stroke recovery enhancing agents: a preclinical
2 systematic review and meta-analysis
3
4 **Abbreviated Title:** CCR5, cerebroprotection, and stroke recovery
5

6 **Authors:** Ayni Sharif[1,2,*], Matthew S. Jeffers[1,2,*], Dean A. Fergusson[1,2,3], Raj Bapuji[1], Stuart G.
7 Nicholls[4], John Humphrey[5], Warren Johnston[5], Ed Mitchell[5], Mary-Ann Speirs[5], Laura
8 Stronghill[5], Michele Vuckovic[5], Susan Wulf[5], Risa Shorr[6], Dar Dowlatshahi[2,7,8], Dale Corbett[8],
9 Manoj M. Lalu[1,2,8,9,#]
10
Affiliations: [1]Clinical Epidemiology Program, Blueprint Translational Research Group, Ottawa
Hospital Research Institute, Ottawa, Canada
[2]School of Epidemiology and Public Health, University of Ottawa, Ottawa, Canada
[3]Department of Medicine, University of Ottawa, Ottawa, Canada
[4]Office for Patient Engagement in Research Activity (OPERA), Ottawa Hospital Research
Institute, Ottawa, Canada
[5]Patient Partner, Blueprint Translational Research Group, Ottawa Hospital Research Institute,
Ottawa, Canada
[6]Ottawa Hospital Library Services, The Ottawa Hospital, Ottawa, Canada
[7]Division of Neurology, Department of Medicine, The Ottawa Hospital, Ottawa, Canada
[8]Department of Cellular and Molecular Medicine, Faculty of Medicine, University of Ottawa,
Ottawa, Canada
[9]Department of Anaesthesiology and Pain Medicine, The Ottawa Hospital, Ottawa, Canada
*These authors contributed equally
25
26 **Corresponding Author:**
27 [#]Manoj M. Lalu, mlalu@toh.ca
28 Department of Cellular and Molecular Medicine, University of Ottawa
29 Clinical Epidemiology and Regenerative Medicine Programs, Ottawa Hospital Research Institute
30 Ottawa, Ontario, Canada, K1Y 4E9
31
32 **X:** @ManojLalu

33
34 **Number of Figures:** 3
35 **Number of Tables:** 3
36 **Word counts:** Abstract (240), Main text (3,785)
37

ABSTRACT

Background: C-C chemokine receptor type 5 (CCR5) antagonists may improve both acute stroke outcome and long-term recovery. Despite their evaluation in ongoing clinical trials, gaps remain in the evidence supporting their use.

Methods: With a panel of patients with lived experiences of stroke, we performed a systematic review of animal models of stroke that administered a CCR5 antagonist and assessed infarct size or behavioural outcomes. MEDLINE, Web of Science, and Embase were searched. Article screening and data extraction were completed in duplicate. We pooled outcomes using random effects meta-analyses. We assessed risk of bias using the Systematic Review Centre for Laboratory Animal Experimentation (SYRCLE) tool and alignment with the Stroke Treatment Academic Industry Roundtable (STAIR) and Stroke Recovery and Rehabilitation Roundtable (SRRR) recommendations.

Results: Five studies representing 10 experiments were included. CCR5 antagonists reduced infarct volume (standard mean difference −1.02; 95% confidence interval −1.58 to −0.46) when compared to stroke-only controls. Varied timing of CCR5 administration (pre- or post-stroke induction) produced similar benefit. CCR5 antagonists significantly improved 11 of 16 behavioural outcomes reported. High risk of bias was present in all studies and critical knowledge gaps in the preclinical evidence were identified using STAIR/SRRR.

Conclusions: CCR5 antagonists demonstrate promise; however, rigorously designed preclinical studies that better align with STAIR/SRRR recommendations and downstream clinical trials are warranted.

Registration: Prospective Register of Systematic Reviews (PROSPERO CRD42023393438)

Key Words: stroke; neuroprotection; impairment; recovery; preclinical; CCR5 antagonists; maraviroc; patient engagement in research; CAMAROS; cerebroprotection

Non-standard Abbreviations and Acronyms:

CAMAROS	Canadian Maraviroc RCT to Augment Rehabilitation Outcomes After Stroke
CCR5	C-C chemokine receptor type 5
DAPTA	D-Ala-Peptide T-Amide
SRRR	Stroke Recovery and Rehabilitation Roundtable
STAIR	Stroke Treatment Academic Industry Roundtable
SYRCLE	Systematic Review Centre for Laboratory Animal Experimentation
TAK-779	Takeda 779

Introduction

C-C chemokine receptor type 5 (CCR5) is expressed across a variety of leukocyte subtypes, endothelial cells, and cell types in the brain (e.g., neurons, microglia, astrocytes), and is thought to play crucial roles in post-stroke neuroinflammation, blood-brain barrier repair, and neuronal survival / repair processes.[1] CCR5 antagonists have emerged as potential therapeutic candidates for stroke, demonstrating both cerebroprotection and improved neural repair/recovery in preclinical animal models.[2–6] However, no CCR5 antagonist drug has an approved indication in the stroke context, necessitating studies to establish safety and efficacy this population. This has led to an ongoing clinical trial to investigate efficacy of CCR5 antagonists in combination with post-stroke rehabilitation.[7] Assessment of the preclinical evidence supporting CCR5's role in stroke is needed to identify areas of potential benefit, and knowledge gaps, that should be addressed by future preclinical research.[8]

The stroke cerebroprotection and recovery communities have advocated for alignment of preclinical and clinical study parameters through publication of consensus recommendations for preclinical research, in an effort to enhance the translation of new stroke therapies.[9–11] Examples include identification of more sensitive and clinically relevant preclinical outcome measures and incorporation of potentially important effect modifiers of treatment efficacy, such as age, sex, and stroke-related comorbidities (hypertension, diabetes, etc.).[9–11] These recommendations aim to improve the translation of novel stroke therapeutics from preclinical to clinical populations, but the degree to which preclinical evidence for CCR5 antagonists satisfy these recommendations is unknown.

We sought to comprehensively evaluate the preclinical evidence for CCR5 antagonist drugs as both cerebroprotective and stroke recovery-promoting agents.[9,11] Both perspectives are necessary to fully understand the therapeutic potential of stroke-related treatments, as distinct biological principles, time windows for treatment, and outcomes of interest underpin each of these treatment domains.[12,13] We conducted a systematic review and meta-analysis of the CCR5 literature in conjunction with a panel of individuals with lived experiences of stroke. This review sought to explore how the preclinical evidence for CCR5 antagonist drugs aligned with guidance for preclinical stroke research provided by previous expert committees and the parameters of an

ongoing clinical trial (The Canadian Maraviroc Randomized Controlled Trial To Augment

Rehabilitation Outcomes After Stroke, CAMAROS, NCT04789616).[9–11]

Materials and Methods

We registered the review protocol on the International Prospective Register of Systematic

Reviews (CRD42023393438).[14] The findings are reported in accordance with the Preferred

Reporting Items for Systematic Reviews and Meta-Analyses (Table S1)[15] and Guidance for

Reporting the Involvement of Patients and the Public (Table S2).[16] A panel of eight patients and

caregivers with lived and living experience of stroke informed project development and were

actively involved throughout research conduct (see Table S2 for more information).

Eligibility criteria

- *Animals:* Any preclinical *in vivo* animal models of adult stroke were included. Human,

invertebrate, *in vitro*, *ex vivo*, and neonatal animal studies were excluded.

- *Model:* Focal ischaemic or intracerebral haemorrhagic stroke models were included,

while animal models of global ischaemia were excluded.

- *Intervention:* Studies administering a CCR5 antagonist drug (e.g., maraviroc, D-Ala-

Peptide T-Amide (DAPTA), Takeda 779 (TAK-779)). Study arms in which CCR5 was

genetically manipulated (e.g., CCR5 knockout strain) were excluded.

- *Comparator:* Vehicle-treated control groups where stroke was induced. CCR5 antagonist

control groups without stroke were excluded.

- *Outcome:* Studies reporting at least one of the following: infarct size, behavioural tests,

mortality, adverse events, and spasticity were included.

- *Study design and publication characteristics:* Controlled interventional studies

(randomized, pseudo-randomized, or non-randomized) published as full journal articles

in any language or year were included. Abstracts, review articles, opinion-based

letters/editorials, and unpublished grey literature were excluded.

Information sources and search strategy

An information specialist with experience in systematic searches of the preclinical literature

developed a comprehensive search strategy based on a previously published strategy for

139 identifying animal experimentation studies.[17] The search strategy underwent peer-review using

140 the Peer Review of Electronic Search Strategies (PRESS) checklist.[18] We searched MEDLINE

141 (OVID interface, including In-Process and Epub Ahead of Print), Web of Science, and Embase

142 (OVID interface). The search was conducted October 25, 2022, to align with the listed launch

143 date of the CAMAROS trial (September 15, 2022). Our intention in doing so was to collate and

144 assess all preclinical evidence that could have feasibly informed the clinical trial. We sought to

145 assess the comprehensiveness of evidence and readiness for translation of CCR5 antagonist

146 drugs at the time of their actual translation into human clinical trials, as well as the alignment of

147 the CAMAROS trial design to the existing preclinical evidence base. See the Supplemental

148 Methods for the full search strategy and exact search dates for each database.

150 *Selection process and data collection*

151 We deduplicated citations and uploaded them into DistillerSR® (Evidence Partners, Ottawa,

152 Canada). Two reviewers independently screened citations by title and abstract using an

153 accelerated method (one reviewer required to include, two reviewers required to exclude). We

154 then screened and extracted data from full-text articles in duplicate. Graphical data was extracted

155 using Engauge Digitizer.[19] A third reviewer with content expertise in preclinical stroke studies

156 audited all data extraction. Conflicts between reviewers were resolved by consensus discussion.

157 See Supplemental Methods for the complete list of data extraction elements.

159 *Effect measures and data synthesis*

160 We performed quantitative analyses using the R (version 4.1.2) "metafor" package (version

161 4.0.0)[20] with inverse variance random effects modelling. We expressed continuous outcome

162 measures as standardized mean differences (SMDs) with 95% confidence intervals (CIs) and

163 assessed statistical heterogeneity of effect sizes using the Cochrane I^2 statistic.[21] This was

164 necessary due to the variety of outcome measures and measurement scales used across studies,

165 which is a common feature of preclinical systematic reviews. Sensitivity analyses were

166 performed using original measurement scales where possible. From patient partner input,

167 subgroups were analyzed based on timing / dose / route of intervention, stroke model, stroke

168 type, species, type of behavioural outcome (i.e., motor, cognitive), and comorbidities.[22] Our

169 planned subgroup analyses on study quality, specific regions/areas of the brain, and post-stroke

rehabilitation paradigms were not performed due to insufficient number of studies. We did not

assess publication bias using Egger plots due to fewer than 10 studies being included in the

analysis, as per Cochrane recommendations.[23]

Risk of bias assessment

Two independent reviewers used the Systematic Review Centre for Laboratory Animal

Experimentation (SYRCLE) risk of bias tool to assess each study as having a "Low Risk",

"Unclear Risk", "Some Concerns of Risk", or "High Risk" across domains such as

randomization, blinding and outcome reporting.[24] "Some Concerns of Risk" indicated reporting

of a domain (i.e., randomization), but lacking methodological details. This differed from

"Unclear Risk" where there was no mention of the domain in the study.

Comprehensiveness of preclinical evidence and alignment with clinical trials

We assessed comprehensiveness of the *overall body* of preclinical evidence for CCR5

antagonists as a cerebroprotective treatment using The Stroke Treatment Academic Industry

Roundtable (STAIR) I, VI, and XI Consolidated Recommendations.[11,25,26] These

recommendations encompass "candidate treatment qualification" (e.g., dose, timing of dose,

outcomes) and "preclinical assessment and validation" (e.g., age, sex, sample size, animal

type).[11] We excluded domains that were redundant with risk of bias (e.g., randomization,

blinding, etc.) and included additional items relevant to stroke recovery studies from the Stroke

Recovery and Rehabilitation Roundtable (SRRR) Translational Working Group consensus

recommendations.[9] Two reviewers extracted data to determine if the overall evidence across

studies satisfied each of the STAIR and SRRR recommendations. A third reviewer audited this

analysis. We then assessed alignment of existing preclinical evidence with an ongoing clinical

trial of CCR5 antagonists for stroke (CAMAROS, NCT04789616).[7]

Protocol deviations

We incorporated an additional assessment using the PRIMED[2] tool for assessing the readiness of

stroke cerebroprotection therapies to be translated to clinical trials based on the body of

preclinical evidence.[27] Additionally, a list of outcomes used to determine CCR5's potential

mechanisms of action were extracted (Figure S1, Table S3) based on feedback from individuals

that reviewed the initial manuscript drafts.

Results

Study selection

Our search identified 263 citations, which was reduced to 166 unique studies after deduplication.

Five studies representing 10 experiments met the eligibility criteria (Figure 1).[2–6] Herein,

"studies" refer to the published articles as a unit, while "experiments" refer to distinct

investigations within each published article used to test various hypotheses (i.e., a subunit of

"studies" comprised of a select cohort of animals).

Study and animal model characteristics

Most studies used ischaemic stroke (n=4/5). This was induced via intraluminal suture (n=2),

cauterization (n=1), or photothrombosis techniques (n=1; Table 1). Haemorrhagic stroke was

induced in one study via autologous whole blood injection. All studies used mouse (n=4) or rat

(n=1) models, comprised of exclusively male animals (n=5). Relevant stroke comorbidities

highlighted by patient partners and STAIR/SRRR recommendations (e.g., hypertension,

diabetes) were not used in any study.

Intervention characteristics

Maraviroc was used in six of the experiments (n=6/10), TAK-779 in three of the experiments

(n=3), and D-Ala-Peptide T-Amide (DAPTA) in one (n=1; Table 2). These CCR5 antagonists

were delivered intraperitoneally (n=4), intranasally (n=2), intracerebroventricularly (n=2),

subcutaneously (n=1), and intravenously (n=1) at a dose range from 0.01 to 100 mg/kg. Most

studies delivered a single dose of the drug (n=6); experiments with multiple administrations

(n=4) ranged from 3–63 doses. Time of initial treatment administration varied widely. Two

studies (3 experiments) administered treatment pre-stroke (10-15 minutes),[2,3] and 4 studies (6

experiments) in the acute, potentially cerebroprotective, post-stroke period (50 minutes – 24

hours post-stroke).[3–6] One study (1 experiment) was conducted in the late sub-acute/early chronic

period beginning at three to four weeks post stroke, which would be oriented towards recovery,

rather than cerebroprotective, effects.[6] Patient partners had identified several *a priori* interests

(physical therapy alongside CCR5 administration, spasticity), which were also aligned with

SRRR recommended considerations for preclinical stroke recovery studies (see Supplemental

Methods).[9] These were not reported in any included studies.

Meta-analysis of infarct volume

Infarct volume was reported in six experiments (n=6/10) from four different studies with an

overall pooled analysis demonstrating marked cerebroprotection with CCR5 antagonists (SMD -

1.02, 95% CI -1.58 to -0.46, p < 0.0001, I^2=34%; Figure 2). Five of these experiments measured

infarct volume at 1-3 days post-stroke, and one experiment measured infarct volume at a delayed

time point of 63 days post-stroke. No significant differences between pre- or post-stroke

administration were observed (*P*=0.47). Post-hoc sensitivity analysis removing one experiment

with extreme values[2] demonstrated that cerebroprotection in the remaining two experiments

remained statistically significant while reducing heterogeneity (SMD -0.81, 95% CI -1.25 to -

0.37, p < 0.001, I^2=0%). A second sensitivity analysis excluded one study that measured infarct

volume in mm^3 so that all other studies could be meta-analyzed using mean differences on the

percentage scale (Figure S2; MD -9.1%, 95% CI -11.6 to -6.7%, p < 0.001, I^2=0%). This

demonstrated a similar cerebroprotective effect as the other analyses. Further sub-group analyses

by route of administration, time of administration, stroke model, species, CCR5 antagonist, dose,

and whether behaviour tests were assessed are described in Figure S3-S5.

Synthesis of behavioural outcomes without meta-analysis

Motor behavioural outcomes were reported in six experiments from three studies and represented

seven different behavioural tasks. An additional study reported motor behavioural outcomes

without standard deviations or standard errors, and thus could not be included (authors did not

respond to email requests for data).[4] A cognitive outcome (Morris Water Maze) was measured in

one study. Overall, CCR5 inhibition was effective in 11 of 16 behavioural outcomes tested

(Figure 3). Meta-analysis and planned subgroup analysis were contra-indicated due to an

inadequate number of studies for each given outcome measure, which necessitated the synthesis

without meta-analysis presented below.[28]

261 Behavioural outcomes are presented by time of CCR5 antagonist administration, as discussed in

262 the Intervention Characteristics section above, as administration time directly relates to the

263 treatment context and patient population to which the results apply (i.e., acute cerebroprotection

264 vs. late sub-acute/early chronic neural repair). Li et al. reported that pre-stroke administration

265 (relevant to surgical contexts with high risk of thrombosis) of DAPTA did not result in

266 significantly greater performance on the neurological deficit score.[2]

268 Regarding acute post-stroke administration times with the potential for *cerebroprotective* effects

269 (up to 24 hours post-stroke based on observed infarct reductions in Figure 2), Yan et al. observed

270 that CCR5 antagonist (maraviroc) administration one-hour following stroke resulted in greater

271 motor performance on the corner test, limb placement, modified Garcia, foot fault, and rotarod

272 tasks compared to vehicle-treated controls. Cognitive outcomes in the Morris Water Maze task

273 (proportion of time spent in the probe quadrant) were also improved by this one-hour post-stroke

274 administration. Significantly improved motor performance was not observed if CCR5 antagonist

275 was administered 24 hours post-stroke, using this model of *intracerebral haemorrhage* (Figure

276 3).[5] In contrast, using a *focal ischaemia* model, Joy et al. observed that CCR5 antagonist

277 (maraviroc) administration initiated 24 hours post-stroke and continued daily for 3 weeks (which

278 could potentially represent a mix of cerebroprotective and recovery-induced effects) resulted in

279 greater performance on the cylinder and grid walk tasks compared to vehicle-treated controls.

281 Joy et al., also performed one experiment with *late sub-acute/early chronic* administration (a

282 potentially critical time for neural repair) initiated at three to four weeks post-stroke and

283 continued daily for 11 weeks, that demonstrated significantly improved performance for the grid

284 walk, but not cylinder, task.[6] This experiment was initiated outside the plausible range for a

285 cerebroprotective effect, implying behavioural improvement involved *recovery-promoting*

286 *mechanisms*. However, equivalent infarct volumes were not demonstrated between the treated

287 and control groups in this cohort, which could potentially lead to confounding effects.

289 *Risk of bias*

290 All articles had a 'high' risk of bias in at least one domain of the SYRCLE tool.[24] Most domains

291 within each study demonstrated an 'unclear' risk of bias. All studies reported randomizing

animals; however, as commonly observed in the preclinical literature,[29–31] only one of these

studies provided sufficient detail to ensure that the randomization method had a low risk of bias.

The SYRCLE domains with the highest risk of bias were incomplete outcome data, with 80% of

studies (n=4/5) failing to provide complete data for all animals initially included in the study, as

well as selective outcome reporting, with 60% (n=3/5; all studies of maraviroc) not providing

complete data for all expected outcomes discussed in the methods (Figure S6).

Comprehensiveness of preclinical evidence and alignment with clinical trials

We assessed comprehensiveness of the preclinical evidence using the STAIR and SRRR

recommendations[9,11,25,26] as well as alignment with study parameters of the CAMAROS trial.[7] A

summary of assessment items from STAIR XI is provided in Table 3, with additional items from

STAIR I, VI, and SRRR recommendations included in Table S4.

For CCR5 antagonists as a post-stroke cerebroprotectant, the overall body of evidence satisfied

all five STAIR XI domains assessing 'candidate treatment qualification' (Table 3). Overall, a

range of doses and clinically relevant administration times for cerebroprotection were evaluated

across a variety of motor and cognitive behavioural domains. All studies tested both behavioural

and histological outcomes and demonstrated cerebroprotective effects, but most studies failed to

measure and control post-stroke temperature, which could potentially confound the observed

cerebroprotection (Table S4).[32] Most histological measurements were also assessed at <72 hours,

which could confound the observed cerebroprotective effects if cell death was merely delayed.[32]

For CCR5 antagonists as a post-stroke recovery-inducing treatment, one experiment assessed the

effects of initiating CCR5 administration in a similar post-stroke phase as the CAMAROS trial.

This experiment (Joy et al.)[6] did not demonstrate that each treatment group had equivalent

baseline stroke volumes, which may potentially confound observed behavioural effects.

Furthermore, the maximum dose used in mice (100 mg/kg) was not sufficient to attain

cerebrospinal fluid levels of maraviroc observed in humans using the CAMAROS dosing regime

(mice: 13.8±5.4 ng/mL; humans 33.6–60 ng/mL).[6]

Areas of concern were identified in all STAIR XI domains assessing 'preclinical assessment and

validation' (Table 3). Although adult animals were used in most of the preclinical studies, the

reported weights and ages of these animals corresponded to young adults, rather than the aged

adults that better represent the stroke population. All experiments used only male rodents that

were free of common stroke comorbidities. It was also unclear if behavioural testing was

performed during the inactive circadian phase or active (dark) phase, which could result in

confounding if CCR5 antagonists affect arousal of animals during their inactive period.[33]

Furthermore, none of the studies had their protocols or results directly replicated by an

independent laboratory, or across multiple sites in a multilaboratory study.[9,34] Regarding stroke

recovery, no studies assessed behavioural effects on upper extremity skilled reaching / grasping

or potential interactions of CCR5 antagonists with rehabilitative therapies or established

recanalization procedures (Table S4).[35–38] These elements are highly relevant to the CAMAROS

trial, as one of the primary outcomes of this trial is upper extremity performance on the Fugl-

Meyer and maraviroc administration will be paired with an 8-week exercise program. These

findings were supported by the PRIMED[2] tool, which resulted in a Readiness for Translation

Score of "medium" (on a scale of "low", "medium", "high"). This tool highlighted similar

promising elements, as well as weaknesses, as our analysis above (Table 3, Table S4),

identifying the limited preclinical evidence for the effects of CCR5 antagonists in clinically

relevant sexes, ages, species, and disease comorbidities, without sufficient dose-response

information to inform trials (Table S5). Overall, our assessments highlight a variety of

knowledge gaps, potential confounding factors, and areas of misalignment between the

preclinical evidence and clinical trial parameters that could be improved with further preclinical

experimentation.

Discussion

The overall body of preclinical evidence for CCR5 antagonists in stroke demonstrates potential

acute cerebroprotection with corresponding impairment reduction, as well as improved

functional recovery in the sub-acute/early chronic phase. Our systematic review also highlights

evidence gaps that could impact successful clinical translation of CCR5 antagonists. Our analysis

of 10 independent experiments, identified that acute administration of CCR5 antagonists within

the first 24 hours post-stroke was associated with a marked reduction in infarct volume. This

cerebroprotective reduction of infarct volume did not significantly vary based on treatment dose

or any other experimental characteristics.[3,4,6] Overall, the majority of behavioural effects

appeared to be in a positive direction, but the low number of included studies precluded meta-

analysis of these results. Indeed, no individual behavioural experiment included more than ten

CCR5-treated animals, and given the wide range of dosages, timings, routes, stroke models, and

rodent strains involved, the certainty of these findings is limited and should be interpreted

cautiously. Pooling data across heterogenous experimental designs, animal/stroke models, and

treatment parameters, as we have done with the infarct volume analysis in the present study, can

introduce variability that increases the risk of overestimating or underestimating the true effect of

the intervention.[39] Treatment effects observed across model systems and therapeutic compounds

may represent different biological mechanisms. Despite this potential limitation, meta-analysis

can provide valuable insights, especially in preclinical settings where the sample sizes of

individual studies may be too small to detect significant effects on their own. In these cases,

pooling data across studies can help identify overarching estimates of benefits and harm,

highlight subgroups of interest, and help guide areas of future research. As described in the

results above, we attempted to mitigate the risks of inappropriate data pooling through careful

investigation of heterogeneity, subgroup analyses, and differentiation between outcomes where

we felt that meta-analytic pooling was (infarct volume) and was not (behavioural outcomes)

appropriate. Overall, we believe that our results indicate that further investigation is warranted to

determine the optimal timing of administration and behavioural domains under which CCR5

antagonists exhibit the strongest post-stroke cerebroprotective and recovery-inducing effects.

Despite the positive direction of treatment effects across all studies of CCR5 antagonists, we

found a substantial risk of bias in the underlying studies.[24] As is commonly observed in the

preclinical literature, all studies either did not adequately report their randomization / blinding

methods and exhibited evidence of selective / incomplete reporting.[29–31] Such features are

associated with biased overestimations of preclinical treatment efficacy, which raises further

concerns about the reliability and validity of the present findings.[40–42] Future studies should

carefully incorporate all elements of the ARRIVE 2.0 guidelines to help ensure that results are

transparently reported and improve confidence in the findings.[43]

Comprehensiveness of the preclinical evidence for CCR5 antagonists was assessed in relation to

STAIR and SRRR consensus recommendations.[9,11,25,26] These recommendations aim to provide

investigators and regulators with "assurance that the candidate treatment shows signals of

efficacy and safety, before embarking on an expensive clinical development program".[11] The

included studies provide good initial evidence for acute cerebroprotection, as well as mechanistic

and behavioural evidence for enhanced recovery in the late sub-acute/early chronic post-stroke

phase. However, demonstration of efficacy under a wider range of conditions, such as in aged

animals, females, animals with stroke-related comorbidities, more clinically relevant timing of

dose administrations, or in conjunction with rehabilitative therapies are necessary to provide

further confidence in these findings. In addition, all studies used unique doses, timings, and

outcomes, so independent replication of the most promising study parameters would further

increase certainty in the evidence.[9,11] Future preclinical research should aim to address these

evidence gaps to further increase the clinical relevance and comprehensiveness of evidence for

CCR5 antagonists in stroke.

In relation to the ongoing CAMAROS trial assessing maraviroc in the subacute post-stroke

phase,[7] the most relevant preclinical evidence comes from one experiment within the Joy et al.

study[6] where maraviroc was initially administered at three to four weeks post-stroke and

continued daily for 11 weeks. This experiment demonstrated that administration of maraviroc in

the late sub-acute/early chronic post-stroke phase improved functional recovery on the grid walk,

but not cylinder, task. These experimental conditions could potentially align with the putative

therapeutic window and outcomes of interest being assessed in CAMAROS (e.g. 10-minute walk

test co-primary outcome).[7] However, caution is warranted as this pivotal supporting preclinical

evidence is based on a low sample size (n=9). Moreover, potential differences in dosing, severity

of infarct, concomitant rehabilitative therapy, and other factors discussed above could influence

the degree to which these results successfully translate to the clinical environment. Finally,

clinically relevant secondary outcomes in the CAMAROS trial, such as cognitive and emotional

domains as measured by the Montreal Cognitive Assessment (MoCA) and Stroke Aphasia

Depression Questionnaire (SADQ) were not modelled in the preclinical literature. Although one

study included a cognitive outcome, the other treatment parameters of this study were not

aligned to the CAMAROS trial.[5] Future preclinical studies should assess a more diverse and

comprehensive battery of clinically relevant behavioural tasks, which could be based on the

range of outcomes employed in the CAMAROS trial, or those found in the SRRR

recommendations.[9] Nevertheless, the Joy et al., study provides a plausible biological mechanism

and "proof of concept" for how CCR5 antagonism might enhance neuroplasticity that improves

functional recovery after stroke.

Our present synthesis of the preclinical evidence for CCR5 antagonists used novel approaches to

increase its utility for assessing certainty of the findings and identification of knowledge gaps.

First, we engaged patients with lived experiences of stroke throughout the review process to

ensure that our research questions, outcomes, and interpretations aligned with the priorities of the

ultimate end-user of stroke research. Second, we incorporated consensus recommendations for

both preclinical cerebroprotection and recovery research as an evidence evaluation tool, which

we found often aligned with the priorities of our patient panel.[9,11,25,26] This guided assessment of

the alignment of preclinical evidence with parameters of ongoing clinical trials, as well as

appraisal of comprehensiveness of the preclinical evidence with a focus on translational

validity[44] rather than only internal validity and risk of bias.[24] We also used this method to

provide concrete avenues for future preclinical studies to close knowledge gaps and improve

certainty in the effects of CCR5 antagonists under clinically relevant experimental conditions.

Similar approaches should be considered by future preclinical systematic reviews to improve

interpretation of the preclinical evidence from a translational perspective.

In conclusion, CCR5 antagonists show promise in preclinical studies for stroke

cerebroprotection, corresponding reduction in impairment, as well as improved functional

recovery related to neural repair in the late sub-acute/early chronic phase. However, high risk of

bias and the limited (or no) evidence in clinically relevant domains underscore the need for more

rigorous and transparent preclinical research to further strengthen the current preliminary

evidence available in the literature. Addressing these concerns will not only enhance the

reliability of preclinical evidence but also better inform the design and execution of clinical trials

of CCR5 antagonists, such as the ongoing CAMAROS trial.[7] The integration of expert

recommendations, such as STAIR and SRRR, should guide future preclinical investigations and

synthesis of the body of preclinical evidence in stroke recovery research.[9,11] Our present

approach serves as a template by which the preclinical evidence supporting translation to clinical

trials can be weighed when justifying early clinical trials of novel interventions for stroke

cerebroprotection and recovery.

Acknowledgements: The authors thank Hannah Laquerre for her assistance with figure

generation. Figure S1 was created with BioRender.com.

Sources of Funding: This study was funded by a Social Accountability Grant from the

University of Ottawa's Faculty of Medicine. MSJ was supported by the Canadian Institutes of

Health Research (CIHR) and Vanier Canada Graduate Scholarship (CGV-186957). The funders

were not involved in the study design, collection, analysis and interpretation of data, writing the

manuscript, or in the decision to submit the article for publication. MML is supported by

The Ottawa Hospital Anesthesia Alternate Funds Association, a University of Ottawa Junior

Research Chair, and the Canadian Anesthesiologists' Society Career Scientist Award.

Disclosures: None.

Supplemental Material:

Supplemental Methods

Figures S1-S6

Tables S1-S5

References:

1. Jing K, Chen F, Shi X, Guo J, Liu X. Dual effect of C–C motif chemokine receptor 5 on ischemic stroke: More harm than benefit? *European journal of pharmacology*. 2023;953:175857–175857.

2. Li L, Zhi D, Shen Y, Liu K, Li H, Chen J. Effects of CC-chemokine receptor 5 on ROCK2 and P-MLC2 expression after focal cerebral ischaemia-reperfusion injury in rats. *Brain injury*. 2016;30:468–473.

3. Takami S, Minami M, Katayama T, Nagata I, Namura S, Satoh M. TAK-779, A Nonpeptide CC Chemokine Receptor Antagonist, Protects the Brain Against Focal Cerebral Ischemia in Mice. *Journal of cerebral blood flow and metabolism*. 2002;22:780–784.

4. Chen B, Cao P, Guo X, Yin M, Li X, Jiang L, Shao J, Chen X, Jiang C, Tao L, et al. Maraviroc, an inhibitor of chemokine receptor type 5, alleviates neuroinflammatory response after cerebral Ischemia/reperfusion injury via regulating MAPK/NF-κB signaling. *International immunopharmacology*. 2022;108:108755–108755.

5. Yan J, Xu W, Lenahan C, Huang L, Wen J, Li G, Hu X, Zheng W, Zhang JH, Tang J. CCR5 Activation Promotes NLRP1-Dependent Neuronal Pyroptosis via CCR5/PKA/CREB Pathway After Intracerebral Hemorrhage. *Stroke*. 2021;52:4021–4032.

6. Joy MT, Ben Assayag E, Shabashov-Stone D, Liraz-Zaltsman S, Mazzitelli J, Arenas M, Abduljawad N, Kliper E, Korczyn AD, Thareja NS, et al. CCR5 Is a Therapeutic Target for Recovery after Stroke and Traumatic Brain Injury. *Cell*. 2019;176:1143-1157.e13.

7. Dukelow S, Clinicaltrials.gov NCT04789616. The Canadian Maraviroc RCT To Augment Rehabilitation Outcomes After Stroke (CAMAROS) [Internet]. [cited 2023 Oct 20];Available from: https://clinicaltrials.gov/study/NCT04789616

8. O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW. 1,026 experimental treatments in acute stroke. *Ann Neurol*. 2006;59:467–477.

9. Corbett D, Carmichael ST, Murphy TH, Jones TA, Schwab ME, Jolkkonen J, Clarkson AN, Dancause N, Weiloch T, Johansen-Berg H, et al. Enhancing the Alignment of the Preclinical and Clinical Stroke Recovery Research Pipeline: Consensus-Based Core Recommendations From the Stroke Recovery and Rehabilitation Roundtable Translational Working Group. *Neurorehabilitation and neural repair*. 2017;31:699–707.

10. Bosetti F, Koenig JI, Ayata C, Back SA, Becker K, Broderick JP, Carmichael ST, Cho S, Cipolla MJ, Corbett D, et al. Translational Stroke Research: Vision and Opportunities. *Stroke*. 2017;48:2632–2637.

11. Lyden P, Buchan A, Boltze J, Fisher M. Top Priorities for Cerebroprotective Studies-A Paradigm Shift: Report From STAIR XI. *Stroke*. 2021;52:3063–3071.

12. Carmichael ST. Emergent properties of neural repair: elemental biology to therapeutic
concepts. *Annals of neurology*. 2016;79:895–906.

13. Murphy TH, Corbett D. Plasticity during stroke recovery: from synapse to behaviour. *Nature*
reviews. Neuroscience. 2009;10:861–872.

14. Lalu MM, Sharif A, Nicholls S, Dowlatshahi D, Fergusson DA, Corbett D, Bapuji R, Jeffers
MS, Humphrey J, Johnston W, et al. Evaluating the effect of CCR5 inhibitors in animal
models of stroke: A systematic review co-designed with patient partners [Internet]. 2023
[cited 2023 Oct 20];Available from:
https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=393438

15. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L,
Tetzlaff JM, Akl EA, Brennan SE, et al. The PRISMA 2020 statement: An updated guideline
for reporting systematic reviews. *PLoS medicine*. 2021;18:e1003583–e1003583.

16. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, Altman DG, Moher D,
Barber R, Denegri S, et al. GRIPP2 reporting checklists: tools to improve reporting of patient
and public involvement in research. *BMJ*. 2017;358:j3453–j3453.

17. Hooijmans CR, Tillema A, Leenaars M, Ritskes-Hoitinga M. Enhancing search efficiency by
means of a search filter for finding all studies on animal experimentation in PubMed.
Laboratory animals (London). 2010;44:170–175.

18. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer
Review of Electronic Search Strategies: 2015 Guideline Statement. *Journal of clinical*
epidemiology. 2016;75:40–46.

19. Mitchell M, Muftakhidinov B, Winchen T. Engauge Digitizer Software [Internet]. [cited
2023 Oct 20];Available from: http://markummitchell.github.io/engauge-digitizer/

20. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *Journal of*
statistical software. 2010;36.

21. Higgins JP, Li T, Deeks J. Chapter 6: Choosing effect measures and computing estimates of
effect [Internet]. In: Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch VA,
editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.4 (updated
August 2023). Cochrane; 2023 [cited 2023 May 1]. Available from:
www.training.cochrane.org/handbook

22. Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, Krakauer JW,
Boyd LA, Carmichael ST, Corbett D, et al. Agreed definitions and a shared vision for new
standards in stroke recovery research: The Stroke Recovery and Rehabilitation Roundtable
taskforce. *International journal of stroke*. 2017;12:444–450.

23. Page MJ, Higgins JP, Sterne JAC. Chapter 13: Assessing risk of bias due to missing results
in a synthesis [Internet]. In: Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch
VA, editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.4

(updated August 2023). Cochrane; 2023 [cited 2023 May 1]. Available from:
www.training.cochrane.org/handbook

24. Hooijmans CR, Rovers MM, de Vries RB, Leenaars M, Ritskes-Hoitinga M, Langendam
MW. SYRCLE's risk of bias tool for animal studies. *BMC Medical Research Methodology*.
2014;14:43.

25. Finklestein SP, Fisher M, Furlan AJ, Goldstein LB. Recommendations for standards
regarding preclinical neuroprotective and restorative drug development. *Stroke*.
1999;30:2752–2758.

26. Fisher M, Feuerstein G, Howells DW, Hum PD, Kent TA, Savitz SI, Lo EH. Update of the
Stroke Therapy Academic Industry Roundtable Preclinical Recommendations. *Stroke*.
2009;40:2244–2250.

27. Bahr-Hosseini M, Bikson M, Iacoboni M, Liebeskind DS, Hinman JD, Carmichael ST, Saver
JL. PRIMED2 Preclinical Evidence Scoring Tool to Assess Readiness for Translation of
Neuroprotection Therapies. *Translational stroke research*. 2022;13:222–227.

28. Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-
Boyce J, Ryan R, Shepperd S, Thomas J, et al. Synthesis without meta-analysis (SWiM) in
systematic reviews: reporting guideline. *BMJ*. 2020;368:l6890.

29. Avey MT, Moher D, Sullivan KJ, Fergusson D, Griffin G, Grimshaw JM, Hutton B, Lalu
MM, Macleod M, Marshall J, et al. The Devil Is in the Details: Incomplete Reporting in
Preclinical Animal Research. *PLoS One*. 2016;11:e0166733.

30. Fergusson DA, Wesch NL, Leung GJ, MacNeil JL, Conic I, Presseau J, Cobey KD, Diallo J-
S, Auer R, Kimmelman J, et al. Assessing the Completeness of Reporting in Preclinical
Oncolytic Virus Therapy Studies. *Mol Ther Oncolytics*. 2019;14:179–187.

31. Fergusson DA, Avey MT, Barron CC, Bocock M, Biefer KE, Boet S, Bourque SL, Conic I,
Chen K, Dong YY, et al. Reporting preclinical anesthesia study (REPEAT): Evaluating the
quality of reporting in the preclinical anesthesiology literature. *PLoS One*.
2019;14:e0215221.

32. Corbett D, Nurse S. The problem of assessing effective neuroprotection in experimental
cerebral ischemia. *Progress in neurobiology*. 1998;54:531–548.

33. MacLellan CL, Keough MB, Granter-Button S, Chernenko GA, Butt S, Corbett D. A Critical
Threshold of Rehabilitation Involving Brain-Derived Neurotrophic Factor Is Required for
Poststroke Recovery. *Neurorehabilitation and neural repair*. 2011;25:740–748.

34. Hunniford VT, Grudniewicz A, Fergusson DA, Montroy J, Grigor E, Lansdell C, Lalu MM.
A systematic assessment of preclinical multilaboratory studies and a comparison to single
laboratory studies. *eLife*. 2023;12.

35. Balbinot G, Schuch CP, Jeffers MS, McDonald MW, Livingston-Thomas JM, Corbett D.
Post-stroke kinematic analysis in rats reveals similar reaching abnormalities as humans.
Scientific reports. 2018;8:8738–13.

36. Jeffers MS, Corbett D. Synergistic Effects of Enriched Environment and Task-Specific
Reach Training on Poststroke Recovery of Motor Function. *Stroke*. 2018;49:1496–1503.

37. Jeffers MS, Karthikeyan S, Gomez-Smith M, Gasinzigwa S, Achenbach J, Feiten A, Corbett
D. Does Stroke Rehabilitation Really Matter? Part B: An Algorithm for Prescribing an
Effective Intensity of Rehabilitation. *Neurorehabilitation and neural repair*. 2018;32:73–83.

38. Wechsler LR, Adeoye O, Alemseged F, Bahr-Hosseini M, Deljkich E, Favilla C, Fisher M,
Grotta J, Hill MD, Kamel H, et al. Most Promising Approaches to Improve Stroke
Outcomes: The Stroke Treatment Academic Industry Roundtable XII Workshop. *Stroke
(1970)*. 2023;54:3202–3213.

39. Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch V. Chapter 15:
Interpreting results and drawing conclusions [Internet]. In: Cochrane Handbook for
Systematic Reviews of Interventions version 6.4 (updated August 2023). Cochrane; 2023.
Available from: www.training.cochrane.org/handbook

40. Holman C, Piper SK, Grittner U, Diamantaras AA, Kimmelman J, Siegerink B, Dirnagl U.
Where Have All the Rodents Gone? The Effects of Attrition in Experimental Research on
Cancer and Stroke. *PLoS Biology*. 2016;14:e1002331–e1002331.

41. Sena ES, van der Worp HB, Bath PMW, Howells DW, Macleod MR. Publication bias in
reports of animal stroke studies leads to major overstatement of efficacy. *PLoS Biology*.
2010;8:e1000344–e1000344.

42. Macleod MR, Bart Van Der Worp H, Sena ES, Howells DW, Dirnagl U, Donnan GA.
Evidence for the Efficacy of NXY-059 in Experimental Focal Cerebral Ischaemia Is
Confounded by Study Quality. *Stroke*. 2008;39:2824–2829.

43. Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A,
Cuthill IC, Dirnagl U, et al. The ARRIVE guidelines 2.0: Updated guidelines for reporting
animal research. *PLoS Biol*. 2020;18:e3000410.

44. Drude NI, Martinez Gamboa L, Danziger M, Dirnagl U, Toelch U. Improving preclinical
studies through replications. *eLife*. 10:e62101.

Figure

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) flow
diagram.
610

Figure 2. CCR5 antagonists reduce infarct volume. Data is presented as a forest plot with standardized mean differences and 95% confidence intervals. Effect sizes <0 favours drug treatment and >0 favours control. Data is stratified by timing of CCR5 antagonist administration (pre- or post-stroke induction). The 'RE Model for All Studies' represents a pooled estimate of the CCR5 antagonist drug effect on infarct volume from all studies combined. Separate pooled estimates are also reported for post-stroke and pre-stroke CCR5.

Figure 3. Synthesis without meta-analysis for all included preclinical CCR5 antagonist studies that reported motor and/or cognitive behavioural outcomes. Data is presented as a forest plot with a standardized mean difference and 95% confidence intervals. Effect sizes <0 favours drug treatment and >0 favours control.

Tables

Table 1. Summary of study and animal model characteristics of included articles.

Author	Year	Country	Species	Strain	Stroke Type	Stroke Model	Sex	Weight	Age
Li et al.[2]	2016	China	Rat	Wistar	Ischaemic	Intraluminal suture	Male	260–300 g	N/A
Takami et al.[3]	2002	Japan	Mice	ddY	Ischaemic	Intraluminal suture	Male	N/A	4 weeks
Chen et al.[4]	2022	China	Mice	C57/BL6	Ischaemic	Permanent middle cerebral artery occlusion	Male	25–30 g	8–10 weeks
Yan et al.[5]	2021	China	Mice	CD1	Haemorrhagic	Intracerebral haemorrhagic	Male	30–40 g	N/A
Joy et al.[6]	2019	USA	Mice	C57/BL6	Ischaemic	Photothrombotic	Male	25–30 g	8–20 weeks

Table 2. Summary of intervention characteristics.

Author	Drug	Dose (mg/kg)	Route	Timing	Doses (total #)	Outcomes Measured (Treatment n/Control n)	Outcome Window (post-stroke))
Li et al.[2]	DAPTA (D-Ala-Peptide T-Amide)	0.01	SC	15 min pre-stroke	1	- Infarct volume (5/5) - Neurological deficit score (5/5)	24 h
Takami et al.[3]	TAK-779	5	ICV	10 min pre-stroke	1	- Infarct volume (10/6)	48 h
		50	ICV	10 min pre-stroke		- Infarct volume (13/6)	
		0.25	IV	50 min post-stroke		- Infarct volume (17/18)	
Chen et al.[4]	Maraviroc	20	IP	1.5 h, 24 h, and 48 h post-stroke	3	- Infarct volume (5/5) - Longa score (5/5) - Neurological deficit score (5/5)	72 h
Yan et al.[5]	Maraviroc	0.15	IN	1 h post-stroke	1	- Garcia test (6/6) - Limb placement (6/6) - Corner turn test (6/6)	72 h
						- Foot fault (8/8) - Rotarod (8/8)	3 weeks
						- Probe quadrant duration (8/8)	25 days
				24 h post-stroke		- Garcia test (6/6) - Limb placement (6/6) - Corner turn test (6/6)	72 h
Joy et al.[6]	Maraviroc	100	IP	24 h post-stroke through daily injections for 9 weeks	63	- Infarct volume (5/5)	9 weeks
						- Grid walk (10/10) - Forelimb (10/10)	8 weeks
				24 h post-stroke then daily for 3 weeks	21	- Grid walk (10/10) - Cylinder test (9/8)	9 weeks
				3 to 4 weeks post-stroke then daily for 11 weeks	56	- Grid walk (9/9) - Cylinder test (9/9)	11 weeks

Abbreviations: ICV = intracerebroventricular; IN = intranasal; IP = intraperitoneal; IV = intravenous; SC = subcutaneous

Table 3. Alignment of included preclinical studies with STAIR recommendations and CAMAROS trial parameters.

Recommendation	Overall preclinical evidence	CAMAROS trial alignment	Notes
Candidate treatment qualification			
1) Establish treatment dose-response	Yes	No	• **Preclinical** – doses across all studies from 0.01 to 100 mg/kg • 6 doses tested for infarct volume, 3 doses for behavioural effects • Joy et al. demonstrated that 100 mg/kg of maraviroc in mice results in cerebrospinal fluid levels lower than in humans (13.8±5.4 vs. 33.6–60 ng/mL) • **CAMAROS** – participants take 300 mg doses twice daily for 8 weeks • **Interpretation** – maximum preclinical dose may not represent clinical levels
2) Treatment given after clinically relevant delayed times (one to 4.5 hours post-stroke)	Yes	Yes	• **Preclinical** – 3/5 studies (all maraviroc) administered drug from 1 to 24 hours post-stroke (acute phase) • Joy et al. initiated maraviroc administration at 20 days post-stroke for 11 weeks (subacute phase) • **CAMAROS** – participants recruited between 5 days and 6 weeks post-stroke (acute / early subacute) • **Interpretation** – one experiment in a time window relevant to clinical trial
3) Both histologic and behavioural outcomes at acute (1 to 3 days) and chronic (7 to 30 days) time points	Yes	Yes	• **Preclinical** – 2/5 studies included both histologic and behavioural outcomes • Mainly tasks of spontaneous movement (e.g., cylinder) or motor coordination (e.g., grid walk) • Tests ranged from 3 days (acute) to 11 weeks (chronic) post-stroke for all included studies, but short term assessment may only delay cell death[32] • **CAMAROS** - motor learning (Fugl-Meyer Upper Extremity Assessment Score and 10-Meter Walk Test Score) measured as the primary outcome at baseline, 4-week (late subacute), 8-week (late subacute), and 6-month (chronic) • **Interpretation** – challenging to directly compare preclinical and clinical motor tasks, but testing range is comparable. Most relevant Joy et al. experiment (subacute administration) did not measure histologic outcomes
4) Treatment reaches presumed target and causes expected physiological effects that can be assessed with a clinically relevant biomarker	Yes	NA	• **Preclinical** – 5/5 studies demonstrated trends for CCR5 antagonism to reduce infarct volume. Plausible that CCR5 influences stroke-relevant mechanisms • **CAMAROS** – does not include outcomes to assess treatment mechanisms
5) Treatment able to pass the blood-brain barrier	Yes	NA	• **Preclinical** – 1/5 studies used mass spectrometry to demonstrate that maraviroc is present in the brain and cerebrospinal fluid • **CAMAROS** – does not include outcomes to assess presence of drug in the brain

Table 3 (continued). Alignment of included preclinical studies with STAIR recommendations and CAMAROS trial parameters.

Recommendation	Overall preclinical evidence	CAMAROS trial alignment	Notes
Preclinical assessment / validation			
1) Aging/adult age	No	No	• **Preclinical** – 4/5 studies used rodents with weights/ages consistent with young adulthood; 0/5 studies used rodents ≥10 months old. • **CAMAROS** – Age ≥18 year old adult participants eligible for recruitment • **Interpretation** – Preclinical studies consistent with CAMAROS eligibility criteria; however, no preclinical study examined middle-aged or elderly animals that would be more representative of the trial and overall stroke populations
2) Male and female animals	No	No	• **Preclinical** – 0/5 studies included female animals • **CAMAROS** – both male and female sexes eligible for recruitment • **Interpretation** – effects of CCR5 antagonists for stroke recovery in female animals represents a critical knowledge gap
3) Animals with co-morbidities	No	No	• **Preclinical** – 0/5 studies included animals with common stroke co-morbidities (e.g., diabetes, hypertension, etc.) • **CAMAROS** - participants with stroke co-morbidities eligible for recruitment • **Interpretation -** effects of CCR5 antagonists for stroke recovery in animals with common stroke comorbidities represents a critical knowledge gap
4) Evidence from two or more laboratories	Yes	No	• **Preclinical** – 5/5 studies demonstrated stroke-related benefits; however, no study had their results replicated by independent laboratories • **CAMAROS** – trial protocol will be replicated by multiple study sites • **Interpretation** – future preclinical experiments could consider assessing effects using a multilaboratory approach, or independently replicate the effects of existing preclinical studies
5) Gyrencephalic species	No	No	• **Preclinical** – 5/5 studies conducted in lissencephalic (rodent) species. • **CAMAROS** – human participants • **Interpretation** – effects of CCR5 antagonists for stroke recovery in gyrencephalic species represents a critical knowledge gap
6) Tests during the awake phase of animals	No	No	• **Preclinical** – 0/5 studies reported assessing outcomes at night (i.e. awake phase of rodents) • **CAMAROS** – follow-up occurs during daytime hours (i.e. awake phase of humans) • **Interpretation** – future preclinical studies should conduct behavioural testing during the awake phase to better align with human testing conditions

4