1 **Title:** Preclinical systematic review of CCR5 antagonists as cerebroprotective and stroke
2 recovery enhancing agents
3
4 **Abbreviated Title:** CCR5, cerebroprotection, and stroke recovery
5

6 **Authors:** Ayni Sharif[1,2,*], Matthew S. Jeffers[1,2,*], Dean A. Fergusson[1,2,3], Raj Bapuji[1], Stuart G.
7 Nicholls[4], John Humphrey[5], Warren Johnston[5], Ed Mitchell[5], Mary-Ann Speirs[5], Laura
8 Stronghill[5], Michele Vuckovic[5], Susan Wulf[5], Risa Shorr[6], Dar Dowlatshahi[2,7,8], Dale Corbett[8],
9 Manoj M. Lalu[1,2,8,9,#]
10
Affiliations: [1]Clinical Epidemiology Program, Blueprint Translational Research Group, Ottawa
Hospital Research Institute, Ottawa, Canada
[2]School of Epidemiology and Public Health, University of Ottawa, Ottawa, Canada
[3]Department of Medicine, University of Ottawa, Ottawa, Canada
[4]Office for Patient Engagement in Research Activity (OPERA), Ottawa Hospital Research
Institute, Ottawa, Canada
[5]Patient Partner, Blueprint Translational Research Group, Ottawa Hospital Research Institute,
Ottawa, Canada
[6]Ottawa Hospital Library Services, The Ottawa Hospital, Ottawa, Canada
[7]Division of Neurology, Department of Medicine, The Ottawa Hospital, Ottawa, Canada
[8]Department of Cellular and Molecular Medicine, Faculty of Medicine, University of Ottawa,
Ottawa, Canada
[9]Department of Anaesthesiology and Pain Medicine, The Ottawa Hospital, Ottawa, Canada
*These authors contributed equally
25
26 **Corresponding Author:**
27 [#]Manoj M. Lalu, mlalu@toh.ca
28 Department of Cellular and Molecular Medicine, University of Ottawa
29 Clinical Epidemiology and Regenerative Medicine Programs, Ottawa Hospital Research Institute
30 Ottawa, Ontario, Canada, K1Y 4E9
31
32 **X:** @ManojLalu

33
34 **Number of Figures:** 4
35 **Number of Tables:** 4
36 **Word counts:** Abstract (219), Main text (3,785)
37

ABSTRACT

C-C chemokine receptor type 5 (CCR5) antagonists may improve both acute stroke outcome and

long-term recovery. Despite their evaluation in ongoing clinical trials, gaps remain in the

evidence supporting their use. With a panel of patients with lived experiences of stroke, we

performed a systematic review of animal models of stroke that administered a CCR5 antagonist

and assessed infarct size or behavioural outcomes. MEDLINE, Web of Science, and Embase

were searched. Article screening and data extraction were completed in duplicate. We pooled

outcomes using random effects meta-analyses. We assessed risk of bias using the Systematic

Review Centre for Laboratory Animal Experimentation (SYRCLE) tool and alignment with the

Stroke Treatment Academic Industry Roundtable (STAIR) and Stroke Recovery and

Rehabilitation Roundtable (SRRR) recommendations. Five studies representing 10 experiments

were included. CCR5 antagonists reduced infarct volume (standard mean difference -1.02; 95%

confidence interval -1.58 to -0.46) when compared to stroke-only controls. Varied timing of

CCR5 administration (pre- or post-stroke induction) produced similar benefit. CCR5 antagonists

significantly improved 11 of 16 behavioural outcomes reported. High risk of bias was present in

all studies and critical knowledge gaps in the preclinical evidence were identified using

STAIR/SRRR. CCR5 antagonists demonstrate promise; however, rigorously designed preclinical

studies that better align with STAIR/SRRR recommendations and downstream clinical trials are

warranted. Prospective Register of Systematic Reviews (PROSPERO CRD42023393438).

57
58 **Key Words:** stroke; neuroprotection; impairment; recovery; preclinical; CCR5 antagonists;
59 maraviroc; patient engagement in research; CAMAROS; cerebroprotection
60
Non-standard Abbreviations and Acronyms:
CAMAROS   Canadian Maraviroc RCT to Augment Rehabilitation Outcomes After Stroke
CCR5   C-C chemokine receptor type 5
DAPTA   D-Ala-Peptide T-Amide
SRRR   Stroke Recovery and Rehabilitation Roundtable
STAIR   Stroke Treatment Academic Industry Roundtable
SYRCLE   Systematic Review Centre for Laboratory Animal Experimentation
TAK-779   Takeda 779

Introduction

C-C chemokine receptor type 5 (CCR5) is expressed across a variety of leukocyte subtypes,

endothelial cells, and cell types in the brain (e.g., neurons, microglia, astrocytes), and is thought

to play crucial roles in post-stroke neuroinflammation, blood-brain barrier repair, and neuronal

survival / repair processes.[1] CCR5 antagonists have emerged as potential therapeutic candidates

for stroke, demonstrating both cerebroprotection and improved neural repair/recovery in

preclinical animal models.[2–6] However, no CCR5 antagonist drug has an approved indication in

the stroke context, necessitating studies to establish safety and efficacy this population. This has

led to an ongoing clinical trial to investigate efficacy of CCR5 antagonists in combination with

post-stroke rehabilitation.[7] Assessment of the preclinical evidence supporting CCR5's role in

stroke is needed to identify areas of potential benefit, and knowledge gaps, that should be

addressed by future preclinical research.[8]

The stroke cerebroprotection and recovery communities have advocated for alignment of

preclinical and clinical study parameters through publication of consensus recommendations for

preclinical research, in an effort to enhance the translation of new stroke therapies.[9–11] Examples

include identification of more sensitive and clinically relevant preclinical outcome measures and

incorporation of potentially important effect modifiers of treatment efficacy, such as age, sex,

and stroke-related comorbidities (hypertension, diabetes, etc.).[9–11] These recommendations aim

to improve the translation of novel stroke therapeutics from preclinical to clinical populations,

but the degree to which preclinical evidence for CCR5 antagonists satisfy these

recommendations is unknown.

We sought to comprehensively evaluate the preclinical evidence for CCR5 antagonist drugs as

both cerebroprotective and stroke recovery-promoting agents.[9,11] Both perspectives are necessary

to fully understand the therapeutic potential of stroke-related treatments, as distinct biological

principles, time windows for treatment, and outcomes of interest underpin each of these

treatment domains.[12,13] We conducted a systematic review and meta-analysis of the CCR5

literature in conjunction with a panel of individuals with lived experiences of stroke. This review

sought to explore how the preclinical evidence for CCR5 antagonist drugs aligned with guidance

for preclinical stroke research provided by previous expert committees and the parameters of an

ongoing clinical trial (The Canadian Maraviroc Randomized Controlled Trial To Augment

Rehabilitation Outcomes After Stroke, CAMAROS, NCT04789616).[9–11]

Results

Study selection

Our search identified 263 citations, which was reduced to 166 unique studies after deduplication.

Five studies representing 10 experiments met the eligibility criteria (Figure 1).[2–6] Herein,

"studies" refer to the published articles as a unit, while "experiments" refer to distinct

investigations within each published article used to test various hypotheses (i.e., a subunit of

"studies" comprised of a select cohort of animals).

Study and animal model characteristics

Most studies used ischaemic stroke (n=4/5). This was induced via intraluminal suture (n=2),

cauterization (n=1), or photothrombosis techniques (n=1; Table 1). Haemorrhagic stroke was

induced in one study via autologous whole blood injection. All studies used mouse (n=4) or rat

(n=1) models, comprised of exclusively male animals (n=5). Relevant stroke comorbidities

highlighted by patient partners and STAIR/SRRR recommendations (e.g., hypertension,

diabetes) were not used in any study.

Intervention characteristics

Maraviroc was used in six of the experiments (n=6/10), TAK-779 in three of the experiments

(n=3), and D-Ala-Peptide T-Amide (DAPTA) in one (n=1; Table 2). These CCR5 antagonists

were delivered intraperitoneally (n=4), intranasally (n=2), intracerebroventricularly (n=2),

subcutaneously (n=1), and intravenously (n=1) at a dose range from 0.01 to 100 mg/kg. Most

studies delivered a single dose of the drug (n=6); experiments with multiple administrations

(n=4) ranged from 3–63 doses. Time of initial treatment administration varied widely. Two

studies (3 experiments) administered treatment pre-stroke (10-15 minutes),[2,3] and 4 studies (6

experiments) in the acute, potentially cerebroprotective, post-stroke period (50 minutes – 24

hours post-stroke).[3–6] One study (1 experiment) was conducted in the late sub-acute/early chronic

period beginning at three to four weeks post stroke, which would be oriented towards recovery,

rather than cerebroprotective, effects.[6] Patient partners had identified several *a priori* interests

(physical therapy alongside CCR5 administration, spasticity), which were also aligned with

SRRR recommended considerations for preclinical stroke recovery studies.[9] These were not

reported in any included studies. We also extracted a list of outcomes used to determine CCR5's

potential mechanisms of action (Appendix 1-figure 1, Appendix 1-table 1).

Meta-analysis of infarct volume

Infarct volume was reported in six experiments (n=6/10) from four different studies with an

overall pooled analysis demonstrating marked cerebroprotection with CCR5 antagonists (SMD -

1.02, 95% CI -1.58 to -0.46, p < 0.0001, I^2=34%; Figure 2). Five of these experiments measured

infarct volume at 1-3 days post-stroke, and one experiment measured infarct volume at a delayed

time point of 63 days post-stroke. No significant differences between pre- or post-stroke

administration were observed (P=0.47). Post-hoc sensitivity analysis removing one experiment

with extreme values[2] demonstrated that cerebroprotection in the remaining two experiments

remained statistically significant while reducing heterogeneity (SMD -0.81, 95% CI -1.25 to -

0.37, p < 0.001, I^2=0%). A second sensitivity analysis excluded one study that measured infarct

volume in mm^3 so that all other studies could be meta-analyzed using mean differences on the

percentage scale (Figure 2-figure supplement 1; MD -9.1%, 95% CI -11.6 to -6.7%, p < 0.001,

I^2=0%). This demonstrated a similar cerebroprotective effect as the other analyses. Further sub-

group analyses by route of administration, time of administration, stroke model, species, CCR5

antagonist, dose, and whether behaviour tests were assessed are described in Figure 2-figure

supplement 2, Figure 2-figure supplement 3, and Figure 2-figure supplement 4.

Synthesis of behavioural outcomes without meta-analysis

Motor behavioural outcomes were reported in six experiments from three studies and represented

seven different behavioural tasks. An additional study reported motor behavioural outcomes

without standard deviations or standard errors, and thus could not be included (authors did not

respond to email requests for data).[4] A cognitive outcome (Morris Water Maze) was measured in

one study. Overall, CCR5 inhibition was effective in 11 of 16 behavioural outcomes tested

(Figure 3). Meta-analysis and planned subgroup analysis were contra-indicated due to an

inadequate number of studies for each given outcome measure, which necessitated the synthesis

without meta-analysis presented below.[14]

Behavioural outcomes are presented by time of CCR5 antagonist administration, as discussed in

the Intervention Characteristics section above, as administration time directly relates to the

treatment context and patient population to which the results apply (i.e., acute cerebroprotection

vs. late sub-acute/early chronic neural repair). Li et al. reported that pre-stroke administration

(relevant to surgical contexts with high risk of thrombosis) of DAPTA did not result in

significantly greater performance on the neurological deficit score.[2]

Regarding acute post-stroke administration times with the potential for *cerebroprotective* effects

(up to 24 hours post-stroke based on observed infarct reductions in Figure 2), Yan et al. observed

that CCR5 antagonist (maraviroc) administration one-hour following stroke resulted in greater

motor performance on the corner test, limb placement, modified Garcia, foot fault, and rotarod

tasks compared to vehicle-treated controls. Cognitive outcomes in the Morris Water Maze task

(proportion of time spent in the probe quadrant) were also improved by this one-hour post-stroke

administration. Significantly improved motor performance was not observed if CCR5 antagonist

was administered 24 hours post-stroke, using this model of *intracerebral haemorrhage* (Figure

3).[5] In contrast, using a *focal ischaemia* model, Joy et al. observed that CCR5 antagonist

(maraviroc) administration initiated 24 hours post-stroke and continued daily for 3 weeks (which

could potentially represent a mix of cerebroprotective and recovery-induced effects) resulted in

greater performance on the cylinder and grid walk tasks compared to vehicle-treated controls.

Joy et al., also performed one experiment with *late sub-acute/early chronic* administration (a

potentially critical time for neural repair) initiated at three to four weeks post-stroke and

continued daily for 11 weeks, that demonstrated significantly improved performance for the grid

walk, but not cylinder, task.[6] This experiment was initiated outside the plausible range for a

cerebroprotective effect, implying behavioural improvement involved *recovery-promoting*

mechanisms. However, equivalent infarct volumes were not demonstrated between the treated

and control groups in this cohort, which could potentially lead to confounding effects.

Risk of bias

All articles had a 'high' risk of bias in at least one domain of the SYRCLE tool.[15] Most domains

within each study demonstrated an 'unclear' risk of bias. All studies reported randomizing

animals; however, as commonly observed in the preclinical literature,[16–18] only one of these

studies provided sufficient detail to ensure that the randomization method had a low risk of bias.

The SYRCLE domains with the highest risk of bias were incomplete outcome data, with 80% of

studies (n=4/5) failing to provide complete data for all animals initially included in the study, as

well as selective outcome reporting, with 60% (n=3/5; all studies of maraviroc) not providing

complete data for all expected outcomes discussed in the methods (Figure 4).

Comprehensiveness of preclinical evidence and alignment with clinical trials

We assessed comprehensiveness of the preclinical evidence using the STAIR and SRRR

recommendations[9,11,19,20] as well as alignment with study parameters of the CAMAROS trial.[7] A

summary of assessment items from STAIR XI is provided in Table 3, with additional items from

STAIR I, VI, and SRRR recommendations included in Table 4.

For CCR5 antagonists as a post-stroke cerebroprotectant, the overall body of evidence satisfied

all five STAIR XI domains assessing 'candidate treatment qualification' (Table 3). Overall, a

range of doses and clinically relevant administration times for cerebroprotection were evaluated

across a variety of motor and cognitive behavioural domains. All studies tested both behavioural

and histological outcomes and demonstrated cerebroprotective effects, but most studies failed to

measure and control post-stroke temperature, which could potentially confound the observed

cerebroprotection (Table 4).[21] Most histological measurements were also assessed at <72 hours,

which could confound the observed cerebroprotective effects if cell death was merely delayed.[21]

For CCR5 antagonists as a post-stroke recovery-inducing treatment, one experiment assessed the

effects of initiating CCR5 administration in a similar post-stroke phase as the CAMAROS trial.

This experiment (Joy et al.)[6] did not demonstrate that each treatment group had equivalent

baseline stroke volumes, which may potentially confound observed behavioural effects.

Furthermore, the maximum dose used in mice (100 mg/kg) was not sufficient to attain

cerebrospinal fluid levels of maraviroc observed in humans using the CAMAROS dosing regime

(mice: 13.8±5.4 ng/mL; humans 33.6–60 ng/mL).[6]

Areas of concern were identified in all STAIR XI domains assessing 'preclinical assessment and

validation' (Table 3). Although adult animals were used in most of the preclinical studies, the

reported weights and ages of these animals corresponded to young adults, rather than the aged

adults that better represent the stroke population. All experiments used only male rodents that

were free of common stroke comorbidities. It was also unclear if behavioural testing was

performed during the inactive circadian phase or active (dark) phase, which could result in

confounding if CCR5 antagonists affect arousal of animals during their inactive period.[22]

Furthermore, none of the studies had their protocols or results directly replicated by an

independent laboratory, or across multiple sites in a multilaboratory study.[9,23] Regarding stroke

recovery, no studies assessed behavioural effects on upper extremity skilled reaching / grasping

or potential interactions of CCR5 antagonists with rehabilitative therapies or established

recanalization procedures (Table 4).[24–27] These elements are highly relevant to the CAMAROS

trial, as one of the primary outcomes of this trial is upper extremity performance on the Fugl-

Meyer and maraviroc administration will be paired with an 8-week exercise program. These

findings were supported by the PRIMED[2] tool, which resulted in a Readiness for Translation

Score of "medium" (on a scale of "low", "medium", "high"). This tool highlighted similar

promising elements, as well as weaknesses, as our analysis above (Table 3, Table 4), identifying

the limited preclinical evidence for the effects of CCR5 antagonists in clinically relevant sexes,

ages, species, and disease comorbidities, without sufficient dose-response information to inform

trials (Appendix 2-table 1). Overall, our assessments highlight a variety of knowledge gaps,

potential confounding factors, and areas of misalignment between the preclinical evidence and

clinical trial parameters that could be improved with further preclinical experimentation.

Discussion

The overall body of preclinical evidence for CCR5 antagonists in stroke demonstrates potential

acute cerebroprotection with corresponding impairment reduction, as well as improved

functional recovery in the sub-acute/early chronic phase. Our systematic review also highlights

evidence gaps that could impact successful clinical translation of CCR5 antagonists. Our analysis

of 10 independent experiments, identified that acute administration of CCR5 antagonists within

the first 24 hours post-stroke was associated with a marked reduction in infarct volume. This

cerebroprotective reduction of infarct volume did not significantly vary based on treatment dose

or any other experimental characteristics.[3,4,6] Overall, the majority of behavioural effects

appeared to be in a positive direction, but the low number of included studies precluded meta-

analysis of these results. Indeed, no individual behavioural experiment included more than ten

CCR5-treated animals, and given the wide range of dosages, timings, routes, stroke models, and

rodent strains involved, the certainty of these findings is limited and should be interpreted

cautiously. Pooling data across heterogenous experimental designs, animal/stroke models, and

treatment parameters, as we have done with the infarct volume analysis in the present study, can

introduce variability that increases the risk of overestimating or underestimating the true effect of

the intervention.[28] Treatment effects observed across model systems and therapeutic compounds

may represent different biological mechanisms. Despite this potential limitation, meta-analysis

can provide valuable insights, especially in preclinical settings where the sample sizes of

individual studies may be too small to detect significant effects on their own. In these cases,

pooling data across studies can help identify overarching estimates of benefits and harm,

highlight subgroups of interest, and help guide areas of future research. As described in the

results above, we attempted to mitigate the risks of inappropriate data pooling through careful

investigation of heterogeneity, subgroup analyses, and differentiation between outcomes where

we felt that meta-analytic pooling was (infarct volume) and was not (behavioural outcomes)

appropriate. Overall, we believe that our results indicate that further investigation is warranted to

determine the optimal timing of administration and behavioural domains under which CCR5

antagonists exhibit the strongest post-stroke cerebroprotective and recovery-inducing effects.

Despite the positive direction of treatment effects across all studies of CCR5 antagonists, we

found a substantial risk of bias in the underlying studies.[15] As is commonly observed in the

preclinical literature, all studies either did not adequately report their randomization / blinding

methods and exhibited evidence of selective / incomplete reporting.[16–18] Such features are

associated with biased overestimations of preclinical treatment efficacy, which raises further

concerns about the reliability and validity of the present findings.[29–31] Future studies should

carefully incorporate all elements of the ARRIVE 2.0 guidelines to help ensure that results are

transparently reported and improve confidence in the findings.[32]

Comprehensiveness of the preclinical evidence for CCR5 antagonists was assessed in relation to

STAIR and SRRR consensus recommendations.[9,11,19,20] These recommendations aim to provide

investigators and regulators with "assurance that the candidate treatment shows signals of

efficacy and safety, before embarking on an expensive clinical development program".[11] The

included studies provide good initial evidence for acute cerebroprotection, as well as mechanistic

and behavioural evidence for enhanced recovery in the late sub-acute/early chronic post-stroke

phase. However, demonstration of efficacy under a wider range of conditions, such as in aged

animals, females, animals with stroke-related comorbidities, more clinically relevant timing of

dose administrations, or in conjunction with rehabilitative therapies are necessary to provide

further confidence in these findings. In addition, all studies used unique doses, timings, and

outcomes, so independent replication of the most promising study parameters would further

increase certainty in the evidence.[9,11] Future preclinical research should aim to address these

evidence gaps to further increase the clinical relevance and comprehensiveness of evidence for

CCR5 antagonists in stroke.

In relation to the ongoing CAMAROS trial assessing maraviroc in the subacute post-stroke

phase,[7] the most relevant preclinical evidence comes from one experiment within the Joy et al.

study[6] where maraviroc was initially administered at three to four weeks post-stroke and

continued daily for 11 weeks. This experiment demonstrated that administration of maraviroc in

the late sub-acute/early chronic post-stroke phase improved functional recovery on the grid walk,

but not cylinder, task. These experimental conditions could potentially align with the putative

therapeutic window and outcomes of interest being assessed in CAMAROS (e.g. 10-minute walk

test co-primary outcome).[7] However, caution is warranted as this pivotal supporting preclinical

evidence is based on a low sample size (n=9). Moreover, potential differences in dosing, severity

of infarct, concomitant rehabilitative therapy, and other factors discussed above could influence

the degree to which these results successfully translate to the clinical environment. Finally,

clinically relevant secondary outcomes in the CAMAROS trial, such as cognitive and emotional

domains as measured by the Montreal Cognitive Assessment (MoCA) and Stroke Aphasia

Depression Questionnaire (SADQ) were not modelled in the preclinical literature. Although one

study included a cognitive outcome, the other treatment parameters of this study were not

aligned to the CAMAROS trial.[5] Future preclinical studies should assess a more diverse and

comprehensive battery of clinically relevant behavioural tasks, which could be based on the

range of outcomes employed in the CAMAROS trial, or those found in the SRRR

recommendations.[9] Nevertheless, the Joy et al., study provides a plausible biological mechanism

and "proof of concept" for how CCR5 antagonism might enhance neuroplasticity that improves

functional recovery after stroke.

Our present synthesis of the preclinical evidence for CCR5 antagonists used novel approaches to

increase its utility for assessing certainty of the findings and identification of knowledge gaps.

First, we engaged patients with lived experiences of stroke throughout the review process to

ensure that our research questions, outcomes, and interpretations aligned with the priorities of the

ultimate end-user of stroke research. Second, we incorporated consensus recommendations for

both preclinical cerebroprotection and recovery research as an evidence evaluation tool, which

we found often aligned with the priorities of our patient panel.[9,11,19,20] This guided assessment of

the alignment of preclinical evidence with parameters of ongoing clinical trials, as well as

appraisal of comprehensiveness of the preclinical evidence with a focus on translational

validity[33] rather than only internal validity and risk of bias.[15] We also used this method to

provide concrete avenues for future preclinical studies to close knowledge gaps and improve

certainty in the effects of CCR5 antagonists under clinically relevant experimental conditions.

Similar approaches should be considered by future preclinical systematic reviews to improve

interpretation of the preclinical evidence from a translational perspective.

In conclusion, CCR5 antagonists show promise in preclinical studies for stroke

cerebroprotection, corresponding reduction in impairment, as well as improved functional

recovery related to neural repair in the late sub-acute/early chronic phase. However, high risk of

bias and the limited (or no) evidence in clinically relevant domains underscore the need for more

rigorous and transparent preclinical research to further strengthen the current preliminary

evidence available in the literature. Addressing these concerns will not only enhance the

reliability of preclinical evidence but also better inform the design and execution of clinical trials

of CCR5 antagonists, such as the ongoing CAMAROS trial.[7] The integration of expert

recommendations, such as STAIR and SRRR, should guide future preclinical investigations and

synthesis of the body of preclinical evidence in stroke recovery research.[9,11] Our present

approach serves as a template by which the preclinical evidence supporting translation to clinical

trials can be weighed when justifying early clinical trials of novel interventions for stroke

cerebroprotection and recovery.

Materials and methods

We registered the review protocol on the International Prospective Register of Systematic

Reviews (CRD42023393438).[34] The findings are reported in accordance with the Preferred

Reporting Items for Systematic Reviews and Meta-Analyses (see attached Reporting Standards

Document)[35] and Guidance for Reporting the Involvement of Patients and the Public (Appendix

3-table 1).[36]

Engagement with patient panel - individuals with lived experience of stroke

A panel of eight patients and caregivers with lived experience of a stroke informed project

development (e.g., research question development, review protocols, search strategy

development) and were actively involved in the research conduct (screening, data extraction,

analysis, interpretation). Monthly meetings occurred with the patients and caregivers to provide

educational sessions of background knowledge of preclinical stroke, systematic review conduct,

and discuss research findings as the review progressed. We co-developed a terms of reference

document *a priori* to document details of the engagement (i.e., roles, responsibilities,

expectations, project goals, etc.). The patient partners co-developed the research question to

address patient interests, including chronic stroke recovery (i.e., the panel was particularly

interested in evaluating the effects of extended drug administration), consideration of physical

therapy in tandem with drug administration, inclusion of stroke-relevant comorbidities,

spasticity, and motor and cognitive outcomes. Co-authorship and financial compensation were

agreed upon with the patients and caregivers and offered as a method of acknowledgement

according to the Canadian Institutes of Health Research (CIHR) Strategies for Patient Oriented

Research (SPOR) Evidence Alliance Patient Partner Appreciation Policy.

Eligibility criteria

- *Animals:* Any preclinical *in vivo* animal models of adult stroke were included. Human,

invertebrate, *in vitro*, *ex vivo*, and neonatal animal studies were excluded.

- *Model:* Focal ischaemic or intracerebral haemorrhagic stroke models were included,

while animal models of global ischaemia were excluded.

- *Intervention:* Studies administering a CCR5 antagonist drug (e.g., maraviroc, D-Ala-

Peptide T-Amide (DAPTA), Takeda 779 (TAK-779)). Study arms in which CCR5 was

genetically manipulated (e.g., CCR5 knockout strain) were excluded.

- *Comparator:* Vehicle-treated control groups where stroke was induced. CCR5 antagonist

control groups without stroke were excluded.

- *Outcome:* Studies reporting at least one of the following: infarct size, behavioural tests,

mortality, adverse events, and spasticity were included.

- *Study design and publication characteristics:* Controlled interventional studies

(randomized, pseudo-randomized, or non-randomized) published as full journal articles

in any language or year were included. Abstracts, review articles, opinion-based

letters/editorials, and unpublished grey literature were excluded.

Information sources and search strategy

An information specialist with experience in systematic searches of the preclinical literature

developed a comprehensive search strategy based on a previously published strategy for

identifying animal experimentation studies (Appendix 4).[37] The search strategy underwent peer-

review using the Peer Review of Electronic Search Strategies (PRESS) checklist.[38] We searched

MEDLINE (OVID interface, including In-Process and Epub Ahead of Print), Web of Science,

and Embase (OVID interface). The search was conducted October 25, 2022, to align with the

listed launch date of the CAMAROS trial (September 15, 2022). Our intention in doing so was to

collate and assess all preclinical evidence that could have feasibly informed the clinical trial. We

sought to assess the comprehensiveness of evidence and readiness for translation of CCR5

antagonist drugs at the time of their actual translation into human clinical trials, as well as the

alignment of the CAMAROS trial design to the existing preclinical evidence base.

Selection process and data collection

We deduplicated citations and uploaded them into DistillerSR® (Evidence Partners, Ottawa,

Canada). Two reviewers independently screened citations by title and abstract using an

accelerated method (one reviewer required to include, two reviewers required to exclude). We

then screened and extracted data from full-text articles in duplicate. Graphical data was extracted

using Engauge Digitizer.[39] A third reviewer with content expertise in preclinical stroke studies

audited all data extraction. Conflicts between reviewers were resolved by consensus discussion.

See Appendix 5 for the complete list of data extraction elements.

Effect measures and data synthesis

We performed quantitative analyses using the R (version 4.1.2) "metafor" package (version

4.0.0)[40] with inverse variance random effects modelling. We expressed continuous outcome

measures as standardized mean differences (SMDs) with 95% confidence intervals (CIs) and

assessed statistical heterogeneity of effect sizes using the Cochrane I^2 statistic.[41] This was

necessary due to the variety of outcome measures and measurement scales used across studies,

which is a common feature of preclinical systematic reviews. Sensitivity analyses were

performed using original measurement scales where possible. From patient partner input,

subgroups were analyzed based on timing / dose / route of intervention, stroke model, stroke

type, species, type of behavioural outcome (i.e., motor, cognitive), and comorbidities.[42] Our

planned subgroup analyses on study quality, specific regions/areas of the brain, and post-stroke

rehabilitation paradigms were not performed due to insufficient number of studies. We did not

assess publication bias using Egger plots due to fewer than 10 studies being included in the

analysis, as per Cochrane recommendations.[43]

Risk of bias assessment

Two independent reviewers used the Systematic Review Centre for Laboratory Animal

Experimentation (SYRCLE) risk of bias tool to assess each study as having a "Low Risk",

"Unclear Risk", "Some Concerns of Risk", or "High Risk" across domains such as

randomization, blinding and outcome reporting.[15] "Some Concerns of Risk" indicated reporting

of a domain (i.e., randomization), but lacking methodological details. This differed from

"Unclear Risk" where there was no mention of the domain in the study.

Comprehensiveness of preclinical evidence and alignment with clinical trials

We assessed comprehensiveness of the *overall body* of preclinical evidence for CCR5

antagonists as a cerebroprotective treatment using The Stroke Treatment Academic Industry

Roundtable (STAIR) I, VI, and XI Consolidated Recommendations.[11,19,20] These

recommendations encompass "candidate treatment qualification" (e.g., dose, timing of dose,

outcomes) and "preclinical assessment and validation" (e.g., age, sex, sample size, animal

type).[11] We excluded domains that were redundant with risk of bias (e.g., randomization,

blinding, etc.) and included additional items relevant to stroke recovery studies from the Stroke

Recovery and Rehabilitation Roundtable (SRRR) Translational Working Group consensus

recommendations.[9] Two reviewers extracted data to determine if the overall evidence across

studies satisfied each of the STAIR and SRRR recommendations. A third reviewer audited this

analysis. We then assessed alignment of existing preclinical evidence with an ongoing clinical

trial of CCR5 antagonists for stroke (CAMAROS, NCT04789616).[7]

Protocol deviations

Additionally, a list of outcomes used to determine CCR5's potential mechanisms of action were

extracted (Appendix 1-figure 1, Appendix 1-table 1) based on feedback from individuals that

reviewed the initial manuscript drafts. We also incorporated an additional assessment using the

PRIMED[2] tool for assessing the readiness of stroke cerebroprotection therapies to be translated to

clinical trials based on the body of preclinical evidence (Appendix 2-table 1).[44]

Acknowledgements: The authors thank Hannah Laquerre for her assistance with figure

generation. Appendix 1-figure 1 was created with BioRender.com.

Sources of Funding: This study was funded by a Social Accountability Grant from the

University of Ottawa's Faculty of Medicine. MSJ was supported by the Canadian Institutes of

Health Research (CIHR) and Vanier Canada Graduate Scholarship (CGV-186957). The funders

were not involved in the study design, collection, analysis and interpretation of data, writing the

manuscript, or in the decision to submit the article for publication. MML is supported by

The Ottawa Hospital Anesthesia Alternate Funds Association, a University of Ottawa Junior

Research Chair, and the Canadian Anesthesiologists' Society Career Scientist Award.

Disclosures: None.

Data Availability: All data generated or analysed during this study are included in the

manuscript and supporting files. Source data and code to generate Figures 2 and 3 have been

provided at the following DOI: https://doi.org/10.17605/OSF.IO/5KGT6

References:

1. Jing K, Chen F, Shi X, Guo J, Liu X. Dual effect of C-C motif chemokine receptor 5 on ischemic stroke: More harm than benefit? *European Journal of Pharmacology*. 2023;953:175857.

2. Li L, Zhi D, Shen Y, Liu K, Li H, Chen J. Effects of CC-chemokine receptor 5 on ROCK2 and P-MLC2 expression after focal cerebral ischaemia-reperfusion injury in rats. *Brain injury*. 2016;30:468–473.

3. Takami S, Minami M, Katayama T, Nagata I, Namura S, Satoh M. TAK-779, A Nonpeptide CC Chemokine Receptor Antagonist, Protects the Brain Against Focal Cerebral Ischemia in Mice. *Journal of cerebral blood flow and metabolism*. 2002;22:780–784.

4. Chen B, Cao P, Guo X, Yin M, Li X, Jiang L, Shao J, Chen X, Jiang C, Tao L, et al. Maraviroc, an inhibitor of chemokine receptor type 5, alleviates neuroinflammatory response after cerebral Ischemia/reperfusion injury via regulating MAPK/NF-κB signaling. *International immunopharmacology*. 2022;108:108755–108755.

5. Yan J, Xu W, Lenahan C, Huang L, Wen J, Li G, Hu X, Zheng W, Zhang JH, Tang J. CCR5 activation promotes NLRP1-dependent neuronal pyroptosis via CCR5/PKA/CREB pathway after intracerebral hemorrhage. *Stroke*. 2021;52:4021 EP – 4032.

6. Joy MT, Ben Assayag E, Shabashov-Stone D, Liraz-Zaltsman S, Mazzitelli J, Arenas M, Abduljawad N, Kliper E, Korczyn AD, Thareja NS, et al. CCR5 Is a Therapeutic Target for Recovery after Stroke and Traumatic Brain Injury. *Cell*. 2019;176:1143-1157.e13.

7. Dukelow S, Clinicaltrials.gov NCT04789616. The Canadian Maraviroc RCT To Augment Rehabilitation Outcomes After Stroke (CAMAROS) [Internet]. [cited 2023 Oct 20];Available from: https://clinicaltrials.gov/study/NCT04789616

8. O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW. 1,026 experimental treatments in acute stroke. *Ann Neurol*. 2006;59:467–477.

9. Corbett D, Carmichael ST, Murphy TH, Jones TA, Schwab ME, Jolkkonen J, Clarkson AN, Dancause N, Weiloch T, Johansen-Berg H, et al. Enhancing the Alignment of the Preclinical and Clinical Stroke Recovery Research Pipeline: Consensus-Based Core Recommendations From the Stroke Recovery and Rehabilitation Roundtable Translational Working Group. *Neurorehabilitation and neural repair*. 2017;31:699–707.

10. Bosetti F, Koenig JI, Ayata C, Back SA, Becker K, Broderick JP, Carmichael ST, Cho S, Cipolla MJ, Corbett D, et al. Translational Stroke Research: Vision and Opportunities. *Stroke*. 2017;48:2632–2637.

11. Lyden P, Buchan A, Boltze J, Fisher M. Top Priorities for Cerebroprotective Studies-A Paradigm Shift: Report From STAIR XI. *Stroke*. 2021;52:3063–3071.

12. Carmichael ST. Emergent properties of neural repair: elemental biology to therapeutic
concepts. *Annals of neurology*. 2016;79:895–906.

13. Murphy TH, Corbett D. Plasticity during stroke recovery: from synapse to behaviour. *Nature
reviews. Neuroscience*. 2009;10:861–872.

14. Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-
Boyce J, Ryan R, Shepperd S, Thomas J, et al. Synthesis without meta-analysis (SWiM) in
systematic reviews: reporting guideline. *BMJ (Online)*. 2020;368:l6890–l6890.

15. Hooijmans CR, Rovers MM, de Vries RB, Leenaars M, Ritskes-Hoitinga M, Langendam
MW. SYRCLE's risk of bias tool for animal studies. *BMC Medical Research Methodology*.
2014;14:43.

16. Avey MT, Moher D, Sullivan KJ, Fergusson D, Griffin G, Grimshaw JM, Hutton B, Lalu
MM, Macleod M, Marshall J, et al. The Devil Is in the Details: Incomplete Reporting in
Preclinical Animal Research. *PLoS One*. 2016;11:e0166733.

17. Fergusson DA, Wesch NL, Leung GJ, MacNeil JL, Conic I, Presseau J, Cobey KD, Diallo J-
S, Auer R, Kimmelman J, et al. Assessing the Completeness of Reporting in Preclinical
Oncolytic Virus Therapy Studies. *Mol Ther Oncolytics*. 2019;14:179–187.

18. Fergusson DA, Avey MT, Barron CC, Bocock M, Biefer KE, Boet S, Bourque SL, Conic I,
Chen K, Dong YY, et al. Reporting preclinical anesthesia study (REPEAT): Evaluating the
quality of reporting in the preclinical anesthesiology literature. *PLoS One*.
2019;14:e0215221.

19. Finklestein SP, Fisher M, Furlan AJ, Goldstein LB. Recommendations for standards
regarding preclinical neuroprotective and restorative drug development. *Stroke*.
1999;30:2752–2758.

20. Fisher M, Feuerstein G, Howells DW, Hum PD, Kent TA, Savitz SI, Lo EH. Update of the
Stroke Therapy Academic Industry Roundtable Preclinical Recommendations. *Stroke*.
2009;40:2244–2250.

21. Corbett D, Nurse S. The problem of assessing effective neuroprotection in experimental
cerebral ischemia. *Progress in neurobiology*. 1998;54:531–548.

22. MacLellan CL, Keough MB, Granter-Button S, Chernenko GA, Butt S, Corbett D. A Critical
Threshold of Rehabilitation Involving Brain-Derived Neurotrophic Factor Is Required for
Poststroke Recovery. *Neurorehabilitation and neural repair*. 2011;25:740–748.

23. Hunniford VT, Grudniewicz A, Fergusson DA, Montroy J, Grigor E, Lansdell C, Lalu MM.
A systematic assessment of preclinical multilaboratory studies and a comparison to single
laboratory studies. *eLife*. 2023;12.

24. Balbinot G, Schuch CP, Jeffers MS, McDonald MW, Livingston-Thomas JM, Corbett D. Post-stroke kinematic analysis in rats reveals similar reaching abnormalities as humans. *Scientific reports*. 2018;8:8738–13.

25. Jeffers MS, Corbett D. Synergistic Effects of Enriched Environment and Task-Specific Reach Training on Poststroke Recovery of Motor Function. *Stroke*. 2018;49:1496–1503.

26. Jeffers MS, Karthikeyan S, Gomez-Smith M, Gasinzigwa S, Achenbach J, Feiten A, Corbett D. Does Stroke Rehabilitation Really Matter? Part B: An Algorithm for Prescribing an Effective Intensity of Rehabilitation. *Neurorehabilitation and neural repair*. 2018;32:73–83.

27. Wechsler LR, Adeoye O, Alemseged F, Bahr-Hosseini M, Deljkich E, Favilla C, Fisher M, Grotta J, Hill MD, Kamel H, et al. Most Promising Approaches to Improve Stroke Outcomes: The Stroke Treatment Academic Industry Roundtable XII Workshop. *Stroke (1970)*. 2023;54:3202–3213.

28. Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M, Welch V. Chapter 15: Interpreting results and drawing conclusions [Internet]. In: Cochrane Handbook for Systematic Reviews of Interventions version 6.4 (updated August 2023). Cochrane; 2023. Available from: www.training.cochrane.org/handbook

29. Holman C, Piper SK, Grittner U, Diamantaras AA, Kimmelman J, Siegerink B, Dirnagl U. Where Have All the Rodents Gone? The Effects of Attrition in Experimental Research on Cancer and Stroke. *PLoS Biology*. 2016;14:e1002331–e1002331.

30. Sena ES, van der Worp HB, Bath PMW, Howells DW, Macleod MR. Publication bias in reports of animal stroke studies leads to major overstatement of efficacy. *PLoS Biology*. 2010;8:e1000344–e1000344.

31. Macleod MR, Bart Van Der Worp H, Sena ES, Howells DW, Dirnagl U, Donnan GA. Evidence for the Efficacy of NXY-059 in Experimental Focal Cerebral Ischaemia Is Confounded by Study Quality. *Stroke*. 2008;39:2824–2829.

32. Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A, Cuthill IC, Dirnagl U, et al. The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research. *PLoS Biol*. 2020;18:e3000410.

33. Drude NI, Martinez Gamboa L, Danziger M, Dirnagl U, Toelch U. Improving preclinical studies through replications. *eLife*. 10:e62101.

34. Lalu MM, Sharif A, Nicholls S, Dowlatshahi D, Fergusson DA, Corbett D, Bapuji R, Jeffers MS, Humphrey J, Johnston W, et al. Evaluating the effect of CCR5 inhibitors in animal models of stroke: A systematic review co-designed with patient partners [Internet]. 2023 [cited 2023 Oct 20];Available from: https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=393438

35. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L,
Tetzlaff JM, Akl EA, Brennan SE, et al. The PRISMA 2020 statement: An updated guideline
for reporting systematic reviews. *PLoS medicine*. 2021;18:e1003583–e1003583.

36. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, Altman DG, Moher D,
Barber R, Denegri S, et al. GRIPP2 reporting checklists: tools to improve reporting of patient
and public involvement in research. *BMJ*. 2017;358:j3453–j3453.

37. Hooijmans CR, Tillema A, Leenaars M, Ritskes-Hoitinga M. Enhancing search efficiency by
means of a search filter for finding all studies on animal experimentation in PubMed.
Laboratory animals (London). 2010;44:170–175.

38. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer
Review of Electronic Search Strategies: 2015 Guideline Statement. *Journal of clinical
epidemiology*. 2016;75:40–46.

39. Mitchell M, Muftakhidinov B, Winchen T. Engauge Digitizer Software [Internet]. [cited
2023 Oct 20];Available from: http://markummitchell.github.io/engauge-digitizer/

40. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *Journal of
statistical software*. 2010;36.

41. Higgins JP, Li T, Deeks J. Chapter 6: Choosing effect measures and computing estimates of
effect [Internet]. In: Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch VA,
editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.4 (updated
August 2023). Cochrane; 2023 [cited 2023 May 1]. Available from:
www.training.cochrane.org/handbook

42. Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, Krakauer JW,
Boyd LA, Carmichael ST, Corbett D, et al. Agreed definitions and a shared vision for new
standards in stroke recovery research: The Stroke Recovery and Rehabilitation Roundtable
taskforce. *International journal of stroke*. 2017;12:444–450.

43. Page MJ, Higgins JP, Sterne JAC. Chapter 13: Assessing risk of bias due to missing results
in a synthesis [Internet]. In: Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch
VA, editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.4
(updated August 2023). Cochrane; 2023 [cited 2023 May 1]. Available from:
www.training.cochrane.org/handbook

44. Bahr-Hosseini M, Bikson M, Iacoboni M, Liebeskind DS, Hinman JD, Carmichael ST, Saver
JL. PRIMED2 Preclinical Evidence Scoring Tool to Assess Readiness for Translation of
Neuroprotection Therapies. *Translational stroke research*. 2022;13:222–227.

Figure Legends

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) flow

diagram.

Figure 2. CCR5 antagonists reduce infarct volume. Data is presented as a forest plot with

standardized mean differences and 95% confidence intervals. Effect sizes <0 favours drug

treatment and >0 favours control. Data is stratified by timing of CCR5 antagonist administration

(pre- or post-stroke induction). The 'RE Model for All Studies' represents a pooled estimate of

the CCR5 antagonist drug effect on infarct volume from all studies combined. Separate pooled

estimates are also reported for post-stroke and pre-stroke CCR5.

Figure 2-figure supplement 1. Sensitivity analysis for all included studies that reported infarct

volume on a percentage scale. Overall, both the original analysis using standardized mean

differences (Figure 2) and the sensitivity analysis demonstrate a significant cerebroprotective

effect of CCR5 antagonists. Data is presented as a forest plot with mean differences and 95%

confidence intervals. Effect sizes <0 favours CCR5 antagonist treatment and >0 favours control.

Data is stratified by timing of CCR5 antagonist administration (pre- or post-stroke induction).

The 'RE Model for All Studies' represents a pooled estimate of the CCR5 antagonist drug effect

on infarct volume from all studies combined. Separate pooled estimates are also reported for

post-stroke and pre-stroke CCR5. T (n) = number of animals in the CCR5 treatment group. C (n)

= number of animals in the control group.

Figure 2-figure supplement 2. Subgroup analysis for all included studies of pre-stroke and post-

stroke CCR5 antagonist administration that reported infarct volume. Each row represents pooled

estimate data from studies within that subgroup. Data is presented as a forest plot with a

standardized mean difference and 95% confidence intervals. The I^2 value represents the

statistical heterogeneity within each subgroup. Effect sizes <0 favours drug treatment and >0

favours control. The 'RE Model for All Studies' represents a pooled estimate of the CCR5

antagonist drug effect on infarct volume from all studies combined. Considering all six

experiments (irrespective of administration timing of CCR5 antagonist), subgroup analysis

demonstrated no difference in effect size when considering route of administration, stroke model,

species type, drug, dose, or whether behaviour tests were assessed.

Figure 2-figure supplement 3. Subgroup analysis for all included studies of post-stroke drug

administration of a CCR5 antagonist that reported infarct volume. Each row represents pooled

estimate data from studies within that subgroup. Data is presented as a forest plot with a

standardized mean difference and 95% confidence intervals. The I^2 value represents the

statistical heterogeneity within each subgroup. Effect sizes <0 favours drug treatment and >0

favours control. The 'RE Model for All Studies' represents a pooled estimate of the CCR5

antagonist drug effect on infarct volume from all post-stroke drug administration of a CCR5

antagonist studies combined. Subgroup analysis considering only studies that administered

CCR5 antagonists post-stroke induction demonstrated no difference in effect size when

considering any of the subgroups.

Figure 2-figure supplement 4. Subgroup analysis for all included studies of pre-stroke drug

administration of a CCR5 antagonist that reported infarct volume. Each row represents pooled

estimate data from studies within that subgroup. Data is presented as a forest plot with a

standardized mean difference and 95% confidence intervals. The I^2 value represents the

statistical heterogeneity within each subgroup. Effect sizes <0 favours drug treatment and >0

favours control. The 'RE Model for All Studies' represents a pooled estimate of the CCR5

antagonist drug effect on infarct volume from all pre-stroke drug administration of a CCR5

antagonist studies combined. Subgroup analysis of studies that administered CCR5 pre-stroke

induction suggested that infarct volume was reduced to a greater extent by the intraluminal

suture stroke model versus other models (P=0.04), in rats versus mice (P=0.01), with DAPTA

versus TAK-779 (P=0.01), and when behaviour tests were performed versus not. (P=0.01).

Figure 3. Synthesis without meta-analysis for all included preclinical CCR5 antagonist studies

that reported motor and/or cognitive behavioural outcomes. Data is presented as a forest plot

with a standardized mean difference and 95% confidence intervals. Effect sizes <0 favours drug

treatment and >0 favours control.

Figure 4. Modified risk of bias traffic light plot in accordance with the SYRCLE tool. Yellow represents an unclear risk of bias, green represents a low risk of bias, and red represents a high risk of bias. Blue represents some concerns of a risk of bias. The risk of bias was 'unclear' across all studies for the domains of baseline characteristics because of missing data in the studies, random housing because no details on this domain were reported, and random outcome assessment because no details of how cohorts of animals were selected to perform certain outcomes nor how the order of outcome assessment proceeded. Four studies did not report on allocation concealment, and two studies did not report on blinding investigators and outcome assessors and were deemed as having an 'unclear' risk of bias. 80% of studies exhibited a 'high' risk for incomplete outcome data. Similarly, three studies (60%) had a 'high' risk of selective outcome reporting since all expected outcomes discussed in the methods of the articles did not align with their results. Other potential sources of bias considered included the source of funding (industry funded), contamination of pooling drugs (additional treatment which might influence or bias the result), unit error analysis (all animals receiving the same intervention are caged together, but analysis was conducted as if every single animal was one experimental unit), design-specific risks of bias (reporting details of which animals performed the same or different outcomes), and the addition of new animals to replace dropouts from the original population. Two studies (40%) had a 'high' risk in at least one of these additional categories.

Tables

Table 1. Summary of study and animal model characteristics of included articles.

Author	Year	Country	Species	Strain	Stroke Type	Stroke Model	Sex	Weight	Age
Li et al.[2]	2016	China	Rat	Wistar	Ischaemic	Intraluminal suture	Male	260–300 g	N/A
Takami et al.[3]	2002	Japan	Mice	ddY	Ischaemic	Intraluminal suture	Male	N/A	4 weeks
Chen et al.[4]	2022	China	Mice	C57/BL6	Ischaemic	Permanent middle cerebral artery occlusion	Male	25–30 g	8–10 weeks
Yan et al.[5]	2021	China	Mice	CD1	Haemorrhagic	Intracerebral haemorrhagic	Male	30–40 g	N/A
Joy et al.[6]	2019	USA	Mice	C57/BL6	Ischaemic	Photothrombotic	Male	25–30 g	8–20 weeks

Table 2. Summary of intervention characteristics.

Author	Drug	Dose (mg/kg)	Route	Timing	Doses (total #)	Outcomes Measured (Treatment n/Control n)	Outcome Window (post-stroke)
Li et al.[2]	DAPTA (D-Ala-Peptide T-Amide)	0.01	SC	15 min pre-stroke	1	- Infarct volume (5/5) - Neurological deficit score (5/5)	24 h
Takami et al.[3]	TAK-779	5	ICV	10 min pre-stroke	1	- Infarct volume (10/6)	48 h
		50	ICV	10 min pre-stroke		- Infarct volume (13/6)	
		0.25	IV	50 min post-stroke		- Infarct volume (17/18)	
Chen et al.[4]	Maraviroc	20	IP	1.5 h, 24 h, and 48 h post-stroke	3	- Infarct volume (5/5) - Longa score (5/5) - Neurological deficit score (5/5)	72 h
Yan et al.[5]	Maraviroc	0.15	IN	1 h post-stroke	1	- Garcia test (6/6) - Limb placement (6/6) - Corner turn test (6/6)	72 h
						- Foot fault (8/8) - Rotarod (8/8)	3 weeks
						- Probe quadrant duration (8/8)	25 days
				24 h post-stroke		- Garcia test (6/6) - Limb placement (6/6) - Corner turn test (6/6)	72 h
Joy et al.[6]	Maraviroc	100	IP	24 h post-stroke through daily injections for 9 weeks	63	- Infarct volume (5/5)	9 weeks
						- Grid walk (10/10) - Forelimb (10/10)	8 weeks
				24 h post-stroke then daily for 3 weeks	21	- Grid walk (10/10) - Cylinder test (9/8)	9 weeks
				3 to 4 weeks post-stroke then daily for 11 weeks	56	- Grid walk (9/9) - Cylinder test (9/9)	11 weeks

Abbreviations: ICV = intracerebroventricular; IN = intranasal; IP = intraperitoneal; IV = intravenous; SC = subcutaneous

Table 3. Alignment of included preclinical studies with STAIR recommendations and CAMAROS trial parameters.

Recommendation	Overall preclinical evidence	CAMAROS trial alignment	Notes
Candidate treatment qualification			
(1) Establish treatment dose-response	Yes	No	• **Preclinical** – doses across all studies from 0.01 to 100 mg/kg • 6 doses tested for infarct volume, 3 doses for behavioural effects • Joy et al. demonstrated that 100 mg/kg of maraviroc in mice results in cerebrospinal fluid levels lower than in humans (13.8±5.4 vs. 33.6–60 ng/mL) • **CAMAROS** – participants take 300 mg doses twice daily for 8 weeks • **Interpretation** – maximum preclinical dose may not represent clinical levels
(2) Treatment given after clinically relevant delayed times (one to 4.5 hours post-stroke)	Yes	Yes	• **Preclinical** – 3/5 studies (all maraviroc) administered drug from 1 to 24 hours post-stroke (acute phase) • Joy et al. initiated maraviroc administration at 20 days post-stroke for 11 weeks (subacute phase) • **CAMAROS** – participants recruited between 5 days and 6 weeks post-stroke (acute / early subacute) • **Interpretation** – one experiment in a time window relevant to clinical trial
(3) Both histologic and behavioural outcomes at acute (1 to 3 days) and chronic (7 to 30 days) time points	Yes	Yes	• **Preclinical** – 2/5 studies included both histologic and behavioural outcomes • Mainly tasks of spontaneous movement (e.g., cylinder) or motor coordination (e.g., grid walk) • Tests ranged from 3 days (acute) to 11 weeks (chronic) post-stroke for all included studies, but short term assessment may only delay cell death[21] • **CAMAROS** - motor learning (Fugl-Meyer Upper Extremity Assessment Score and 10-Meter Walk Test Score) measured as the primary outcome at baseline, 4-week (late subacute), 8-week (late subacute), and 6-month (chronic) • **Interpretation** – challenging to directly compare preclinical and clinical motor tasks, but testing range is comparable. Most relevant Joy et al. experiment (subacute administration) did not measure histologic outcomes
(4) Treatment reaches presumed target and causes expected physiological effects that can be assessed with a clinically relevant biomarker	Yes	NA	• **Preclinical** – 5/5 studies demonstrated trends for CCR5 antagonism to reduce infarct volume. Plausible that CCR5 influences stroke-relevant mechanisms • **CAMAROS** – does not include outcomes to assess treatment mechanisms
(5) Treatment able to pass the blood-brain barrier	Yes	NA	• **Preclinical** – 1/5 studies used mass spectrometry to demonstrate that maraviroc is present in the brain and cerebrospinal fluid • **CAMAROS** – does not include outcomes to assess presence of drug in the brain
Preclinical assessment / validation			
(1) Aging/adult age	No	No	• **Preclinical** – 4/5 studies used rodents with weights/ages consistent with young adulthood; 0/5 studies used rodents ≥10 months old. • **CAMAROS** – Age ≥18 year old adult participants eligible for recruitment • **Interpretation** – Preclinical studies consistent with CAMAROS eligibility criteria; however, no preclinical study examined middle-aged or elderly animals that would be more representative of the trial and overall stroke

			populations
(2) Male and female animals	No	No	• **Preclinical** – 0/5 studies included female animals • **CAMAROS** – both male and female sexes eligible for recruitment • **Interpretation** – effects of CCR5 antagonists for stroke recovery in female animals represents a critical knowledge gap
(3) Animals with co-morbidities	No	No	• **Preclinical** – 0/5 studies included animals with common stroke co-morbidities (e.g., diabetes, hypertension, etc.) • **CAMAROS** - participants with stroke co-morbidities eligible for recruitment • **Interpretation -** effects of CCR5 antagonists for stroke recovery in animals with common stroke comorbidities represents a critical knowledge gap
(4) Evidence from two or more laboratories	Yes	No	• **Preclinical** – 5/5 studies demonstrated stroke-related benefits; however, no study had their results replicated by independent laboratories • **CAMAROS** – trial protocol will be replicated by multiple study sites • **Interpretation** – future preclinical experiments could consider assessing effects using a multilaboratory approach, or independently replicate the effects of existing preclinical studies
(5) Gyrencephalic species	No	No	• **Preclinical** – 5/5 studies conducted in lissencephalic (rodent) species. • **CAMAROS** – human participants • **Interpretation** – effects of CCR5 antagonists for stroke recovery in gyrencephalic species represents a critical knowledge gap
(6) Tests during the awake phase of animals	No	No	• **Preclinical** – 0/5 studies reported assessing outcomes at night (i.e. awake phase of rodents) • **CAMAROS** – follow-up occurs during daytime hours (i.e. awake phase of humans) • **Interpretation** – future preclinical studies should conduct behavioural testing during the awake phase to better align with human testing conditions

3　**Table 4.** Alignment of included preclinical studies with additional STAIR / SRRR items and CAMAROS trial parameters.

Recommendation	Preclinical evidence	CAMAROS alignment	Notes
(1) Testing in both permanent and transient occlusion models	Yes	Yes	• **Preclinical** – 2/5 studies permanent occlusion, 2/5 transient occlusion, 1/5 hemorrhagic • **CAMAROS** – participants with ischemic anterior circulation stroke, with or without reperfusion treatments eligible for enrollment
(2) Monitoring of treatment effects on physiological parameters, including temperature, both during and after ischemia	No	NA	• **Preclinical** – 1/5 studies (Takami 2002; TAK-779) monitored and provided data on temperature for an extended period post-stroke • **CAMAROS** – participants will not be administered maraviroc in a cerebroprotective context
(3) Testing interaction with thrombolytics and other medications commonly administered in acute stroke	No	No	• **Preclinical** – 0/5 studies included assessed drug interactions • **CAMAROS** – participants may or may not be exposed to thrombolytics or other concomitant medications that could interact with recovery-inducing effects of maraviroc
(4) Animal model should produce infarction similar in relative size and location to that observed in humans	Yes	Yes	• **Preclinical** – 1/5 studies (Joy 2019; maraviroc) produced infarcts in motor regions of control animals that were less than 25% of hemispheric volume • **CAMAROS** - participants with ischemic anterior circulation stroke with hemiparesis requiring rehabilitation eligible for enrollment
(5) For studies claiming recovery effects, analysis of infarct volume should be performed to show equivalency of injury in the treated and stroke control groups	No	NA	• **Preclinical** – 0/1 experiments in late sub-acute/early chronic post-stroke phase (Joy 2019) demonstrated that each treatment group had equivalent baseline stroke volumes. This could potentially confound observed behavioral effects
(6) For studies claiming recovery effects, tissue repair / neuroplasticity processes should be quantified and directly related to behavioral recovery	Yes	NA	• **Preclinical** – 1/5 studies (Joy 2019) demonstrated maraviroc association with dendritic spine preservation, axonal projections to contralateral cortex, reduced inflammatory response, and upregulation of CREB / DLK signalling • **CAMAROS** – outcomes to assess mechanism of recovery are not included
(7) For studies claiming recovery effects, initial impairment with spontaneous recovery that plateaus significantly below pre-stroke performance	No	NA	• **Preclinical** – 1/5 studies assessed long-term outcomes (Joy 2019). Large variability in initial levels of impairment across experiments. Control animals demonstrate limited spontaneous recovery or atypical worsening of performance across time in multiple experiments. • **CAMAROS** – placebo participants will receive 8-week exercise program making assessment of spontaneous recovery difficult
(8) Behavioral effects should be assessed across a battery of domains, including both skilled and spontaneous upper limb and hindlimb use	No	No	• **Preclinical** – 2/5 studies included battery of behavioral tasks, but no study assessed effects on upper extremity skilled reaching / grasping or alteration of movement kinematics • **CAMAROS** – two primary outcomes: 1) upper extremity assessment using Fugl-Meyer; 2) change in 10-meter walk test score. Preclinical outcomes have limited relevance to Fugl-Meyer
(9) Mechanism of action should be assessed through gain of function / loss of function studies and directly associated to behavioral effects	Yes	NA	• **Preclinical** – 1/5 studies (Joy 2019) demonstrated that knockdown of DLK signalling using small hairpin DLK is able to block beneficial behavioral effects of CCR5 knockdown, providing a possible mechanism for stroke recovery-inducing effects • **CAMAROS** – outcomes to assess mechanism of recovery are not included
(10) Testing interactions of treatment with clinically inspired best practice, such as training or enrichment	No	No	• **Preclinical** – 0/5 studies included behavioral therapy / rehabilitative interventions • **CAMAROS** – all participants (including placebo) will participate in an 8-week

			exercise program

5   **Appendix 1.** Potential mechanisms of action.

6
7   **Appendix 1-figure 1.** Potential mechanistic pathways and proposed domains of biological
8   activity underlying CCR5 antagonist's cerebroprotective and neural repair effects post-stroke
9   described in the included studies. A list of mechanistic evidence supporting these pathways was
10   extracted from included studies in Appendix 1-table 1. Created with BioRender.com.

11 **Appendix 1-table 1.** List of outcomes used to determine potential mechanisms of action, by study.

Paper	Mechanistic evidence presented
Takami 2002 – *TAK-779*	- Infarct volume reduction (reduced by TAK-779) - MPO + Mac-1-alpha antibody used to stain infiltrating neutrophils/macrophages/microglia (Mac-1-alpha reduced by TAK-779 / MPO not decreased) – neutrophils the same, macrophages/microglia reduced
Li 2016 – *DAPTA*	- Cell death reduction (TUNEL / H&E) - Western blot – ROCK2 / P-MLC2 reduced (by DAPTA)
Joy 2019 – *maraviroc*	- CCR5 knockdown FACS isolated neuronal assay -> CREB / pCREB / DLK significantly increased. Erk(p44), GAP43, SAP/JNK1+2 increased, but not significantly. P38 decreased, but not significantly - 2-photon imaging - CCR5 knockdown + maraviroc -> increased dendritic spine count, fewer lost - BDA axonal quantification – CCR5 knockdown + maraviroc -> increased bihemispheric sprouting - GFAP / IBA-1 immunostaining – CCR5 knockdown + maraviroc -> reduced immunoreactivity. Decreased Ly6C (neutrophil) and Ly6Clow (macrophage) recruitment
Yan 2021 – *maraviroc*	- Cell death reduction (TUNEL / FJC / cleaved caspase 1 / Nissl) - Western blot – maraviroc -> increased PK A-Calpha / pCreb -> decreased NLRp1 / cleaved caspase 1 / IL-1B / apoptosis-associated speck protein / N-gasdermin D / IL-18 - Western blot -> effects reversed by 666-15 - Western blot -> rCCL5 shows reversed effects from maraviroc -> rCCL5 + cAMP reverses those effects
Chen 2022 – *maraviroc*	- Infarct volume / Cell death reduction (TTC / TUNEL) - Western blot – maraviroc -> increased Bcl2:BAX ratio -> increased IκBα, -> decreased P-IκBα / P-P65 / P-P38 / P-JNK - Western blot – P-JNK decreased, Anisomycin reverses the effect on P-JNK qRT-PCR, ELISA -> reduced IL-1B / IL-6 / TNF-alpha

14 **Appendix 2-table 1.** Readiness of CCR5 antagonists for translation based on the PRIMED[2] tool.

PRIMED[2] Domain	TAK-779	DAPTA	Maraviroc	CCR5 antagonism (all agents)
Sex of animals (0 – male OR female; 2 – both sexes)	0	0	0	**0**
Age of animals (0 – young only; 2 – older adults)	0	0	0	**0**
Species and strains (0 – one rodent species / strain; 1 - ≥2 rodent species / strains; 2 – rodents AND primates)	0	0	1	**1**
Reproducibility (0 – one species, one laboratory; 1 - ≥2 species, one lab OR ≥2 labs, one species; 2 - ≥2 species AND ≥2 labs)	0	0	2	**2**
Treatment time epoch (0 – no significant benefit; 1 – benefit in one epoch; 2 – benefit in ≥2 epochs)	1	1	2	**2**
Baseline comorbidities (0 – healthy animals only; 1 – one comorbid condition; 2 - ≥2 comorbid conditions)	0	0	0	**0**
Feasible time window (0 – treatment <45 minutes after ischemia onset; 2 - ≥45 minutes after ischemia onset)	0	0	2	**2**
Dose-response (0 – one dose; 2 – multiple doses with dose-response relationship)	0	0	0	**0**
Feasible route of delivery (0 – infeasible route (e.g., intraventricular injection); 1 – semi-infeasible route (e.g., intraarterial injection); 2 – feasible route (e.g., intravenous injection))	2	1	1	**2**
Behavioral and/or long-term outcome assessment (0 – no benefit; 1 – behavioral OR other outcome ≥30 days post-stroke; 2 – behavioral AND outcomes ≥30 days post-stroke)	0	0	2	**2**
Typical infarct volume reduction magnitude (0 – small effect (Cohen's d 0.2-0.39); 1 – medium effect (Cohen's d 0.4-0.69); 2 – large effect (Cohen's d ≥ 0.7)	2	2	2	**2**
Readiness for translation score (0-7 – low; 8-15 - medium; 16-22 – high)	Low (5)	Low (4)	Medium (12)	**Medium (13)**

16   **Appendix 3-table 1.** GRIPP2 Short Form Checklist.
17

Section and topic	Item
1: Aim	To conduct a preclinical systematic review assessing the effects of C-C chemokine receptor type 5 (CCR5) antagonists on motor and cognitive impairment following stroke. To collaborate with a panel of patients and caregivers with lived experience of stroke throughout the development and conduct of the preclinical systematic review.
2: Methods	A panel of eight patients and caregivers with lived experience of stroke was recruited to join the research team through the Heart & Stroke Foundation and the Patient and Family Advocacy Program at The Ottawa Hospital. Recruitment advertisements were distributed to both organizations. The patients and caregivers were involved in developing the research question and the protocol (i.e. elements of the AMICO question presented in the Methods section); identifying data items for extraction; conduct of the review including screening and extraction; interpreting the review results; and contributed to edits to the final manuscript. Patients and caregivers attended monthly virtual meetings. These meetings provided background knowledge of preclinical stroke, systematic review conduct, and discussed research findings as the review progressed. Patients provided insights into their lived experiences with stroke and helped identify priority areas that they were interested in. Patients and caregivers were offered financial compensation and co-authorship in recognition of their contributions to the research project.
3: Results	Patient engagement contributed to the study in several ways, including: - Informing the research question with the patient partner experience: patient partners have lived experience of stroke and provided patient priorities to analyze in our review - Refining our primary and secondary outcomes to include patient priorities - Developing of the protocol to include non-technical language - Editing screening and extraction forms - Participating as reviewers in abstract and full-text level screening - Participating as reviewers in extraction - Attending an international stroke conference and participating as co-authors of an abstract presented there - Interpretating analyzed data, including identifying patient important barriers and limitations to incorporate in the discussion
4: Discussion	Overall, patient engagement was successful in informing review development and conduct. Additionally, the researchers on the team learned from the patient panel. The patient panel brought a unique perspective to the planning and conduct of a preclinical systematic review. Additionally, members of the panel stated that the experience was useful for them as they gained new insights into preclinical research.
5: Reflections	Engagement was embedded within the research project from the inception to dissemination, where patient partners were members of the larger team. Patient partners directed us to outcomes for the systematic review that critical clinically to patients in the chronic phase of stroke recovery. Their participation in the conduct of the review facilitated meaningful collaborations and discussion.

20   **Appendix 4.** Full search strategies for MEDLINE / Embase and Web of Science.
21
22   Medline: 58
23   Embase: 118
24   Web of Science: 87
25
26   Total: 230
27   Total after duplicates: 166
28   Duplicates: 64
29
30   ***
31
32   Embase Classic+Embase <1947 to 2022 October 24>
33   Ovid MEDLINE(R) ALL <1946 to October 24, 2022>
34
1   exp animal experimentation/ or exp models, animal/ or animals/ or exp animal population
groups/ or chordata/ or vertebrates/ or exp amphibians/ or exp birds/ or exp fishes/ or exp
reptiles/ or mammals/ or primates/ or eutheria/ or exp artiodactyla/ or exp carnivore/ or exp
cephalopoda/ or exp cetacea/ or exp chiroptera/ or exp elephants/ or exp hyraxes/ or exp
insectivora/ or exp lagomorpha/ or marsupialia/ or exp monotremata/ or exp perissodactyla/ or
proboscidea mammal/ or exp rodentia/ or exp scandentia/ or exp sirenia/ or exp cingulata/ or
haplorhini/ or exp strepsirhini/ or exp platyrrhini/ or exp tarsii/ or catarrhini/ or exp
cercopithecidae/ or exp hylobatidae/ or hominidae/ or exp gorilla gorilla/ or exp pan paniscus/ or
exp pan troglodytes/ or exp pongo/   38941083
2   (rat or rats or animal or animals or mice or in vivo or mouse or rabbit or rabbits or murine
or pig or pigs or dog or dogs or bovine or fish or vertebrate or vertebrates or cat or cats or rodent
or rodents or mammal or mammals or chicken or chickens or monkey or monkeys or sheep or
canine or canines or porcine or cattle or bird or birds or hamster or hamsters or primate or
primates or cow or cows or chick or horse or horses or avian or avians or calf or swine or swines
or xenopus or turkeys or bear or bears or frog or frogs or zebrafish or goat or goats or equine or
calves or poultry or macaque or macaques or mole or moles or ovine or lamb or lambs or fishes
or diptera or amphibian or amphibians or snake or snakes or ruminant or ruminants or hen or
hens or piglet or piglets or feline or felines or simian or simians or laevis or trout or trouts or
teleost or teleosts or salmon or salmons or seal or seals or bull or bulls or ewe or ewes or
hedgehog or hedgehogs or macaca or macacas or proteus or pigeon or pigeons or bat or bats or
duck or ducks or chimpanzee or chimpanzees or baboon or baboons or deer or rana or ranas or
carp or carps or heifer or swallow or swallows or lizard or lizards or canis or sow or sows or
cynomolgus or quail or quails or reptile or reptiles or turtle or turtles or buffalo or gerbil or
gerbils or boar or boars or squirrel or squirrels or oncorhynchus or mus or toad or toads or fowl
or fowls or rerio or danio or ara or aras or musculus or tadpole or tadpoles or mulatta or salmo or
ram or eagle or eagles or ferret or ferrets or goldfish or catfish or whale or whales or fox or foxes
or ape or apes or elephant or elephants or bos or marmoset or marmosets or cod or cods or shark
or sharks or wolf or eel or eels or auratus or rattus or zebra or zebras or tilapia or tilapias or gilt
or camel or camels or squid or gallus or marsupial or marsupials or vole or voles or fascicularis
or ovis or salmonid or salmonids or tiger or tigers or dolphin or dolphins or robin or robins or
carpio or opossum or opossums or cyprinus or salamander or salamanders or felis or mink or

minks or swan or swans or norvegicus or bufo or torpedo or bass or lamprey or lampreys or sus
or python or pythons or tetrapod or tetrapods or shrew or shrews or lion or lions or hog or hogs
or songbird or songbirds or oreochromis or starling or starlings or caprine or carassius or owl or
owls or newt or newts or papio or scrofa or hare or hares or gorilla or gorillas or flounder or
flounders or goose or herring or herrings or therian or buffaloes or canary or sparrow or sparrows
or microtus or octopus or troglodytes or tuna or amphibia or chinchilla or chinchillas or ide or
oryzias or cervus or kangaroo or kangaroos or armadillo or armadillos or callithrix or pan
troglodytes or saimiri or cichlid or cichlids or donkey or donkeys or bream or char or chars or
finch or raccoon or raccoons or bothrops or anguilla or perch or cricetus or seabird or seabirds or
buck or bucks or naja or coturnix or salmonids or geese or minnow or minnows or raptor or
raptors or merione or meriones or rodentia or elaphus or amniote or amniotes or elasmobranch or
emu or emus or peromyscus or hominid or hominids or bubalus or crotalus or gull or gulls or
anas or anura or lemur or lemurs or crow or crows or camelus or gibbon or gibbons or waterfowl
or parrot or parrots or eels or cob or stickleback or sticklebacks or columba or mesocricetus or
ambystoma or raven or ravens or gadus or penguin or penguins or orangutan or orangutans or
sturgeon or sturgeons or cuniculus or aves or virginianus or cephalopod or cephalopods or cebus
or sparus or tortoise or tortoises or guttata or morhua or unguiculatus or dogfish or vulpes or
mallard or mallards or apodemus or alligator or alligators or oryctolagus or llama or llamas or
reindeer or mustela or duckling or ducklings or wolves or sander or amazona or zebu or badger
or badgers or dove or doves or ictalurus or capra or capras or equus or camelid or camelids or
poecilia or mule or mules or perciformes or salvelinus or labrax or cyprinidae or ariidae or
crocodile or crocodiles or fundulus or dicentrarchus or clarias or cercopithecus or chiroptera or
alpaca or alpacas or pike or pikes or paralichthys or puma or pumas or didelphis or pisces or
macropus or triturus or bison or bisons or epinephelus or gasterosteus or panthera or acipenser or
mackerel or mackerels or tamarin or tamarins or ostrich or anolis or vervet or vervets or wallaby
or glareolus or beaver or beavers or dromedary or catus or killifish or pimephales or promelas or
aotus or phoca or panda or pandas or porpoise or porpoises or myotis or yak or yaks or
agkistrodon or vipera or otter or otters or turbot or turbots or squamate or carnivora or mullet or
mullets or hawk or hawks or taeniopygia or seahorse or seahorses or poecilia reticulata or falcon
or falcons or prosimian or prosimians or parus or perca or fingerling or fingerlings or antelope or
antelopes or tupaia or passeriformes or sepia or saguinus or coyote or coyotes or pongo or
meleagris or reptilia or lepus or psittacine or hagfish or warbler or warblers or russell's viper or
russell's vipers or smolt or smolts or budgerigar or sardine or sardines or cavia or cavias or hyla
or pleurodeles or siluriformes or great tit or great tits or guppy or bonobo or bonobos or rutilus or
trichosurus or muridae or phodopus or channa or squalus or lynx or sturnus or petromyzon or
vitulina or monodelphis or cuttlefish or adder or adders or lepomis or canaria or gambusia or
guppies or xiphophorus or flatfish or koala or koalas or labeo or stingray or stingrays or chelonia
or lampetra or spermophilus or crocodilian or passer domesticus or sciurus or artiodactyla or
ranidae or corvus or necturus or platypus or canaries or bovid or lagopus or trimeresurus or
gariepinus or marten or martens or drosophilidae or mugil or sunfish or porcellus or
cypriniformes or alouatta or scophthalmus or anser or electrophorus or putorius or iguana or
iguanas or lama or lamas or takifugu or circus or eptesicus or flycatcher or galago or galagos or
trachemys or lungfish or characiformes or shorebird or shorebirds or giraffe or giraffes or
micropterus or scyliorhinus or cichlidae or loligo or porcupine or porcupines or chub or chubs or
solea or pleuronectes or hylidae or viperidae or echis or sorex or anchovy or lagomorph or
ostriches or vulture or vultures or whitefish or araneus or jird or jirds or tern or esox or drake or

drakes or elapidae or gallopavo or chordata or myodes or caretta or serinus or grouse or
misgurnus or meles or blackbird or blackbirds or coregonus or bobwhite or bobwhites or
heteropneustes or mammoth or mammoths or turdus or rhinella or ateles or characidae or clupea
or bungarus or brill or struthio camelus or sloth or sloths or pteropus or sculpin or anthropoids or
pollock or pollocks or morone or pan paniscus or litoria or chipmunk or chipmunks or
balaenoptera or marmota or melopsittacus or hyrax or lemming or lemmings or halibut or
hylobates or lates or caiman or caimans or sigmodon or stenella or barbel or barbels or sterna or
parakeet or parakeets or phocoena or leptodactylus or canidae or buteo or harengus or gopher or
gophers or marmot or marmots or gosling or goslings or platichthys or gar or gars or sebastes or
marsupialia or notophthalmus or gazelle or gazelles or insectivora or paridae or felidae or russula
or galliformes or bombina or colobus or echidna or echidnas or seabass or syncerus or plaice or
blue tit or blue tits or pagrus or catfishes or cetacea or barbus or cygnus or ficedula or chamois or
colubridae or perches or coelacanth or fitch or urodela or cynops or martes or halichoerus or aix
or salmonidae or leuciscus or magpie or magpies or silurus or whiting or whitings or
anseriformes or colinus or rhea or chlorocebus or octodon or acinonyx or mouflon or mouflons
or ibex or tetraodon or bufonidae or equidae or jackal or cephalopoda or dendroaspis or glama or
muskrat or muskrats or sable or sables or wildebeest or streptopelia or albifrons or
vespertilionidae or woodpecker or woodpeckers or muntjac or muntjacs or archosaur or branta or
cricetulus or megalobrama or poeciliidae or desmodus or snakehead or snakeheads or tench or
teal or teals or bandicoot or bandicoots or apteronotus or phyllostomidae or crocidura or buzzard
or buzzards or larimichthys or cercocebus or pipistrellus or erithacus or impala or impalas or
rousettus or haddock or haddocks or tinca or ratite or calidris or cynoglossus or
hypophthalmichthys or bullock or bullocks or dromedaries or alectoris or filly or salamandra or
cingulata or bitis or grus or ammodytes or macaw or macaws or hypoleuca or sapajus or
cyprinodontiformes or hippopotamus or pelophylax or capybara or capybaras or weasel or
weasels or cairina or cynomys or lutra or cockatoo or cockatoos or lachesis or lagomorpha or
rupicapra or daboia or orang utan or orang utans or platyrrhini or charadriiformes or micrurus or
psittaciformes or spalax or loris or mustelidae or sylvilagus or vitticeps or cockatiel or mustelus
or cottus or erythrocebus or dipodomys or platessa or callicebus or loricariidae or catostomus or
cuneata or cyanistes or cyprinodon or sigmodontinae or elasmobranchii or trichechus or
sauropsid or xenarthra or dormouse or perissodactyla or nautilus or cirrhinus or gulo or
tragelaphus or merula or numida or sciaenidae or cerastes or sciuridae or gibbosus or octopuses
or eland or elands or phyllomedusa or pogona or walrus or agamidae or leptodactylidae or
ridibundus or leontopithecus or anteater or anteaters or pelodiscus or cebidae or columbianus or
pelteobagrus fulvidraco or hominoidea or mandrillus or zonotrichia leucophrys or agama or
gobiocypris or bearded dragon or bearded dragons or sarotherodon or talpa or discoglossus or
hagfishes or sphenodon or gudgeon or amphiuma or aythya or tenrec or tenrec or hominidae or
risoria or salamandridae or camelidae or columbiformes or latimeria or plover or plovers or
afrotheria or falco sparverius or polecat or polecats or crotalinae or salvadora or tarsier or
lucioperca or anchovies or lungfishes or terrapin or dromaius novaehollandiae or lateolabrax or
eigenmannia or pelamis or theropithecus or murinae or gander or gymnotus or pseudacris or
gymnophiona or gymnotiformes or laticauda or falconiformes or dugong or dugongs or pintail or
pintails or rook or rooks or lasiurus or catshark or catsharks or micropogonias or red junglefowl
or paddlefish or ophiophagus or hollandicus or nymphicus or pimelodidae or aepyceros or
cobitidae or strigiformes or cobitis or dormice or alytes or calloselasma or guanaco or
phasianidae or round goby or trichogaster or catarrhini or eelpout or eelpouts or galaxias or gaur

or pungitius or suslik or susliks or flatfishes or percidae or caprinae or todarodes or osmerus or
ameiurus or anthropoidea or castor canadensis or pouting or poutings or tetraodontiformes or
arvicolinae or siamang or siamangs or castor fiber or nomascus or red knot or red knots or
syngnathidae or iguanidae or eretmochelys or ursidae or callimico or columbidae or
microhylidae or anaxyrus or menidia or pipistrelle or greylag or pipidae or scandentia or bowfin
or bowfins or dendrobatidae or zenaida or bushbaby or harrier or harriers or macropodidae or
pygerythrus or clupeidae or odorrana or corvidae or jerboa or jerboas or canutus or hylobatidae
or clupeiformes or great cormorant or great cormorants or scorpaeniformes or chondrostean or
garfish or proboscidea or psetta or diapsid or serotinus or tetrao or walruses or carcharhiniformes
or leucoraja or pumpkinseed or dosidicus or acipenseriformes or daubentonii or emberizidae or
gadiformes or hyraxes or stizostedion or wolverine or wolverines or lissotriton or acanthurus or
centrarchidae or gloydius or laurasiatheria or limosa or psittacula or leporidae or proteidae or
zander or zanders or arapaima or bagridae or cyprinodontidae or mithun or pandion or jackdaw
or jackdaws or procyonidae or carus or jaculus or salmoniformes or common sole or common
soles or protobothrops or calamita or brachyteles or trionyx or turdidae or boidae or luscinia or
pugnax or euarchontoglires or saithe or saithes or symphalangus or aardvark or aardvarks or
oystercatcher or oystercatchers or arius or corydoras or poacher or poachers or aurochs or
cebuella or crecca or lemuridae or sirenia or lemmus or perdix or glires or lepidosaur or muskox
or deinagkistrodon or pholidota or holocephali or cercopithecinae or clariidae or agapornis or
doryteuthis or tyrannidae or dicroglossidae or godwit or godwits or monedula or pongidae or
atheriniformes or colobinae or lophocebus or atelidae or cottidae or leucopsis or acanthuridae or
didelphimorphia or elver or elvers or lapponica or dermoptera or european hake or european
hakes or gerbillinae or banteng or hartebeest or hartebeests or hogget or haematopus or anguis
fragilis or grey heron or grey herons or blue whiting or blue whitings or furnariidae or
macrovipera or esocidae or lapwing or lapwings or mylopharyngodon or wallabia or
beloniformes or potoroo or potoroos or athene noctua or pleuronectidae or bushbabies or
muscicapidae or alligatoridae or fuligula or bush baby or guineafowl or spoonbill or spoonbills
or viverridae or catostomidae or zebrafishes or ibexes or vendace or estrildidae or monotremata
or sepiella or ambystomatidae or shelduck or shelducks or treeshrew or treeshrews or
hoplobatrachus or pochard or hoolock or hoolocks or lynxes or antilope or antilopes or
blackbuck or blackbucks or cricetinae or paramisgurnus or skylark or skylarks or soleidae or
allobates or northern wheatear or northern wheatears or pitheciidae or takin or theria or vanellus
or galaxiidae or lorisidae or ostralegus or palaeognathae or stone loach or alauda or callitrichinae
or caniformia or duttaphrynus or ictaluridae or osteoglossiformes or poultries or curema or ruddy
turnstone or ruddy turnstones or sheatfish or sunfishes or centropomidae or hemachatus or
platalea or thamnophilidae or song thrush or atherinopsidae or siluridae or tadorna or
chroicocephalus or ermine or ermines or gavialis or ruff or tupaiidae or diprotodontia or
hyaenidae or antilopinae or crocodylidae or herpestidae or hippopotamidae or northern shoveler
or round gobies or cheirogaleidae or indriidae or fundulidae or pythonidae or rhynchocephalia or
anodorhynchus or red-backed shrike or red-backed shrikes or triakidae or phalangeridae or
aoudad or boreoeutheria or eurasianjay or eurasian jays or feliformia or haplorhini or
osteoglossidae or paenungulata or struthioniformes or ferina or sanderling or sanderlings or
spheniscidae or cuttlefishes or cygnet or dasycneme or gadwall or gadwalls or pelobates fuscus
or wryneck or wrynecks or afrosoricida or culaea or dover sole or dover soles or paralichthyidae
or passeridae or osteolaemus or song thrushes or bluethroat or bluethroats or hydrophiidae or
megrim or mephitidae or strepsirhini or tomistoma or epidalea or osmeriformes or bush babies or

tarsiiform or atelinae or bufotes or eurasian coot or eurasian coots or galagidae or geopelia or philomachus or tubulidentata or bombinatoridae or pelobatidae or tachysurus or ailuridae or woodlark or woodlarks or alcelaphinae or redshank or redshanks or salientia or sand smelt or sand smelts or woodmice or woodmouse or dasyproctidae or eurasian wigeon or eurasian wigeons or garganey or garganeys or lemon sole or lemon soles or common dab or common dabs or graylag or graylags or leucorodia or osphronemidae or bewickii or common moorhen or common moorhens or decapodiformes or gobbler or gobblers or odontophoridae or paddlefishes or eutheria or salmonine or esociformes or eurasian woodcock or eurasian woodcocks or european smelt or european smelts or goldfishes or tenches or tyranni or common chaffinch or common chaffinchs or common redstart or common redstarts or common roach or common roachs or great knot or great knots or potoroidae or alytidae or coregonine or dipteral or leveret or poeciliopsis gracilis or amphiumidae or batrachoidiformes or bighead goby or heteropneustidae or lullula or norway pout or norway pouts or sipunculida or dogfishes or sebastidae or tarsiidae or alethinophidia or common nase or common nases or common sandpiper or common sandpipers or eurasian blackcap or eurasian blackcaps or pterocnemia or syngnathiformes or common chaffinches or eupleridae or octopodiformes or phascolarctidae or scophthalmidae or starry smooth-hound or starry smooth-hounds or whitefishes or cuniculidae or european sprat or european sprats or rosy bitterling or rosy bitterlings or common dace or common daces or lesser weever or lesser weevers or scaldfish or water rail or water rails or alouattinae or centrarchiformes or common whitethroat or common whitethroats or gavialidae or grey gurnard or grey gurnards or lateolabracidae or rheiformes or tubgurnard or tub gurnards or common chiffchaff or common chiffchaffs or garfishes or lesser whitethroat or lesser whitethroats or myoxidae or seabasses or spariformes or umbridae or yellow boxfish or anabantiformes or aotidae or common bleak or common bleaks or common rudd or common rudds or greater pipefish or hapale or nandiniidae or stone loaches or whinchat or whinchats or acanthuriformes or brotula barbata or common ling or common lings or common roaches or cottonrat or cottonrats or douroucoulis or dromaiidae or fitches or fitchew or galaxiiformes or laprine or saimiriinae or solenette or tarsii or tompot blenny or common dragonet or common dragonets or longspined bullhead or longspined bullheads or monotremate or monotremates or pempheriformes or perdicinae or presbytini or smegmamorpha or bighead gobies or carangaria incertae sedis or coiidae or fivebeard rockling or foulmart or foumart or grasskeet or greater pipefishes or ibices or millionfish or muguliformes or norwegian topknot or peewit or red sea sailfin tang or rupicapras or sheatfishes or tompot blennies or twait shad or yellow boxfishes).tw. 14176107

medline.st. 29904626

2 not 3 8398417

1 or 4 40352035

"Receptors, CCR5"/ 18466

(cc chemokine receptor* 5 or cc-ckr5 or ccr5 or cd195 antigen* or ckr5 receptor*).tw,kf. 22382

exp CCR5 Receptor Antagonists/ 8211

C-C chemokine receptor type 5.tw,kf. 434

(md6p741w8a or maraviroc or selzentry or Celsentri or uk 427,857 or uk-427,857 or uk427,857).tw,kf. 3408

Vicriviroc.mp. 728

(Aplaviroc or ancriviroc or aplaviroc or leronlimab or mavorixafor).mp. 866

13   INCB009471.mp.   2
14   TBR 652.mp. 37
15   Pro 140.mp.   297
16   HGS004.mp. 15
17   6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16   33160
18   5 and 17   23661
19   exp cerebrovascular disorders/ or stroke/   1287736
20   stroke*.tw,kf. 792358
21   transient isch?em* attack*.tw,kf.   42566
22   brain injur*.tw,kf.   199884
23   ((brain or cereb* or intracereb*) adj2 (isch?em* or infarct* or h?emorrhag*)).tw,kf.
261843
24   or/19-23   1714778
25   18 and 24   373
26   25 use medall 58
27   exp animal experiment/ or exp animal model/ or exp experimental animal/ or exp
transgenic animal/ or exp male animal/ or exp female animal/ or exp juvenile animal/ or animal/
or chordata/ or vertebrate/ or tetrapod/ or exp fish/ or amniote/ or exp amphibia/ or mammal/ or
exp reptile/ or exp sauropsid/ or therian/ or exp monotreme/ or placental mammal/ or exp
marsupial/ or Euarchontoglires/ or exp Afrotheria/ or exp Boreoeutheria/ or exp Laurasiatheria/
or exp Xenarthra/ or primate/ or exp Dermoptera/ or exp Glires/ or exp Scandentia/ or
Haplorhini/ or exp prosimian/ or simian/ or exp tarsiiform/ or Catarrhini/ or exp Platyrrhini/ or
ape/ or exp Cercopithecidae/ or hominid/ or exp hylobatidae/ or exp chimpanzee/ or exp gorilla/
or exp orang utan/ or exp cephalopod/   14616185
28   (rat or rats or animal or animals or mice or "in vivo" or mouse or rabbit or rabbits or
murine or pig or pigs or dog or dogs or bovine or fish or vertebrate or vertebrates or cat or cats or
rodent or rodents or mammal or mammals or chicken or chickens or monkey or monkeys or
sheep or canine or canines or porcine or cattle or bird or birds or hamster or hamsters or primate
or primates or cow or cows or chick or horse or horses or avian or avians or calf or swine or
swines or xenopus or turkeys or bear or bears or frog or frogs or zebrafish or goat or goats or
equine or calves or poultry or macaque or macaques or mole or moles or ovine or lamb or lambs
or fishes or diptera or amphibian or amphibians or snake or snakes or ruminant or ruminants or
hen or hens or piglet or piglets or feline or felines or simian or simians or laevis or trout or trouts
or teleost or teleosts or salmon or salmons or seal or seals or bull or bulls or ewe or ewes or
hedgehog or hedgehogs or macaca or macacas or proteus or pigeon or pigeons or bat or bats or
duck or ducks or chimpanzee or chimpanzees or baboon or baboons or deer or deers or rana or
ranas or carp or carps or heifer or swallow or swallows or lizard or lizards or canis or sow or
sows or cynomolgus or quail or quails or reptile or reptiles or turtle or turtles or buffalo or gerbil
or gerbils or boar or boars or squirrel or squirrels or oncorhynchus or mus or toad or toads or
fowl or fowls or rerio or danio or ara or aras or musculus or tadpole or tadpoles or mulatta or
salmo or ram or eagle or eagles or ferret or ferrets or goldfish or catfish or whale or whales or
fox or foxes or ape or apes or elephant or elephants or bos or marmoset or marmosets or cod or
cods or shark or sharks or wolf or eel or eels or auratus or rattus or zebra or zebras or tilapia or
tilapias or gilt or camel or camels or squid or gallus or marsupial or marsupials or vole or voles
or fascicularis or ovis or salmonid or salmonids or tiger or tigers or dolphin or dolphins or robin
or robins or carpio or opossum or opossums or cyprinus or salamander or salamanders or felis or

mink or minks or swan or swans or norvegicus or bufo or torpedo or bass or lamprey or lampreys
or sus or python or pythons or tetrapod or tetrapods or shrew or shrews or lion or lions or hog or
hogs or songbird or songbirds or oreochromis or starling or starlings or caprine or carassius or
owl or owls or newt or newts or papio or scrofa or hare or hares or gorilla or gorillas or flounder
or flounders or goose or herring or herrings or therian or buffaloes or canary or sparrow or
sparrows or microtus or octopus or troglodytes or tuna or amphibia or chinchilla or chinchillas or
ide or oryzias or cervus or kangaroo or kangaroos or armadillo or armadillos or callithrix or "pan
troglodytes" or saimiri or cichlid or cichlids or donkey or donkeys or bream or char or chars or
finch or raccoon or raccoons or bothrops or anguilla or perch or cricetus or seabird or seabirds or
buck or bucks or naja or coturnix or salmonids or geese or minnow or minnows or raptor or
raptors or merione or meriones or rodentia or elaphus or amniote or amniotes or elasmobranch or
emu or emus or peromyscus or hominid or hominids or bubalus or crotalus or gull or gulls or
anas or anura or lemur or lemurs or crow or crows or camelus or gibbon or gibbons or waterfowl
or parrot or parrots or eels or cob or stickleback or sticklebacks or columba or mesocricetus or
ambystoma or raven or ravens or gadus or penguin or penguins or orangutan or orangutans or
sturgeon or sturgeons or cuniculus or aves or virginianus or cephalopod or cephalopods or cebus
or sparus or tortoise or tortoises or guttata or morhua or unguiculatus or dogfish or vulpes or
mallard or mallards or apodemus or alligator or alligators or oryctolagus or llama or llamas or
reindeer or mustela or duckling or ducklings or wolves or sander or amazona or zebu or badger
or badgers or dove or doves or ictalurus or capra or capras or equus or camelid or camelids or
poecilia or mule or mules or perciformes or salvelinus or labrax or cyprinidae or ariidae or
crocodile or crocodiles or fundulus or dicentrarchus or clarias or cercopithecus or chiroptera or
alpaca or alpacas or pike or pikes or paralichthys or puma or pumas or didelphis or pisces or
macropus or triturus or bison or bisons or epinephelus or gasterosteus or panthera or acipenser or
mackerel or mackerels or tamarin or tamarins or ostrich or anolis or vervet or vervets or wallaby
or glareolus or beaver or beavers or dromedary or catus or killifish or pimephales or promelas or
aotus or phoca or panda or pandas or porpoise or porpoises or myotis or yak or yaks or
agkistrodon or vipera or otter or otters or turbot or turbots or squamate or carnivora or mullet or
mullets or hawk or hawks or taeniopygia or seahorse or seahorses or "poecilia reticulata" or
falcon or falcons or prosimian or prosimians or parus or perca or fingerling or fingerlings or
antelope or antelopes or tupaia or passeriformes or sepia or saguinus or coyote or coyotes or
pongo or meleagris or reptilia or lepus or psittacine or hagfish or warbler or warblers or "russell s
viper" or "russell s vipers" or smolt or smolts or budgerigar or sardine or sardines or cavia or
cavias or hyla or pleurodeles or siluriformes or "great tit" or "great tits" or guppy or bonobo or
bonobos or rutilus or trichosurus or muridae or phodopus or channa or squalus or lynx or sturnus
or petromyzon or vitulina or monodelphis or cuttlefish or adder or adders or lepomis or canaria
or gambusia or guppies).tw.   14111285
29   (xiphophorus or flatfish or koala or koalas or labeo or stingray or stingrays or chelonia or
lampetra or spermophilus or crocodilian or "passer domesticus" or sciurus or artiodactyla or
ranidae or corvus or necturus or platypus or canaries or bovid or lagopus or trimeresurus or
gariepinus or marten or martens or drosophilidae or mugil or sunfish or porcellus or
cypriniformes or alouatta or scophthalmus or anser or electrophorus or putorius or iguana or
iguanas or lama or lamas or takifugu or circus or eptesicus or flycatcher or galago or galagos or
trachemys or lungfish or characiformes or shorebird or shorebirds or giraffe or giraffes or
micropterus or scyliorhinus or cichlidae or loligo or porcupine or porcupines or chub or chubs or
solea or pleuronectes or hylidae or viperidae or echis or sorex or anchovy or lagomorph or

342 ostriches or vulture or vultures or whitefish or araneus or jird or jirds or tern or esox or drake or
343 drakes or elapidae or gallopavo or chordata or myodes or caretta or serinus or grouse or
344 misgurnus or meles or blackbird or blackbirds or coregonus or bobwhite or bobwhites or
345 heteropneustes or mammoth or mammoths or turdus or rhinella or ateles or characidae or clupea
346 or bungarus or brill or "struthio camelus" or sloth or sloths or pteropus or sculpin or anthropoids
347 or pollock or pollocks or morone or "pan paniscus" or litoria or chipmunk or chipmunks or
348 balaenoptera or marmota or melopsittacus or hyrax or lemming or lemmings or halibut or
349 hylobates or lates or caiman or caimans or sigmodon or stenella or barbel or barbels or sterna or
350 parakeet or parakeets or phocoena or leptodactylus or canidae or buteo or harengus or gopher or
351 gophers or marmot or marmots or gosling or goslings or platichthys or gar or gars or sebastes or
352 marsupialia or notophthalmus or gazelle or gazelles or insectivora or paridae or felidae or russula
353 or galliformes or bombina or colobus or echidna or echidnas or seabass or syncerus or plaice or
354 "blue tit" or "blue tits" or pagrus or catfishes or cetacea or barbus or cygnus or ficedula or
355 chamois or colubridae or perches or coelacanth or fitch or urodela or cynops or martes or
356 halichoerus or aix or salmonidae or leuciscus or magpie or magpies or silurus or whiting or
357 whitings or anseriformes or colinus or rhea or chlorocebus or octodon or acinonyx or mouflon or
358 mouflons or ibex or tetraodon or bufonidae or equidae or jackal or cephalopoda or dendroaspis
359 or glama or muskrat or muskrats or sable or sables or wildebeest or streptopelia or albifrons or
360 vespertilionidae or woodpecker or woodpeckers or muntjac or muntjacs or archosaur or branta or
361 cricetulus or megalobrama or poeciliidae or desmodus or snakehead or snakeheads or tench or
362 teal or teals or bandicoot or bandicoots or apteronotus or phyllostomidae or crocidura or buzzard
363 or buzzards or larimichthys or cercocebus or pipistrellus or erithacus or impala or impalas or
364 rousettus or haddock or haddocks or tinca or ratite or calidris or cynoglossus or
365 hypophthalmichthys or bullock or bullocks or dromedaries or alectoris or filly or salamandra or
366 cingulata or bitis or grus or ammodytes or macaw or macaws or hypoleuca or sapajus or
367 cyprinodontiformes or hippopotamus or pelophylax or capybara or capybaras or weasel or
368 weasels or cairina or cynomys or lutra or cockatoo or cockatoos or lachesis or lagomorpha or
369 rupicapra or daboia or "orang utan" or "orang utans" or platyrrhini or charadriiformes or
370 micrurus or psittaciformes or spalax or loris or mustelidae or sylvilagus or vitticeps or cockatiel
371 or mustelus or cottus or erythrocebus or dipodomys or platessa or callicebus or loricariidae or
372 catostomus or cuneata or cyanistes or cyprinodon or sigmodontinae or elasmobranchii or
373 trichechus or sauropsid or xenarthra or dormouse or perissodactyla or nautilus or cirrhinus or
374 gulo or gulos or tragelaphus or merula or numida or sciaenidae or cerastes or sciuridae or
375 gibbosus or octopuses or eland or elands or phyllomedusa or pogona or walrus or agamidae or
376 leptodactylidae or ridibundus or leontopithecus or anteater or anteaters or pelodiscus or cebidae
377 or columbianus or "pelteobagrus fulvidraco" or hominoidea or mandrillus or "zonotrichia
378 leucophrys" or agama or gobiocypris or "bearded dragon" or "bearded dragons" or sarotherodon
379 or talpa or discoglossus or hagfishes or sphenodon or gudgeon or amphiuma or aythya or tenrec
380 or tenrec or hominidae or risoria or salamandridae or camelidae or columbiformes or latimeria or
381 plover or plovers or afrotheria or "falco sparverius" or polecat or polecats or crotalinae or
382 salvadora or tarsier or lucioperca or anchovies or lungfishes or terrapin or "dromaius
383 novaehollandiae" or lateolabrax or eigenmannia or pelamis or theropithecus or murinae or
384 gander or gymnotus or pseudacris or gymnophiona or gymnotiformes or laticauda or
385 falconiformes or dugong or dugongs or pintail or pintails or rook or rooks or lasiurus or catshark
386 or catsharks or micropogonias or "red junglefowl" or paddlefish or ophiophagus or hollandicus
387 or nymphicus or pimelodidae or aepyceros or cobitidae or strigiformes or cobitis or dormice or

alytes or calloselasma or guanaco or guanacos or phasianidae or "round goby" or trichogaster or
catarrhini or eelpout or eelpouts or galaxias or gaur or pungitius or suslik or susliks or flatfishes
or percidae or caprinae or todarodes or osmerus or ameiurus or anthropoidea or "castor
canadensis" or pouting or poutings or tetraodontiformes or arvicolinae or siamang or siamangs or
"castor fiber" or nomascus or "red knot" or "red knots" or syngnathidae or iguanidae or
eretmochelys or ursidae or callimico or columbidae or microhylidae or anaxyrus or menidia or
pipistrelle or greylag or pipidae or scandentia or bowfin or bowfins or dendrobatidae or zenaida
or bushbaby or harrier or harriers or macropodidae or pygerythrus or clupeidae or odorrana or
corvidae or jerboa or jerboas or canutus or hylobatidae or clupeiformes or "great cormorant" or
"great cormorants" or scorpaeniformes or chondrostean or garfish or proboscidea or psetta or
diapsid or serotinus or tetrao or walruses or carcharhiniformes or leucoraja or pumpkinseed or
dosidicus or acipenseriformes or daubentonii or emberizidae or gadiformes or hyraxes or
stizostedion or wolverine or wolverines or lissotriton or acanthurus or centrarchidae or gloydius
or laurasiatheria or limosa or psittacula or leporidae or proteidae or zander or zanders or
arapaima or bagridae or cyprinodontidae or mithun or pandion or jackdaw or jackdaws or
procyonidae or carus or jaculus or salmoniformes or "common sole" or "common soles" or
protobothrops or calamita or brachyteles or trionyx or turdidae or boidae or luscinia or pugnax or
euarchontoglires or saithe or saithes or symphalangus or aardvark).tw.   272058
30   (aardvarks or oystercatcher or oystercatchers or arius or corydoras or poacher or poachers
or aurochs or cebuella or crecca or lemuridae or sirenia or lemmus or perdix or glires or
lepidosaur or muskox or deinagkistrodon or pholidota or holocephali or cercopithecinae or
clariidae or agapornis or doryteuthis or tyrannidae or dicroglossidae or godwit or godwits or
monedula or pongidae or atheriniformes or colobinae or lophocebus or atelidae or cottidae or
leucopsis or acanthuridae or didelphimorphia or elver or elvers or lapponica or dermoptera or
"european hake" or "european hakes" or gerbillinae or banteng or hartebeest or hartebeests or
hogget or haematopus or "anguis fragilis" or "grey heron" or "grey herons" or "blue whiting" or
"blue whitings" or furnariidae or macrovipera or esocidae or lapwing or lapwings or
mylopharyngodon or wallabia or beloniformes or potoroo or potoroos or "athene noctua" or
pleuronectidae or bushbabies or muscicapidae or alligatoridae or fuligula or "bush baby" or
guineafowl or spoonbill or spoonbills or viverridae or catostomidae or zebrafishes or ibexes or
vendace or estrildidae or monotremata or sepiella or ambystomatidae or shelduck or shelducks or
treeshrew or treeshrews or hoplobatrachus or pochard or hoolock or hoolocks or lynxes or
antilope or antilopes or blackbuck or blackbucks or cricetinae or paramisgurnus or skylark or
skylarks or soleidae or allobates or "northern wheatear" or "northern wheatears" or pitheciidae or
takin or theria or vanellus or galaxiidae or lorisidae or ostralegus or palaeognathae or "stone
loach" or alauda or callitrichinae or caniformia or duttaphrynus or ictaluridae or
osteoglossiformes or poultries or curema or "ruddy turnstone" or "ruddy turnstones" or sheatfish
or sunfishes or centropomidae or hemachatus or platalea or thamnophilidae or "song thrush" or
atherinopsidae or siluridae or tadorna or chroicocephalus or ermine or ermines or gavialis or ruff
or tupaiidae or diprotodontia or hyaenidae or antilopinae or crocodylidae or herpestidae or
hippopotamidae or "northern shoveler" or "round gobies" or cheirogaleidae or indriidae or
fundulidae or pythonidae or rhynchocephalia or anodorhynchus or "red-backed shrike" or "red-
backed shrikes" or triakidae or phalangeridae or aoudad or boreoeutheria or "eurasian jay" or
"eurasian jays" or feliformia or haplorhini or osteoglossidae or paenungulata or struthioniformes
or ferina or sanderling or sanderlings or spheniscidae or cuttlefishes or cygnet or dasycneme or
gadwall or gadwalls or "pelobates fuscus" or wryneck or wrynecks or afrosoricida or culaea or

13 eLife-VOR-RA-2024-103245R1

"dover sole" or "dover soles" or paralichthyidae or passeridae or osteolaemus or "song thrushes"
or bluethroat or bluethroats or hydrophiidae or megrim or mephitidae or strepsirhini or
tomistoma or epidalea or osmeriformes or "bush babies" or tarsiiform or atelinae or bufotes or
"eurasian coot" or "eurasian coots" or galagidae or geopelia or philomachus or tubulidentata or
bombinatoridae or pelobatidae or tachysurus or ailuridae or woodlark or woodlarks or
alcelaphinae or redshank or redshanks or salientia or "sand smelt" or "sand smelts" or woodmice
or woodmouse or dasyproctidae or "eurasian wigeon" or "eurasian wigeons" or garganey or
garganeys or "lemon sole" or "lemon soles" or "common dab" or "common dabs" or graylag or
graylags or leucorodia or osphronemidae or bewickii or "common moorhen" or "common
moorhens" or decapodiformes or gobbler or gobblers or odontophoridae or paddlefishes or
eutheria or salmonine or esociformes or "eurasian woodcock" or "eurasian woodcocks" or
"european smelt" or "european smelts" or goldfishes or tenches or tyranni or "common
chaffinch" or "common chaffinchs" or "common redstart" or "common redstarts" or "common
roach" or "common roachs" or "great knot" or "great knots" or potoroidae or alytidae or
coregonine or dipteral or leveret or "poeciliopsis gracilis" or amphiumidae or batrachoidiformes
or "bighead goby" or heteropneustidae or lullula or "norway pout" or "norway pouts" or
sipunculida or dogfishes or sebastidae or tarsiidae or alethinophidia or "common nase" or
"common nases" or "common sandpiper" or "common sandpipers" or "eurasian blackcap" or
"eurasian blackcaps" or pterocnemia or syngnathiformes or "common chaffinches" or eupleridae
or octopodiformes or phascolarctidae or scophthalmidae or "starry smooth-hound" or "starry
smooth-hounds" or whitefishes or cuniculidae or "european sprat" or "european sprats" or "rosy
bitterling" or "rosy bitterlings" or "common dace" or "common daces" or "lesser weever" or
"lesser weevers" or scaldfish or "water rail" or "water rails" or alouattinae or centrarchiformes or
"common whitethroat" or "common whitethroats" or gavialidae or "grey gurnard" or "grey
gurnards" or lateolabracidae or rheiformes or "tub gurnard" or "tub gurnards" or "common
chiffchaff" or "common chiffchaffs" or garfishes or "lesser whitethroat" or "lesser whitethroats"
or myoxidae or seabasses or spariformes or umbridae or "yellow boxfish" or anabantiformes or
aotidae or "common bleak" or "common bleaks" or "common rudd" or "common rudds" or
"greater pipefish" or hapale or nandiniidae or "stone loaches" or whinchat or whinchats or
acanthuriformes or "brotula barbata" or "common ling" or "common lings" or "common roaches"
or cottonrat or cottonrats or douroucoulis or dromaiidae or fitches or fitchew or galaxiiformes or
laprine or saimiriinae or solenette or tarsii or "tompot blenny" or "common dragonet" or
"common dragonets" or "longspined bullhead" or "longspined bullheads" or monotremate or
monotremates or pempheriformes or perdicinae or presbytini or smegmamorpha or "bighead
gobies" or "carangaria incertae sedis" or coiidae or "fivebeard rockling" or foulmart or foumart
or grasskeet or "greater pipefishes" or ibices or millionfish or muguliformes or "norwegian
topknot" or peewit or "red sea sailfin tang" or rupicapras or sheatfishes or "tompot blennies" or
"twait shad" or "yellow boxfishes").tw.   21723
31   or/27-30   18002118
32   exp chemokine receptor CCR5 antagonist/   6682
33   (cc chemokine receptor* 5 or cc-ckr5 or ccr5 or cd195 antigen* or ckr5 receptor* or C-C
chemokine receptor type 5).tw.   22155
34   (md6p741w8a or maraviroc or selzentry or Celsentri or uk 427,857 or uk-427,857 or
uk427,857).tw.   3329
35   Vicriviroc.mp. 728
36   (Aplaviroc or ancriviroc or aplaviroc or leronlimab or mavorixafor).mp.   866

37   INCB009471.mp.   2
38   TBR 652.mp. 37
39   Pro 140.mp.   297
40   HGS004.mp.   15
41   chemokine receptor CCR5/   12698
42   or/32-41   31956
43   31 and 42   10275
44   exp cerebrovascular disease/ or exp cerebrovascular accident/   1207681
45   stroke*.tw.   771477
46   ((brain or cereb* or intracereb*) adj2 (isch?em* or infarct* or h?emorrhag*)).tw. 247502
47   (transient isch?em* attack* or brain injur*).tw.   231020
48   44 or 45 or 46 or 47   1675247
49   43 and 48   172
50   49 use emczd 118
51   26 or 50   176
495

****

Web of Science – October 25, 2022

Web of Science Search Strategy (v0.1)

Entitlements:

- WOS.SSCI: 1900 to 2022

- WOS.AHCI: 1975 to 2022

- WOS.ISTP: 1990 to 2022

- WOS.ESCI: 2005 to 2022

- WOS.SCI: 1900 to 2022

- WOS.ISSHP: 1990 to 2022

Searches:

1: CCR5 (Topic)   Results: 11678

2: TS=(cc chemokine receptor* 5) OR TS=(C-C chemokine receptor type 5)
    Results: 1796

3: TS=(cc-ckr5)   Results: 9

4: TS=(Vicriviroc)   Results: 158

5: TS=(((md6p741w8a OR maraviroc OR selzentry OR Celsentri)))
    Results: 1542

6: ALL=((Aplaviroc or ancriviroc or aplaviroc or leronlimab or mavorixafor))
        Results: 97

7: (((TS=(INCB009471)) OR TS=(TBR 652)) OR TS=(Pro 140)) OR TS=(HGS004)
            Results: 972

8: #7 OR #6 OR #5 OR #4 OR #3 OR #2 OR #1   Results: 14241

9: (TS=(stroke*)) OR TS=("transient isch?em* attack*")   Results: 430668

10: TS=((brain or cereb* or intracereb*) NEAR/2 ischem*)   Results: 77919

11: TS=(((brain or cereb* or intracereb*) NEAR/2 ischaem*))
        Results: 5389

12: TS=(((brain or cereb* or intracereb*) NEAR/2 hemorrhag*))
        Results: 33959

13: TS=(((brain or cereb* or intracereb*) NEAR/2 haemorrhag*))
Results: 4867

14: TS="cerebrovascular disorder*"   Results: 3993

15: TS="brain injur*"   Results: 121601

16: #15 OR #14 OR #13 OR #12 OR #11 OR #10 OR #9   Results:
591495

17: #8 AND #16   Results: 136

18: TS=(rat OR rats OR animal OR animals OR mice OR "in vivo" OR mouse OR rabbit OR
rabbits OR murine OR pig OR pigs OR dog OR dogs OR bovine OR fish OR vertebrate
OR vertebrates OR cat OR cats OR rodent OR rodents OR mammal OR mammals OR
chicken OR chickens OR monkey OR monkeys OR sheep OR canine OR canines OR porcine
OR cattle OR bird OR birds OR hamster OR hamsters OR primate OR primates OR cow
OR cows OR chick OR horse OR horses OR avian OR avians OR calf OR swine OR
swines OR xenopus OR turkeys OR bear OR bears OR frog OR frogs OR zebrafish OR
goat OR goats OR equine OR calves OR poultry OR macaque OR macaques OR mole OR
moles OR ovine OR lamb OR lambs OR fishes OR diptera OR amphibian OR amphibians
OR snake OR snakes OR ruminant OR ruminants OR hen OR hens OR piglet OR
piglets OR feline OR felines OR simian OR simians OR laevis OR trout OR trouts OR
teleost OR teleosts OR salmon OR salmons OR seal OR seals OR bull OR bulls OR
ewe OR ewes OR hedgehog OR hedgehogs OR macaca OR macacas OR proteus OR
pigeon ORpigeons OR bat OR bats OR duck OR ducks OR chimpanzee OR chimpanzees OR
baboon OR baboons OR deer OR rana OR ranas OR carp OR carps OR heifer OR swallow OR
swallows OR lizard OR lizards OR canis OR sow OR sows OR cynomolgus OR quail OR quails
OR reptile OR reptiles OR turtle OR turtles OR buffalo OR gerbil OR gerbils OR boar OR boars
OR squirrel OR squirrels OR oncorhynchus OR mus OR toad OR toads OR fowl OR fowls OR
rerio OR danio OR ara OR aras OR musculus OR tadpole OR tadpoles OR mulatta OR
salmo OR ram OR eagle OR eagles OR ferret OR ferrets OR goldfish OR catfish OR
whale OR whales OR fox OR foxes OR ape OR apes OR elephant OR elephants OR bos OR
marmoset OR marmosets OR cod OR cods OR shark OR sharks OR wolf OR eel OR eels OR
auratus OR rattus OR zebra OR zebras OR tilapia OR tilapias OR gilt OR camel OR camels
OR squid OR gallus OR marsupial OR marsupials OR vole OR voles OR fascicularis OR
ovis OR salmonid OR salmonids OR tiger OR tigers OR dolphin OR dolphins OR robin OR

robins OR carpio OR opossumOR opossums OR cyprinus OR salamander OR salamanders
OR felis OR mink OR minks OR swan OR swans OR norvegicus OR bufo OR torpedo OR
bass OR lamprey OR lampreys OR sus OR python OR pythons OR tetrapod OR tetrapods OR
shrew OR shrews OR lion OR lions OR hogOR hogs OR songbird OR songbirds OR
oreochromis OR starling OR starlings OR caprine OR carassius OR owl OR owls OR newt OR
newts OR papio OR scrofa OR hare OR hares OR gorilla OR gorillas OR flounder OR flounders
OR goose OR herring OR herrings OR therianOR buffaloes OR canary OR sparrow OR
sparrows OR microtus OR octopus OR troglodytes OR tuna OR amphibia OR chinchilla
OR chinchillas OR ide OR oryzias OR cervus OR kangaroo OR kangaroos OR armadillo
OR armadillos OR callithrix OR "pan troglodytes" OR saimiri OR cichlid OR cichlids OR
donkey OR donkeys OR bream OR char OR chars OR finch OR raccoon OR raccoons OR
bothrops OR anguilla OR perch OR cricetus OR seabird OR seabirds OR buck OR bucks OR
naja OR coturnix OR salmonids OR geese OR minnow OR minnows ORraptor OR raptors OR
merione OR meriones OR rodentia OR elaphus OR amniote OR amniotes OR
elasmobranch OR emu OR emus OR peromyscus OR hominid OR hominids OR bubalus OR
crotalus OR gull OR gulls OR anas OR anura OR lemur OR lemurs OR crow OR crows OR
camelus OR gibbon OR gibbons OR waterfowl OR parrot OR parrots OR eels OR cob
OR stickleback OR sticklebacks OR columba OR mesocricetus OR ambystoma OR raven
OR ravens OR gadus OR penguin OR penguins OR orangutan OR orangutans OR sturgeon
OR sturgeons OR cuniculus OR aves OR virginianus OR cephalopod OR cephalopods OR cebus
OR sparus OR tortoise OR tortoises OR guttata OR morhua OR unguiculatus OR dogfish OR
vulpes OR mallard OR mallards OR apodemus OR alligator OR alligators OR oryctolagus
OR llama OR llamas OR reindeer OR mustela OR duckling OR ducklings OR wolves OR
sander OR amazona OR zebu OR badger OR badgers OR dove OR doves OR ictalurus OR
capra OR capras OR equus OR camelid OR camelids OR poecilia OR mule OR mules
OR perciformes OR salvelinus OR labrax OR cyprinidae OR ariidae OR crocodile OR
crocodiles OR fundulus OR dicentrarchus OR clarias OR cercopithecus OR chiroptera OR
alpaca OR alpacas OR pike OR pikes OR paralichthys OR puma OR pumas OR didelphis OR
pisces OR macropus OR triturus OR bison OR bisons OR epinephelus OR gasterosteus OR
panthera OR acipenser OR mackerel OR mackerels OR tamarin OR tamarins OR ostrich OR
anolis OR vervet OR vervets OR wallaby OR glareolus OR beaver OR beavers OR
dromedary OR catus OR killifish OR pimephales OR promelas OR aotus OR phoca OR
panda OR pandas OR porpoise OR porpoises OR myotis OR yak OR yaks OR agkistrodon
OR vipera OR otter OR otters OR turbot OR turbots OR squamate OR carnivora OR mullet OR
mullets OR hawk OR hawks OR taeniopygia OR seahorse OR seahorses OR "poecilia reticulata"
OR falcon OR falcons OR prosimian OR prosimians OR parus OR perca OR fingerling
OR fingerlings OR antelope OR antelopes OR tupaia OR passeriformes OR sepia OR
saguinus OR coyote OR coyotes OR pongo OR meleagris OR reptilia OR lepus OR psittacine
OR hagfish OR warbler OR warblers OR "russell s viper" OR "russell s vipers" OR smolt OR
smolts OR budgerigar OR sardine OR sardines OR cavia OR cavias OR hyla OR pleurodeles OR
siluriformes OR "great tit" OR "great tits" OR guppy OR bonobo OR bonobos OR rutilus OR
trichosurus OR muridae OR phodopus OR channa OR squalus OR lynx OR sturnus OR
petromyzon OR vitulina OR monodelphis OR cuttlefish OR adder OR adders OR lepomis OR
canaria OR gambusia OR guppies OR xiphophorus OR flatfish OR koala ORkoalas OR

labeo OR stingray OR stingrays OR chelonia OR lampetra OR spermophilus OR
crocodilian OR "passer domesticus" OR sciurus OR artiodactyla OR ranidae OR corvus
OR necturus OR platypus OR canaries OR bovid OR lagopus OR trimeresurus OR
gariepinus ORmarten OR martens OR drosophilidae OR mugil OR sunfish OR porcellus OR
cypriniformes OR alouatta OR scophthalmus OR anser OR electrophorus OR putorius OR
iguana OR iguanas OR lama OR lamas OR takifugu OR circus OR eptesicus OR flycatcher OR
galago OR galagos OR trachemys OR lungfish OR characiformes OR shorebird OR
shorebirds OR giraffe OR giraffes OR micropterus OR scyliorhinus OR cichlidae OR
loligo OR porcupine OR porcupines OR chub OR chubs OR solea OR pleuronectes OR hylidae
OR viperidae OR echis OR sorex OR anchovy OR lagomorph OR ostriches OR vulture OR
vultures OR whitefish OR araneus OR jird OR jirds OR tern OR esox OR drake OR drakes OR
elapidae OR gallopavo OR chordata OR myodes OR caretta OR serinus OR grouse OR
misgurnus OR meles OR blackbird OR blackbirds OR coregonus OR bobwhite OR
bobwhites OR heteropneustes OR

mammoth OR mammoths OR turdus OR rhinella OR ateles OR characidae OR clupea
OR bungarus OR brill OR "struthio camelus" OR sloth OR sloths OR pteropus OR sculpin
OR anthropoids OR pollock OR pollocks OR morone OR "pan paniscus" OR litoria OR
chipmunk OR chipmunks OR balaenoptera OR marmota OR melopsittacus OR hyrax OR
lemming OR lemmings OR halibut OR hylobates OR lates OR caiman OR caimans OR
sigmodon OR stenella OR barbel OR barbels ORsterna OR parakeet OR parakeets OR
phocoena OR leptodactylus OR canidae OR buteo OR harengus OR gopher OR gophers
OR marmot OR marmots OR gosling OR goslings OR platichthys OR gar OR gars OR sebastes
OR marsupialia OR notophthalmus OR gazelle OR gazelles OR insectivora OR paridae OR
felidae OR russula OR galliformes OR bombina OR colobus OR echidna OR echidnas OR
seabass OR syncerus OR plaice OR "blue tit" OR "blue tits" OR pagrus OR catfishes OR cetacea
OR barbus OR cygnus OR ficedula OR chamois OR colubridae OR perches OR coelacanth OR
fitch OR urodela OR cynops OR martes OR halichoerus OR aix OR salmonidae OR leuciscus
OR magpie OR magpies OR silurus OR whiting OR whitings OR anseriformes OR colinus OR
rhea OR chlorocebus OR octodon OR acinonyx OR mouflon OR mouflons OR ibex OR
tetraodon OR bufonidae OR equidae OR jackal OR cephalopoda OR dendroaspis OR
glama OR muskrat OR muskrats OR sable OR sables OR wildebeest OR streptopelia OR
albifrons OR vespertilionidae OR woodpecker OR woodpeckers OR muntjac OR muntjacs OR
archosaur OR branta OR cricetulus OR megalobrama OR poeciliidae OR desmodus OR
snakehead OR snakeheads OR tench OR teal OR teals OR bandicoot OR bandicoots OR
apteronotus OR phyllostomidae OR crocidura OR buzzard OR buzzards OR larimichthys OR
cercocebus OR pipistrellus OR erithacus OR impala OR impalas OR rousettus OR haddock OR
haddocks OR tinca OR ratite OR calidris OR cynoglossus OR hypophthalmichthys OR bullock
OR bullocks OR dromedaries OR alectoris OR filly OR salamandra OR cingulata OR bitis
OR grus OR ammodytes OR macaw OR macaws OR hypoleuca OR sapajus OR
cyprinodontiformes OR hippopotamus OR pelophylax OR capybara OR capybaras OR
weasel OR weasels OR cairina OR cynomys OR lutra OR cockatoo OR cockatoos OR
lachesis OR lagomorpha OR rupicapra OR daboia OR "orang utan" OR "orang utans" OR
platyrrhini OR charadriiformes OR micrurus OR psittaciformes OR spalax OR loris OR
mustelidae OR sylvilagus OR vitticeps OR cockatiel OR mustelus OR cottus OR

erythrocebus OR dipodomys OR platessa OR callicebus OR loricariidae OR catostomus
OR cuneata OR cyanistes OR cyprinodon OR sigmodontinae OR elasmobranchii OR
trichechus OR sauropsid OR xenarthra OR dormouse OR perissodactyla OR nautilus OR
cirrhinus OR gulo OR gulos OR tragelaphus OR merula OR numidaOR sciaenidae OR
cerastes OR sciuridae OR gibbosus OR octopuses OR eland OR elands OR
phyllomedusa OR pogona OR walrus OR agamidae OR leptodactylidae OR ridibundus
OR leontopithecus OR anteater OR anteaters OR pelodiscus OR cebidae OR columbianus
OR "pelteobagrus fulvidraco" OR hominoidea OR mandrillus OR "zonotrichia leucophrys"
OR agama OR gobiocypris OR "bearded dragon" OR "bearded dragons" OR sarotherodon OR
talpa OR discoglossus OR hagfishes OR sphenodon OR gudgeon OR amphiuma OR aythya OR
tenrec OR tenrec OR hominidae OR risoria OR salamandridae OR camelidae OR
columbiformes OR latimeria OR plover OR plovers OR afrotheria OR "falco sparverius"
OR polecat OR polecats OR crotalinae OR salvadora OR tarsier OR lucioperca OR
anchovies OR lungfishes OR terrapin OR "dromaius novaehollandiae" OR lateolabrax OR
eigenmannia OR pelamis OR theropithecus OR murinae OR gander OR gymnotus OR
pseudacris OR gymnophiona OR gymnotiformes OR laticauda OR falconiformes OR
dugong OR dugongs OR pintail OR pintails OR rook ORrooks OR lasiurus OR catshark
OR catsharks OR micropogonias OR "red junglefowl" OR paddlefish OR eutheria OR
ophiophagus OR hollandicus OR nymphicus OR pimelodidae OR aepyceros OR cobitidae
OR strigiformes OR cobitis OR dormice OR alytes OR calloselasma OR guanaco OR
guanacos OR phasianidae OR "round goby" OR trichogaster OR catarrhini OR eelpout OR
eelpouts OR galaxias OR gaur OR pungitius OR suslik OR susliks OR flatfishes OR percidae
OR caprinae OR todarodes OR osmerus OR ameiurus OR anthropoidea OR "castor canadensis"
OR pouting OR poutings OR tetraodontiformes OR arvicolinae OR siamang OR siamangs OR
"castor fiber" OR nomascus OR "red knot" OR "red knots" OR syngnathidae OR iguanidae
OR eretmochelys OR ursidae OR callimico OR columbidae OR microhylidaeOR anaxyrus
OR menidia OR pipistrelle OR greylag OR pipidae OR scandentia OR bowfin OR
bowfins OR dendrobatidae OR zenaida OR bushbaby OR harrier OR harriers OR
macropodidae OR pygerythrus OR clupeidae OR odorrana OR corvidae OR jerboa OR
jerboas OR canutus OR hylobatidae OR clupeiformes OR "great cormorant" OR "great
cormorants" OR scorpaeniformes OR chondrostean OR garfish OR proboscidea OR psetta OR
diapsid OR serotinus OR tetrao OR walruses OR carcharhiniformes OR leucoraja OR
pumpkinseed OR dosidicus OR acipenseriformes OR daubentonii OR emberizidae OR
gadiformes OR hyraxes OR stizostedion OR wolverine OR wolverines OR lissotriton OR
acanthurus OR centrarchidae OR gloydius OR laurasiatheria OR limosa OR psittacula OR
leporidae OR proteidae OR zander ORzanders OR arapaima OR bagridae OR cyprinodontidae
OR mithun OR pandion OR jackdaw OR jackdaws OR procyonidae OR carus OR jaculus OR
salmoniformes OR "common sole" OR "common soles" OR protobothrops OR calamita OR
brachyteles OR trionyx OR turdidae OR boidae OR luscinia OR pugnax OR euarchontoglires
OR saithe OR saithes OR symphalangus OR aardvark OR aardvarks OR oystercatcher OR
oystercatchers OR arius OR corydoras OR poacher OR poachers OR aurochs OR cebuella OR
crecca OR lemuridae OR sirenia OR lemmus OR perdix OR glires OR lepidosaur OR muskox
OR deinagkistrodon OR pholidota OR holocephali OR cercopithecinae OR clariidae OR
agapornis OR doryteuthis OR tyrannidae OR dicroglossidae OR godwit OR godwits OR

monedula OR pongidae OR atheriniformes OR colobinae OR lophocebus OR atelidae OR cottidae OR leucopsis OR acanthuridae OR didelphimorphia OR elver OR elvers OR lapponica OR dermoptera OR "european hake" OR "european hakes" OR gerbillinae OR banteng OR hartebeest OR hartebeests OR hogget OR haematopus OR "anguis fragilis" OR "grey heron" OR "grey herons" OR "blue whiting" OR "blue whitings" OR furnariidae OR macrovipera OR esocidae OR lapwing OR lapwings OR mylopharyngodon OR wallabia OR beloniformes OR potoroo OR potoroos OR "athene noctua" OR pleuronectidae OR bushbabies OR muscicapidae OR alligatoridae OR fuligula OR "bush baby" OR guineafowl OR spoonbill OR spoonbills OR viverridae OR catostomidae OR zebrafishes OR ibexes OR vendace OR estrildidae OR monotremata OR sepiella OR ambystomatidae OR shelduck OR shelducks OR treeshrew OR treeshrews OR hoplobatrachus OR pochard OR hoolock OR hoolocks OR lynxes OR antilope OR antilopes OR blackbuck OR blackbucks OR cricetinae OR paramisgurnus OR skylark OR skylarks OR soleidae OR allobates OR "northern wheatear" OR "northern wheatears" OR pitheciidae OR takin OR theria OR vanellus OR galaxiidae OR lorisidae OR ostralegus OR palaeognathae OR "stone loach" OR alauda OR callitrichinae OR caniformia OR duttaphrynus OR ictaluridae OR osteoglossiformes OR poultries OR curema OR "ruddy turnstone" OR

"ruddy turnstones" OR sheatfish OR sunfishes OR centropomidae OR hemachatus OR platalea OR thamnophilidae OR "song thrush" OR atherinopsidae OR siluridae OR tadorna OR chroicocephalus OR ermine OR ermines OR gavialis OR ruffe OR tupaiidae OR diprotodontia OR hyaenidae OR antilopinae OR crocodylidae OR herpestidae OR hippopotamidae OR "northern shoveler" OR "round gobies" OR cheirogaleidae OR indriidae OR fundulidae OR pythonidae OR rhynchocephalia OR anodorhynchus OR "red-backed shrike" OR "red-backed shrikes" OR triakidae OR phalangeridae OR aoudad OR boreoeutheria OR "eurasian jay" OR "eurasian jays" OR feliformia OR haplorhini OR osteoglossidae OR paenungulata OR struthioniformes OR ferina OR sanderling OR sanderlings OR spheniscidae OR cuttlefishes OR cygnet OR dasycneme OR gadwall OR gadwalls OR "pelobates fuscus" OR wryneck OR wrynecks OR afrosoricida OR culaea OR "dover sole" OR "dover soles" OR paralichthyidae OR passeridae OR osteolaemus OR "song thrushes" OR bluethroat OR bluethroats OR hydrophiidae OR megrim OR mephitidae OR strepsirhini OR tomistoma OR epidalea OR osmeriformes OR "bush babies" OR tarsiiform OR atelinae OR bufotes OR "eurasian coot" OR "eurasian coots" OR galagidae OR geopelia OR philomachus OR tubulidentata OR bombinatoridae OR pelobatidae OR tachysurus OR ailuridae OR woodlark OR woodlarks OR alcelaphinae OR redshank OR redshanks OR salientia OR "sand smelt" OR "sand smelts" OR woodmice OR woodmouse OR dasyproctidae OR "eurasian wigeon" OR "eurasian wigeons" OR garganey OR garganeys OR "lemon sole" OR "lemon soles" OR "common dab" OR "common dabs" OR graylag OR graylags OR leucorodia OR osphronemidae OR bewickii OR "common moorhen" OR "common moorhens" OR decapodiformes OR gobbler OR gobblers OR odontophoridae OR paddlefishes OR salmonine OR esociformes OR "eurasian woodcock" OR "eurasian woodcocks" OR "european smelt" OR "european smelts" OR goldfishes OR tenches OR tyranni OR "common chaffinch" OR "common chaffinchs" OR "common redstart" OR "common redstarts" OR "common roach" OR "common roachs" OR "great knot" OR "great knots" OR potoroidae OR alytidae OR coregonine OR dipteral OR leveret OR "poeciliopsis

gracilis" OR amphiumidae OR batrachoidiformes OR "bighead goby" OR heteropneustidae OR lullula OR "norway pout" OR "norway pouts" OR sipunculida OR dogfishes OR sebastidae OR tarsiidae OR alethinophidia OR "common nase" OR "common nases" OR "common sandpiper" OR "common sandpipers" OR "eurasian blackcap" OR "eurasian blackcaps" OR pterocnemia OR syngnathiformes OR "common chaffinches" OR eupleridae OR octopodiformes OR phascolarctidae OR scophthalmidae OR "starry smooth-hound" OR "starry smooth-hounds" OR whitefishes OR cuniculidae OR "european sprat" OR "european sprats" OR "rosy bitterling" OR "rosy bitterlings" OR "common dace" OR "common daces" OR "lesser weever" OR "lesser weevers" OR scaldfish OR "water rail" OR "water rails" OR alouattinae OR centrarchiformes OR "common whitethroat" OR "common whitethroats" OR gavialidae OR "grey gurnard" OR "greygurnards" OR lateolabracidae OR rheiformes OR "tub gurnard" OR "tub gurnards" OR "common chiffchaff" OR "common chiffchaffs" OR garfishes OR "lesser whitethroat" OR "lesser whitethroats" OR myoxidae OR seabasses OR spariformes OR umbridae OR "yellow boxfish" OR anabantiformes OR aotidae OR "common bleak" OR "common bleaks" OR "common rudd" OR "common rudds" OR "greater pipefish" OR hapale OR nandiniidae OR "stone loaches" OR whinchat OR whinchats OR acanthuriformes OR "brotula barbata" OR "common ling" OR "common lings" OR "common roaches" OR cottonrat OR cottonrats OR douroucoulis OR dromaiidae OR fitches OR fitchew OR galaxiiformes OR laprine OR saimiriinae OR solenette OR tarsii OR "tompot blenny" OR "common dragonet" OR "common dragonets" OR "longspinedbullhead" OR "longspined bullheads" OR monotreme OR monotremates OR pempheriformes OR perdicinae OR presbytini OR smegmamorpha OR "bighead gobies" OR "carangaria incertae sedis" OR coiidae OR "fivebeard rockling" OR foulmart OR foumart OR grasskeet OR "greater pipefishes" OR ibices OR millionfish OR muguliformes OR "norwegian topknot" OR peewit OR "red sea sailfin tang" OR rupicapras OR sheatfishes OR "tompot blennies" OR "twait shad" OR "yellow boxfishes")
                    Results: 10177702

19: #17 AND #18   Results: 87

****

Appendix 5. Data Extraction Forms

<u>Study Characteristics</u>

1. What is the year of publication?
2. What journal is the article published in?
3. Name of the first author? (Full last name, initial of first name - e.g. Smith, K)
4. Name and email of the corresponding author? (e.g. Smith K, smithk89@uottawa.ca)
5. What is the source of funding?
a. Not Reported
b. Government
c. Industry
d. Academic Institution
e. Charity/Foundation
f. Other
g. Unclear
6. What country is the corresponding author from (look at corresponding address)?

<u>Animal Model</u>

1. What is the species?
a. Mouse
b. Rat
c. Monkey
d. Rabbit
e. Other
2. What type of stroke is being studied?
a. Ischemic
b. Hemorrhagic
c. Other
3. What type of stroke model is being used?
a. Intraluminal suture (e.g. Filament, distal, and transient middle cerebral artery
814   occlusion)
815   i. What is the number of minutes of occlusion time (relevant to lesion
816   severity)?
817   b. Permanent middle cerebral artery occlusion (i.e. cauterization, permanent clip,
818   permanent distal middle cerebral artery ligation)
c. Photothrombotic
d. Embolism
e. Modified 3 vessel occlusion
f. Endothelin-1
g. Microvascular embolic
h. L-NIO
i. Intracerebral hemorrhagic
j. Other (please specify)
4. What are the sexes of the animals?

a. Male
b. Female
c. Male and female
d. Not reported
5. What is the weight in grams and, if applicable, specify when the weight(s) was
measured?
6. What is the age in weeks of the species?
7. Are there comorbidities in the animals have (e.g. aged, obesity, diabetes/hyperglycemia,
hypertension)?
a. Yes (please specify)
b. No
8. What specific area of the brain was the stroke induced in?
a. Frontal lobe
b. Parietal lobe
c. Temporal lobe
d. Occipital lobe
e. Cerebellum
f. Thalamus
g. Basal ganglia
h. Other (please describe)
i. Not reported
9. Were there any animals excluded?
a. Yes (please explain why i.e. death, no impairment from stroke, etc.)
b. No
10. What was the total number of animals excluded from the study?

<u>Study Intervention</u>

1. What agent is used?
a. Maraviroc
b. Selzentry
c. Celsentri
d. Leronlimab
e. Aplaviroc
f. Vicriviroc
g. Ancriviroc
h. Other (please specify)
2. What is the dose (numeric value)?
3. What is the unit used for the dose?
4. What is the route of administration?
5. What was the agent prepared/diluted in?
6. When was the treatment administered?
a. Pre-stroke
b. Post-stroke

7. What is the timing in hours of intervention post-stroke (first adminsteration if
applicable)?
8. Was the treatment administered at multiple timepoints?
873   a. Yes (please specify the other timepoints in hours)
874   b. No
875   9. How is time zero defined in the study?
876   10. Was the intervention paired with any other therapy (e.g. post-stroke rehabilitation
877   paradigms)?
878   a. Yes (please describe)
879   b. No

880   <u>Outcomes</u>

881   1. Are motor behaviour outcomes (e.g. motor skills) being assessed? Please check all that
882   apply
883   a. Yes
884   i. Neurological deficit score
885   ii. Rotarod
886   iii. Adhesive removal test
887   iv. Cylinder task
888   v. Foot fault & paw placement tests (e.g. tapered beam, ladder, grid walking)
889   vi. Montoya staircase
890   vii. Reaching task (e.g. tray, pellet, pasta matrix)
891   viii. Corner test
892   ix. Staircase test
893   x. Forelimb placing test
894   xi. Wire hanging test
895   xii. Other
896   b. No
897   2. Are cognitive behavioural outcomes (e.g. cognitive skills) being assessed? Please check
898   all that apply
899   a. Yes
900   i. Morris water maze
901   ii. Y-maze test
902   iii. Novel object recognition test
903   iv. Elevated plus maze
904   v. Sucrose preference test
905   vi. Tail suspension test
906   vii. Open field test
907   viii. Forced swim test
908   ix. Other
909   b. No
910   3. Was infarct volume/size measured?
911   a. Yes

i. What is the mean infarct size in mm3 or % of the hemisphere for the treatment animals?

ii. What is the mean infarct size in mm3 or % of the hemisphere for the control animals?

iii. What is the method for measuring infarct size? Please check all that apply
1. Triphenyltetrazolium chloride (TTC)
2. Cresyl violet (CV)
3. Hematoxylin and eosin (H&E)
4. Magnetic resonance imaging (MRI)
5. Other (please specify)

iv. What is the (first, if applicable) time of the post-stroke measure of infarct size in days?

v. Was infarct size measured multiple times?
1. Yes (please list the other times in days)
2. No

b. No

4. How many animals are in the treatment and control groups?

5. What number of animals died for control and treatment groups?

6. What was the adverse effect (e.g. muscle fatigue, reduced mobility) for control and treatment groups

7. Were tissue outcomes (e.g. immunohistochemical staining, axonal neurofilament staining, pharmacogenetic techniques, axonal tracing techniques, genetic markers identification) measured or visualized?
a. Yes, please explain
b. No

8. Were brain imaging or biomarkers used to measure animal neural connectivity?
a. Yes
i. Diffusion tensor imaging (DTI)
ii. Resting state functional MRI (rsFMRI)
iii. Task-based functional MRI
iv. Other (please specify)
b. No

Risk of Bias

"Yes" indicates low risk of bias; "no" indicates high risk of bias; and "unclear" indicates an unclear risk of bias. If one of the relevant signaling questions is answered with "no," this indicates high risk of bias for that specific entry.

1. Was the allocation sequence adequately generated and applied?
*Did the investigators describe a random component in the sequence generation process such as:
- Referring to a random number table;
- Using a computer random number generator.
a. Yes
b. No

954   c. Unclear
955
Additional info:
Examples of a non-random approach:
• Allocation by judgement or by investigator's preference;
• Allocation based on the results of a laboratory test or a series of tests;
• Allocation by availability of the intervention;
• Sequence generated by odd or even date of birth;
• Sequence generated by some rule based on animal number or cage number.
963
964   2. Were the groups similar at baseline or were they adjusted for confounders in the analysis?
965   *Was the distribution of relevant baseline characteristics balanced for the intervention
966   and control groups?
967   a. Yes
968   b. No
969   c. Unclear
970
971   *If relevant, did the investigators adequately adjust for unequal distribution of some
972   relevant baseline characteristics in the analysis?
973   a. Yes
974   b. No
975   c. Unclear
976
977   *Was the timing of disease induction adequate?
978   a. Yes
979   b. No
980   c. Unclear
981
Additional info:
The number and type of baseline characteristics are dependent on the review question. Before
starting their risk of bias assessment, therefore, reviewers need to discuss which baseline
characteristics need to be comparable between the groups. In an SR investigating the effects of
hypothermia on infarct size, for example, gender distribution, left ventricular weight and heart
rate and blood pressure should be similar between the groups at the start of the study.
A description of baseline characteristics and/or confounders usually contains:
• The sex, age and weight of the animals
• Baseline values of the outcomes which are of interest in the study
991
992   Timing of disease induction:
993   In some prevention studies, the disease is induced after allocation of the intervention. For
994   example, in an experiment on preventive probiotic supplementation in acute pancreatitis,
995   pancreatitis is induced after allocation of the animals to the probiotic or control group. To reduce
996   baseline imbalance, the timing of disease induction should be equal for both treatment groups.

Examples of adequate timing of disease induction:

997   •   The disease was induced before randomization of the intervention.

998   •   The disease was induced after randomization of the intervention, but the timing of disease

999   induction was at random, and the individual inducing the disease was adequately blinded

1000   from knowing which intervention each animal received.

1001

1002   3. Was the allocation to the different groups adequately concealed during?

1003   *Could the investigator allocating the animals to intervention or control group not foresee

1004   assignment due to one of the following or equivalent methods?

1005   •   ☐Third-party coding of experimental and control group allocation Central

1006   randomization by a third party; Sequentially numbered opaque, sealed envelopes

1007   a. Yes

1008   b. No

1009   c. Unclear

1010

Additional info:

Examples of investigators allocating the animals being possibly able to foresee

assignments:

•   Open randomization schedule

•   Envelopes without appropriate safeguard

•   Alternation or rotation

•   Allocation based on date of birth

•   Allocation based on animal number

•   Any other explicitly unconcealed procedure of a non-random approach

1020

4. Were the animals randomly housed during the experiment?

*Did the authors randomly place the cages or animals within the animal room/facility?

a. Yes

b. No

c. Unclear

•   Animals were selected at random during outcome assessment (use signaling

questions of entry 6).

1028

1029   *Is it unlikely that the outcome or the outcome measurement was influenced by not

1030   randomly housing the animals?

1031   a. Yes

1032   b. No

1033   c. Unclear

1034

1035   The animals from the various experimental groups live together in one cage/pasture (e.g.,

1036   housing conditions are identical).

1037   Additional info:

1038   Examples of investigators using a non-random approach when placing the cages:

- Experimental groups were studied on various locations (e.g., group A in lab A or on shelf A; Group B in Lab B or on shelf B).

5. Were the caregivers and/or investigators blinded from knowledge which intervention each animal received during the experiment?
   *Was blinding of caregivers and investigators ensured, and was it unlikely that their blinding could have been broken?
   a. Yes
   b. No
   c. Unclear

   - ID cards of individual animals, or cage/animal labels are coded and identical in appearance.
   - Sequentially numbered drug containers are identical in appearance.
   - The circumstances during the intervention are specified and similar in both groups (#).
   - Housing conditions of the animals during the experiment are randomized within the room (use criteria of entry 4).

   Additional info:
   Examples of inappropriate blinding:
   - Colored cage labels (red for group A, yellow group B)
   - Expected differences in visible effects between control and experimental groups
   - Housing conditions of the animals are not randomized within the room during the experiment; use criteria of entry 4
   - The individual who prepares the experiment is the same as the one who conducts and analyses the experiment
   - Circumstances during the intervention are not similar in both groups (#)
   - Examples where circumstances during the intervention were not similar:
   - Timing of administration of the placebo and exp drug was different.
   - Instruments used to conduct experiment differ between experimental and control group (e.g., experiment about effects abdominal pressure; exp group receives operation and needle to increase pressure, while control group only has the operation).

   **The relevance of the above-mentioned items depends on the experiment. Authors of the review need to judge for themselves which of the above-mentioned items could cause bias in the results when not similar. These should be assessed.

6. Were animals selected at random for outcome assessment?
   *Did the investigators randomly pick an animal during outcome assessment, or did they use a random component in the sequence generation for outcome assessment?
   a. Yes
   b. No
   c. Unclear

- Referring to a random number table;
- Using a computer random number generator;
- Etc.

7. Was the outcome assessor blinded?
   *Was blinding of the outcome assessor ensured, and was it unlikely that blinding could have been broken?
     a. Yes
     b. No
     c. Unclear

   - Outcome assessment methods were the same in both groups.
   - Animals were selected at random during outcome assessment (use signaling questions of entry 6).

   *Was the outcome assessor not blinded, but do review authors judge that the outcome is not likely to be influenced by lack of blinding?
     a. Yes
     b. No
     c. Unclear

   (e.g., mortality)
   Additional info:
   This item needs to be assessed for each main outcome.

8. Were incomplete outcome data adequately addressed? (*)
   *Were all animals included in the analysis?
     a. Yes
     b. No
     c. Unclear

   *Were the reasons for missing outcome data unlikely to be related to true outcome? (e.g., technical failure)
     a. Yes
     b. No
     c. Unclear

   *Are missing outcome data balanced in numbers across intervention groups, with similar reasons for missing data across groups?
     a. Yes
     b. No
     c. Unclear

1126   *Are missing outcome data imputed using appropriate methods?
1127   a. Yes
1128   b. No
1129   c. Unclear
1130
1131

9. Are reports of the study free of selective outcome reporting? (*)
*Was the study protocol available and were all of the study's pre-specified primary and
secondary outcomes reported in the current manuscript?
a. Yes
b. No
c. Unclear
1138

1139   *Was the study protocol not available, but was it clear that the published report included
1140   all expected outcomes (i.e. comparing methods and results section)?
1141   a. Yes
1142   b. No
1143   c. Unclear
1144

Additional info:
Selective outcome reporting:
☐- Not all of the study's pre-specified primary outcomes have been reported;
☐- One or more primary outcomes have been reported using measurements, analysis
methods or data subsets (e.g., subscales) that were not pre-specified in the protocol;
☐- One or more reported primary outcomes were not pre-specified (unless clear
justification for their reporting has been provided, such as an unexpected adverse effect);
☐- The study report fails to include results for a key outcome that would be expected
to have been reported for such a study.
1154

1155 10. Was the study apparently free of other problems that could result in high risk of bias? (*)
1156   *Was the study free of contamination (pooling drugs)?
1157   a. Yes
1158   b. No
1159   c. Unclear
1160

1161   *Was the study free of inappropriate influence of funders?
1162   a. Yes
1163   b. No
1164   c. Unclear
1165

1166   *Was the study free of unit of analysis errors?
1167   a. Yes
1168   b. No
1169   c. Unclear

1170
1171   *Were design-specific risks of bias absent?
1172   a. Yes
1173   b. No
1174   c. Unclear
1175
1176   *Were new animals added to the control and experimental groups to replace drop-outs
1177   from the original population?
1178   a. Yes
1179   b. No
1180   c. Unclear
1181
Additional info:
The relevance of the signaling questions (Table 3) depends on the experiment. Review
authors need to judge for themselves which of the items could cause bias in their results
and should be assessed.
Contamination/pooling drugs:
Experiments in which animals receive □ besides the intervention drug □ additional
treatment or drugs which might influence or bias the result.
Unit of analysis errors:
• Interventions to parts of the body within one participant (i. e., one eye exp; one
eye control).
• All animals receiving the same intervention are caged together, but analysis was
conducted as if every single animal was one experimental unit.
• Design-specific risks of bias:
• Crossover design that was not suitable (intervention with no temporary effect, or
the disease is not stable over time)
• Crossover design with risk of carry-over effect
• Crossover design with only first period data being available
• Crossover design with many animals not receiving 2nd or following treatment due
to large number of drop-outs probably due to longer duration of study
• Crossover design in which all animals received same order of interventions
• Multi-arm study in which the same comparisons of groups are not reported for all
outcomes (selective outcome reporting)
• Multi-arm study in which results of different arms are combined (all data should
be presented per group)
• Cluster randomized trial not taking clustering into account during statistical
analysis (unit of analysis error)
• Crossover design in which paired analysis of the results is not taken into account

Validity

1. Does the treatment effect vary with dose?
a. Yes

            b. No
            c. Not applicable (only tested one dose)
            d. Unsure
    2. Does the treatment remain effective when administered at clinically relevant delayed
        times?
            a. Yes (tested after 6 hours and the CCR5 antagonist was still effective)
            b. No (tested after 6 hours, but the CCR5 antagonist was not effective)
            c. Not applicable (not tested after 6 hours)
            d. Unsure
    3. Does the treatment cause expected physiological effects?
            a. Yes
            b. No
            c. Unsure
    4. Does the treatment penetrate the blood brain barrier?
            a. Yes
            b. No
            c. Unsure
    5. What the tests done across multiple laboratories?
            a. Yes
            b. No
            c. Unsure
    6. Was testing done on gyrencephalic species (non-human primates i.e. monkeys, dogs,
        pigs, etc.)
            a. Yes
            b. No
            c. Unsure
    7. Did the preclinical testing of therapy occur during the awake phase for the animal model
        (during the dark phase for rodents i.e. was the test done during the nighttime)?
            a. Yes
            b. No
            c. Unsure

    ****

Author	Year	Dose (mg/kg)	Time of Drug Administration	Sacrifice (days)	T (n)	C (n)	MD [95% CI]
Post-stroke administration of CCR5 antagonist							
Takami et al.	2002	0.25	50 min post-stroke	2	17	18	-10.2 [-19.6, -0.8]
Chen et al.	2022	20	1.5 h post-stroke	3	5	5	-26.3 [-54.0, 1.4]
RE Model for Post-Stroke (Q = 1.16, df = 1, p = 0.28; I^2 = 13.8%)							-12.8 [-24.2, -1.3]
Pre-stroke administration of CCR5 antagonist							
Li et al.	2016	0.01	15 min pre-stroke	1	5	5	-8.7 [-11.4, -6.0]
Takami et al.	2002	5	10 min pre-stroke	2	10	6	-9.4 [-22.5, 3.7]
Takami et al.	2002	50	10 min pre-stroke	2	13	6	-13.3 [-26.0, -0.5]
RE Model for Pre-Stroke (Q = 0.48, df = 2, p = 0.79; I^2 = 0.0%)							-8.9 [-11.5, -6.3]

Favours Treatment   Favours Control

RE Model for All Studies (Q = 2.03, df = 4, p = 0.73; I^2 = 0.0%)   -9.1 [-11.6, -6.7]
Test for Subgroup Differences: Q_M = 0.39, df = 1, p = 0.53

-60   -40   -20   0   10

Estimated Effect Size (Mean Difference)

Subgroup	No. of Experiments		SMD [95% CI]
Time of Administration (I^2 = 0 %)			
≤24 post-stroke	N=3		-0.80 [-1.35, -0.25]
Pre-Stroke	N=3		-1.48 [-2.83, -0.14]
Route of Administration (I^2 = 0 %)			
Intraperitoneally	N=2		-0.98 [-1.91, -0.06]
Subcutaneously	N=1		-3.62 [-5.64, -1.61]
Intracerebroventricular	N=2		-0.83 [-1.56, -0.11]
Intravenously	N=1		-0.70 [-1.38, -0.02]
Stroke Model (I^2 = 0 %)			
Permanent MCAO	N=1		-1.06 [-2.39, 0.26]
Photothrombotic	N=1		-0.91 [-2.21, 0.39]
Intraluminal Suture	N=4		-1.14 [-1.98, -0.29]
Species (I^2 = 0 %)			
Mouse	N=5		-0.81 [-1.25, -0.37]
Rat	N=1		-3.62 [-5.64, -1.61]
Drug (I^2 = 0 %)			
Maraviroc	N=2		-0.98 [-1.91, -0.06]
DAPTA	N=1		-3.62 [-5.64, -1.61]
TAK-779	N=3		-0.76 [-1.26, -0.26]
Dose (I^2 = 0 %)			
<10 mg/kg	N=3		-1.33 [-2.60, -0.07]
10-60 mg/kg	N=2		-1.00 [-1.80, -0.19]
>60 mg/kg	N=1		-0.91 [-2.21, 0.39]
Behaviour Tests (I^2 = 27.2 %)			
Motor	N=3		-1.67 [-3.12, -0.23]
None	N=3		-0.76 [-1.26, -0.26]

Favours Treatment   Favours Control

RE Model for All Studies (Q = 7.55, df = 5, p = 0.18; I^2 = 33.8%)   -1.02 [-1.58, -0.46]

-6 -5 -4 -3 -2 -1 0 1 2 3 4 5 6

Estimated Effect Size (Standardized Mean Difference)

Author	Year	Behaviour Test	Dose (mg/kg)	Time of Drug Administration	Time of Test (days)	Treatment (n)	Control (n)	SMD [95% CI]
Motor - Post-stroke administration of CCR5 antagonist								
Yan et al.	2021	Corner test	0.15	1 h post stroke	3	6	6	-4.22 [-6.25, -2.19]
Yan et al.	2021	Limb placement	0.15	1 h post stroke	3	6	6	-2.92 [-4.55, -1.29]
Yan et al.	2021	Modified Garcia	0.15	1 h post stroke	3	6	6	-2.70 [-4.27, -1.14]
Yan et al.	2021	Foot fault	0.15	1 h post stroke	21	8	8	-2.31 [-3.57, -1.04]
Yan et al.	2021	Rotarod	0.15	1 h post stroke	21	8	8	-1.88 [-3.06, -0.70]
Yan et al.	2021	Corner test	0.15	24 h post-stroke	3	6	6	-0.53 [-1.68, 0.62]
Yan et al.	2021	Limb placement	0.15	24 h post-stroke	3	6	6	-0.42 [-1.56, 0.72]
Yan et al.	2021	Modified Garcia	0.15	24 h post-stroke	3	6	6	-0.19 [-1.33, 0.94]
Joy et al.	2019	Cylinder task	100	24 h post-stroke + daily for 3 weeks	63	9	8	-1.60 [-2.69, -0.51]
Joy et al.	2019	Grid walk	100	24 h post-stroke + daily for 3 weeks	63	9	8	-1.19 [-2.23, -0.16]
Joy et al.	2019	Cylinder task	100	24 h post-stroke + daily for 9 weeks	56	10	10	-0.83 [-1.74, 0.08]
Joy et al.	2019	Grid walk	100	24 h post-stroke + daily for 9 weeks	56	10	10	-1.01 [-1.94, -0.08]
Joy et al.	2019	Cylinder task	100	21-28 d post-stroke + daily for 11 weeks	77	9	9	-0.27 [-1.20, 0.66]
Joy et al.	2019	Grid walk	100	21-28 d post-stroke + daily for 11 weeks	77	9	9	-1.66 [-2.73, -0.59]
Motor - Pre-stroke administration of CCR5 antagonist								
Li et al.	2016	Neurological deficit score	0.01	15 min pre-stroke	1	5	5	-1.04 [-2.36, 0.28]
Cognitive - Post-stroke administration of CCR5 antagonist								
Yan et al.	2021	Probe quadrant duration	0.15	1 h post stroke	25	8	8	-2.43 [-3.73, -1.14]

Favours Treatment   *Favours Control*

-5   -4   -3   -2   -1   0   1   2   3   4   5

Estimated Effect Size (Standardized Mean Difference)

	Randomization	Balanced Baselines Characteristics	Allocation Concealment	Random Housing	Investigators Blinded	Random Outcome Assessment	Outcome Assessors Blinded	Incomplete Outcome Data	Selective Reporting	Contamination	Funding	Unit Analysis Errors	Design Specific Risks	Addition of Animals	Overall
Chen et al. (2022)	?	−	−	−	−	−	?	X	X	+	+	−	−	+	X
Yan et al. (2021)	+	−	+	−	−	−	?	+	X	+	+	−	+	X	X
Joy et al. (2019)	?	−	−	−	−	−	?	X	X	+	+	−	−	+	X
Li et al. (2016)	?	−	−	−	−	−	−	X	+	+	+	−	+	+	X
Takami et al. (2002)	?	−	−	−	−	−	−	X	−	+	X	−	+	+	X

Judgement:

X High

− Unclear

+ Low

? Some concerns