Title: CCR5 antagonists as neuroprotective and stroke recovery enhancing agents: a preclinical systematic review and meta-analysis

Abbreviated Title: CCR5, neuroprotection, and stroke recovery

Authors: Ayni Sharif[1,2,*], Matthew S. Jeffers[1,2,*], Dean A. Fergusson[1,2,3], Raj Bapuji[1], Stuart G. Nicholls[4], John Humphrey[5], Warren Johnston[5], Ed Mitchell[5], Mary-Ann Speirs[5], Laura Stronghill[5], Michele Vuckovic[5], Susan Wulf[5], Risa Shorr[6], Dar Dowlatshahi[2,7,8], Dale Corbett[8], Manoj M. Lalu[1,2,8,9,#]

Affiliations: [1]Clinical Epidemiology Program, Blueprint Translational Research Group, Ottawa Hospital Research Institute, Ottawa, Canada
[2]School of Epidemiology and Public Health, University of Ottawa, Ottawa, Canada
[3]Department of Medicine, University of Ottawa, Ottawa, Canada
[4]Office for Patient Engagement in Research Activity (OPERA), Ottawa Hospital Research Institute, Ottawa, Canada
[5]Patient Partner, Blueprint Translational Research Group, Ottawa Hospital Research Institute, Ottawa, Canada
[6]Ottawa Hospital Library Services, The Ottawa Hospital, Ottawa, Canada
[7]Division of Neurology, Department of Medicine, The Ottawa Hospital, Ottawa, Canada
[8]Department of Cellular and Molecular Medicine, Faculty of Medicine, University of Ottawa, Ottawa, Canada
[9]Department of Anaesthesiology and Pain Medicine, The Ottawa Hospital, Ottawa, Canada
*These authors contributed equally

Corresponding Author:
[#]Manoj M. Lalu, mlalu@toh.ca
Department of Cellular and Molecular Medicine, University of Ottawa
Clinical Epidemiology and Regenerative Medicine Programs, Ottawa Hospital Research Institute
Ottawa, Ontario, Canada, K1Y 4E9

X: @ManojLalu

Number of Figures: 3
Number of Tables: 3
Word counts: Abstract (240), Main text (3,785)

ABSTRACT

Background: C-C chemokine receptor type 5 (CCR5) antagonists may improve both acute stroke outcome and long-term recovery. Despite their evaluation in ongoing clinical trials, gaps remain in the evidence supporting their use.

Methods: With a panel of patients with lived experiences of stroke, we performed a systematic review of animal models of stroke that administered a CCR5 antagonist and assessed infarct size or behavioral outcomes. MEDLINE, Web of Science, and Embase were searched. Article screening and data extraction were completed in duplicate. We pooled outcomes using random effects meta-analyses. We assessed risk of bias using the Systematic Review Centre for Laboratory Animal Experimentation (SYRCLE) tool and alignment with the Stroke Treatment Academic Industry Roundtable (STAIR) and Stroke Recovery and Rehabilitation Roundtable (SRRR) recommendations.

Results: Five studies representing 10 experiments were included. CCR5 antagonists reduced infarct volume (standard mean difference −1.02; 95% confidence interval −1.58 to −0.46) when compared to stroke-only controls. Varied timing of CCR5 administration (pre- or post-stroke induction) produced similar benefit. CCR5 antagonists significantly improved 11 of 16 behavioral outcomes reported. High risk of bias was present in all studies and critical knowledge gaps in the preclinical evidence were identified using STAIR/SRRR.

Conclusions: CCR5 antagonists demonstrate promise; however, rigorously designed preclinical studies that better align with STAIR/SRRR recommendations and downstream clinical trials are warranted.

Registration: Prospective Register of Systematic Reviews (PROSPERO CRD42023393438)

Key Words: stroke; neuroprotection; impairment; recovery; preclinical; CCR5 antagonists; maraviroc; patient engagement in research; CAMAROS

Non-standard Abbreviations and Acronyms:
CAMAROS   Canadian Maraviroc RCT to Augment Rehabilitation Outcomes After Stroke
CCR5   C-C chemokine receptor type 5
DAPTA   D-Ala-Peptide T-Amide
SRRR   Stroke Recovery and Rehabilitation Roundtable
STAIR   Stroke Treatment Academic Industry Roundtable
SYRCLE   Systematic Review Centre for Laboratory Animal Experimentation
TAK-779   Takeda 779

Introduction

C-C chemokine receptor type 5 (CCR5) is expressed across a variety of leukocyte subtypes, endothelial cells, and cell types in the brain (e.g., neurons, microglia, astrocytes), and is thought to play crucial roles in post-stroke neuroinflammation, blood-brain barrier repair, and neuronal survival / repair processes.[1] CCR5 antagonists have emerged as potential therapeutic candidates for stroke, demonstrating both neuroprotection and improved neural repair/recovery in preclinical animal models.[2–6] However, no CCR5 antagonist drug has an approved indication in the stroke context, necessitating studies to establish safety and efficacy this population. This has led to an ongoing clinical trial to investigate efficacy of CCR5 antagonists in combination with post-stroke rehabilitation.[7] Assessment of the preclinical evidence supporting CCR5's role in stroke is needed to identify areas of potential benefit, and knowledge gaps, that should be addressed by future preclinical research.[8]

The stroke neuroprotection and recovery communities have advocated for alignment of preclinical and clinical study parameters through publication of consensus recommendations for preclinical research, in an effort to enhance the translation of new stroke therapies.[9–11] Examples include identification of more sensitive and clinically relevant preclinical outcome measures and incorporation of potentially important effect modifiers of treatment efficacy, such as age, sex, and stroke-related comorbidities (hypertension, diabetes, etc.).[9–11] These recommendations aim to improve the translation of novel stroke therapeutics from preclinical to clinical populations, but the degree to which preclinical evidence for CCR5 antagonists satisfy these recommendations is unknown.

We sought to comprehensively evaluate the preclinical evidence for CCR5 antagonist drugs as both neuroprotective and stroke recovery-promoting agents.[9,11] Both perspectives are necessary to fully understand the therapeutic potential of stroke-related treatments, as distinct biological principles, time windows for treatment, and outcomes of interest underpin each of these treatment domains.[12,13] We conducted a systematic review and meta-analysis of the CCR5 literature in conjunction with a panel of individuals with lived experiences of stroke. This review sought to explore how the preclinical evidence for CCR5 antagonist drugs aligned with guidance for preclinical stroke research provided by previous expert committees and the parameters of an

ongoing clinical trial (The Canadian Maraviroc Randomized Controlled Trial To Augment

Rehabilitation Outcomes After Stroke, CAMAROS, NCT04789616).[9–11]

Materials and Methods

We registered the review protocol on the International Prospective Register of Systematic

Reviews (CRD42023393438).[14] The findings are reported in accordance with the Preferred

Reporting Items for Systematic Reviews and Meta-Analyses (Table S1)[15] and Guidance for

Reporting the Involvement of Patients and the Public (Table S2).[16] A panel of eight patients and

caregivers with lived and living experience of stroke informed project development and were

actively involved throughout research conduct (see Table S2 for more information).

Eligibility criteria

- *Animals:* Any preclinical *in vivo* animal models of adult stroke were included. Human,

invertebrate, *in vitro*, *ex vivo*, and neonatal animal studies were excluded.

- *Model:* Focal ischemic or intracerebral hemorrhagic stroke models were included, while

animal models of global ischemia were excluded.

- *Intervention:* Studies administering a CCR5 antagonist drug (e.g., maraviroc, D-Ala-

Peptide T-Amide (DAPTA), Takeda 779 (TAK-779)). Study arms in which CCR5 was

genetically manipulated (e.g., CCR5 knockout strain) were excluded.

- *Comparator:* Vehicle-treated control groups where stroke was induced. CCR5 antagonist

control groups without stroke were excluded.

- *Outcome:* Studies reporting at least one of the following: infarct size, behavioral tests,

mortality, adverse events, and spasticity were included.

- *Study design and publication characteristics:* Controlled interventional studies

(randomized, pseudo-randomized, or non-randomized) published as full journal articles

in any language or year were included. Abstracts, review articles, opinion-based

letters/editorials, and unpublished grey literature were excluded.

Information sources and search strategy

An information specialist with experience in systematic searches of the preclinical literature

developed a comprehensive search strategy on October 25, 2022, based on a previously

published strategy for identifying animal experimentation studies.[17] The search strategy

underwent peer-review using the Peer Review of Electronic Search Strategies (PRESS)

checklist.[18] We searched MEDLINE (OVID interface, including In-Process and Epub Ahead of

Print), Web of Science, and Embase (OVID interface). See the Supplemental Methods for the

full search strategy and exact search dates for each database.

Selection process and data collection

We deduplicated citations and uploaded them into DistillerSR® (Evidence Partners, Ottawa,

Canada). Two reviewers independently screened citations by title and abstract using an

accelerated method (one reviewer required to include, two reviewers required to exclude). We

then screened and extracted data from full-text articles in duplicate. Graphical data was extracted

using Engauge Digitizer.[19] A third reviewer with content expertise in preclinical stroke studies

audited all data extraction. Conflicts between reviewers were resolved by consensus discussion.

See Supplemental Methods for the complete list of data extraction elements.

Effect measures and data synthesis

We performed quantitative analyses using the R (version 4.1.2) "metafor" package (version

4.0.0)[20] with inverse variance random effects modelling. We expressed continuous outcome

measures as standardized mean differences (SMDs) with 95% confidence intervals (CIs) and

assessed statistical heterogeneity of effect sizes using the Cochrane I^2 statistic.[21] This was

necessary due to the variety of outcome measures and measurement scales used across studies,

which is a common feature of preclinical systematic reviews. Sensitivity analyses were

performed using original measurement scales where possible. From patient partner input,

subgroups were analyzed based on timing / dose / route of intervention, stroke model, stroke

type, species, type of behavioral outcome (i.e., motor, cognitive), and comorbidities.[22] Our

planned subgroup analyses on study quality, specific regions/areas of the brain, and post-stroke

rehabilitation paradigms were not performed due to insufficient number of studies. We did not

assess publication bias using Egger plots due to fewer than 10 studies being included in the

analysis, as per Cochrane recommendations.[23]

Risk of bias assessment

Two independent reviewers used the Systematic Review Centre for Laboratory Animal
Experimentation (SYRCLE) risk of bias tool to assess each study as having a "Low Risk",
"Unclear Risk", "Some Concerns of Risk", or "High Risk" across domains such as
randomization, blinding and outcome reporting.[24] "Some Concerns of Risk" indicated reporting
of a domain (i.e., randomization), but lacking methodological details. This differed from
"Unclear Risk" where there was no mention of the domain in the study.

Comprehensiveness of preclinical evidence and alignment with clinical trials
We assessed comprehensiveness of the *overall body* of preclinical evidence for CCR5
antagonists as a neuroprotective treatment using The Stroke Treatment Academic Industry
Roundtable (STAIR) I, VI, and XI Consolidated Recommendations.[11,25,26] These
recommendations encompass "candidate treatment qualification" (e.g., dose, timing of dose,
outcomes) and "preclinical assessment and validation" (e.g., age, sex, sample size, animal
type).[11] We excluded domains that were redundant with risk of bias (e.g., randomization,
blinding, etc.) and included additional items relevant to stroke recovery studies from the Stroke
Recovery and Rehabilitation Roundtable (SRRR) Translational Working Group consensus
recommendations.[9] Two reviewers extracted data to determine if the overall evidence across
studies satisfied each of the STAIR and SRRR recommendations. A third reviewer audited this
analysis. We then assessed alignment of existing preclinical evidence with an ongoing clinical
trial of CCR5 antagonists for stroke (CAMAROS, NCT04789616).[7]

Protocol deviations
We incorporated an additional assessment using the PRIMED[2] tool for assessing the readiness of
stroke neuroprotection therapies to be translated to clinical trials based on the body of preclinical
evidence.[27] Additionally, a list of outcomes used to determine CCR5's potential mechanisms of
action were extracted (Figure S1, Table S3) based on feedback from individuals that reviewed
the initial manuscript drafts.

Results
Study selection

Our search identified 263 citations, which was reduced to 166 unique studies after deduplication.

Five studies representing 10 experiments met the eligibility criteria (Figure 1).[2–6]

Study and animal model characteristics

Most studies used ischemic stroke (n=4/5). This was induced via intraluminal suture (n=2),

cauterization (n=1), or photothrombosis techniques (n=1; Table 1). Hemorrhagic stroke was

induced in one study via autologous whole blood injection. All studies used mouse (n=4) or rat

(n=1) models, comprised of exclusively male animals (n=5). Relevant stroke comorbidities

highlighted by patient partners and STAIR/SRRR recommendations (e.g., hypertension,

diabetes) were not used in any study.

Intervention characteristics

Maraviroc was used in six of the experiments (n=6/10), TAK-779 in three of the experiments

(n=3), and D-Ala-Peptide T-Amide (DAPTA) in one (n=1; Table 2). These CCR5 antagonists

were delivered intraperitoneally (n=4), intranasally (n=2), intracerebroventricularly (n=2),

subcutaneously (n=1), and intravenously (n=1) at a dose range from 0.01 to 100 mg/kg. Most

studies delivered a single dose of the drug (n=6); experiments with multiple administrations

(n=4) ranged from 3–63 doses. Time of initial treatment administration varied widely. Two

studies (3 experiments) administered treatment pre-stroke (10-15 minutes),[2,3] and 4 studies (6

experiments) in the acute, potentially neuroprotective, post-stroke period (50 minutes – 24 hours

post-stroke).[3–6] One study (1 experiment) was conducted in the late sub-acute/early chronic

period beginning at three to four weeks post stroke, which would be oriented towards recovery,

rather than neuroprotective, effects.[6] Patient partners had identified several *a priori* interests

(physical therapy alongside CCR5 administration, spasticity), which were also aligned with

SRRR recommended considerations for preclinical stroke recovery studies (see Supplemental

Methods).[9] These were not reported in any included studies.

Meta-analysis of infarct volume

Infarct volume was reported in six experiments (n=6/10) from four different studies with an

overall pooled analysis demonstrating marked neuroprotection with CCR5 antagonists (SMD -

1.02, 95% CI -1.58 to -0.46, p < 0.0001, I^2=34%; Figure 2). Five of these experiments measured

infarct volume at 1-3 days post-stroke, and one experiment measured infarct volume at a delayed

time point of 63 days post-stroke. No significant differences between pre- or post-stroke

administration were observed (P=0.47). Post-hoc sensitivity analysis removing one experiment

with extreme values[2] demonstrated that neuroprotection in the remaining two experiments

remained statistically significant while reducing heterogeneity (SMD -0.81, 95% CI -1.25 to -

0.37, p < 0.001, I^2=0%). A second sensitivity analysis excluded one study that measured infarct

volume in mm^3 so that all other studies could be meta-analyzed using mean differences on the

percentage scale (Figure S2; MD -9.1%, 95% CI -11.6 to -6.7%, p < 0.001, I^2=0%). This

demonstrated a similar neuroprotective effect as the other analyses. Further sub-group analyses

by route of administration, time of administration, stroke model, species, CCR5 antagonist, dose,

and whether behavior tests were assessed are described in Figure S3-S5.

Synthesis of behavioral outcomes without meta-analysis

Motor behavioral outcomes were reported in six experiments from three studies and represented

seven different behavioral tasks. An additional study reported motor behavioral outcomes

without standard deviations or standard errors, and thus could not be included (authors did not

respond to email requests for data).[4] A cognitive outcome (Morris Water Maze) was measured in

one study. Overall, CCR5 inhibition was effective in 11 of 16 behavioral outcomes tested (Figure

3). Meta-analysis and planned subgroup analysis were contra-indicated due to an inadequate

number of studies for each given outcome measure, which necessitated the synthesis without

meta-analysis presented below.[28]

Behavioral outcomes are presented by time of CCR5 antagonist administration, as discussed in

the Intervention Characteristics section above, as administration time directly relates to the

treatment context and patient population to which the results apply (i.e., acute neuroprotection

vs. late sub-acute/early chronic neural repair). Li et al. reported that pre-stroke administration

(relevant to surgical contexts with high risk of thrombosis) of DAPTA did not result in

significantly greater performance on the neurological deficit score.[2]

Regarding acute post-stroke administration times with the potential for *neuroprotective* effects

(up to 24 hours post-stroke based on observed infarct reductions in Figure 2), Yan et al. observed

that CCR5 antagonist (maraviroc) administration one-hour following stroke resulted in greater

motor performance on the corner test, limb placement, modified Garcia, foot fault, and rotarod

tasks compared to vehicle-treated controls. Cognitive outcomes in the Morris Water Maze task

(proportion of time spent in the probe quadrant) were also improved by this one-hour post-stroke

administration. Significantly improved motor performance was not observed if CCR5 antagonist

was administered 24 hours post-stroke, using this model of *intracerebral hemorrhage* (Figure

3).[5] In contrast, using a *focal ischemia* model, Joy et al. observed that CCR5 antagonist

(maraviroc) administration initiated 24 hours post-stroke and continued daily for 3 weeks (which

could potentially represent a mix of neuroprotective and recovery-induced effects) resulted in

greater performance on the cylinder and grid walk tasks compared to vehicle-treated controls.

Joy et al., also performed one experiment with *late sub-acute/early chronic* administration (a

potentially critical time for neural repair) initiated at three to four weeks post-stroke and

continued daily for 11 weeks, that demonstrated significantly improved performance for the grid

walk, but not cylinder, task.[6] This experiment was initiated outside the plausible range for a

neuroprotective effect, implying behavioral improvement involved *recovery-promoting*

mechanisms. However, equivalent infarct volumes were not demonstrated between the treated

and control groups in this cohort, which could potentially lead to confounding effects.

Risk of bias

All articles had a 'high' risk of bias in at least one domain of the SYRCLE tool.[24] Most domains

within each study demonstrated an 'unclear' risk of bias. All studies reported randomizing

animals; however, as commonly observed in the preclinical literature,[29–31] only one of these

studies provided sufficient detail to ensure that the randomization method had a low risk of bias.

The SYRCLE domains with the highest risk of bias were incomplete outcome data, with 80% of

studies (n=4/5) failing to provide complete data for all animals initially included in the study, as

well as selective outcome reporting, with 60% (n=3/5; all studies of maraviroc) not providing

complete data for all expected outcomes discussed in the methods (Figure S6).

Comprehensiveness of preclinical evidence and alignment with clinical trials

We assessed comprehensiveness of the preclinical evidence using the STAIR and SRRR

recommendations[9,11,25,26] as well as alignment with study parameters of the CAMAROS trial.[7] A

summary of assessment items from STAIR XI is provided in Table 3, with additional items from

STAIR I, VI, and SRRR recommendations included in Table S4.

For CCR5 antagonists as a post-stroke neuroprotectant, the overall body of evidence satisfied all

five STAIR XI domains assessing 'candidate treatment qualification' (Table 3). Overall, a range

of doses and clinically relevant administration times for neuroprotection were evaluated across a

variety of motor and cognitive behavioral domains. All studies tested both behavioral and

histological outcomes and demonstrated neuroprotective effects, but most studies failed to

measure and control post-stroke temperature, which could potentially confound the observed

neuroprotection (Table S4).[32] Most histological measurements were also assessed at <72 hours,

which could confound the observed neuroprotective effects if cell death was merely delayed.[32]

For CCR5 antagonists as a post-stroke recovery-inducing treatment, one experiment assessed the

effects of initiating CCR5 administration in a similar post-stroke phase as the CAMAROS trial.

This experiment (Joy et al.)[6] did not demonstrate that each treatment group had equivalent

baseline stroke volumes, which may potentially confound observed behavioral effects.

Furthermore, the maximum dose used in mice (100 mg/kg) was not sufficient to attain

cerebrospinal fluid levels of maraviroc observed in humans using the CAMAROS dosing regime

(mice: 13.8±5.4 ng/mL; humans 33.6–60 ng/mL).[6]

Areas of concern were identified in all STAIR XI domains assessing 'preclinical assessment and

validation' (Table 3). Although adult animals were used in most of the preclinical studies, the

reported weights and ages of these animals corresponded to young adults, rather than the aged

adults that better represent the stroke population. All experiments used only male rodents that

were free of common stroke comorbidities. It was also unclear if behavioral testing was

performed during the inactive circadian phase or active (dark) phase, which could result in

confounding if CCR5 antagonists affect arousal of animals during their inactive period.[33]

Furthermore, none of the studies had their protocols or results directly replicated by an

independent laboratory, or across multiple sites in a multilaboratory study.[9,34] Regarding stroke

recovery, no studies assessed behavioral effects on upper extremity skilled reaching / grasping or

potential interactions of CCR5 antagonists with rehabilitative therapies (Table S4).[35–37] Both

elements are highly relevant to the CAMAROS trial, as one of the primary outcomes of this trial

is upper extremity performance on the Fugl-Meyer and maraviroc administration will be paired

with an 8-week exercise program. These findings were supported by the PRIMED[2] tool, which

resulted in a Readiness for Translation Score of "medium" (on a scale of "low", "medium",

"high"). This tool highlighted similar promising elements, as well as weaknesses, as our analysis

above (Table 3, Table S4), identifying the limited preclinical evidence for the effects of CCR5

antagonists in clinically relevant sexes, ages, species, and disease comorbidities, without

sufficient dose-response information to inform trials (Table S5). Overall, our assessments

highlight a variety of knowledge gaps and areas of misalignment between the preclinical

evidence and clinical trial parameters that could be improved with further preclinical

experimentation.

Discussion

The overall body of preclinical evidence for CCR5 antagonists in stroke demonstrates potential

acute neuroprotection with corresponding impairment reduction, as well as improved functional

recovery in the sub-acute/early chronic phase. Our systematic review also highlights evidence

gaps that could impact successful clinical translation of CCR5 antagonists. Our analysis of 10

independent experiments, identified that acute administration of CCR5 antagonists within the

first 24 hours post-stroke was associated with a marked reduction in infarct volume. This

neuroprotective reduction of infarct volume did not significantly vary based on treatment dose or

any other experimental characteristics.[3,4,6] Overall, the majority of behavioral effects appeared to

be in a positive direction, but the low number of included studies precluded meta-analysis of

these results. Indeed, no individual behavioral experiment included more than ten CCR5-treated

animals, and given the wide range of dosages, timings, routes, stroke models, and rodent strains

involved, the certainty of these findings is limited and should be interpreted cautiously. Further

investigation is warranted to determine the optimal timing of administration and behavioral

domains under which CCR5 antagonists exhibit the strongest post-stroke neuroprotective and

recovery-inducing effects.

Despite the positive direction of treatment effects across all studies of CCR5 antagonists, we

found a substantial risk of bias in the underlying studies.[24] As is commonly observed in the

preclinical literature, all studies either did not adequately report their randomization / blinding

methods and exhibited evidence of selective / incomplete reporting.[29–31] Such features are

associated with biased overestimations of preclinical treatment efficacy, which raises further

concerns about the reliability and validity of the present findings.[38–40]

Comprehensiveness of the preclinical evidence for CCR5 antagonists was assessed in relation to

STAIR and SRRR consensus recommendations.[9,11,25,26] These recommendations aim to provide

investigators and regulators with "assurance that the candidate treatment shows signals of

efficacy and safety, before embarking on an expensive clinical development program".[11] The

included studies provide good initial evidence for acute neuroprotection, as well as mechanistic

and behavioral evidence for enhanced recovery in the late sub-acute/early chronic post-stroke

phase. However, demonstration of efficacy under a wider range of conditions, such as in aged

animals, females, animals with stroke-related comorbidities, more relevant timing of dose

administrations, or in conjunction with rehabilitative therapies are necessary to provide further

confidence in these findings. In addition, all studies used unique doses, timings, and outcomes,

so independent replication of the most promising study parameters would further increase

certainty in the evidence.[9,11]

In relation to the ongoing CAMAROS trial assessing maraviroc in the subacute post-stroke

phase,[7] the most relevant preclinical evidence comes from one experiment within the Joy et al.

study[6] where maraviroc was initially administered at three to four weeks post-stroke and

continued daily for 11 weeks. This experiment demonstrated that administration of maraviroc in

the late sub-acute/early chronic post-stroke phase improved functional recovery on the grid walk,

but not cylinder, task. These experimental conditions could potentially align with the putative

therapeutic window and outcomes of interest being assessed in CAMAROS (e.g. 10-minute walk

test co-primary outcome).[7] However, caution is warranted as this pivotal supporting preclinical

evidence is based on a low sample size (n=9). Moreover, potential differences in dosing, severity

of infarct, concomitant rehabilitative therapy, and other factors discussed above could influence

the degree to which these results successfully translate to the clinical environment. Nevertheless,

this study provides a plausible biological mechanism and "proof of concept" for how CCR5

antagonism might enhance neuroplasticity that improves functional recovery after stroke.

Our present synthesis of the preclinical evidence for CCR5 antagonists used novel approaches to

increase its utility for assessing certainty of the findings and identification of knowledge gaps.

First, we engaged patients with lived experiences of stroke throughout the review process to

ensure that our research questions, outcomes, and interpretations aligned with the priorities of the

ultimate end-user of stroke research. Second, we incorporated consensus recommendations for

both preclinical neuroprotection and recovery research as an evidence evaluation tool, which we

found often aligned with the priorities of our patient panel.[9,11,25,26] This guided assessment of the

alignment of preclinical evidence with parameters of ongoing clinical trials, as well as appraisal

of comprehensiveness of the preclinical evidence with a focus on translational validity[41] rather

than only internal validity and risk of bias.[24] We also used this method to provide concrete

avenues for future preclinical studies to close knowledge gaps and improve certainty in the

effects of CCR5 antagonists under clinically relevant experimental conditions. Similar

approaches should be considered by future preclinical systematic reviews to improve

interpretation of the preclinical evidence from a translational perspective.

In conclusion, CCR5 antagonists show promise in preclinical studies for stroke neuroprotection,

corresponding reduction in impairment, as well as improved functional recovery related to neural

repair in the late sub-acute/early chronic phase. However, high risk of bias and the limited (or

no) evidence in clinically relevant domains underscore the need for more rigorous and

transparent preclinical research to further strengthen the overall evidence base. Addressing these

concerns will not only enhance the reliability of preclinical evidence but also better inform the

design and execution of clinical trials of CCR5 antagonists, such as the ongoing CAMAROS

trial.[7] The integration of expert recommendations, such as STAIR and SRRR, should guide

future preclinical investigations and synthesis of the body of preclinical evidence in stroke

recovery research.[9,11] Our present approach serves as a template by which the preclinical

evidence supporting translation to clinical trials can be weighed when justifying early clinical

trials of novel interventions for stroke neuroprotection and recovery.

Acknowledgements: The authors thank Hannah Laquerre for her assistance with figure
generation. Figure S1 was created with BioRender.com.

Sources of Funding: This study was funded by a Social Accountability Grant from the
University of Ottawa's Faculty of Medicine. MSJ was supported by the Canadian Institutes of
Health Research (CIHR) and Vanier Canada Graduate Scholarship (CGV-186957). The funders
were not involved in the study design, collection, analysis and interpretation of data, writing the
manuscript, or in the decision to submit the article for publication. MML is supported by
The Ottawa Hospital Anesthesia Alternate Funds Association, a University of Ottawa Junior
Research Chair, and the Canadian Anesthesiologists' Society Career Scientist Award.

Disclosures: None.

Supplemental Material:

Supplemental Methods

Figures S1-S6

Tables S1-S5

References:

1. Jing K, Chen F, Shi X, Guo J, Liu X. Dual effect of C–C motif chemokine receptor 5 on ischemic stroke: More harm than benefit? *European journal of pharmacology*. 2023;953:175857–175857.

2. Li L, Zhi D, Shen Y, Liu K, Li H, Chen J. Effects of CC-chemokine receptor 5 on ROCK2 and P-MLC2 expression after focal cerebral ischaemia-reperfusion injury in rats. *Brain injury*. 2016;30:468–473.

3. Takami S, Minami M, Katayama T, Nagata I, Namura S, Satoh M. TAK-779, A Nonpeptide CC Chemokine Receptor Antagonist, Protects the Brain Against Focal Cerebral Ischemia in Mice. *Journal of cerebral blood flow and metabolism*. 2002;22:780–784.

4. Chen B, Cao P, Guo X, Yin M, Li X, Jiang L, Shao J, Chen X, Jiang C, Tao L, et al. Maraviroc, an inhibitor of chemokine receptor type 5, alleviates neuroinflammatory response after cerebral Ischemia/reperfusion injury via regulating MAPK/NF-κB signaling. *International immunopharmacology*. 2022;108:108755–108755.

5. Yan J, Xu W, Lenahan C, Huang L, Wen J, Li G, Hu X, Zheng W, Zhang JH, Tang J. CCR5 Activation Promotes NLRP1-Dependent Neuronal Pyroptosis via CCR5/PKA/CREB Pathway After Intracerebral Hemorrhage. *Stroke*. 2021;52:4021–4032.

6. Joy MT, Ben Assayag E, Shabashov-Stone D, Liraz-Zaltsman S, Mazzitelli J, Arenas M, Abduljawad N, Kliper E, Korczyn AD, Thareja NS, et al. CCR5 Is a Therapeutic Target for Recovery after Stroke and Traumatic Brain Injury. *Cell*. 2019;176:1143-1157.e13.

7. Dukelow S, Clinicaltrials.gov NCT04789616. The Canadian Maraviroc RCT To Augment Rehabilitation Outcomes After Stroke (CAMAROS) [Internet]. [cited 2023 Oct 20];Available from: https://clinicaltrials.gov/study/NCT04789616

8. O'Collins VE, Macleod MR, Donnan GA, Horky LL, van der Worp BH, Howells DW. 1,026 experimental treatments in acute stroke. *Ann Neurol*. 2006;59:467–477.

9. Corbett D, Carmichael ST, Murphy TH, Jones TA, Schwab ME, Jolkkonen J, Clarkson AN, Dancause N, Weiloch T, Johansen-Berg H, et al. Enhancing the Alignment of the Preclinical and Clinical Stroke Recovery Research Pipeline: Consensus-Based Core Recommendations From the Stroke Recovery and Rehabilitation Roundtable Translational Working Group. *Neurorehabilitation and neural repair*. 2017;31:699–707.

10. Bosetti F, Koenig JI, Ayata C, Back SA, Becker K, Broderick JP, Carmichael ST, Cho S, Cipolla MJ, Corbett D, et al. Translational Stroke Research: Vision and Opportunities. *Stroke*. 2017;48:2632–2637.

11. Lyden P, Buchan A, Boltze J, Fisher M. Top Priorities for Cerebroprotective Studies-A Paradigm Shift: Report From STAIR XI. *Stroke*. 2021;52:3063–3071.

12. Carmichael ST. Emergent properties of neural repair: elemental biology to therapeutic
concepts. *Annals of neurology*. 2016;79:895–906.

13. Murphy TH, Corbett D. Plasticity during stroke recovery: from synapse to behaviour. *Nature*
reviews. Neuroscience. 2009;10:861–872.

14. Lalu MM, Sharif A, Nicholls S, Dowlatshahi D, Fergusson DA, Corbett D, Bapuji R, Jeffers
MS, Humphrey J, Johnston W, et al. Evaluating the effect of CCR5 inhibitors in animal
models of stroke: A systematic review co-designed with patient partners [Internet]. 2023
[cited 2023 Oct 20];Available from:
https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=393438

15. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L,
Tetzlaff JM, Akl EA, Brennan SE, et al. The PRISMA 2020 statement: An updated guideline
for reporting systematic reviews. *PLoS medicine*. 2021;18:e1003583–e1003583.

16. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, Altman DG, Moher D,
Barber R, Denegri S, et al. GRIPP2 reporting checklists: tools to improve reporting of patient
and public involvement in research. *BMJ*. 2017;358:j3453–j3453.

17. Hooijmans CR, Tillema A, Leenaars M, Ritskes-Hoitinga M. Enhancing search efficiency by
means of a search filter for finding all studies on animal experimentation in PubMed.
Laboratory animals (London). 2010;44:170–175.

18. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer
Review of Electronic Search Strategies: 2015 Guideline Statement. *Journal of clinical*
epidemiology. 2016;75:40–46.

19. Mitchell M, Muftakhidinov B, Winchen T. Engauge Digitizer Software [Internet]. [cited
2023 Oct 20];Available from: http://markummitchell.github.io/engauge-digitizer/

20. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *Journal of*
statistical software. 2010;36.

21. Higgins JP, Li T, Deeks J. Chapter 6: Choosing effect measures and computing estimates of
effect [Internet]. In: Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch VA,
editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.4 (updated
August 2023). Cochrane; 2023 [cited 2023 May 1]. Available from:
www.training.cochrane.org/handbook

22. Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, Krakauer JW,
Boyd LA, Carmichael ST, Corbett D, et al. Agreed definitions and a shared vision for new
standards in stroke recovery research: The Stroke Recovery and Rehabilitation Roundtable
taskforce. *International journal of stroke*. 2017;12:444–450.

23. Page MJ, Higgins JP, Sterne JAC. Chapter 13: Assessing risk of bias due to missing results
in a synthesis [Internet]. In: Higgins JPT, Chandler J, Cumpston M, Li T, Page MJ, Welch
VA, editors. Cochrane Handbook for Systematic Reviews of Interventions version 6.4

(updated August 2023). Cochrane; 2023 [cited 2023 May 1]. Available from:
www.training.cochrane.org/handbook

24. Hooijmans CR, Rovers MM, de Vries RB, Leenaars M, Ritskes-Hoitinga M, Langendam
MW. SYRCLE's risk of bias tool for animal studies. *BMC Medical Research Methodology*.
2014;14:43.

25. Finklestein SP, Fisher M, Furlan AJ, Goldstein LB. Recommendations for standards
regarding preclinical neuroprotective and restorative drug development. *Stroke*.
1999;30:2752–2758.

26. Fisher M, Feuerstein G, Howells DW, Hum PD, Kent TA, Savitz SI, Lo EH. Update of the
Stroke Therapy Academic Industry Roundtable Preclinical Recommendations. *Stroke*.
2009;40:2244–2250.

27. Bahr-Hosseini M, Bikson M, Iacoboni M, Liebeskind DS, Hinman JD, Carmichael ST, Saver
JL. PRIMED2 Preclinical Evidence Scoring Tool to Assess Readiness for Translation of
Neuroprotection Therapies. *Translational stroke research*. 2022;13:222–227.

28. Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-
Boyce J, Ryan R, Shepperd S, Thomas J, et al. Synthesis without meta-analysis (SWiM) in
systematic reviews: reporting guideline. *BMJ*. 2020;368:l6890.

29. Avey MT, Moher D, Sullivan KJ, Fergusson D, Griffin G, Grimshaw JM, Hutton B, Lalu
MM, Macleod M, Marshall J, et al. The Devil Is in the Details: Incomplete Reporting in
Preclinical Animal Research. *PLoS One*. 2016;11:e0166733.

30. Fergusson DA, Wesch NL, Leung GJ, MacNeil JL, Conic I, Presseau J, Cobey KD, Diallo J-
S, Auer R, Kimmelman J, et al. Assessing the Completeness of Reporting in Preclinical
Oncolytic Virus Therapy Studies. *Mol Ther Oncolytics*. 2019;14:179–187.

31. Fergusson DA, Avey MT, Barron CC, Bocock M, Biefer KE, Boet S, Bourque SL, Conic I,
Chen K, Dong YY, et al. Reporting preclinical anesthesia study (REPEAT): Evaluating the
quality of reporting in the preclinical anesthesiology literature. *PLoS One*.
2019;14:e0215221.

32. Corbett D, Nurse S. The problem of assessing effective neuroprotection in experimental
cerebral ischemia. *Progress in neurobiology*. 1998;54:531–548.

33. MacLellan CL, Keough MB, Granter-Button S, Chernenko GA, Butt S, Corbett D. A Critical
Threshold of Rehabilitation Involving Brain-Derived Neurotrophic Factor Is Required for
Poststroke Recovery. *Neurorehabilitation and neural repair*. 2011;25:740–748.

34. Hunniford VT, Grudniewicz A, Fergusson DA, Montroy J, Grigor E, Lansdell C, Lalu MM.
A systematic assessment of preclinical multilaboratory studies and a comparison to single
laboratory studies. *eLife*. 2023;12.

35. Balbinot G, Schuch CP, Jeffers MS, McDonald MW, Livingston-Thomas JM, Corbett D.
Post-stroke kinematic analysis in rats reveals similar reaching abnormalities as humans.
Scientific reports. 2018;8:8738–13.

36. Jeffers MS, Corbett D. Synergistic Effects of Enriched Environment and Task-Specific
Reach Training on Poststroke Recovery of Motor Function. *Stroke*. 2018;49:1496–1503.

37. Jeffers MS, Karthikeyan S, Gomez-Smith M, Gasinzigwa S, Achenbach J, Feiten A, Corbett
D. Does Stroke Rehabilitation Really Matter? Part B: An Algorithm for Prescribing an
Effective Intensity of Rehabilitation. *Neurorehabilitation and neural repair*. 2018;32:73–83.

38. Holman C, Piper SK, Grittner U, Diamantaras AA, Kimmelman J, Siegerink B, Dirnagl U.
Where Have All the Rodents Gone? The Effects of Attrition in Experimental Research on
Cancer and Stroke. *PLoS Biology*. 2016;14:e1002331–e1002331.

39. Sena ES, van der Worp HB, Bath PMW, Howells DW, Macleod MR. Publication bias in
reports of animal stroke studies leads to major overstatement of efficacy. *PLoS Biology*.
2010;8:e1000344–e1000344.

40. Macleod MR, Bart Van Der Worp H, Sena ES, Howells DW, Dirnagl U, Donnan GA.
Evidence for the Efficacy of NXY-059 in Experimental Focal Cerebral Ischaemia Is
Confounded by Study Quality. *Stroke*. 2008;39:2824–2829.

41. Drude NI, Martinez Gamboa L, Danziger M, Dirnagl U, Toelch U. Improving preclinical
studies through replications. *eLife*. 10:e62101.

Figure

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) flow

diagram.

Figure 2. CCR5 antagonists reduce infarct volume. Data is presented as a forest plot with standardized mean differences and 95% confidence intervals. Effect sizes <0 favours drug treatment and >0 favours control. Data is stratified by timing of CCR5 antagonist administration (pre- or post-stroke induction). The 'RE Model for All Studies' represents a pooled estimate of the CCR5 antagonist drug effect on infarct volume from all studies combined. Separate pooled estimates are also reported for post-stroke and pre-stroke CCR5.

Figure 3. Meta-analysis for all included preclinical CCR5 antagonist studies that reported motor and/or cognitive behavioral outcomes. Data is presented as a forest plot with a standardized mean difference and 95% confidence intervals. Effect sizes <0 favours drug treatment and >0 favours control.

574 **Tables**

576 **Table 1.** Summary of study and animal model characteristics of included articles.

Author	Year	Country	Species	Strain	Stroke Type	Stroke Model	Sex	Weight	Age
Li et al.[2]	2016	China	Rat	Wistar	Ischemic	Intraluminal suture	Male	260–300 g	N/A
Takami et al.[3]	2002	Japan	Mice	ddY	Ischemic	Intraluminal suture	Male	N/A	4 weeks
Chen et al.[4]	2022	China	Mice	C57/BL6	Ischemic	Permanent middle cerebral artery occlusion	Male	25–30 g	8–10 weeks
Yan et al.[5]	2021	China	Mice	CD1	Hemorrhagic	Intracerebral hemorrhagic	Male	30–40 g	N/A
Joy et al.[6]	2019	USA	Mice	C57/BL6	Ischemic	Photothrombotic	Male	25–30 g	8–20 weeks

577
578

579   **Table 2.** Summary of intervention characteristics.

Author	Drug	Dose (mg/kg)	Route	Timing	Doses (total #)	Outcomes Measured (Treatment n/Control n)	Outcome Window (post-stroke))
Li et al.[2]	DAPTA (D-Ala-Peptide T-Amide)	0.01	SC	15 min pre-stroke	1	- Infarct volume (5/5) - Neurological deficit score (5/5)	24 h
Takami et al.[3]	TAK-779	5	ICV	10 min pre-stroke	1	- Infarct volume (10/6)	48 h
		50	ICV	10 min pre-stroke		- Infarct volume (13/6)	
		0.25	IV	50 min post-stroke		- Infarct volume (17/18)	
Chen et al.[4]	Maraviroc	20	IP	1.5 h, 24 h, and 48 h post-stroke	3	- Infarct volume (5/5) - Longa score (5/5) - Neurological deficit score (5/5)	72 h
Yan et al.[5]	Maraviroc	0.15	IN	1 h post-stroke	1	- Garcia test (6/6) - Limb placement (6/6) - Corner turn test (6/6)	72 h
						- Foot fault (8/8) - Rotarod (8/8)	3 weeks
						- Probe quadrant duration (8/8)	25 days
				24 h post-stroke		- Garcia test (6/6) - Limb placement (6/6) - Corner turn test (6/6)	72 h
Joy et al.[6]	Maraviroc	100	IP	24 h post-stroke through daily injections for 9 weeks	63	- Infarct volume (5/5)	9 weeks
						- Grid walk (10/10) - Forelimb (10/10)	8 weeks
				24 h post-stroke then daily for 3 weeks	21	- Grid walk (10/10) - Cylinder test (9/8)	9 weeks
				3 to 4 weeks post-stroke then daily for 11 weeks	56	- Grid walk (9/9) - Cylinder test (9/9)	11 weeks

580   Abbreviations: ICV = intracerebroventricular; IN = intranasal; IP = intraperitoneal; IV = intravenous; SC = subcutaneous

Table 3. Alignment of included preclinical studies with STAIR recommendations and CAMAROS trial parameters.

Recommendation	Overall preclinical evidence	CAMAROS trial alignment	Notes
Candidate treatment qualification			
(1) Establish treatment dose-response	✓	X	• **Preclinical** – doses across all studies from 0.01 to 100 mg/kg • 6 doses tested for infarct volume, 3 doses for behavioral effects • Joy et al. demonstrated that 100 mg/kg of maraviroc in mice results in cerebrospinal fluid levels lower than in humans (13.8±5.4 vs. 33.6–60 ng/mL) • **CAMAROS** – participants take 300 mg doses twice daily for 8 weeks • **Interpretation** – maximum preclinical dose may not represent clinical levels
(2) Treatment given after clinically relevant delayed times (one to 4.5 hours post-stroke)	✓	✓	• **Preclinical** – 3/5 studies (all maraviroc) administered drug from 1 to 24 hours post-stroke (acute phase) • Joy et al. initiated maraviroc administration at 20 days post-stroke for 11 weeks (subacute phase) • **CAMAROS** – participants recruited between 5 days and 6 weeks post-stroke (acute / early subacute) • **Interpretation** – one experiment in a time window relevant to clinical trial
(3) Both histologic and behavioral outcomes at acute (1 to 3 days) and chronic (7 to 30 days) time points	✓	✓	• **Preclinical** – 2/5 studies included both histologic and behavioral outcomes • Mainly tasks of spontaneous movement (e.g., cylinder) or motor coordination (e.g., grid walk) • Tests ranged from 3 days (acute) to 11 weeks (chronic) post-stroke for all included studies, but short term assessment may only delay cell death[32] • **CAMAROS** - motor learning (Fugl-Meyer Upper Extremity Assessment Score and 10-Meter Walk Test Score) measured as the primary outcome at baseline, 4-week (late subacute), 8-week (late subacute), and 6-month (chronic) • **Interpretation** – challenging to directly compare preclinical and clinical motor tasks, but testing range is comparable. Most relevant Joy et al. experiment (subacute administration) did not measure histologic outcomes
(4) Treatment reaches presumed target and causes expected physiological effects that can be assessed with a clinically relevant biomarker	✓	NA	• **Preclinical –** 5/5 studies demonstrated trends for CCR5 antagonism to reduce infarct volume. Plausible that CCR5 influences stroke-relevant mechanisms • **CAMAROS** – does not include outcomes to assess treatment mechanisms
(5) Treatment able to pass the blood-brain barrier	✓	NA	• **Preclinical** – 1/5 studies used mass spectrometry to demonstrate that maraviroc is present in the brain and cerebrospinal fluid • **CAMAROS** – does not include outcomes to assess presence of drug in the brain

Table 3 (continued). Alignment of included preclinical studies with STAIR recommendations and CAMAROS trial parameters.

Recommendation	Overall preclinical evidence	CAMAROS trial alignment	Notes
Preclinical assessment / validation			
(1) Aging/adult age	X	X	• **Preclinical** – 4/5 studies used rodents with weights/ages consistent with young adulthood; 0/5 studies used rodents ≥10 months old. • **CAMAROS** – Age ≥18 year old adult participants eligible for recruitment • **Interpretation** – Preclinical studies consistent with CAMAROS eligibility criteria; however, no preclinical study examined middle-aged or elderly animals that would be more representative of the trial and overall stroke populations
(2) Male and female animals	X	X	• **Preclinical** – 0/5 studies included female animals • **CAMAROS** – both male and female sexes eligible for recruitment • **Interpretation** – effects of CCR5 antagonists for stroke recovery in female animals represents a critical knowledge gap
(3) Animals with co-morbidities	X	X	• **Preclinical** – 0/5 studies included animals with common stroke co-morbidities (e.g., diabetes, hypertension, etc.) • **CAMAROS** - participants with stroke co-morbidities eligible for recruitment • **Interpretation -** effects of CCR5 antagonists for stroke recovery in animals with common stroke comorbidities represents a critical knowledge gap
(4) Evidence from two or more laboratories	✓	X	• **Preclinical** – 5/5 studies demonstrated stroke-related benefits; however, no study had their results replicated by independent laboratories • **CAMAROS** – trial protocol will be replicated by multiple study sites • **Interpretation** – future preclinical experiments could consider assessing effects using a multilaboratory approach, or independently replicate the effects of existing preclinical studies
(5) Gyrencephalic species	X	X	• **Preclinical** – 5/5 studies conducted in lissencephalic (rodent) species. • **CAMAROS** – humans participants • **Interpretation** – effects of CCR5 antagonists for stroke recovery in gyrencephalic species represents a critical knowledge gap
(6) Tests during the awake phase of animals	X	X	• **Preclinical** – 0/5 studies reported assessing outcomes at night (i.e. awake phase of rodents) • **CAMAROS** – follow-up occurs during daytime hours (i.e. awake phase of humans) • **Interpretation** – future preclinical studies should conduct behavioral testing during the awake phase to better align with human testing conditions

584