Co-occurrence of diverse defense systems shapes complex microbe-virus relationships in deep-sea cold seeps

Yingchun Han[1][#], Jing Liao[1][#], Chengpeng Li[2], Fengmin Xing[3],

Jiaxue Peng[4], Xinyue Liu[1], Wentao Xie[5], Fabai Wu[6], Huahua Jian[5],

Rui Cheng[7, 8][*], Xiyang Dong[1][*]

[1] Key Laboratory of Marine Genetic Resources, Third Institute of Oceanography, Ministry of Natural Resources, Xiamen 361005, China

[2] Key Laboratory of Advanced Marine Materials, Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

[3] Institute of Marine Biology and Pharmacology, Ocean College, Zhejiang University, Zhoushan 316021, China

[4] College of Environmental Science and Engineering, Dalian Maritime University, Dalian 116026, China

[5] State Key Laboratory of Microbial Metabolism, Joint International Research Laboratory of Metabolic & Development Sciences, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China

[6] School of Life Sciences, College of Science, Eastern Institute of Technology, Ningbo 315200, China

[7] National Key Laboratory of Agricultural Microbiology, College of Biomedicine and Health, Huazhong Agricultural University, Wuhan 430070, China

[8] National Key Laboratory of Agricultural Microbiology, College of Life Science and Technology, Huazhong Agricultural University, Wuhan 430070, China

[#] These authors contributed equally to this work.

[*] Correspondence can be addressed to Xiyang Dong (dongxiyang@tio.org.cn) and Rui Cheng (ruicheng@hzau.edu.cn).

Abstract

Cold seeps host diverse microbes and viruses with numerous unexplored defense and anti-defense systems. Analysis of 3,813 microbial and 13,336 viral genomes from 191 metagenomes across 17 cold seep sites reveals extensive microbial defense repertoires, with over 60% representing candidate systems. Experimental validation confirms that several candidates protect against viral infection. These defense systems frequently co-occur, suggesting potential synergistic interactions, and are broadly distributed across sediments. In response, viruses have evolved diverse anti-defense genes, and the concurrent presence of multiple viral and microbial systems highlights intricate coevolution. Functionally critical lineages, such as anaerobic methanotrophic archaea, sulfate-reducing bacteria, and diazotrophs, appear to modify their defensive strategies under ecological and environmental pressures; for example, sulfate-reducing bacteria harbor multiple Gabija systems while corresponding viruses carry anti-Gabija genes, illustrating specific coevolutionary adaptations. Overall, these findings underscore the critical role of virus-microbe interactions in shaping microbial metabolic functions and environmental adaptation in deep-sea ecosystems.

Introduction

Cold seeps are extreme deep-sea environments located along continental margins worldwide, characterized by the migration of fluids rich in methane and other hydrocarbons from the deep subsurface to the sediment-seawater interface[1, 2]. The continuous release of these fluids enhances local biodiversity and microbial activity, making cold seeps thriving deep-sea oases[2]. These ecosystems support diverse microbial communities, including anaerobic methanotrophic archaea (ANME) and their syntrophic sulfate-reducing bacterial partners[3, 4]. Moreover, the sediments harbor a rich diversity of previously understudied prokaryotes and viruses[5, 6], suggesting the existence of complex and novel microbe-virus interactions within cold seep ecosystems. Prokaryotes in these environments constantly face viral threats, leading to the evolution of multiple defense mechanisms that regulate the flow of genetic information spread by mobile genetic elements (MGEs) through horizontal gene transfer (HGT)[7-9]. Cold seep viruses also employ diverse strategies for environmental adaptation, including counter-defense systems, auxiliary metabolic genes (AMGs), reverse transcriptases, and alternative genetic code assignments[10]. Among these, anti-defense systems have evolved to counteract the defense mechanisms of host microbes. Despite these insights, the diversity of prokaryotic defense genes and the intricate evolutionary interactions between prokaryotes and viruses in cold seep ecosystems remain largely unexplored.

Prokaryotes have developed an arsenal of defense systems to provide partial or full resistance against viral and MGE infections[7, 8, 11]. The most widespread prokaryotic immune systems are the restriction-modification (RM) systems and the clustered regularly interspaced short palindromic repeats CRISPR-Cas system. RM systems recognize specific DNA sequences in invading phages and distinguish self from non-self DNA through methylation[12], while CRISPR-Cas provides acquired immunity by incorporating short viral and mobile genetic element (MGE) sequences as spacers in the host genome[9, 13]. In recent years, many novel prokaryotic antiviral systems have

been revealed. For example, Gabija system, composed of GajA and GajB proteins, degrades viral and host DNA through site-specific double-stranded DNA nicking by GajA upon activation[14]. The cyclic oligonucleotide-based antiphage signaling system (CBASS) uses cyclic oligonucleotides to activate antiviral immunity[15]. Gabija and CBASS systems defend against phages by inducing cell death or metabolic disturbance upon infection, an abortive infection mechanism that is similarly adopted by various other defense systems[14, 16]. Recent efforts have uncovered several defense systems with diverse genetic architectures[7, 8], reflecting the strong selective pressure imposed by viruses and MGEs on microbial communities in various ecosystems such as soil, marine, human gut and rumen[9, 17]. However, in the deep sea, particularly in cold seep environments, there is a lack of corresponding studies on microbial defense systems, despite these environments potentially equipped with vast and untapped wealth of immune systems and mechanisms[5, 6].

Defense genes encoding components from multiple distinct systems tend to cluster together in specific genomic regions referred to as defense islands[9, 18, 19]. This co-localization, especially within MGEs, facilitates joint HGT, which is thought to confer fitness advantages to recipient microbes, particularly in virus-rich environments[20]. Synergistic interactions between different defense systems may drive their co-localization and favor their joint transfer[21, 22], as evidenced by the conservation of certain sets of defense systems across different microbial species. Synergistic systems enhance protective effects against specific phages beyond the sum of individual systems, providing microbes with an evolutionary advantage. Recent research indicated that non-random synergistic interactions between defense systems in *Escherichia coli* are adaptive responses to specific selective forces from the virome, shaping unique fitness landscapes in their specific ecological niches[23]. Given that cold seep viruses in extreme deep-sea subseafloor conditions are endemic and unique compared to those in other ecosystems[6, 10], the specific virus-driven synergistic interactions between defense systems may differ and warrant investigation.

Viruses can evade microbial defenses both indirectly, by mutating immune system

activators or disrupting defense pathways, and directly, by deploying anti-defense

genes that target immune system components[24]. In this ongoing arms race, viruses

have evolved sophisticated anti-defense mechanisms to overcome microbial defense

systems[24, 25], such as anti-CRISPRs (Acr)[26, 27], anti-RM[28], anti-CBASS[29] and anti-

Gabija[30]. These viral anti-defense systems not only highlight the complexity of viral

adaptations but also reveal evolutionary connections within microbial immunity[24].

Beyond defense evasion, viruses play a crucial role in shaping microbial communities

and influencing ecosystem functioning, including key biogeochemical cycles such as

methane oxidation, sulfate reduction and nitrogen fixation, often mediated through

AMGs[31, 32]. Despite their importance, the interactions between viral anti-defense

mechanisms and functionally critical lineages in the cold seep environment[25]—with

its darkness, low temperatures, and hydrocarbon-rich conditions—is not well

understood. Understanding these dynamics in extreme deep-sea ecosystem is vital for

unraveling the full impact of viruses on microbial ecology and biogeochemical

processes.

To explore the diversity of cold seep microbial defense systems, their interactions, and

the evolutionary associations between these systems and viral anti-defense strategies

in extreme deep-sea subseafloor habitats, we analyze 3,813 microbial genomes and

13,336 viral genomes from 191 metagenomic samples collected at 17 global sites

(Supplementary Fig. 1). Our findings reveal that cold seep microbes possess a vast

and diverse repertoire of defense systems, which may work synergistically against

viral infections. In response, cold seep viruses deploy diverse anti-defense systems to

counter microbial defenses. This complex virus-microbe interaction drives microbial

evolution and enhances environmental adaptation. Overall, this study enriches our

understanding of the ecological and evolutionary dynamics of virus-microbe

interactions in deep sea cold seep subsurface ecosystems.

Results and Discussion

Cold seep prokaryotes possess diverse and putative defense systems

We investigated defense strategies of cold seep microbiome by analyzing 3,813 species-level metagenome-assembled genomes (MAGs), comprising 639 archaeal and 3,174 bacterial genomes, spanning 107 bacterial and 16 archaeal phyla **(Supplementary Fig. 2)**. The bacterial phyla with the largest diversity of recovered species included *Pseudomonadota* (n = 471), *Chloroflexota* (n = 399), *Desulfobacterota* (n = 298), *Planctomycetota* (n = 225), and *Bacteroidota* (n = 219). Among the archaeal phyla, *Halobacteriota* (n = 153), *Thermoplasmatota* (n = 119), *Thermoproteota* (n = 110), and *Asgardarchaeota* (n = 105) were the most diverse.

We identified defense genes and systems using DefenseFinder[19] and PADLOC[33] and detected a total of 36,783 defense genes in 65% of cold seep microbial genomes, which were assigned to 26,389 defense systems **(Fig. 1a and Supplementary Data 1-3)**. Among these, 63% (n = 16,520) were annotated as candidate systems, primarily including HECs (Hma-Embedded Candidates) and PDCs (Phage Defense Candidates). These candidates were predicted by PADLOC based on their recurrent genetic embedding within known defense loci (e.g., Hma, RM, BREX, DISARM) rather than defense-island enrichment[34]. Together, these candidates comprised 18,195 defense genes, highlighting that cold seep microbes harbor a diverse array of putative antiviral strategies. Experimentally verified defense systems, comprising 37% (n = 9,869), included 106 types, with RM systems (n = 3,778) being the most prevalent, followed by CRISPR-Cas (n = 977), SoFic (n = 715), and AbiE (n = 612). This distribution aligns with previous findings that RM and CRISPR-Cas systems are predominant in microbiomes from soil, human gut, and Haima cold seep environments[9, 10]. In contrast, seawater microbial genomes, while also rich in RM systems, exhibit significant enrichment of CBASS[9], suggesting distinct distribution patterns of defense systems in cold seep sediments compared to other marine environments.

To validate whether these candidate systems confer antiviral activity, we randomly

selected eight candidate systems from the top 12 most abundant system types (each

accounting for more than 1% of the total), including PDC-S01, PDC-S09, PDC-S05,

PDC-S70, PDC-S27, PDC-S02, PDC-M05, and HEC-01. Within each selected type,

one representative gene cluster was randomly chosen for experimental validation.

Recombinant vectors containing these candidate systems were transformed into the

host strain *E. coli* B (ATCC® 11303™), which naturally lacks these systems. The

transformed strains were then challenged with three representative lytic phages: T7

(*Podoviridae*), T5 (*Siphoviridae*), and T4 (*Myoviridae*). Among them, four candidate

defense systems (PDC-S05, PDC-S70, PDC-M05, and HEC-01) conferred protection

against at least one phage **(Fig. 1b)**. Especially, the PDC-S05 and PDC-S70 systems

exhibited obvious and broad-spectrum phage resistance. The PDC-S05 system

showed an approximately 1,000-fold reduction in efficiency of plating (EOP) for

phages T7 and T5, and a ~10-fold decrease for phage T4. Similarly, the PDC-S70

system caused a ~100-fold decrease in EOP for phages T7 and T4, and an ~1,000-fold

decrease for phage T5. Although these results confirmed the antiviral activity of four

systems, the vast, yet candidate defense systems require further experimental

validation. Predicted protein domain analysis of these four systems **(Supplementary**

Data 4) revealed that HEC-01 contains an AAA_21 domain, which is associated with

ATP binding and hydrolysis activity. PDC-M05A harbors ArsR-like_HTH and

NT_KNTase_like domains, suggesting DNA-binding transcription factor activity.

Additionally, PDC-S05 features a Virulence_RhuM domain, which is linked to

virulence and pathogenicity **(Fig. 1b)**. These findings indicate that most of these

candidate defense systems may function through mechanisms such as nucleic acid

degradation and replication inhibition.

Defense system mechanisms can be categorized in three types based on their

fundamental mechanisms of action: nucleic acid degradation, abortive infection,

inhibition of replication, and other unknown mechanisms[19, 35]. For instance, RM and

CRISPR-Cas systems function through viral nucleic acid degradation, CBASS and

retrons through abortive infection, and Viperins and dCTP deaminases through

inhibition of replication. Among the experimentally confirmed defense systems in

cold seep microbes **(Fig. 1a)**, the most prevalent mechanism was nucleic acid

degradation (n = 5,416, 55%), followed by abortive infection (n = 1,867, 19%) and

replication inhibition (n = 722, 7%). Notably, a substantial proportion of defense

mechanisms remain unidentified (n = 1,864, 19%), encompassing 59 types that

represent 56% of all detected defense system types (n = 106).

Cold seep defensive patterns are associated with microbial taxonomy

The distribution of verified defense systems differed across taxonomical domains **(Fig.**

2a and Supplementary Fig. 3), suggesting that archaea and bacteria possess diverse

and distinct defense strategies against viral or MGE infections[20, 33, 36]. Most bacterial

and archaeal MAGs harbored a limited number of defense systems, averaging four per

genome **(Fig. 2b and 2c)**, consistent with observations from other marine

environments[9, 10]. However, this value is lower than the average numbers reported for

bacterial genomes (5.6-5.8)[8, 19] and archaeal genomes (4.55)[37]. This discrepancy may

be attributed to the completeness of MAGs used in this study, differences in detection

methods and parameter settings, and the unique ecological pressures of cold seep

environments. On average, bacterial genomes in cold seep environments still encode

more defense systems per genome than archaeal genomes (4.1 vs 3.5, **Supplementary**

Data 5). Despite this average, several genomes (n = 40) encoded more than 20

defense systems. For instance, a genome belonging to the *Bacteroidales* order encodes

54 defense systems, followed by 48 in a *Methylococcales* genome

(*Gammaproteobacteria*) and 39 in a *Desulfuromonadales* genome (*Desulfobacterota*)

(Supplementary Data 3). Additionally, we found that the majority of defense systems

(~99.88%) are encoded in chromosomes, several systems—especially AbiV, AbiD,

DRT, and AbiL—are carried by plasmids, and others like AbiZ, PT_Dnd, and

Lamassu are carried by proviruses **(Fig. 3 and Supplementary Fig. 4a)**. Furthermore,

the number of defense systems in prokaryotes increases with the number of proviruses

and plasmids, a trend particularly evident in bacteria **(Supplementary Fig. 4b)**, suggesting that defense genes in cold seep prokaryotes could be disseminated between microbes through MGEs[38, 39].

The density of defense systems (per MAG and per Mb) varied widely among different clades **(Fig. 3, Supplementary Figs. 5-6, and Supplementary Data 3)**. In archaeal MAGs, the density ranged from as low as 0.11 in certain phyla to over 10.5 in the *Aenigmatarchaeota* phylum (**Fig. 3a and Supplementary Fig. 5**). In bacterial MAGs, the density can be as high as 15.1 in the *Desulfobacterota* phylum (**Fig. 3b and Supplementary Fig. 6**). While previous studies have suggested that deeper phylogenetic lineages accumulate more horizontally transferred genes, potentially leading to a buildup of defense systems[23], our linear regression analyses indicate that phylogenetic depth has a relatively minor impact on the number of defense systems **(Supplementary Data 6)**. This suggests that other factors, such as environmental pressures, viral diversity, and ecological interactions, may play a more significant role in shaping the abundance and distribution of defense systems.

Cold seep defense systems tend to co-occur and may interact synergistically

Previous studies have demonstrated that certain defense systems interact to enhance or broaden protection against viruses[23, 40, 41]. Here, based on the relative abundances of defense systems, we used Pearson correlation analysis to investigate the relationships among defense systems in cold seep prokaryotes to explore their potential interactions. We considered only statistically significant correlations ($p < 0.05$), following the approach of Wu et al.[42]. We identified 182 pairs of defense systems exhibiting significant positive correlations and 10 pairs with significant negative correlations **(Fig. 4a and Supplementary Data 7)**. Among these significantly correlated pairs, 94.8% were positive, suggesting that defense systems are more likely to co-occur rather than to be negatively associated. This pattern is consistent with observations in certain bacterial genomes, such as *E. coli*, *Enterobacterales, Bacillales,*

Burkholderiales, and *Pseudomonadales*[23]. Notably, prevalent defense systems such as

Gabija (20 pairs), RM (19 pairs), CRISPR-Cas (19 pairs), and SoFic (18 pairs)

exhibited a higher number of significant co-occurrences. Additionally, less common

systems such as Septu (18 pairs), Druantia (15 pairs), and Menshen (15 pairs), also

demonstrated a tendency to co-occur with other defense systems. This suggests that

the significant co-occurrence of these systems is not merely a result of their higher

overall frequency but may also stem from functional complementarities[22, 41]. For

instance, CRISPR-Cas and RM systems can work synergistically by targeting

different sites on invading phage DNA, thereby enhancing overall protection[41].

Similarly, defense islands encoding both BREX systems and type IV restriction

enzymes have been shown to provide layered protection[22]. Furthermore, RM-

mediated cleavage of viral DNA can stimulate spacer acquisition in CRISPR-Cas

systems, reinforcing their cooperative function[43]. Interestingly, different subtypes of

the same defense system exhibited distinct co-occurrence patterns with other systems

(Supplementary Fig. 7 and Supplementary Data 8). For example, Wadjet I co-

occurred with multiple other defense systems, showing 11 pairs of co-occurrences. In

contrast, Wadjet II had only 4 co-occurrences, which included CRISPR-Cas I-G,

CBASS III, Eleos, and PD-T4-3; notably, none of these systems co-occurred with

Wadjet I. These specific co-occurrence patterns may indicate unique cooperative

interactions among different subtypes of defense systems.

Defense islands are sometimes associated with MGEs integrated into distinct

hotspots—regions of high genetic turnover—that may serve as catalysts for novel

defensive strategies[9, 23]. The clustering of defense systems in these islands, especially

within integrated MGEs, enhances the probability of horizontal co-transfer. If the

repertoire of defense systems is primarily shaped by HGT, rather than functional

synergies, we would expect the systems co-localizing in defense islands to correspond

with those co-occurring in microbial genomes[23]. To explore the potential link between

the co-occurrence and co-localization of defense systems, we selected 34 defense

systems clustered in defense islands and analyzed the correlations among them.

Among the clustered systems, 105 pairs were identified with significant correlations

($p < 0.05$), which is lower than the 182 significant pairs observed when considering

all defense systems. Notably, only one pair among the clustered systems exhibited a

negative correlation **(Fig. 4a and Supplementary Data 9)**. Furthermore, there

appears to be no obvious relationship between the number of defense systems

clustered in defense islands and the number of significantly correlated pairs among

those systems **(Fig. 4a and Supplementary Fig. 8)**. For instance, although RM and

CRISPR-Cas systems were frequently found in defense islands, they did not exhibit

an obvious higher number of correlations with other systems. In contrast, systems like

BREX (13 pairs), Septu (13 pairs), and PT_PbeABCD (12 pairs), which were less

commonly associated with defense islands, showed more significant correlations.

These findings suggest that co-localization represents only a specific instance of co-

occurrence, and the repertoire of defense genes is influenced by various factors

beyond HGT.

The genomic context of defense islands supports the observed correlations among

defense systems. For instance, the CBASS system frequently clusters within these

islands, often flanked by defense genes related to RM and Abi systems **(Fig. 4b and**

Supplementary Data 10). These defense islands are not limited to defense genes

alone; they also encompass a rich array of functional genes involved in transport,

replication, recombination, repair, and transcription. This high genetic turnover within

defense islands creates hotspots of genetic diversity, promoting the evolution of novel

defense strategies in cold seep microorganisms[9]. Thus, the clustering and dynamic

interactions within defense islands highlight their role in the co-localization and co-

evolution of defense systems.

Cold seep defense systems are abundant, actively expressed and site-dependent

To assess the distribution and activity of defense systems in cold seep prokaryotes, we

calculated the relative gene abundance and transcript abundance of various defense

genes and systems. In alignment with the prevalence of defense systems in microbial genomes **(Fig. 1a)**, RM, CRISPR-Cas, AbiE and SoFic systems exhibit higher metagenomic abundances, primarily due to the elevated abundances of restriction endonucleases (REases), methyltransferases (MTases), AbiEi, AbiEii, SoFic proteins, and Cas2 genes associated with these systems **(Fig. 5a and Supplementary Fig. 9)**. These differences are statistically significant at both systems and gene levels ($p < 0.001$). Although there were no significant differences in metatranscriptomic abundances among defense systems, the detection of their expression levels (averaging 3.64 transcripts per million [TPM], with over 1 TPM for 30 systems) indicate microbial defensive responses are active *in situ*. Additionally, most defense systems were detected across geographically diverse cold seep sediments (17 sites) at almost every sediment depth (0-68.55 meters below seafloor [mbsf]; **Supplementary Fig. 10)**, indicating a widespread presence of these systems in cold seeps globally. High abundances of defense systems were particularly notable in archaeal *Halobacteriota*, and bacterial *Desulfobacterota*, *Atribacterota*, and *Pseudomonadota* **(Fig. 5b)**. The relative abundance of defense systems varied considerably across phyla, with RM, CRISPR-Cas, AbiE and SoFic systems being the most predominant.

The distribution patterns of defense systems in global cold seep prokaryotes are shaped not only by host microbial taxa but also by their targets, such as viruses and MGEs. The distribution of cold seep microbes and viruses is influenced by sediment depth and site location[10, 44, 45]; consequently, these factors may also impact the distribution of defense systems. To assess potential depth stratification of defense systems, we categorized each metagenome based on depth ranges: surface (<1 mbsf), shallow (1 to 10 mbsf), and deep (>10 mbsf). Our analyses revealed no significant differences in the abundance of defense systems across these depth categories at either the metagenomic or metatranscriptomic level **(Fig. 5c, 5d and Supplementary Fig. 11)**. Significant differences in defense system abundance were found among different cold seep sites, with more pronounced differences in shallower depths (surface and shallow, $p \leq 0.05$) compared to deeper depths (deep, $p > 0.05$; **Supplementary Fig.**

12a). Metatranscriptomic abundance also varied significantly across different sites, with a more notable difference observed in surface depths ($p < 0.01$; **Supplementary Fig. 12b**). Overall, these results suggest that the distribution of defense systems in cold seep prokaryotes is more strongly influenced by site-specific environmental and ecological factors than by sediment depth.

Cold seep viruses employ diverse anti-defense genes to counteract host defense systems

We retrieved 13,336 single-contig viral genomes with estimated completeness of greater than 50% from 191 cold seep metagenomes **(Supplementary Fig. 13 and Supplementary Data 11)** using three virus identification tools: VIBRANT[46], Virsorter2[47] and geNomad[48]. These viral genomes were clustered into 9,971 species-level viral operational taxonomic units (vOTUs)[49]. Of these, 9,638 vOTUs (~97%) could be taxonomically assigned, predominantly belonging to the *Duplodnaviria* phylum (n = 9,600), with a majority classified under the *Caudoviricetes* class (n = 9,599; **Supplementary Fig. 14 and Supplementary Data 12)**. However, less than 0.4% vOTUs (n = 36) within this class could be annotated further, highlighting an obvious knowledge gap in the taxonomy of deep-sea cold seep viruses[6, 10].

We used the iPHoP v1.3.2 pipeline[50] and CRISPR spacer sequence alignment to infer host-virus linkages, and identified a total of 2,688 connections between prokaryotes (810 genomes) and viruses (1,310 vOTUs; **Fig. 6a, Supplementary Fig. 15 and Supplementary Data 13)**. These host-virus linkages spanned 57 phyla, including 10 archaeal and 47 bacterial phyla. The most common predicted host phyla were *Chloroflexota* (n = 156), followed by *Halobacteriota* (n = 86), *Pseudomonadota* (n = 85), *Planctomycetota* (n = 56), and *Asgardarchaeota* (n = 52). Most of the identified connections involved viruses in the *Caudoviricetes* class infecting the *Chloroflexota* (731 pairs), and archaeal *Asgardarchaeota* (446 pairs) and *Halobacteriota* (284 pairs). The host-virus network revealed that the majority of clusters (60%, n=1,616)

consisted of more than one viral or prokaryotic genome. This suggests that a single

virus could infect multiple prokaryotic populations, or a single prokaryote may be

susceptible to several viral population[51], indicating a complex relationship between

viruses and prokaryotes in cold seeps.

Viruses encode an arsenal of anti-defense proteins that facilitate infection by disabling

various prokaryotic defense mechanisms[18, 24]. A total of 1,197 anti-defense genes were

detected from 11% viral genomes (n = 1,065), predominantly comprising anti-

CRISPR-Cas, anti-Thoeris, anti-Gabija, and anti-RM genes (**Fig. 6b and**

Supplementary Data 14). While most anti-defense genes were detected across a

broad range of viruses, some were found only in specific viral genomes. For example,

anti-CRISPR, anti-RM, anti-Thoeris, and anti-Gabija genes were commonly found,

whereas anti-RecBCD genes and broad-spectrum anti-defense genes (*orf148*[52], a

broad-spectrum anti-defense gene in *Escherichia* phage OSYSP that targets conserved

host barrier defenses and metabolic pathways to weaken the host's multi-layered

antiviral mechanisms) were exclusively detected in certain *Caudoviricetes* viruses.

From the 2,688 host-virus linkages, we identified 155 pairs in which host microbes

contained defense genes while the corresponding viruses possessed anti-defense genes

(Fig. 6c and Supplementary Data 13). These connections were predominantly

observed between viruses in the *Caudoviricetes* class and their hosts, including

archaeal groups such as *Asgardarchaeota* (38 pairs) and *Halobacteriota* (26 pairs), as

well as various bacterial taxa including *Pseudomonadota* (25 pairs) and

Desulfobacterota (24 pairs). Notably, a single virus employing two anti-defense

systems could infect two ETH-SRB1 genomes belonging to the *Desulfobacterota*

phylum, and one *Asgardarchaeota* genome was found to harbor six defense systems

to fend off infection by five viruses **(Fig. 6c and Supplementary Data 13)**. These

findings suggest that the distribution of defense and anti-defense genes appear lineage

dependent, particularly among various archaeal lineages and their viruses[10, 25, 53].

A single virus may encode multiple anti-defense genes, enabling it to counter various

host defense strategies or adapt to different defenses employed by the same host throughout its life cycle[25]. We identified 118 viruses that encode two or more types of anti-defense systems **(Supplementary Data 13)**. For instance, a viral genome from the *Caudoviricetes* family contains AcrIF17, AcrIF24, and Tad1 genes to counteract the CRISPR-Cas and Thoeris defense systems of its host belonging to the phylum *Pseudomonadota* **(Fig. 6c)**. Additionally, two viral genomes from the same family harbor DarB and Tad2 genes to counteract the RM and Thoeris defense systems of their host, ETH-SRB1 **(Fig. 6d and Supplementary Data 13 in green)**. The predicted dimeric structure of AcrIF24 from *Caudoviricetes* exhibits high similarity (TM-score = 0.86) to the anti-CRISPR protein form *Pseudomonas aeruginosa* **(Fig. 6e)**, which has been experimentally confirmed to inhibit Cas7f[54]. AcrIF24 interacts with immune system components through steric or allosteric effects in a stoichiometric, non-enzymatic manner[54]. Many other anti-defense genes that function similarly were also detected in cold seeps, including most Acrs (e.g., AcrIIA7, AcrIIA2, AcrIIA4, AcrIIIB1, and AcrIIC1). In contrast, enzymatic inhibitors, which effectively suppress host defenses by permanently neutralizing microbial immunity[24, 55, 56], were also identified, such as Acb1, NTases, AcrIII-1, and AcrVA1-5. Additionally, several anti-defense systems with unknown molecular mechanisms, such as Anti-Dnd, Gad2, Abc1, and KlcA, were also present. Overall, viral anti-defense systems engage in complex evolutionary interactions with host defense systems, employing enzymatic, non-enzymatic, and unknown molecular mechanisms in cold seep ecosystems.

Defense systems protect key metabolic microbial groups from viral infections

Anaerobic methane oxidation, sulfate reduction, and nitrogen fixation, associated with the activities of the *mcrA*, *dsrA*, and *nifH* genes, respectively, are key metabolic processes in cold seeps[57-59]. The significant correlation between the abundance of defense systems and these key metabolic genes **(Supplementary Fig. 16)** underscores the role of defense systems in protecting functional microbes from viral and MGE

infections. To further explore this relationship, we analyzed 35 anaerobic methanotrophic archaea (ANME), 47 sulfate-reducing bacteria (SRB), and 57 diazotrophs identified from 3,813 species-level MAGs. The total number of systems and the diversity of families varied significantly across individual microbial groups, particularly among SRB and diazotrophs **(Fig. 7b)**. While the overall repertoire of defense systems remain relatively conserved across major microbial groups **(Fig. 7a, Supplementary Fig. 17, and Supplementary Data 15)**, their abundance and diversity differ markedly among groups, underscoring the complexity of defense system distribution across different microbial groups. For example, ANME generally encode fewer defense systems, with a limited diversity of types. Within SRB, ETH-SRB1 encode on average 17.6 defense systems across 7.3 types, whereas another SRB strain in *Gemmatimonadota* order (SY6_cobin_101) encodes only 3 systems of just 1 type (belonging to AbiE; **Fig. 7b and Supplementary Data 15**).

In detail, SRB predominantly encode type IIG and type II RM systems, followed by type I RM systems. ANME and diazotrophs primarily encode type IIG and type I systems, with type II RM systems following (**Supplementary Fig. 18 and Supplementary Data 15**). The number of these systems in ANME is obvious smaller compared to the other two functional groups. RM systems primarily function in the early stages of viral infection to eliminate invading nucleic acids[11], indicating that cold seep prokaryotes possess the ability to respond rapidly and clear viruses promptly. CRISPR-Cas systems are categorized into two classes and six types based on the set of Cas genes they encode and whether their effector is a single protein or a protein complex[13, 60, 61]. Both classes of CRISPR-Cas were present in cold seep microbes, with class I being more abundant in both number and diversity (**Supplementary Fig. 19 and Supplementary Data 15**). Most archaea, including ANME, primarily encode type I CRISPR-Cas systems, while the majority of bacteria, including SRB and diazotrophs, mainly encode both type I and type III CRISPR-Cas systems. CRISPR-Cas has been shown to contribute to shaping a highly dynamic network of interactions between viruses/MGEs and their prokaryotic hosts in various ecosystems[62-64].

Additionally, SRB tend to harbor two or more Gabija systems, particularly in ETH-
SRB1 strains, with some encoding as many as four Gabija systems (**Supplementary**
Fig. 20 and Supplementary Data 15). Overall, these findings highlight the diversity
and complexity of defense systems across various microbial groups, likely shaped by
distinct evolutionary pressures and ecological niches[19].

Gabija is effective in providing bacterial immunity against certain bacteriophages, as
shown by its strong defense against phi29, rho14, phi105, and SpBeta in *Bacillus*
cereus VD045 [7]. Through sequence alignment and both sequence- and structure-based
phylogenetic analyses (**Supplementary Fig. 21**), we found that Gabija is widely
present in cold seep microbes, spanning 45 phyla (including 9 archaeal and 36
bacterial phyla) and encompassing the three key metabolic microbial groups. We
discovered that one ETH-SRB1 strain encodes two Gabija systems, and the virus
infecting it carries the *gad1* anti-defense gene (**Fig. 7c**). This ETH-SRB1 strain
harbored both a complete and an incomplete Gabija system. In the complete system,
the GajA protein contains an N-terminal ATPase domain and a C-terminal Toprim
(topoisomerase-primase) domain, while the GajB protein includes an AAA domain
and a UvrD-like helicase[30, 65]. In contrast, the GajB of the incomplete Gabija system
lacks the UvrD-like helicase (**Supplementary Fig. 20**). The predicted structure shows
similarity (TM-score = 0.70) to the Gabija protein from *Bacillus cereus* VD045[30] (**Fig.**
7d), suggesting potential conservation of functional domains. The viral protein Gad1
(**Fig. 7e**), larger than other phage immune evasion proteins (~35 kDa), appears to bind
and encapsulate the GajAB complex, potentially blocking the degradation of target
DNA. The binding of Gad1 to the GajAB complex encoded by a virus targeting ETH-
SRB1 suggests that Gad1 may inhibit the Gabija system in this strain.

Considering the differences in the number and types of defense systems among
functional microbial groups, we hypothesize that the specific pairs of co-occurring
and negatively associated defense systems vary across these groups. As anticipated,
defense   systems   that   co-occurred   in   one   group   were   absent   in   another

(Supplementary Fig. 22 and Supplementary Data 16). For example, AbiE system

in SRB exhibited 28 co-occurrences, including with CBASS, DarTG, Gabija, and

Mokosh, none of which were observed in ANME or diazotrophs. These findings

highlight the unique defensive strategies employed by different microbial lineages.

Previous research has revealed that the relationships between defense systems likely

depend on environmental and genetic factors that select for a particular anti-virus

immunity strategy, and that synergistic immunity provides an evolutionary advantage

to microbial populations[23]. Overall, our results indicate that cold seep microbial

defense systems are generally mechanistically compatible, allowing microbes to adopt

diverse and flexible strategies for anti-virus defense based on their unique

environmental and genetic contexts.

Implications of microbe-virus relationships in cold seeps

Cold seep ecosystems, characterized by the seepage of hydrocarbon-rich fluids into

deep-sea sediments, harbor a diverse array of previously uncharacterized microbes

and viruses engaged in a continuous evolutionary arms race. In this study, we

identified diverse defense systems in cold seep microbes, including a substantial

proportion of candidate systems. These defense systems frequently co-occur and may

work synergistically to combat infections from MGEs and viruses. Experimental

validation confirmed that several of these candidate defense systems effectively

protect against viral infections. Notably, defense systems such as RM, CRISPR-Cas,

AbiE, and SoFic exhibited high metagenomic abundances, widespread distribution,

and active expression across various microbial genomes from different sediment

depths and geographical locations, underscoring their vital role in microbial survival

within these extreme environments. In response, cold seep viruses have evolved

diverse anti-defense mechanisms, functioning through enzymatic, non-enzymatic, and

as-yet-unknown mechanisms to evade host defenses. The presence of multiple viral

anti-defense genes within single viral genomes, along with various defense genes in

their hosts, highlights the intricate and dynamic interactions between these viruses and

their microbial hosts, reflecting a potential ongoing evolutionary arms race. Functionally critical lineages in cold seeps, such as ANME, SRB, and diazotrophs, adjust the correlations among different defense systems according to their ecological niches and environmental pressures. Overall, these findings emphasize the complexity and flexibility of interactions within microbial defense systems in extreme environments. The ability of cold seep microbes to maintain diverse and adaptable defense mechanisms highlights their resilience and evolutionary success in one of Earth's most challenging habitats. This study deepens our understanding of virus-microbe interactions in cold seep ecosystems and underscores the significance of microbial defense systems in shaping microbial ecology and evolution in extreme environments.

Methods

Dataset and processing

Metagenomic assemblies were generated from 171 deep-sea sediment samples collected from 17 geographically diverse cold seep sites, with sediment depths ranging from 0 to 68.55 mbsf (**Supplementary Fig. 1**). The construction of the genomic inventory followed the methods detailed in our previous study[5]. Raw metagenomic sequence data underwent quality control and were assembled into contigs sequences. Contigs exceeding 1,000 base pairs in length were selected for binning. The resulting MAGs were deduplicated using dRep v3.4.0 [66] (parameters: -comp 50 -con 10) at a 95% average nucleotide identity threshold, yielding a total of 3,813 cold seep prokaryotic MAGs. Initial taxonomic assignment of each MAG was conducted using GTDB-TK v2.1.3 [67] with reference to the Genome Taxonomy Database (GTDB) release R214, followed by validation through a maximum-likelihood phylogenomic tree. For the archaeal MAGs, the phylogenomic tree was inferred based on the concatenation of 53 conserved single-copy genes as marker genes, while for the bacterial MAGs, it was based on 120 marker genes. The trees

were constructed using IQ-TREE v2.2.0.3 (parameters: -m MFP -B 1000)[68] and visualized by using iTOL v6 [69]. Sankey plots were constructed using Pavian (https://fbreitwieser.shinyapps.io/pavian/)[70]. Protein-coding sequences were predicted from the MAGs using Prodigal v2.6.3 (parameter: -p meta)[71], resulting in a gene catalog annotated from all MAGs.

Thirty-three metatranscriptomes, originating from cold seeps of the South China Sea in regions of Jiaolong, Haima, Qiongdongnan Basin, and the Shenhu area[59, 72-74] **(Supplementary Fig. 1)**, were subjected to quality control using the Read_QC module of the metaWRAP v1.3.2 pipeline (parameters: --skip-bmtagger)[75]. Following this, SortMeRNA v2.1 was utilized to remove ribosomal RNA[76]. The resulting clean reads were then aligned to the gene catalog annotated from all MAGs, employing Salmon v.1.9.0 (parameters: -validateMappings -meta)[77], to quantify the transcript abundance of prokaryotic defense genes. Transcript levels were normalized to transcripts per million (TPM).

Identification of defense genes, systems, and islands

Defense genes and systems were identified using DefenseFinder v1.2.1 (parameters: --db-type gembase)[19] and PADLOC v2.0.0 (parameter: --fix-prodigal)[33]. DefenseFinder detects 710 genes related to 154 defense system families by matching sequences to 845 Hidden Markov Models (HMMs) representing experimentally validated antiviral proteins. It classifies defense systems based on predefined genetic architecture rules, which require mandatory proteins to be genomically adjacent and exclude prohibited components. PADLOC detects 385 defense subsystems using HMM libraries and classification files, enforcing gene proximity via the max_separation parameter and validating associations through co-occurrence frequency statistics (one-sample proportion test, $p < 0.001$). Candidate defense systems were annotated according to the PADLOC database, which includes HECs (Hma-Embedded Candidates) and PDCs (Phage Defense Candidates)[33]. These

candidates are predicted using a "guilt-by-embedding" strategy, where uncharacterized genes are recurrently embedded within known defense systems (e.g., Hma, RM, BREX, DISARM)[34]. The classification of proteins within a single defense system was determined based on parameters established by DefenseFinder and PADLOC, including maximum intergenic distances, domain architecture consistency, and functional annotation rules. Results from the two tools were merged, with duplicate results compared based on e-values to retain those with higher confidence. Defense islands were defined as arrays of defense genes (or defense systems) separated by no more than ten genes and containing at least five genes from at least three different defense systems[9].

Gene abundances of gene catalog annotated from all MAGs were quantified using Salmon v.1.9.0 in mapping-based mode (parameters: --meta --validateMappings)[77] and read counts were normalized to GPM (genes per million). The abundance of a defense system is represented by the sum of the abundances of the genes within that system divided by the average number of genes in the system.

Identification of viruses and virus-host linkages

Potential viral sequences were identified from metagenomic assemblies (contig lengths greater than 10 kb) derived from the aforementioned 171 cold seep site samples and an additional 20 cold seep samples. Three methodologies were used: geNomad v1.7.0 (parameters: end-to-end)[48], Virsorter2 v2.2.3 (parameters: default)[47], and VIBRANT v1.2.1 (parameters: default)[46]. The completeness and contamination of these sequences were assessed using CheckV v1.0.1 with the database v1.5 [78]. Sequences with an estimated completeness of 50% or more were selected for clustering into species-level viral operational taxonomic units (vOTUs). Clustering parameters followed CheckV guidelines: 95% average nucleotide identity and a minimum aligned fraction of 85%. This process resulted in the identification of 9,971 species-level vOTUs. Taxonomic assignment of these vOTUs was conducted using

geNomad v1.7.0. The resulting taxonomic lineages adhere to the classification outlined in the International Committee on Taxonomy of Viruses (ICTV) Virus Metadata Resource (VMR) number 19. Open reading frames (ORFs) within each vOTU were predicted using Prodigal v2.6.3 (parameter: -p meta)[71]. Functional annotation of the viral sequences was performed utilizing the virome model in DRAM v1.4.0 [79].

Linkages between viruses and their hosts were primarily determined using the iPHoP v1.3.2 pipeline[50], utilizing a database constructed from 3,813 prokaryotic MAGs identified in this study and the iPHoP_db_Sept21_rw. iPHoP employs a variety of methods to predict virus-host linkages, including Blastn v2.12.0 [80] alignment of viral genomes against the iPHoP_db_Sept21_rw database and its CRISPR spacer database; The k-mer composition similarity between viral and host genomes was analyzed using four methods: RaFAH v0.3 [81], WIsH v1.0 [82], VirHostMatcher-Net[83], and PHP[84]. In addition to the iPHop methods, for contigs identified with CRISPR-Cas systems, CRISPRCasFinder v4.2.21 [85] was used to identify CRISPR arrays flanking the Cas genes. Local alignments of extracted spacers (with lengths greater than 25 base pairs) against viral genomes were performed using Blastn. Only BLAST matches with 100% alignment coverage and at most two mismatches were considered high-confidence matches. The network graph between hosts and viruses was visualized using Gephi v0.9.2 [86]. The workflow for this part was illustrated using Draw.io v23.1.5 (https://app.diagrams.net/).

Identification of anti-defense genes

The anti-defense genes of viruses were primarily identified through two approaches. (1) APIS database search. We utilized the Anti-Prokaryotic Immune System (APIS) database[87], which contains experimentally validated viral proteins that counteract prokaryotic immune systems and their associated protein families. All viral proteins from our dataset were compared against the APIS database using DIAMOND blastp

v2.0.8 (parameters: -e 1e-10 --id 30)[88] and HMMER v3.3.2 (parameters: -E 1e-10). (2)

Reference sequence comparison. We compiled sequences of viral anti-defense

proteins as documented by Mayo-Muñoz et al. in their review[24]. These 36 reference

sequences were downloaded from the NCBI database and used to construct a custom

database using the DIAMOND makedb command. Our viral proteins were then

compared against this database using DIAMOND BLASTp v2.0.8 (parameters: -e 1e-

10 --id 30). The results were obtained using two methods, with the more reliable

results selected based on e-values to derive the final set of anti-defense genes.

Experimental validation of candidate defense systems

Coding sequences of the candidate defense systems were commercially synthesized

and cloned into the pQE82L expression vector. The recombinant vectors were

transformed into *E. coli* B (ATCC® 11303™). Cells transformed with the empty

vector served as controls. Plaque assays were conducted following established

protocols[14, 89]. A single colony from a fresh LB agar plate was inoculated into LB

broth containing ampicillin (100 µg/mL) and grown at 37°C to an OD_{600} of

approximately 0.6. Protein expression was induced by adding 0.2 mM isopropyl β-D-

1-thiogalactopyranoside (IPTG). After an additional hour of incubation, 500 µL of the

bacterial culture was mixed with 14.5 mL of 0.5% LB top agar, and the mixture was

poured onto LB plates containing ampicillin (100 µg/mL) and IPTG (0.1 mM). The

plates were then spotted with 4 µL aliquots of phage dilutions prepared in LB medium.

Eight 10-fold serial dilutions were used for each phage: dilutions 10^{-1} to 10^{-8} for

phage T7, and 10^{0} to 10^{-7} for phages T4 and T5. Plates were incubated overnight at

25°C before imaging. All the assays are repeated at least 3 times with similar results.

Functional annotations

622   Functional annotation of MAGs was performed by searching against the KEGG, Pfam,

623   MEROPS, and dbCAN databases using DRAM v1.4.0 (parameters: --min_contig_size

624   1000)[79]. Gene context was visualized with Chiplot (https://www.chiplot.online/).

Plasmid and provirus sequences were identified from prokaryotic MAGs using geNomad v1.7.0 [48]. To classify different types of MGEs, such as IS elements and transposons (IS_Tn), integrons, and conjugative elements (CE), recombinase marker genes, were identified through alignment with the proMGE database[90] using HMMER v3.3.2. For the identification of key metabolic genes in cold seep prokaryotes, *dsrA* genes were detected using DiSco v1.0.0 [91], and *mcrA* genes were identified using METABOLIC v4.0 [92] and DRAM v1.4.0 [79]. Nitrogen fixation marker genes *nifH* genes were detected using three methods: (1) protein sequences in the MAGs were compared against NCycDB[93] using DIAMOND blastp v2.0.8 [88], (2) potential *nifH* sequences were searched based on *nifH* reference sequences (n = 1,271) from the Greening lab metabolic marker gene database using DIAMOND blastp v2.0.8 (parameters: --id 65)[88], and (3) *nifH* genes were extracted using HMMER v3.3.2 with the TIGRFAM model TIGR01287.1.

Protein sequence and structural analysis

To verify the accuracy and phylogenetic placement of the identified defense and anti-defense genes, multiple sequence alignments were performed using MAFFT v7.505 [94]. Alignments were trimmed using TrimAl v1.4.1 under default settings[95]. Maximum likelihood phylogenetic trees were constructed using IQ-TREE v2.2.0.3 (parameters: -m MFP -B 1000)[68]. Trees were visualized and annotated using iTOL v6 [69]. Protein domains were identified using the Batch Protein Annotation plugin in Tbtools-II v2.042 [96], which integrates results from InterProScan[97]. Sequence conservation among proteins was aligned and visualized using Jalview v2.11.3.3 [98]. Structural predictions for proteins, including GajA and GajB, were performed using ColabFold v1.5.5 which leverages AlphaFold2[99-101] on September 14, 2023. Structures of the AcrIF24 dimer, GajAB octamer, and Gad1 protein were predicted using AlphaFold3[102] on August 30, 2024. The predicted structures were evaluated based on pLDDT and Predicted Aligned Error (PAE) scores **(Supplementary Fig. 23)**. The reference structures AcrIF24 dimer and GajAB octamer were obtained from experimentally determined

653 Csy-AcrIF24 complex (PDB code: 7ELM) and GajAB-Gad1 complex (PDB code:

654 8U7I). Homologous and structurally similar proteins were identified using the

655 Foldseek web interface[103]. Protein structures were visualized and aligned using

656 PyMOL v2.5.5 [104].

Statistical analyses

Statistical analyses were conducted using R v4.2.3. Normality of the data was

assessed using the Kolmogorov-Smirnov test, and homogeneity of variances was

tested using Levene's test. The Wilcoxon rank-sum test was employed to compare

characteristics of defense systems between archaeal and bacterial groups. For

comparisons involving more than two groups—such as defense system and gene

densities among different microbial phyla, or differences in gene and transcript

abundances among various types of defense systems or genes—the Kruskal-Wallis

test was used. Pearson correlation analysis was conducted using R's default method

(cor.test function with Pearson method) to evaluate relationships between defense

systems and their subsystems, based on abundance profiles. All correlation analyses

were performed on $\log_{10}$-transformed abundance data to meet normality assumptions.

For multiple comparisons, p-values were adjusted using the Benjamini-Hochberg

false discovery rate (FDR) correction with a significance threshold of 0.05.

Data availability

The non-redundant MAGs catalog has been deposited in the NCBI database under

BioProject PRJNA950938. Sequences of defense systems, viral genomes, viral anti-

defense systems, and phylogenetic trees of GajAB based on amino acid sequences and

protein structures, generated in this study, are available at

https://doi.org/10.6084/m9.figshare.27004438. All additional data supporting the

findings of this study are provided within the article and its Supplementary

Information Files. Source data are provided with this paper.

Code availability

The present study did not generate codes, and mentioned tools used for the data
analysis were applied with default parameters unless specified otherwise.

References

1.   Dong, X. et al. Metabolic potential of uncultured bacteria and archaea associated with
petroleum seepage in deep-sea sediments. *Nature Communications* **10**, 1816 (2019).
2.   Joye, S.B. The geology and biogeochemistry of hydrocarbon seeps. *Annual Review of Earth*
and Planetary Sciences **48**, 205-231 (2020).
3.   Dong, X. et al. Thermogenic hydrocarbon biodegradation by diverse depth-stratified microbial
populations at a Scotian Basin cold seep. *Nature Communications* **11**, 5825 (2020).
4.   Kleindienst, S. et al. Diverse sulfate-reducing bacteria of the *Desulfosarcina/Desulfococcus*
clade are the key alkane degraders at marine seeps. *The ISME Journal* **8**, 2029-2044 (2014).
5.   Yang, X.Y., Shen, Z., Lin, Q. & Fu, T.M. Molecular basis of Gabija anti-phage supramolecular
assemblies. *Nature* **15**, 836 (2023).
6.   Li, Z. et al. Deep sea sediments associated with cold seeps are a subsurface reservoir of viral
diversity. *The ISME Journal* **15**, 2366-2378 (2021).
7.   Doron, S. et al. Systematic discovery of antiphage defense systems in the microbial
pangenome. *Science* **359**, eaar4120 (2018).
8.   Millman, A. et al. An expanded arsenal of immune systems that protect bacteria from phages.
Cell Host & Microbe **30**, 1556-1569 (2022).
9.   Beavogui, A. et al. The defensome of complex bacterial communities. *Nature*
Communications **15**, 2146 (2024).
10.   Peng, Y. et al. Viruses in deep-sea cold seep sediments harbor diverse survival mechanisms
and remain genetically conserved within species. *The ISME Journal* **17**, 1774-1784 (2023).
11.   Georjon, H. & Bernheim, A. The highly diverse antiphage defence systems of bacteria. *Nature*
Reviews Microbiology **21**, 686-700 (2023).
12.   Tock, M.R. & Dryden, D.T.F. The biology of restriction and anti-restriction. *Current Opinion*
in Microbiology **8**, 466-472 (2005).
13.   Hille, F. et al. The biology of CRISPR-Cas: backward and forward. *Cell* **172**, 1239-1259
(2018).
14.   Cheng, R. et al. Prokaryotic Gabija complex senses and executes nucleotide depletion and
DNA cleavage for antiviral defense. *Cell Host & Microbe* **31**, 1331-1344 (2023).
15.   Rousset, F. et al. A conserved family of immune effectors cleaves cellular ATP upon viral
infection. *Cell* **186**, 3619-3631 (2023).
16.   Lopatina, A., Tal, N. & Sorek, R. Abortive infection: bacterial suicide as an antiviral immune
strategy. *Annual Review of Virology* **7**, 371-384 (2020).
17.   Saenz, J.S., Seifert, J. & Rios-Galicia, B. Antiviral defense systems in the rumen microbiome.
mSystems **10**, e01521-24 (2025).
18.   Piel, D. et al. Phage–host coevolution in natural populations. *Nature Microbiology* **7**, 1075-

1086 (2022).

19.   Tesson, F. et al. Systematic and quantitative view of the antiviral arsenal of prokaryotes.
Nature Communications **13**, 2561 (2022).

20.   Bernheim, A. & Sorek, R. The pan-immune system of bacteria: antiviral defence as a
community resource. *Nature Reviews Microbiology* **18**, 113-119 (2020).

21.   Rocha, E.P.C. & Bikard, D. Microbial defenses against mobile genetic elements and viruses:
Who defends whom from what? *PLoS Biology* **20**, e3001514 (2022).

22.   Picton, D.M. et al. The phage defence island of a multidrug resistant plasmid uses both BREX
and type IV restriction for complementary protection from viruses. *Nucleic Acids Research* **49**,
11257-11273 (2021).

23.   Wu, Y. et al. Bacterial defense systems exhibit synergistic anti-phage activity. *Cell Host &*
Microbe **32**, 557-572.e556 (2024).

24.   Mayo-Muñoz, D., Pinilla-Redondo, R., Camara-Wilpert, S., Birkholz, N. & Fineran, P.C.
Inhibitors of bacterial immune systems: discovery, mechanisms and applications. *Nature*
Reviews Genetics **25**, 237-254 (2024).

25.   Hampton, H.G., Watson, B.N.J. & Fineran, P.C. The arms race between bacteria and their
phage foes. *Nature* **577**, 327-336 (2020).

26.   Duan, N., Hand, E., Pheko, M., Sharma, S. & Emiola, A. Structure-guided discovery of anti-
CRISPR and anti-phage defense proteins. *Nature Communications* **15**, 649 (2024).

27.   Gussow, A.B. et al. Machine-learning approach expands the repertoire of anti-CRISPR protein
families. *Nature Communications* **11**, 3784 (2020).

28.   Serfiotis-Mitsa, D. et al. The structure of the KlcA and ArdB proteins reveals a novel fold and
antirestriction activity against Type I DNA restriction systems in vivo but not in vitro. *Nucleic*
Acids Research **38**, 1723-1737 (2010).

29.   Hobbs, S.J. et al. Phage anti-CBASS and anti-Pycsar nucleases subvert bacterial immunity.
Nature **605**, 522-526 (2022).

30.   Antine, S.P. et al. Structural basis of Gabija anti-phage defence and viral immune evasion.
Nature **625**, 360-365 (2024).

31.   An, L. et al. Global diversity and ecological functions of viruses inhabiting oil reservoirs.
Nature Communications **15**, 6789 (2024).

32.   Zhong, Z.-P. et al. Viral potential to modulate microbial methane metabolism varies by habitat.
Nature Communications **15**, 1857 (2024).

33.   Payne, L.J. et al. Identification and classification of antiviral defence systems in bacteria and
archaea with PADLOC reveals new system types. *Nucleic Acids Research* **49**, 10868-10878
(2021).

34.   Payne, L.J., Hughes, T.C.D., Fineran, P.C. & Jackson, S.A. New antiviral defences are
genetically embedded within prokaryotic immune systems. *bioRxiv*,
https://doi.org/10.1101/2024.01.29.577857 (2024).

35.   Mayo-Munoz, D., Pinilla-Redondo, R., Birkholz, N. & Fineran, P.C. A host of armor:
Prokaryotic immune strategies against mobile genetic elements. *Cell Reports* **42**, 112672
(2023).

36.   Tesson, F. et al. A comprehensive resource for exploring antiphage defense: DefenseFinder
webservice, wiki and databases. *Nucleic Acids Research* **53**, 13 (2025).

37.   Leão, P. et al. Asgard archaea defense systems and their roles in the origin of eukaryotic

immunity. *Nature Communications* **15**, 6386 (2024).

38.   Liu, Y., Botelho, J. & Iranzo, J. Timescales and genetic linkage explain the variable impact of defense systems on horizontal gene transfer *Genome Research* **35**, 268-278 (2025).

39.   Kogay, R., Wolf, Y.I. & Koonin, E.V. Defence systems and horizontal gene transfer in bacteria. *Environmental Microbiology* **26** (2024).

40.   Li, M. et al. Toxin-antitoxin RNA pairs safeguard CRISPR-Cas systems. *Science* **372**, eabe5601 (2021).

41.   Dupuis, M.È., Villion, M., Magadán, A.H. & Moineau, S. CRISPR-Cas and restriction–modification systems are compatible and increase phage resistance. *Nature Communications* **4**, 2087 (2013).

42.   Wu, Y. et al. Bacterial defense systems exhibit synergistic anti-phage activity. *Cell Host & Microbe* **32**, 557-572 (2024).

43.   Maguin, P., Varble, A., Modell, J.W. & Marraffini, L.A. Cleavage of viral DNA by restriction endonucleases stimulates the type II CRISPR-Cas immune response. *Molecular Cell* **82**, 907-919.e907 (2022).

44.   Dong, X. et al. Evolutionary ecology of microbial populations inhabiting deep sea sediments associated with cold seeps. *Nature Communications* **14**, 1127 (2023).

45.   Dong, X. et al. A vast repertoire of secondary metabolites potentially influences community dynamics and biogeochemical processes in cold seeps. *Science Advances* **10**, eadl2281 (2024).

46.   Kieft, K., Zhou, Z. & Anantharaman, K. VIBRANT: automated recovery, annotation and curation of microbial viruses, and evaluation of viral community function from genomic sequences. *Microbiome* **8**, 1-23 (2020).

47.   Guo, J. et al. VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses. *Microbiome* **9**, 1-13 (2021).

48.   Camargo, A.P. et al. Identification of mobile genetic elements with geNomad. *Nature Biotechnology* **42**, 1303–1312 (2023).

49.   Roux, S. et al. Minimum Information about an Uncultivated Virus Genome (MIUViG). *Nature Biotechnology* **37**, 29-37 (2019).

50.   Roux, S. et al. iPHoP: An integrated machine learning framework to maximize host prediction for metagenome-derived viruses of archaea and bacteria. *PLoS Biology* **21**, e3002083 (2023).

51.   Zhou, K. et al. Genomic and transcriptomic insights into complex virus-prokaryote interactions in marine biofilms. *The ISME Journal* **17**, 2303–2312 (2023).

52.   Silas, S. et al. Activation of bacterial programmed cell death by phage inhibitors of host immunity. *Molecular Cell* **85**, 1838-1851.e10 (2025).

53.   Yirmiya, E. et al. Phages overcome bacterial immunity via diverse anti-defence proteins. *Nature* **625**, 352-359 (2024).

54.   Yang, L. et al. Insights into the inhibition of type I-F CRISPR-Cas system by a multifunctional anti-CRISPR protein AcrIF24. *Nature Communications* **13**, 1931 (2022).

55.   Suresh, S.K., Murugan, K. & Sashital, D.G. Enzymatic anti-CRISPRs improve the bacteriophage arsenal. *Nature Structural & Molecular Biology* **26**, 250-251 (2019).

56.   Knott, G.J. et al. Broad-spectrum enzymatic inhibition of CRISPR-Cas12a. *Nature Structural & Molecular Biology* **26**, 315-321 (2019).

57.   Müller, A.L., Kjeldsen, K.U., Rattei, T., Pester, M. & Loy, A. Phylogenetic and environmental diversity of DsrAB-type dissimilatory (bi)sulfite reductases. *The ISME Journal* **9**, 1152-1165

(2015).

58.   Gu, Y. et al. Bacterial Shedu immune nucleases share a common enzymatic core regulated by
diverse sensor domains. *Molecular Cell* **85**, 523 - 536 (2024).

59.   Dong, X. et al. Phylogenetically and catabolically diverse diazotrophs reside in deep-sea cold
seep sediments. *Nature Communications* **13**, 4885 (2022).

60.   Sheng, Y. et al. Insertion sequence transposition inactivates CRISPR-Cas immunity. *Nature*
Communications **14**, 4366 (2023).

61.   Koonin, E.V., Makarova, K.S. & Zhang, F. Diversity, classification and evolution of CRISPR-
Cas systems. *Current Opinion in Microbiology* **37**, 67-78 (2017).

62.   Martínez Arbas, S. et al. Roles of bacteriophages, plasmids and CRISPR immunity in
microbial community dynamics revealed using time-series integrated meta-omics. *Nature*
Microbiology **6**, 123-135 (2021).

63.   Sun, C.L., Thomas, B.C., Barrangou, R. & Banfield, J.F. Metagenomic reconstructions of
bacterial CRISPR loci constrain population histories. *The ISME Journal* **10**, 858-870 (2016).

64.   Guerrero, L.D. et al. Long-run bacteria-phage coexistence dynamics under natural habitat
conditions in an environmental biotechnology system. *The ISME Journal* **15**, 636-648 (2021).

65.   Huo, Y. et al. Structural and biochemical insights into the mechanism of the Gabija bacterial
immunity system. *Nature Communications* **15**, 836 (2024).

66.   Olm, M.R., Brown, C.T., Brooks, B. & Banfield, J.F. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes through de-
replication. *The ISME journal* **11**, 2864-2868 (2017).

67.   Parks, D.H. et al. GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic*
Acids Research **50**, D785-D794 (2022).

68.   Nguyen, L.T., Schmidt, H.A., Von Haeseler, A. & Minh, B.Q. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and*
Evolution **32**, 268-274 (2015).

69.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic Acids Research* **49**, W293-W296 (2021).

70.   Breitwieser, F.P. & Salzberg, S.L. Pavian: interactive analysis of metagenomics data for
microbiome studies and pathogen identification. *Bioinformatics* **36**, 1303-1304 (2020).

71.   Hyatt, D. et al. Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 1-11 (2010).

72.   Zhang, C. et al. The majority of microorganisms in gas hydrate-bearing subseafloor sediments
ferment macromolecules. *Microbiome* **11**, 37 (2023).

73.   Xiao, X. et al. Metal-driven anaerobic oxidation of methane as an important methane sink in
methanic cold seep sediments. *Microbiology Spectrum* **11**, e05337-05322 (2023).

74.   Li, W.L. et al. Microbial ecology of sulfur cycling near the sulfate–methane transition of
deep□sea cold seep sediments. *Environmental Microbiology* **23**, 6844-6858 (2021).

75.   Uritskiy, G.V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-
resolved metagenomic data analysis. *Microbiome* **6**, 1-13 (2018).

76.   Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211-3217 (2012).

77.   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast and

850   bias-aware quantification of transcript expression. *Nature Methods* **14**, 417-419 (2017).

851 78. Nayfach, S. et al. CheckV assesses the quality and completeness of metagenome-assembled
852   viral genomes. *Nature Biotechnology* **39**, 578-585 (2021).

853 79. Shaffer, M. et al. DRAM for distilling microbial metabolism to automate the curation of
854   microbiome function. *Nucleic Acids Research* **48**, 8883-8900 (2020).

855 80. Ye, J., McGinnis, S. & Madden, T.L. BLAST: improvements for better sequence analysis.
856   *Nucleic Acids Research* **34**, W6-W9 (2006).

857 81. Coutinho, F.H. et al. RaFAH: Host prediction for viruses of Bacteria and Archaea based on
858   protein content. *Patterns* **2**, 100274 (2021).

859 82. Galiez, C., Siebert, M., Enault, F., Vincent, J. & Söding, J. WIsH: who is the host? Predicting
860   prokaryotic hosts from metagenomic phage contigs. *Bioinformatics* **33**, 3113-3114 (2017).

861 83. Wang, W. et al. A network-based integrated framework for predicting virus–prokaryote
862   interactions. *NAR Genomics and Bioinformatics* **2**, lqaa044 (2020).

863 84. Lu, C. et al. Prokaryotic virus host predictor: a Gaussian model for host prediction of
864   prokaryotic viruses in metagenomics. *BMC Biology* **19**, 5 (2021).

865 85. Couvin, D. et al. CRISPRCasFinder, an update of CRISRFinder, includes a portable version,
866   enhanced performance and integrates search for Cas proteins. *Nucleic Acids Research* **46**,
867   W246-W251 (2018).

868 86. Grandjean, M. Gephi: Introduction to network analysis and visualisation. *International AAAI*
869   *Conference on Weblogs and Social Media* (2015).

870 87. Yan, Y., Zheng, J., Zhang, X. & Yin, Y. dbAPIS: a database of anti-prokaryotic immune
871   system genes. *Nucleic Acids Research* **52**, 419-425 (2023).

872 88. Buchfink, B., Reuter, K. & Drost, H.G. Sensitive protein alignments at tree-of-life scale using
873   DIAMOND. *Nature Methods* **18**, 366-368 (2021).

874 89. Mazzocco, A., Waddell, T.E., Lingohr, E. & Johnson, R.P. Enumeration of bacteriophages
875   using the small drop plaque assay system. *Bacteriophages: methods and protocols, volume 1:*
876   *isolation, characterization, and interactions*, 81-85 (2009).

877 90. Khedkar, S. et al. Landscape of mobile genetic elements and their antibiotic resistance cargo in
878   prokaryotic genomes. *Nucleic Acids Research* **50**, 3155-3168 (2022).

879 91. Neukirchen, S. & Sousa, F.L. DiSCo: a sequence-based type-specific predictor of Dsr-
880   dependent dissimilatory sulphur metabolism in microbial data. *Microbial Genomics* **7**, 000603
881   (2021).

882 92. Zhou, Z. et al. METABOLIC: high-throughput profiling of microbial genomes for functional
883   traits, metabolism, biogeochemistry, and community-scale functional networks. *Microbiome*
884   **10**, 33 (2022).

885 93. Tu, Q., Lin, L., Cheng, L., Deng, Y. & He, Z. NCycDB: a curated integrative database for fast
886   and accurate metagenomic profiling of nitrogen cycling genes. *Bioinformatics* **35**, 1040-1048
887   (2019).

888 94. Nakamura, T., Yamada, K.D., Tomii, K. & Katoh, K. Parallelization of MAFFT for large-scale
889   multiple sequence alignments. *Bioinformatics* **34**, 2490-2492 (2018).

890 95. Capella-Gutiérrez, S., Silla-Martínez, J.M. & Gabaldón, T. trimAl: a tool for automated
891   alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973
892   (2009).

893 96. Chen, C. et al. TBtools-II: A "one for all, all for one" bioinformatics platform for biological

big-data mining. *Molecular Plant* **16**, 1733-1742 (2023).

97.   Paysan-Lafosse, T. et al. InterPro in 2022. *Nucleic Acids Research* **51**, D418-D427 (2023).

98.   Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M. & Barton, G.J. Jalview Version
2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189-
1191 (2009).

99.   Mirdita, M., Steinegger, M. & Söding, J. MMseqs2 desktop and local web server app for fast,
interactive sequence searches. *Bioinformatics* **35**, 2856-2858 (2019).

100.   Mirdita, M. et al. ColabFold: making protein folding accessible to all. *Nature Methods* **19**,
679-682 (2022).

101.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *nature* **596**, 583-
589 (2021).

102.   Abramson, J. et al. Accurate structure prediction of biomolecular interactions with AlphaFold
3. *Nature* **630**, 493-500 (2024).

103.   Van Kempen, M. et al. Fast and accurate protein structure search with Foldseek. *Nature
biotechnology* **42**, 243-246 (2024).

104.   DeLano, W.L. Pymol: An open-source molecular graphics tool. *CCP4 Newsl. Protein
Crystallogr* **40**, 82-92 (2002).

Acknowledgements

The work was supported by National Key R&D Program of China (No.

2024YFC2816200 to X.D.), National Science Foundation of China (No. 92351304 to

X.D., No. 42376115 to X.D., No. 42406109 to Y.H., No. 32470036 to R.C. and No.

32100025 to R.C.), Natural Science Foundation Project of Xiamen City (No.

3502Z202373076 to X.D.), Natural Science Foundation of Fujian Province (No.

2023J06042 to X.D.), Scientific Research Foundation of Third Institute of

Oceanography, MNR (No. 2022025 to X.D., No. 2023022 to X.D. and No. 2025013

to Y.H.), Open Funding Project of State Key Laboratory of Microbial Metabolism (No.

MMLKF23-05 to X.D.) and Fundamental Research Funds for the Central Universities

(No. 2662025SYPY003 to R.C.).

Author contributions

X.D. conceived and designed the study. Y.H. and J.L. performed the omics analyses.

C.L. contributed to viral identification and classification. R.C. designed and

conducted the experimental implementations. X.L. participated in the analysis of
some metagenomic data. C.L., F.X., J.P. and W.X. actively participated in discussions
and data interpretation. F.W. and H.J. provided valuable suggestions for manuscript
revision. Y.H., J.L. and X.D. wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

 Figure Legends

Figure 1. Distribution and activities of defense systems in cold seep prokaryotes.

(a) Types and proportions of verified defense systems and candidate defense systems

in cold seep microbial genomes, including their primary mechanisms of action and

corresponding system counts. "DS" denotes defense systems and "num" represents

the total count. "Type num" refers to the number of distinct defense system types

within each mechanism category. **(b)** Plaque assays demonstrating phage infection of

Escherichia coli B (ATCC® 11303™) transformed with plasmids carrying various

candidate defense systems or an empty vector (control) using a small volume drop

method. The bacteria described above were mixed in the molten agar matrix and ten-

fold serial dilutions of phages T7, T5, and T4 were spotted onto bacterial lawns. All

the assays are repeated at least 3 times with similar results. Predicted protein domains

of the tested systems are shown on the right. Detailed data on annotated defense

systems are provided in **Supplementary Data 1-3**.

Figure 2. Characteristics of defense systems in cold seep prokaryotes. (a)

Proportion of metagenome-assembled genomes (MAGs) containing defense systems

among cold seep prokaryotes, and the proportions of various types of defense systems

within all prokaryotes, archaeal, and bacterial MAGs. **(b)** Distribution of the number

of defense systems (DS) per MAG for all prokaryotes (blue), archaea (orange), and

bacteria (red). **(c)** Number of defense systems (DSs) per MAG for all prokaryotes

(blue, n = 2,466), archaea (orange, n = 395), and bacteria (red, n = 2,071), with black

lines indicating the average number. Detailed statistical data are provided in

Supplementary Data 5. Detailed data on the proportion of genomes carrying defense

systems are provide as a Source Data file.

Figure 3. Abundance and distribution of defense systems in cold seep

prokaryotes. Phylogenetic representation of archaeal **(a)** and bacterial **(b)** MAGs,

categorized and color-coded by their corresponding phyla. The heatmap illustrates the

density of defense systems (DSs per Mb, blue), defense genes (DGs per Mb×10^{-1}, orange), candidate defense systems (CDs per Mb×10^{-1}, green), and the number of defense islands (DIs, red). Circles and triangles denote MAGs with defense systems located on plasmids and proviruses, respectively. The outer bar charts display the genome sizes of MAGs (Mb). Detailed statistical data for this figure are provided in **Supplementary Data 3**.

Figure 4. Co-occurrence patterns among defense systems in cold seep prokaryotes. (a) Correlations among 34 defense systems in cold seep prokaryotes. The lower-left matrix (blue-red gradient) displays correlations among all defense systems across prokaryotic MAGs, while the upper-right matrix (green-pink gradient) shows correlations within defense islands. Pearson correlation coefficients were calculated based on system abundance using two-sided tests. Coefficients are color-coded, and larger squares indicate stronger correlations. Significance levels are denoted by asterisks: * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$. **(b)** Genomic collinearity of eight contiguous fragments containing defense islands. Defense genes are highlighted in green, candidate defense genes in yellow, and associated genes are color-coded: transport-related genes (orange), mobile genetic elements (MGEs; pink), other annotated functional genes (purple), and unknown functions (gray). Detailed statistics for correlation are provided in **Supplementary Data 7 and 9**.

Figure 5. Relative abundance and expression of microbial defense systems in cold seep sediments. (a) Relative abundance (genes per million, GPM, n = 171, $p < 2.2e-16$) and transcript abundance (transcripts per million, TPM, n = 33, $p = 0.522$) of the top ten defense systems across different cold seep sediment samples. Different colors represent different types of defense systems. Statistical differences among defense system types were assessed using a two-sided Kruskal-Wallis rank-sum test. Boxplot: center line, median; box limits, upper and lower quartiles; whiskers, 1.5 × interquartile range; points, individual data values. **(b)** The heatmap displays the average gene abundance of various defense systems within each phylum in cold seep

sediments, expressed in GPM. Only phyla with more than 20 defense systems are shown (n = 51). Gray squares indicate a GPM value of zero. Green labels represent bacterial phyla, while orange denotes archaeal phyla. The bar graph illustrates the average relative abundance of defense systems across phyla ($p < 2.2e\text{-}16$). Data are presented as mean ± SD. **(c)** Gene abundance of defense system genes at different sediment depths in cold seeps, measured in GPM. Sediment depths are categorized into three groups: surface (<1 meter below seafloor [mbsf], n = 143), shallow (1-10 mbsf, n = 14), and deep (>10 mbsf, n = 13). Differences were computed using a two-sided Wilcoxon rank-sum test. Boxplot: center line, median; box limits, upper and lower quartiles; whiskers, 1.5 × interquartile range; points, individual data values. **(d)** Expression levels of defense system genes at different sediment depths in cold seeps, measured in TPM. Sediment depths are categorized into three groups: surface (<1 mbsf, n = 20), shallow (1-10 mbsf, n = 4), and deep (>10 mbsf, n = 9). Differences were computed using a two-sided Wilcoxon rank-sum test. Boxplot: center line, median; box limits, upper and lower quartiles; whiskers, 1.5 × interquartile range; points, individual data values. Source data are provided as a Source Data file.

Figure 6. Interactions between cold seep viral anti-defense systems and host defense systems. (a) The network diagram illustrates 2,688 host-virus pairs identified in cold seeps. Prokaryotes are colored according to their phyla. **(b)** Anti-defense genes identified in cold seep viruses, showing the types of anti-defense systems and corresponding viral taxonomy. **(c)** Illustration of 155 host-virus pairs where host prokaryotes contain defense genes and the corresponding viruses possess anti-defense genes. **(d)** Genomic organization of viral gene clusters containing multiple types of anti-defense systems, along with the corresponding host defense systems. Viral genes are categorized and color-coded: anti-defense genes (yellow), metabolic genes (blue), structural genes (brown), attachment genes (pink), and unknown genes (grey). Host defense genes are highlighted in green, transport-related genes in orange, and other functional genes in purple. Green lines indicate interactions between host defense systems and viral anti-defense systems. **(e)** Structural comparison of the AcrIF24

protein dimer from a *Caudoviricetes* virus infecting *Halomonas* sp. (blue) with the known cryo-electron microscopy structure from *Pseudomonas aeruginosa* (pink, PDB code: 7ELM). A detailed view highlights key sites involved in AcrIF24 protein dimerization and binding to the Csy complex.

Figure 7. Defense competition between key metabolic microbial groups in cold seeps and their viruses. (a) Type of defense systems in key metabolic microbial groups, including anaerobic methanotrophic archaea (ANME), sulfate-reducing bacteria (SRB), and diazotrophs. Colored squares indicate the presence of key functional genes: *mcrA* for ANME, *dsrA* for SRB, and *nifH* for diazotrophs. Colored circles represent the types of defense systems detected in these microbial groups. Different phyla are denoted by distinct colors, and different defense mechanisms are indicated by circles of various colors. To address potential overlap, regions where *mcrA* and *nifH* genes coexist have been highlighted, reflecting functional versatility within the same microenvironments. **(b)** Relationship between the number of defense systems and the diversity of defense types in key metabolic microbial groups, categorized into three groups based on quantity: few systems, diverse systems, and similar systems. **(c)** Genomic organization of the Gabija defense system and the viral *gad1* gene in a virus-host pair involving ETH-SRB1 and a *Caudoviricetes* virus. Genes are color-coded: anti-defense genes (green), structural genes (purple), metabolic genes (pink), attachment genes (orange), transport genes (yellow), defense genes (blue), and genes of unknown function (gray). Green lines indicate interactions between the host defense system and the viral anti-defense system. **(d)** The structure of GajAB complex in cold seep ETH-SRB1 predicted by Alphafold3, including detailed diagrams of the main interaction sites between GajA proteins and between GajB proteins. **(e)** The structure of Gad1 protein in a cold seep *Caudoviricetes* virus predicted by Alphafold3.

a

Verified(37.40%)
9,869

Candidate(62.60%)
16,520

Verified Defense Systems

others(17.60%)
Tiamat(0.90%)
Pycsar(0.93%)
DRT(1.05%)
SEFIR(1.17%)
VSPR(1.18%)
Wadjet(1.28%)
Gabija(1.61%)
Septu(1.85%)
AbiL(1.94%)
CBASS(2.19%)
viperin(2.59%)
dXTPase(3.98%)
AbiE(6.30%)
SoFic(7.24%)
RM(38.28%)
CRISPR-Cas(9.90%)

Candidate Defense Systems

others(19.15%)
HEC-01(1.39%)
HEC-09(1.47%)
PDC-S03(2.45%)
PDC-M05(2.64%)
PDC-S07(2.68%)
PDC-S13(3.37%)
PDC-S02(4.88%)
PDC-S27(5.15%)
PDC-S70(7.26%)
PDC-S01(31.57%)
PDC-S09(9.35%)
PDC-S05(8.63%)

n=9,869   n=106

19%
(n=1,864)   56%
(n=59)

55%
(n=5,416)

7% (n=722)   8% (n=9)
7% (n=7)

19%
(n=1,867)   29%
(n=31)

System num   Type num

Mechanisms

- Nucleic acid degradation
- Inhibition of replication
- Abortive infection
- Unknown

b

Phage T7   Phage T5   Phage T4   Predicted protein domains

Empty vector

PDC-S05   1 ▸ Virulence_RhuM 356

PDC-S70   1 ▸ DUF3368 166
ArsR−like_HTH

PDC-M05   1 ▸ NT_KNTase_like / Nucleotidyltransferases domain2 364

HEC-01   1 ▸ AAA_21 391

a

b

Type
- Defense genes
- PDC
- Transport
- MGE
- Function
- Unknown