Diverse defense systems synergistically drive the microbe-virus arms race in deep sea cold seeps

Yingchun Han[1][#], Jing Liao[1][#], Chengpeng Li[2], Fengmin Xing[3],

Jiaxue Peng[4], Xinyue Liu[1], Wentao Xie[5], Fabai Wu[6], Huahua Jian[5],

Rui Cheng[7][*], Xiyang Dong[1][*]

[1] Key Laboratory of Marine Genetic Resources, Third Institute of Oceanography, Ministry of Natural Resources, Xiamen 361005, China

[2] Key Laboratory of Advanced Marine Materials, Key Laboratory of Marine Environmental Corrosion and Bio-fouling, Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

[3] Institute of Marine Biology and Pharmacology, Ocean College, Zhejiang University, Zhoushan 316021, China

[4] College of Environmental Science and Engineering, Dalian Maritime University, Dalian 116026, China

[5] State Key Laboratory of Microbial Metabolism, Joint International Research Laboratory of Metabolic & Development Sciences, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai 200240, China

[6] School of Life Sciences, College of Science, Eastern Institute of Technology, Ningbo 315200, China

[7] Key Laboratory of Molecular Biophysics, the Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, China

[#] These authors contributed equally to this work.

[*] Correspondence can be addressed to Xiyang Dong (dongxiyang@tio.org.cn) and Rui Cheng (rui_cheng@hust.edu.cn).

Abstract

Cold seeps are unique deep-sea ecosystems that host diverse uncharacterized microbes and viruses, many of which potentially possess novel defense and counter-defense systems. Despite their ecological significance, the diversity of microbial defense systems, their interactions, and the evolutionary arms race with viral anti-defense strategies in cold seeps remain largely unexplored. To address this, we analyzed 3,813 microbial and 13,336 viral genomes from 193 metagenomic samples collected across 17 cold seep sites. Cold seep microbes harbor an extensive repertoire of `defense systems (n = 26,389), with 63% representing novel systems. Laboratory experiments confirm that several of these novel systems effectively protect against viral infections. These defense systems frequently co-occur and interact synergistically to combat viral infections. They are widely distributed across different sediment depths and geographical locations, exhibiting active expression. In response, cold seep viruses have evolved a diverse array of anti-defense genes (n = 1,197), employing complex strategies to counteract host defenses. The concurrent presence of multiple viral anti-defense genes and various microbial defense genes in their hosts highlights the intricate and dynamic interactions between viruses and their microbial counterparts. Functionally critical lineages, such as anaerobic methanotrophic archaea (ANME), sulfate-reducing bacteria (SRB), and diazotrophs, adapt their defensive strategies according to ecological niches and environmental pressures. Notably, SRB strains were found to harbor multiple Gabija defense systems, enhancing their resistance to viral attacks, while corresponding viruses carry anti-Gabija genes, illustrating a specific evolutionary adaptation. Overall, these findings underscore the critical role of virus-microbe interactions in shaping microbial metabolic functions and environmental adaptation in extreme deep-sea subseafloor environments.

Introduction

Cold seeps are extreme deep-sea environments located along continental margins worldwide, characterized by the migration of fluids rich in methane and other hydrocarbons from the deep subsurface to the sediment–seawater interface[1, 2]. The continuous release of these fluids enhances local biodiversity and microbial activity, making cold seeps thriving deep-sea oases[2]. These ecosystems support diverse microbial communities, including anaerobic methanotrophic archaea (ANME) and their syntrophic sulfate-reducing bacterial partners[3, 4]. Moreover, the sediments harbor a rich diversity of previously understudied prokaryotes and viruses[5, 6], suggesting the existence of complex and novel microbe–virus interactions within cold seep ecosystems. Prokaryotes in these environments constantly face viral threats, leading to the evolution of multiple defense mechanisms that regulate the flow of genetic information spread by mobile genetic elements (MGEs) through horizontal gene transfer (HGT)[7-9]. Cold seep viruses also employ diverse strategies for environmental adaptation, including counter-defense systems, auxiliary metabolic genes (AMGs), reverse transcriptases, and alternative genetic code assignments[10]. Among these, counter-defense systems are designed to counteract the defense mechanisms of host microbes. Despite these insights, the diversity of prokaryotic defense genes and the intricate evolutionary arms race between prokaryotes and viruses in cold seep ecosystems remain largely unexplored.

Prokaryotes have developed an arsenal of defense systems to provide partial or full resistance against viral and MGE infections[7, 8, 11]. These systems can be broadly categorized into two major groups based on their components and mechanisms of action: innate and adaptive immune systems[12, 13]. Innate immune systems provide rapid, non-specific defense mechanisms and include restriction–modification (RM) systems, which use methylation to distinguish self from non-self DNA[14], and abortive infection (Abi) systems, which lead to cell death or metabolic disturbance upon infection[15], and Gabija systems, composed of GajA and GajB proteins, which degrade

viral and host DNA through site-specific double-stranded DNA nicking by GajA upon

activation[16], and cyclic oligonucleotide-based antiphage signaling systems (CBASS),

which uses cyclic oligonucleotides to activate antiviral immunity[17]. Adaptive immune

systems enable microbes to adapt to evolving viral threats over time by creating a

tailored defense based on past infections; currently, this category is represented by the

clustered regularly interspaced short palindromic repeats (CRISPR)–Cas system[9].

This system provides acquired immunity by incorporating short DNA sequences from

viruses and MGEs into the host genome as spacers[18]. Recent efforts have uncovered

several defense systems with diverse genetic architectures[7, 8], reflecting the strong

selective pressure imposed by viruses and MGEs on microbial communities in various

ecosystems such as soil, marine, human gut and rumen[9, 19]. However, in the deep sea,

particularly in cold seep environments, there is a lack of corresponding studies on

microbial defense systems, despite these environments potentially equipped with vast

and untapped wealth of immune systems and mechanisms[5, 6].

Defense genes encoding components from multiple distinct systems tend to cluster

together in specific genomic regions referred to as defense islands[9, 20, 21]. This co-

localization, especially within MGEs, facilitates joint HGT, which is thought to confer

fitness advantages to recipient microbes, particularly in virus-rich environments[12].

Synergistic interactions between different defense systems may drive their co-

localization and favor their joint transfer[22, 23], as evidenced by the conservation of

certain sets of defense systems across different microbial species. Synergistic systems

enhance protective effects against specific phages beyond the sum of individual

systems, providing microbes with an evolutionary advantage. Recent research

indicated that non-random synergistic interactions between defense systems in

prokaryotes are adaptive responses to specific selective forces from the virome,

shaping unique fitness landscapes in their specific ecological niches[24]. Given that cold

seep viruses in extreme deep-sea subseafloor conditions are endemic and unique

compared to those in other ecosystems[6, 10], the specific virus-driven synergistic

interactions between defense systems may differ and warrant investigation.

Viruses can evade microbial defenses both indirectly, by mutating immune system activators or disrupting defense pathways, and directly, by deploying anti-defense genes that target immune system components[25]. In this ongoing arms race, viruses have evolved sophisticated anti-defense mechanisms to overcome microbial defense systems[25, 26], such as anti-CRISPRs (Acr)[27, 28], anti-RM[29], anti-CBASS[30] and anti-Gabija[31]. These viral anti-defense systems not only highlight the complexity of viral adaptations but also reveal evolutionary connections within microbial immunity[25]. Beyond defense evasion, viruses play a crucial role in shaping microbial communities and influencing ecosystem functioning, including key biogeochemical cycles such as methane oxidation, sulfate reduction and nitrogen fixation, often mediated through AMGs[32, 33]. Despite their importance, the interactions between viral anti-defense mechanisms and functionally critical lineages in the cold seep environment[26]—with its darkness, low temperatures, and hydrocarbon-rich conditions—is not well understood. Understanding these dynamics in extreme deep-sea ecosystem is vital for unraveling the full impact of viruses on microbial ecology and biogeochemical processes.

To explore the diversity of cold seep microbial defense systems, their interactions, and the evolutionary arms race between these systems and viral anti-defense strategies in extreme deep-sea subseafloor habitats, we analyzed 3,813 microbial genomes and 13,336 viral genomes from 193 metagenomic samples collected at 17 global sites (**Supplementary Fig. 1**). Our findings reveal that cold seep microbes possess a vast and diverse repertoire of defense systems, which tend to work synergistically against viral infections. In response, cold seep viruses deploy diverse anti-defense systems to counter microbial defenses. This complex virus–microbe interaction drives microbial evolution and enhances environmental adaptation. Overall, this study enriches our understanding of the ecological and evolutionary dynamics of virus–microbe interactions in deep sea cold seep subsurface ecosystems.

Results and Discussion

Cold seep prokaryotes possess diverse and novel defense systems

We investigated defense strategies of cold seep microbiome by analyzing 3,813 species-level metagenome-assembled genomes (MAGs), comprising 639 archaeal and 3,174 bacterial genomes, spanning 107 bacterial and 16 archaeal phyla **(Supplementary Fig. 2)**. The bacterial phyla with the largest diversity of recovered species included *Pseudomonadota* (n = 471), *Chloroflexota* (n = 399), *Desulfobacterota* (n = 298), *Planctomycetota* (n = 225), and *Bacteroidota* (n = 219). Among the archaeal phyla, *Halobacteriota* (n = 153), *Thermoplasmatota* (n = 119), *Thermoproteota* (n = 110), and *Asgardarchaeota* (n = 105) were the most diverse.

A total of 36,783 defense genes were detected in 65% of cold seep microbial genomes, which were assigned to 26,389 defense systems **(Fig. 1a and Supplementary Tables 1-3)**[8, 34]. Among these, candidate defense systems accounted for 63% (n = 16,520), comprising 18,195 defense genes, indicating that cold seep microbes harbor a diverse array of novel defense strategies. Experimentally confirmed defense systems, comprising 37% (n = 9,869), included 106 types, with RM systems (n = 3,778) being the most prevalent, followed by CRISPR–Cas (n = 977), SoFic (n = 715), and AbiE (n = 612). This distribution aligns with previous findings that RM and Cas systems are predominant in microbiomes from soil, human gut, and Haima cold seep environments[9, 10]. In contrast, seawater microbial genomes, while also rich in RM systems, exhibit significant enrichment of CBASS[9], suggesting distinct distribution patterns of defense systems in cold seep sediments compared to other marine environments.

To validate whether these novel systems confer defense against viruses, we selected four candidate defense systems, namely PDC-S05, PDC-S70, PDC-M05, and HEC-S01, for experimental examination. Recombinant vectors containing these candidate

systems were transformed into the host strain *Escherichia coli* B (ATCC® 11303™),

which naturally lacks these systems. The transformed strains were then challenged

with three representative lytic phages: T7 (*Podoviridae*), T5 (*Siphoviridae*), and T4

(*Myoviridae*). All four systems conferred protection against at least one phage **(Fig.**

1b). Especially, the PDC-S05 and PDC-S70 systems exhibited obvious and broad-

spectrum phage resistance. The PDC-S05 system showed an approximately 1,000-fold

reduction in efficiency of plating (EOP) for phages T7 and T5, and a ~10-fold

decrease for phage T4. Similarly, the PDC-S70 system caused a ~100-fold decrease in

EOP for phages T7 and T4, and an ~1,000-fold decrease for phage T5. These results

confirm the vast, yet uncharacterized, diversity of defense systems present in cold

seep microbes.

Defense system mechanisms can be categorized in three types based on their

fundamental mechanisms of action: nucleic acid degradation, abortive infection,

inhibition of replication, and other unknown mechanisms[13, 21]. For instance, RM and

Cas systems function through viral nucleic acid degradation, CBASS and retrons

through abortive infection, and Viperins and dCTP deaminases through inhibition of

replication. Among the experimentally confirmed defense systems in cold seep

microbes **(Fig. 1a)**, the most prevalent mechanism was nucleic acid degradation

(55%), followed by abortive infection (19%) and replication inhibition (7%). Notably,

a substantial proportion of defense mechanisms remain unidentified (19%),

encompassing 59 types that represent 56% of all detected defense system types.

Cold seep defensive patterns are associated with genome size and taxonomy

The distribution of defense systems differed considerably across taxonomical domains

(Fig. 2a and Supplementary Fig. 3), suggesting that archaea and bacteria possess

diverse and distinct defense strategies against viral or MGE infections[12, 34, 35]. Most

bacterial and archaeal MAGs harbored a limited number of defense systems,

averaging four per genome **(Fig. 2b)**, consistent with observations from other marine

environments[9, 10]. On average, bacterial genomes encode more defense systems per

genome than archaeal genomes (4.1 vs 3.5). Despite this average, several genomes (n

= 40) encoded more than 20 defense systems. For instance, a genome belonging to the

Bacteroidales order encodes 54 defense systems, followed by 48 in a *Methylococcales*

genome (*Gammaproteobacteria*) and 39 in a *Desulfuromonadales* genome

(*Desulfobacterota*) (**Supplementary Table 3**).

A positive correlation was observed between the total number of defense systems and

genome size (R^2 = 0.1678, *P* < 2.2e-16) for each MAG (**Fig. 2c, Fig. 3,**

Supplementary Fig. 4 and Supplementary Table 4). Additionally, a negative

correlation was found between the density of defense systems and genome size (*P* =

0.03). These trends can be attributed to the fact that microbes with larger genomes

typically engage in more HGT, necessitating a more abundant and diverse array of

defense systems[36]. While the majority of defense systems (~99.88%) are encoded in

chromosomes, several systems—especially AbiV, AbiD, DRT, and AbiL—are carried

by plasmids, and others like AbiZ, PT_Dnd, and Lamassu are carried by proviruses

(**Fig. 3 and Supplementary Fig. 5a**). Furthermore, the number of defense systems in

prokaryotes increases with the number of proviruses and plasmids, a trend particularly

evident in bacteria (**Supplementary Fig. 5b**), suggesting that defense genes in cold

seep prokaryotes could be disseminated between microbes through MGEs[37, 38].

The density of defense systems (per MAG and per Mb) varied widely among different

clades (**Fig. 3, Supplementary Figs. 6-7, and Supplementary Table 3**). In archaeal

MAGs, the density ranged from as low as 0.11 in certain phyla to over 10.5 in the

Aenigmatarchaeota phylum (**Fig. 3a and Supplementary Fig. 6**). In bacterial MAGs,

the density can be as high as 15.1 in the *Desulfobacterota* phylum (**Fig. 3b and**

Supplementary Fig. 7). The number of defense genes is influenced by phylogenetic

depth, as deeper lineages tend to accumulate more HGT events, presumably leading to

a buildup of defense systems[24]. This is evident in the high-density defense systems

observed within the phyla *Desulfobacterota*, *Nanoarchaeota*, and *Asgardarchaeota*.

These genomes, which represent greater phylogenetic depth, tend to have a higher density of defense systems **(Supplementary Table 3)**. However, linear regression analyses revealed that phylogenetic depth has a relatively minor impact on the number of defense systems (**Supplementary Table 5**). This suggests that while phylogenetic depth and HGT events contribute to the presence of defense systems, other factors also play important roles in determining their abundance and distribution.

Cold seep defense systems tend to co-occur and interact synergistically

Previous studies have demonstrated that certain defense systems interact to enhance or broaden protection against viruses[24, 39, 40]. Here, we investigated the relationships among defense systems in cold seep prokaryotes to explore their potential interactions. Our analysis identified 182 pairs of defense systems exhibiting significant positive correlations ($P < 0.05$), whereas only 10 pairs showed significant negative correlations ($P < 0.05$; **Fig. 4a and Supplementary Table 6**). These findings suggest a propensity for defense systems to co-occur rather than to be negatively associated, consistent with observations in certain bacterial genomes (e.g., *Escherichia coli*, *Enterobacterales, Bacillales, Burkholderiales,* and *Pseudomonadales*)[24]. Notably, prevalent defense systems such as Gabija (20 pairs), RM (19 pairs), Cas (19 pairs), and SoFic (18 pairs) exhibited a higher number of significant co-occurrences. Additionally, less common systems such as Septu (18 pairs), Druantia (15 pairs), and Menshen (15 pairs), also demonstrated a tendency to co-occur with other defense systems. Interestingly, different subtypes of the same defense system exhibited distinct co-occurrence patterns with other systems **(Supplementary Fig. 8 and Supplementary Table 7)**. For example, Wadjet I co-occurred with multiple other defense systems, showing 11 pairs of co-occurrences. In contrast, Wadjet II had only 4 co-occurrences, which included Cas I-G, CBASS III, Eleos, and PD-T4-3; notably, none of these systems co-occurred with Wadjet I. These specific co-occurrence patterns may indicate unique cooperative interactions among different subtypes of defense systems.

Defense islands are sometimes associated with MGEs integrated into distinct hotspots—regions of high genetic turnover—that may serve as catalysts for novel defensive strategies[9, 24]. The clustering of defense systems in these islands, especially within integrated MGEs, enhances the probability of horizontal co-transfer. If the repertoire of defense systems is primarily shaped by HGT, rather than functional synergies, we would expect the systems co-localizing in defense islands to correspond with those co-occurring in microbial genomes[24]. To explore the potential link between the co-occurrence and co-localization of defense systems, we selected 34 defense systems clustered in defense islands and analyzed the correlations among them. Among the clustered systems, 105 pairs were identified with significant correlations ($P<0.05$), which is lower than the 182 significant pairs observed when considering all defense systems. Notably, only one pair among the clustered systems exhibited a negative correlation (**Fig. 4a and Supplementary Table 8**). Furthermore, there appears to be no obvious relationship between the number of defense systems clustered in defense islands and the number of significantly correlated pairs among those systems (**Fig. 4a and Supplementary Fig. 9**). For instance, although RM and Cas systems were frequently found in defense islands, they did not exhibit an obvious higher number of correlations with other systems. In contrast, systems like BREX (13 pairs), Septu (13 pairs), and PT_PbeABCD (12 pairs), which were less commonly associated with defense islands, showed more significant correlations. These findings suggest that co-localization represents only a specific instance of co-occurrence, and the repertoire of defense genes is influenced by various factors beyond HGT.

The genomic context of defense islands supports the observed correlations among defense systems. For instance, the CBASS system frequently clusters within these islands, often flanked by defense genes related to RM and Abi systems (**Fig. 4b and Supplementary Table 9**). These defense islands are not limited to defense genes alone; they also encompass a rich array of functional genes involved in transport, replication, recombination, repair, and transcription. This high genetic turnover within defense islands creates hotspots of genetic diversity, promoting the evolution of novel

defense strategies in·cold·seep microorganisms[9]. Thus, the clustering and dynamic

interactions within defense islands highlight their role in the co-localization and co-

evolution of defense systems.

Cold seep defense systems are abundant, actively expressed and site-dependent

In alignment with the prevalence of defense systems in microbial genomes **(Fig. 1a)**,

RM, Cas, AbiE and SoFic systems exhibit higher metagenomic abundances, primarily

due to the elevated abundances of restriction endonucleases (REases),

methyltransferases (MTases), AbiEi, AbiEii, SoFic proteins, and Cas2 genes

associated with these systems **(Fig. 5a and Supplementary Fig. 10)**. These

differences are statistically significant at both systems and gene levels ($P < 0.001$).

Although there were no significant differences in metatranscriptomic abundances

among defense systems, the observed expression levels (averaging 3.64 transcripts per

million [TPM], with over 1 TPM for 30 systems) indicate active microbial defensive

responses *in situ*. Additionally, most defense systems were detected across

geographically diverse cold seep sediments (17 sites) at almost every sediment depth

(0-68.55 meters below seafloor [mbsf]; **Supplementary Fig. 11)**, indicating a

widespread presence of these systems in cold seeps globally. High abundances of

defense systems were particularly notable in archaeal *Halobacteriota*, and bacterial

Desulfobacterota, *Atribacterota*, and *Pseudomonadota* **(Fig. 5b)**. The relative

abundance of defense systems varied considerably across phyla, with RM, Cas, AbiE

and SoFic systems being the most predominant.

The distribution patterns of defense systems in global cold seep prokaryotes are

shaped not only by the prokaryotes but also by their targets, such as viruses and

MGEs. The distribution of cold seep microbes and viruses is influenced by sediment

depth and site location[10, 41, 42] ; consequently, these factors may also impact the

distribution of defense systems. To assess depth stratification of defense systems, we

categorized each metagenome based on depth ranges: surface (<1 mbsf), shallow (1 to

10 mbsf), and deep (>10 mbsf). We found that the abundance of defense systems (especially predominant RM and Cas systems) in surface samples was slightly higher than in shallow and deep samples, although these differences were not statistically significant (**Fig. 5c and Supplementary Fig. 12a**). This observation may be attributed to prokaryotes in deeper layers compensating for their lower abundance by increasing the copy number of defense genes, adapting to more stable and extreme environments[42]. The expression patterns of defense genes, with higher transcript abundances in deeper sediment layers (both shallow and deep) compared to surface layers (**Fig. 5d and Supplementary Fig. 12b**), further support this hypothesis. Significant differences in defense system abundance were also found between sites, with more pronounced differences in shallower depths (surface and shallow, $P \leq 0.05$) compared to deeper depths (deep, $P > 0.05$; **Supplementary Fig. 13a**). Metatranscriptomic abundance also varied significantly across different sites, with a more notable difference observed in surface depths ($P < 0.01$; **Supplementary Fig. 13b**). Overall, the distribution of defense systems in cold seep prokaryotes appears to be slightly dependent on depth and more dependent on site-specific characteristics.

Cold seep viruses employ diverse anti-defense genes to counteract host defense systems.

We retrieved 13,336 single-contig viral genomes with estimated completeness of greater than 50% from 193 cold seep metagenomes (**Supplementary Fig. 14 and Supplementary Table 10**) using three virus identification tools: VIBRANT[43], Virsorter2[44] and geNomad[45]. These viral genomes were clustered into 9,971 species-level viral operational taxonomic units (vOTUs)[46]. Of these, 9,638 vOTUs (~97%) could be taxonomically assigned, predominantly belonging to the *Duplodnaviria* phylum (n = 9,600), with a majority classified under the *Caudoviricetes* class (n = 9,599; **Supplementary Fig. 15 and Supplementary Table 11**). However, less than 0.4% vOTUs (n = 36) within this class could be annotated further, highlighting an obvious knowledge gap in the taxonomy of deep-sea cold seep viruses[6, 10].

Host predictions identified a total of 2,688 connections between prokaryotes (810

genomes) and viruses (1,310 vOTUs) **(Fig. 6a, Supplementary Fig. 16 and**

Supplementary Table 12). These host–virus linkages spanned 57 phyla, including 10

archaeal and 47 bacterial phyla. The most common predicted host phyla were

Chloroflexota (n = 156), followed by *Halobacteriota* (n = 86), *Pseudomonadota* (n =

85), *Planctomycetota* (n = 56), and *Asgardarchaeota* (n = 52). Most of the identified

connections involved viruses in the *Caudoviricetes* class infecting the *Chloroflexota*

(731 pairs), and archaeal *Asgardarchaeota* (446 pairs) and *Halobacteriota* (284 pairs).

The host–virus network revealed that the majority of clusters (60%, n=1,616)

consisted of more than one viral or prokaryotic genome. This suggests that a single

virus could infect multiple prokaryotic populations, or a single prokaryote may be

susceptible to several viral population[47], indicating a complex relationship between

viruses and prokaryotes in cold seeps.

Viruses encode an arsenal of anti-defense proteins that facilitate infection by disabling

various prokaryotic defense mechanisms[20, 25]. A total of 1,197 anti-defense genes were

detected from 11% viral genomes (n = 1,065), predominantly comprising anti-Cas,

anti-Thoeris, anti-Gabija, and anti-RM genes (**Fig. 6b and Supplementary Table 13)**.

While most anti-defense genes were detected across a broad range of viruses, some

were found only in specific viral genomes. For example, anti-Cas, anti-RM, anti-

Thoeris, and anti-Gabija genes were commonly found, whereas anti-RecBCD genes

and broad-spectrum anti-defense genes were exclusively detected in certain

Caudoviricetes viruses. From the 2,688 host-virus linkages, we identified 155 pairs in

which host microbes contained defense genes while the corresponding viruses

possessed anti-defense genes (**Fig. 6c and Supplementary Table 12)**. These

interactions were predominantly observed between viruses in the *Caudoviricetes* class

and their hosts, including archaeal groups such as *Asgardarchaeota* (38 pairs) and

Halobacteriota (26 pairs), as well as various bacterial taxa including *Pseudomonadota*

(25 pairs) and *Desulfobacterota* (24 pairs). Notably, a single virus employing two

anti-defense systems could infect two ETH-SRB1 genomes belonging to the

Desulfobacterota phylum, and one *Asgardarchaeota* genome was found to harbor six

defense systems to fend off infection by five viruses (**Fig. 6c and Supplementary**

Table 12). These findings highlight a complex evolutionary arms race between these

prokaryotes and their viral counterparts, particularly among archaea and viruses[10, 26, 48].

A single virus may encode multiple anti-defense genes, enabling it to counter various

host defense strategies or adapt to different defenses employed by the same host

throughout its life cycle[26]. We identified 118 viruses that encode two or more types of

anti-defense systems (**Supplementary Table 12**). For instance, a viral genome from

the *Caudoviricetes* family contains AcrIF17, AcrIF24, and Tad1 genes to counteract

the CRISPR-Cas and Thoeris defense systems of its host belonging to the phylum

Pseudomonadota (**Fig. 6c**). Additionally, two viral genomes from the same family

harbor DarB and Tad2 genes to counteract the RM and Thoeris defense systems of

their host, ETH-SRB1 (**Fig. 6d and Supplementary Table 12 in green**). AcrIF24

form homodimers and induce dimerization of their target effector complexes,

blocking the hybridization of the CRISPR RNA with its target[49]. AcrIF24 engages up

to five Cas7f subunits (Cas7.2f-7.6f), sterically blocking access to the crRNA guide

by the complementary target DNA, and inducing non-sequence-specific DNA binding

to the Csy complex[49]. The dimeric protein structure of AcrIF24 from *Caudoviricetes*

exhibits high structural similarity (TM-score = 0.86) to the anti-CRISPR protein form

Pseudomonas aeruginosa (**Fig. 6e**), which has been experimentally confirmed to

inhibit Cas7f[49]. Additionally, many anti-defense genes interact with immune system

components through steric or allosteric effects in a stoichiometric, non-enzymatic

manner were also detected[25]. For example, the production of AcrIIA7 and most Acrs

(e.g., AcrIIA2, AcrIIA4, AcrIIIB1, AcrIIC1) is often repressed by Aca proteins

following an initial strong surge in expression and binding to the targeted CRISPR–

Cas component[50-52]. Some enzymatic inhibitors effectively suppress host defenses by

specifically and permanently specifically microbial immunity[25, 53, 54]; examples

detected in cold seep viruses include Acb1, NTases, AcrIII-1, and AcrVA1–5. Despite

these insights, many anti-defense systems with unknown molecular mechanisms, such

as Anti-Dnd, Gad2, Abc1, and KlcA, were also present. Overall, viral anti-defense systems engage in a complex evolutionary arms race with host defense systems through both enzymatic, non-enzymatic and unknown molecular mechanisms in cold seep ecosystems.

Defense systems protect key metabolic microbial groups from viral infections

Anaerobic methane oxidation, sulfate reduction, and nitrogen fixation, associated with the activities of the *mcrA*, *dsrA*, and *nifH* genes, respectively, are key metabolic processes in cold seeps[55-57]. The significant correlation between the abundance of defense systems and these key metabolic genes **(Supplementary Fig. 17)** underscores the role of defense systems in protecting functional microbes from viral and MGE infections. To further explore this relationship, we analyzed 35 anaerobic methanotrophic archaea (ANME), 47 sulfate-reducing bacteria (SRB), and 57 diazotrophs identified from 3,813 species-level MAGs. Although the defense systems of these microbes generally lack specificity—most encoding RM systems, followed by Cas, AbiE, and SoFic **(Fig. 7a, Supplementary Fig. 18 and Supplementary Table 13)**—the total number of systems and the diversity of families varied significantly within specific functional groups, particularly among SRB and diazotrophs **(Fig. 7b)**. For example, within SRB, ETH-SRB1 encode on average 17.6 defense systems across 7.3 types, whereas another SRB strain in *Desulfatiglandaceae* order (SY5_S20_sbin_4) encodes only 3 systems of just 1 type. Despite these variations, most ANME encode a limited number of defense systems with fewer types.

In detail, SRB predominantly encode type IIG and type II RM systems, followed by type I RM systems. ANME and diazotrophs primarily encode type IIG and type I systems, with type II RM systems following **(Supplementary Fig. 19 and Supplementary Table 14)**. However, the number of these systems in ANME is obvious smaller compared to the other two functional groups. RM systems primarily function in the early stages of viral infection to eliminate invading nucleic acids[11],

indicating that cold seep prokaryotes possess the ability to respond rapidly and clear

viruses promptly. CRISPR–Cas systems are categorized into two classes and six types

based on the set of Cas genes they encode and whether their effector is a single

protein or a protein complex[18, 58, 59]. Both classes of CRISPR-Cas were present in cold

seep microbes, with class I being more abundant in both number and diversity

(**Supplementary Fig. 20 and Supplementary Table 14**). Most archaea, including

ANME, primarily encode type I CRISPR-Cas systems, while the majority of bacteria,

including SRB and diazotrophs, mainly encode both type I and type III CRISPR-Cas

systems. CRISPR-Cas has been shown to contribute to shaping a highly dynamic

network of interactions between viruses/MGEs and their prokaryotic hosts in various

ecosystems[60-62]. Additionally, SRB tend to harbor two or more Gabija systems,

particularly in ETH-SRB1 strains, with some encoding as many as four Gabija

systems (**Supplementary Fig. 22 and Supplementary Table 14**). Overall, these

findings highlight the diversity and complexity of defense systems across various

microbial groups, likely shaped by distinct evolutionary pressures and ecological

niches[21].

Gabija is effective in providing bacterial immunity against certain bacteriophages, as

shown by its strong defense against phi29, rho14, phi105, and SpBeta in *Bacillus*

cereus VD045 [7]. This system is widespread in prokaryotes, present in about 15% of

all analyzed genomes[7]. Through sequence alignment and both sequence- and

structure-based phylogenetic analyses (**Supplementary Fig. 21**), we found that

Gabija is widely present in cold seep microbes, spanning 45 phyla (including 9

archaeal and 36 bacterial phyla) and encompassing the three key metabolic microbial

groups. We discovered that one ETH-SRB1 strain encodes two Gabija systems, and

the virus infecting it carries the *gad1* anti-defense gene (**Fig. 7c**). This ETH-SRB1

strain harbored both a complete and an incomplete Gabija system. In the complete

system, the GajA protein contains an N-terminal ATPase domain and a C-terminal

Toprim (topoisomerase-primase) domain, while the GajB protein includes an AAA

domain and a UvrD-like helicase[31, 63]. In contrast, the GajB of the incomplete Gabija

system lacks the UvrD-like helicase **(Supplementary Fig. 22)**. The complete Gabija

system forms a supramolecular complex and exhibits high structural similarity (TM-

score = 0.70) to the Gabija protein from *Bacillus cereus* VD045 [31] **(Fig. 7d)**. We also

identified interaction sites between GajA and GajB, which are critical for the

formation of the GajAB supramolecular complex essential for Gabija-mediated anti-

phage defense[31, 63]. The viral protein Gad1 **(Fig. 7e)**, which is larger than other phage

immune evasion proteins (35 kDa), binds and encapsulates the GajAB complex,

blocking the degradation of target DNA. The binding of Gad1 to the GajAB complex

encoded by a virus targeting ETH-SRB1 suggests that Gad1 can effectively inhibit the

Gabija system in the ETH-SRB1 strain.

Considering the differences in the number and types of defense systems among

functional microbial groups, we hypothesize that the specific pairs of co-occurring

and negatively associated defense systems vary across these groups. As anticipated,

defense systems that co-occurred in one group were absent in another

(Supplementary Fig. 23 and Supplementary Table 15). For example, AbiE system

in ANME exhibited 28 co-occurrences, including with CBASS, DarTG, Gabija, and

Mokosh, none of which were observed in SRB or diazotrophs. These findings

highlight the unique defensive strategies employed by different microbial lineages.

Previous research has revealed that the interactions between defense systems likely

depend on environmental and genetic factors that select for a particular anti-virus

immunity strategy, and that synergistic immunity provides an evolutionary advantage

to microbial populations[24]. Overall, our results indicate that cold seep microbial

defense systems are generally mechanistically compatible, allowing microbes to adopt

diverse and flexible strategies for anti-virus defense based on their unique

environmental and genetic contexts.

Conclusions

Cold seep ecosystems, characterized by the seepage of hydrocarbon-rich fluids into deep-sea sediments, harbor a diverse array of previously uncharacterized microbes and viruses engaged in a continuous evolutionary arms race. In this study, we identified a diverse and novel set of defense systems in cold seep microbes, which work synergistically to combat infections from MGEs and viruses. Experimental validation confirmed that several of these novel defense systems effectively protect against viral infections. Defense systems in cold seep microbes frequently co-occur and exhibit significant positive correlations, suggesting cooperative interactions that enhance antiviral defense. Notably, defense systems such as RM, Cas, AbiE, and SoFic exhibited high metagenomic abundances, widespread distribution, and active expression across various microbial genomes from different sediment depths and geographical locations, underscoring their vital role in microbial survival within these extreme environments. In response, cold seep viruses have evolved diverse anti-defense mechanisms, functioning through enzymatic, non-enzymatic, and as-yet-unknown mechanisms to evade host defenses. The presence of multiple viral anti-defense genes within single viral genomes, along with various defense genes in their hosts, highlights the intricate and dynamic interactions between these viruses and their microbial hosts, reflecting an ongoing evolutionary arms race. Functionally critical lineages in cold seeps, such as ANME, SRB, and diazotrophs, adjust the interactions among different defense systems according to their ecological niches and environmental pressures, enabling them to adapt to the evolutionary challenges posed by MGEs and viruses in extreme deep-sea subseafloor environments. Overall, these findings emphasize the complexity and flexibility of interactions within microbial defense systems in extreme environments. The ability of cold seep microbes to maintain diverse and adaptable defense mechanisms highlights their resilience and evolutionary success in one of Earth's most challenging habitats. This study deepens our understanding of virus–microbe interactions in cold seep ecosystems and

underscores the significance of microbial defense systems in shaping microbial ecology and evolution in extreme environments.

Materials and Methods

Dataset and processing

Metagenomic assemblies were generated from 173 deep-sea sediment samples collected from 17 geographically diverse cold seep sites, with sediment depths ranging from 0 to 68.55 mbsf **(Supplementary Fig. 1)**. The construction of the genomic inventory followed the methods detailed in our previous study[5]. Raw metagenomic sequence data underwent quality control and were assembled into contigs sequences. Contigs exceeding 1,000 base pairs in length were selected for binning. The resulting MAGs were deduplicated using dRep v3.4.0 [64] (parameters: -comp 50 -con 10) at a 95% average nucleotide identity threshold, yielding a total of 3,813 cold seep prokaryotic MAGs. Initial taxonomic assignment of each MAG was conducted using GTDB-TK v2.1.3 [65] with reference to the Genome Taxonomy Database (GTDB) release R214, followed by validation through a maximum-likelihood phylogenomic tree. For the archaeal MAGs, the phylogenomic tree was inferred based on the concatenation of 53 conserved single-copy genes as marker genes, while for the bacterial MAGs, it was based on 120 marker genes. The trees were constructed using IQ-TREE v2.2.0.3 (parameters: -m MFP -B 1000)[66] and visualized by using iTOL v6 [67]. Sankey plots were constructed using Pavian (https://fbreitwieser.shinyapps.io/pavian/)[68]. Protein-coding sequences were predicted from the MAGs using Prodigal v2.6.3 (parameter: -p meta)[69], resulting in a gene catalog annotated from all MAGs.

Thirty-three metatranscriptomes, originating from cold seeps of the South China Sea in regions of Jiaolong, Haima, Qiongdongnan Basin, and the Shenhu area[57, 70-72] **(Supplementary Fig. 1)**, were subjected to quality control using the Read_QC

module of the metaWRAP v1.3.2 pipeline (parameters: --skip-bmtagger)[73]. Following

this, SortMeRNA v2.1 was utilized to remove ribosomal RNA[74]. The resulting clean

reads were then aligned to the gene catalog annotated from all MAGs, employing

Salmon v.1.9.0 (parameters: -validateMappings -meta)[75], to quantify the transcript

abundance of prokaryotic defense genes. Transcript levels were normalized to

transcripts per million (TPM).

Identification of defense genes, systems, and islands

Defense genes and systems were identified using DefenseFinder v1.2.1 (parameters: -

-db-type gembase)[21] and PADLOC v2.0.0 (parameter: --fix-prodigal)[34].

DefenseFinder screens for 710 genes related to 154 defense system families, while

PADLOC screens for 385 defense subsystems. The results from the two tools were

merged, with duplicate results compared based on e-values to retain those with higher

confidence. Defense islands were defined as arrays of defense genes (or defense

systems) separated by no more than ten genes and containing at least five genes from

at least three different defense systems[9]. Gene abundances of gene catalog annotated

from all MAGs were quantified using Salmon v.1.9.0 in mapping-based mode

(parameters: --meta --validateMappings)[75] and read counts were normalized to GPM

(genes per million). The abundance of a defense system is represented by the sum of

the abundances of the genes within that system divided by the average number of

genes in the system.

Identification of viruses and virus–host linkages

Potential viral sequences were identified from metagenomic assemblies (contig

lengths greater than 10 kb) derived from the aforementioned 173 cold seep site

samples and an additional 20 cold seep samples. Three methodologies were used:

geNomad v1.7.0 (parameters: end-to-end)[45], Virsorter2 v2.2.3 (parameters: default)[44],

and VIBRANT v1.2.1 (parameters: default)[43]. The completeness and contamination of

these sequences were assessed using CheckV v1.0.1 with the database v1.5 [76].

Sequences with an estimated completeness of 50% or more were selected for
clustering into species-level viral operational taxonomic units (vOTUs). Clustering
parameters followed CheckV guidelines: 95% average nucleotide identity and a
minimum aligned fraction of 85%. This process resulted in the identification of 9,971
species-level vOTUs. Taxonomic assignment of these vOTUs was conducted using
geNomad v1.7.0. The resulting taxonomic lineages adhere to the classification
outlined in the International Committee on Taxonomy of Viruses (ICTV) Virus
Metadata Resource (VMR) number 19. Open reading frames (ORFs) within each
vOTU were predicted using Prodigal v2.6.3 (parameter: -p meta)[69]. Functional
annotation of the viral sequences was performed utilizing the virome model in DRAM
v1.4.0 [77].

Linkages between viruses and their hosts were primarily determined using the iPHoP
v1.3.2 pipeline[78], utilizing a database constructed from 3,813 prokaryotic MAGs
identified in this study and the iPHoP_db_Sept21_rw. iPHoP employs a variety of
methods to predict virus–host linkages, including Blastn v2.12.0 [79] alignment of viral
genomes against the iPHoP_db_Sept21_rw database and its CRISPR spacer database;
The k-mer composition similarity between viral and host genomes was analyzed using
four methods: RaFAH v0.3 [80], WIsH v1.0 [81], VirHostMatcher-Net[82], and PHP[83]. In
addition to the iPHop methods, for contigs identified with CRISPR-Cas systems,
CRISPRCasFinder v4.2.21 [84] was used to identify CRISPR arrays flanking the Cas
genes. Local alignments of extracted spacers (with lengths greater than 25 base pairs)
against viral genomes were performed using Blastn. Only BLAST matches with 100%
alignment coverage and at most two mismatches were considered high-confidence
matches. The network graph between hosts and viruses was visualized using Gephi
v0.9.2 [85]. The workflow for this part was illustrated using Draw.io v23.1.5
(https://app.diagrams.net/).

Identification of anti-defense genes

The anti-defense genes of viruses were primarily identified through two approaches. (1) APIS database search. We utilized the Anti-Prokaryotic Immune System (APIS) database[86], which contains experimentally validated viral proteins that counteract prokaryotic immune systems and their associated protein families. All viral proteins from our dataset were compared against the APIS database using DIAMOND blastp v2.0.8 (parameters: -e 1e-10 --id 30)[87] and HMMER v3.3.2 (parameters: -E 1e-10). (2) Reference sequence comparison. We compiled sequences of viral anti-defense proteins as documented by Mayo-Muñoz et al. in their review[25]. These 36 reference sequences were downloaded from the NCBI database and used to construct a custom database using the DIAMOND makedb command. Our viral proteins were then compared against this database using DIAMOND BLASTp v2.0.8 (parameters: -e 1e-10 --id 30). The results were obtained using two methods, with the more reliable results selected based on e-values to derive the final set of anti-defense genes.

Experimental validation of candidate defense systems

Coding sequences of the candidate defense systems were commercially synthesized and cloned into the pQE82L expression vector. The recombinant vectors were transformed into *Escherichia coli* B (ATCC® 11303™). Cells transformed with the empty vector served as controls. Plaque assays were conducted following established protocols[16, 88]. A single colony from a fresh LB agar plate was inoculated into LB broth containing ampicillin (100µg/mL) and grown at 37°C to an OD_{600} of approximately 0.6. Protein expression was induced by adding 0.2mM isopropyl β-D-1-thiogalactopyranoside (IPTG). After an additional hour of incubation, 500µL of the bacterial culture was mixed with 14.5mL of 0.5% LB top agar, and the mixture was poured onto LB plates containing ampicillin (100µg/mL) and IPTG (0.1mM). The plates were then spotted with 4µL aliquots of phage dilutions prepared in LB medium. Eight 10-fold serial dilutions were used for each phage: dilutions 10^{-1} to 10^{-8} for

601 phage T7, and 10 to 10^{-7} for phages T4 and T5. Plates were incubated overnight at

602 25°C before imaging.

Functional annotations

Functional annotation of MAGs was performed by searching against the KEGG, Pfam,

MEROPS, and dbCAN databases using DRAM v1.4.0 (parameters: --min_contig_size

1000)[77]. Gene context was visualized with Chiplot (https://www.chiplot.online/).

Plasmid and provirus sequences were identified from prokaryotic MAGs using

geNomad v1.7.0 [45]. To classify different types of MGEs, such as IS elements and

transposons (IS_Tn), integrons, and conjugative elements (CE), recombinase marker

genes, were identified through alignment with the proMGE database[89] using HMMER

v3.3.2. For the identification of key metabolic genes in cold seep prokaryotes, *dsrA*

genes were detected using DiSco v1.0.0 [90], and *mcrA* genes were identified using

METABOLIC v4.0 [91] and DRAM v1.4.0 [77]. Nitrogen fixation marker genes *nifH*

genes were detected using three methods: (1) protein sequences in the MAGs were

compared against NCycDB[92] using DIAMOND blastp v2.0.8 [87], (2) potential *nifH*

sequences were searched based on *nifH* reference sequences (n = 1271) from the

Greening lab metabolic marker gene database using DIAMOND blastp v2.0.8

(parameters: --id 65)[87], and (3) *nifH* genes were extracted using HMMER v3.3.2 with

the TIGRFAM model TIGR01287.1.

Protein sequence and structural analysis

To verify the accuracy and phylogenetic placement of the identified defense and anti-

defense genes, multiple sequence alignments were performed using MAFFT v7.505 [93].

Alignments were trimmed using TrimAl v1.4.1 under default settings[94]. Maximum

likelihood phylogenetic trees were constructed using IQ-TREE v2.2.0.3 (parameters: -

m MFP -B 1000)[66]. Trees were visualized and annotated using iTOL v6 [67]. Protein

domains were identified using the Batch Protein Annotation plugin in Tbtools-II

v2.042 [95], which integrates results from InterProScan[96]. Sequence conservation among

proteins was aligned and visualized using Jalview v2.11.3.3 [97]. Structural predictions

for proteins, including GajA and GajB, were performed using ColabFold v1.5.5 which

leverages AlphaFold2 [98-100]. Structures of the AcrIF24 dimer, GajAB octamer, and

Gad1 protein were predicted using AlphaFold3 [101]. Homologous and structurally

similar proteins were identified using the Foldseek web interface[102]. Protein structures

were visualized and aligned using PyMOL v2.5.5 [103].

Statistical analyses

Statistical analyses were conducted using R v4.2.3. Normality of the data was

assessed using the Kolmogorov–Smirnov test, and homogeneity of variances was

tested using Levene's test. The Wilcoxon rank-sum test was employed to compare

characteristics of defense systems between archaeal and bacterial groups. For

comparisons involving more than two groups—such as defense system and gene

densities among different microbial phyla, or differences in gene and transcript

abundances among various types of defense systems or genes—the Kruskal–Wallis

test was used. Pearson's correlation coefficients were calculated to assess correlations

among defense systems and subsystems present on defense islands, based on their

abundances.

Data availability

Sequences of defense systems, viral genomes, viral anti-defense systems, and

phylogenetic trees of GajAB based on amino acid sequences and protein structures are

available at figshare (https://doi.org/10.6084/m9.figshare.27004438).

Code availability

The present study did not generate codes, and mentioned tools used for the data

analysis were applied with default parameters unless specified otherwise.

References

1. Dong, X. et al. Metabolic potential of uncultured bacteria and archaea associated with petroleum seepage in deep-sea sediments. *Nature Communications* **10**, 1816 (2019).

2. Joye, S.B. The geology and biogeochemistry of hydrocarbon seeps. *Annual Review of Earth and Planetary Sciences* **48**, 205-231 (2020).

3. Dong, X. et al. Thermogenic hydrocarbon biodegradation by diverse depth-stratified microbial populations at a Scotian Basin cold seep. *Nature Communications* **11**, 5825 (2020).

4. Kleindienst, S. et al. Diverse sulfate-reducing bacteria of the *Desulfosarcina/Desulfococcus* clade are the key alkane degraders at marine seeps. *The ISME Journal* **8**, 2029-2044 (2014).

5. Han, Y. et al. A comprehensive genomic catalog from global cold seeps. *Scientific Data* **10**, 596 (2023).

6. Li, Z. et al. Deep sea sediments associated with cold seeps are a subsurface reservoir of viral diversity. *The ISME Journal* **15**, 2366-2378 (2021).

7. Doron, S. et al. Systematic discovery of antiphage defense systems in the microbial pangenome. *Science* **359**, eaar4120 (2018).

8. Millman, A. et al. An expanded arsenal of immune systems that protect bacteria from phages. *Cell Host & Microbe* **30**, 1556-1569 (2022).

9. Beavogui, A. et al. The defensome of complex bacterial communities. *Nature Communications* **15**, 2146 (2024).

10. Peng, Y. et al. Viruses in deep-sea cold seep sediments harbor diverse survival mechanisms and remain genetically conserved within species. *The ISME Journal* **17**, 1774-1784 (2023).

11. Georjon, H. & Bernheim, A. The highly diverse antiphage defence systems of bacteria. *Nature Reviews Microbiology* **21**, 686-700 (2023).

12. Bernheim, A. & Sorek, R. The pan-immune system of bacteria: antiviral defence as a community resource. *Nature Reviews Microbiology* **18**, 113-119 (2020).

13. Mayo-Munoz, D., Pinilla-Redondo, R., Birkholz, N. & Fineran, P.C. A host of armor: Prokaryotic immune strategies against mobile genetic elements. *Cell Reports* **42**, 112672 (2023).

14. Tock, M.R. & Dryden, D.T.F. The biology of restriction and anti-restriction. *Current Opinion in Microbiology* **8**, 466-472 (2005).

15. Lopatina, A., Tal, N. & Sorek, R. Abortive infection: bacterial suicide as an antiviral immune strategy. *Annual Review of Virology* **7**, 371-384 (2020).

16. Cheng, R. et al. Prokaryotic Gabija complex senses and executes nucleotide depletion and DNA cleavage for antiviral defense. *Cell Host & Microbe* **31**, 1331-1344 (2023).

17. Rousset, F. et al. A conserved family of immune effectors cleaves cellular ATP upon viral infection. *Cell* **186**, 3619-3631 (2023).

18. Hille, F. et al. The biology of CRISPR-Cas: backward and forward. *Cell* **172**, 1239-1259 (2018).

19. Saenz, J.S., Seifert, J. & Rios-Galicia, B. Antiviral defence systems in the rumen microbiome. *bioRxiv* (2024). doi: https://doi.org/10.1101/2024.04.04.588093.

20. Piel, D. et al. Phage–host coevolution in natural populations. *Nature Microbiology* **7**, 1075-1086 (2022).

21.   Tesson, F. et al. Systematic and quantitative view of the antiviral arsenal of prokaryotes.
Nature communications **13**, 2561 (2022).
22.   Rocha, E.P.C. & Bikard, D. Microbial defenses against mobile genetic elements and viruses:
Who defends whom from what? *PLoS Biology* **20**, e3001514 (2022).
23.   Picton, D.M. et al. The phage defence island of a multidrug resistant plasmid uses both BREX
and type IV restriction for complementary protection from viruses. *Nucleic Acids Research* **49**,
11257-11273 (2021).
24.   Wu, Y. et al. Bacterial defense systems exhibit synergistic anti-phage activity. *Cell Host &*
Microbe **32**, 557-572.e556 (2024).
25.   Mayo-Muñoz, D., Pinilla-Redondo, R., Camara-Wilpert, S., Birkholz, N. & Fineran, P.C.
Inhibitors of bacterial immune systems: discovery, mechanisms and applications. *Nature*
Reviews Genetics **25**, 237-254 (2024).
26.   Hampton, H.G., Watson, B.N.J. & Fineran, P.C. The arms race between bacteria and their
phage foes. *Nature* **577**, 327-336 (2020).
27.   Duan, N., Hand, E., Pheko, M., Sharma, S. & Emiola, A. Structure-guided discovery of anti-
CRISPR and anti-phage defense proteins. *Nature Communications* **15**, 649 (2024).
28.   Gussow, A.B. et al. Machine-learning approach expands the repertoire of anti-CRISPR protein
families. *Nature Communications* **11**, 3784 (2020).
29.   Serfiotis-Mitsa, D. et al. The structure of the KlcA and ArdB proteins reveals a novel fold and
antirestriction activity against Type I DNA restriction systems in vivo but not in vitro. *Nucleic*
Acids Research **38**, 1723-1737 (2010).
30.   Hobbs, S.J. et al. Phage anti-CBASS and anti-Pycsar nucleases subvert bacterial immunity.
Nature **605**, 522-526 (2022).
31.   Antine, S.P. et al. Structural basis of Gabija anti-phage defence and viral immune evasion.
Nature **625**, 360-365 (2024).
32.   An, L. et al. Global diversity and ecological functions of viruses inhabiting oil reservoirs.
Nature Communications **15**, 6789 (2024).
33.   Zhong, Z.-P. et al. Viral potential to modulate microbial methane metabolism varies by habitat.
Nature Communications **15**, 1857 (2024).
34.   Payne, L.J. et al. Identification and classification of antiviral defence systems in bacteria and
archaea with PADLOC reveals new system types. *Nucleic Acids Research* **49**, 10868-10878
(2021).
35.   Tesson, F. et al. A comprehensive resource for exploring antiphage defense: DefenseFinder
webservice, wiki and databases. *bioRxiv* (2024). doi:
https://doi.org/10.1101/2024.01.25.577194.
36.   Brito, I.L. Examining horizontal gene transfer in microbial communities. *Nature Reviews*
Microbiology **19**, 442-453 (2021).
37.   Liu, Y., Botelho, J. & Iranzo, J. Timescales and genetic linkage explain the variable impact of
defense systems on horizontal gene transfer *bioRxiv* (2024). doi:
https://doi.org/10.1101/2024.02.29.582795.
38.   Kogay, R., Wolf, Y.I. & Koonin, E.V. Defence systems and horizontal gene transfer in bacteria.
Environmental Microbiology **26** (2024).
39.   Li, M. et al. Toxin-antitoxin RNA pairs safeguard CRISPR-Cas systems. *Science* **372**,
eabe5601 (2021).

40.   Dupuis, M.-È., Villion, M., Magadán, A.H. & Moineau, S. CRISPR-Cas and restriction–modification systems are compatible and increase phage resistance. *Nature Communications* **4**, 2087 (2013).

41.   Dong, X. et al. Evolutionary ecology of microbial populations inhabiting deep sea sediments associated with cold seeps. *Nature Communications* **14**, 1127 (2023).

42.   Dong, X. et al. A vast repertoire of secondary metabolites potentially influences community dynamics and biogeochemical processes in cold seeps. *Science Advances* **10**, eadl2281 (2024).

43.   Kieft, K., Zhou, Z. & Anantharaman, K. VIBRANT: automated recovery, annotation and curation of microbial viruses, and evaluation of viral community function from genomic sequences. *Microbiome* **8**, 1-23 (2020).

44.   Guo, J. et al. VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses. *Microbiome* **9**, 1-13 (2021).

45.   Camargo, A.P. et al. Identification of mobile genetic elements with geNomad. *Nature Biotechnology* **42**, 1303–1312 (2023).

46.   Roux, S. et al. Minimum Information about an Uncultivated Virus Genome (MIUViG). *Nature Biotechnology* **37**, 29-37 (2019).

47.   Zhou, K. et al. Genomic and transcriptomic insights into complex virus-prokaryote interactions in marine biofilms. *The ISME Journal* **17**, 2303–2312 (2023).

48.   Yirmiya, E. et al. Phages overcome bacterial immunity via diverse anti-defence proteins. *Nature* **625**, 352-359 (2024).

49.   Yang, L. et al. Insights into the inhibition of type I-F CRISPR-Cas system by a multifunctional anti-CRISPR protein AcrIF24. *Nature Communications* **13**, 1931 (2022).

50.   Birkholz, N., Fagerlund, R.D., Smith, L.M., Jackson, S.A. & Fineran, P.C. The autoregulator Aca2 mediates anti-CRISPR repression. *Nucleic Acids Research* **47**, 9658-9665 (2019).

51.   Stanley, S.Y. et al. Anti-CRISPR-associated proteins are crucial repressors of anti-CRISPR transcription. *Cell* **178**, 1452-1464.e1413 (2019).

52.   Uribe, R.V. et al. Discovery and characterization of Cas9 inhibitors disseminated across seven bacterial phyla. *Cell Host & Microbe* **25**, 233-241 (2019).

53.   Suresh, S.K., Murugan, K. & Sashital, D.G. Enzymatic anti-CRISPRs improve the bacteriophage arsenal. *Nature Structural & Molecular Biology* **26**, 250-251 (2019).

54.   Knott, G.J. et al. Broad-spectrum enzymatic inhibition of CRISPR-Cas12a. *Nature Structural & Molecular Biology* **26**, 315-321 (2019).

55.   Müller, A.L., Kjeldsen, K.U., Rattei, T., Pester, M. & Loy, A. Phylogenetic and environmental diversity of DsrAB-type dissimilatory (bi)sulfite reductases. *The ISME Journal* **9**, 1152-1165 (2015).

56.   Zhang, C. et al. Unexpected genetic and microbial diversity for arsenic cycling in deep sea cold seep sediments. *npj Biofilms and Microbiomes* **9**, 13 (2023).

57.   Dong, X. et al. Phylogenetically and catabolically diverse diazotrophs reside in deep-sea cold seep sediments. *Nature Communications* **13**, 4885 (2022).

58.   Sheng, Y. et al. Insertion sequence transposition inactivates CRISPR-Cas immunity. *Nature Communications* **14**, 4366 (2023).

59.   Koonin, E.V., Makarova, K.S. & Zhang, F. Diversity, classification and evolution of CRISPR-Cas systems. *Current Opinion in Microbiology* **37**, 67-78 (2017).

60.   Martínez Arbas, S. et al. Roles of bacteriophages, plasmids and CRISPR immunity in

microbial community dynamics revealed using time-series integrated meta-omics. *Nature*
Microbiology **6**, 123-135 (2021).

61.   Sun, C.L., Thomas, B.C., Barrangou, R. & Banfield, J.F. Metagenomic reconstructions of
bacterial CRISPR loci constrain population histories. *The ISME Journal* **10**, 858-870 (2016).

62.   Guerrero, L.D. et al. Long-run bacteria-phage coexistence dynamics under natural habitat
conditions in an environmental biotechnology system. *The ISME Journal* **15**, 636-648 (2021).

63.   Huo, Y. et al. Structural and biochemical insights into the mechanism of the Gabija bacterial
immunity system. *Nature Communications* **15**, 836 (2024).

64.   Olm, M.R., Brown, C.T., Brooks, B. & Banfield, J.F. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes through de-
replication. *The ISME journal* **11**, 2864-2868 (2017).

65.   Parks, D.H. et al. GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic*
Acids Research **50**, D785-D794 (2022).

66.   Nguyen, L.-T., Schmidt, H.A., Von Haeseler, A. & Minh, B.Q. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and*
Evolution **32**, 268-274 (2015).

67.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic Acids Research* **49**, W293-W296 (2021).

68.   Breitwieser, F.P. & Salzberg, S.L. Pavian: interactive analysis of metagenomics data for
microbiome studies and pathogen identification. *Bioinformatics* **36**, 1303-1304 (2020).

69.   Hyatt, D. et al. Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 1-11 (2010).

70.   Zhang, C. et al. The majority of microorganisms in gas hydrate-bearing subseafloor sediments
ferment macromolecules. *Microbiome* **11**, 37 (2023).

71.   Xiao, X. et al. Metal-driven anaerobic oxidation of methane as an important methane sink in
methanic cold seep sediments. *Microbiology Spectrum* **11**, e05337-05322 (2023).

72.   Li, W.L. et al. Microbial ecology of sulfur cycling near the sulfate–methane transition of
deep☐sea cold seep sediments. *Environmental Microbiology* **23**, 6844-6858 (2021).

73.   Uritskiy, G.V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-
resolved metagenomic data analysis. *Microbiome* **6**, 1-13 (2018).

74.   Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211-3217 (2012).

75.   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast and
bias-aware quantification of transcript expression. *Nature Methods* **14**, 417-419 (2017).

76.   Nayfach, S. et al. CheckV assesses the quality and completeness of metagenome-assembled
viral genomes. *Nature Biotechnology* **39**, 578-585 (2021).

77.   Shaffer, M. et al. DRAM for distilling microbial metabolism to automate the curation of
microbiome function. *Nucleic Acids Research* **48**, 8883-8900 (2020).

78.   Roux, S. et al. iPHoP: An integrated machine learning framework to maximize host prediction
for metagenome-derived viruses of archaea and bacteria. *PLoS Biology* **21**, e3002083 (2023).

79.   Ye, J., McGinnis, S. & Madden, T.L. BLAST: improvements for better sequence analysis.
Nucleic Acids Research **34**, W6-W9 (2006).

80.   Coutinho, F.H. et al. RaFAH: Host prediction for viruses of Bacteria and Archaea based on

protein content. *Patterns* **2**, 100274 (2021).

81. Galiez, C., Siebert, M., Enault, F., Vincent, J. & Söding, J. WIsH: who is the host? Predicting
prokaryotic hosts from metagenomic phage contigs. *Bioinformatics* **33**, 3113-3114 (2017).

82. Wang, W. et al. A network-based integrated framework for predicting virus–prokaryote
interactions. *NAR Genomics and Bioinformatics* **2**, lqaa044 (2020).

83. Lu, C. et al. Prokaryotic virus host predictor: a Gaussian model for host prediction of
prokaryotic viruses in metagenomics. *BMC Biology* **19**, 5 (2021).

84. Couvin, D. et al. CRISPRCasFinder, an update of CRISRFinder, includes a portable version,
enhanced performance and integrates search for Cas proteins. *Nucleic Acids Research* **46**,
W246-W251 (2018).

85. Grandjean, M. Gephi: Introduction to network analysis and visualisation. *International AAAI*
Conference on Weblogs and Social Media (2015).

86. Yan, Y., Zheng, J., Zhang, X. & Yin, Y. dbAPIS: a database of anti-prokaryotic immune
system genes. *Nucleic Acids Research* **52**, 419-425 (2023).

87. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale using
DIAMOND. *Nature Methods* **18**, 366-368 (2021).

88. Mazzocco, A., Waddell, T.E., Lingohr, E. & Johnson, R.P. Enumeration of bacteriophages
using the small drop plaque assay system. *Bacteriophages: methods and protocols, volume 1:*
isolation, characterization, and interactions, 81-85 (2009).

89. Khedkar, S. et al. Landscape of mobile genetic elements and their antibiotic resistance cargo in
prokaryotic genomes. *Nucleic Acids Research* **50**, 3155-3168 (2022).

90. Neukirchen, S. & Sousa, F.L. DiSCo: a sequence-based type-specific predictor of Dsr-
dependent dissimilatory sulphur metabolism in microbial data. *Microbial Genomics* **7**, 000603
(2021).

91. Zhou, Z. et al. METABOLIC: high-throughput profiling of microbial genomes for functional
traits, metabolism, biogeochemistry, and community-scale functional networks. *Microbiome*
10, 33 (2022).

92. Tu, Q., Lin, L., Cheng, L., Deng, Y. & He, Z. NCycDB: a curated integrative database for fast
and accurate metagenomic profiling of nitrogen cycling genes. *Bioinformatics* **35**, 1040-1048
(2019).

93. Nakamura, T., Yamada, K.D., Tomii, K. & Katoh, K. Parallelization of MAFFT for large-scale
multiple sequence alignments. *Bioinformatics* **34**, 2490-2492 (2018).

94. Capella-Gutiérrez, S., Silla-Martínez, J.M. & Gabaldón, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973
(2009).

95. Chen, C. et al. TBtools-II: A "one for all, all for one" bioinformatics platform for biological
big-data mining. *Molecular Plant* **16**, 1733-1742 (2023).

96. Paysan-Lafosse, T. et al. InterPro in 2022. *Nucleic Acids Research* **51**, D418-D427 (2023).

97. Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M. & Barton, G.J. Jalview Version
2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189-
1191 (2009).

98. Mirdita, M., Steinegger, M. & Söding, J. MMseqs2 desktop and local web server app for fast,
interactive sequence searches. *Bioinformatics* **35**, 2856-2858 (2019).

99. Mirdita, M. et al. ColabFold: making protein folding accessible to all. *Nature Methods* **19**,

679-682 (2022).

100.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *nature* **596**, 583-
872   589 (2021).

873   101.   Abramson, J. et al. Accurate structure prediction of biomolecular interactions with AlphaFold
874   3. *Nature*, 1-3 (2024).

875   102.   Van Kempen, M. et al. Fast and accurate protein structure search with Foldseek. *Nature*
876   *biotechnology* **42**, 243-246 (2024).

877   103.   DeLano, W.L. Pymol: An open-source molecular graphics tool. *CCP4 Newsl. Protein*
878   *Crystallogr* **40**, 82-92 (2002).

Acknowledgements

The work was supported by National Science Foundation of China (No. 92351304, No. 42376115, No. 42406109 and No. 32100025), Natural Science Foundation Project of Xiamen City (No. 3502Z202373076), Natural Science Foundation of Fujian Province (No. 2023J06042), Scientific Research Foundation of Third Institute of Oceanography, MNR (No. 2022025 and No. 2023022), China Postdoctoral Science Foundation (No. 2023M734096) and Open Funding Project of State Key Laboratory of Microbial Metabolism (No. MMLKF23-05).

Author contributions

X.D. conceived and designed the study. Y.H. and J.L. performed the omics analyses. C.L. contributed to viral identification and classification. R.C. designed and conducted the experimental implementations. X.L. participated in the analysis of some metagenomic data. C.L., F.X., J.P. and W.X. actively participated in discussions and data interpretation. Y.H., J.L. and X.D. wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

 Figure Legends

 Figure 1. Distribution and activities of defense systems in cold seep prokaryotes.

 (a) Types and proportions of defense systems and candidate defense systems in cold

 seep microbial genomes, including their primary mechanisms of action and

 corresponding system counts. **(b)** Plaque assays demonstrating phage infection of

 Escherichia coli B (ATCC® 11303™) transformed with plasmids carrying various

 candidate defense systems or an empty vector (control). Ten-fold serial dilutions of

 phages T7, T5, and T4 were spotted onto bacterial lawns. "DS" denotes defense

 systems. Detailed data on annotated defense systems are provided in **Supplementary**

 Tables 1-3.

 Figure 2. Characteristics of defense systems in cold seep prokaryotes. (a)

 Proportion of metagenome-assembled genomes (MAGs) containing defense systems

 among cold seep prokaryotes, and the proportions of various types of defense systems

 within all prokaryotes, archaeal, and bacterial MAGs. **(b)** Distribution and average

 number of defense systems (DS) per MAG. **(c)** Relationship between the number and

 density of defense systems (DS per Mb per MAG) and MAG genome size (Mb).

 Figure 3. Abundance and distribution of defense systems in cold seep

 prokaryotes. Phylogenetic representation of archaeal **(a)** and bacterial **(b)** MAGs,

 categorized by their corresponding phyla. Shown are the density of defense systems

 (DSs per Mb, blue), defense genes (DGs per Mb$\times10^{-1}$, orange), candidate defense

 systems (CDs per Mb$\times10^{-1}$, green), and the number of defense islands (DIs, red).

 Circles and triangles denote MAGs with defense systems located on plasmids and

 proviruses, respectively. The outer bar charts display the genome sizes of MAGs (Mb).

 Detailed statistics for defense systems in each MAG are provided in **Supplementary**

 Table 3.

 Figure 4. Interactions among defense systems in cold seep prokaryotes. (a)

 Correlations among 34 defense systems in cold seep prokaryotes. Overall correlations

are displayed on the left (blue), while correlations within defense islands are on the right (green). Pearson correlation coefficients were calculated based on system abundance; coefficients are color-coded, and larger squares indicate stronger correlations. Significance levels are denoted by asterisks: $* P \leq 0.05$, $** P \leq 0.01$, $*** P \leq 0.001$. Mechanisms of action for each defense systems are indicated by color: nucleic acid degradation (green), inhibition of replication (blue), abortive infection (pink), and unknown (red). **(b)** Genomic collinearity of eight contiguous fragments containing defense islands. Defense genes are highlighted in green, candidate defense genes in yellow, and associated genes are color-coded: transport-related genes (orange), mobile genetic elements (MGEs; pink), other annotated functional genes (purple), and unknown functions (gray). Detailed statistics for correlation are provided in **Supplementary Tables 6 and 8**.

Figure 5. Relative abundance and expression of microbial defense systems in cold seep sediments. (a) Relative abundance (genes per million, GPM) and transcript abundance (transcripts per million, TPM) of the top ten defense systems across different cold seep sediment samples. Different colors represent different types of defense systems. Statistical differences among defense system types were assessed using the Kruskal–Wallis rank-sum test. **(b)** The heatmap displays the average gene abundance of various defense systems within each phylum in cold seep sediments, expressed in genes per million (GPM). Gray squares indicate a GPM value of zero. The bar graph illustrates the average relative abundance of defense systems across phyla. Green labels represent bacterial phyla, while orange denotes archaeal phyla; different defense mechanisms are distinguished by varying colors. **(c)** Gene abundance of defense system genes at different sediment depths in cold seeps, measured in GPM. **(d)** Expression levels of defense system genes at different sediment depths in cold seeps, measured in TPM. Sediment depths are categorized into three groups: surface (<1 meter below seafloor [mbsf]), shallow (1−10 mbsf), and deep (>10 mbsf).

Figure 6. Interactions between cold seep viral anti-defense systems and host defense systems. (a) The network diagram illustrates 2,688 host-virus pairs identified in cold seeps. prokaryotes are colored according to their phyla. **(b)** Anti-defense genes identified in cold seep viruses, showing the types of anti-defense systems and corresponding viral taxonomy. **(c)** Illustration of 155 host–virus pairs where host prokaryotes contain defense genes and the corresponding viruses possess anti-defense genes. **(d)** Genomic organization of viral gene clusters containing multiple types of anti-defense systems, along with the corresponding host defense systems. Viral genes are categorized and color-coded: anti-defense genes (yellow), metabolic genes (blue), structural genes (brown), attachment genes (pink), and unknown genes (grey). Host defense genes are highlighted in green, transport-related genes in orange, and other functional genes in purple. Green lines indicate interactions between host defense systems and viral anti-defense systems. **(e)** Structural comparison of the AcrIF24 protein dimer from a *Caudoviricetes* virus infecting *Halomonas* sp. (blue) with the known cryo-electron microscopy structure from *Pseudomonas aeruginosa* (pink). A detailed view highlights key sites involved in AcrIF24 protein dimerization and binding to the Csy complex.

Figure 7. Defense competition between key metabolic microbial groups in cold seeps and their viruses. (a) Types of defense systems in key metabolic microbial groups, including ANME, SRB and diazotrophs. Colored squares indicate the presence of key functional genes, while colored circles indicate the presence of defense systems. Different phyla are indicated by distinct colors, and defense mechanisms are represented by circles of different colors. **(b)** Relationship between the number of defense systems and the diversity of defense types in key metabolic microbial groups, categorized into three groups based on quantity: few systems, diverse systems, and similar systems. **(c)** Genomic organization of the Gabija defense system and the viral *gad1* gene in a virus–host pair involving ETH-SRB1 and a *Caudoviricetes* virus. Genes are color-coded: anti-defense genes (green), structural genes (purple), metabolic genes (pink), attachment genes (orange), transport genes

(yellow), defense genes (blue), and genes of unknown function (gray). Green lines

indicate interactions between the host defense system and the viral anti-defense

system. **(d)** The structure of GajAB complex in cold seep ETH-SRB1 predicted by

Alphafold3, including detailed diagrams of the main interaction sites between GajA

proteins and between GajB proteins. **(e)** The structure of Gad1 protein in a cold seep

Caudoviricetes virus predicted by Alphafold3.

a

others(17.60%)

Tiamat(0.90%)
Pycsar(0.93%)
DRT(1.05%)
SEFIR(1.17%)
VSPR(1.18%)
Wadjet(1.28%)
Gabija(1.61%)
Septu(1.85%)
AbiL(1.94%)
CBASS(2.19%)
viperin(2.59%)
dXTPase(3.98%)
AbiE(6.30%)
SoFic(7.24%)

Defense Systems

RM(38.28%)

Cas(9.90%)

DS(37.40%)
9,869

Candidate(62.60%)
16,520

others(19.15%)

HEC-01(1.39%)
HEC-09(1.47%)
PDC-S03(2.45%)
PDC-M05(2.64%)
PDC-S07(2.68%)
PDC-S13(3.37%)
PDC-S02(4.88%)
PDC-S27(5.15%)
PDC-S70(7.26%)

Candidate Systems

PDC-S01(31.57%)

PDC-S09(9.35%)

PDC-S05(8.63%)

1,864

5,416

1,867

System num

Type num

Mechanisms

Nucleic acid degradation   Abortive infection

Inhibition of replication   Unknown

b

	Phage T7	Phage T5	Phage T4
Empty vector			
PDC-S05			
PDC-S70			
PDC-M05			
HEC-01			

a

Tree scale: 0.5

Phylum
- Aenigmatarchaeota
- Altiarchaeota
- Asgardarchaeota
- B1Sed10-29
- EX4484-52
- Hadarchaeota
- Halobacteriota
- Hydrothermarchaeota
- Iainarchaeota
- Methanobacteriota_B
- Micrarchaeota
- Nanoarchaeota
- SpSt-1190
- Thermoplasmatota
- Thermoproteota
- Undinarchaeota

Density DSs (Mb⁻¹)
- 0
- 5
- 10

Density DGs (Mb⁻¹)(10⁻¹)
- 0
- 1.5
- 3

Density CDs (Mb⁻¹)(10⁻¹)
- 0
- 1.5
- 3

DI Number
- 0
- 1

bootstrap
- ≥ 90

DG on MGE
- DG on plasmids
- DG on proviruses

b

Tree scale: 0.5

Phylum
- Acidobacteriota
- Actinomycetota
- Aerophobota
- Atribacterota
- Bacteroidota
- Campylobacterota
- Chloroflexota
- Desulfobacterota
- Desulfobacterota_F
- Gemmatimonadota
- Krumholzibacteriota
- Myxococcota
- Omnitrophota
- Patescibacteria
- Planctomycetota
- Pseudomonadota
- Spirochaetota
- Verrucomicrobiota
- WOR-3
- Zixibacteria
- Others

Density DSs (Mb⁻¹)
- 0
- 7.5
- 15

Density DGs (Mb⁻¹)(10⁻¹)
- 0
- 1.6
- 3.2

Density CDs (Mb⁻¹)(10⁻¹)
- 0
- 1.1
- 2.2

DI Number
- 0
- 1
- 2

bootstrap
- ≥ 90

DG on MGE
- DG on plasmids
- DG on proviruses