1

The effects of iron limitation on the small chlorophyte *Micromonas* from the Northeast Pacific Ocean

Meredith G. Meyer[1], Vincent J. White[1], Olivia Torano[1], Heidi Hannoush[1], Margarita Lankford[1], and Adrian Marchetti[1]

[1]Department of Earth, Marine, and Environmental Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Corresponding Author: A. Marchetti (amarchetti@unc.edu)

The authors declare no competing interests.

Abstract

Small eukaryotic phytoplankton can account for a considerable amount of biomass and primary production in high nutrient, low chlorophyll (HNLC) regions of the ocean where iron-limitation is pronounced. However, the physiological and metabolic strategies these cells invoke to cope under low iron conditions and the extent to which they are responsible for new production (i.e., the fraction of primary production supported by nutrients from outside of the euphotic zone) is unclear. Here, we examine how a representative picoeukaryote, the chlorophyte *Micromonas* sp., recently isolated from the iron-limited subarctic Northeast Pacific Ocean, responds to iron-limitation when grown on nitrate as a nitrogen source. Iron-limited *Micromonas* exhibits reductions in growth rate, cell volume, and elemental quotas along with a restructuring of cellular metabolism. Gene expression and pathway analyses show evidence of strategies to mitigate iron limitation with constitutive expression of genes related to nitrogen uptake and utilization. Additionally, cellular carbon and nitrogen quotas ranged from $20 - 170$ fmol C cell^{-1} and $3.3 - 20$ fmol N cell^{-1}, respectively, as a function of iron status. Based on the measured cellular quotas, we estimate that representative picoeukaryotes (<2 μm), such as *Micromonas,* in HNLC Northeast Pacific waters can account for a significant proportion of new production, supporting the need for a reconsideration of the role small eukaryotic phytoplankton play in the global carbon cycle.

1. Introduction

In open-ocean ecosystems, high surface area-to-volume ratios and generally lower nutrient quotas make picophytoplankton strong competitors for nutrients, thus enabling them to routinely dominate eukaryotic phytoplankton communities in these oligotrophic environments (Worden and Not, 2008). While much research has focused on cyanobacteria, specifically *Prochlorococcus* (Chisholm et al., 1991) and *Synechococcus* (Moore et al., 2002), fewer studies have investigated the growth strategies of picoeukaryotes which have been shown to potentially play key ecological roles (Fowler et al., 2020). Previous studies have suggested chlorophytes and other small eukaryotic groups primarily rely on ammonium and other forms of recycled nitrogen to support their growth (Greene et al., 1992; Marchetti et al, 2006; Meyer et al., 2022). Compared to cyanobacteria, diatoms, and dinoflagellates, these picoeukaryotes are poorly characterized and are less likely to be available in culture. Thus, there are fewer monocultured representatives sequenced for genomic and/or transcriptomic analyses compared to other phytoplankton groups (Le Gall et al., 2008). This overall lack of knowledge has led to these organisms commonly being overlooked in whole community analyses despite their high abundances.

Given the potentially outsized role these eukaryotic picophytoplankton could play in oligotrophic and nutrient-limited regions, understanding the nutrient acquisition strategies, requirements and metabolisms employed by these phytoplankton and the resultant effect for general ecosystem productivity and carbon export potential is of substantial importance. Here we present measurements obtained from the chlorophyte, *Micromonas* sp., that was recently isolated from Ocean Station Papa (Station P) in the subarctic Northeast Pacific during the NASA EXport Processes in the Ocean from RemoTe Sensing (EXPORTS) program in 2018. *Micromonas* sp. is an abundant member of the *Mamiellales* order and is closely related to *Ostreococcus* and *Bathycoccus* (Monier et al., 2016; Vannier et al., 2016), two other

ecologically important picoeukaryotic chlorophytes. Studies have shown that *Micromonas*

can occupy a wide range of habitats, from the tropics to the poles as well as offshore and

coastal environments (Demory et al., 2018). The group often exhibits an "occupy-abundance

pattern" where *Micromonas*, and other members of the *Mamiellophyceae* lineage, frequently

become the most abundant genera in regions they are present (Monier et al., 2016). Yet, the

metabolic capabilities and the ecological role of *Micromonas* in iron-limited regions is still

poorly understood.

Station P is the site of a long-term oceanographic monitoring program which resides

within a High Nutrient (predominantly nitrate (NO_3^-)), Low Chlorophyll, or HNLC, region.

This region is chronically limited by the bioavailability of iron which, despite replete

macronutrient concentrations, inhibits substantial phytoplankton biomass standing stock and

higher rates of primary production (Martin et al., 1988; Marchetti et al., 2006). Iron is used as

a crucial co-factor in photosynthesis and nitrogen assimilation (LaRoche et al.,1993; Peers

and Price, 2006). Many groups of phytoplankton that experience iron limitation have been

shown to utilize regenerated nitrogen sources (e.g., ammonium (NH_4^+) or urea) which

theoretically have a lower iron requirement relative to oxidized forms of nitrogen or nitrogen

gas (Price et al., 1994; Varela and Harrison, 1999). In addition, recent findings suggest that

the most dominant ecotype of the cyanobacteria in the Northeast Pacific, *Synechococcus*,

lacks the ability to assimilate NO_3^-, thus being restricted to regenerated production (i.e., using

recycled nitrogen sources such as ammonium and urea) (Sharpe et al., 2023). This finding

presents a knowledge gap in our understanding of which organisms are responsible for the

vast majority of new production in the region, which has routinely been measured to be

driven predominantly by small phytoplankton groups (i.e., < 5 μm in cell diameter) (Meyer et

al. 2022).

Here, we sought to examine why *Micromonas* can be found in high abundances in the

Northeast Pacific and their possible role in carbon export. We had two primary research

objectives: a) examine the distinct physiological and metabolic strategies of *Micromonas* to

grow under low iron conditions and b) quantify the extent to which *Micromonas* can

contribute substantially to new production in this region. We performed these objectives

through determining rates of cellular growth, photophysiology, carbon and nitrogen cellular

content, and trends in gene expression as a function of iron status. The combination of

molecular and physiological analyses provides a more wholistic approach to examining

ecologically important phytoplankton groups and their influence on marine biogeochemical

cycles.

2. Methods

2.1 Sampling and Cell Isolation

Whole seawater was collected from approximately 5 m depth via a trace metal clean

rosette during the NASA EXPORTS field campaign to Ocean Station Papa (Station P) in the

subarctic North Pacific. This campaign occurred from August 16[th] to September 7[th], 2018. To

keep storage conditions as similar to the ambient environment as possible, sampled seawater

was placed in culture flasks and incubated in on-deck incubators before being transported

back to the University of North Carolina for phytoplankton isolation. An abundant, small (< 5

µm) chlorophyte was isolated via single-cell isolation using a 10 µL pipette. DNA was

extracted using standard protocols with a DNEasy Plant Mini Kit (Qiagen). The V4 region of

the 18S rRNA gene was sequenced (National Center for Biotechnology Information (NCBI)

ID PRJNA1072555), and the consensus sequence of forward and reverse reads was searched

against the NCBI repository using BLAST, revealing the highest sequence similarity to

Micromonas pusilla (accession number KY980282.1) with a 100% percent ID and 76%

query cover.

2.2 Culture conditions

139   *Micromonas sp*. (UNC1837) cells were maintained in Aquil media in an incubator

under constant 120-150 μmol photons m^{-2} s^{-1} light at 12ºC to maintain light saturation and

ambient temperatures close to environmental conditions at Station P. To examine the effects

of differing nitrogen forms under variable iron states on *Micromonas* growth and physiology,

four trace metal clean culture treatments were initiated. Aquil medium was sterilized in a

microwave, cooled, and supplemented with filter-sterilized (0.2-μm Acrodisc)

ethylenediamine tetraacetic acid (EDTA) trace metals (minus iron), vitamins (B_{12}, thiamine,

and biotin), and chelexed macronutrients (NO_3^-, phosphate) in accordance with Aquil

medium concentrations (Price et al. 1989). Trace-metal concentrations were buffered using

100 μM of EDTA. Two treatments contained NO_3^- (300 μM) and two treatments contained

NH_4^+ (30 μM) as their nitrogen source. Within each nitrogen treatment, there was an iron

replete (1370 nM total Fe) and an iron deplete (3.1 nM total Fe) treatment. Iron

concentrations were selected after initial testing, and the chosen concentrations were able to

achieve the most consistent growth rates after a minimum of three transfers with transfers

occurring approximately every 7-10 days. Thus, the four treatments can be described as NO_3^-

Fe+, NO_3^- Fe-, NH_4^+ Fe+, and NH_4^+ Fe-. Each treatment was first cultured in triplicate acid-

washed 28 mL clear, polycarbonate centrifuge tubes (Nalgene, Rochester, NY) for

acclimation to their respective treatments. Cells were acclimated to their treatment for a

minimum of three transfers before experiments began. During acclimation, 200 μL of dense

late exponential-phase culture was transferred to new tubes with fresh media to maintain

growth, thus representing a 125-fold dilution. Following acclimation, cultures were

transferred to a trace metal clean, 1 L polycarbonate bottle that was approximately 700 mL

full. We were not able to maintain steady state growth rates in NH_4^+ Fe+ and NH_4^+ Fe-

cultures, thus confining our experiment to the NO_3^- Fe+ and NO_3^- Fe-, referred to as Fe+ and

Fe- treatments herein. All culture preparation, maintenance, and cell transfers were conducted

in a trace metal clean lab under a positive-pressure, laminar flow hood.

During the experiments, approximate cell growth and concentrations were monitored

via relative fluorescence units (RFUs) on a Turner 10AU fluorometer from lag phase through

stationary phase (Brand et al. 1981). Once samples were determined to be in mid-exponential

phase, cultures were measured for RFUs and other photophysiological parameters using a

Satlantic FIRe fluorometer with blue light diode (450 nm) and 60 iterations per triplicate

measurement (see below) and harvested for the following parameters: cell counts,

chlorophyll a (Chl a) concentration, particulate carbon (C) and nitrogen (N) concentration,

and RNA.

Specific growth rates (μ) were calculated as the slope of the natural log of relative

fluorescence units over time, according to Brand et al. (1981). Assuming Fe+ represents a

maximum growth rate (μ_{max}), the ratio of average μ/μ_{max} (i.e., Fe-/Fe+) can be calculated.

The maximum quantum yield of photochemistry in photosystem II (PSII; F_v/F_m) and the

functional absorption cross section of PSII (σ_{PSII}) were measured as metrics to evaluate

photosynthetic efficiency (Gorbunov and Falkowski; 2004).

Chl a was analyzed fluorometrically following Graff and Rynearson (2011).

Particulate carbon (PC) and nitrogen (PN) were analyzed on a Carlo Erba NC 2500 elemental

analyzer at University of Maryland Center for Environmental Science Appalachian

Laboratory in Frostburg, Maryland. Cultures for RNA were filtered onto an Isopore 0.4 μm

pore size, 47 mm polycarbonate filter under gentle vacuum pressure, flash frozen in liquid

nitrogen, and stored at -80 ºC before extraction. RNA was extracted using standard protocols

with a RNAqueous-4PCR kit according to the manufacturer's protocols. Extracted RNA was

checked for quality (A280/260, A260/230, and nucleic acid concentration) on a Nanodrop

spectrophotometer and cleaned using a RNeasy MinElute Clean-up kit. RNA was sent to the

Azenta Genewiz Sequencing Facility in South Plainfield, New Jersey for Illumina high

throughput analysis with ~350M raw paired-end reads per lane and single index, 2 x 150 bp

per lane, resulting in a read depth of 62-65 million reads per sample.

2.3 Cell counts and size estimates

193   A small aliquot (1.8 mL) of each culture (both treatments and all replicates) were

preserved in 200 µL of paraformaldehyde in 2 mL cryovials at the time of cell harvest, the

day following the harvest and several (2-3) days preceding the harvest. Cryovials were

immediately flash frozen in liquid nitrogen and stored at -80 ºC until they were analyzed on a

Guava easyCyte flow cytometer equipped with a red laser in the Paerl Lab at North Carolina

State University. Data were processed by InCyte software to provide cell counts based on red

fluorescence. Estimates of cell size (diameter) were also calculated using calibration beads of

known size based on the relationships of Paerl et al. (2020). Cell volume was calculated by

assuming cells were spherical. Cellular Chl *a*, PC and PN concentrations were calculated by

normalizing concentrations by cell counts and by cell biovolume.

2.4 North Pacific new production calculations

205   If we assume *Micromonas* is a good representative (i.e., in the top 20% most

represented species in 18S rRNA gene community composition data; Jones and Rynearson,

2022) of the picoeukaryote population at Station P, upper bound estimates of the contribution

of picoeukaryotes to total new production under Fe-replete and Fe-limiting conditions can be

calculated according to Equation 2:

$New\ production = cell\ concentration * \mu * N\ quota$ (Eq. 1)

Where the nitrogen quota is in units of fmol N cell^{-1}. Here, the N quota and growth rate (μ; d^{-}

^{1}) come from our laboratory results with the *Micromonas* culture. The cell concentration used

in Fe+ and Fe- culture new production rates is the picoeukaryote cell counts (3.3 x 10^6 cells

L^{-1}) measured from the EXPORTS Northeast Pacific cruise in 2018 (Graff, 2023). These are

average picoeukaryote counts collected from surface waters and measured via flow cytometry

during the observation period.

2.5 Gene expression analysis

Transcripts were trimmed of sequence adapters and poly-A tails via Trim_galore

(v.0.6.2; Altos Labs) and underwent quality control analysis via FastQC (v.0.11.9) where

sequences with a quality score below 20 and/or length less than 50 base pairs were removed.

Merged sequence files were assembled de novo via Trinity (v.2.8.6) before being combined

into grand assemblies based on iron treatment via cdhit (v.4.8.1; Fu et al., 2012). Sequences

underwent functional annotation via the Kyoto Encyclopedia of Genes and Genomes (KEGG;

Release 88.2). Additional manual annotation of genes associated with Fe metabolism, but are

not included in KEGG (e.g., protin-pumping rhodopsins, iron starvation induced proteins

(ISIPs), and phytotransferrins), was conducted (see Moreno et al., 2017 for more information

on manual annotation and sequence acquisition). Independent end-to-end read alignment

occurred via Salmon (v.10.9.1; Patro et al., 2017) and annotation and count files were

combined via tximport, providing counts per million (CPM) values per gene per sample

(Soneson et al., 2015). Differential expression of genes was analyzed at the KEGG Ortholog

(KO) level, averaging values for multiple contigs with the same KO, according to the

DESeq2 package in RStudio (v.4.2.2). The Log2Fold change (adjusted p-value < 0.05) of

specific genes of interest, particularly those associated with photosynthesis, N metabolism,

and Fe metabolism, were investigated. Additionally, trends in KEGG metabolic pathways

were analyzed among samples. Figures were generated in Matlab R2022b. Sequences from

this analysis are available at the National Center for Biotechnology Information (Ascension

numbers: SAMN39858273 - 284).

3. Results

3.1 Growth rates and photosynthetic efficiency

The Fe+ cultures had an average growth rate of 0.96 ± 0.01 d^{-1} (Fig. 1A). The average

growth rate of Fe- cultures was substantially lower at 0.48 ± 0.07 d^{-1} resulting in a μ/μ_{max}

(Fe-/Fe+) of 0.50, suggesting substantial limitation of iron in the Fe- treatment relative to the

Fe+ treatment (Fig. 1A). Estimates of maximum photochemical yield of PSII (F_v/F_m) confirm

that the Fe+ cultures had an increased efficiency of PSII compared to Fe-, which was

experiencing iron-limited conditions. The mean F_v/F_m for the Fe+ treatment was 0.73 ± 0.02,

whereas values for the Fe- cultures were 0.37 ± 0.03, consistent with a decline in

photosynthetic efficiency of PSII due to iron limitation (Fig. 1B). The cross-sectional

absorption area of photosystem II (σ_{PSII}) was also consistent with other iron-limited

phytoplankton cultures with an increase in σ_{PSII} under iron-limited conditions (Ellis, 2015;

Moreno et al., 2020). For Fe- cultures, σ_{PSII} averaged 1036 ± 264, and for Fe+, σ_{PSII} averaged

372 ± 12 (data not shown), resulting in a 2.8-fold increase under iron-limited conditions.

There was a substantially lower F_v/F_m value for the Fe+ replicate A sample compared

to the other replicates, suggesting that this culture may have entered stationary phase,

experiencing N depletion at the time of sampling. This triplicate was therefore removed from

the averages of Fe+ sample data presented in Figures 1 and 2 as well as from the gene

expression analysis. Although with all replicates included, there was still a significant

difference in growth rates, cellular Chl *a* concentrations and cellular C and N quotas between

the Fe+ and Fe- treatments (Supplemental Table 1).

3.2 Cell Size and chlorophyll a *content*

The Fe+ cultures had an average cell diameter of 1.85 ± 0.02 µm (data not shown)

and corresponding cell volume of 3.35 ± 0.10 µm^3 (Fig. 1C). The Fe- cultures had an average

cell diameter of 1.58 ± 0.00 µm and cell volume of 2.07 ± 0.00 µm^3 (Fig. 1C).

Cellular Chl *a* contents exhibited an order of magnitude difference between the Fe+

and Fe- samples, with the Fe+ cultures having an average of 6.42 ± 0.50 fg cell^{-1} and the Fe-

cultures averaging only 0.53 ± 0.09 fg cell^{-1}. Chl *a* concentration, when normalized to cell

biovolume, exhibited a similar trend to cellular Chl *a* content, but the difference in the

magnitude between the Fe+ and Fe- cultures was substantially lower (Fig. 1D). The average

biovolume-normalized Chl *a* content was 1.92 ± 0.09 fg µm^{-3} in the Fe+ treatment and 7.5-

fold lower at 0.25 ± 0.04 fg µm^{-3} in the Fe- treatment.

3.4 Cellular carbon and nitrogen quotas

Like Chl *a*, the cellular C content exhibited an order of magnitude difference between

the Fe+ and Fe- treatments. The Fe+ cultures had a high cellular C quota of 158 ± 20 fmol C

cell^{-1} whereas the Fe- cultures had an 8-fold lower C quota of 23 ± 0.0 fmol C cell^{-1} (Fig. 2A).

As is seen throughout the parameters measured, the Fe- sample quotas were consistent, only

ranging from 22 – 24 fmol C cell^{-1} among triplicates. When normalized to cell volume, the

biovolume-normalized C quotas were more similar between treatments due to the cells in the

Fe+ culture being ~1.5-fold larger than the cells in the Fe- culture. The Fe+ treatment had a C

quota of 47 ± 3.8 fmol C µm^{-3} whereas the quota for the Fe- treatment was 11.1 ± 0.38 fmol

C µm^{-3} (Fig. 2B). Thus, when considering the difference in cell volumes (i.e., calculated per

μm^{-3} rather than per cell), the difference in C quota decreases from 8.0- to 4.3-fold higher in

the Fe+ treatment relative to the Fe- treatment.

The cellular N contents likewise exhibited orders of magnitude differences between

iron treatments when normalized per cell and per unit volume. The average N quota for the

Fe+ cultures was 21.1 ± 1.5 fmol N cell^{-1} whereas the average for the Fe- cultures was $3.37 \pm$

0.13 fmol N cell^{-1} (Fig. 2C). The per unit volume average for Fe+ samples was 6.30 ± 0.26

fmol N μm^{-3}, and for the Fe- samples was $1.62 \pm 6.44 \times 10^{-2}$ fmol N μm^{-3} (Fig. 2D). The

change in cellular N contents as a function of iron status was similar to that of C at

approximately 6- and 4-fold larger for the Fe+ treatment when normalized per cell and per

unit volume, respectively. Both treatments exhibited slightly higher particulate carbon :

nitrogen (C:N) ratios than the Redfield ratio of 6.6 with averages of 7.49 ± 0.30 and $6.84 \pm$

0.05 for Fe+ and Fe-, respectively (Fig. 2E). C:Chl *a* ratios exhibited the opposite pattern

with a higher average ratio (537 ± 111 µg : µg) in the Fe- treatment than in the Fe+ treatment

(296 ± 9.74 µg : µg; Fig. 2F).

3.5 Potential contributions to new production in the NE Pacific

The average picoeukaryote cell concentration during the EXPORTS observation

period used in these calculations was 3.34×10^6 cells L^{-1} (Graff, 2023). Based on these

picoeukaryote cell densities, rates of potential contributions to new production varied by an

order of magnitude between iron treatments, averaging 5.4 ± 1.0 nmol N L^{-1} d^{-1} for Fe-

cultures and 67.5 ± 4.3 nmol N L^{-1} d^{-1} for Fe+ cultures (Fig. 3). Rates were highly variable,

with Fe- values ranging from 3.7 to 8.4 nmol N L^{-1} d^{-1} and Fe+ values ranging from 45.9 to

105.1 nmol N L^{-1} d^{-1}. Rates of small-celled new production at Station P were always between

the Fe- and Fe+ rates, except for on YD 240 when the EXPORTS rate was higher than Fe+

(55.4 nmol N L^{-1} d^{-1}) at 58.5 nmol N L^{-1} d^{-1}.

3.6 Gene expression

Molecular sequencing of the *Micromonas* sp. (UNC1837) transcriptome indicated a higher GC content for the Fe+ (61%) vs. Fe- (49%) cultures. Total sequences and sequence length were comparable between treatments for the Fe+ and Fe-, respectively.

The impact of Fe availability was analyzed broadly on the metabolic pathway level (Fig. 4A) and specifically on a targeted gene level (Fig. 4B). Of the 17 metabolic pathways of interest analyzed, on average, 10 displayed overrepresentation in Fe- treatment (i.e., negative Log2Fold change). Key pathways exhibiting this trend include those related to iron limitation mitigation strategies or general physiological stress responses, including iron starvation induced protein 3 (ISIP3) expression, ferredoxin expression, higher expression of photosystem I relative to photosystem II, and zeaxanthin epoxidase (Fig. 4A). Unsurprisingly, many gene transcripts involved in pathways related to general cell maintenance, including nitrogen metabolism, arginase, flavin adenine dinucleotide synthesis, etc., were overrepresented in the Fe+ treatment.

Evaluating these physiological trends further, our findings appear consistent with those previously reported for the closely-related chlorophyte, *Ostreococcus* (Lelandis et al., 2016), with *Micromonas* appearing to contain far fewer transcribed genes involved in nitrogen and iron metabolism than are commonly reported in diatoms and other phytoplankton taxa. Transcripts for the nitrogen assimilation genes nitrate/nitrite transporters (*NRT*), nitrate reductase (*NR*), nitrite reductase (*NiR*), ammonium transporters (*AMT*), and urea transporters (*Urt*) were identified in our *Micromonas* isolate (Fig. 4B, 4C). Most genes associated with NO_3^- uptake (*NRT, NiR*) increased expression (Log2Fold changes > 0) in the Fe+ treatment relative to Fe- treatment (Fig. 4B). Urea transporters were also overrepresented (Log2Fold change = 4.21; adjusted p-value < 0.01) under Fe+ conditions. Conversely,

ammonium transporters showed a slight negative Log2Fold change (-0.47), suggesting

increased expression in the Fe- treatment (Fig. 4B, 4C). Overall, genes associated with the

expression of photosystems I and II (*Psa*, *Psb*) exhibited a stronger Log2Fold change (-2.93)

in Fe- conditions, in contrast to the common assumption that these photosynthetic genes

should be more highly expressed under iron-replete conditions (Fig. 4B).

Our data suggests that *Micromonas* may lack genes commonly involved in iron

regulation and homeostasis in diatoms residing in HNLC regions, including proton-pumping

rhodopsins and ferritin (Marchetti et al., 2009; Andrew et al., 2023). However, we did

identify the gene for iron starvation induced protein 3 (*ISIP3*) which was strongly expressed

(Log2Fold change = -4.62; adjusted p-value <0.01) under Fe- conditions (Fig. 4B). The

percent of overall contigs without a functional annotation was high (87.9%), suggesting

additional genes involved in both nitrogen and iron metabolism may be present but are not

currently identified in the KEGG database.

4. Discussion

4.1 Cellular and metabolic characterization

While less is known about the ecology and physiology of *Micromonas* relative to

other picoeukaryotic chlorophytes such as *Ostreoccocus*, our findings reported here are

consistent with and build upon some of what has been reported on *Micromonas* (Cochlan and

Harrison, 1991a; 1991b). Our estimates of cell size are similar to the previously reported

values of <2 μm (McKie-Krisberg and Sanders, 2014). Additionally, *Ostreoccocus* has been

reported to exhibit fast growth rates under nitrogen replete conditions (μ_{max} of 1.11 d^{-1};

Cochlan and Harrison, 1991a), consistent with our measured growth rates for our

Micromonas sp. isolate in iron-replete conditions. However, a substantial lag phase has not

been previously reported. Additionally, our inability to achieve steady state growth in our

isolate using NH_4^+ media is noteworthy. In fact, following nitrogen starvation, Cochlan and

Harrison (1991a; 1991b) reported a preference for NH_4^+ and urea uptake relative to NO_3^-

uptake. While we were unable to identify what was inhibiting growth, it is likely that our

Micromonas sp. requires specific conditions for NH_4^+ utilization that were not met within our

culture conditions. Our findings suggest that our strain of *Micromonas* may exhibit a

preference for NO_3^- over other forms of nitrogen (such as NH_4^+), but further research is

warranted.

Our study quantified changes in cellular Chl *a* concentration and C and N quotas in

iron-replete and deplete *Micromonas sp.* cells. Understanding these iron limitation-driven

differences is critical as changes in growth rates and elemental quotas of *Micromonas* may

have ecosystem and biogeochemical implications in HNLC regions. Consistent with current

paradigms of diatoms, iron-replete *Micromonas* sp. were larger, had higher growth rates,

higher cellular Chl *a* content, and appreciably higher C and N quotas relative to their iron-

limited counterparts. Although the Fe-limited cells were smaller with lower growth rates and

cellular quotas, they were able to reach higher cell densities under similar culturing

conditions. These findings suggest *Micromonas* can achieve numerically dominance in iron-

limited environments through reducing their cellular elemental quotas.

4.2 Iron limitation strategies

Our findings are in line with previous work (Martin and Fitzwater, 1988; Boyd et al.,

1998; Marchetti et al., 2006) suggesting low Fe concentrations limit nitrate uptake as is

supported by a low relative growth rate (μ/μ_{max}). This leads to substantially curtailed growth

rates, and trends in gene expression where genes involved in nitrate metabolism (e.g., *NRT*,

NiR) are more highly expressed under iron-replete conditions. The enhanced expression of

genes related to nitrogen metabolism under iron replete conditions as well as our Fe+

modeled rates consistent with those small celled new production rates measured during

EXPORTS suggests that picoeukaryotes such as *Micromonas* may be an important

contributor to new production in the iron-limited waters of the Northeast Pacific. Our data

also demonstrates clear expression patterns in genes involved in NO_3^- metabolism coinciding

with their ability to grow effectively using NO_3^-.

Micromonas also appears capable of urea metabolism by expressing urea transporters

in addition to *NiR* and *NRT*. A similar finding was reported by Piganeau et al. (2011) who

exhibited selective pressure for nitrate and urea uptake, but not ammonium uptake. This

finding is consistent with our inability to establish steady-state growth when cells were grown

in NH_4^+ based media. However, this is surprising given the need to reduce NO_3^- to NH_4^+ for

utilization in protein synthesis and the lower energetic cost of NH_4^+ utilization relative to

NO_3^- utilization (Raven, 1988). Cultures were able to grow sporadically on NH_4^+, but steady-

state growth was not achieved. This, combined with the presence of *AMT* in the

transcriptome, supports previous studies (Cochlan and Harrison, 1991a; 1991b) and suggests

Micromonas have the physiological capabilities for NH_4^+ utilization but were perhaps

inhibited by some other factor, such as a vitamin or a micronutrient.

The lack of information on iron regulation is consistent with findings from previous

studies of the closely related *Ostreococcus* (Sutak et al., 2012; Lelandais et al., 2016). The

majority of what is known about phytoplankton iron metabolism comes from studies of

diatoms (Peers and Price, 2006; Marchetti et al., 2012). It is therefore not surprising that

many of the iron transport and homeostasis genes identified in diatoms were not readily

identified in *Micromonas* given the substantial physiological differences between the two

phytoplankton groups. The only iron-related gene identified and present in the *Micromonas*

transcriptome is ISIP3, an iron-starvation induced protein with ferritin-like domains (Behnke

and LaRoche, 2020). However, as is noted by Lelandais et al. (2016) for *Ostreococcus*, the

high percentage of unannotated genes in our strain suggests that chlorophytes likely have

novel methods for dealing with iron limitation which have not yet been well characterized.

Sutak et al. (2012) also found both *Ostreococcus* and *Micromonas* were able to take up ferric

and ferrous iron without prior iron reduction but unable to store it. The authors hypothesize

ferritin may be present in *Ostreococcus* but not in *Micromonas* (Sutak et al., 2012). Our study

represents the first analysis of *Micromonas* isolated from an HNLC region. The cosmopolitan

nature and wide thermal niche of *Micromonas* suggests it can adapt to new or changing

environments and proliferate quickly (Vannier et al., 2016; Demory et al., 2018). This may

suggest the isolate studied here exhibits physiological characteristics and adaptations that

may be quite different from what is currently characterized in the scientific literature using

isolates from other regions.

An additional, noteworthy finding is that our data appears to suggest that *Micromonas*

exhibit a form of nitrogen metabolism conservation even under Fe replete conditions. The

gene expression analysis demonstrates most protein-encoding genes related to nitrogen

metabolism (*NRT*, *NR*, *AMT*, *nirA*) have some of the highest constitutive transcript

abundances of any pathway, exhibiting some significant directionality of gene expression

under variable Fe conditions, but only to a minimal extent (Fig. 4B). Some studies have

suggested diatoms exhibit substantial changes in expression of genes involved in nitrogen

metabolism, particularly in the natural environment (Marchetti et al., 2012), whereas others

have likewise shown a lack of substantial change in gene expression (but not protein

synthesis) in diatoms grown in culture (Cohen et al., 2018). This finding supports previous

studies which suggest picoeukaryotes, and other small phytoplankton, provide a "baseline"

rate of primary production under limiting conditions with a minimal physiological response

to dramatic shifts in environmental conditions (Meyer et al., 2022). It is possible that

Micromonas expresses additional genomic adaptation strategies for iron limitation that were

not identified from our transcriptome analysis. Additional research is needed to fully

elucidate the strategies *Micromonas* uses to subsist in iron-limited environments.

4.3 Oceanographic context

439   Our study provides valuable cellular characteristics of an abundant, picoeukaryote in

a climatologically important oceanographic context. Meyer et al. (2022) found that small

cells (<5 μm) accounted for the majority of biomass (68% of total biomass) and primary

production (89% of total primary production), including new production (72% of total new

production), at Station P during the EXPORTS observation period. Furthermore, *Micromonas*

appears to be highly represented within the small-celled phytoplankton community, with 18S

rDNA data indicating that *Micromonas* was within the top 20% of the most abundant

phytoplankton genera and one of the top five most abundant picoeukaryotes (Jones and

Rynearson, 2022).

448   Our estimated rates of new production by picoeukaryotes such as *Micromonas*, taken

as a representative species, under Fe- and Fe+ conditions were compared to the average

small-celled new production rates of 33.3 ± 14.4 nmol N L^{-1} d^{-1} from the NP EXPORTS

campaign (Fig. 3). While higher rates of NO_3^- uptake may occur within a culture relative to a

mixed, natural phytoplankton community, our analysis supports the idea that *Micromonas*

contributes substantially to total new production at Station P, especially when iron

concentrations can support higher growth rates and cellular nitrogen quotas. Inherently, other

phytoplankton groups, both within the picoplankton and nanoplankton size ranges, present at

St. P during the EXPORTS observation period also contribute appreciably to new production.

Our culture-based analysis is not meant to suggest *Micromonas* accounted for all of this new

production, but rather, that this chlorophyte should at least be considered as a significant

contributor to new production (and by inference carbon export) as estimated from in situ

picoeukaryote cell densities along with our measured cellular nitrogen quotas and growth

rates as a function of iron status.

With projected decreases in euphotic zone nutrient concentrations in many parts of

the oceans as a result of climate change (Kwiatkowski et al., 2020), induced changes in water

mass stratification and overall circulation, models of future projections predict an overall

shift towards dominance by smaller phytoplankton groups such as chlorophytes (Henson et

al., 2021). Abundant, small-cell phytoplankton with streamlined genomes and enhanced

capabilities to adapt to changing environmental conditions such as *Micromonas* hold distinct

advantages in future climate conditions. Further studies on *Micromonas* grazing rates, sinking

speeds, etc. are needed to fully understand its ecology moving forward and how this will

impact net carbon export rates and trophic level interactions.

Conclusion

In low iron environments, *Micromonas sp.* exhibits physiological strategies, including

reduced growth rates, small cell size, and metabolic restructuring, to combat Fe limitation

and maintain high rates of new production and substantial abundances. This study furthers

our understanding of this poorly characterized picoeukaryote and verifies a potential role for

chlorophytes in performing new production in iron-limited regions. Here, we add to the body

of knowledge reaffirming the need to shift our thinking on key phytoplankton groups that

have the potential to contribute to carbon export and pay additional attention to the role

picoeukaryotes play in ecosystem dynamics both in current and future ocean climate

scenarios.

Acknowledgements

We thank the EXPORTS Chief Scientists, D. Steinberg and J. Graff, and the Captain and crew of the *R/V* Roger Revelle for their roles in the initial sample collection, R. Paerl for use of his flow cytometer, and J. Graff for NP EXPORTS flow cytometry data. We also thank R. Flynn for her helpful comments. This work was supported by NASA Grant 80NSSC17K0552 and NSF Grant OCE2219973 to AM.

Competing Interests

The authors declare no competing interests.

Data Availability Statement

All data presented here are publicly available at the National Center for Biotechnology Information (Accession number PRJNA1072555) or the Biological and Chemical Oceanography Data Management Office (project ID: 914269).

499
500 **References**
501

Andrew, S.M., Moreno, C.M., Plumb, K., Hassanzadeh, B., Gomez-Consarnau, L., Smith,
S.N., Schofield, O., Fujiwara, T., Sunda, W.G., Hopkinson, B.M., Septer, A.N., &
Marchetti, A. (2023). Widespread use of proton-pumping rhodopsin in Antarctic
phytoplankton. *Proceedings of the National Academy of Science* 120 (39).
doi.org/10.1073/pnas.2307638120.

Behnke, J., & La Roche, J. (2020). Iron uptake proteins in algae and the role of Iron
Starvation-Induced Proteins (ISIPs). European Journal of Phycology 55(3), 339-360.
doi.org/10.1080/09670262.2020.1744039.

Boyd, P.W., Berges, J.A., & Harrison, P.J. (1998). In vitro enrichment experiments at iron-
rich and poor sites in the NE subarctic Pacific. *Journal of Experimental Marine
Biology and Ecology* 227: 133–151. doi.org/10.1029/98GB00745.

Brand, L.E., Guillard, R.R.L., & Murphy, L.S. (1981). A method for rapid and precise
determination of acclimated phytoplankton reproduction rates. *Journal of Plankton
Research* 3(2): 193-201. doi.org/10.1093/plankt/3.2.193.

Chisholm, S.W., Frankel, S.L., Goericke, R., Olson, R.J., Palenik, B., Waterbury, J.B., West-
Johnsrud, L., & Zettler, E.R. (1991). *Prochlorococcus marinus* nov. gen. nov. sp.: an
oxyphototrophic marine prokaryote containing divinyl chlorophyll *a* and *b*. *Arch.
Microbiol.* 157:297-300.

Cochlan, W.P., & Harrison, P.J. (1991a). Uptake of nitrate, ammonium, and urea by nitrogen-
starved cultures of Micromonas pusilla (Prasinophyceae): transient responses. *Journal
of Phycology* 27: 673 – 679.

Cochlan, W.P., & Harrison, P.J. (1991b). Kinetics of nitrogen (nitrate, ammonium and urea)
uptake by the picoflagellate Micromonas pusilla (Prasinophyceae). *Journal of
Experimental Marine Biology and Ecology* 153: 129 – 141.

Cohen, N.R., Gong, W., Moran, D.M., McIlvin, M.R., Saito, M.A., & Marchetti, A. (2018).
Transcriptomic and proteomic responses of the oceanic diatom Pseudo-nitzschia
granii to iron limitation. *Environmental Microbiology* 20(8), 3109-3126.
doi:10.1111/1462-2920.15386.

Demory, D., Baudoux, A.-C., Monier, A., Simon, N., Six, C., Ge, P., Rigaut-Jalabert, F.,
Marie, D., Sciandra, A., Bernard, O., & Rabouille, S. (2019). Picoeukaryotes of the
Micromonas genus: sentinels of a warming ocean. *The ISME Journal* 13:132-146.
doi.org/10.1038/s41396-018-0248-0.

Ellis, K. (2015). The coeffects of iron and vitamin B12 on marine diatoms. ProQuest
Dissertations & Theses Global. (1758252386).

Fowler, B.L., Neubert, M.G., Hunter-Cevera, K.R., Olson, R.J., Shalapyonok, A., Solow, A.R., & Sosik, H. (2020). Dynamics and functional diversity of the smallest phytoplankton on the Northeast US Shelf. *Proceedings of the National Academy of Science* 117(22), 12215-12221. https://doi.org/10.1073/pnas.1918439117.

Fu, L., Beifang, N., Zhu, Z., Wu, S., Weizhong, L. (2012). CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 28(23; 3150-3152. doi:10.1093/bioinformatics/bts565.

Gorbunov, M.Y., & Falkowski, P.G. (2004).Fluorescence induction and relaxation (FIRe) technique and instrumentation for monitoring photosynthetic processes and primary production in aquatic ecosystems. *Photosynthesis: Fundamental aspects to global perspectives*. 1029-1031.

Graff, J.R. (2023). Data from the EXport Processes in the Ocean from RemoTe Sensing North Pacific Campaign. NASA SeaWiFS Bio-optical Archive and Storage System (SeaBASS). (Version 1) https://seabass.gsfc.nasa.gov/cruise/EXPORTSNP/EXPORTS_EXPORTSNP_RR181 3_Graff_Cphyto_Analytical_R0/ [accessed 25/02/2025].

Graff, J.R., & Rynearson, T.A. (2011). Extraction method influences the recovery of phytoplankton pigments from natural assemblages. *Limnology and Oceanography* 9: 129–139. doi.org/10.4319/lom.2011.9.129.

Greene, R.M., Geider, R.J., Kolber, Z., & Falkowski, P.G. (1992). Iron-induced changes in light harvesting and photochemical energy conversion processes in eukaryotic marine algae. *Plant Physiology* 100: 565–575. doi:0032-0889/92/100/0565/11/$01.00/0

Henson, S.A., Cael, B.B., Allen, S.R., & Dutkiewicz, S. (2021). Future phytoplankton diversity in a changing climate. *Nature Communications* 12:5372. https://doi.org/10.1038/s41467-021-25699-w.

Jones, E., & Rynearson, T. (2022). Data from the EXport Processes in the Ocean from RemoTe Sensing North Pacific Campaign. National Center for Biotechnology Information (NCBI). (Version 1) https://www.ncbi.nlm.nih.gov/bioproject/796004 [accessed 08/10/2024].

Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J.R., Dunne, J.P., Gehlen, M., Ilyina, T., John, J.G., Lenton, A., Li, H., Lovenduski, N.S., Orr, J.C., Palmieri, J., Santana-Falcon, Y., Schwinger, J., Seferian, R., Stock, C.A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A., & Ziehn, T. (2020). Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections. *Biogeosciences* 17(13), 3439-3470. https://doi.org/10.5194/bg-17-3439-2020.

La Roche, J., Geider, R.J., Graziano, L.M., Murray, H., & Lewis, K. (1993). Induction of specific proteins in eukaryotic algae grown under iron-, phosphorus-, or nitrogen-

594 deficient conditions. *Journal of Phycology* 29, 767-777.
595 doi:http://dx.doi.org/10.1111/j.0022-3646.1993.00767.x.
596

597 Le Gall, F., Rigaut-Jalabert, F., Marie, D., Garczarek, L., Viprey, M., Gobet, A., & Vaulot,
598 D. (2008). Picoplankton diversity in the South-East Pacific Ocean from cultures.
599 *Biogeosciences* 5, 203-214. https://doi.org/10.5194/bg-5-203-2008.
600

601 Lelandais, G., Scheiber, I., Paz-Yepes, J., Lozano, J., Botebol, H., Pilatova, J., Zarsky, V.,
602 Leger, T., Blaiseau, P., Bowler, C., Bouget, F., Camadro, J., Sutak, R., & Lesuisse, E.
603 (2016). Ostreococcus tauri is a new model green alga for studying iron metabolism in
604 eukaryotic phytoplankton. *BMC Genomics* 17:319. doi:10.1186/s12864-016-2666-6.
605

606 Marchetti, A., Juneau, P., Whitney, F.A., Wong, C., & Harrison, P.J. (2006). Phytoplankton
607 processes during a mesoscale iron enrichment in the NE subarctic Pacific: Part II-
608 Nutrient utilization. *Deep Sea Research II* 53, 2114-2130.
609 doi:http://dx.doi.org/10.1016/j.dsr2.2006.05.038.
610

611 Marchetti, A., Parker, M.S., Moccia, L.P., Lin, E.O., Arrieta, A.L., Ribalet, F., Murphy,
612 M.E.P., Maldonado, M.T., & Armbrust, E.V. (2009). Ferritin is used for iron storage
613 in bloom-forming marine pennate diatoms. *Nature* 457, 467-470.
614 doi.org/10.1038/nature07539.
615

616 Marchetti, A., Schruth, D.M., Durkin, C.A., Parker, M.S., Kodner, R.B., Berthiaume, C.T.,
617 Morales, R., Allen, A.E., & Armbrust, E.V. (2012). Comparative metatranscriptomics
618 identifies molecular bases for the physiological responses of phytoplankton to varying
619 iron availability. *Proceedings of the National Academy of Sciences* 109(6), E317-
620 E325. doi.org/10.1073/pnas.1118408109.
621

622 Martin, J.H., & Fitzwater, S. (1988). Iron deficiency limits phytoplankton growth in the
623 north-east Pacific subarctic. *Nature* 331, 341-343.
624 doi:http://dx.doi.org/10.1038/331341a0.
625

McNair, H., Meyer, M.G., Lerch, S., Maas, A.E., Stephens, B., Fox, J., Buck, K.N., Burns,
S.M., Cetinic, I., Cohn, M., Durkin, C., Gifford, S., Gong, W., Graff, J.R., Jones,
E.L., Santoro, A.E., Shea, C.H., Stamieszkin, K., Steinberg, D.K., Marchetti, A.,
Carlson, C.A., Menden-Deuer, S., Brzezinski, M.A., Siegel, D.A., Rynearson, T.A.
(2019). Quantitative analysis food web dynamics in a low export ecosystem. *BioRvix*.
doi.org/10.1101/2023.03.17.532807.
632

633 Meyer, M.G., Gong, W., Kafrissen, S.M., Torano, O., Varela, D.E., Santoro, A.E., Cassar,
634 N., Gifford, S., Niebergall, A.K., Sharpe, G., & Marchetti, A. (2022). Phytoplankton
635 size-class contributions to new and regenerated production during the EXPORTS
636 Northeast Pacific Ocean field deployment. *Elementa: Science of the Anthropocene* 10.
637 https://doi.org/10.1525/elementa.2021.00068.
638

639 Moore, L.R., Post, A.F., Rocap, G., & Chisholm, S.W. (2002). Utilization of different
640 nitrogen sources by the marine cyanobacteria, *Prochlorococcus* and *Synechococcus*.
641 *Limnology and Oceanography* 47(4):989-996.
642

643   Monier, A., Worden, A.Z., & Richards, T.A. (2016). Phylogenetic diversity and
644   biogeography of the Mamiellophyceae lineage of eukaryotic phytoplankton across the
645   oceans. *Environmental Microbiology Reports* 8(4), 461-469. doi:10.1111/1758-
646   2229.12390.
647

648   Moreno, C.M., Gong, W., Cohen, N.R., DeLong, K., & Marchetti, A. (2020). Interactive
649   effects of iron and light limitation on the molecular physiology of the Southern Ocean
650   diatom Fragilariopsis kerguelensis. Limnology and Oceanography 65(7), 1511-1531.
651   doi.org/10.101002/lno.11404.
652

653   Moreno, C.M., Lin, Y., Davies, S., Monbureau, E., Cassar, N., & Marchetti, A. (2017).
654   Examination of gene repertoires and physiological responses to iron and light
655   limitation in Southern Ocean diatoms. *Polar Biology* 41, 679-696.
656   doi.org/10.1007/s00300-017-2228-7.
657

658   Paerl, R.W., Venezia, R.E., Sanchez, J.J., & Paerl, H.W. (2020). Picophytoplankton
659   dynamics in a large temperate estuary and impacts of extreme storm events. *Scientific*
660   *Reports* 10. https://doi.org/10.1038/s41598-020-79157-6.
661

662   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., & Kingsford, C. (2017). Salmon provides
663   fast and bias-aware quantification of transcript expression. *Nature Methods*.
664

665   Peers, G., & Price, N.M. (2006). Copper-containing plastocyanin used for electron transport
666   by an oceanic diatom. *Nature* 441:341-344.
667   doi:http://dx.doi.org/10.1038/nature04630.
668

669   Piganeau, G., Grimsley, N., & Moreau, H. (2011). Genome diversity in the smallest marine
670   photosynthetic eukaryotes. *Research in Microbiology* 162(6), 570-577.
671   doi.org/10.1016/j.resmic.2011.04.005.
672

673   Price, N.M., Ahner, B.A., & Morel, F.M.M. (1994). The equatorial Pacific Ocean: Grazer-
674   controlled phytoplankton populations in an iron-limited ecosystem. *Limnology and*
675   *Oceanography* 39, 520-534. http:doi.org/10.4319/lo.1994.39.3.0520.
676

677   Price, N.M., Harrison, G.I., Hering, J.G., Hudson, R.J., Nirel, P.M.V., Palenik, B., & Morel,
678   F.M.M. (1989). Preparation and chemistry of the artificial algal culture medium
679   Aquil. *Biological Oceanography* 6, 443-461.
680   doi.org/10.1080/10965581.1988.10749544.
681

682   Raven, J.A. (1988). The iron and molybdenum use efficiencies of plant growth with different
683   energy, carbon, and nitrogen sources. *New Phytologist* 279–287.
684   doi.org/10.1111/j.1469-8137.1988.tb04196.x
685

686   Sharpe, G., Zhao, L., Meyer, M.G., Gong, W., Burns, S.M., Tagliabue, A., Buck, K.N.,
687   Santoro, A.E., Graff, J.R., Marchetti, A., & Gifford, S. (2023). *Synechococcus*
688   nitrogen gene loss in iron-limited ocean regions. *ISME Communications* 3, 107.
689   https://doi.org/10.1038/s43705-023-00314-9.
690

691   Soneson, C., Love, M.I., & Robinson, M.D. (2015). Differential analyses for RNA-seq:
692   transcript-level estimates improve gene-level inferences. *F1000Research*. 4.
693   Doi:10.12688/f1000research.7563.1.
694
695   Sutak, R., Botebol, H., Blaiseau, P.-L., Leger, T., Bouget, F.-Y., Camadro, J.-M., & Lesuisse,
696   E. (2012). A comparative study of iron uptake mechanisms in marine microalgae: iron
697   binding at the cell surface is a critical step. *Plant Physiology* 160, 2271-2284.
698   doi:10.1104/pp.112.204156.
699
700   Varela, D.E., & Harrison, P.J. (1999). Seasonal variability in nitrogenous nutrition of
701   phytoplankton assemblages in the northeastern subarctic Pacific Ocean. *Deep Sea*
702   *Research II* 46, 2505-2538. http://dx.doi.org/10.1016/S0967-0645(99)00074-0.
703
704   Vannier, T., Leconte, J., Seeleuthner, Y., Mondy, S., Pelletier, E., Aury, J., de Vargas, C.,
705   Sierack, M., Iudicone, D., Vaulot, D., Wincker, P., & Jaillon, O. (2016). Survey of the
706   green picoalga Bathycoccus genomes in the global ocean. *Scientific Reports* 6:37900.
707   doi:10.1038/srep37900.
708
709   Worden, A.Z., Nolan, J.K., & Palenik, B. (2004). Assessing the dynamics and ecology of
710   marine picophytoplankton: The importance of the eukaryotic component. *Limnology*
711   *and Oceanography* 49(1), 168-179. doi.org/10.4319/lo.2004.49.1.0168.
712
713   Worden, A.Z., & Not, F. (2008). Ecology and diversity of picoeukaryotes. *Microbial ecology*
714   *of the Oceans* 2, 159-205.
715

Fig. 1. Specific growth rates (A; d^{-1}), maximum photochemical efficiencies of PSII (B;
F$_v$/F$_m$), cell volumes (C; μm^3), and cellular chlorophyll a quotas (D; fg μm^{-3}) for *Micromonas*
grown under iron-replete (Fe+) and iron-limited (Fe-) conditions. Error bars represent
standard deviations with n = 3 for Fe- and n = 2 for Fe+.

Fig. 2. Cellular carbon (A,B) and nitrogen (C,D) quotas and C:N (E) and C:Chl *a* (F) ratios
for *Micromonas* grown under iron-replete (Fe+) and iron-limited (Fe-) conditions. Quotas are
represented per cell (A,C; fmol cell^{-1}) and per cell volume (B,D; fmol µm^{-3}). Error bars
represent standard deviations with n = 3 Fe- and n = 2 for Fe+.

Fig. 3. Estimates of *Micromonas sp.* contributions to new production during the North Pacific EXPORTS cruise. *Micromonas sp.* new production rate estimates for NO_3^- Fe- and NO_3^- Fe+, are calculated from culture-based growth rates, cellular N quotas, and in situ NP EXPORTS picoeukaryotic cell counts obtained from flow cytometry. The gray shaded region represents the upper bounds (NO_3^- Fe+) and lower bounds (NO_3^- Fe-) for the culture-based rate estimates. The blue line represents the average mixed-layer rates of new production of the small the size-fraction (< 5 μm) during EXPORTS by yearday during the observation period.

724
725

Fig. 4. *Micromonas sp.* gene expression as a function of iron status. Heatmap (A) of average Log2Fold change of KEGG pathways, RA plot (B) of average log counts per million (CPM) vs. average Log2Fold change, and nitrogen assimilation pathway map (C) of *Micromonas sp.* Fe+ vs. Fe- culture RNA transcripts. Transcripts were annotated and evaluated on the KO level using a significance threshold of $p < 0.05$. For all plots, positive Log2Fold change indicates overrepresentation in Fe+ cultures and negative Log2Fold change indicates overrepresentation in Fe- samples. Astericks on the heatmap and pathway map indicate significant changes ($p < 0.05$). The RA highlights protein-encoding genes of interest. The cellular pathway map indicates the sign and relative strength of Log2Fold change expression of genes. Grey genes are absent from transcript data. Abbreviations are: nitrate transporter (*NRT*), nitrite reductase (*NiR*), nitrate reductase (*NR*), ammonium transporter (*AMT*), ornithine carbamolytransferase (*OTC*), urea transport (*Urt*), arginase (*Arg*), citrate synthase (*Glta*), photosystem I (*Psa*), photosystem II (*Psb*), cytochrome c oxidase (*COX*), cytochrome b (*PetB*), and cytochrome c (*PetC*).

731