Weak, specific chemical interactions dictate barnase stability in diverse cellular environments

Ume Tahir[1] and Caitlin M. Davis[1*]

[1]Department of Chemistry, Yale University, New Haven, CT 06511, USA

*Corresponding Author:

Caitlin M. Davis, Department of Chemistry, Yale University, New Haven, CT 06511, USA

/ c.davis@yale.edu

Running Title: Weak, specific chemical interactions dictate barnase stability in diverse cellular environments

Total Number of Manuscript Pages: 30

Total Number of Supplementary Material Pages: 9

Number of Tables: 0

Number of Figures: 5

Description of Supplementary Material: Supplementary materials include a representative circular dichroism and fluorescence melt of unlabeled barnase, representative temperature-dependent behavior of aggregated in-cell populations, a table of protein sequences, and tables of melting temperatures collected in vitro and in living cells.

Abstract

It is well-established that in vitro measurements do not reflect protein behaviors in-cell, where macromolecular crowding and chemical interactions modulate protein stability and kinetics. Recent work suggests that peptides and small proteins experience the cellular environment differently from larger proteins, as their small sizes leaves them primarily susceptible to chemical interactions. Here, we investigate this principle in diverse cellular environments, different intracellular compartments and host organisms. Our model protein is barnase, a bacterial ribonuclease that has been extensively characterized in vitro. We find that barnase is stabilized in the cytoplasm and destabilized in the nucleus of U2-OS cells. These trends could not be reproduced in vitro by Ficoll and M-PER^{TM}, which mimic macromolecular crowding and non-specific chemical interactions, respectively. Instead, in-cell trends were best replicated by cytoplasmic and nuclear lysates, indicating that weak specific interactions with proteins in either compartment are responsible for the in-cell observations. Interestingly, in the cytoplasm barnase's unfolded state is unstable and prone to aggregation, while in the nucleus a stable unfolded state exists prior to aggregation. We hypothesize that the reduced thermal stability and stable unfolded state in the nucleus correlates with increased conformational flexibility in a functionally relevant context for a ribonuclease. Indeed, in the biologically relevant environment of bacterial cells barnase folding resembled that in the nucleus, but with no aggregation at higher temperatures. These findings show that protein interactions are evolved for their native environment, which highlights the importance of studying and designing proteins in situ.

Keywords

barnase, chemical interactions, fast relaxation imaging (FReI), Förster resonance energy transfer (FRET), laser-induced temperature jump, macromolecular crowding, protein folding

1 Introduction

It is well-established that macromolecular crowding and chemical interactions between biomolecules inside cells can substantially tune both protein stability and folding kinetics (Danielsson et al., 2015; Davis & Gruebele, 2018; Ebbinghaus et al., 2010; Knab & Davis, 2023; Monteith et al., 2015). Furthermore, given the heterogeneity of the cell interior, these properties may vary depending on the local microenvironment. For example, it has been demonstrated that two model proteins, phosphoglycerate kinase (PGK) and variable major protein-like sequence expressed (VlsE), exhibit different stabilities and folding kinetics in different cellular compartments (Dhar et al., 2011; Tai et al., 2016). Thus, in-cell measurements contain information in a functionally-relevant context that cannot be simply extrapolated from in vitro data.

Currently, much of our understanding of how the cellular environment modulates protein behavior, especially in the context of eukaryotic cells, is derived from large proteins, like PGK and VlsE. There is comparatively less known about how peptides and small proteins behave in eukaryotes, despite their diverse biological functions – including as enzymes, chaperones, and regulators of cell division (Steinberg & Koch, 2021; Storz et al., 2014). Given their small sizes relative to the surrounding macromolecules, peptides and small proteins are expected to experience the cellular environment differently from larger proteins. While the average $\approx$400 amino acid eukaryotic protein is susceptible to both macromolecular crowding and chemical interactions, peptides and small proteins are unlikely to be sterically constrained by larger macromolecules (Brocchieri & Karlin, 2005; Christiansen et al., 2010; Chu et al., 2023). Indeed, recent work on the 80 amino acid λ-repressor protein fragment ($\lambda_{6\text{-}85}$), suggests that the stability of small proteins in

eukaryotic cells is dictated solely by chemical interactions (Knab & Davis, 2023). This aligns with earlier results for SH3, a small protein folding model that is destabilized by chemical interactions in both prokaryotic and eukaryotic cells, despite the vastly different concentrations of biomolecules in these environments (Smith et al., 2016; Thole et al., 2021). However, there is still a need for additional studies to determine if this principle applies to other small proteins and to further explore the effects of different types of cellular microenvironments on small protein and peptide stability.

A

AQVINTFDGV ADG**L**QTYHKL PDNYITKSEA QALGWVASKG
NLADVAPGKS IGGDIFSNRE GKLPGKSGRT WREADINYTS
GFRNSDRILY SSDWLIYKTT D**A**YQTFTKIR

B

Figure 1. (A) Structure and sequence of barnase with Y13G and H102A mutations highlighted in yellow (PDB: 1BNR). Figure created using PyMOL (The PyMOL Molecular Graphics System, n.d.). (B) Representative fluorescence melts for 10 μM unlabeled (blue) and 0.25 μM FRET-labeled (magenta) barnase in 20 mM sodium phosphate, pH 7.4. The melts were normalized and fit to a two-state denaturation (E1).

This work focuses on barnase (Figure 1A), a 110-residue extracellular ribonuclease from *Bacillus amyloliquefaciens* (Hartley, 1989). Barnase has an extensive history of being used as a protein folding model, with a thoroughly characterized in vitro

folding mechanism (Alemany et al., 2016; Fersht, 1993, 2000; Kippen et al., 1994). The abundance of in vitro data makes it an interesting candidate for in-cell work. For example, a collection of barnase mutants with differing stabilities has been used as biosensors to report on in-cell chaperone activity (Wood et al., 2018). With Förster resonance energy transfer (FRET) reporting on the folding state of the protein, it was found that the barnase mutants have different distributions between the folded, unfolded, and aggregated states in the cytoplasm and nucleus (Raeburn et al., 2022). These differences in behavior across different cellular compartments raise the question of how the local microenvironment modulates the folding of small proteins and peptides.

Here, we explore the effect of different cellular compartments on barnase stability. We find that compared to in vitro, barnase is slightly stabilized in the cytoplasm and significantly destabilized in the nucleus of mammalian U2-OS cells. Interestingly, a stable unfolded state exists in the nucleus, whereas barnase is more aggregation-prone in the cytoplasm. We hypothesize that this may be attributed to barnase, a ribonuclease, having greater conformational flexibility in a more functionally relevant environment, rich in both proteins and nucleic acids. To understand the differences between in vitro and in-cell stabilities, we used Ficoll and mammalian protein extraction reagent (M-PER^{TM}) to mimic crowding and non-specific chemical interactions, respectively, and found that neither could replicate cellular trends. Instead, these trends were best reproduced by cytoplasmic and nuclear lysates, suggesting that weak specific interactions with proteins in either compartment are most important for determining cellular stability. Lastly, we investigated how the conclusions drawn from eukaryotic U2-OS cells translate to the more biologically relevant context of bacterial cells. We found that unlike in mammalian cells, barnase does

not aggregate upon unfolding in bacterial cells demonstrating that evolution has carefully optimized protein interactions in their native environment. This has important implications for functional protein design, which is typically performed in vitro, and interpretation of phylogenic reconstructions of ancestral proteins, which is often performed protein by protein.

2 Results and Discussion

2.1 The addition of FRET labels to barnase is minimally perturbative

For this work, we used the Y13G H102A mutant of barnase (Figure 1A). The H102A mutation is commonly used to diminish the toxic ribonuclease activity of barnase in the absence of its intracellular inhibitor (Meiering et al., 1992), and the Y13G mutation further destabilizes the construct to yield a melting temperature (T_M) that is compatible with our in-cell temperature-dependent experiments (36.7 $^{\circ}$C) (Vu et al., 2004). We expressed and purified this construct and found its structure and T_M to be consistent with the literature using circular dichroism (Figure S1) (Vu et al., 2004; Vuilleumier et al., 1993).

To track the folding state of the protein in-cell, we designed a FRET-labeled construct with AcGFP1 as the donor fluorophore on the N-terminus and mCherry as the acceptor fluorophore on the C-terminus. The donor to acceptor fluorescence ratio (D/A) reports on the end-to-end distance of the protein, with higher values representing a more extended state. However, prior to using this construct in-cell, it was important to verify that the labels did not perturb the folding behavior of barnase. Thus, we derived T_M values for the unlabeled and labeled constructs using tryptophan fluorescence and FRET melts, respectively. We found that the T_M of the unlabeled construct was 38.16 ± 0.07 $^{\circ}$C, and

the T_M of the FRET-labeled construct was 39 ± 2 °C (Figure 1B). We therefore concluded that the introduction of FRET labels was minimally thermodynamically perturbative.

2.2. Barnase is slightly stabilized in the cytoplasm and significantly destabilized in the nucleus of U2-OS cells

To examine the effect of different local cellular environments on barnase stability, we transfected U2-OS cells with FRET-labeled barnase containing either a nuclear export or nuclear localization sequence in U2-OS cells (Table S1). The stability in the cytoplasm and nucleus of U2-OS cells was probed by fast relaxation imaging (FReI), which couples the spatial resolution of fluorescence microscopy with perturbations in the form of laser-induced temperature jumps; thermodynamic information is derived from the 10 second equilibrium phase following the temperature jump (Ebbinghaus et al., 2010).

We observed distinct behaviors in the cytoplasm and nucleus. Interestingly, the D/A in the cytoplasm consistently trended downwards with increasing temperature, which is indicative of aggregation without an observable, stable unfolded state (Figure 2). Thus, we report an aggregation temperature (T_A) in place of T_M for the cytoplasm. The T_A was 41.4 ± 0.5 °C across 70 cells (Table S2). This reflects that the folded state is slightly stabilized by ≈2 °C relative to in vitro.

In the nucleus, there is a clear unfolding transition prior to the onset of aggregation at a T_A similar to the cytoplasm (Figure 2). The T_M was 26.8 ± 0.3 °C across 75 cells (Table S3). This reflects a significant destabilization of ≈12 °C relative to in vitro, and ≈15 °C relative to the cytoplasm. A subset of cells (44%) exhibited only aggregation at T_A; these cells tended to have a lower or greater than average D/A, suggesting that the initial

conformation was either misfolded or unfolded (Figure S2). This is not surprising as the melting temperature in the nucleus is low, barnase is likely unfolded but below T_A at normal cell culture temperatures (37 °C), and therefore our experiments are conducted on cells with populations of barnase that can be reversibly folded.

Figure 2. Representative donor/acceptor (D/A) melting curves extracted from FRel of barnase in the cytoplasm (top) and nucleus (bottom) of U2-OS cells. The data are fit to a two-state denaturation (E1) with dashed vertical lines indicating the melting temperature in the nucleus (T_M) and midpoint of aggregation in the cytoplasm (T_A), respectively. Insets display representative cell images. In the bottom image the cell membrane is highlighted by a white dashed line. Scale bar is set to 10 μm.

2.3. Cellular stability trends cannot be explained by macromolecular crowding or non-specific chemical interactions alone

The cell interior is primarily water, and the remaining 30-40% consists of ions, various small organics and metabolites, and macromolecules (Davis & Gruebele, 2018; Davis et al., 2018; Zimmerman & Trach, 1991). Therefore, protein stabilities in-cell may be modulated by factors like macromolecular crowding and chemical interactions. Macromolecular crowding entropically destabilizes the extended unfolded state of a protein, but may have minimal impact on smaller proteins, like barnase, that are not sterically confined by larger macromolecules (Knab & Davis, 2023). Additionally, the differing stabilities of barnase in the cytoplasm and nucleus of U2-OS cells are unlikely to be explained by macromolecular crowding theory, given the relatively conserved concentrations of macromolecules across these cellular compartments (Guigas et al., 2007). Instead, since the cytoplasm and nucleus vary in their chemical compositions it is more probable that stability differences arise from differences in chemical interactions.

Cellular mimetic reagents can be used in vitro to untangle the origin of in-cell stability trends. Because of its relatively inert surface, Ficoll is often used to isolate crowding effects (Benton et al., 2012; Davis et al., 2020; Knab & Davis, 2023). Here, we derived melting temperatures for unlabeled barnase in the presence of increasing concentrations of Ficoll (Figure 3). We find that although there is a small stabilization trend between 0-35% Ficoll, which could account for the slight $\approx$2 °C stabilization of barnase in the cytoplasm, it does not explain the $\approx$12 °C destabilization in the similarly crowded nucleus. Additionally, the previously validated cytomimetic concentration of Ficoll (15%) (Davis et al., 2020; Knab & Davis, 2023), near the physiological concentration

of macromolecules in eukaryotic cells, produces only a mild stabilization of ≈ 0.8 °C. Therefore, as predicted, it is unlikely that macromolecular crowding plays a large role in regulating barnase stability in-cell.

Other than crowding, proteins also experience non-specific chemical interactions in-cell with ions, small organic molecules, fatty acids, amino acids, etc. Ionic strength is important because the presence of ions may electrostatically screen a protein from interactions with surrounding molecules; here, we replicate intracellular ionic strength using 150 mM KCl, as K^+ is the most abundant ion in mammalian cells (Ames & Nesbett, 1966). We find that although K^+ slightly stabilizes barnase by 1.66 ± 0.06 °C, it does not fully recapitulate our in-cell observations. In particular, the concentration of K^+ is likely higher in the nucleus than in the cytoplasm (Kellermayer, 1981; Paine et al., 1981), so this stabilizing effect does not account for the significant destabilization observed in the nucleus relative to the cytoplasm.

We use M-PER™ to account for non-specific chemical interactions with small molecules in the cell. M-PER™ is a mild cell lysis reagent formulated to minimally interfere with downstream biological applications by preserving native protein properties. Specifically, M-PER™ contains a proprietary zwitterionic detergent in 25 mM bicine buffer (pH 7.6) that mimic interactions with lipids and amino acids, respectively (Knab & Davis, 2023; Yoo & Davis, 2022). Converse from Ficoll, we find that there is a destabilization trend with increasing concentrations of M-PER™ (Figure 3). However, at the previously validated cytomimetic concentration of 20% M-PER™ (Knab & Davis, 2023; Yoo & Davis, 2022), the destabilization is only 0.5 °C compared to a ≈ 12 °C destabilization in the nucleus. Thus, while chemical interactions with small molecules may account for some of

the nuclear destabilization of barnase, the observed cellular trends cannot be fully reproduced by macromolecular crowding or non-specific chemical interactions alone.

Figure 3. Change in the melting temperature of unlabeled barnase relative to 20 mM sodium phosphate, pH 7.4 with increasing concentrations of Ficoll and M-PER™. Melting temperatures derived from tryptophan melts with 10 μM barnase. Error bars display standard deviation of 3 measurements. Data is fit to a line.

2.4. Differences in stability between the cytoplasm and nucleus arise from weak specific interactions with proteins in either compartment

Although Ficoll and M-PER™ can reproduce non-specific crowding and chemical interaction effects, they do not capture behaviors that arise from weak, transient but specific interactions with macromolecules in the local microenvironment in-cell. The cytoplasm and nucleus differ in the types and quantities of macromolecules present, with the cytoplasm containing machinery for diverse metabolic functions, including protein synthesis (Kırlı et al., 2015), and the nucleus having more specialized proteins, including highly charged histones and transcriptional machinery, in addition to nucleic acids (van der Zanden et al., 2023). Thus, to account for weak specific chemical interactions in the

cytosol we mimic the cytoplasmic environment in vitro using cytoplasmic lysate, which contains cytoplasmic proteins extracted from U2-OS cells; likewise, we mimic the specific-protein and non-specific nucleic acid components of the nucleus using nuclear lysate, which contains nuclear proteins extracted from U2-OS cells, and salmon sperm DNA, respectively.

We find that FRET-labeled barnase experiences a substantial stabilization of 13 ± 6 °C in the presence of 10 mg/ml salmon sperm DNA, concentrations that are insufficient to induce macromolecular crowding (Figure 4, Table S4). Barnase is a ribonuclease that catalyzes the cleavage of single-stranded RNA. Like most ligands, RNA binding stabilizes barnase (Martinez et al., 1994). Positively charged residues at the binding interface of barnase bind and stabilize interactions with the negatively charged RNA substrate (Spaar et al., 2006). This positively charged patch is likely also poised to interact with other negatively charged biomolecules like DNA. Indeed, we observe that barnase is stabilized by non-specific interactions with DNA. This suggests that chemical interactions with nucleic acids are not responsible for the destabilization of barnase in the nucleus of U2-OS cells.

Rather, we find that specific protein interactions reproduce cellular trends; while barnase is slightly stabilized in cytoplasmic lysate, it is destabilized in nuclear lysate. This suggests that weak specific chemical interactions with proteins in either compartment are ultimately responsible for the observed in-cell stability trends. These differences may result from electrostatic interactions between barnase and proteins in each compartment; barnase has a pI of 9.2 and is therefore positively charged and poised to interact at physiological pH (Shankar et al., 2021). In eukaryotes, cytoplasmic proteins are typically

slightly acidic, resulting in an overall negative charge at physiological pH, whereas nuclear proteins span the entire range from acidic to basic, having overall negative, neutral or positive charges (Schwartz et al., 2001). Thus, attractive electrostatic interactions in the cytosol likely stabilize the native state of barnase, whereas the mixture of attractive and repulsive electrostatic interactions in the nucleus is destabilizing to barnase. It is important to note that lysates reproduce the observed cellular trends, but not their magnitudes; this is likely because the protein concentrations in these lysates are significantly lower than in-cell concentrations.

Figure 4. Change in the melting temperature of 0.25 µM FRET-labeled barnase relative to 20 mM sodium phosphate buffer, pH 7.4 in different conditions. The concentrations of in vitro buffer additive are 150 mg/ml Ficoll, 150 mM KCl, 20% M-PER^{TM}, 10 mg/ml salmon sperm DNA (ssDNA), $\approx$1 mg/ml cytoplasmic lysate (Cyt. Lte.), and $\approx$1 mg/ml nuclear lysate (Nuc. Lte.). Error bars display standard error of at least 3 measurements. *Reporting T_A for the cytoplasm instead of T_M

2.5. Barnase is destabilized and less aggregation prone in bacterial cells

Barnase is an extracellular bacterial protein that is synthesized and secreted by *B. amyloliquefaciens*. We hypothesize that though the thermal stability of barnase is reduced in the nucleus, this may represent enhanced conformational flexibility in a functionally relevant context for a secretory protein. Because it must be inhibited by barstar to protect the host from its ribonuclease activity (Shilova et al., 2021), it is evident that barnase is folded in the *B. amyloliquefaciens* cytosol. However, in Gram-positive bacteria like *B. amyloliquefaciens,* most secretory proteins are unfolded before they are translocated across the cytoplasmic membrane (Sarvas et al., 2004). This would also serve as a functional mechanism to release the tightly bound intracellular inhibitor, barstar, prior to excretion. Therefore, the host cytosol likely promotes more extended unfolded conformations. We have previously observed that the outer membrane protein VlsE is more extended and destabilized in the cytosol compared to in vitro (Davis et al., 2020; Guzman et al., 2014). Indeed, barnase is in a more extended conformation in the nucleus, with an average D/A of 5.0 ± 0.1 compared to 3.0 ± 0.1 in the cytoplasm.

Figure 5. Representative donor/acceptor (D/A) melting curve extracted from FReI of barnase Y13G H102A and barnase H102A overexpressed in BL21 DE3 cells. The data are fit to a two-state denaturation (E1) with dashed vertical lines indicating the melting temperature (T_M).

To further investigate the folding of barnase in a more biologically relevant environment we study its folding in BL21 DE3 *E. coli*, a bacterial prokaryote with no compartmentalization: the cytoplasm is a heterogeneous mixture of highly charged bacterial proteins and nucleic acids (Wennerström et al., 2020; Zimmerman & Trach, 1991). As anticipated, we find that barnase is destabilized in *E. coli* by 15.0 ± 0.7 °C (Figure 5, Table S5). Similar to the nucleus of U2-OS cells, the D/A trends upwards with increasing temperature, unfolding to a stable state that can be observed without aggregation. However, unlike in the nucleus of U2-OS cells, barnase does not typically aggregate within the temperature range of our experiments. These trends hold for pseudo-wild type barnase (barnase H102A), which is destabilized in *E. coli* by 9.5 ± 0.9 °C (Figure 5, Table S6). Taken together with the conclusions from U2-OS cells, this suggests that barnase becomes less thermally stable and more conformationally flexible

when approaching more native-like environments; however, this increase in conformational flexibility is less likely to be accompanied by unfavorable aggregation that can inhibit function. Proteins have evolved weak interactions to stabilize functional unfolded states in their native environment, like those necessary for the transport of extracellular proteins. However, such native interactions can be easily disrupted when proteins are moved to a nonnative environment (Mu et al., 2017).

3 Conclusion

We speculate that the decreased thermal stability of barnase in-cell relative to in vitro is representative of increased conformational flexibility in a more functionally relevant context for a ribonuclease, where it must be unfolded prior to excretion. We find that this flexibility is promoted by weak but specific chemical interactions with proteins in more native-like environments. That barnase does not aggregate upon unfolding in bacterial cells but does aggregate upon unfolding in eukaryotic cells demonstrates that proteins have carefully evolved protein interactions in their native environment. Such interactions may be crucial for functional protein design and interpretation of the evolution of proteomes, which are typically performed on an individual protein or protein-binding partner.

It is not yet possible to computationally predict how protein stability and kinetics change in-cell. Peptides and small proteins offer a path towards bridging this gap between theory and experiment, as experimental data can viably be coupled with all-atom molecular dynamics simulations (Rickard et al., 2020; Rickard et al., 2019). In the future, barnase offers the potential to explore in depth how in vitro protein folding principles

translate to the cellular environment. Barnase has an extensive history of being used as a protein folding model, and there exists a large library of mutants designed to probe the contributions of different types of intramolecular interactions to protein folding and stability (Serrano et al., 1992). However, it is not yet clear how these interactions may be modulated by the crowded cellular environment.

4 Materials and Methods

Protein Expression and Purification

The barnase Y13G H102A gene was inserted into the pET-15b vector for the unlabeled construct and the pDream 2.1 vector for the FRET-labeled construct, both of which contained His tags that could be cleaved by thrombin (GenScript Biotech, Piscataway, NJ). The proteins were expressed and purified from BL21(DE3) competent *E. coli* (New England Biolabs, Ipswich, MA). The cells were grown at 37°C until the OD_{600} reached 0.6, at which point expression was induced with 1 mM isopropyl-β-D-thiogalactopyranoside (Inalco, San Luis Obispo, CA). The induced cells grew overnight at 20°C. The cells were harvested by centrifugation, suspended in lysis buffer, and then lysed by sonication. The resulting lysate was centrifuged and filtered prior to being passed through a Ni-NTA column. Fractions containing protein were confirmed by running SDS-PAGE gels. The resulting fractions were then dialyzed into 20 mM sodium phosphate, pH 7.4. The unlabeled construct further underwent a thrombin cleavage to remove the His tag, using biotinylated thrombin (Millipore Sigma, Burlington, MA). This was also purified using a Ni-NTA column and dialyzed into 20 mM sodium phosphate, pH 7.4. The resulting concentration of the unlabeled construct was determined by absorbance at 280 nm (ε =

26,930 L^{-1} mol^{-1} cm^{-1}), and the concentration of the labeled construct was determined by mCherry absorption at 587 nm (ε = 72,000 L^{-1} mol^{-1} cm^{-1}). The unlabeled construct was characterized using circular dichroism, to be comparable to the literature (Vu et al., 2004; Vuilleumier et al., 1993), in addition to tryptophan fluorescence (Figure S3). The labeled construct was characterized using FRET.

Cell Culture and Transfection

Human bone osteosarcoma epithelial cells (U2-OS, ATCC, Manassas, VA) were transfected with the plasmid for either FRET-labeled barnase with a nuclear export sequence or nuclear localization sequence (Table S1). The cells used for transfection were grown in DMEM (Corning, Corning, NY) supplemented with 10% Fetal Bovine Serum (FBS, Gibco, Waltham, MA) and 1% Penicillin-Streptomycin (PS, ThermoFisher Scientific) to a confluency of 70-80%. These cells were split into dishes containing coverslips for subsequent temperature jumps. Initially, these cells are in DMEM with 10% FBS only. To the cells, we added a plasmid-Lipofectamine 2000 (Invitrogen, Waltham, MA) complex prepared in OptiMEM media (Gibco). After 6 hours of incubation at 37°C, the media is exchanged for DMEM with 10% FBS and 1% PS. Temperature jumps were performed after 16 hours of incubation at 37°C. For experiments, the cells were imaged in FluroBrite^{TM} DMEM (Gibco) with 10% FBS.

Fast Relaxation Imaging

The FReI instrument and methodology have been described elsewhere (Knab & Davis, 2023; Yoo & Davis, 2022). Briefly, the sample is heated by a continuous-wave 2

µm laser (AdValue Photonics, Tucson, AZ). The heating occurs in the form of a step-shaped function, where the temperature is rapidly raised, followed by a 10 second equilibration period at the higher temperature. The sample starts at room temperature and is raised to 50-60°C in ≈2-3°C increments. The temperature at each step is determined based on calibration with mCherry, which has a well-established temperature-dependent quantum yield. For in-cell FRET measurements, AcGFP1 was excited with blue light selected from a white light LED source (X-Cite Mini +, Excelitas Technology Corporation, Waltham, MA), using a ET480/40x bandpass filter (Chroma, Bellows Falls, VT) and a FF509-FDi01 dichroic mirror (Semrock, West Henrietta, NY). To avoid detection of excitation light, an ET510lp long pass filter (Chroma) selects for emission light. The emission is then separated into green and red light by T510lpxrxt-UF3 and ZT594rdc-UF3 dichroic mirrors (Chroma). The collected light is detected by a CMOS camera (Moment, Teledyne, Thousand Oaks, CA) with an integration time of 200 ms for cells.

Buffer Preparation

All in vitro measurements were performed in 20 mM sodium phosphate, pH 7.4. All in vitro tryptophan melts for the unlabeled protein were performed at a concentration of 10 µM, and all in vitro FRET melts for the labeled protein were performed at a concentration of 0.25 µM. Table S4 contains data for all in vitro melts.

Ficoll PM 70 (Sigma-Aldrich, St. Louis, MO) and M-PER™ (ThermoFisher Scientific, Waltham, MA) melts were performed using tryptophan fluorescence of the unlabeled protein. The Ficoll solutions were prepared by first creating a 400 mg/mL stock solution in 20 mM sodium phosphate, pH 7.4 and then diluting to the final working

concentration. The stock solution was left in a 55°C water bath overnight. Working solutions were further sonicated for 15 minutes prior to addition of the protein.

Melts in salmon sperm DNA (Invitrogen, Waltham, MA) were performed using the FRET-labeled protein. The DNA was dialyzed from its storage buffer into 20 mM sodium phosphate, pH 7.4 for experiments. The final concentration of salmon sperm DNA was 10 mg/mL.

Melts in cytoplasmic and nuclear lysate were performed using the FRET-labeled protein. The lysates were prepared from U2-OS cells at 70-80% confluency, following the manufacturer's protocol for the NE-PER^{TM} Nuclear and Cytoplasmic Extraction Reagents (Thermo Scientific, Waltham, MA). The concentration of protein in the extracts was determined using a Bradford assay. Concentrations were in the range of ≈600-4000 μg/mL, with generally lower yields for nuclear proteins.

Fluorescence Melts

Fluorescence melts were performed on a Jasco FP-8500 (Easton, MD). For tryptophan melts, the excitation wavelength was 280 nm and emission was monitored across 275 to 450 nm. For FRET melts, the excitation wavelength was 475 nm and emission was monitored across 470 to 700 nm. Melts were performed from 20 to 71 °C in 3 °C increments, with an equilibration period of 3 minutes at each temperature.

Preparation of *E. Coli*

The plasmid for FRET-labeled barnase was transformed in BL21(DE3) *E. coli*. The cells were grown at 37°C until the OD$_{600}$ reached 0.6, at which point expression was

induced with 1 mM isopropyl-β-D-thiogalactopyranoside (Inalco, San Luis Obispo, CA). The induced cells grew overnight at 20°C. *E. coli* were prepared for temperature jumps by initially spinning down 10-mL of culture at 4000 rpm for 10 minutes at 4 °C. The resulting pellet was then resuspended in 10-mL of PBS (Corning, Corning, NY) that had been refrigerated overnight. The PBS-suspended cells were then centrifuged at 4000 rpm for 10 minutes at 4 °C, and this process was repeated for a total of 3 times. After the last wash, the pellet was resuspended in 200 μL of PBS. This was used for temperature jumps, with an exposure time of 100-200 ms.

Analysis of Thermodynamic Data

For tryptophan melts, the intensity at the wavelength of maximum emission was plotted against temperature; for FRET melts and temperature jumps, the donor/acceptor ratio (D/A) was plotted against temperature. The resulting melting curve was fit to the following two-state denaturation equation:

$$S(T) = \frac{[\alpha_D + \beta_D(T - 273.15)] + [\alpha_N + \beta_N(T - 273.15)]e^{\frac{-\Delta G}{RT}}}{\left(1 + e^{\frac{-\Delta G}{RT}}\right)} \tag{E1}$$

Where N and D denote the native and denatured states, α_N and α_D are the signals from each state at 0°C, β_N and β_D are the slopes of the baselines for each state, and ΔG is the free energy of folding.

5 Supplementary Material Description

Supplementary materials include a representative circular dichroism and fluorescence melt of unlabeled barnase, representative temperature-dependent behavior of aggregated in-cell populations, a table of protein sequences, and tables of melting temperatures collected in vitro and in living cells.

6 Acknowledgements

This work was supported by National Institutes of Health (NIH) grant R35 GM151146. U.T. was partially supported by the NIH under Biophysics Training Grant T32 GM008283.

7. Conflict of Interests

The authors declare no conflict of interest

References

Alemany, A., Rey-Serra, B., Frutos, S., Cecconi, C., & Ritort, F. (2016). Mechanical

    Folding and Unfolding of Protein Barnase at the Single-Molecule Level.

    Biophysical Journal, *110*(1), 63–74. doi: 10.1016/j.bpj.2015.11.015

Ames, A., & Nesbett, F. B. (1966). Intracellular and extracellular compartments of

    mammalian central nervous tissue. *The Journal of Physiology*, *184*(1), 215–238.

Benton, L. A., Smith, A. E., Young, G. B., & Pielak, G. J. (2012). Unexpected Effects of

    Macromolecular Crowding on Protein Stability. *Biochemistry*, *51*(49), 9773–9775.

    doi: 10.1021/bi300909q

Brocchieri, L., & Karlin, S. (2005). Protein length in eukaryotic and prokaryotic

    proteomes. *Nucleic Acids Research*, *33*(10), 3390–3400. doi: 10.1093/nar/gki615

Christiansen, A., Wang, Q., Samiotakis, A., Cheung, M. S., & Wittung-Stafshede, P.

    (2010). Factors Defining Effects of Macromolecular Crowding on Protein Stability:

    An in Vitro/in Silico Case Study Using Cytochrome c. *Biochemistry*, *49*(31),

    6519–6530. doi: 10.1021/bi100578x

Chu, I.-T., Hutcheson, B. O., Malsch, H. R., & Pielak, G. J. (2023). Macromolecular

    Crowding by Polyethylene Glycol Reduces Protein Breathing. *The Journal of*

    Physical Chemistry Letters, *14*(10), 2599–2605. doi: 10.1021/acs.jpclett.3c00271

Danielsson, J., Mu, X., Lang, L., Wang, H., Binolfi, A., Theillet, F.-X., … Oliveberg, M.

    (2015). Thermodynamics of protein destabilization in live cells. *Proceedings of*

    the National Academy of Sciences, *112*(40), 12402–12407. doi:

    10.1073/pnas.1511308112

Davis, C. M., Deutsch, J., & Gruebele, M. (2020). An in vitro mimic of in-cell solvation
for protein folding studies. *Protein Science: A Publication of the Protein Society*,
29(4), 1060–1068. doi: 10.1002/pro.3833

Davis, C. M., & Gruebele, M. (2018). Non-Steric Interactions Predict the Trend and
Steric Interactions the Offset of Protein Stability in Cells. *ChemPhysChem*,
19(18), 2290–2294. doi: 10.1002/cphc.201800534

Davis, C. M., Gruebele, M., & Sukenik, S. (2018). How does solvation in the cell affect
protein folding and binding? *Current Opinion in Structural Biology*, *48*, 23–29. doi:
10.1016/j.sbi.2017.09.003

Dhar, A., Girdhar, K., Singh, D., Gelman, H., Ebbinghaus, S., & Gruebele, M. (2011).
Protein Stability and Folding Kinetics in the Nucleus and Endoplasmic Reticulum
of Eucaryotic Cells. *Biophysical Journal*, *101*(2), 421–430. doi:
10.1016/j.bpj.2011.05.071

Ebbinghaus, S., Dhar, A., McDonald, J. D., & Gruebele, M. (2010). Protein folding
stability and dynamics imaged in a living cell. *Nature Methods*, *7*(4), 319–323.
doi: 10.1038/nmeth.1435

Fersht, A. R. (1993). Protein folding and stability: The pathway of folding of barnase.
FEBS Letters, *325*(1–2), 5–16. doi: 10.1016/0014-5793(93)81405-O

Fersht, A. R. (2000). A kinetically significant intermediate in the folding of barnase.
Proceedings of the National Academy of Sciences, *97*(26), 14121–14126. doi:
10.1073/pnas.260502597

Guigas, G., Kalla, C., & Weiss, M. (2007). The degree of macromolecular crowding in

    the cytoplasm and nucleoplasm of mammalian cells is conserved. *FEBS Letters*,

    581(26), 5094–5098. doi: 10.1016/j.febslet.2007.09.054

Guzman, I., Gelman, H., Tai, J., & Gruebele, M. (2014). The Extracellular Protein VlsE

    Is Destabilized Inside Cells. *Journal of Molecular Biology*, *426*(1), 11–20. doi:

    10.1016/j.jmb.2013.08.024

Hartley, R. W. (1989). Barnase and barstar: Two small proteins to fold and fit together.

    Trends in Biochemical Sciences, *14*(11), 450–454. doi: 10.1016/0968-

    0004(89)90104-7

Kellermayer, M. (1981). Soluble and "Loosely Bound" Nuclear Proteins in Regulation of

    the Ionic Environment in Living Cell Nuclei. In H. G. Schweiger (Ed.),

    International Cell Biology 1980–1981 (pp. 915–924). Berlin, Heidelberg: Springer

    Berlin Heidelberg. doi: 10.1007/978-3-642-67916-2_103

Kippen, A. D., Sancho, J., & Fersht, A. R. (1994). Folding of Barnase in Parts.

    Biochemistry, *33*(12), 3778–3786. doi: 10.1021/bi00178a039

Kırlı, K., Karaca, S., Dehne, H. J., Samwer, M., Pan, K. T., Lenz, C., … Görlich, D.

    (2015). A deep proteomics perspective on CRM1-mediated nuclear export and

    nucleocytoplasmic partitioning. *eLife*, *4*, e11466. doi: 10.7554/eLife.11466

Knab, E., & Davis, C. M. (2023). Chemical interactions modulate λ6-85 stability in cells.

    Protein Science, *32*(7), e4698. doi: 10.1002/pro.4698

Martinez, J. C., Harrous, M. E., Filimonov, V. V., Mateo, P. L., & Fersht, A. R. (1994). A

    Calorimetric Study of the Thermal Stability of Barnase and Its Interaction with

    3'GMP. *Biochemistry*, *33*(13), 3919–3926. doi: 10.1021/bi00179a018

Meiering, E. M., Serrano, L., & Fersht, A. R. (1992). Effect of active site residues in

    barnase on activity and stability. *Journal of Molecular Biology*, *225*(3), 585–589.

    doi: 10.1016/0022-2836(92)90387-Y

Monteith, W. B., Cohen, R. D., Smith, A. E., Guzman-Cisneros, E., & Pielak, G. J.

    (2015). Quinary structure modulates protein stability in cells. *Proceedings of the*

    National Academy of Sciences, *112*(6), 1739–1742. doi:

    10.1073/pnas.1417415112

Mu, X., Choi, S., Lang, L., Mowray, D., Dokholyan, N. V., Danielsson, J., & Oliveberg,

    M. (2017). Physicochemical code for quinary protein interactions in Escherichia

    coli. *Proceedings of the National Academy of Sciences*, *114*(23), E4556–E4563.

    doi: 10.1073/pnas.1621227114

Paine, P. L., Pearson, T. W., Tluczek, L. J. M., & Horowitz, S. B. (1981). Nuclear

    sodium and potassium. *Nature*, *291*(5812), 258–261. doi: 10.1038/291258a0

Raeburn, C. B., Ormsby, A. R., Cox, D., Gerak, C. A., Makhoul, C., Moily, N. S., …

    Hatters, D. M. (2022). A biosensor of protein foldedness identifies increased

    "holdase" activity of chaperones in the nucleus following increased cytosolic

    protein aggregation. *Journal of Biological Chemistry*, *298*(8). doi:

    10.1016/j.jbc.2022.102158

Rickard, M. M., Zhang, Y., Pogorelov, T. V., & Gruebele, M. (2020). Crowding, Sticking,

    and Partial Folding of GTT WW Domain in a Small Cytoplasm Model. *The*

    Journal of Physical Chemistry B, *124*(23), 4732–4740. doi:

    10.1021/acs.jpcb.0c02536

Rickard, Meredith M., Zhang, Y., Gruebele, M., & Pogorelov, T. V. (2019). In-Cell

  Protein–Protein Contacts: Transient Interactions in the Crowd. *The Journal of*

  Physical Chemistry Letters, *10*(18), 5667–5673. doi: 10.1021/acs.jpclett.9b01556

Sarvas, M., Harwood, C. R., Bron, S., & van Dijl, J. M. (2004). Post-translocational

  folding of secretory proteins in Gram-positive bacteria. *Biochimica et Biophysica*

  Acta (BBA) - Molecular Cell Research, *1694*(1), 311–327. doi:

  10.1016/j.bbamcr.2004.04.009

Schwartz, R., Ting, C. S., & King, J. (2001). Whole Proteome pI Values Correlate with

  Subcellular Localizations of Proteins for Organisms within the Three Domains of

  Life. *Genome Research*, *11*(5), 703–709. doi: 10.1101/gr.158701

Serrano, L., Kellis, J. T., Cann, P., Matouschek, A., & Fersht, A. R. (1992). The folding

  of an enzyme: II. Substructure of barnase and the contribution of different

  interactions to protein stability. *Journal of Molecular Biology*, *224*(3), 783–804.

  doi: 10.1016/0022-2836(92)90562-X

Shankar, R., Schäffer, N., Schmeer, M., Risse, J. M., Friehs, K., & Schleef, M. (2021).

  Recombinant expression of Barnase in Escherichia coli and its application in

  plasmid purification. *Microbial Cell Factories*, *20*, 171. doi: 10.1186/s12934-021-

  01642-y

Shilova, O., Kotelnikova, P., Proshkina, G., Shramova, E., & Deyev, S. (2021). Barnase-

  Barstar Pair: Contemporary Application in Cancer Research and

  Nanotechnology. *Molecules*, *26*(22), 6785. doi: 10.3390/molecules26226785

Smith, A. E., Zhou, L. Z., Gorensek, A. H., Senske, M., & Pielak, G. J. (2016). In-cell

    thermodynamics and a new role for protein surfaces. *Proceedings of the National*

    Academy of Sciences, *113*(7), 1725–1730. doi: 10.1073/pnas.1518620113

Spaar, A., Dammer, C., Gabdoulline, R. R., Wade, R. C., & Helms, V. (2006).

    Diffusional Encounter of Barnase and Barstar. *Biophysical Journal*, *90*(6), 1913–

    1924. doi: 10.1529/biophysj.105.075507

Steinberg, R., & Koch, H.-G. (2021). The largely unexplored biology of small proteins in

    pro- and eukaryotes. *The FEBS Journal*, *288*(24), 7002–7024. doi:

    10.1111/febs.15845

Storz, G., Wolf, Y. I., & Ramamurthi, K. S. (2014). Small Proteins Can No Longer Be

    Ignored. *Annual Review of Biochemistry*, *83*(Volume 83, 2014), 753–777. doi:

    10.1146/annurev-biochem-070611-102400

Tai, J., Dave, K., Hahn, V., Guzman, I., & Gruebele, M. (2016). Subcellular modulation

    of protein VlsE stability and folding kinetics. *FEBS Letters*, *590*(10), 1409–1416.

    doi: 10.1002/1873-3468.12193

The PyMOL Molecular Graphics System. (n.d.). Schrödinger, LLC.

Thole, J. F., Fadero, T. C., Bonin, J. P., Stadmiller, S. S., Giudice, J. A., & Pielak, G. J.

    (2021). Danio rerio Oocytes for Eukaryotic In-Cell NMR. *Biochemistry*, *60*(6),

    451–459. doi: 10.1021/acs.biochem.0c00922

van der Zanden, S. Y., Jongsma, M. L. M., Neefjes, A. C. M., Berlin, I., & Neefjes, J.

    (2023). Maintaining soluble protein homeostasis between nuclear and

    cytoplasmic compartments across mitosis. *Trends in Cell Biology*, *33*(1), 18–29.

    doi: 10.1016/j.tcb.2022.06.002

Vu, N.-D., Feng, H., & Bai, Y. (2004). The Folding Pathway of Barnase: The Rate-Limiting Transition State and a Hidden Intermediate under Native Conditions. *Biochemistry*, *43*(12), 3346–3356. doi: 10.1021/bi0362267

Vuilleumier, S., Sancho, J., Loewenthal, R., & Fersht, A. R. (1993). Circular dichroism studies of barnase and its mutants: Characterization of the contribution of aromatic side chains. *Biochemistry*, *32*(39), 10303–10313. doi: 10.1021/bi00090a005

Wennerström, H., Vallina Estrada, E., Danielsson, J., & Oliveberg, M. (2020). Colloidal stability of the living cell. *Proceedings of the National Academy of Sciences*, *117*(19), 10113–10121. doi: 10.1073/pnas.1914599117

Wood, R. J., Ormsby, A. R., Radwan, M., Cox, D., Sharma, A., Vöpel, T., … Hatters, D. M. (2018). A biosensor-based framework to measure latent proteostasis capacity. *Nature Communications*, *9*(1), 287. doi: 10.1038/s41467-017-02562-5

Yoo, H., & Davis, C. M. (2022). An in Vitro Cytomimetic of In-Cell RNA Folding. *ChemBioChem*, *23*(20), e202200406. doi: 10.1002/cbic.202200406

Zimmerman, S. B., & Trach, S. O. (1991). Estimation of macromolecule concentrations and excluded volume effects for the cytoplasm of Escherichia coli. *Journal of Molecular Biology*, *222*(3), 599–620. doi: 10.1016/0022-2836(91)90499-v