Combined PARP14 Inhibition and PD-1 Blockade Promotes Cytotoxic T Cell Quiescence and

Modulates Macrophage Polarisation in Relapsed Melanoma

Rotem Leshem[1,2,$], Kieran N. Sefton[1,2,$], Chun Wai Wong[1,2,3,4], I-Hsuan Lin[5], Dervla T. Isaac[6],

Mario Niepel[6], and Adam F.L. Hurlstone[1,2,#]

[1]School of Biological Sciences, Faculty of Biology, Medicine and Health, The

University of Manchester, Manchester, M13 9PT, UK

[2]Lydia Becker Institute of Immunology, The University of Manchester, Manchester,

M13 9PT, UK

[3]Now at Massachusetts General Hospital Cancer Center, Department of Medicine,

Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

[4]Now at Broad Institute of MIT and Harvard, Cambridge, MA, USA

[5]Bioinformatics Core Facility, The University of Manchester, Manchester, M13 9PT,

UK

[6]Ribon Therapeutics Inc., 35 Cambridge Park Drive, Suite 300, Cambridge, MA

02140, USA

[$]These authors contributed equally

[#]To whom correspondence should be addressed: Dr Adam Hurlstone, Michael Smith

Building, The University of Manchester, Dover Street, Manchester M13 9PT, UK; tel:

+44 161 2751574; email: adam.hurlstone@manchester.ac.uk

Abstract

Programmed Cell Death 1 (PD-1) signalling blockade is effective in restoring immune surveillance and treating melanoma. However, immune homeostatic mechanisms driven by chronic interferon-gamma (IFNγ) signalling in melanoma cells reduce efficacy, leading to persistent immune evasion and the development of acquired treatment resistance. Previously, we demonstrated that poly ADP ribosyl polymerase 14 (PARP14), encoded by an IFNγ-responsive gene, mediates IFNγ-driven resistance, as inhibition of PARP14 prolonged responsiveness to PD-1 blockade in preclinical models. Nevertheless, inhibition of PARP14 alone was insufficient to fully restore immune clearance, suggesting the presence of redundant resistance mechanisms. The findings here reveal a robust PARP14 catalytic inhibitor (PARP14i) gene signature associated with improved patient survival. We further elucidate the response and adaptation of the immune infiltrate and melanoma cells to PARP14 inhibition using preclinical models. Combining PARP14i with PD-1 blockade therapy revitalised cytotoxic T cells, driving them into a quiescent state while reducing alternatively activated macrophages and increasing pro-inflammatory memory macrophages. Despite this initial enhancement of immune responses through combination therapy, adaptive resistance mechanisms emerged within tumour cells, engaging alternative exhaustion pathways. Our findings indicate that this novel combination therapy leaves immune cells primed for further therapeutic intervention, offering a strategy to address resistance development and optimise treatment outcomes comprehensively.

Keywords: PARP14, T cell exhaustion, quiescent T cells, memory macrophages, adaptive resistance

Introduction

Nivolumab and pembrolizumab, drugs antagonising the PD-1 signalling axis by interfering with PD-1's interaction with Programmed Cell Death Ligand 1 (PD-L1) on exhausted T cells, have vastly improved outcomes for patients with high-risk and late-stage melanoma. However, among responders to PD-1 blockade therapy, approximately 25% will relapse within two years of initiating treatment [1]. Little can be done at present for these patients. Analysis of relapsing tumours has revealed that 30% are characterised by intrinsic—that is, ligand-independent—IFNγ signalling [2]. Intrinsic IFNγ signalling in cancer cells has been shown to follow chronic IFNγ stimulation and to entail epigenetic remodelling wherein signal transducer and activator of transcription 1 (STAT1) and Interferon Regulatory Factor 3 (IRF3) permanently occupy the regulatory sites of a subset of interferon-stimulated genes (ISG), including notably those promoting T cell dysfunction [3]. Indeed, the ISG selectively upregulated in chronically stimulated cancer cells (ISG resistance signature or ISG.RS) predicts poor survival in patients and poor response to PD-1 blockade [4].

Our previous work revealed a contribution from PARP14 in mediating the ability of chronic IFNγ stimulation to drive PD-1 blockade resistance in preclinical models [5]. One model we developed entailed implantation of the mouse rapidly accelerated fibrosarcoma homolog B (BRAF) and Phosphatase and tensin homolog (PTEN) mutant melanoma cell line YUMM2.1 [6] into the hypodermis of syngeneic C57/BL6 mice, and then treating these mice with a short course of anti-PD-1 (α-PD-1) antibodies when tumours became palpable. In tumours that resumed growth following α-PD-1 treatment, we observed increased expression of multiple ISG, including ISG.RS genes, relative to control (untreated) tumours and to tumours still responding to PD-1 blockade therapy [5]. AI-assisted immune profiling, based on

deconvolving gene expression, indicated that relapsing tumours were also relatively more T

cell infiltrated, including by T_{EM} and $T_{CM.}$ However, T cells were also more dysfunctional

compared to these other two tumour groups [5]. The regrowth of such tumours was

significantly delayed—but not abolished— and survival lengthened by adding a highly potent

and selective PARP14 inhibitor RBN012759 (PARP14i, Ribon Therapeutics)[7] to the

treatment regime [5].

This previous study also highlighted that the gene encoding PARP14 is a STAT1 target and

that PARP14 itself is a STAT1 interacting protein [5]. Moreover, in macrophages, PARP14

interaction with STAT1 inhibits the expression of proinflammatory genes [8] but regulates

IRF3 recruitment of RNA polII to promoters of IRF3 target genes [9] and is thus positioned to

influence the expression of ISG. However, the details are far from clear. Certainly, our

previous analysis of gene expression indicated that PARP14 inhibitor treatment of melanoma

cells chronically stimulated with IFNγ increased IFN-, TNF-α-, and IL-6-driven inflammatory

signalling [5].

Intrigued by the potential of PARP14 to influence the expression of ISG in tumour cells but

also the failure of PARP14 inhibitor treatment to permanently flatten tumour rebound after

α-PD-1 treatment, we undertook a detailed analysis of tumour composition and gene

expression changes attending PARP14 inhibitor treatment to identify other targets for

intervention which we now present below.

Materials and Methods

Reagents

Recombinant human IFNγ (PHC4031, Gibco) and recombinant mouse IFNγ (PMC4031,

Gibco). PARP14 inhibitor RBN012759 [7] and PARP14 PROTAC RBN012811 (Ribon

Therapeutics) [10].

Cell lines

Human melanoma cell lines A375 and 501-Mel (provided by Claudia Wellbrock, The

University of Manchester), MC38 cells (provided by Santiago Zelenay, The University of

Manchester) and YUMM2.1 cells (provided by Richard Marais, The University of Manchester)

were maintained in RPMI-1640 (Sigma–Aldrich) supplemented with 10% v/v foetal bovine

serum (Life Technologies) and 1% w/v penicillin-streptomycin (Sigma–Aldrich), at 37 °C in a

5% v/v CO_2 humidified incubator. Cell lines authenticated by STR profiling; cultures routinely

tested for mycoplasma. Cells were treated for two weeks with 50 IU/mL IFNγ, refreshed

every 2-3 days, and subsequently treated with 500nM DMSO (Sigma–Aldrich); 500nM

PARP14i; or 100nM PARP14 PROTAC for 48h.

Mouse tumour implants

Mice were housed in the Biological Services Facility of The University of Manchester on a

12/12 h light/dark cycle and given ad libitum food (Bekay, B&K Universal, Hull, UK) and

water. All procedures were approved by the University of Manchester's animal welfare

ethical review board and performed under relevant Home Office licences according to the

UK Animals (Scientific Procedures) Act, 1986.

8–12-week-old C57BL/6 female mice (ENVIGO) were allowed 1-week to acclimatise.

YUMM2.1 cells (5×10^6 cells) in 100 µL serum-free RPMI-1640 were subcutaneously injected

into the left flank. Tumour volumes (height x width x length calliper measurements) and

mouse weights were monitored every 2–3 days. When tumours reached a volume of 80

mm^3, mice were administered two doses of 300 µg α-PD-1 antibody (BioXCell) in 100 µL

InVivoPure pH 7.0 Dilution Buffer (BioXCell) via intraperitoneal (i.p.) injection administered

at 3-day intervals. Mice were culled if tumours remained under 100 mm^3 (total responders)

or conversely reached 800 mm3 (non-responders). Otherwise, mice with tumours recovering

to 140 mm^3 were randomised on to a second round of treatments (re-treatment): 2 doses of

300 µg of rat isotype control antibody IgG2b (BioXCell) (IgG group) or α-PD-1 antibody

(BioXCell) (α-PD-1 group) in 100 µL InVivoPure pH 7.0 Dilution Buffer (BioXCell) via i.p.

injection at three-day intervals, 14 doses of 500 mg/Kg of PARP14I by oral gavage twice a day

(PARP14i group), or a combination of α-PD-1 i.p. and PARP14i gavage in the same doses

(Combination group). PARP14i was dissolved in 0.5% w/v methylcellulose (Sigma–Aldrich) +

0.2% v/v Tween 80 (Sigma–Aldrich). Dose volume - 10 mL/kg.

Tumour immune infiltrate analysis by flow cytometry

Tumours were dissected upon reaching 250-400mm^3 following retreatment and incubated

for 45 minutes with 100 µg/mL Liberase (Sigma–Aldrich) in serum-free media at 37 °C, then

pushed through a 100 µM nylon cell strainer. The cell suspension was stained for 20 min

with Zombie UV™ Fixable Viability Kit (BioLegend) in PBS. Subsequently, Fc receptors were

blocked, and cells were stained with a surface stain antibody mix. Cells were fixed and

permeabilised using the Foxp3/Transcription Factor Staining Buffer Set (eBioscience)

following manufacturer instructions. Samples were measured on a BD Fortessa flow

cytometer (BDBiosciences), and data was collected using BD FACSDiva™ software. For all

antibodies, a non-stained cell sample and appropriate fluorescence minus one control were

analysed as well. Data were analysed using FlowJo version 8.7.

Antibodies: CD16/32 (clone 93), CD45 (Cat. 103132), CD3ε (Cat. 100306), CD4 (Cat. 100552),

CD8α (Cat. 100742), PD-1 (Cat. 135231), LAG-3 (Cat. 125224), TIM-3 (Cat. 119721), Ki-67

(Cat. 652420), Granzyme B (Cat. 372214), CD69 (Cat. 104531), and CTLA-4 (Cat. 106338)

from Biolegend. Tox (Cat. 12-6502-82) was purchased from eBioscience. Gating strategy as

described in SUPPLEMENTARY FIGURE 5.

Single-cell RNAseq sequencing

Tumour processing

Tumours were collected on day 7 of retreatment into serum-free RPMI on ice, then minced

into 1mm pieces and incubated at 37 °C up to 60 min in Liberase TL (Millipore Sigma) with

rocking according to manufacturer instructions. Digestion was stopped using foetal bovine

serum, and tissue was further dissociated using a wide-bore pipette and filtered through a

cell strainer to achieve a single-cell solution. Immune population enrichment was done by

CD45 mouse Microbeads (Miltenyi) according to manufacturer instructions. Following

sorting, both populations were counted, and ~5% of CD45 negative cells were spiked back

into the CD45 positive fraction.

Cells were fixed for 18 hours, then quenched and stored at -80°C in Enhancer according to

10x Genomics Tissue Fixation & Dissociation for Chromium Fixed RNA Profiling protocol

(CG000553 Rev A) until all samples were collected.

Single-cell isolation and library construction

Gene expression libraries were prepared from formaldehyde-fixed samples using the

Chromium X and Chromium Fixed RNA kit, Mouse Transcriptome, 4 rxns x 16 BC (10x

Genomics, Inc.) according to the manufacturer's protocol (CG000527 Rev C).

Illumina-compatible sequencing libraries were constructed by adding P5, P7, i5 and i7

sample indexes and Illumina Small Read 2 sequences to the 10x barcoded, ligated probe

products via Sample Index PCR followed by a SPRIselect size-selection.

Sequencing

The resulting sequencing library comprised standard Illumina paired-end constructs flanked

with P5 and P7 sequences. The 16 bp 10x Barcode and 12 bp UMI were encoded in Read 1,

while Read 2 sequenced the ligated probe insert, constant sequence and the 8bp Probe

Barcode. Sample indexes were incorporated as the i5 and i7 index reads. Paired-end

sequencing (28:90) was performed on the Illumina NovaSeq6000 platform.

Data Analysis

Raw data processing and cell filtering

Raw sequencing data were processed using the 10x Genomics Cell Ranger pipeline (v7.0.0).

Base call (BCL) files generated by the sequencer were demultiplexed and converted to

FASTQ files using "cellranger mkfastq". The FASTQ files were then mapped against the pre-

built Mouse reference package from 10X Genomics (mm10-2020-A) using "cellranger multi"

to demultiplex and produce the gene-cell barcode matr-ix for individual samples.

The single-cell data were processed in R environment (v4.1) following the workflow

documented in Orchestrating Single-Cell Analysis with Bioconductor [11]. Briefly, for each

sample, the HDF5 file generated by Cell Ranger was imported into R to create a

SingleCellExperiment object. A combination of median absolute deviation (MAD), as

implemented by the "isOutlier" function in the scuttle R package (v1.4.0) and exact

thresholds were used to identify and subsequently remove low-quality cells before data

integration.

Data integration, visualisation and cell clustering

The log-normalised expression values of the combined data were re-computed using the

"multiBatchNorm" function from the bachelor R package (v1.10.0). The per-gene variance of

the log-expression profile was modelled using the "modelGeneVarByPoisson" function from

the scran R package (v1.22.1), and the top 5000 highly variable genes were selected. The

mutual nearest neighbours (MNN) approach implemented by the "fastMNN" function from

the bachelor R package was used to perform batch correction.

The first 50 dimensions of the MNN low-dimensional corrected coordinates for all cells were

used as input to produce the t-stochastic neighbour embedding (t-SNE) projection and

uniform manifold approximation and projection (UMAP) using the "runTSNE" and

"runUMAP" functions from the scater R package (v1.22.0) respectively.

Putative cell clusters were identified using the Leiden algorithm from the igraph R package

(v1.3.0).

Marker genes and differential expression analysis

Cluster-specific markers were identified using the "findMarkers" function from the scran R

package, which performs pairwise t-tests between clusters. Differential expression analysis

between treatment and control (IgG) was performed on pseudo-bulk samples using the

quasi-likelihood pipeline from the edgeR R package (v3.36.0). Genes with a false discovery

rate (FDR) below 5% were considered differentially expressed.

Pathway analysis

Pathway enrichment analysis of differentially expressed genes was performed using the R

package enrichR (v3.0). Pathway results were ranked by the combined scores (calculated as

log(P-value) multiplied by z-score), and pathways with an adjusted P-value < 0.05 were

considered significant.

Further gene set enrichment analysis GSEA was done using Metascape web tool [12] and

plotted as -$\log_{10}$P.

Subclustering and further analysis

Cells were divided into functional cell type groups and further subclustered using the

Louvain method for community detection [13].

Trajectory inference was made using the Slingshot R package [14], and the association of

our PARP14i signature with survival in immune checkpoint blockade therapy(ICBT) cohorts,

as well as a comparison of high and low-expressed genes in specific clusters, was assessed

using Tumour Immune Dysfunction and Exclusion (TIDE) web tool [15]. Sample RL1i

projection on UMAP was done using ProjecTILs R package [16].

Data Availability

Single-cell RNAseq data have been deposited in the ArrayExpress database at EMBL-EBI

under accession number E-MTAB-14471. Bulk RNA sequencing was performed as

previously described [5], with data available from the ArrayExpress database under

accession numbers E-MTAB-12195 and E-MTAB-12872.

Results

A genetic signature reflecting the modulating effect of PARP14i on chronic IFNγ

stimulation predicts response to α-PD-1 in clinical cohorts

We and others have previously shown the role of PARP14 in modulating type II interferon

signalling [5,8,17]. Following chronic stimulation of several mouse and human cancer cell

lines with Interferon-γ (IFNγ) and then subsequent treatment for 48 hours with a PARP14

inhibitor (PARP14i), PARP14 PROTAC or DMSO (drug solvent control), we assessed

differences in gene expression by bulk RNA sequencing (RNAseq). Comparison of the top 50

differentially expressed genes within each cell model revealed a shared expression pattern,

which we narrowed down to a PARP14i signature consisting of 30 genes **(FIGURE 1A,**

SUPPLEMENTARY FIGURE 1 and SUPPLEMENTARY TABLES). Pathway enrichment analysis

performed by Metascape [12] and ARCHS4 [18] revealed that among this signature were

genes strongly correlated with response to IFNγ signalling, with a focus on STAT/IRF

regulation **(FIGURE 1B and SUPPLEMENTARY FIGURE 2).** However, we found this signature is

distinct from the genes contained with the IFNγ signalling GO term, with less than half of the

PARP14i signature displaying overlap. We then investigated whether this PARP14i signature

could predict response to α-PD-1 therapy by employing Tumour Immune Dysfunction and

Exclusion (TIDE) analysis [15]. High expression of genes within this signature was associated

with substantial improvement in overall patient survival across several clinical cohorts

receiving immune checkpoint blockade therapy [19–21] **(FIGURE 1C)**, underscoring the

prognostic value of this signature and supporting the potential therapeutic benefit of

PARP14 inhibition.

Furthermore, we performed RNAseq analysis on subcutaneous tumours comprising

YUMM2.1 cells dissected from mice treated with either IgG or α-PD-1, dividing the latter

into those either relapsing or responding to the treatment. Although the PARP14i signature

was upregulated in α-PD-1-treated samples, it did not distinguish between treatment

response and relapse (**FIGURE 1D**). For further comparison, we also analysed IgG-treated

tumours arising from YUMM2.1 cells pre-treated *ex vivo* with chronic IFNγ before

implantation and found that the PARP14i signature was significantly downregulated,

indicating a Tumour Microenvironment (TME) which is T cell cold and inherently resistant to

α-PD-1.

Analysis of changes to the cellular composition of YUMM2.1 tumours relapsing following

α-PD-1 therapy driven by subsequent PARP14 inhibitor treatment

To investigate further the ability of PARP14i to improve α-PD-1 response, we turned to our

previously described mouse model of acquired resistance [5]: YUMM2.1 cells were

implanted subcutaneously, and upon tumours reaching ~80mm^3 in volume, two doses of α-

PD-1 antibodies were given three days apart **(FIGURE 2A and SUPPLEMENTARY FIGURE 3A)**.

Tumour volumes were routinely recorded and at ~100-150mm^3, animals were randomly

allocated into the following re-treatment groups: IgG2a (n=4); α-PD-1 (n=3); PARP14i alone

(n=3); and a combination of α-PD-1 + PARP14i (n=4). Animals were sacrificed following seven

days of re-treatment, and tumours were subsequently processed for single-cell RNA

sequencing (scRNAseq). Following tumour dissociation and enrichment for CD45$^+$ cells, 5%

from the CD45$^-$ cell fraction was added to ensure tumour cell representation in our analyses.

Cells were fixed and frozen until all tumour samples were collected; then, samples were

barcoded, multiplexed, and libraries prepared and sequenced.

In total, fourteen samples of tumour immune cells were sequenced. Initial analysis of

sequencing data identified one of the IgG samples (RL11i) to be anomalous, which upon

further review of its corresponding tumour growth curve was attributed to it being, in

reality, a non-responder to the initial α-PD-1 treatment, and therefore, it was excluded from

subsequent analyses **(SUPPLEMENTARY FIGURE 3A-C)**. Leiden unsupervised clustering

across the remaining thirteen treated samples identified twenty-eight clusters based upon

gene expression patterns, with a diverse range of cell types represented across samples and

treatment groups **(FIGURE 2B and SUPPLEMENTARY FIGURES 3C and D)**. The following cell

types were identified: Monocytes (Clusters 1-5); Macrophages (6-8); B cells (9-10); Plasma

cells (11); Natural Killer (NK) cells (12); T cells (13-17); Dendritic cells (18-21); Granulocytes

(22-25); Cancer-Associated Fibroblasts (26); and Tumour cells (27-28). To characterise the

individual clusters within these groups, we manually validated cell type identity by

comparing the expression of known markers across clusters **(SUPPLEMENTARY TABLES)**.

Macrophages, Monocytes and T cells were the most abundant leukocyte populations across

all treatment groups, with distinct subpopulations displaying treatment-specific enrichment

(SUPPLEMENTARY FIGURE 3E). Notably, T cells were particularly abundant in combination-

treated samples, with increased prevalence of clusters 14 and 16 and depletion of cluster 13,

which was highly enriched in IgG-treated samples **(FIGURE 2C)**. Similarly, analysis by cell

density plots revealed a shift from cluster 7 in macrophages, prominent in α-PD-1 -treated

samples, towards cluster 6 in the PARP14i and combination treatments **(FIGURE 2D)**.

Furthermore, we observed IgG-specific enrichment and depletion of B cells and monocytes,

respectively. Similarly, across our treatment cohort, few neutrophils and eosinophils were

identified, and these could not be well separated, yet we found the sequenced tumour

sample, which did not respond to α-PD-1 to be highly enriched in neutrophils

(**SUPPLEMENTARY FIGURE 3B**), consistent with a documented role of neutrophils in

promoting rapid tumour growth [22].

PARP14 inhibitor treatment in combination with α-PD-1 therapy drives activated cytotoxic

T cells and limits their transition to dysfunctional states

To further characterise specific T cell identities and assess their functionality, we conducted

subclustering of clusters 12-17 from the initial clustering **(FIGURE 3A).** By combining NK and

T cells for further subclustering, we can be confident that our new T cell clusters are free

from any contaminating readouts due to the presence of NK T cells. This subclustering can

be split broadly into NK cells (1-5) and T cells (6-19), with key subpopulations including naïve

T cells (7 and 19), regulatory T cells (Tregs) (10 and 11), and cytotoxic T cells (13-18).

Comparing the proportions of each T cell subcluster revealed treatment-specific changes in

cellular identity and function **(FIGURE 3B).** Cell density mapping **(FIGURE 3C)** highlighted a

treatment-dependent decrease in naïve T cells (subcluster 7), which were highly abundant in

IgG-treated samples, suggesting their transition into differentiated or activated states. These

naïve T cells also exhibited increased Sell expression compared to other T cell subclusters,

indicative of early activation **(SUPPLEMENTARY FIGURE 4A)**. This suggests that IgG-treated

tumours suffer from inefficient antigen presentation due to dysfunctional priming by

Antigen-Presenting Cells (APCs), which fail to deliver essential differentiation signals to naïve

T cells.

Corroborating our hypothesis that the reduction in naïve T cells is due to their differentiation

into more active T cell states, we identified a significant enrichment amongst all cytotoxic T

cell clusters in the combination treatment group. Detailed gene expression analysis revealed

that T cell subclusters 13-16 exhibit distinct characteristics associated with T cell exhaustion,

quiescence, hyperactivation and T cell response, respectively (**FIGURE 3D & E)**. In the

context of anti-tumour activity, these states are closely interconnected in driving the T cell

effector response. T cell exhaustion is associated with poor clinical outcomes due to loss of T

cell functionality, which fails to control tumour progression [23]. Our slingshot trajectory

analysis (**FIGURE 3F)** indicates that by entering a state of dormancy, quiescent cytotoxic T

cells (subcluster 14) avoid a state of hyperactivity (subcluster 15), also evading exhaustion

(subcluster 13) in the presence of chronic antigen stimulation. Herein, these quiescent cells

can either undergo reactivation that avoids terminal exhaustion (subcluster 16) or persist as

a reservoir of memory T cells with retained effector function, poised for re-activation.

TIDE analysis was performed using gene expression signatures arising from each T cell

subcluster from subclusters 13-16, revealing that high expression of the gene signature for

subcluster 16 (reactivated T cells) was associated with significantly improved survival of

patients receiving immune checkpoint blockade therapy (**FIGURE 3G**). Furthermore, we

observed this was also true, albeit to a lesser degree, for the gene signatures arising from

subclusters 14 (quiescent T cells) and 15 (hyperactive T cells). On the other hand, the gene

signature of T cell subcluster 13 (exhausted T cells) did not associate with positive patient

outcomes, possibly due to heterogeneity in T cell exhaustion phenotypes within this cluster.

This suggests that cells within subclusters 14–16 are critical in mounting an effective anti-

tumour response through mechanisms which sustain effector function without succumbing

to terminal exhaustion.

We next employed Single-Cell rEgulatory Network Inference and Clustering (SCENIC) [24] to

highlight transcriptional regulatory systems that are most pertinent to the functional and

prognostic characteristics of these T cell populations and revealed distinct regulatory

patterns across the T cell subclusters, highlighting the differential activity of several key

transcription factors (**FIGURE 3H and SUPPLEMENTARY FIGURE 4D)**. *Jdp2*, an AP-1 repressor,

exhibited differential regulon activity across the T cell subclusters, being elevated in

subcluster 13 compared to subclusters 14-16. Higher *Jdp2* expression contributes to

impaired cellular functionality by T cell exhaustion and anergy [25,26]. Therefore, lower

activity of the *Jdp2* regulon in subclusters 14 and 15 is consistent with a more functional and

active T cell state and, considering the enrichment of these subclusters with Combination

treatment supports the notion that decreased *Jdp2* activity facilitates better immune

responses. Furthermore, elevated activity of the *Mxd1* regulon in subcluster 13 suggests

this, too, has a crucial role in maintaining the exhausted state characteristic of this

subcluster by repressing cellular growth, contributing to its adverse clinical outcomes [27].

Antagonistic roles of *Mxd1* and *Myc* have been widely reported [28]. It is consistent,

therefore, that the *Myc* regulon was notably active in T cell subclusters 14 and 15, which are

linked to better prognoses and effective anti-tumour responses. Elevated *Myc* activity in

these clusters indicates a role that supports T cell activation and functionality via increased

metabolic and proliferative activities that promote more robust and favourable immune

responses. Taken together, these relationships aid our understanding of ways in which

transcriptional networks influence T cell states and responses within the TME.

The findings from our cytotoxic T cell subclustering and gene expression analyses are further

supported by bulk RNAseq data obtained from the YUMM2.1 subcutaneous tumours

described earlier. Focussing mostly on α-PD-1 relapsing versus responding tumours, these

analyses provide a broader context for understanding the immune dynamics at play in the

TME. In agreement with other studies, IFNγ pre-treatment led to the development of an

'immune desert' in the TME, characterised by a marked reduction in immune cell infiltration.

In contrast, IgG-treated naïve tumours exhibited higher levels of T cell activation and

exhaustion, reflecting an active yet ultimately ineffective immune response

(SUPPLEMENTARY FIGURE 4B). Interestingly, tumours relapsing on α-PD-1 therapy displayed

even higher levels of T cell activation and exhaustion than those treated with IgG alone,

which suggests that despite the robust initial immune response, these tumours managed to

evade sustained immune control. Conversely, α-PD-1 responding tumours displayed lower

levels of exhaustion and elevated markers of T cell hyperactivation, consistent with a robust

anti-tumour response. These bulk RNAseq findings align with our scRNAseq data, indicating

that Combination therapy may successfully replicate the T-cell activation patterns seen in α-

PD-1 responders. As the literature suggests activities which both elicit pro-inflammatory

responses and mediate tumour progression for hyperactivated T cells [29,30], our TIDE

analysis suggests the correlation with positive responses to α-PD-1 may be transient and

insufficient for long-term tumour control, emphasising the need for more durable

therapeutic interventions.

To further explore the immune landscape shaped by Combination therapy, we performed

flow cytometry analysis of α-PD-1 relapsing tumours treated with either α-PD-1 or

Combination therapy. Building on our observations of T cell dynamics, this revealed shifts in

T cell populations following Combination treatment compared to α-PD-1 treated tumours

(FIGURE 3I). Specifically, Combination treatment reduced T cell cytotoxicity **(FIGURE 3J)** and

was accompanied by nuanced changes in the expression of immune checkpoint markers. We

observed a decrease in LAG-3 positivity in PD-1$^+$ CD8$^+$ T cells, associated with reduced T cell

exhaustion and an enhancement of immune response. However, this was counterbalanced

by increased TIM-3 expression, another marker linked to T-cell dysfunction and exhaustion

(FIGURE 3K) [31,32]. Further examination of PD-1$^+$CTLA-4$^+$ CD8$^+$ T cells showed that

Combination treatment had limited impact on reversing terminal exhaustion, likely due to

the combined inhibitory effects of multiple markers **(FIGURE 3L)**. Further analysis, focused

on the PD-1$^+$CTLA-4$^-$ population, suggested that LAG-3 and CTLA-4 are co-dependent in this

context **(FIGURE 3M)**. This suggests that targeting each alone as a further intervention may

prove ineffective. Together, these findings highlight the complexity and heterogeneity of T

cell states in the TME that influence therapeutic efficacy.

422 **PARP14 inhibitor treatment increases M1 macrophage polarisation to generate a memory-**

423 **like state**

424 Given the previous demonstration that PARP14i can suppress M2 macrophage

425 differentiation and mimic α-PD-1 treatment in renal tumour explants [7], we expanded our

426 investigation to the macrophage populations in the TME. By further subclustering clusters 1-

427 8 from the initial analysis, we created a new MonoMac UMAP representation detailing a

428 clear division of these subclusters into either those which exhibited characteristics of M1

429 macrophages (Left) or M2 macrophages (Right) (**FIGURE 4A** and **SUPPLEMENTARY TABLES)**.

430

431 Analysis of cell type proportions within this subclustering revealed treatment-specific

432 changes. Combination treatment notably shifted macrophage polarisation from M2 towards

433 M1 (**FIGURE 4B**). Further inspection of gene enrichment analysis of DEGs, applied alongside

434 cell density mapping, allowed us to robustly identify subclusters most affected by treatment

435 **(FIGURE 4C)**. This demonstrated that subcluster 10— the most abundant cluster in the IgG-

436 treated samples— is effectively depleted by Combination treatment **(FIGURE 4D)**. This

437 subcluster consists primarily of tumour-associated macrophages (TAMs), which are known to

438 facilitate tumour progression and metastasis, and as such, their reduction by Combination

439 treatment disrupts pro-tumoural mechanisms.

440

441 These anti-tumoural effects are further supported by the expansion of specific macrophage

442 clusters. The increase in M1 macrophages (subcluster 2) signifies an enhanced pro-

443 inflammatory response through Th1 pathways that bolsters anti-tumour immunity. Similarly,

444 the increased abundance of a transitory M1/M2 macrophage state (subcluster 4) further

445 illustrates the dynamic changes in macrophage populations, highlighting cellular plasticity

modulated by Combination treatment to influence the balance between inflammatory and

immunoregulatory activities. Concurrently, macrophage subcluster 8, representing M2a

macrophages, exhibits an expression pattern that supports tissue repair and dampening

excessive inflammation in the tumour.

The overall increase in M1 macrophages following Combination treatment is linked to

reduced *Tgm2*, *Vegfa* and *Cebpb* expression, key drivers of M2 macrophage polarisation

(FIGURE 4E) [33]. Their expression patterns observed in subclusters 8 and 10 align with the

depletion of M2-related clusters and the relative enrichment of M2a macrophages induced

by Combination treatment, reinforcing the transition towards a more hostile tumour

environment. Additionally, PARP14i treatment results in a marked downregulation of *Nfatc1*,

Nfatc3 and *Smad4* across subclusters (**FIGURE 4E)**, reducing *Tgfb1* signalling in M2

macrophages, as evidenced by lower *Sox2* and *Sox4* expression [34]. This corresponds with

diminished pro-inflammatory gene expression, while the decreased expression of *Stat3*,

Stat6 and *Jun* in subclusters 2 and 4 correlates with reduced transcriptional activity and

cellular activation. These changes suggest subclusters 2 and 4 are similar to T cell subcluster

14 in that they may represent memory-like cells with the potential for re-activation [35].

Crucially, the interaction between immune cell subsets largely defines their activities.

Contact between PILRA and CD8α was previously shown to maintain cytotoxic T cells in a

quiescent state [36]. Therefore, elevated *Pilra* expression in M1-like subclusters likely serves

as a barrier to prevent premature re-activation (**FIGURE 4E)**. Given that *Pilra* correlates

positively with tumour immune infiltration and prognosis [37], its elevated expression in

subclusters enriched by Combination treatment supports the establishment of quiescent T

cells.

To further explore the molecular underpinnings of macrophage polarisation, we applied

SCENIC analysis to evaluate transcriptional regulatory programs within these subclusters

(FIGURE 4F). Here, we identified regulons which effectively delineated M1 (*Runx3* and *Cux1*)

and M2 (*Irf4*) macrophages. Additionally, six regulons associated with M2 macrophages

(*Nfia, Nr1d1, Klf2, Klf3, Etv1, Dbp*) were most strongly associated with subcluster 10. In the

context of Combination treatment, the depletion of subcluster 10 resulted in a significant

reduction in the activity of these regulons. However, this reduction may be offset by the

enrichment of subcluster 8, indicative of a directional shift in this treatment towards an M1-

like phenotype. These findings suggest that the overall balance between M1 and M2

macrophages is shifting towards a pro-inflammatory, anti-tumour state in Combination

treatment. Notably, subclusters 2 and 4, despite their relatively low regulon activity beyond

the M1-defining features, may represent more memory-like macrophage states.

Combined α-PD-1 and PARP14 inhibitor treatment provokes dynamic adaptive resistance

strategies in tumour cells

To further understand the impact of PARP14 inhibition on the TME, we further subclustered

tumour cells and Cancer-Associated Fibroblasts (CAFs)— covering clusters 26-28— into six

distinct subclusters **(FIGURE 5A)**, each exhibiting different characteristics which reflect their

roles in tumour growth, progression and adaptive responses to therapy. For instance,

compared with IgG, α-PD-1 treatment alone effectively doubled the size of tumour

subcluster 3, comprising stem-like cells (**FIGURE 5B and C**). This tumour subcluster was

found by slingshot trajectory analysis [14] to be the origin of tumour cell types associated

with immunomodulation and immune evasion— subclusters 4 and 5— (**FIGURE 5D**). Their

stem-like qualities are particularly evident when surveying Gene Set Enrichment Analysis

(GSEA) outputs for this tumour subcluster (**FIGURE 5E**). Therefore, suppression of this

subcluster by combination treatment indicates how a more potent treatment response is

associated with reducing the stemness of tumour cells, thereby limiting self-renewal

capacity and disrupting cell survival pathways.

In contrast, tumour subclusters 1, 2 and 4 were enriched by Combination treatment.

Tumour-subcluster 2, the most prominent of these and notable for its highly invasive and

adhesive cells, plays a crucial role in tumour metastasis through extracellular matrix

remodelling. Its enrichment demonstrates, we infer, an adaptive response to evade immune

attack under intense therapeutic pressure, as evidenced by increased activation of

biosynthetic pathways (**FIGURE 5E**). Similarly, the angiogenic and epithelial-mesenchymal

transition (EMT) phenotypes shared by cells within tumour subcluster 1 reflect their

migratory and metastatic capacity. As shown by pathway enrichment analyses, DEGs in this

subcluster support tumour growth, vascularisation and metastasis whilst also dampening

immune responses (**FIGURE 5E**). The increased activation of metabolic pathways in tumour

subcluster 4 suggests that the Combination treatment enhances proliferative potential and

cellular turnover, which could inadvertently support aggressive tumour growth. This

behaviour is characteristic of tumours with high resistance to therapy, which sustain growth

and repair mechanisms despite treatment. Furthermore, GSEA reveals these strategies likely

involve elevated angiogenesis, cell expansion and interference with immune cell recognition

machinery (**FIGURE 5E**). Therefore, while treatment suppresses key tumour-promoting

pathways such as stemness, it can also select tumour adaptations which permit immune

escape.

Across these tumour subclusters, we evaluated the top 200 upregulated genes and extracted

those uniquely upregulated in each subcluster. We then visualised the expression of each

subcluster's gene list in our bulk RNAseq data for the previously described YUMM2.1

subcutaneous tumours. Here, we found that the gene lists for subclusters 1, 2 and 5 were

more strongly associated with tumours relapsing on α-PD-1 treatment, whereas genes from

subclusters 3 and 4 correlated with tumours that responded to α-PD-1 (**SUPPLEMENTARY**

FIGURE 6A). Furthermore, when evaluating these unique gene lists using TIDE, these were

largely in agreement with our identification of relapse or response outcomes with subcluster

4, which responds to α-PD-1 treatment, resulting in improved patient survival

(**SUPPLEMENTARY FIGURE 6B).** Similarly, signatures for subclusters relapsing on α-PD-1

therapy were associated with worse clinical outcome. However, concerning genes uniquely

upregulated in subcluster 3, the inability of this signature to predict clinical response

suggests that this subcluster of cells can follow different paths under selective pressure,

transitioning into states that are either susceptible to immune attack or capable of evading

immune defences. It is this balance between vulnerability and evasion that ultimately

determines the overall treatment outcome.

When next we queried the previously identified PARP14i signature within the subclusters,

we observed treatment- and subcluster-specific regulation of many genes within this

signature **(FIGURE 5F)**. In tumour subcluster 1, Combination treatment upregulated many of

these genes, indicating elevated interferon signalling, while in subcluster 4, these are

downregulated, reflecting reduced metabolic and proliferative activity. Similarly, most genes

within this signature are enriched in tumour subcluster three following α-PD-1 treatment

alone, yet Combination treatment limits PD-1-driven upregulation for many of these,

potentially to prevent chronic stimulation. In subcluster 2, responses were more gene-

specific, with both PARP14i and combination treatments reducing *Ifit3* and elevating *Gbp7*

expression, suggesting a shift towards increased Type II interferon signalling.

Discussion

The therapeutics currently used for melanoma, particularly immune checkpoint inhibitors

(ICIs) such as α-PD-1, have made a significant impact but still face considerable challenges.

One of the most persistent problems is tumour resistance. Although combination therapies

and targeted agents have been investigated, they have only been partially successful in

overcoming resistance mechanisms [38,39]. This ongoing challenge underscores the

necessity for novel interventions that can enhance treatment efficacy by more effectively

addressing resistance. Our previous research demonstrated that PARP14, a crucial regulator

of immune responses and chromatin dynamics, can modulate the TME and that its inhibition

boosts the activity of infiltrating T cells [5]. However, these findings were limited by an

incomplete understanding of how PARP14 impacts immune cell dynamics. In this study, we

address these gaps by investigating the effects of combining PARP14 inhibition with α-PD-1

therapy on the tumour immune infiltrate at the onset of α-PD-1 resistance.

A central challenge in overcoming resistance is T cell exhaustion, where prolonged antigen

exposure causes T cells to lose their functional capacity [40,41]. This state is characterised by

the upregulation of inhibitory receptors and a corresponding decline in cytokine production

and cytotoxic activity. Rather than inducing terminal exhaustion, we found that the

combination therapy triggered a unique strategy to evade exhaustion characterised by

hyperactivated T cells entering a state of quiescence. Elevated expression of genes

associated with T cell hyperactivation contribute to a pro-inflammatory environment;

however, this may exacerbate tumour progression if inflammation is excessive [29] rather

than effectively controlling it [30]. This implies that to maintain a productive anti-tumour

response, T cells must employ an exit strategy, such as entering quiescence, which prevents

terminal T-cell exhaustion but also limits hyperactivation. Unlike classical exhaustion, where

T cells lose all activation markers, the quiescent T cells in our study retained a gene

expression pattern associated with longevity (eg. *Bcl2*, *Il-6*) compared with exhausted states.

This reservoir could potentially be leveraged by integrating further therapeutics designed to

reactivate these T cells.

The role of Pilra interactions as a driver of T cell quiescence presents a potential axis for such

an intervention. Intriguingly, the Pilra ligand CD8α was highly expressed in T cell subcluster

14 and even more so in subcluster 16, suggesting that overcoming PILRA inhibition may be

necessary to exit the quiescent state. Predictions of better patient prognosis in tumours

highly expressing *Pilra* support a strategy to evade T cell exhaustion by entering quiescence

[37]. Although the use of a monoclonal antibody therapeutic specific to PILRA has not yet

been explored in cancer treatment, preclinical models have shown promising results in the

context of autoinflammatory diseases [42,43]. Targeting the PILRA-CD8α axis could thus

reinvigorate quiescent T cells to generate T cell states with a more active anti-tumour

response (e.g. T cell subcluster 16).

Moreover, PARP14 inhibition reduced immunosuppressive signals, fostering a more pro-

inflammatory environment that enhanced T cell activation and function. This effect is

particularly noteworthy as a potential means of counteracting the T cell dysfunction often

associated with markers of ICI resistance, such as chronic IFNγ exposure. We observed that

inhibiting PARP14 induced a significant shift from an immunosuppressive TME, typically

linked with M2 macrophages, to a more inflammatory state characterised by increased

polarisation towards an M1 phenotype, in keeping with previous findings [7,8]. M2

macrophages are typically exploited by tumours for their roles in tissue repair and

immunosuppression to create a microenvironment that facilitates tumour progression

[44,45]. Conversely, M1 macrophages, known for their production of pro-inflammatory

cytokines such as TNF-α, IL-1β, and IL-6, are instrumental in orchestrating and sustaining

effective anti-tumour immune responses [46,47]. Therefore, by generating a more

immunostimulatory and anti-tumour environment, PARP14 inhibition can be utilised to

overcome immune evasion strategies employed by tumours [48]. In addition to their direct

anti-tumour effects, enrichment of M1 macrophages by PARP14i also promotes the

production of cytokines and chemokines that recruit and activate T cells [49,50].

Consequently, M1 macrophages can enhance the effectiveness of PD-1 blockade therapy by

increasing the infiltration and activation of cytotoxic T cells within the tumour.

The activity of PARP14 in chromatin remodelling and gene expression regulation impacts key

immune pathways through its interaction with *STAT1* [8], *STAT3* [51] and *STAT6* [52].

Following cytokine stimulation, PARP14 modulates the accessibility of these promoter

regions through its interaction with HDACs, facilitating transcription factor recruitment [52].

Recent studies highlight the critical role of epigenetic regulation for macrophage

polarisation, which determines their functional state, whilst quiescent T cells actively

manage their epigenetic landscape to remain poised for rapid reactivation while preserving

their long-term functionality [53]. In the context of combination therapy, our PARP14i

intervention occurs following this stimulation. Thus, we speculate that PARP14 inhibition

prevents the re-recruitment of HDACs to the promoter regions, leading to a "dormant"

memory state in T cells and macrophages while retaining essential readiness for reactivation

of immune cell responsiveness. This illustrates the influence of PARP14 upon the regulation

of chromatin dynamics, with the capacity to alter immune cell responsiveness and

functionality in the TME, emphasising the therapeutic potential of modulating such

pathways.

Despite the positive effects of our novel therapeutic combination, our study also

underscores the concept of dynamic adaptive resistance, wherein tumour cells rapidly

evolve in response to therapeutic pressure. Although the combination of α-PD-1 and PARP14

inhibition initially enhances immune responses, tumour cells can engage alternative

pathways and modulate immune checkpoints, allowing them to evade immunotherapy and

persist despite intensified immune pressure. This observation highlights the remarkable

adaptability within the TME and aligns with findings from other studies that have reported

similar resistance phenomena in response to targeted therapies [54,55].

Our investigation revealed a robust PARP14 inhibition (PARP14i) gene signature that is

strongly associated with IFNγ signalling and STAT/IRF pathway regulation. While this

signature was upregulated in α-PD-1-treated samples, it did not differentiate between

treatment response and relapse in our preclinical model. Nonetheless, the high expression

of these genes correlates with improved patient survival, supporting the potential of

PARP14i to enhance the efficacy of PD-1 blockade. This suggests that, despite an initial

sensitivity to PD-1 blockade resulting in total tumour eradication, residual YUMM2.1 cells

that have developed resistance to α-PD-1 adapt rapidly to further therapeutic

interventions—possibly by acquiring more stem-like characteristics over time, which

subsequently enable tumour growth even in the presence of a re-established immune

response. Consequently, the subcluster-specific effects of combination therapy in tumour

cells indicate that, although combination therapy enhances T cell activation and

inflammation, alterations in the expression of the PARP14i gene signature combined with

mounting immune pressure when combined with α-PD-1 treatment may simultaneously

facilitate the development of resistance mechanisms that allow tumour persistence. Notably,

the upregulation of *Stat1*, *Atf3*, *Irf1*, and *Cd274* in the EMT angiogenic tumour cells within

tumour subcluster 1 suggests that combination treatment may inadvertently enhance EMT-

associated properties, thereby promoting more invasive and immune-evasive tumour cells

[56]. Conversely, given the role of IFNγ in regulating many genes within this signature, the

downregulation of these genes in tumour cells of tumour subcluster 4 suggests reduced

intrinsic IFN signalling, which could potentially increase their susceptibility to immune-

mediated destruction despite their high proliferative activity [57]. This dual effect

underscores the complexity of targeting regulatory pathways involved in tumour immunity

and illustrates the delicate balance between tumour cell expansion and immune attack.

The observed adaptive resistance underscores the necessity for a multifaceted approach to

cancer treatment. As adaptive resistance mechanisms can evolve rapidly, our findings

indicate that the initial benefits of combining α-PD-1 therapy with PARP14 inhibition may be

counteracted by subsequent resistance. Therefore, future research must focus on

developing multi-modal strategies that integrate immune checkpoint inhibitors with agents

that both target the tumour and enhance a pro-inflammatory immune infiltrate within the

TME, potentially leading to more durable patient responses. Moreover, our identification of

a PARP14i gene signature offers therapeutic promise, particularly for patients in whom

expression of these genes is low. By utilising this gene signature as a biomarker, clinicians

can more accurately identify patients who are likely to benefit from PARP14 inhibition,

facilitating a more personalised and targeted treatment approach. This not only optimises

treatment efficacy but also helps avoid unnecessary interventions in patients who are less

likely to respond. Additionally, there is potential to further refine this approach by

identifying supplementary interventions that could be combined with PARP14 inhibition. The

precise mechanism that unlocks T cell potential in the Combination treatment is unclear,

requiring further investigation to determine whether combination treatment prevents

tumour cells from down-regulating their antigens through dedifferentiation or reverses an

inhibitory mechanism impairing T cell-APC interactions. If future studies can identify agents

that either target complementary pathways or prevent the adaptive resistance observed in

tumour cells, it may be possible to develop more robust combination therapies.

Interventions that synergise with PARP14i and α-PD-1 could potentially overcome even the

most resilient tumour populations, ultimately providing a more durable and comprehensive

therapeutic response.

Declarations

Acknowledgements

This study was funded by a research sponsorship agreement between Ribon Therapeutics

and The University of Manchester, grant 821034 from the Melanoma Research Alliance

(MRA), grant 22-0091 from Worldwide Cancer Research (WCR), and an award from the Skin

Cancer Research Fund (SCaRF). C.W.W. was funded by a scholarship from the Hong Kong

Scholarship for Excellence Scheme (HKSES). We thank Dr Gareth Howell and The University

of Manchester Flow Cytometry Core Facility for facilitating flow cytometry analysis, Dr Andy

Hayes and Claire Morrisroe in the Genomic Technologies Core Facility for assistance with

single-cell RNA sequencing, Dr Leo Zeef and the Bioinformatics Core facility for their help

with bioinformatic analysis, and members of the Biological Services Facility at The University

of Manchester for help with animal work. FIGURE 2A and Graphical Abstract were created

using BioRender.

Author Contributions Statement

A.H. conceived the study. R.L., K.N.S., and C.W.W. performed experiments. I.L. and R.L.

analysed the scRNAseq data. K.N.S. analysed the flow cytometry and bulk RNAseq data.

D.T.I., M.N., and A.H. contributed to experimental design and data analysis. R.L., K.N.S., and

A.H. drafted the original version of this manuscript, with all authors reviewing subsequent

drafts. A.H. obtained funding for the study.

Competing Interests Statement

The authors declare the following competing interests:

D.T.I. and M.N. were employees and shareholders of Ribon Therapeutics at the time of data

collection. A.H. received research sponsorship from Ribon Therapeutics. All other authors

declare no competing interests.

References

[1] Ribas A, Hamid O, Daud A, Hodi FS, Wolchok JD, Kefford R, et al. Association of Pembrolizumab With Tumor Response and Survival Among Patients With Advanced Melanoma. JAMA 2016;315:1600. https://doi.org/10.1001/jama.2016.4059.

[2] Lim J, Kang I, La J, Ku KB, Kang BH, Kim Y, et al. Harnessing type I interferon-mediated immunity to target malignant brain tumors. Front Immunol 2023;14:1203929. https://doi.org/10.3389/fimmu.2023.1203929.

[3] Qiu J, Xu B, Ye D, Ren D, Wang S, Benci JL, et al. Cancer cells resistant to immune checkpoint blockade acquire interferon-associated epigenetic memory to sustain T cell dysfunction. Nat Cancer 2023. https://doi.org/10.1038/s43018-022-00490-y.

[4] Benci JL, Johnson LR, Choa R, Xu Y, Qiu J, Zhou Z, et al. Opposing Functions of Interferon Coordinate Adaptive and Innate Immune Responses to Cancer

Immune Checkpoint Blockade. Cell 2019;178:933-948.e14.
https://doi.org/10.1016/j.cell.2019.07.019.
[5] Wong CW, Evangelou C, Sefton KN, Leshem R, Zhang W, Gopalan V, et al.
PARP14 inhibition restores PD-1 immune checkpoint inhibitor response
following IFNγ-driven acquired resistance in preclinical cancer models. Nat
Commun 2023;14:5983. https://doi.org/10.1038/s41467-023-41737-1.
[6] Meeth K, Wang JX, Micevic G, Damsky W, Bosenberg MW. The YUMM lines: a
series of congenic mouse melanoma cell lines with defined genetic alterations.
Pigment Cell Melanoma Res 2016;29:590–7.
https://doi.org/10.1111/pcmr.12498.
[7] Schenkel LB, Molina JR, Swinger KK, Abo R, Blackwell DJ, Lu AZ, et al. A
potent and selective PARP14 inhibitor decreases protumor macrophage gene
expression and elicits inflammatory responses in tumor explants. Cell Chem
Biol 2021;28:1158-1168.e13. https://doi.org/10.1016/j.chembiol.2021.02.010.
[8] Iwata H, Goettsch C, Sharma A, Ricchiuto P, Goh WWB, Halu A, et al. PARP9
and PARP14 cross-regulate macrophage activation via STAT1 ADP-ribosylation.
Nat Commun 2016;7:12849. https://doi.org/10.1038/ncomms12849.
[9] Caprara G, Prosperini E, Piccolo V, Sigismondo G, Melacarne A, Cuomo A, et
al. PARP14 Controls the Nuclear Accumulation of a Subset of Type I IFN–
Inducible Proteins. J Immunol 2018;200:2439–54.
https://doi.org/10.4049/jimmunol.1701117.
[10] Wigle TJ, Ren Y, Molina JR, Blackwell DJ, Schenkel LB, Swinger KK, et al.
Targeted Degradation of PARP14 Using a Heterobifunctional Small Molecule.
ChemBioChem 2021;22:2107–10. https://doi.org/10.1002/cbic.202100047.
[11] Amezquita RA, Lun ATL, Becht E, Carey VJ, Carpp LN, Geistlinger L, et al.
Orchestrating single-cell analysis with Bioconductor. Nat Methods 2020;17:137–
45. https://doi.org/10.1038/s41592-019-0654-x.
[12] Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al.
Metascape provides a biologist-oriented resource for the analysis of systems-
level datasets. Nat Commun 2019;10:1523. https://doi.org/10.1038/s41467-019-
09234-6.
[13] Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. Fast unfolding of
communities in large networks. J Stat Mech Theory Exp 2008;2008:P10008.
https://doi.org/10.1088/1742-5468/2008/10/P10008.
[14] Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell
lineage and pseudotime inference for single-cell transcriptomics. BMC
Genomics 2018;19:477. https://doi.org/10.1186/s12864-018-4772-0.
[15] Fu J, Li K, Zhang W, Wan C, Zhang J, Jiang P, et al. Large-scale public data
reuse to model immunotherapy response and resistance. Genome Med
2020;12:21. https://doi.org/10.1186/s13073-020-0721-z.
[16] Andreatta M, Corria-Osorio J, Müller S, Cubas R, Coukos G, Carmona SJ.
Interpretation of T cell states from single-cell transcriptomics data using
reference atlases. Nat Commun 2021;12:2965. https://doi.org/10.1038/s41467-
021-23324-4.
[17] Higashi H, Maejima T, Lee LH, Yamazaki Y, Hottiger MO, Singh SA, et al. A
Study into the ADP-Ribosylome of IFN-γ-Stimulated THP-1 Human
Macrophage-like Cells Identifies ARTD8/PARP14 and ARTD9/PARP9 ADP-
Ribosylation. J Proteome Res 2019;18:1607–22.
https://doi.org/10.1021/acs.jproteome.8b00895.

[18] Lachmann A, Torre D, Keenan AB, Jagodnik KM, Lee HJ, Wang L, et al.
Massive mining of publicly available RNA-seq data from human and mouse. Nat
Commun 2018;9:1366. https://doi.org/10.1038/s41467-018-03751-6.
[19] Nathanson T, Ahuja A, Rubinsteyn A, Aksoy BA, Hellmann MD, Miao D, et al.
Somatic Mutations and Neoepitope Homology in Melanomas Treated with
CTLA-4 Blockade. Cancer Immunol Res 2017;5:84–91.
https://doi.org/10.1158/2326-6066.CIR-16-0019.
[20] Mariathasan S, Turley SJ, Nickles D, Castiglioni A, Yuen K, Wang Y, et al. TGFβ
attenuates tumour response to PD-L1 blockade by contributing to exclusion of T
cells. Nature 2018;554:544–8. https://doi.org/10.1038/nature25501.
[21] Gide TN, Quek C, Menzies AM, Tasker AT, Shang P, Holst J, et al. Distinct
Immune Cell Populations Define Response to Anti-PD-1 Monotherapy and Anti-
PD-1/Anti-CTLA-4 Combined Therapy. Cancer Cell 2019;35:238-255.e6.
https://doi.org/10.1016/j.ccell.2019.01.003.
[22] Wang X, Qiu L, Li Z, Wang X-Y, Yi H. Understanding the Multifaceted Role of
Neutrophils in Cancer and Autoimmune Diseases. Front Immunol 2018;9.
https://doi.org/10.3389/fimmu.2018.02456.
[23] Wherry EJ, Kurachi M. Molecular and cellular insights into T cell exhaustion. Nat
Rev Immunol 2015;15:486–99. https://doi.org/10.1038/nri3862.
[24] Aibar S, González-Blas CB, Moerman T, Huynh-Thu VA, Imrichova H,
Hulselmans G, et al. SCENIC: single-cell regulatory network inference and
clustering. Nat Methods 2017;14:1083–6. https://doi.org/10.1038/nmeth.4463.
[25] Yuanhong X, Feng X, Qingchang L, Jianpeng F, Zhe L, Kejian G.
Downregulation of Ap-1 Repressor Jdp2 is Associated with Tumor Metastasis
and Poor Prognosis in Patients with Pancreatic Carcinoma. Int J Biol Markers
2010;25:136–40. https://doi.org/10.1177/172460081002500303.
[26] Atsaves V, Leventaki V, Rassidakis GZ, Claret FX. AP-1 Transcription Factors
as Regulators of Immune Responses in Cancer. Cancers 2019;11:1037.
https://doi.org/10.3390/cancers11071037.
[27] Iritani BM, Delrow J, Grandori C, Gomez I, Klacking M, Carlos LS, et al.
Modulation of T-lymphocyte development, growth and cell size by the Myc
antagonist and transcriptional repressor Mad1. EMBO J 2002;21:4820.
https://doi.org/10.1093/emboj/cdf492.
[28] Lüscher B. MAD1 and its life as a MYC antagonist: An update. Eur J Cell Biol
2012;91:506–14. https://doi.org/10.1016/j.ejcb.2011.07.005.
[29] Kalfaoglu B, Almeida-Santos J, Tye CA, Satou Y, Ono M. T-Cell Hyperactivation
and Paralysis in Severe COVID-19 Infection Revealed by Single-Cell Analysis.
Front Immunol 2020;11. https://doi.org/10.3389/fimmu.2020.589380.
[30] Palmer DC, Webber BR, Patel Y, Johnson MJ, Kariya CM, Lahr WS, et al.
Internal checkpoint regulates T cell neoantigen reactivity and susceptibility to
PD1 blockade. Med 2022;3:682-704.e8.
https://doi.org/10.1016/j.medj.2022.07.008.
[31] Avery L, Filderman J, Szymczak-Workman AL, Kane LP. Tim-3 co-stimulation
promotes short-lived effector T cells, restricts memory precursors, and is
dispensable for T cell exhaustion. Proc Natl Acad Sci 2018;115:2455–60.
https://doi.org/10.1073/pnas.1712107115.
[32] Minnie SA, Waltner OG, Zhang P, Takahashi S, Nemychenkov NS, Ensbey KS,
et al. TIM-3+ CD8 T cells with a terminally exhausted phenotype retain
functional capacity in hematological malignancies. Sci Immunol
2024;9:eadg1094. https://doi.org/10.1126/sciimmunol.adg1094.

[33] Sestito C, Brevé JJP, Bol JGJM, Wilhelmus MMM, Drukarch B, van Dam A-M.
860   Tissue Transglutaminase contributes to myelin phagocytosis in interleukin-4-
861   treated human monocyte-derived macrophages. Cytokine 2020;128:155024.
862   https://doi.org/10.1016/j.cyto.2020.155024.
863   [34] Liu Z, Kuang W, Zhou Q, Zhang Y. TGF-β1 secreted by M2 phenotype
864   macrophages enhances the stemness and migration of glioma cells via the
865   SMAD2/3 signalling pathway. Int J Mol Med 2018;42:3395.
866   https://doi.org/10.3892/ijmm.2018.3923.
867   [35] Kloc M, Kubiak JZ, Zdanowski R, Ghobrial RM. Memory Macrophages. Int J Mol
868   Sci 2022;24:38. https://doi.org/10.3390/ijms24010038.
869   [36] Zheng L, Han X, Yao S, Zhu Y, Klement J, Wu S, et al. The CD8α-PILRα
870   interaction maintains CD8+ T cell quiescence. Science 2022;376:996–1001.
871   https://doi.org/10.1126/science.aaz8658.
872   [37] Li Q, Yang Z, He X, Yang X. Comprehensive analysis of PILRA's association
873   with the prognosis, tumor immune infiltration, and immunotherapy in pan-
874   cancer. Sci Rep 2023;13:14334. https://doi.org/10.1038/s41598-023-41649-6.
875   [38] Huang AC, Zappasodi R. A decade of checkpoint blockade immunotherapy in
876   melanoma: understanding the molecular basis for immune sensitivity and
877   resistance. Nat Immunol 2022;23:660–70. https://doi.org/10.1038/s41590-022-
878   01141-1.
879   [39] Rager T, Eckburg A, Patel M, Qiu R, Gantiwala S, Dovalovsky K, et al.
880   Treatment of Metastatic Melanoma with a Combination of Immunotherapies and
881   Molecularly Targeted Therapies. Cancers 2022;14:3779.
882   https://doi.org/10.3390/cancers14153779.
883   [40] Wherry EJ, Kurachi M. Molecular and cellular insights into T cell exhaustion. Nat
884   Rev Immunol 2015;15:486–99. https://doi.org/10.1038/nri3862.
885   [41] Collier JL, Weiss SA, Pauken KE, Sen DR, Sharpe AH. Not-so-opposite ends of
886   the spectrum: CD8+ T cell dysfunction across chronic infection, cancer and
887   autoimmunity. Nat Immunol 2021;22:809–19. https://doi.org/10.1038/s41590-
888   021-00949-7.
889   [42] Sun Y, Caplazi P, Zhang J, Mazloom A, Kummerfeld S, Quinones G, et al.
890   PILRα Negatively Regulates Mouse Inflammatory Arthritis. J Immunol
891   2014;193:860–70. https://doi.org/10.4049/jimmunol.1400045.
892   [43] Monroe K, Weerakkody T, Sabelström H, Tatarakis D, Suh J, Chin M, et al.
893   PILRA regulates microglial neuroinflammation and lipid metabolism as a
894   candidate therapeutic target for Alzheimer's disease 2024.
895   https://doi.org/10.21203/rs.3.rs-3954863/v1.
896   [44] Wang H, Yung MMH, Ngan HYS, Chan KKL, Chan DW. The Impact of the
897   Tumor Microenvironment on Macrophage Polarization in Cancer Metastatic
898   Progression. Int J Mol Sci 2021;22:6560. https://doi.org/10.3390/ijms22126560.
899   [45] Chen S, Saeed AFUH, Liu Q, Jiang Q, Xu H, Xiao GG, et al. Macrophages in
900   immunoregulation and therapeutics. Signal Transduct Target Ther 2023;8:207.
901   https://doi.org/10.1038/s41392-023-01452-1.
902   [46] Sica A, Erreni M, Allavena P, Porta C. Macrophage polarization in pathology.
903   Cell Mol Life Sci 2015;72:4111–26. https://doi.org/10.1007/s00018-015-1995-y.
904   [47] Mantovani A, Marchesi F, Malesci A, Laghi L, Allavena P. Tumour-associated
905   macrophages as treatment targets in oncology. Nat Rev Clin Oncol
906   2017;14:399–416. https://doi.org/10.1038/nrclinonc.2016.217.
907   [48] Qian B-Z, Pollard JW. Macrophage Diversity Enhances Tumor Progression and
908   Metastasis. Cell 2010;141:39–51. https://doi.org/10.1016/j.cell.2010.03.014.

[49] Zhang Y, Guan X, Jiang P. Cytokine and Chemokine Signals of T-Cell Exclusion
in Tumors. Front Immunol 2020;11. https://doi.org/10.3389/fimmu.2020.594609.
[50] Kohli K, Pillarisetty VG, Kim TS. Key chemokines direct migration of immune
cells in solid tumors. Cancer Gene Ther 2022;29:10–21.
https://doi.org/10.1038/s41417-021-00303-x.
[51] Mehrotra P, Krishnamurthy P, Sun J, Goenka S, Kaplan MH. Poly-ADP-ribosyl
polymerase-14 promotes T helper 17 and follicular T helper development.
Immunology 2015;146:537–46. https://doi.org/10.1111/imm.12515.
[52] Mehrotra P, Riley JP, Patel R, Li F, Voss L, Goenka S. PARP-14 Functions as a
Transcriptional Switch for Stat6-dependent Gene Activation. J Biol Chem
2011;286:1767–76. https://doi.org/10.1074/jbc.M110.157768.
[53] Bennett TJ, Udupa VAV, Turner SJ. Running to Stand Still: Naive CD8+ T Cells
Actively Maintain a Program of Quiescence. Int J Mol Sci 2020;21:9773.
https://doi.org/10.3390/ijms21249773.
[54] Anderson KG, Stromnes IM, Greenberg PD. Obstacles Posed by the Tumor
Microenvironment to T cell Activity: A Case for Synergistic Therapies. Cancer
Cell 2017;31:311–25. https://doi.org/10.1016/j.ccell.2017.02.008.
[55] Yang L, Li A, Lei Q, Zhang Y. Tumor-intrinsic signaling pathways: key roles in
the regulation of the immunosuppressive tumor microenvironment. J Hematol
OncolJ Hematol Oncol 2019;12:125. https://doi.org/10.1186/s13045-019-0804-
8.
[56] Liu Z, Yu X, Xu L, Li Y, Zeng C. Current insight into the regulation of PD-L1 in
cancer. Exp Hematol Oncol 2022;11:44. https://doi.org/10.1186/s40164-022-
00297-8.
[57] Castro F, Cardoso AP, Gonçalves RM, Serre K, Oliveira MJ. Interferon-Gamma
at the Crossroads of Tumor Immune Surveillance or Evasion. Front Immunol
2018;9. https://doi.org/10.3389/fimmu.2018.00847.
936

Figure 1 – Identification of a genetic signature associated with PARP14
inhibition.
YUMM2.1 and 501MEL cells treated for 48h *in vitro* with either DMSO, 100nM PARP14
inhibitor (RBN012759), or 500nM PARP14 PROTAC (RBN012811); *n* = 3. **A.** Bulk RNAseq
analysis of a PARP14 inhibition signature identified across human and mouse cell lines.
Heatmap of treatment vs control per cell line, colours representing row-scaled z-score
expression high (red) to low (blue). **B.** Gene set enrichment analysis (GSEA) of 30 PARP14i
signature genes identified by comparison of bulk RNAseq from human and mouse cell lines,
chronically treated with IFN-γ followed by 48 hours of PARP14 inhibition (Metascape
analysis, values presented as -$\log_{10}$ of P value). **C.** Kaplan-Meier (KM) curves showing cancer
patients with high Tumour Immune Dysfunction and Exclusion (TIDE) scores of PARP14i
signature genes having better overall survival (OS) rates in representative studies. Red
indicates patients with high (top 50%) TIDE scores and blue indicates low (bottom 50%) TIDE
scores. **D.** Bulk RNAseq analysis on naïve YUMM2.1 tumours treated with IgG or α-PD-1. α-
PD-1-treated cohort was split into relapsed and responders. Compared to YUMM2.1
tumours pre-treated with chronic IFN-γ stimulation.
953
Figure 2 – Analysis of the cellular composition of YUMM2.1 tumours relapsing
following α-PD-1 antibodies with and without PARP14 inhibitor treatment. **A.**
YUMM2.1 cells were implanted subcutaneously into the flank of female C57BL/6 mice. Once

tumours reached ~80mm^3 in volume, treatment of α-PD-1 antibodies was commenced for a
total of two doses, 3 days apart. Tumour shrinkage and regrowth was monitored. When
tumours reached a volume between ~100-150mm^3, a second round of treatment
commenced, lasting 7 days. Tumours were subsequently processed for single-cell RNAseq.
Graphical summary of experimental timeline, showing tumour inoculation and
treatment regime (top), and illustration of tumour growth (bottom). **B.** Mutual nearest
neighbour (MNN) corrected Uniform Manifold Approximation and Projection (UMAP)
of 13 tumour samples (*n*=65,476 cells). Clustered using Leiden unsupervised
clustering and coloured by cell types. Cluster identifies below. **C.** Cell number
changes in each cluster by treatment: three mice IgG2a; three mice α-PD-1; three mice
PARP14i (RBN012759) alone; and four mice a combination of α-PD-1 + PARP14i
(RBN012759). **D.** Cell density projections by treatment.
969
970
Figure 4 - PARP14 inhibitor treatment in combination with α-PD-1 therapy
drives activated cytotoxic T cells but limits transition to dysfunctional states.
A. MNN corrected UMAP of T cell and natural killer (NK) cells (clusters 12-17,
n=18,298). Subclustered using Louvain method of community ordering and coloured
by subclusters. **B.** Cell number changes in each T and NK cell subcluster by
treatment: IgG2a; α-PD-1; PARP14i (RBN012759); combination treatment α-PD-1 + PARP14i
(RBN012759). **C.** Cell density projections of T and NK cells by treatment. **D.** T cell
subcluster identity- clusters 13-16. GSEA relevant terms of up- (red, FDR=0.05, top
25 genes) and downregulated (blue, FDR=0.05, bottom 25 genes) genes
(Metascape analysis, values presented as -log$_{10}$ of P value). **E.** Gene expression
presented as high (red) to low (blue) average expression and ratio of cells expressing (circle
size), across T cell subclusters 13-16. **F.** Slingshot trajectory inference analysis of CD8$^+$ T
cell subclusters 13-18 (*n=6,194*) with trajectory direction indicated by arrows (black).
G. KM curves showing cancer patients with high TIDE scores of genes from T cell
subclusters 13-16 and their effect on OS rates in bladder cancer cohort. Red
indicates patients with high (top 50%) TIDE scores and blue indicates low (bottom
50%) TIDE scores. **H.** Binarised heatmap of each T cell subcluster by relative
regulon activity (SCENIC analysis). **I - N.** Flow cytometry analysis of YUMM2.1
tumour xenografts in C57BL/6 mice relapsing on an initial dose of α-PD-1 and
harvested on-treatment following subsequent dosing with either α-PD-1 (Blue; *n* = 6)
or α-PD-1 + PARP14i (Red; *n* = 6). **I.** Percentage of total immune cells and individual
T cell subsets as a proportion of CD45$^+$ cells. **J.** Percentage of cytotoxic CD8$^+$ T cells
evaluated via   their expression of GZMB. Percentage of exhaustion subsets in
cytotoxic T cells measured by LAG-3 and TIM-3 levels for PD-1$^+$ **(K)**, PD-1$^+$CTLA-4$^+$
(L), or PD-1$^+$CTLA-4$^-$ **(M)** background. The p-value was assessed by two-sided
unpaired t-test and the data were presented as the mean□±□SEM. *p0.05,
***p□<□0.01, ***p□<□0.001, ****p□<□0.0001.
998
Figure 5 - PARP14 inhibitor treatment increases M1 macrophage polarisation
to generate a memory-like state. A. MNN corrected UMAP of monocytes and
macrophages (clusters 1-8, *n*=34,705). Subclustered using Louvain method of
community ordering and coloured by subclusters. **B.** Cell number changes in each
subcluster by treatment: IgG2a; α-PD-1; PARP14i (RBN012759); combination treatment α-
PD-1 + PARP14i (RBN012759). **C.** Cell density projections of both monocytes and
macrophages (colloquially referred to as MonoMacs) by treatment. **D.** Macrophage

subcluster identity- subclusters 2, 4, 8 and 10. GSEA of up- (FDR=0.05, logFC > 1,
marked in red) and downregulated (FDR=0.05, logFC < -1, marked in blue) genes
(Metascape analysis, values presented as -$\log_{10}$ of P value). **E.** Gene expression
presented as high (red) to low (blue) average expression and ratio of cells expressing (circle
size), across Macrophage subclusters. **F.** Binarised heatmap of 10% random subsampling
of each MonoMac subcluster by relative regulon activity (SCENIC analysis).
#
A. YUMM2.1 cells were implanted subcutaneously into the flank of female C57BL/6 mice.
Once tumours reached ~80mm^3 in volume, treatment of α-PD-1 antibodies was commenced
for a total of two doses, 3 days apart. Tumour shrinkage and regrowth was monitored. When
tumours reached a volume between ~100-150mm^3, a second round of treatment
commenced, lasting 7 days. Tumours were subsequently processed for single-cell RNAseq.
Graphical summary of experimental timeline, showing tumour inoculation and
treatment regime (top), and illustration of tumour growth (bottom). **B.** Mutual nearest
neighbour (MNN) corrected Uniform Manifold Approximation and Projection (UMAP)
of 13 tumour samples (n=65,476 cells). Clustered using Leiden unsupervised
clustering and coloured by cell types. Cluster identifies below. **C.** Cell number
changes in each cluster by treatment: three mice IgG2a; three mice α-PD-1; three mice
PARP14i (RBN012759) alone; and four mice a combination of α-PD-1 + PARP14i
(RBN012759). **D.** Cell density projections by treatment.
1026
Figure 3 - PARP14 inhibitor treatment in combination with α-PD-1 therapy
drives activated cytotoxic T cells but limits transition to dysfunctional states.
A. MNN corrected UMAP of T cell and natural killer (NK) cells (clusters 12-17,
$n=18,298$). Subclustered using Louvain method of community ordering and coloured
by subclusters. **B.** Cell number changes in each T and NK cell subcluster by
treatment: IgG2a; α-PD-1; PARP14i (RBN012759); combination treatment α-PD-1 + PARP14i
(RBN012759). **C.** Cell density projections of T and NK cells by treatment. **D.** T cell
subcluster identity- clusters 13-16. GSEA relevant terms of up- (red, FDR=0.05, top
25 genes) and downregulated (blue, FDR=0.05, bottom 25 genes) genes
(Metascape analysis, values presented as -$\log_{10}$ of P value). **E.** Gene expression
presented as high (red) to low (blue) average expression and ratio of cells expressing (circle
size), across T cell subclusters 13-16. **F.** Slingshot trajectory inference analysis of CD8$^+$ T
cell subclusters 13-18 ($n=6,194$) with trajectory direction indicated by arrows (black).
G. KM curves showing cancer patients with high TIDE scores of genes from T cell
subclusters 13-16 and their effect on OS rates in bladder cancer cohort. Red
indicates patients with high (top 50%) TIDE scores and blue indicates low (bottom
50%) TIDE scores. **H.** Binarised heatmap of each T cell subcluster by relative
regulon activity (SCENIC analysis). **I - N.** Flow cytometry analysis of YUMM2.1
tumour xenografts in C57BL/6 mice relapsing on an initial dose of α-PD-1 and
harvested on-treatment following subsequent dosing with either α-PD-1 (Blue; $n = 6$)
or α-PD-1 + PARP14i (Red; $n = 6$). **I.** Percentage of total immune cells and individual
T cell subsets as a proportion of CD45$^+$ cells. **J.** Percentage of cytotoxic CD8$^+$ T cells
evaluated via   their expression of GZMB. Percentage of exhaustion subsets in
cytotoxic T cells measured by LAG-3 and TIM-3 levels for PD-1$^+$ **(K)**, PD-1$^+$CTLA-4$^+$
(L), or PD-1$^+$CTLA-4$^-$ **(M)** background. The p-value was assessed by two-sided
unpaired t-test and the data were presented as the mean□±□SEM. *p < 0.05, ***p <
0.01, ***p < 0.001, ****p < 0.0001.
1054

Figure 4 - PARP14 inhibitor treatment increases M1 macrophage polarisation to generate a memory-like state.
A. MNN corrected UMAP of monocytes and macrophages (clusters 1-8, n=34,705). Subclustered using Louvain method of community ordering and coloured by subclusters. **B.** Cell number changes in each subcluster by treatment: IgG2a; α-PD-1; PARP14i (RBN012759); combination treatment α-PD-1 + PARP14i (RBN012759). **C.** Cell density projections of both monocytes and macrophages (colloquially referred to as MonoMacs) by treatment. **D.** Macrophage subcluster identity- subclusters 2, 4, 8 and 10. GSEA of up- (FDR=0.05, logFC > 1, marked in red) and downregulated (FDR=0.05, logFC < -1, marked in blue) genes (Metascape analysis, values presented as -$\log_{10}$ of P value). **E.** Gene expression presented as high (red) to low (blue) average expression and ratio of cells expressing (circle size), across Macrophage subclusters. **F.** Binarised heatmap of 10% random subsampling of each MonoMac subcluster by relative regulon activity (SCENIC analysis).

Figure 5 - Combined treatment of α-PD-1 with PARP14 inhibitor promotes dynamic adaptive resistance strategies in tumour cells which permit their persistence.
A. MNN corrected UMAP of tumour and cancer associated fibroblasts (CAFs) cells (clusters 26-28, n=3,445). Subclustered using Louvain method of community ordering and coloured by subclusters. **B.** Cell number changes in each subcluster by treatment: IgG2a; α-PD-1; PARP14i (RBN012759); combination treatment α-PD-1 + PARP14i (RBN012759). **C.** Cell density projections of tumour and CAF cells by treatment. **D.** Slingshot trajectory inference analysis of tumour cells with trajectory direction indicated by arrows (black). **E.** Tumour subcluster identity, GSEA relevant terms of up- (red, FDR=0.05, logFC > 1) and downregulated (blue, FDR=0.05, logFC < -1) genes in tumour subclusters 1-4 (Metascape analysis, values presented as -$\log_{10}$ of P value). **F.** PARP14i signature gene expression presented as high (red) to low (blue) average expression and ratio of cells expressing (circle size), across tumour and CAF subclusters in different treatments.

Supplementary Figure 1 – Identification of a genetic signature associated with PARP14 inhibition.
MC38 or A375 cells treated for 48h *in vitro* with either DMSO, 100nM PARP14 inhibitor (RBN012759), or 500nM PARP14 PROTAC (RBN012811); n = 3. Bulk RNAseq analysis of a PARP14 inhibition signature identified across human and mouse cell lines. Heatmap of treatment vs control per cell line, colours representing row-scaled z-score expression high (red) to low (blue).

Supplementary Figure 2 –
Clustergram of gene-TF associations derived from EnrichR analysis using the ARCHS4_TFs_Coexp database. Heatmap displays associations between genes (rows) and transcription factor terms (columns). Colour-coded to represent the -log10 transformed adjusted p-values of enrichment, darker colours indicating stronger statistical significance. No normalisation is applied, ensuring raw association strengths are visualised.

Supplementary Figure 3 -

A. YUMM2.1 cells were implanted subcutaneously into the flank of female C57BL/6 mice.
Once tumours reached ~80mm^3 in volume, treatment of α-PD-1 antibodies was commenced
for a total of two doses, 3 days apart (horizontal dotted line). Tumour shrinkage and
regrowth was monitored and individual growth curves plotted. When tumours reached a
volume between ~100-150mm^3, a second round of treatment commenced (day 0);
consisting of four IgG2a, α-PD-1, PARP14i or a combination of α-PD-1 + PARP14i. Treatment
lasted 7 days. Tumours were subsequently processed for single-cell RNAseq. **B.** RNAseq data
of excluded IgG2a control treated tumour sample (black triangles and high-density lines)
projected on the sample UMAP (colour) using projecTILs R package. **C.** Sample
representation in each cluster – histogram and **D.** UMAP. **E.** Cell number changes in each
cell type by treatment: three mice IgG2a; three mice α-PD-1; three mice PARP14i
(RBN012759) alone; and four mice a combination of α-PD-1 + PARP14i (RBN012759).
1115

Supplementary Figure 4 – -
A. Bulk RNAseq analysis on naïve YUMM2.1 tumours treated with IgG or α-PD-1. α-PD-1-
treated cohort was split into relapsed and responders. Compared to YUMM2.1 tumours pre-
treated with chronic IFN-γ stimulation. Heatmap representations of uniquely upregulated
genes per each subcluster within the top 200 expressed genes. Boxes represent relevant
clinical outcome per cluster. **B.** KM curves showing cancer patients with high TIDE
scores of uniquely upregulated genes per tumour subcluster and their effect on OS rates
in bladder cancer cohort. Red indicates patients with high (top 50%) TIDE scores
and blue indicates low (bottom 50%) TIDE scores.
1125

Supplementary Figure 5
A. Selected T cell marker gene expression presented as high (red) to low (blue) average
expression and ratio of cells expressing (circle size), across T cell subclusters. **B.** Bulk RNAseq
analysis on naïve YUMM2.1 tumours treated with IgG or α-PD-1. α-PD-1-treated cohort was
split into relapsed and responders. Compared to YUMM2.1 tumours pre-treated with
chronic IFN-γ stimulation. Heatmap representations of activation and exhaustion markers.
C. Representative image of Arid3a, Cebpb, Irf5, Mxd1 regulon activity AUC score in
MNN corrected T cell subcluster UMAP. Calculated by SCENIC. **D.** Binarised
heatmap of CD8$^+$ T cell subclusters by relative regulon activity (SCENIC analysis).
Supplementary Figure 6 – Flow cytometry gating strategy for T cell subsets
and exhaustion phenotypes
1137

1138
1139   **Supplementary Figure 5 – Flow cytometry gating strategy for T cell subsets**
1140   **and exhaustion phenotypes**
1141

Supplementary Figure 6 –
A. Bulk RNAseq analysis on naïve YUMM2.1 tumours treated with IgG or α-PD-1. α-PD-1-
treated cohort was split into relapsed and responders. Compared to YUMM2.1 tumours pre-
treated with chronic IFN-γ stimulation. Heatmap representations of uniquely upregulated
genes per each subcluster within the top 200 expressed genes. Boxes represent relevant
clinical outcome per cluster. **B.** KM curves showing cancer patients with high TIDE
scores of uniquely upregulated genes per tumour subcluster and their effect on OS rates

in bladder cancer cohort. Red indicates patients with high (top 50%) TIDE scores
and blue indicates low (bottom 50%) TIDE scores.

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

A MNN-UMAP 2 vs MNN-UMAP 1 plot with clusters 1–6

B Number of cells per treatment by Cluster (IgG, α-PD-1, PARP14i, α-PD-1 + PARP14i)

D UMAP2 vs UMAP1 plot with clusters 1–5

C Density Levels plots (IgG, α-PD-1, PARP14i, α-PD-1 + PARP14i)
- Highly invasive and adhesive
- CAFs
- Angiogenic EMT
- Stem-like
- Immunoregulatory
- Hyper-proliferative

E

Cluster 1 (−log$_{10}$P):
- MET activates PTK2 signaling
- Comprehensive IL 17A signaling
- regulation of lipid biosynthetic process
- skin development
- artery morphogenesis
- negative regulation of cell differentiation
- regulation of fat cell differentiation
- negative regulation of secretion by cell
- regulation of cell division
- Post-translational protein phosphorylation
- negative regulation of immune response
- regulation of angiogenesis
- negative regulation of cell cycle

Cluster 2 (−log$_{10}$P):
- regulation of epidermis development
- regulation of ATP-dependent activity
- negative regulation of mRNA catabolic process
- negative regulation of catalytic activity
- negative regulation of DNA-binding transcription factor activity
- organophosphate biosynthetic process

Cluster 3 (−log$_{10}$P):
- transmembrane receptor protein tyrosine kinase signaling...
- protein maturation
- chaperone cofactor-dependent protein refolding
- negative regulation of chemotaxis
- extracellular matrix organization
- negative regulation of cell differentiation
- Signaling by PDGF
- nuclear export
- regulation of epithelial cell migration
- Transport of inorganic cations/anions and amino...
- regulation of leukocyte chemotaxis

Cluster 4 (−log$_{10}$P):
- endocytosis
- intracellular protein transport
- cytoplasmic pattern recognition receptor signaling pathway
- negative regulation of angiogenesis
- semaphorin-plexin signaling pathway
- negative regulation of cell adhesion
- regulation of ERK1 and ERK2 cascade
- regulation of calcium ion import
- antigen processing and presentation of exogenous peptide...
- regulation of cellular response to growth factor stimulus
- regulation of smooth muscle cell migration
- regulation of cell growth
- regulation of lipid transport

■ Upregulated
■ Downregulated

F Dot plots (IgG, αPD1, PARP14i, α-PD-1 + PARP14i)

Genes: Gbp4, Batf2, Ifit3, Gbp5, Ciita, Psmb9, Cd274, Tap1, Apol6, Nlrc5, Gbp7, Gbp3, Gbp2, Stat1, Psmb8, Psmb10, Cd74, Serpina3f, A2m, Hapln3, Parp14, Irf1, Atf3

NumDetected: 0.00, 0.25, 0.50, 0.75, 1.00

z-score: 2, 1, 0, −1, −2