1 **Bioinformatic discovery of type 11 secretion system (T11SS) cargo across the Proteobacteria**
2

Authors: Alex S. Grossman[1,2], Nicholas C. Mucci[1], Heidi Goodrich-Blair[1,*]

[1]Department of Microbiology, University of Tennessee, Knoxville, Knoxville, TN 37996-0845

[2]Current address: The ADA Forsyth Institute, 245 First St, Cambridge, MA 02142

[*]To whom correspondence should be addressed: hgblair@utk.edu

Keywords: Co-occurrence; T11SS, Host-association; Cargo detection; Plasmin-sensitive protein;

DUF1194.

Graphical abstract:

Abstract:

<u>T</u>ype <u>11</u> <u>s</u>ecretion <u>s</u>ystems (T11SS) are broadly distributed among proteobacteria, with more

than 3000 T11SS family outer membrane proteins (OMPs) comprising 10 major sequence similarity

network (SSN) clusters. Of these, only 7, all from animal-associated cluster 1, have been

experimentally verified as secretins of cargo, including adhesins, hemophores, and metal binding

proteins. To identify novel cargo of a more diverse set of T11SS, we identified gene families co-

occurring in gene neighborhoods with either cluster 1 or marine microbe-associated cluster 3 T11SS

OMP genes. We developed bioinformatic controls to ensure perceived co-occurrences are specific

to T11SS, and not general to OMPs. We found that both cluster 1 and cluster 3 T11SS OMPs

frequently co-occur with single carbon metabolism and nucleotide synthesis pathways, but that only

cluster 1 T11SS OMPs had significant co-occurrence with metal and heme pathways, as well as with

mobile genetic islands, potentially indicating diversified function of this cluster. Cluster 1 T11SS co-

occurrences included 2556 predicted cargo proteins, unified by the presence of a C-terminal β-barrel

domain, which fall into 141 predicted UniRef50 clusters and approximately 10 different

architectures: 4 similar to known cargo and 6 uncharacterized types. We experimentally demonstrate

T11SS-dependent secretion of an uncharacterized cargo type with homology to <u>P</u>lasmin <u>s</u>ensitive

protein (Pls). Unexpectedly, genes encoding marine cluster 3 T11SS OMPs only rarely co-occurred

with the C-terminal β-barrel domain and instead frequently co-occurred with DUF1194-containing

genes. Overall, our results show that with sufficiently large-scale and controlled genomic data,

T11SS-dependent cargo proteins can be accurately predicted.

Impact Statement: We describe a novel method for controlling genomic neighborhood analyses

and use this controlled co-occurrence technique to examine two distinct clusters of genes from the

recently described type 11 secretion system (T11SS): Cluster 1, predominantly encoded by host-

associated microbes and Cluster 3, encoded by marine microbes. We found that both clusters of

predicted T11SS frequently co-occur with single carbon metabolism and nucleotide synthesis

pathways, but only the host-associated cluster co-occurred with iron uptake pathways. Using these

datasets, we predicted 2687 T11SS dependent cargo with approximately ten unique architectures, six

of which have not previously been linked to T11SS. Finally, we validate our results by demonstrating

T11SS-dependent secretion of a cargo protein with one of the novel architectures, plasmin-sensitive

surface protein Pls from *Haemophilus parahaemolyticus*.

Introduction

Identifying protein function remains a difficult task in the postgenomic age. As of the Pfam

2021 update v33.1, there were 4244 <u>d</u>omains of <u>u</u>nknown <u>f</u>unction (DUF) families, which

constitutes around 23% of the database (1). This means roughly one quarter of the Pfam database

remains uninvestigated, limiting our understanding of the breadth of biological processes occurring

on Earth. Functional characterization of proteins with DUF domains enables their annotation and

removal from the DUF list, improving our datasets permanently. *In silico* approaches applied to

protein domains are an important component of this process, as they can focus attention on the

likely functions of a particular protein family (2). Recently, a combination of *in silico* and

experimental approaches has yielded insight into the DUF560 protein family of outer membrane

proteins (OMPs), defining it as the type 11 bacterial secretion system (T11SS), represented by 10

distinct sequence-level clusters (3). T11SS homologs are present among a diverse array of

Proteobacteria, including human pathogens in the *Neisseria*, *Haemophilus*, and *Moraxella* genera (4–6).

In the gamma-proteobacterium *Xenorhabdus nematophila*, a T11SS system encoded within the

nematode intestinal localization locus (*nil*) is necessary for colonization of the mutualistic nematode

host *Steinernema carpocapsae* (7). In this system the T11SS, NilB, regulates the surface exposure of a

surface lipoprotein, NilC, both of which are necessary for colonization. Other known lipidated

T11SS-dependent cargo include co-receptors for metal-containing compounds heme, transferrin,

and lactoferrin (4–6,8,9). These cargo are composed of two types of domains, variable N-terminal

effector domains and 8-stranded C-terminal β-barrel domains (such as TbpBBD and Lipoprotein C).

T11SS can transport unlipidated, soluble cargo proteins across the outer membrane. For example,

T11SS-dependent hemophilin homologs have been characterized from *X. nematophila* (HrpC),

Acinetobacter baumannii (HphA), and *Haemophilus haemolyticus* (Hpl) (3,8,10,11). However, the majority

of T11SS OMPs do not have a known or predicted cognate cargo. A sequence similarity network

analysis suggested that experimentally characterized T11SS only capture a small portion of the

functional diversity present in this family (3).

The availability and accessibility of sequencing data have skyrocketed since the first bacterial

genome was sequenced in 1995, with over 200,000 bacterial and archaeal genomes currently

deposited on public databases (12). This tremendous repository of genomic sequence data allows

computational biologists to probe these genomes for patterns in genetic co-occurrence, as measured

through proximity to a gene of interest. Genes within a common functional pathway often cluster

together within a genome. Such patterns can be useful when forming hypotheses about the function

of uncharacterized proteins. Gene neighborhood studies analyze the genomic regions upstream and

downstream of any given gene of interest with the assumption that nearby genetic partners could

illuminate the role(s) of the unknown gene (13).

However, one flaw in the current methodology used to derive co-occurrence lists is the lack

of bioinformatic controls capable of removing non-specific co-occurring genes. For example,

commonly inherited, deeply ancestral domains, such as transcription and translation machinery,

often appear as the top result of a co-occurrence analysis, but do not yield useful information about

the specific query gene since their appearance is a coincidence of ubiquity. Also, meaningful results

can be obscured by co-occurrence based on structural or other common features present within the

query protein, such as those that target the gene product to specific cellular locations, which are

expected to interact commonly with the relevant protein trafficking machinery. Herein, we use

computational approaches to elucidate potential functional roles played by T11SS-dependent cargo

and predict novel cargo. Using these results, we then generated a co-expression system in *Escherichia*

coli to experimentally demonstrate the secretion of one novel, bioinformatically predicted cargo

family using the p̲lasmin s̲ensitive protein (Pls) from *Haemophilus parahaemolyticus* and the t̲ype e̲leven

P̲ls S̲ecretor (TepS).

Methods

Gene neighborhood analysis

The T11SS genome neighborhoods (±6 genes around a T11SS gene) from cluster 1 (1111

queries) and cluster 3 (145 queries) (3) were analyzed. T11SS genome neighborhoods were generated

through Rapid ORF Detection & Evaluation Online (RODEO) (14). As described in the main text,

a control was developed to identify and remove non-specific genes from among those flagged as co-

inherited or syntenic with T11SS (Fig. 2). The primary control list was generated by extracting all Go

term outer membrane proteins from proteobacteria within 1 amino acid of the median T11SS length

and removing all DUF560 containing proteins. For both T11SS cluster analyses a random subset of

control sequences were chosen equal to the number of queries being analyzed. Controlled protein

sequences were downloaded from UniProt (15) in February 2021. Co-occurring protein domains

were annotated and summed up to determine frequency among queries and controls. False

Discovery Rate was calculated by dividing frequency among T11SS queries and control sequences.

All domains with and FDR > 0.1 were excluded from analysis. For cluster 1 all domains with fewer

than 25 occurrences were excluded. For cluster 3 all domains with fewer than 5 co-occurrences were

excluded.

108   The Joint Genome Institute Integrated Microbial Genomes and Microbiomes platform

(https://img.jgi.doe.gov/) (16,17) was used to examine genomic regions surrounding the T11SS

OMPs-encoding genes of select cluster 3 bacteria. Briefly, the "find genes" function was used to

identify T11SS OMP homologs from selected representatives of the Roseobacteriaceae and

Rhodobacteraceae. The "gene neighborhoods" function was then used to visualize the regions

surrounding the T11SS OMP homologs and to identify syntenic loci (see Neighborhood Alignment

Tab of Supplemental File 3 for an example). Predicted structures were downloaded as .pdb files

from AlphaFold 2.0 (18,19) except in one instance (*Amylibacter cionae* CGMCC 1.15880

IEX10_RS12250 for which a Phyre 2.0 was used to predict the structure, and a .pdb file of the top ranked

prediction was used for Fig. 8G (18,20). This prediction was ranked with 98.51 confidence only over

16% identity over the C-terminal portion of the predicted protein, on the crystal structure of

lactoferrin binding protein b (LbpB) template 7jrd. Using each .pdb file, structural motifs were

highlighted using Protean 3D®, Version 17.4.3, DNASTAR, Madison, WI.

Functional and structural analyses of co-occurring gene neighborhood domains

123   Protein sequences that contain domains that co-occur with T11SS from the gene

neighborhood analysis were obtained via UniProt (15). KofamKOALA (21) and BlastKOALA (22)

were used to assess function. KofamKOALA assigns KEGG orthology to user sequence data by

HMMER/HMMSEARCH against the KEGG database. BlastKOALA is similar but uses a BLAST

search to assign KEGG orthology. Genes were matched to brite hierarchies to assess general cellular

function and KEGG pathway to assess potential interactions.

To identify potential cargo proteins all downloaded protein sequences were matched to

UniRef50 clusters to compress potential homologs (23). The reference sequences from these

UniRef50 clusters where then subjected to a BLAST search against known T11SS cargo β-barrel and

handle domains (nematode intestine localization protein (NilC) in *Xenorhabdus nematophila*, transferrin

binding protein (TbpB) N-terminal and C-terminal in *Xenorhabdus nematophila*, heme receptor protein

(HrpC) in *Xenorhabdus nematophila*, Cobalt receptor protein (CrpC) in *Xenorhabdus cabanillasii*,

Hemophilin (Hpl) in *Haemophilus haemolyticus*, hemophilin (HphA) in *Acinetobacter baumanii*, factor H

binding protein (fHbp) *Neisseria meningitidis*, and lactoferrin binding protein (LbpB) N-terminal and

C-terminal in *Neisseria meningitidis* to search for homologs of T11SS cargo with a low stringency E-

value filter of <0.05 in order to find even distantly related homologs, generating a list of 141

potential cargo proteins from cluster 1 and a single potential cargo from cluster 3. MSA alignment

was used to group the predicted cargo proteins into distinct architectures. 10 groups emerged,

TbpB-like, LbpB-like, Hemophilin-like 1 and 2, fHbp-like, Pls-like, highly disordered N-terminus 1

and 2, *Psychrobacter* surface protein, and *Sphingomonas* surface protein. Structural data predictions for

the 9 architectures which contained more than one sequence were generated through PSIPRED (24)

for secondary structure and AlphaFold2 (18,19) for tertiary structure.

Co-expression of Pls and its cognate T11SS protein TepS

All cultures were grown in glucose minimal media supplemented with 1% LB. Plate based

cultures were grown on glucose minimal plates (25). For plasmid-based expression, chemically

competent *Escherichia coli* strain BL21-DE3 (C43) were chosen for ease of transformation and their

ability to non-toxically express membrane proteins (26,27). Strains of *E coli* were grown at 37°C.

Where appropriate media was supplemented with ampicillin at a concentration of 150 µg/ml.

Protein expression was induced at the midlog point of bacterial growth via addition of isopropyl β-

d-1-thiogalactopyranoside (henceforth IPTG) at a concentration of 0.5mM.

Using sequences from *H. parahaemolyticus* strain NCTC10794, GenScript synthesized genes

for Plasmin sensitive surface protein (Pls-STO63610.1) with an attached C-terminal FLAG tag and

its genomically associated T11SS OMP (TepS-STO63613.1) with an N-terminal 6xHis tag. Both

genes were codon optimized for expression in *E. coli* Bl21DE3. GenScript cloned Pls-FLAG into

multiple cloning site 1 of pETDuet-1 and 6xHis-TepS into multiple cloning site 2, resulting in

expression plasmids pETDuet-1/Pls-FLAG and pETDuet-1/6xHis-TepS/Pls-FLAG. Expression

plasmids were transformed into *E. coli* via electroporation. Strains were grown in defined medium

with 150 µg/ml ampicillin (28). Bacteria were subcultured into 100 ml of broth at an initial optical

density at 600 nm (OD_{600}) of 0.1, grown for 6 h at 37°C to reach late logarithmic growth, and

induced with 500 µM isopropyl-β-d-thiogalactopyranoside (IPTG) for 1 hour. To monitor surface

exposure of Pls whole cells normalized to an OD600 of 1, rinsed 3x, and spotted in technical

triplicate on nitrocellulose membranes. The remaining cells were lysed open via 30 seconds of

sonication at ~500 root mean square volts (V_{rms}). Cell lysates were also spotted in technical triplicate

on nitrocellulose membranes to enumerate total cellular Pls. To monitor extracellular secretion of

Pls supernatant was collected from cultures after induction and filtered sterilized to remove all cells.

Supernatant samples were ultracentrifuged at 150,000 relative centrifugal force (RCF) for 3 h to

separate the soluble fraction from the insoluble component, composed of cell membrane

components and membrane vesicles. The membrane vesicle fraction was suspended in protein

sample buffer. 700 µl of each soluble supernatant fraction was precipitated via 10% trichloroacetic

acid (TCA) and the resulting pellet was suspended in protein sample buffer (29). Samples were

analyzed by 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and

immunoblotting. All dot blots and Western blots were probed with rat anti-FLAG primary antibody

and goat anti-rat secondary antibody conjugated to a 680nm fluorophore. Intensities were recorded

for FLAG reactive bands. For every supernatant sample, a band from the Coomassie blue-stained

gel was used as a loading control to normalize intensities of supernatant samples prior to analysis.

Fold change of secretion was determined by dividing the amount of supernatant Pls detected in the

Pls/TepS co-expression treatment by the amount detected in the Pls alone treatment. A Tukey's

honestly significant difference (HSD) test was used for comparing fold change of secretion (30).

Results:

Establishing a control for domain of unknown function (DUF) gene neighborhood co-

occurrence analysis

Characterized T11SS-cargo pairs encoded by Proteobacteria typically have conserved roles in

host-metal acquisition and immune evasion (3,6,8,10). Only a small portion of predicted T11SS-

cargo pairs has been experimentally characterized, leaving open the possibility that as-yet

uncharacterized systems may be participating in diverse symbiotic and metabolic pathways. To gain

insights into the potential breadth of biological processes in which T11SS-cargo pairs may function,

we collected all T11SS sequences identified as part of the largest T11SS sequence similarity network

cluster described in Grossman *et al.*, 2021 (cluster 1, containing 1111 genes) and utilized the Rapid

ORF Description & Evaluation Online (RODEO) software to identify all protein domains encoded

within six open reading frames upstream or downstream of the query genes, resulting in a data set of

1504 domains, co-occurring between 2 and 928 times (Cluster1Co tab of Supplemental file 1) (14).

To avoid spurious correlations, we chose to exclude any domains that co-occurred fewer than 25

times (~2.7% of the maximum frequency co-occurrence). This number was arrived at by generating

a histogram of co-occurrence frequency and identifying the inflection point between the rare co-

occurring sequences which made up the bulk of the dataset (1331 or 88.5%) and those domains

whose co-occurrence frequency was greater than expected randomly (173 or 11.5%).

To ensure that the observed co-occurrences were specific to the T11SS protein family and

not due to common features of genomic loci encoding outer membrane proteins, we developed and

implemented a control process. Briefly, RODEO co-occurrence neighborhoods were generated

using randomly selected proteins with biophysical characteristics similar to our DUF560 query

proteins such as protein length and outer membrane localization (GO term: outer membrane

GO0019867), and then used as a point of comparison for gene neighborhoods from T11SS genes.

The control sequences obtained were limited to the Proteobacteria phylum, where T11SS

predominantly occur (3). Due to a left-skewed distribution, the median length of all DUF560

domain containing proteins in Pfam (accessed February 2021) was used to reduce bias from

pseudogenes and gene fragments (Fig. S1) (1). The median size of all predicted T11SS proteins was

481 amino acids, so we extracted all outer membrane ORFs between 480 and 482 amino acids in

size. Sequences that appeared in the T11SS database were removed from the control database to

ensure unbiased selection. A random subsample of this control database was taken equal to the

number of sequences in the original query database (ControlQueries tab of Supplemental file 1) (Fig.

1).

OMP sequences from the subsampled control dataset were then submitted as a query to the

genome neighborhood network function of RODEO using the same parameters as the T11SS query

list. The output of this control co-occurrence analysis was used to generate a false discovery rate

(FDR) for any given domain by dividing frequency of co-occurrence with random OMPs by

frequency of co-occurrences with T11SS genes. We then gated the dataset for domains with an FDR

less than 0.1 (10%) to exclude non-specific correlations, resulting in a final set of 51 significantly co-

occurring domains. (Cluster1FDR_Threshold tab of Supplemental file 1). The most common

significant co-occurring domains alongside cluster 1 T11SS OMPs were other DUF560 proteins

(928/1111), the TbpBBD β barrel domain (503/1111), the TonB C-terminal domains (364/1111),

heme oxygenase (137/1111), and DUF454 (118/1111). Next, all genes that co-occurred with a

cluster 1 T11SS gene and possessed one of the identified domains were extracted resulting in 3405

significantly co-occurring genes (Cluster1SigGenes tab of Supplemental file 1). These genes were

submitted to KofamKOALA and BlastKOALA in parallel for annotation into molecular pathways

and brite hierarchies (21,22).

Functional assessment of T11SS co-occurring proteins identifies a role of iron uptake,

protein export, and single carbon metabolism

We examined the co-occurrence gene set for those with annotations in the curated Kyoto

Encyclopedia of Genes and Genomes (KEGG) ortholog (KO) database (21,31). A hidden Markov

model search using KofamKOALA (HMMER/HMMSEARCH) was used to annotate the genes

according to the KO database. The software found matches for 1488 of the 3405 significant co-

occurring genes (Cluster1KofamBrite tab of Supplemental file 2). When matched to cellular

functions (brite hierarchies) the most common categories were transporters (periplasmic TonB,

ExbD, Signal peptidase II, TolA, etc.), enzymes (heme oxygenase, formate dehydrogenase,

exopolyphosphatase, dihydrofolate reductase, etc.), and tRNA biogenesis (tRNA modifying GTPase,

tryptophanyl-tRNA synthase, Ribonuclease P, aminoacyl tRNA synthase) (Fig. 2A). Additionally,

cluster 1 T11SS genes were found to significantly co-occur with transposase genes. Of the 1488

matched sequences, 259 had known pathway association, most commonly biosynthesis of secondary

metabolites (heme oxygenase, cobalamin-dependent methionine synthase, vitamin K biosynthesis

protein MenH, etc.), porphyrin metabolism (heme oxygenase), and protein export (signal peptidase

II and YidC insertase) (Fig. 2B) (Cluster1KofamPath tab of Supplemental file 2). The list of

pathways also included ribosome components (RpmB, RpmG, RpmH) and aminoacyl tRNA

biosynthesis (TyrS, TrpS). A high frequency of co-occurrence with transposases, ribosome

components, and tRNA synthesis genes is characteristic of pathogenicity islands, phage regions, and

other mobile genetic elements, since many mobile genetic elements use universally conserved

ribosome and tRNA genes as anchors for genomic insertion (32). The frequency at which T11SS

genes are encoded within mobile genetic element islands suggests that T11SS may frequently be

horizontally acquired. Finally, the list included several carbohydrate biosynthesis pathways

(hsa00670, one carbon pool by folate; map00230, purine metabolism).

To further functionally annotate co-occurrences, we used BLAST alignment, rather than

hidden Markov searching, to establish sequence similarity using BlastKOALA, which compares

query sequences to the non-redundant KEGG dataset (22). The software found matches for 1387 of

the 3405 significantly co-occurring genes. Categorization into cellular functions (brite hierarchies)

again revealed the largest categories to be enzymes, transporters, and tRNA biogenesis (Fig. S2A)

(Cluster1BlastBrite tab of Supplemental file 2). BlastKOALA identified more co-occurring amino

acid importers (arginine, lysine, tyrosine, tryptophan, and serine transporters) in the dataset than

KofamKOALA. Of the 1387 identified genes, 260 had known pathways. This analysis identified

similar pathways to those identified by KofamKOALA, including secondary metabolite biosynthesis,

protein export, porphyrin metabolism, and carbohydrate biosynthesis (Fig. S2B) (Cluster1BlastPath

tab of Supplemental file 2).

Predicted structure analysis of putative cluster 1 T11SS cargo reveals variable N-termini and shared C-terminal domains

To bioinformatically identify potential cluster 1 T11SS-dependent cargo we searched co-occurring proteins for homology to domains of known cargo. The 3405 significantly cluster 1 co-occurring sequences were first consolidated using UniProt ID mapper into 666 UniRef50 clusters with 50% or greater identity at an amino acid level (Cluster1Uniref tab of Supplemental file 1) (23). The reference sequences of these UniRef50 were next searched using BLASTp for sequence similarity with N-terminal domains and C-terminal β-barrel domains from experimentally verified T11SS cargo proteins: NilC, TbpB, LbpB, HrpC, CrpC, Hpl, and Factor H binding protein, with a low stringency E-value cutoff of 0.05 (3,5,6). No matches were found for NilC or the N-terminal ligand binding domain of fHbp, but all other queries matched at least one cluster. This analysis revealed 141 clusters of potential T11SS-dependent cargo proteins representing 2656 ORFs across 1048 species/taxa (Cluster1PutativeCargo tab of Supplemental file 1). Each sequence was also annotated with SignalP 6 (33,34) which predicted 81 Sec secreted lipoproteins, 33 Sec secreted soluble proteins, and 27 proteins with no predicted signal peptide. Several of the proteins with no signal peptide were annotated as fragments or pseudogenes, supporting the previous observations of T11SS transporting Sec dependent lipoproteins and soluble proteins.

We utilized multiple sequence alignment to separate the predicted cargo protein clusters into distinct, annotated architectures (Cluster1PutativeCargo tab of Supplemental file 1). This analysis identified two distinctive groups of Hemophilin-like proteins based on homology to known hemophilins (Hpl, HrpC, HsmA), the slightly larger and predominantly soluble group 1 (96 lipoproteins, 912 soluble) and the predominantly lipidated group 2 (65 lipoproteins, 7 soluble), a group of TbpB-like proteins (638 lipoproteins, 7 soluble), a group of proteins annotated as fHbp-

like proteins by virtue of their Lipoprotein C domain (54 lipoproteins, 285 soluble), and a group of

lactoferrin-binding protein-like proteins (105 lipoproteins). In addition, lipoproteins with an

elongated, disordered N-terminal region occurred as two groups: disordered N-terminus 1 (196

lipoproteins, 8 soluble) and 2 (1 lipoprotein only). Another group had an elongated N-terminal

domain with repeat rich (EEARKA motif) α-helical regions homologous to the Pls surface

glycoprotein in *Staphylococcus aureus*, termed Pls-like proteins (173 lipoproteins, 20 soluble). Two

phylogenetically restricted architectures were observed with multiple TbpBBD domains,

Sphingomonas surface proteins, which have an N-terminal sequence of proline-threonine repeats (18

lipoproteins, 12 soluble) and *Psychrobacter* surface proteins which have an N-terminal sequence of

~12 proline-threonine-aspartic acid repeats (PTD motif) (55 lipoproteins, 4 soluble). Proline-

threonine repeats occur in *Xanthomonas campestris* endoglucanase, where they provide elasticity and

facilitate enzyme positioning around its substrate (35). The PTD motif is found in a family of

Gracilibacteria (BD1-5) surface proteins and in the eukaryotic microbe *Plasmodium falciparum* and are

hypothesized to act as adhesins facilitating interactions with hosts (36,37).

Using PsiPred 4.0 (24) and AlphaFold2 (18) we predicted the secondary and tertiary

structures of representative proteins from the largest UniRef50 cluster from each architecture with

more than one representative (Fig. 3 and Fig. 4). All the architectures were predicted to have at least

one domain, generally at the C-terminus, composed of β-strands linked by disordered loops that

form a C-terminal β-barrel domain similar to the TbpBBD domain or the Lipoprotein C domain.

The N-terminal domains of most cargo had β-strands predicted to form handle domains like those

seen in hemophilin or TbpB (3,5). When present, the disordered regions and long α-helical repeat

regions preceded this handle domain.

When examining secondary and tertiary structures in parallel we noticed that the predicted

cargo form three distinctive structural regions: a common C-terminal 8-stranded β-barrel domain, a

common central region formed by β-strands, which we have termed a "pedestal" region based on its

positioning as a platform for a variable N-terminal domain "effector" domain, predicted to serve the

functional role specific to each cargo protein type. Using solved crystal structures where available

(4,8,38), and Alphafold2 or Phyre 2.0 models for representative cargo proteins (18,20), we divided

the predicted cargo structures into these three regions and categorized them by recurrent patterns

(Fig. 5). The predicted central pedestal-domains fell into six visibly distinguishable categories; 4-

stranded groove, wing, and sheet, 5-stranded sheet, and 6-stranded/extended sheets (Fig. 5). None

of the examined pedestal regions featured complex, variable loops, suggesting they may play a

structural role (Fig. 5). In contrast, the C-terminal 8-stranded β-barrel core structures are

embellished within each protein by diverse inter-strand loops including disordered regions, extended

β-strands, and α-helices that could be playing functional roles, as do the loops of the N-lobe

TbpBBD domain of TbpB (39).

The N-terminal effector domains displayed a variety of structural motifs including

disordered loops, α-helices, β-strands, and β-barrels (Fig. 5). The examined effector regions ranged

in size, with some barely extending beyond the pedestal domain and others including duplications of

the 8-stranded β-barrel domain. The latter is typified by the solved structure of TbpB, which

comprises two lobes, each containing the hemophilin handle-β-barrel pair architecture. The TbpB

N-lobe pair comprises the effector region, while the C-lobe handle comprises the pedestal and the

C-lobe barrel is the putative T11SS targeting domain (38,39). This domain duplication and

functional specialization may also explain the architecture of *Sphingomonas* and *Psychrobacter* surface

proteins that feature 2-3 copies of the handle-barrel pair. However, the long, disordered loops

present in those proteins likely generate a distinctive structural architecture relative to TbpB.

Demonstrating T11SS-dependent secretion of Plasmin sensitive surface protein

To assess the accuracy of our bioinformatic predictions of T11SS cargo, we experimentally

tested levels of secretion in *E. coli* of a representative novel cargo protein with or without its

predicted cognate T11SS protein. Several of our predicted cargo possess a large α-helix repeat region

absent among known T11SS cargo. We chose to test a representative of this group, the 525 amino

acid P̲lasmin s̲ensitive p̲rotein homolog (Pls) (STO63610.1) and its genomically associated putative

t̲ype e̲leven P̲ls s̲ecretor (TepS) (STO63613.1) from *H. parahaemolyticus* strain NCTC10794. The *H.*

parahaemolyticus Pls α-helix repeat region shares between 49 and 62% identity with the repeat regions

of surface glycoprotein Pls from *Staphylococcus aureus* (WP_256928426, WP_256927786,

WP_258808513, WP_256933640, and WP_257570445) that contributes to virulence by stimulating

biofilm formation (40). *Staphylococcus* and *H. parahaemolyticus* Pls differ in that homologs of the former

are larger and lack the C-terminal β-barrel domain.

To experimentally test for TepS-mediated secretion of *H. parahaemolyticus* Pls, we used

addition of a FLAG-tag epitope to allow Pls detection and expressed this protein with or without

TepS using pETDuet-1 based plasmids in *E. coli* BL21 DE3 C43. After protein expression

induction, anti-FLAG immunodotblots were performed on intact (to measure surface exposed

protein) and lysed cells (to measure total protein) to compare among treatments. Co-expression with

TepS did not significantly impact surface exposure of Pls or the intracellular concentration of Pls

(Fig. 6 and S3). This result was initially surprising, since Pls was predicted to be a surface anchored

lipoprotein. However, since *S. aureus* Pls can be cleaved from the cell surface spontaneously and by

proteinases such as plasmin (40,41) we next monitored free and extracellular-vesicle fractions of cell-

free supernatants for the presence of Pls via SDS-PAGE. These data revealed that Pls was present in

the supernatant, but only when co-expressed with the T11SS protein TepS. Some Pls was located in

extracellular vesicles independent of TepS expression, but this amount was higher in the presence of

TepS (Fig. 6 and Fig. S3). In the immunoblot, FLAG-tagged Pls is visible as a single clear band in

individual replicates, but varied in apparent size between biological replicates from 82.3 to 121.3 kD

with an average of 103kD. This is almost twice the value predicted from sequence alone (56.8kD).

This large and variable size may suggest that Pls is either being significantly slowed by glycan

moieties as it travels through polyacrylamide, or that it exists as a dimer when expressed in *E. coli*.

Bioinformatic analysis of cluster 3 T11SS proteins from marine environments reveals distinct

co-occurrence patterns

To test the robustness of our analysis we then utilized our co-occurrence technique to

explore cluster 3 T11SS proteins, predominantly containing sequences from marine microbes. This

cluster almost exclusively contains sequences from the families *Roseobacteriaceae* and *Rhodobacteraceae*

(42), including both pelagic microbes and symbionts of algae (*Silicimonas algicola*), mollusks

(*Aliiroseovarius crassostreae*), tunicates (*Ascidiaceihabitans donghaensis*), echinoderms (*Sulfitobacter delicatus*),

and corals (*Roseivivax isoporae*) (43–47). All genes from cluster 3 (145 sequences) were used as queries

in the RODEO genome neighborhood network function, using the same parameters as the cluster 1

analysis, resulting in 203 co-occurring domains with frequencies between 2 and 102 (Cluster3Co tab

of Supplemental file 1). Domains which co-occurred fewer than 5 times were excluded from analysis

(122 or ~60.0%). Co-occurring domains were compared to another random subset of the non-

specific control co-occurrence dataset to assign FDR values. All domains with an FDR greater than

0.1 were excluded from analysis, resulting in 42 significantly co-occurring domains

(Cluster3FDR_Threshold tab of Supplemental file 1). The most common co-occurring domains

were additional DUF560/T11SS domains (102/145 loci), DUF1194 (102/145 loci), glyoxalase

domains (89/145 loci), and thymidylate synthase/thymidylate synthase complementing protein

(77/145 loci). The TbpBBD domain passed the FDR threshold, however relative to the analysis of

cluster 1 it was exceptionally rare (8/145 loci). All genes that co-occurred with a marine cluster 3

T11SS OMP and contained any of the significant co-occurring domains were extracted, resulting in

860 significant co-occurring genes which we submitted to KofamKOALA and BlastKOALA for

functional analysis (21,22).

KofamKOALA found matches for 595 of the 860 significant co-occurring genes

(Cluster3KofamBrite tab of Supplemental file 2). When matched to cellular functions (brite

hierarchies) the most common categories were enzymes (lactoylglutathione lyase, thymidylate

synthase complementing proteins, aspartate aminotransferase, etc.), prokaryotic defense systems

(antitoxins CptB and HigA-1, topoisomerase IV B), and transcription factors (Cell division repressor

DicA, cold shock protein) (Fig. 7A). Of the 595 matched sequences, 270 had known pathway

association, the most common pathways being biosynthesis of one carbon pool by folate

(dihydrofolate reductase, thymidylate synthase), pyrimidine/nucleotide metabolism (thymidylate

synthase), pyruvate metabolism (lactoylglutathione lyase), and biosynthesis of amino acids (aspartate

aminotransferase) (Cluster3KofamPath tab of Supplemental file 2). Unlike cluster 1, transposases

were not among the significantly co-occurring genes of cluster 3. However, aminoacyl-tRNA

synthesis functions, which can be associated themselves with mobile genetic elements (32,48), do

appear on the co-occurrence list, including alanyl-tRNA, prolyl-tRNA, and threonyl-tRNA synthases

(Fig. 7B). Parallel analysis with BlastKOALA identified 514 of the 860 significant co-occurring

genes, of which 267 had known pathway association (Cluster3BlastBrite and Path tabs of

Supplemental file 2). BlastKOALA did not identify any additional cellular functions or pathways not

already revealed by KofamKOALA (Fig. S4).

Bioinformatic investigation of potential cluster 3 T11SS cargo

Initial attempts to identify new cargo proteins within the co-occurrence datasets from cluster

3 used the same methods used for cluster 1. UniRef50 clustering with ≥50% identity resulted in 218

homologous groups (Cluster3Uniref tab of Supplemental file 1) that were assessed with BLASTp for

sequence level homology to experimentally characterized T11SS cargo domains. Only a few sets of

homologs were identified, even when including all TbpBBD-domain-containing UniRef50 clusters.

Homologs of six UniRef50 clusters, specific to the *Rhodobacteraceae/Roseobacteriaceae* families,

resembled the Hemophilin-like 1 architecture, and homologs from another UniRef50 resembled the

disordered N-terminus 1 architecture noted previously (see Figs. 3 and 4). Collectively these 7

UniRef50 clusters encompass a total of only 31 ORFs across 27 species/taxa (Cluster3PutativeCargo

tab of Supplemental file 1). The ratio of predicted cargo per T11SS OMP was far lower in cluster 3

(0.21 Cargo/input OMP) than it was in cluster 1 (2.39 Cargo/input OMP). This dearth of co-

occurring putative cargo proteins suggests that marine cluster 3 T11SS OMPs seldom co-occur with

their cognate cargo proteins, utilize cargo proteins with below-threshold sequence similarity to those

of cluster 1 T11SS, or have evolved distinct functions and no longer have partner cargo proteins.

To further examine this potential transition in function, we comparatively annotated a

syntenic T11SS OMP-encoding locus in 11 representative *Rhodobacteraceae/Roseobacteriaceae* strains

(Clus3SyntenicPairs tab of Supplemental file 3). Each locus encodes all four of the top co-occurring

domains from our co-occurrence analysis, including flavin-dependent thymidylate synthase,

lactoylglutathione lyase, and one or more DUF1194 (NeighborhoodAlignment tab of Supplemental

file 3 and Fig. 8A). All but one (*Pseudooceanicola marinus* LMG 23705) of the loci also encodes a protein

that is predicted to be a T11SS cargo lipoprotein (282-375 aa) based on the presence of an N-

terminal SPII type signal sequence and a C-terminal 8-stranded β-barrel domain (Fig. 5F; Fig. 8B-K)

(18,20). The presence of both cluster 1 and cluster 3 co-occurring domains indicates that this

conserved locus is a hybrid of the two. It is tempting to speculate that this locus represents a

transitional state in the divergence of these systems, with at least one analyzed genome appearing to

have lost the cluster 1 co-occurring, 8-stranded β-barrel domain.

We next examined the cluster 3 marine T11SS co-occurrence dataset to consider the

possibility of alternative putative cargo types. In contrast to cluster 1 T11SS, co-occurrence with

TbpBBD/lipoprotein C domains, TonB-dependent receptors, and TonB domains are rare within

the marine cluster 3 T11SS loci. Instead, the DUF1194 domain (PF06707) was highly prevalent

(102/145 queries) (Supplemental file 4). Of the 102 DUF1194-containing sequences found in the

cluster 3 co-occurrence list, 7 are predicted to be lipoproteins, 76 are predicted to be soluble Sec-

secreted proteins, and 19 have no detectable signal peptide according to SignalP 6 (33,34). Many

cluster 3 T11SS loci encoded 2 or 3 distinct DUF1194 genes in close proximity (e.g., Fig. 8),

predicted to encode both lipidated and non-lipidated proteins (e.g., algal symbiont *Rhodobacteraceae*

PD-2 ETA49263.2 and ETA49262.2, respectively) (Supplemental file 4). The predicted

RoseTTAFold structure (49) of the DUF1194 domain is a mixed β-strand/α-helix. According to the

Pfam database, the DUF1194 domain occurs in 31 distinct protein architectures with other domains,

including C-terminal autotransporter domain (PF03797) predicted to function in T5SS secretion, the

CARDB domain (PF07705), which adopts a 7 β-strand structure, DUF11 (PF01345), predicted to

have an 8-stranded β-barrel structure, and PF18911, the known crystal structures of which also

adopt a β-barrel (e.g., 1wgo and 2y72).

Of the co-occurring DUF1194 sequences, 6 are annotated as having homology to von

Willebrand factor type A domain or vWA (PF00092). This homology is reflected in the structural

similarity between these two domains, including the presence of a Rossman fold, which may hint at

similar molecular roles. vWA protein architectures can include divalent-cation-binding metal ion-

dependent adhesion sites (MIDAS), with a characteristic DxSxS…T…D motif (50,51). Almost all

the DUF1194 proteins that co-occurred with DUF560 proteins contained at least one MIDAS motif

(98/102). To examine potential structural features of cluster 3 T11SS co-occurring DUF1194-

containing homologs, we focused on three adjacently encoded DUF1194 genes from

Ascidiaceihabitans donghaensis (SPH20589/SPH20588/SPH20587) for structural analysis. PsiPred 4.0

predicted that these proteins have at least 5 predominantly hydrophobic β-strands, all separated by

α-helix regions in a structure that also appears to be a variation on the classical α/β Rossman fold

(Fig. 9A). RoseTTAFold predictions of the tertiary structure of these DUF1194 proteins reveal

globular proteins wherein 6 hydrophobic β-strands form a β-sheet at the core of the protein, which

is protected by three α-helices per side. All three representatives had a MIDAS motif starting 10-11

residues after the predicted signal peptide. For each protein, the last 11-13 C-terminal residues

appeared to be disordered, and their positions could not be estimated with a high degree of

confidence (Fig. 9B).

Discussion

The controlled bioinformatics analyses we report here reinforce the idea that cluster 1

T11SS-dependent cargo have roles in metal uptake, single carbon metabolism, and nutrient

provision. Further, our data extend the experimentally verified T11SS cargo to include *Haemophilus*

parahaemolyticus Pls and lend credence to the concept that T11SS cargo can be bioinformatically

predicted. The structural prediction that this cargo protein is a fusion between a functional Pls

domain, typical of Gram-positive bacteria, and a C-terminal 8-stranded β-barrel domain lends

further evidence of the function of the barrel domain in directing cargo to the T11SS, and for the

function of T11SS in facilitating transport across an outer membrane. However, our analysis of the

more distantly related cluster 3 T11SS OMPs revealed that known barrel domains (TbpBBD and

lipoprotein C) are rare among the significantly co-occurring genes identified, raising the intriguing

possibility that novel structural domains may exist that mediate targeting to cluster 3 T11SS.

Our bioinformatic approach relied upon gene co-occurrence analysis, which uses genomic

proximity as a proxy for coordinate regulation and potential interactions, since genes within a

common functional pathway often cluster together within a genome. While this is not a foolproof

method, the use of large genomic datasets increases the power of co-occurrence analysis to identify

potential co-interacting partners. One of the methodological refinements we developed was the use

of *in silico* controls for genome neighborhood co-occurrence to reduce false positives and noise in

the output datasets of putative T11SS co-occurring protein families. For instance, activities

associated with protein translation and RNA metabolism, which are essential cellular processes

expected to be present in every cell, are not always useful co-occurrence signatures for elucidating

T11SS function and components. Such general functions are not indicative of the specific functions

of T11SS relative to any other OMP and can be considered a form of false negative, Type II error;

such co-occurrences are not biologically relevant, but are not being excluded from the analysis

because they do co-occur frequently with the query gene. However, removal of such co-occurrences

can also lead to false positive, Type I error, because they can also result in removal of some

biologically relevant co-occurrences. For example, there could be protein families that are important

functional partners of the query protein, but that are removed from the dataset if they were

important for both query and control proteins. This type of error occurred in our analysis. For

example, TonB-dependent receptors, with which many T11SS-dependent cargo interact, were

excluded from the cluster 1 T11SS analyses since these domains associate with other outer

membrane proteins unrelated to T11SS. In a sense, this control narrows the focus from total co-

occurrence to unique co-occurrence, and as such it is best suited towards applications in which the

researcher seeks to differentiate protein families. Additionally, the technique can be further adapted

and refined, depending on application. For instance, multiple non-specific gene neighborhood

controls could be generated by randomly subsampling the list of proteins that are biophysically

similar to the query proteins, allowing for increased stringency and measures of variance.

The cluster 1 T11SS significantly co-occurring genes identified here are consistent with

previous observations (3,5). Iron uptake from the environment is a common function among

experimentally verified T11SS-dependent cargo proteins and this theme was extended within the

cluster 1 T11SS co-occurrence dataset, which included heme oxygenases and heme transporters that

facilitate iron uptake (52). Similarly, the protein export genes identified as co-occurring with cluster 1

T11SS included TonB and ExbD, which are essential to energize uptake of nutrients from known

T11SS-dependent cargo (6,8,53). TolA is essential to bring periplasmic nutrients into the cytoplasm

of the cell (54). Signal peptidase II is essential for the generation of bacterial lipoproteins, which

constitute a substantial percentage of T11SS-dependent cargo (55). Finally, the observed co-

occurrence with transposases, ribosome components, and tRNA-synthases is consistent with the

hypothesis that T11SS are regularly associated with mobile genetic islands and can be horizontally

acquired and contribute to fitness within a host environment (32).

Significantly co-occurring pathways were commonly identified for both cluster 1 and cluster

3 T11SS, and represent pathways not previously associated with T11SS. These included folate

(vitamin B9)-related pathways, including folate biosynthesis, folate-dependent enzymes, and one

carbon metabolism via folate. Folate, like heme, is an enzyme cofactor required for many diverse

biological pathways. It is essential for the function of thymidylate synthase complementing protein,

also known as flavin-dependent thymidylate synthase, in nucleotide biosynthesis (56). This protein is

the third most common cluster 3 marine T11SS co-occurring gene, suggesting coordinate regulation

or inheritance. Since, unlike heme, folate does not incorporate a metallic ion and most bacteria

synthesize their folate instead of scavenging it from the environment (57), we do not favor the

hypothesis that folate itself is acquired through a T11SS-dependent uptake mechanism. Instead, the

close association of folate metabolism and T11SS may reflect the fact that folate biosynthesis, as well

as iron-sulfur cluster biosynthesis, and methylotrophic metabolism by formate dehydrogenase

(which also significantly co-occurred with T11SS) depend on the metal-bearing enzyme cofactors,

cobalamin (vitamin B12) and molybdopterin cofactor (MoCo) (58–61). Our data suggest that some

T11SS OMPs may function in cobalamin or molybdopterin uptake pathways. Regardless, our data

clearly demonstrate an association between T11SS and single carbon metabolism and methyltropism.

Cluster 1 T11SS co-occurrences featured TonB energization or heme/iron uptake that were lacking

from cluster 3 T11SS co-occurrences. In turn, the most frequent cluster 3 T11SS co-occurrences were

DUF1194 and lactoylglutathione lyase (GloA), neither of which appeared in the Cluster 1 T11SS co-

occurrence datasets, but both of which have connections to metal ions. The function of DUF1194 is

currently unknown but it is predicted to fold into a globular protein with an α/β Rossman fold.

Several of the DUF1194 proteins (6/103) we detected also had homology to a structurally similar

α/β Rossman fold domain called von Willebrand factor A (vWA), which is most often associated

with serum glycoproteins and large multimeric protein complexes in mammalian systems (62,63). In

Myxococcus xanthus bacteria, the vWA domain-containing surface lipoprotein CglB acts as a surface

adhesin essential for gliding motility (64). Upon binding a ligand, vWA domain-containing adhesins

such as CglB can stabilize their bond via a conformational change induced by MIDAS motif

allosteric binding of a divalent metal cation, often Mg^{2+}(51,64). The presence of a MIDAS motif in

the vast majority of DUF1194 identified within our dataset suggests that these proteins may also

play a role in adhesion or may participate in metal homeostasis in previously unpredicted ways. An

exciting future direction will be to determine if DUF1194 proteins are a type of T11SS-dependent

cargo, or if they contribute to a multimeric protein complex that interacts with T11SS proteins.

Lactoylglutathione lyase is an enzyme with a coordinated metal ion (typically nickel) and functions in

the regeneration step of the glyoxalase-mediated aldehyde detoxification system (65). It is possible

that the nickel requirement of this enzyme reflects a role for cluster 3 T11SS in facilitating nickel

uptake. Alternatively, this co-occurrence may indicate that cluster 3 T11SS are more generally

participating in an aldehyde producing process. Notably, aldehydes like methylglyoxal are

exceptionally cytotoxic and can damage DNA and proteins, so the cluster 3 T11SS co-occurrence of

nucleotide and amino acid biosynthesis may reflect a more general relationship between T11SS and

the aldehyde repair response (65,66).

In addition to providing insights into functional pathways associated with T11SS, our

analysis also allowed us to identify 148 homology groups of putative T11SS-dependent cargo,

including those with domain architectures distinct from those found in known cargo. We verified

that at least one of these, the plasmin-sensitive surface protein (Pls) from the Gram-negative

organism *H. parahaemolyticus,* is a bona fide T11SS-dependent cargo protein, demonstrating the

efficacy of our bioinformatics approach. Pls represents a novel N-terminal effector domain among

T11SS cargo. This domain is homologous to glycosylated Pls proteins found on or spontaneously

cleaved from the surfaces of Gram-positive organisms such as *Staphylococcus aureus* and *S. epidermidis*

where function in biofilm formation and modulating adhesion to host proteins (40,41). Given that

the function of Pls requires surface exposure, its presence in a Gram-negative organism is

complicated by the existence of the outer membrane. In *H. parahaemolyticus*, this hurdle seems to

have been overcome via fusion of the Pls repeat domain to the C-terminal 8-stranded β-barrel

domain necessary for cluster 1 T11SS-dependent secretion across the outer membrane (3,5).

When co-expressed in *E. coli* with its T11SS partner, TepS, Pls predominantly localizes to the

supernatant and is not tethered to the cell surface as would be expected of a lipoprotein. Our

working model is that some Pls can reach the cell surface in the absence of T11SS, as has been

observed for factor H binding protein (67,68), and that in the presence of the T11SS TepS, Pls is

secreted but then cleaved, releasing it into the extracellular milieu Given that our experiments were

conducted in a protease-deficient strain of *E. coli*, it will be exciting in future studies to determine if

TepS alters the topology of Pls in order to make it susceptible to proteolysis, or if it is directly

responsible for Pls cleavage, which would be a novel function for T11SS. The possibility that Gram-

negative Pls are secreted via T11SS and may be released from the cell surface in the process will be

an important factor to consider in ongoing efforts to target Pls homologs for vaccine development,

such as that being pursued for *Actinobacillus pleuropneumoniae* Pls (YP_001652736.1) (69). Also, the

Pls-like proteins have repeat rich regions and the possible glycosylation of *H. parahaemolyticus* Pls is

supported by our observations of its aberrant mobility in SDS-PAGE. It is possible that other

T11SS-dependent cargo, including lipoproteins, may also be glycosylated. If so, such glycoproteins

tethered to the cell surface would display glycans as surface antigens that may modulate host-

microbe recognition and immune evasion (70–72).

In summary, our findings have enhanced our understanding of potential effectors of the

novel T11SS and expanded the list of experimentally verified T11SS secreted cargo. We also have

expanded the field of genomic co-occurrence analysis by establishing a protocol that can serve as a

basis for more controlled and specific experimental designs. Exciting avenues of future research

could focus on generating an automated pipeline for controlling co-occurrences analyses,

determining if Pls is required for host-colonization by *Haemophilus parahaemolyticus*, investigating if

DUF1194 represents a novel T11SS cargo type, and exploring the relationship between one carbon

metabolism and T11SS.

Conflicts of interest

The author(s) declare that there are no conflicts of interest.

Funding information

This work was funded, in part, by funds awarded to H.G.-B. from the University of Tennessee-Knoxville (David and Sandra White Endowment) and the National Science Foundation (IOS-2128266).

Author contributions

ASG: Conceptualization, data curation, formal analysis, investigation, methodology, validation, writing original draft and review/editing

NCM: Conceptualization, data curation, investigation, formal analysis, methodology, writing original draft

HGB: Conceptualization, data curation, funding acquisition, investigation, methodology, project administration, resources, supervision, visualization, writing original draft and review/editing

Acknowledgements

We thank past and present members of the Goodrich-Blair lab for feedback on the development of controls for bioinformatic co-occurrence analyses.

References:

1.   Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar G, Sonnhammer ELL, et al. Pfam: The protein families database in 2021. Nucleic Acids Res [Internet]. 2021 Jan 8;49(D1):D412–9. Available from: https://doi.org/10.1093/nar/gkaa913

2.   Mudgal R, Sandhya S, Chandra N, Srinivasan N. De-DUFing the DUFs: Deciphering distant evolutionary relationships of Domains of Unknown Function using sensitive homology detection methods. Biol Direct [Internet]. 2015;10(1):38. Available from: https://doi.org/10.1186/s13062-015-0069-2

3.   Grossman AS, Mauer TJ, Forest KT, Goodrich-Blair H, Comstock L. A widespread bacterial secretion system with diverse substrates. mBio [Internet]. 2022 Apr 19;12(4):e01956-21. Available from: https://doi.org/10.1128/mBio.01956-21

4.   Shin HE, Pan C, Curran DM, Bateman TJ, Chong DHY, Ng D, et al. Prevalence of Slam-dependent hemophilins in Gram-negative bacteria. J Bacteriol [Internet]. 2024 May 30;206(6):e00027-24. Available from: https://doi.org/10.1128/jb.00027-24

5.   Hooda Y, Lai CCL, Moraes TF. Identification of a large family of Slam-dependent surface lipoproteins in Gram-negative bacteria. Front Cell Infect Microbiol [Internet]. 2017;7. Available from: http://dx.doi.org/10.3389/fcimb.2017.00207

6.   Hooda Y, Lai CCL, Judd A, Buckwalter CM, Shin HE, Gray-Owen SD, et al. Slam is an outer membrane protein that is required for the surface display of lipidated virulence factors in Neisseria. Nat Microbiol [Internet]. 2016;1(4). Available from: http://dx.doi.org/10.1038/nmicrobiol.2016.9

7.   Cowles CE, Goodrich-Blair H. The Xenorhabdus nematophila nilABC genes confer the ability of Xenorhabdus spp. to colonize Steinernema carpocapsae nematodes. J Bacteriol [Internet]. 2008 Jun 15;190(12):4121–8. Available from: https://doi.org/10.1128/JB.00123-08

8.   Bateman TJ, Shah M, Ho TP, Shin HE, Pan C, Harris G, et al. A Slam-dependent hemophore contributes to heme acquisition in the bacterial pathogen Acinetobacter baumannii. Nat Commun [Internet]. 2021;12(1):6270. Available from: https://doi.org/10.1038/s41467-021-26545-9

9. Latham RD, Torrado M, Atto B, Walshe JL, Wilson RK, Guss JM, et al. A heme-binding
    protein produced by Haemophilus haemolyticus inhibits non-typeable Haemophilus
    influenzae. Mol Microbiol [Internet]. 2020 Feb 1;113(2):381–98. Available from:
    https://doi.org/10.1111/mmi.14426

10. Grossman AS, Gell DA, Wu DG, Carper DL, Hettich RL, Goodrich-Blair H. Bacterial
    hemophilin homologs and their specific type eleven secretor proteins have conserved roles in
    heme capture and are diversifying as a family. J Bacteriol [Internet]. 2024 Mar
    20;206(6):e00444-23. Available from: https://doi.org/10.1128/jb.00444-23

11. Sam F, Brianna A, Arianna M, J HK, S RJ, Elan E, et al. Heme sequestration by hemophilin
    from Haemophilus haemolyticus reduces respiratory tract colonization and infection with
    non-typeable Haemophilus influenzae. mSphere [Internet]. 2024 Feb 21;9(3):e00006-24.
    Available from: https://doi.org/10.1128/msphere.00006-24

12. Zhang Z, Wang J, Wang J, Wang J, Li Y. Estimate of the sequenced proportion of the global
    prokaryotic genome. Microbiome. 2020 Sep;8(1):134.

13. Rogozin IB, Makarova KS, Murvai J, Czabarka E, Wolf YI, Tatusov RL, et al. Connected
    gene neighborhoods in prokaryotic genomes. Nucleic Acids Res. 2002 May;30(10):2212–23.

14. Tietz JI, Schwalen CJ, Patel PS, Maxson T, Blair PM, Tai HC, et al. A new genome-mining
    tool redefines the lasso peptide biosynthetic landscape. Nat Chem Biol. 2017 May;13(5):470–
    8.

15. The UniProt Consortium. UniProt: The universal protein knowledge base in 2021. Nucleic
    Acids Res [Internet]. 2021 Jan 8;49(D1):D480–9. Available from:
    https://doi.org/10.1093/nar/gkaa1100

16. Chen IMA, Chu K, Palaniappan K, Ratner A, Huang J, Huntemann M, et al. The IMG/M
    data management and analysis system v.7: content updates and new features. Nucleic Acids
    Res [Internet]. 2023 Jan 6;51(D1):D723–32. Available from:
    https://doi.org/10.1093/nar/gkac976

17. Mukherjee S, Stamatis D, Li CT, Ovchinnikova G, Bertsch J, Sundaramurthi JC, et al.
Twenty-five years of Genomes OnLine Database (GOLD): data updates and new features in
v.9. Nucleic Acids Res [Internet]. 2023 Jan 6;51(D1):D957–63. Available from:
https://doi.org/10.1093/nar/gkac974

18. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021 Aug;596(7873):583–9.

19. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making
protein folding accessible to all. Nat Methods. 2022 Jun;19(6):679–82.

20. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. The Phyre2 web portal for
protein modeling, prediction and analysis. Nat Protoc [Internet]. 2015;10(6):845–58. Available
from: https://doi.org/10.1038/nprot.2015.053

21. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al.
KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score
threshold. Bioinformatics [Internet]. 2020 Apr 1;36(7):2251–2. Available from:
https://doi.org/10.1093/bioinformatics/btz859

22. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for
functional characterization of genome and metagenome sequences. J Mol Biol. 2016
Feb;428(4):726–31.

23. Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH. UniRef: comprehensive and non-
redundant UniProt reference clusters. Bioinformatics [Internet]. 2007 May 15;23(10):1282–8.
Available from: https://doi.org/10.1093/bioinformatics/btm098

24. Buchan DWA, Jones DT. The PSIPRED protein analysis workbench: 20 years on. Nucleic
Acids Res. 2019 Jul;47(W1):W402–7.

25. Bhasin A, Chaston JM, Goodrich-Blair H. Mutational analyses reveal overall topology and
functional regions of NilB, a bacterial outer membrane protein required for host association
in a model of animal-microbe mutualism. J Bacteriol [Internet]. 2012 Apr 1;194(7):1763–76.
Available from: https://doi.org/10.1128/JB.06711-11

26. Miroux B, Walker JE. Over-production of proteins in Escherichia coli: mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. J Mol Biol. 1996 Jul;260(3):289–98.

27. Dumon-Seignovert L, Cariot G, Vuillard L. The toxicity of recombinant proteins in Escherichia coli: a comparison of overexpression in BL21(DE3), C41(DE3), and C43(DE3). Protein Expr Purif. 2004 Sep;37(1):203–6.

28. Orchard SS, Goodrich-Blair H. Identification and functional characterization of a Xenorhabdus nematophila oligopeptide permease. Appl Environ Microbiol [Internet]. 2004 Sep 1;70(9):5621–7. Available from: https://doi.org/10.1128/AEM.70.9.5621-5627.2004

29. Koontz L. Chapter one - TCA precipitation. In: Lorsch JBTM in E, editor. Laboratory methods in enzymology [Internet]. Academic Press; 2014. p. 3–10. Available from: http://www.sciencedirect.com/science/article/pii/B978012420119400001X

30. Tukey JW. Comparing individual means in the analysis of variance. Biometrics [Internet]. 1949 May 25;5(2):99–114. Available from: http://www.jstor.org/stable/3001913

31. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res. 2017 Jan;45(D1):D353–61.

32. Dobrindt U, Reidl J. Pathogenicity islands and phage conversion: evolutionary aspects of bacterial pathogenesis. International Journal of Medical Microbiology [Internet]. 2000;290(6):519–27. Available from: https://www.sciencedirect.com/science/article/pii/S143842210080017X

33. Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, et al. SignalP 5.0 improves signal peptide predictions using deep neural networks. Nat Biotechnol [Internet]. 2019;37(4):420–3. Available from: http://dx.doi.org/10.1038/s41587-019-0036-z

34. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, et al. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat

Biotechnol [Internet]. 2022;40(7):1023–5. Available from: https://doi.org/10.1038/s41587-
021-01156-3

35. Skaf MS, Polikarpov I, Stanković IM. A linker of the proline-threonine repeating motif
sequence is bimodal. J Mol Model. 2020 Jun;26(7):178.

36. Swearingen KE, Lindner SE, Shi L, Shears MJ, Harupa A, Hopp CS, et al. Interrogating the
Plasmodium sporozoite surface: identification of surface-exposed proteins and demonstration
of glycosylation on CSP and TRAP by mass spectrometry-based proteomics. PLoS Pathog.
2016 Apr;12(4):e1005606.

37. Sieber CMK, Paul BG, Castelle CJ, Hu P, Tringe SG, Valentine DL, et al. Unusual
Metabolism and Hypervariation in the Genome of a Gracilibacterium (BD1-5) from an Oil-
Degrading Community. mBio. 2019 Nov;10(6).

38. Calmettes C, Alcantara J, Yu RH, Schryvers AB, Moraes TF. The structural basis of
transferrin sequestration by transferrin-binding protein B. Nat Struct Mol Biol [Internet].
2012;19(3):358–60. Available from: https://doi.org/10.1038/nsmb.2251

39. Ostberg K, DeRocco A, Mistry S, Dickinson Mary K, Cornelissen CN. Conserved regions of
gonococcal TbpB are critical for surface exposure and transferrin iron utilization. Infect
Immun [Internet]. 2013 Sep 1;81(9):3442–50. Available from:
https://doi.org/10.1128/IAI.00280-13

40. Bleiziffer I, Eikmeier J, Pohlentz G, McAulay K, Xia G, Hussain M, et al. The Plasmin-
Sensitive Protein Pls in Methicillin-Resistant Staphylococcus aureus (MRSA) is a glycoprotein.
PLoS Pathog [Internet]. 2017 Jan 12;13(1):e1006110. Available from:
https://doi.org/10.1371/journal.ppat.1006110

41. Savolainen K, Paulin L, Westerlund-Wikström B, Foster TJ, Korhonen TK, Kuusela P.
Expression of pls, a gene closely associated with the mecA gene of methicillin-resistant
Staphylococcus aureus, prevents bacterial adhesion in vitro. Infect Immun. 2001
May;69(5):3013–20.

42. Liang KYH, Orata FD, Boucher YF, Case RJ. Roseobacters in a Sea of Poly- and Paraphyly:
Whole Genome-Based Taxonomy of the Family Rhodobacteraceae and the Proposal for the
Split of the "Roseobacter Clade" Into a Novel Family, Roseobacteraceae fam. nov. Front
Microbiol [Internet]. 2021 [cited 2024 Aug 18];12. Available from:
https://doi.org/10.3389/fmicb.2021.683109

43. Chen MH, Sheu SY, Chen CA, Wang JT, Chen WM. Roseivivax isoporae sp. nov., isolated
from a reef-building coral, and emended description of the genus Roseivivax. Int J Syst Evol
Microbiol. 2012 Jun;62(Pt 6):1259–64.

44. Crenn K, Serpin D, Lepleux C, Overmann J, Jeanthon C. Silicimonas algicola gen. nov., sp.
nov., a member of the Roseobacter clade isolated from the cell surface of the marine diatom
Thalassiosira delicatula. Int J Syst Evol Microbiol. 2016 Nov;66(11):4580–8.

45. Kessner L, Spinard E, Gomez-Chiarri M, Rowley DC, Nelson DR. Draft genome sequence
of Aliiroseovarius crassostreae CV919-312, the causative agent of Roseovarius Oyster Disease
(formerly Juvenile Oyster Disease). Genome Announc. 2016 Mar;4(2).

46. Kim YO, Park S, Nam BH, Lee C, Park JM, Kim DG, et al. Ascidiaceihabitans donghaensis
gen. nov., sp. nov., isolated from the golden sea squirt Halocynthia aurantium. Int J Syst Evol
Microbiol. 2014 Dec;64(Pt 12):3970–5.

47. Ivanova EP, Gorshkova NM, Sawabe T, Zhukova N V, Hayashi K, Kurilenko V V, et al.
Sulfitobacter delicatus sp. nov. and Sulfitobacter dubius sp. nov., respectively from a starfish
(Stellaster equestris) and sea grass (Zostera marina). Int J Syst Evol Microbiol. 2004 Mar;54(Pt
2):475–80.

48. Alamos P, Tello M, Bustamante P, Gutiérrez F, Shmaryahu A, Maldonado J, et al.
Functionality of tRNAs encoded in a mobile genetic element from an acidophilic bacterium.
RNA Biol. 2018;15(4–5):518–27.

49. Baek M, DiMaio F, Anishchenko I, Dauparas J, Ovchinnikov S, Lee GR, et al. Accurate
prediction of protein structures and interactions using a three-track neural network. Science.
2021 Aug;373(6557):871–6.

50.   Bergelson JM, Hemler ME. Integrin- Ligand Binding: Do integrins use a 'MIDAS touch' to
823   grasp an Asp? Current Biology [Internet]. 1995;5(6):615–7. Available from:
824   https://www.sciencedirect.com/science/article/pii/S0960982295001242

826   51.   Cantí C, Nieto-Rostro M, Foucault I, Heblich F, Wratten J, Richards MW, et al. The metal-
827   ion-dependent adhesion site in the Von Willebrand factor-A domain of alpha2delta subunits
828   is key to trafficking voltage-gated Ca2+ channels. Proc Natl Acad Sci U S A. 2005
829   Aug;102(32):11230–5.

831   52.   Hantke K. Is the bacterial ferrous iron transporter FeoB a living fossil? Trends Microbiol.
832   2003 May;11(5):192–5.

834   53.   Ollis AA, Postle K. ExbD mutants define initial stages in TonB energization. J Mol Biol.
835   2012 Jan;415(2):237–47.

837   54.   Levengood SK, Beyer WFJ, Webster RE. TolA: a membrane protein involved in colicin
838   uptake contains an extended helical region. Proc Natl Acad Sci U S A. 1991 Jul;88(14):5939–
839   43.

841   55.   Tjalsma H, Kontinen VP, Prágai Z, Wu H, Meima R, Venema G, et al. The role of
842   lipoprotein processing by signal peptidase II in the Gram-positive eubacterium bacillus
843   subtilis. Signal peptidase II is required for the efficient secretion of alpha-amylase, a non-
844   lipoprotein. J Biol Chem. 1999 Jan;274(3):1698–707.

846   56.   Agrawal N, Lesley SA, Kuhn P, Kohen A. Mechanistic studies of a flavin-dependent
847   thymidylate synthase. Biochemistry [Internet]. 2004 Aug 1;43(32):10295–301. Available from:
848   https://doi.org/10.1021/bi0490439

850   57.   Dawadi S, Kordus SL, Baughn AD, Aldrich CC. Synthesis and analysis of bacterial folate
851   metabolism intermediates and antifolates. Org Lett [Internet]. 2017 Oct 6;19(19):5220–3.
852   Available from: https://doi.org/10.1021/acs.orglett.7b02487

854   58.   Luk LYP, Javier Ruiz-Pernía J, Dawson WM, Roca M, Loveridge EJ, Glowacki DR, et al.
855   Unraveling the role of protein dynamics in dihydrofolate reductase catalysis. Proc Natl Acad
856   Sci U S A. 2013 Oct;110(41):16344–9.

59.   Mendel RR. The molybdenum cofactor. J Biol Chem. 2013 May;288(19):13165–72.

60.   Zhong Q, Kobe B, Kappler U. Molybdenum enzymes and how they support virulence in
        pathogenic bacteria [Internet]. Vol. 11, Frontiers in Microbiology. 2020. Available from:
        https://www.frontiersin.org/articles/10.3389/fmicb.2020.615860
860
861
862

61.   Van Bibber M, Bradbeer C, Clark N, Roth JR. A new class of cobalamin transport mutants
        (btuF) provides genetic evidence for a periplasmic binding protein in Salmonella
        typhimurium. J Bacteriol. 1999 Sep;181(17):5539–41.
864
865
866

62.   Peyvandi F, Garagiola I, Baronciani L. Role of von Willebrand factor in the haemostasis.
        Blood Transfus. 2011 May;9 Suppl 2(Suppl 2):s3-8.
868
869

63.   Von Willebrand EA. Hereditary pseudohaemophilia. Haemophilia. 1999 May;5(3):223–31;
        discussion 222.
871
872

64.   Islam ST, Jolivet NY, Cuzin C, Belgrave AM, My L, Fleuchot B, et al. Unmasking of the von
        Willebrand A-domain surface adhesin CglB at bacterial focal adhesions mediates
        myxobacterial gliding motility. Sci Adv [Internet]. 2023 May 8;9(8):eabq0619. Available from:
        https://doi.org/10.1126/sciadv.abq0619
874
875
876
877

65.   Lee C, Park C. Bacterial responses to glyoxal and methylglyoxal: Reactive electrophilic
        species. Int J Mol Sci. 2017 Jan;18(1).
879
880

66.   Anaya-Sanchez A, Feng Y, Berude JC, Portnoy DA. Detoxification of methylglyoxal by the
        glyoxalase system is required for glutathione availability and virulence activation in Listeria
        monocytogenes. PLoS Pathog. 2021 Aug;17(8):e1009819.
882
883
884

67.   Fantappiè L, Irene C, De Santis M, Armini A, Gagliardi A, Tomasi M, et al. Some Gram-
        negative lipoproteins keep their surface topology when transplanted from one species to
        another and deliver foreign polypeptides to the bacterial surface. Molecular & Cellular
        Proteomics [Internet]. 2017 Jul 1;16(7):1348 LP – 1364. Available from:
        http://www.mcponline.org/content/16/7/1348.abstract

68. Konar M, Rossi R, Walter H, Pajon R, Beernink PT. A mutant library approach to identify improved Meningococcal Factor H Binding Protein vaccine antigens. PLoS One [Internet]. 2015 Jun 9;10(6):e0128185. Available from: https://doi.org/10.1371/journal.pone.0128185

69. Cao Y, Gao L, Zhang L, Zhou L, Yang J, Deng L, et al. Genome-wide screening of lipoproteins in Actinobacillus pleuropneumoniae identifies three antigens that confer protection against virulent challenge. Sci Rep. 2020 Feb;10(1):2343.

70. Schmidt MA, Riley LW, Benz I. Sweet new world: glycoproteins in bacterial pathogens. Trends Microbiol. 2003 Dec;11(12):554–61.

71. Tortorelli G, Rautengarten C, Bacic A, Segal G, Ebert B, Davy SK, et al. Cell surface carbohydrates of symbiotic dinoflagellates and their role in the establishment of cnidarian-dinoflagellate symbiosis. ISME J. 2022 Jan;16(1):190–9.

72. Zhou M, Wu H. Glycosylation and biogenesis of a family of serine-rich bacterial adhesins. Microbiology (Reading). 2009 Feb;155(Pt 2):317–27.

909
910

Supplemental Figure 1. **T11SS (DUF560) protein size distribution (in amino acids).** After
examination of the size distribution, median was chosen as an appropriate measure of representative
protein size over mean due to left-skewed distribution. The blue dataset was used to perform the
T11SS co-occurrence analysis described in this manuscript. In the intervening time more homologs
have been added to the Pfam database, the red dataset reflects these additions and their impact on
the size distribution.
918

Figure 1. **Design for a novel control technique for genomic neighborhood co-occurrence analysis.** Schematic diagram outlining the process we used to remove nonspecific co-occurrences from a genomic neighborhood analysis, with commentary highlighting how we used this outline to refine our T11SS co-occurrence database.

Figure 2. **Host-associated T11SS genome neighborhoods reveal conserved association with iron/heme uptake, export, and one-carbon metabolism**. KofamKOALA uses hidden Markov models to assign functions to query sequences and reveal shared pathways. Cellular functions were estimated using brite hierarchies and assigned to known pathways.

932 Supplemental Figure 2. **BlastKOALA supports T11SS association with iron/heme uptake,**
933 **protein export, and one-carbon metabolism**. BlastKOALA uses the BLAST alignment algorithm
934 to assign functions to query sequences and reveal shared pathways. Cellular functions were estimated
935 using brite hierarchies and assigned to known pathways.

Figure 3. **Secondary structure of predicted T11SS-dependent cargo reveals diverse and novel gene architectures and effector domains**. PsiPred 4 and DisoPred 3 use amino acid sequence to annotate secondary structures highlighting a diversity of N-terminal domains with proline rich repeats, intrinsically disordered regions, α-helical repeats, and ligand binding handles. For each of the detected cargo architectures the representative sequence from the largest UniRef50 cluster was trimmed of it signal peptide to reflect a mature protein.

Figure 4. **AlphaFold2 annotation of predicted T11SS-dependent cargo reveals conservation of C-terminal or centrally located β-barrel domains**. For each of the detected cargo architectures the representative sequence from the largest UniRef50 cluster was used to predict a representative structure. Green residues indicate C-terminal β-barrel domains. Blue residues indicate putative ligand binding handle domains. Orange residues indicate glycosylation motif repeats within the Pls-like protein architecture. Yellow residues indicate regions predicted to be disordered.

Figure 5. **AlphaFold2 annotation of predicted T11SS-dependent cargo reveals three distinct**

domains ubiquitous to cargo. Solved **(D, G, J)** or predicted **(A, B, C, E, F, H, I, K)** structures

from representative UniRefF50 cluster sequences of predicted **(A, B, D, E, F, H, K)** or known

cargo **(D, G, I, J)**. Each structure was separated into three sections representing common structural

domains: N-terminal effector (blue), pedestal (red), and C-terminal β-barrel (gray barrel residues and

purple loop residues). N-terminal effector domains varied in their predicted structural motifs and

included disordered **(B, C, D, H, I)**, α-helical **(A, E, K)**, β-strand **(D, F, J)**, and barrel **(G)**. Pedestal

domains comprised 4-stranded groove **(A)**, 4-stranded "wing" **(B-D),** 4-stranded sheet **(E, F)**, 5-

stranded sheet **(G-I)**, or 6-stranded/extended β-sheet **(J, K)**. Some of these β-strands are donated

by primary sequences from either the N-terminus or the C-terminus of the protein. Barrel domains

exhibited variation in the presence of extended loops, helices, or strands in predicted surface- or

periplasmic-exposed regions, as indicated by purple coloration.

Figure 6. **Localization of FLAG tagged Pls in the absence and presence of its cognate T11SS secretor.** *E. coli* cells expressing Pls-FLAG alone or Pls-FLAG alongside its cognate T11SS (TepS) were probed using dot blots (cell surfaces and lysates) or Western blotting (supernatant and extracellular vesicles) to see how expression of the T11SS protein impacted protein localization. Supernatant secretion of Pls was only detectable in the presence of the T11SS TepS. The extracellular vesicle fraction collected via ultracentrifugation showed some Pls localization in the absence of TepS, however that localization was significantly increased by TepS. Pls was detected on the surface of both treatments equally, and cell lysates showed that the level of Pls expression between the treatments was equivalent.

974
975
Supplemental Figure 3. **Localization of FLAG tagged Pls in the absence and presence of its**
cognate T11SS secretor. *E. coli* cells expressing Pls-FLAG alone or Pls-FLAG alongside its cognate
T11SS (TepS) were probed using dot blots (cell surfaces and lysates) or Western blotting
(supernatant and extracellular vesicles) to see how expression of the T11SS protein impacted protein
localization. **(A)** a representative Western blot demonstrating TepS dependent secretion of Pls. **(B)**
A representative Coomassie stain showing total supernatant protein content. **(C)** Immuno-dot blots
demonstrating that TepS was not required for surface exposure of Pls when expressed in *E. coli*.
983

986 Figure 7. **Marine T11SS genome neighborhoods reveal association with one-carbon**
987 **metabolism, nucleotide metabolism, and the glyoxalase-detoxification pathway**.
988 KofamKOALA uses hidden Markov models to assign functions to query sequences and reveal
989 shared pathways. Cellular functions were estimated using brite hierarchies and assigned to known
990 pathways.

Fig. 8. **A conserved marine cluster 3, T11SS OMP locus encodes predicted cargo proteins.** A schematic representation of select examples of a T11SS OMP-encoding, syntenic locus found with genomes of indicated strains of *Rhodobacteraceae* and *Roseobacteriaceae* **(A)**. Each gene is represented by a block arrow (not to scale), and the *Ruegeria pomeroyi* locus tags of each homolog are indicated at the top. In all 11 genomes examined, the syntenic locus included genes predicted to encode a T11SS OMP (turquoise arrows) as well as the top cluster 3 co-occurring functions: flavin-dependent thymidylate synthase (gold arrows), lactoylglutathione lyase (neon green arrows), and one or more DUF1194/von Willebrand factor proteins (pink arrows). In all but one, the locus includes a putative T11SS cargo protein (light blue arrows), identified based on the presence of a predicted C-terminal 8-stranded β-barrel domain. Predicted Alphafold2 or Phyre2 predicted structures of putative cargo proteins encoded within these loci is shown in **(B-K)**. Color coding is used to indicate the putative effector (dark blue), pedestal (red), barrel (gray), and surface or periplasmic loops (purple) of each protein. The top five appear to have similar structural motifs, including an α-helix in one of the β-barrel loops, antiparallel β-strands in the N-terminal domain (blue), and a 4-stranded sheet pedestal domain (red). The remaining five structures show the overall domains of N-terminal (blue), pedestal (red) and β-barrel (gray), but the N-terminal domains are predicted to have different structural motifs relative to the others examined.

Supplemental Figure 4. **BlastKOALA supports T11SS association with one-carbon metabolism, nucleotide metabolism, and the glyoxalase-detoxification pathway.** BlastKOALA uses the BLAST alignment algorithm to assign functions to query sequences and reveal shared pathways. Cellular functions were estimated using brite hierarchies and assigned to known pathways.

1015

Figure 9 **Representative structures of DUF1194 proteins located within a T11SS genomic**
neighborhood in *Ascidiaceihabitans donghaensis*. These three DUF1194 homologs are
encoded adjacent to a T11SS OMP-encoding gene within a microbe isolated from a tunicate host.
These proteins appear to be paralogs of each other based on sequence similarity. Signal peptides
were predicted with SignalP 6 and trimmed from the sequences prior to submission to PsiPred **(A)**
and RoseTTAFold **(B).**
1022

1023 Supplemental Files:

1024 Supplemental File 1: **T11SSGenomicCo-occurrence.xlsx**
1025 Controlled co-occurrence analysis of all domains and genes significantly co-occurring with T11SS in
1026 a cluster of host-associated organisms (cluster 1) or a cluster of marine organisms (cluster 3).
1027 Significant genes were then clustered into UniRef50 clusters to combine close homologs and mined
1028 for predicted T11SS cargo proteins via homology to known cargo proteins.
1029
1030 Supplemental File 2: **Co-occurrenceFunctionalAnnotations.xlsx**
1031 KEGG based analysis genes significantly co-occurring with host associated T11SS (cluster 1) and a
1032 marine T11SS (cluster 3).
1033
1034 Supplemental File 3: **Cluster3CargoLoci.xlsx**
1035 Selected Cluster 3 predicted cargo proteins co-occurring with T11SS.
1036
1037 Supplemental File 4: **T11SS-DUF1194Co-occurrence.xlsx**
1038 Co-occurrence of Marine associated T11SS alongside DUF1194 domain encoding genes, color
1039 coded to indicate host-associated organisms.

DUF560, our co-occurrence query, comprises outer membrane proteins conserved among Proteobacteria. OMPs were determined using the GO term

1. Search for proteins that fit similar criteria of co-occurrence query on Uniprot

2. Determine median protein size from the original query list

This step is necessary to narrow the control list. Mean protein size was considered, but due to left-skewed protein size distribution, the median was utilized to decrease influence of outliers.

4. Remove any duplicates between the control and query list and take a random sample of this control list to the original query list

3. Download protein accession IDs within +/1 X% of median protein size

X% will be determined by how many +/- amino acids results in a # of proteins that is close to, yet over, the query list

We utilized Rapid ORF Description & Evaluation Online (RODEO).

5. Run this control list through the same protein co-occurrence software as the original query list

6. Take the ratio of the number of co-occurrences in the control list/# of co-occurrences in the original query list

This will determine the False Discovery Rate (FDR), where the lower the ratio, the more confident the user can be that the co-occurrence is unique to the original query.

8. Determine a cutoff # of co-occurrences to which any # above is a significant amount of protein co-occurrences

7. Divide the original query list into 2 categories: co-occurrences that pass the FDR and those that do not

We utilized an FDR of 10%, which is a common rate used to determine the amount of false positive results present.

9. Extract all proteins with the PFam IDs for both lists on UniProt

10. Run both of these new protein lists through KOALA to determine metabolic function

This will result in two KEGG annotated datasets that describe the broad metabolic functions of the types of proteins that are unique or common to the original query list.

Effector

A_eff B_eff C_eff D_eff E_eff F_eff G_eff H_eff I_eff J_eff K_eff

Pedestal

A_ped B_ped C_ped D_ped E_ped F_ped G_ped H_ped I_ped J_ped K_ped

Barrel

A_barr B_barr C_barr D_barr E_barr F_barr G_barr H_barr I_barr J_barr K_barr

A. *Haemophilus haemolyticus* A0A52GK7Y1, Pllis

B. *Moraxella plurainimalium* A0A17ECV7, Hhp

C. *Elikenella* sp. NML080894 A0A1B6VSV4, Hhp

D. *Neisseria meningitidis* 5T5F, Hhp

E. *Haemophilus haemolyticus* A0A1B8PSS3

F. *Ruegeria pomeroyi* Q5LTY8, SPO1272

G. *Neisseria meningitidis* 3v8u, Tps

H. *Haemophilus* A0A024A3X0

I. *Xenorhabdus nematophila* I3V0Y9, NilC

J. *Haemophilus haemolyticus* 6CM6, Hpl

K. *Kingella negevensis* A0A23BT9F7

A

Marine-associated T11SS Co-occurence KofamKOALA cellular function analysis

Category	Value
ENZYMES (KO01000)	276
AMINO ACID RELATED ENZYMES (KO01007)	75
PROKARYOTIC DEFENSE SYSTEM (KO02048)	68
TRANSCRIPTION FACTORS (KO03000)	59
TRANSPORTERS (KO02000)	16
TRANSFER RNA BIOGENESIS (KO03016)	15
CHROMOSOME AND ASSOCIATED PROTEINS (KO03036)	6
DNA REPLICATION PROTEINS (KO03032)	6
PEPTIDOGLYCAN BIOSYNTHESIS/DEGRADATION (KO01011)	6
TRANSLATION FACTORS (03012)	5
PEPTIDASES AND INHIBITORS (KO01002)	2
TWO-COMPONENT SYSTEM (KO02022)	1
PROTEIN KINASES (KO01001)	1

(x-axis: 0, 50, 100, 150, 200, 250, 300)

B

Marine-associated T11SS Co-occurence KofamKOALA pathway analysis

Category	Value
ONE CARBON POOL BY FOLATE (00670)	104
PYRIMIDINE METABOLISM (00240)	88
NUCLEOTIDE METABOLISM (01232)	88
PYRUVATE METABOLISM (00620)	82
BIOSYNTHESIS OF AMINO ACIDS (01230)	60
ANTIFOLATE RESISTANCE (01523)	33
FOLATE BIOSYNTHESIS (00790)	16
BIOSYNTHESIS OF COFACTORS (01240)	16
AMINOACYL-TRNA BIOSYNTHESIS (00970)	15
PEPTIDOGLYCAN BIOSYNTHESIS (00550)	6
BIOSYNTHESIS OF NUCLEOTIDE SUGARS (01250)	6
AMINO SUGAR AND NUCLEOTIDE SUGAR METABOLISM (00520)	6
CATIONIC ANTIMICROBIAL PEPTIDE RESISTANCE (01503)	2
TWO-COMPONENT SYSTEM (02020)	1

(x-axis: 0, 20, 40, 60, 80, 100, 120)

A.

	SPO1070	SPO1071	SPO1072	SPO1073	SPO1089

Ruegeria pomeroyi DSS-3

Rhodovulum sp. NI22

Aliiroseovarius crassostreae CV919-312

Roseivivax isoporae LMG 25204

Pseudoceanicola marinus LMG 23705

Amylibacter cionae CGMCC 1.15880

Rhodobacteraceae bacterium PD-2

Ascidiaceihabitans donghaensis CECT8599

Sulfitobacter sp. NAS-14.1

Roseovarius sp. THAF27

Silicimonas algicola DSM 103371

Legend:
- thymidylate synthase
- 8-standed barrel protein
- T1188 OMP
- Isctoylglutathione lyase
- DUF1194

B. Q5LTY8
R. pomeroyi DSS-3

C. A0A095DSB7
Rhodovulum sp. NI22

D. A0A0P7IGY3
A. crassostreae CV919-312

E. X7F6N0
R. isoporae LMG 25204

F. A0A1X6Z1X8
P. marinus LMG23705

G. c7JrdA
A. cionae CGMCC 1.15880

H. A0A5P9H0N7
Roseovarius sp. THAF27

I. X6KVJ4
R. bacterium PD-2

J. A0A2R8BC28
A. donghaensis CECT8599

K. A3SXD3
Sulfitobacter NAS14.1

```
Q C R Q A L A L G L D V S G S V D A R E Y R L Q L D G V A A L N A P T V Q A A L W D T P G S H V S
L L I Y E W S G P A D Q A L I L P W T P I M S V A D L D D V T E I L Q Q T V R R A T S P G T A L G D
A M I L G A R H L A M Q P E C W K H T L D I S G D G K S N L G A R P R D V K P V L L H D D I T I N A
L V I G A D A P Q S G D N R Q A D I A G L S A Y F H A E V L G P D S F V E T A L G F D A Y A Q A M
Q R K L E R E L Q T L M S E H D L D A G T Q K R D L R
```

```
C R L A L L L A V D V S S S V D A A E D Q L Q R G G I V S A L I A P E V Q N A F F A S D V P V A L G
I Y E W S G R Y N Q E V L L D W V I I D T P A D L L A A A E T V A D S K R S H N D F P T A M G Y A L
G F G A K M L A R A P K C D T K T L D M A G D G Q N N E G F G P Q A A Y R E F P F D D V T V N G L V
V N G A D F E A E T G L I A F Y E G Q V M H G P G A F I V V A Q G F D D Y A R A M R R K L V R E L S
L P I L G A L P L P G P G
```

```
D C R L A L L L A V D V S N S V N A T E D A L Q R K G L S N A L N A A N V K R A V F A T D Q T V A L
A A F E W S G R H H Q Q V M L D W T I L K T P A D L S R A A H T I A H S K R G Q I D F P T A M G H A
L T F G A E M F A K A P D C L F K T L D V A G D G K S N T G Y P P K L A Y A D P A Y D E I T V N G L
V V R V L D L G M D L G V T A Y Y E A H V K H G P G A F I E V A D G F E D Y T R A M R T K L M R E L
S P T V L S A R P A S D N P G
```