Potentiation of active locomotor state by spinal-projecting serotonergic neurons

Sara J. Fenstermacher[1]*, Annie Vonasek[1], Hannah Gattuso[1], Corryn Chaimowitz[1], Susan M. Dymecki[2], Thomas M. Jessell, Jeremy S. Dasen[1]*

[1]Neuroscience Institute, Department of Neuroscience and Physiology, NYU School of Medicine
[2]Department of Genetics, Harvard Medical School
*Corresponding authors: sara.fenstermacher@nyulangone.org, jeremy.dasen@nyulangone.org

Abstract

Animals produce diverse motor actions that enable expression of context-appropriate behaviors.
Neuromodulators facilitate behavioral flexibility by altering the temporal dynamics and output of
neural circuits. Discrete populations of serotonergic (5-HT) neurons target circuits in the
brainstem and spinal cord, but their role in the control of motor behavior is unclear. Here we
define the pre- and post-synaptic organization of the spinal-projecting serotonergic system and
define a role in locomotor control. We show that while forebrain-targeting 5-HT neurons
decrease their activity during locomotion, subpopulations of spinal projecting neurons increase
their activity in a context-dependent manner. Optogenetic activation of ventrally projecting 5-HT
neurons does not trigger initiation of movement, but rather enhances the speed and duration of
ongoing locomotion over extended time scales. These findings indicate that the descending
serotonergic system potentiates locomotor output and demonstrate a role for serotonergic
neurons in modulating the temporal dynamics of motor circuits.

Introduction

Neuromodulators act throughout the central nervous system to enable behavioral flexibility in changing environments[1,2]. Neuromodulators alter the dynamics of neural circuits, often leading to profound changes in network output. In the motor system, rhythm-generating circuits that produce essential behaviors such as walking, breathing, and swimming, are under strong neuromodulatory control[3,4]. While studies of rhythm-generating circuits have yielded insights into the mechanisms of neuromodulation, how neuromodulators are integrated with instructive motor commands and effector neurons to control behavior is less understood.

The monoamine serotonin (5-HT) is an evolutionarily-conserved modulator of motor circuits, including the rhythmically active circuits required to produce locomotion[5-20]. In vitro and pharmacological manipulations have shown serotonin is a potent modulator of spinal motor circuits[21]. Application of 5-HT to ex vivo neonatal spinal cord induces rhythmic activity similar to that observed during locomotion[14,22]. In vivo, 5-HT agonists and antagonists significantly alter the magnitude and timing of muscle activity during locomotion, as well as the strength of spinal reflexes[7,15,23,24]. At the cellular level, 5-HT promotes the excitability of spinal motor neurons (MNs) and ventral interneurons (INs)[25,26], in part mediated by the generation of calcium-dependent plateau potentials[11,18,27]. Despite a rich history of investigating 5-HT actions in the spinal cord, a major challenge has been to resolve how descending serotonergic pathways operate during movement and to determine their role in locomotor behavior.

The majority of serotonergic neurons reside within the raphe nuclei near the midline[28,29] and are divided into two discrete clusters: a rostral group which largely projects to the forebrain and a caudal group that projects to the brainstem and spinal cord[30]. Within the caudal group, raphe obscurus (ROb) and raphe pallidus (RPa) project into ventral regions of the spinal cord, while raphe magnus (RMg) targets dorsal laminae[31-37]. Recent studies combining genetic fate-mapping and molecular profiling have defined multiple anatomically and functionally distinct serotonergic neuron populations[36-40]. In particular, genetically-distinct serotonergic nuclei in the ventral medulla target either sensory or motor regions with the brainstem and spinal cord. Serotonergic neurons with a history of expressing the transcription factor encoding gene *Egr2* project dorsally while those expressing the neuropeptide-encoding gene *Tac1* terminate ventrally[36,37].

Physiological studies revealed that the activity of 5-HT neurons correlates with movement. Extracellular recordings in cat showed that the firing rates of ROb and RPa neurons increase during treadmill-induced locomotion, indicating the activity of spinal-projecting 5-HT neurons is highly correlated to motor activity. By contrast, these neurons are relatively silent during REM sleep, a behavioral state characterized by cessation of movement and decreased muscle tone[41]. These observations suggest that descending modulation of spinal circuits by serotonergic neurons is critical for regulating motor function. Whether spinal-projecting serotonergic neurons affect the initiation, intensity, or duration of locomotor output is unresolved.

Here, we define the cellular and synaptic architecture of spinal cord-projecting serotonergic neurons and examine their function in motor control. We find that ROb/Pa provides direct input to spinal MNs and receives input from locomotor control regions, indicating the descending serotonergic system is recruited in parallel with motor command systems. The activity of ROb/Pa neurons corelates directly with locomotor activity and is distinct in nature and timing from other serotonergic populations. We show that activation of ROb/Pa neurons targeting

ventral spinal circuits produce long-lasting increases in running duration and speed. Our studies
reveal that anatomically and functionally distinct serotonergic neuron populations targeting
ventral spinal motor circuits act to regulate locomotor behavior over long time scales.

Results

Spinal motor circuits receive biased 5-HT input from medullary raphe nuclei

To examine the spinal cord targets of subtypes of medullary serotonergic neurons, we first
analyzed the distribution of spinal cord 5-HT in adult mice. Punctate 5-HT immunoreactivity was
detected throughout the rostrocaudal extent of the spinal cord, with strongest labeling observed
in the superficial dorsal horn, intermediolateral column, and ventral horn (Fig1.a,b,c), confirming
earlier work[28]. In the ventral horn, serotonin fibers densely targeted ChAT$^+$ motor neurons
(Fig1d), including both limb-innervating lateral motor column (LMC) and axial-innervating medial
motor column (MMC) neurons. Serotonin puncta were also observed in proximity to the cell
bodies of ventral interneurons, including excitatory Chx10$^+$ V2a neurons (Fig.1e).

Building from prior serotonergic neuron subtype identification and efferent mapping[36,37], we
performed anterograde labeling of genetically defined 5-HT populations to visualize serotonergic
projections to the spinal cord. We used an intersectional genetic-approach to visualize
synaptophysinGFP-labelled terminals[42] of two caudal 5-HT populations defined by co-
expression of *Pet1* with *Egr2*[36] or *Tac1*[37] (Extended Data Fig.1a,b). We observed that for *Egr2-*
Pet1 neurons, whose soma reside predominantly in Raphe Magnus (RMg)[36] (Extended Data
Fig.1d), synaptophysinGFP-labeled terminals are exclusively in the dorsal horn of cervical,
thoracic, and lumbar segments (Fig.1f,g,h,i, Extended Data Fig.1c). By contrast, *Tac1-Pet1*
neurons, whose soma reside within the Raphe Obscurus (ROb), Raphe Pallidus (RPa), and
Lateral Paragigantocellularis (LPGi) nuclei of the caudal medulla[37] (Extended Data Fig.1d),
selectively target the ventral horn throughout rostrocaudal levels and densely surround MNs
(Fig.1j,k,l,m, Extended Data Fig.1c). Genetically-distinct sets of *Pet1* neurons, therefore
innervate non-overlapping domains along the dorsoventral axis of the spinal cord (Fig.1n), likely
providing separate channels for serotonergic modulation of sensory input and motor output.

While neuromodulators such as serotonin are often released extrasynaptically to signal via
volume transmission, MNs receive numerous 5-HT synaptic contacts upon their soma and
dendrites[43]. To determine the origin of this direct and synaptic serotonergic neuron input to MNs,
we performed retrograde tracing from spinal MNs using G-deleted rabies virus[44], which travels
exclusively through synapses (Fig1o). To restrict expression of the avian TVA receptor and
rabies glycoprotein to spinal MNs we injected AAV-FLEX-TVA-G into the lateral ventricle of
Chat-Cre mice at P0 (Fig.1p)[45]. Eight weeks later, we infected spinal MNs by delivery of EnvA
pseudotyped, G-deleted N2c rabies virus fused to GFP (EnvA-N2cΔG-GFP) into the spinal cord
(Fig.1o,p). We identified serotonin neurons by presence of tryptophan hydroxylase 2 (TPH2)
and quantified GFP$^+$TPH2$^+$ neurons in the brainstem (Fig.1q, Extended Data Fig.2a,b). Rabies-
labeled TPH2$^+$ cells were found almost exclusively within the medullary raphe (ROb, RPa, RMg)
and lateral paragigantocellularis (LPGi) nuclei, with over 60% of GFP$^+$TPH2$^+$ cells located in
ROb (Fig.1r). Over 75% of GFP$^+$TPH2$^+$ cells were found within the neighboring ROb and RPa
nuclei, thus we will henceforth refer to this population as 'ROb/Pa'. Only a few cells were
labeled in the dorsal raphe nucleus (DRN)(Fig.1r). By comparison, similar transsynaptic tracing
assays from V2a INs (using *Chx10-Cre* mice), labeled very few 5-HT neurons (Extended Data
Fig.2c-f), suggesting that excitatory spinal INs do not receive substantial direct synaptic input
from serotonergic neurons, despite local innervation.

We next examined whether serotonergic neurons target specific spinal populations, or instead
broadly innervate multiple segments. We injected AAV2r-FLEX-tdTomato into the cervical spinal
cord of *Pet1-Cre* mice to retrogradely label serotonergic neurons (Extended Data Fig.3).
Following this, we observed tdTomato[+] fibers throughout both thoracic and lumbar levels,
demonstrating that descending 5-HT neurons exhibit a highly collateralized structure with
widespread targets across rostrocaudal spinal segments. Together, these studies showed that
within the spinal cord, fibers from ROb and Pa serotonergic neurons target neurons in the
ventral horn, provide direct input to spinal MNs, and that these inputs distribute to more than
one axial level.

112 Activity of ventrally-projecting serotonin neurons increases during locomotion

The activity of serotonergic neurons can depend on behavioral context, with specific populations
increasing or decreasing their activity during movement[41,46,47]. However, the activity of
serotonergic populations during locomotion are just beginning to be defined. We therefore
measured the activity of three serotonergic neuron populations (ROb/Pa, RMg, or DRN) in mice
during both wheel and treadmill running. To record the activity of specific subpopulations of 5-HT
neurons, we performed fiber photometry in *Pet1-Cre* transgenic mice that were infected to
express the calcium indicator GCaMP6s (AAV-FLEX-GCaMPs6, Fig.2c,g,k, Extended Data
Fig.5b,d,f). One week following viral injection, a low-profile running wheel was placed into each
mouse cage for introduction and practice during the remainder of the experiment. One week
later, mice were acclimated to the recording area and allowed to run on the wheel and treadmill.
Starting 3 weeks after the viral injection, photometry signals were recorded during 30-minute
sessions on the wheel (Fig.2a,b) and for 10s intervals on the treadmill (Fig.2o) with delivery of
470nm excitation light to monitor neural activity and 405nm control light to assess motion
artifact.

We observed that ROb/Pa *Pet1* neurons display a sustained increase in activity during periods
of running (Fig.2b,d, Extended Data Fig.4a). Averaged dF/F during the start and stop of run
bouts shows an increase in activity at running onset and decrease at offset (Fig2e,q, Extended
Data Fig.5g). A scatterplot of averaged dF/F values during run bouts vs. rest periods just prior to
the run shows the neural activity increased during locomotion (Fig.2f). Similarly, we observed a
robust increase in GCaMP signal during treadmill-evoked locomotion (Fig.2o,p, Extended Data
Fig.5a). During both spontaneous and evoked locomotion, ROb/Pa *Pet1* neurons displayed a
rapid increase activity at the onset of movement (Fig.2p,q). Thus, ROb/Pa *Pet1* neurons that
project to the spinal ventral column exhibit activity that positively correlates with locomotion.

By contrast, the activity of DRN *Pet1* neurons were anti-correlated with locomotion (Fig.2k,
Extended Data Fig.4c). DRN activity decreased during running (Fig.2l), with a consistent
reduction in signal at the start of running and an increase in activity when the animal stops
(Fig.2m,w). The scatterplot of averaged dF/F values during run bouts vs. rests shows the neural
activity during run bouts shifted below zero, reflecting decreased activity during locomotor bouts
(Fig.2n). We also observed a similar decrease in DRN activity during treadmill locomotion
(Fig.2u,v, Extended Data Fig.5e). Thus, DRN and ROb/Pa *Pet1* neurons display reciprocal
patterns of activity during locomotion.

Finally, to determine whether movement-correlated activity in ROb/Pa is unique to ventral spinal
cord-projecting 5-HT neurons, we compared GCaMP signal to the neighboring dorsal spinal
cord-projecting RMg (Fig.2g). We observed that RMg *Pet1* neuron activity did not strongly
correlate with locomotion when animals ran freely on the wheel (Fig.2h-j, Extended Data

Fig.4b). When animals were forced to perform treadmill running however, RMg *Pet1* neuron
activity was positively correlated to movement, with a slow rise in activity during locomotion and
decrease during rest (Fig.2r,s, Extended Data Fig.5c), suggesting the activity of RMg neurons
during movement depends upon context. Altogether, these observations indicate that the activity
of serotonergic neurons that send their projections ventral spinal motor circuits is distinct from 5-
HT populations targeting dorsal spinal cord and brain regions.

Locomotor state is a strong predictor of ROb/Pa activity

To determine whether locomotor state could be used to predict the activity of serotonin neurons,
we used a linear-non-linear model to examine the relationship between neural activity and
locomotor behavior[48,49] We generated linear filters for ROb/Pa, RMg, and DRN, which illustrate
the temporal relationship between activity and wheel speed (Fig. 3). This analysis was
performed using total wheel running data to assess whether the patterns of activity observed
during running onset and offset extend throughout running behavior. Indeed, the model
generated unique filters for each of the three serotonergic populations that were consistent with
the activity patterns we observed in the previous analysis (Fig.2).

The ROb/Pa filter contains a single, positive peak, indicating that increases in ROb/Pa *Pet1*
neuron activity occur during increases in wheel speed (Fig.3a). By contrast, the DRN filter
contains a negative peak (Fig.3e), indicating that increases in DRN *Pet1* neuron activity occur
during decreases in wheel speed. The position of each peak around 0 implies that changes in
ROb/Pa and DRN *Pet1* neuron activity are coincident to changes in speed, as we observed in
the aligned dF/F and wheel speed traces (Fig.2q,w). To assess the quality of these models, we
used each filter to predict neural activity from wheel speed and compared this to their actual
dF/F signals (Fig.3b,f) We find that wheel running is a good predictor of ROb/Pa *Pet1* neuron
activity (Fig.3b,CC=0.41,SEM=0.08) and DRN *Pet1* neuron activity (Fig.3f, CC=0.28,sem=0.02).

By comparison, the RMg filter is noisier and contains less pronounced structure (Fig.3c), likely
due to the varied signal we observed during wheel running (Fig.2h). We find using this model
that wheel speed is a poor predictor of RMg *Pet1* neuron activity (Fig.3d, CC=0.12,sem=0.05).
The RMg filter does however contain a small negative peak left of 0, suggesting that changes in
speed precede change in RMg *Pet1* neuron activity. This is consistent with the slower RMg
dynamics we observed during both wheel and treadmill locomotion (Fig.2s,t). Thus, in
comparison to other 5-HT populations, ROb/Pa exhibits a distinct pattern of activity that is highly
correlated to locomotor activity.

Brain-wide inputs to raphe *Pet1* neurons targeting ventral spinal cord

The tight coupling between ROb/Pa *Pet1* neuron activity and locomotion suggests that ROb/Pa
neurons may be part of a broader network involved in locomotor control. Descending command
pathways within the midbrain and brainstem are known to regulate the start, stop and speed of
locomotion[50-53]. The cuneiform nucleus (CnF), a component of the mesencephalic locomotor
region (MLR), is important for regulating locomotor initiation and speed[51,52]. The brainstem LPGi,
acts directly upon spinal locomotor networks to control motor output[50]. To determine whether
ROb/Pa neuron neurons receive input from these locomotor control areas, we performed
monosynaptic rabies tracing to identify brain regions that provide direct input to ventrally-
projecting serotonin neurons (Fig.4a,b).

To restrict rabies infection to *Tac1-Pet1* ROb/Pa neurons, we used an intersectional strategy by
generating a knock-in mouse line allowing for Cre- and Flpe-dependent expression of the TVA
receptor and rabies N2c glycoprotein (Extended Data Fig.6a,b). To label inputs to the *Tac1-Pet1*
subpopulation, we injected *Tac1-Cre:Pet1-Flpe:N2cG-TVA* mice with N2c rabies virus (EnvA-
N2cΔG-tdTomato) (Fig.4c). The rabies glycoprotein was tagged with HA, allowing for
identification of the initially infected starter population (HA⁺tdT⁺). We observe no HA expression
or RV infection in *N2cG-TVA* mice in the absence of Cre and Flpe (Extended Data Fig.6c),
indicating that rabies tracing depends on cell-type specific expression of N2cΔG and TVA.

Following rabies injection of *Tac1-Cre:Pet1-Flpe:N2cG-TVA* mice, we observed labeled tdT⁺
cells in several distinct brain areas. We identified dense bilateral clusters of tdT⁺ cells within
LPGi, IRt, PAG, and hypothalamus (Fig.4d,f,j,k,l). Additionally, we observed sparse labeling
within CnF (Fig.4e), the medullary reticular formation (MRF, including Gi, GiV, GiA, MdV), and
superior colliculus. We found comparable labeling of these regions after tracing from ROb/Pa
neurons in *Pet1-Cre* mice (Extended Data Fig.6d-g).

Several of the pre-ROb/Pa *Pet1* neuron populations, including CnF, LPGi, MRF and PAG, have
known roles in motor control[50,51,54,55]. The highest percentage of rabies-labeled cells (~18%)
were located in LPGi (Fig.4m). Recent work identified Vglut2+ neurons within the LPGi that
activate and are required for high-speed locomotion[50,56]. Therefore, we performed in situ
hybridization for *vglut2* mRNA and identified Vglut2-expressing RV-labeled cells in LPGi
(Fig.4g,h,i). These observations suggest a model whereby descending serotonergic pathways
are recruited in conjunction with locomotor command neurons to modulate spinal motor circuits
(Fig.4n).

Activation of ROb/Pa potentiates ongoing locomotor behavior

The activity of ROb/Pa *Pet1* neurons during wheel running and connectivity with multiple
locomotor control regions suggests that serotonergic input may be important for regulation of
locomotion. However, it is unknown whether spinal cord-projecting 5-HT neurons regulate
specific features of locomotion, such as initiation, duration, speed, and termination. To test this,
we measured the effect of optogenetically activating ROb/Pa neurons during wheel running. We
injected Cre-dependent channel rhodopsin (AAV-DIO-ChR-EYFP) into ROb/Pa of *Pet1-Cre*
mice (Fig.5a, Extended Data Fig.7a). Two weeks post injection mice were acclimated to the
wheel and fiber attachment. Three weeks post injection, mice were placed on running wheels
for 30 minutes, during which they received alternating 5-minute periods of light delivery (5 sec
20Hz pulses) and no light (Fig.5b). Control Cre-negative littermates were injected with ChR
virus and did not exhibit any ChR expression (Extended Data Fig.7c).

Because serotonin acts through both ionotropic and metabotropic receptors, changes in
locomotor parameters might emerge over both long and short times scales. We therefore
selected a pattern of blue light delivery that included both short 5-second pulses and longer 5-
minute periods with or without light pulses to enable us to observe how neuronal activation
influenced behavior over short and long-time scales. We did not observe locomotor initiation or
acute changes in speed in any animals upon blue light stimulation of ROb/Pa *Pet1* neurons
(Extended Data Fig.7d,f,g, Fig.5b). This is in stark contrast to locomotor-driving regions CnF
and LPGi, which when stimulated result in short latency activation of locomotion[50,51,56].

We next asked whether activation of ROb/Pa *Pet1* neurons influenced the amount of time
animals spent engaged in locomotion (wheel speed >5cm/sec). On average, mice spent

approximately 66% of their time moving on the wheel (~20min), for both control and ChR animals (Fig.5c). Thus, activation of ROb/Pa *Pet1* neurons did not affect the total duration of locomotion. We next sought to determine whether activation of serotonergic neurons influenced the number of discrete locomotor bouts mice produced. Serotonin increases the excitability of spinal MNs[11,25]. Thus, we predicted that activation of ROb/Pa may lead to an increase in run bouts, as heightened MN excitability could lower the threshold needed for a command region to initiate locomotion. Unexpectedly, we observed the opposite outcome, as ChR animals produced fewer locomotor bouts than control animals (Fig.5d,e). However, since ChR animals run the same total amount of time as control animals (Fig.5c), this observation suggests that ChR animals run bouts are longer. Indeed, we found that activation of ROb/Pa *Pet1* neurons significantly increased the length of locomotor bouts by the third light period (Fig.5f). Furthermore, ChR animals exhibited fewer total stops in locomotion (Fig. 5d) and an increase in the number of runs that were maintained following each light pulse (Extended Data Fig.7h,i), indicating that serotonergic input reduces the probability of terminating locomotor output. Together, these observations suggest that once an animal has initiated a run bout, ROb/Pa *Pet1* neuron activity increases the probability of maintaining the active locomotor state.

We next examined whether activation of ROb/Pa *Pet1* neurons influenced locomotor speed. We measured the fraction of time mice spent in defined 10cm/s speed bins and found that ChR mice spent less time engaged in locomotion at slower speeds (<25cm/s) and an increased time at faster speeds (>35cm/s) (Fig.5g). A histogram of time across wheel speed shows a rightward shift during the light periods as compared to no light periods (Fig5h,i, Extended Data Fig.7e), indicating that faster running by ChR mice occurs during light periods. To more precisely determine when light activation influenced speed, we measured average speed during each of the 5 min time bins. We found that average speed increased significantly in ChR animals compared to control animals during the third light period (Fig.5j, Extended Data Fig.7j,k). We also observed significant increases in the fraction of time spent running >35cm/s and the maximum speed during the third light period compared to the first light period (Fig.5k,l). Interestingly, the smaller increases in speed and duration of locomotion during the second light period are maintained during the subsequent no-light period (Fig.5j,k,l), suggesting that serotonin continues to influence motor circuits beyond periods of increased neuronal activity. The effects of light activation on locomotion therefore accumulate over time and are strongest during the final light activation period. These observations indicate that descending serotonergic modulation of spinal circuits can influence the speed of locomotion and may be important for regulating behavior over longer time scales.

Discussion

Neuromodulators are critical for the production of flexible and adaptive motor behaviors. While serotonin is a known modulator of spinal circuit dynamics[21], how the serotonergic system is integrated with motor circuits and how it influences movement has remained unknown. Our findings reveal that spinal cord-projecting serotonergic neurons are interconnected with motor circuits involved in the initiation and execution of locomotion. Serotonergic neurons targeting rhythm-generating spinal circuits receive input from locomotor control areas, display increased neuronal activity during locomotion, and can produce long-lasting potentiation of locomotor behavior. We discuss these findings in the context of neuromodulatory circuit architecture and the function of the serotonergic system in motor control.

Anatomical and molecular genetic neuronal tracing studies have demonstrated the existence of
distinct serotonergic populations targeting dorsal and ventral spinal cord[31-37]. Using genetic and
viral tracing assays, we provide a more complete picture of the input-output organization of the
descending serotonergic system. Our studies show that ventral spinal cord-projecting *Pet1*
neurons form a highly branched system that innervates multiple spinal segments, yet in a
spatially restricted manner along the dorsoventral domain. The distributed output of descending
ROb/Pa *Pet1* neurons likely coordinates modulation of multiple circuits to enable activation of
muscles across the body required for behavioral control, including the respiratory and
autonomic systems[36,37,57-59].

We found that ROb/Pa *Pet1* neurons receive input from locomotor control regions including
LPGi and CnF. This organization suggests that locomotor command circuits recruit the
descending serotonergic system to facilitate changes in excitability required to modulate motor
behaviors. Consistent with this model, previous work demonstrated that electrical stimulation of
the MLR, including CnF, causes release of 5-HT in the spinal cord[60]. In addition to locomotor-
related inputs, we find that ROb/Pa *Pet1* neurons receive input from additional regions that may
be essential in mediating context-dependent behaviors. For example, input from PAG, a region
known to be activated during fight or flight responses, may ensure spinal motor circuits are in an
appropriate state to produce the required motor response[54,61,62].

Like other neuromodulators, serotonin acts through volume transmission, but also through direct
synaptic contacts[43]. Our transsynaptic tracing studies suggest that spinal MNs receive synaptic
input from 5-HT neurons originating from ROb, and to a lesser extent the neighboring caudal 5-
HT nuclei. Interestingly, results from our retrograde transsynaptic tracing studies suggest that
excitatory ventral INs receive little direct 5-HT synaptic input, as few TPH2+ neurons were
rabies-labeled despite robust IN starter cell infection. This suggests possible distinct
mechanisms for serotonergic modulation of spinal INs and MNs. Additional layers of specificity
are likely to be conferred by differences in receptor expression among spinal neuronal
classes[63,64].

Raphe neurons exhibit activity related to levels of tonic motor activity[65]. During REM sleep when
there is reduced muscle tone the activity of raphe serotonergic neurons is strongly suppressed,
whereas their activity is elevated during waking states. Raphe populations differ however in their
activities during specific motor behaviors, such as locomotion. Single-unit recordings in cat
showed that the activity of ROb and RPa neurons increases during treadmill locomotion[41], while
activity within DRN neurons remains unchanged[47]. Here, we find that during spontaneous bouts
of locomotion, ROb and RPa neurons increase their activity, while DRN neurons decrease
activity. Our filter analysis of these data revealed that these relationships between neural activity
and locomotion are consistent throughout the spontaneous locomotor behavior we observed.

Anti-correlated activity has also been observed in DRN during locomotion within an open field,
yet DRN activity changes in situations where animals perceive a threat, becoming highly
correlated to muscle activity[46]. Our findings suggest that similar context-dependent changes in
activity occur within dorsal spinal cord projecting RMg neurons. During wheel running we
observed that RMg *Pet1* neuron activity was variable with a slight trend towards being
anticorrelated with movement. By contrast, during treadmill running RMg *Pet1* neuron activity
was strongly correlated with locomotor activity. The forced motorized treadmill assay is likely a
more stressful condition for mice compared to unrestrained wheel running. The differences in
RMg activity we observed therefore could reflect differences in the internal state of mice in these

conditions. RMg modulates incoming sensory information[66-68], and therefore may act as a
context-dependent gating system for sensory feedback. One could imagine a high-threat
scenario where high-speed locomotion must be prioritized to promote escape and survival. In
this context, it would be distracting or even life-threatening to respond to a pain signal from your
limb. In contrast, during non-threatening situations, it would be important to notice and respond
to a painful stimulus. In the future it will be important to examine the possible role of RMg in
context-dependent gating of sensory feedback and to gauge its influence on motor behavior.

Serotonin modulates the excitability of spinal MNs and INs, generating changes in the temporal
dynamics of motor output and magnitude and timing of muscle activation. It has been proposed
that serotonin mediates gain control, adjusting the input-output gain of MNs to achieve the
desired activation of muscles for specific movements[8,24]. We find that activation of ventral spinal
cord-projecting *Pet1* neurons increases the speed and length of running. Activation of
serotonergic neurons may function like turning up the gain "knob", where enhanced release of
5-HT increases the excitability of spinal MNs and INs. This could allow specific motor
commands to yield larger or more sustained motor output. Conversely, turning down this knob
would result in changes to locomotor behavior in the opposite direction, perhaps reducing
locomotor duration and speed. Consistent with this model, local delivery of a serotonin receptor
antagonist to the lumbar spinal cord results in impaired hindlimb stepping[7].

Finally, our studies reveal an unexpected feature of a neuromodulatory system's influence on
movement. The descending serotonergic pathway impacts locomotor behavior in a manner that
fundamentally differs from the relatively immediate effects that reticulospinal pathways have on
locomotor initiation, speed, and termination[50,52,55,56,69]. While the activity of ROb/Pa is tightly
synced with locomotor behavior, we find that the neurons do not act as a "go-signal" for
locomotion. Rather, serotonergic input strengthens and promotes maintenance of ongoing
locomotion. We find that activation of ROb/Pa produces increases in locomotor speed and
duration, and that these changes accumulate and are sustained over minutes. The influence
that the serotonergic pathway has on movement is slower and extends well-beyond the current
locomotor bout. These results are reminiscent of work in *C. elegans* showing that 5-HT
mediates a long-lasting effect on locomotor speed[5], suggesting that 5-HT mediates the effects
we have observed on locomotion. This enduring influence on behavior may be facilitated by
metabotropic 5-HT receptors that result in sustained changes in the excitability of spinal motor
circuits. These may be critical for producing sustained locomotor output to meet demands of
context-specific motor behaviors.

Acknowledgements

We thank Kathy Nagel, Michael Long, and Niels Ringstad for valuable discussions and
comments on the manuscript. Thank you to Arkarup Banerjee, Dayu Lin, Andrew Miri, Anders
Nelson, and Nic Tritsch for helpful discussions and technical advice throughout this work. We
thank members of the Lin, Long, Nagel, Schoppik, and Tritsch labs for technical support and
advice. We are grateful to Susan Morton for generating the TPH2 antibody, Sebastian Poliak for
generating the N2cG-TVA mice, Kim Ritola and Julia Sable for virus preparations, Sarah
Pfennig and Anne Cavanagh for technical support, and Bryan Chadwick for coding support.
Thank you to Nikos Balaskas, Joriene De Nooij, and Andy Murray for technical training and
support. Research was supported by the National Institute of Neurological Disorders and Stroke

Materials and Methods

Experimental animals.

Male and female mice aged 8-10 weeks at the time of surgical procedures were used for
experiments. All mice were maintained on a C57BL/6 genetic background. Prior to surgical
procedures, mice were housed communally with littermates. Following surgical procedures and
for the duration of behavioral analysis, mice were single housed. Mice were given *ad libitum*
access to food and water and maintained on a 12-hour light-dark cycle. All experimental and
surgical procedures involving animals were approved by the NYU Grossman School of Medicine
Institutional Animals Care and Use Committee (IACUC) and in compliance with the NIH
guidelines for care and use of animals.

A listing of all mouse strains using in experiments can be found below in Table 1. To perform
monosynaptic retrograde rabies tracing from genetically-defined subpopulations of serotonin
neurons (Fig. 4), we generated two mouse lines to enable recombinase-dependent expression
of the N2c rabies virus glycoprotein (N2cG), and the receptor for the subgroup A avian sarcoma
and leukosis virus (TVA). An HA-tagged N2cG and mutated TVA66T[70,71] were inserted into both
the Ai9 (Addgene plasmid #22799) and Ai65 (Addgene plasmid #61577) targeting vector to
enable either Cre-dependent or Cre- and Flpe-dependent expression of N2cG and TVA,
respectively. Both targeting vectors allow expression of the inserted cassette at the mouse
Rosa26 locus. The targeting vectors were used to generate mice at Kallyope and then
transferred to NYU Langone animal facility.

Table 1. Mouse strains by experiment

Experiment	Mouse Strain	Source
Anterograde labeling of Tac1-Pet1 or Egr2-Pet1 synaptic terminals (Fig. 1)	RC::FPSit (Cre- and Flpe-dependent synaptophysin-EGFP) Pet1-Flpe Tac1-Cre Egr2-Cre	S. Dymecki (Harvard) [36,37,42] JAX 030206
Monosynaptic rabies virus tracing from spinal motor neurons (Fig. 1)	ChAT-Cre	T. Jessell (Columbia) [72] Jax 006410
Monosynaptic rabies virus tracing from Chx10$^+$ interneurons (Ext Data Fig. 1)	Chx10-Cre	T. Jessell (Columbia) [73]
Optogenetics and photometry (Fig. 2&5)	Pet1-Cre	JAX 012712
Monosynaptic rabies tracing from raphe (Fig. 4, Ext Data Fig. 6)	Ai65-N2cG-TVA (Cre- and Flpe-dependent) Ai9-N2cG-TVA (Cre-dependent)	Generated with Sebastian Poliak (Kallyope)

Viral injections.

Anesthesia was induced using vaporized isoflurane at 3% in oxygen (2 L/min). Anesthesia was maintained throughout the duration of procedures at 1.5-2.5% in oxygen (1 L/min) with mice held in stereotaxic frame (Kopf Instruments, Model 940) atop a feedback-controlled heating pad set to 37 °C. Viruses were injected using a Nanoject II (Drummond Scientific Company) and pulled glass capillaries.

To perform rabies tracing from spinal MNs, AAV9-Syn-DIO-TVA66T-tdT-N2cG (Julie Sable, Jessell lab, Columbia) was injected into the lateral ventricle by intracerebroventricular (ICV) injection at P0-1. The ICV injection allows virus to spread with the cerebrospinal fluid and enables infection of spinal MNs[45]. At eight weeks of age, mice were injected with EnvA-N2c(ΔG)-TdT (Janelia, NeuroTools Viral Vector Core) into spinal segments C4 to T1. To perform rabies tracing from Ch10+ INs, Chx10-Cre mice were injected with AAV1-flex-TVA-N2cG (Julie Sable, Jessell lab, Columbia) into spinal segments C5 to C8, and three weeks later EnvA-N2c(ΔG)-TdT (Julie Sable, Jessell lab, Columbia) was injected into segments C5 to C8. Mice were perfused 1-2 weeks post rabies injection. To retrogradely label serotonergic neurons from the spinal cord (Extended Data Fig.3), AAV2retro-CAG-FLEX-tdTomato (Addgene 28306-AAVrg) was injected into spinal segments C5 to C8 of Pet1-cre mice (8 weeks). Mice were perfused 2 weeks post-injection. For rabies tracing in *N2cG-TVA* mice, EnvA-N2c(ΔG)-TdT (Janelia, NeuroTools Viral Vector Core) was injected using ROb coordinates described below and mice were perfused 7-8 days post-injection.

Coordinates for ROb injections were 6.85mm posterior and 0mm mediolateral of bregma with injection at depth of 5.4 to 5.7mm below bregma. Coordinates for RMg injections were 5.3mm posterior and 0mm mediolateral of bregma with injection at depth of 5.7mm. Coordinates for DRN injections were 4.7mm posterior and 0mm mediolateral of bregma with injection at depth of 3mm. Cell-type specific expression of Gcamp6s and channelrhodopsin was achieved using the following Cre-dependent viruses: AAV5-FLEX-GcAMP6s (Addgene #100842-AAV5) for ROb/Pa and RMg, AAV1-FLEX-GcAMP6s (Addgene #100842-AAV1) for DRN, and AAV2-EF1a-DIO-ChR2(H134R)-EYFP (UNC Vector Core). Approximately 300 nL of virus was injected per animal. Following the delivery of viral vector, a 26-gauge guide cannula (P1 Technologies) was positioned overtop of the injection site, descended to a depth of .2mm above desired fiber position and removed. A fiberoptic cannula (400μm core diameter, Doric Lenses) was then lower into the brain and cemented in place (C&B Metabond, Parkell). Lastly, a headplate was cemented to the skull to facilitate handling of mice while attaching and removing the fiber optic cable.

Histology and imaging.

Mice were euthanized via isoflurane overdose delivered by the open drop method and transcardially perfused with 10mL of cold 1X phosphate buffered saline (PBS) followed by 10mL of ice cold 4% paraformaldehyde (PFA). Brain and/or spinal cords were removed and post-fixed overnight in 4% PFA at 4°C. Tissue was sectioned free floating in cold 1X PBS using a vibratome (Leica VT1200S). Brain tissue was sectioned at 70 μm thickness and spinal cord tissue was sectioned at 80 μm thickness.

For immunohistochemistry, sections were permeabilized in 0.3% TritonX-100 in 1X PBS for 15 minutes at room temperature. Following permeabilization, sections were incubated free floating in primary antibodies (Table 2) diluted in a blocking solution of 1% bovine serum albumin (BSA) and 0.3% TritonX-100 in 1X PBS for 72 hours at 4°C. Secondary antibodies (Table 3) diluted in

447 blocking solution were applied overnight at 4°C. Sections were mounted onto superfrost plus
448 microscope slides (Fisherbrand) with Fluoromount-G (SouthernBiotech).
449
450 TPH2 antibody was generated by Covance using a TPH2 peptide (RRGLSLDSAVPEDHQLC,
451 Atlantic Peptides) coupled to KLH (Thermofisher 77605). Reagents were designed and
452 prepared by Susan Brenner-Morton at Columbia University.
453
454
455 Table 2: Primary Antibodies

Ab	Target	Host Species	Dilution	Manufacturer
Gp ∝ TPH2	tryptophan hydroxylase 2	Guinea Pig	1:4000	Covance
Ch ∝ GFP	Green Fluorescent Protein	Chicken	1:2000	ThermoFisher (A10262)
Rat ∝ 5-HT	5-hydroxytryptamine	Rat	1:1000	Enzo (BML-SZ1011-0025)
Rb ∝ DsRed	DsRed	Rabbit	1:1000	Takara (632496)
Rb ∝ HA	Hemagglutinin (HA) tag	Rabbit	1:1000	Abcam (ab9110)
Gt ∝ ChAT	choline acetyltransferase	Goat	1:200	Millipore (AB144P)

457 Table 3: Secondary Antibodies

Ab	Conjugated fluorophore	Dilution	Manufacturer & Cat #
Donkey ∝ Guinea Pig - Cy3	Cyanine 3	1:1000	Jackson (706-165-148)
Donkey ∝ Guinea Pig - 647	Alexa Fluor 647	1:1000	Jackson (706-605-148)
Donkey ∝ Chicken - 488	Alexa Fluor 488	1:1000	Jackson (703-545-155)
Donkey ∝ Rat - Cy3	Cyanine 3	1:1000	Jackson (712-165-150)
Donkey ∝ Rat - Cy5	Cyanine 5	1:1000	Jackson (712-175-150)
Donkey ∝ Rabbit - Cy3	Cyanine 3	1:1000	Jackson (711-165-152)
Donkey ∝ Rabbit - 488	Alexa Fluor 488	1:1000	Jackson (711-545-152)
Donkey ∝ Goat - 647	Alexa Fluor 647	1:1000	Jackson (705-605-147)

Image analysis.

Imaris (Bitplane) software was used to analyze the position of synaptophysinGFP puncta in spinal cord sections and MATLAB was used to generate the position plots and density plots for each population, as previously described in Bikoff et al.[74]. This approach was also used to generate the position plots of RV+ and TPH+ cells following rabies tracing from spinal MNs and Chx10+ INs.

Fluorescent *In Situ* hybridization.

In situ hybridization was performed on brain tissue following the injection of CVS-N2c(ΔG)-tdT rabies virus to identify retrogradely labeled cells expressing the vesicular glutamate transporter 2 (Vglut2). Hybridization chain reaction (HCR) probes for Vglut2 were generated from the sense sequence of Vglut2 cDNA (NCBI) using the HCR 3.0 probe maker[75,76]. Following tissue collection and post-fixation described above, brains were placed in a 30% sucrose solution overnight at 4 °C, then embedded in Tissue-Tek O.C.T. compound. Tissue was cryosectioned at 18μm thickness, collected onto superfrost plus microscope slides, and stored at -80 °C. Fluorescent in situ hybridization was performed using the Hybridization Chain Reaction RNA fluorescent in situ hybridization (HCR RNA-FISH, Molecular Instruments) protocol, described in D'Elia et al.[77].

Locomotor assays and acclimation.

Wheel running experiments were performed using a low-profile running wheel (Fast Trac K3251, Bio-Serve) attached to rotary encoder (A2 optical shaft encoder, US Digital). Encoder output was collected using a USB interface board (RHD2000, #C3100, Intan Technologies) and recorded using RHD2000 Interface software (Intan Technologies). Treadmill experiments were performed using a custom-built motorized rodent treadmill (Model 802, University of Cologne electronics lab).

Mice were acclimated to the running wheel and treadmill for 4 days prior to the first behavioral data collection trials. On the first day of behavioral acclimation, mice were allowed explore the wheel and treadmill environments for 10 minutes each. On the subsequent 3 days, mice were acclimated with the fiber optic cable attached. Each of these 3 days, mice were then allowed to run freely on the wheel for 10-30 minutes. On the second day of behavioral acclimation, mice ran on the treadmill for 1 minute at 20 cm/sec and 1 minute at 25 cm/sec with a 1-minute rest period in between. The treadmill speed was ramped up slowly by hand to acclimate the animal to the moving belt. On the third and fourth days of behavioral acclimation, mice on the treadmill performed alternating 10 second intervals of rests and runs: 3X at 20 cm/sec, 3X at 25 cm/sec, 3X at 30 cm/sec, and 3X at 35 cm/sec.

Fiber photometry.

A rig for collecting fiber photometry data was constructed using LEDs (M470F3 & M405FP1, Thorlabs), Fluorescent Mini Cube (FMC Gen1, Doric), and photoreceiver (Model 2151, Newport). A 385 Hz sinusoidal 470 LED light (30μW, bandpass filtered 460-490nm) was delivered by a 400μm optic cable to excite Gcamp6s in the brain. A 315 Hz sinusoidal 405 LED light (30μW, bandpass filtered 400-410nm) was also delivered to the brain as a control for motion artifact. The isosbestic point for Gcamp6s is near 405nm, where emitted light is independent of calcium binding. The emitted light from the brain was bandpass filtered (500-

550nm) and collected by photodetector and recorded using the Intan USB interface board with Intan software described above with locomotor assays.

Recordings were made during 30min wheel sessions and treadmill runs over 14-21 days and collected at 4kHz sampling rate. The signals were bandpass filtered (470nm passing band: 385Hz ± 10, 405 passing band: 315Hz ± 10 and demodulated using a phase sensitive detection method comparing the modulated signal with a recorded reference signal of same frequency. Demodulation extracts the envelope of the 385Hz signal reflecting the intensity of Gcamp6s signal. A baseline was calculated from demodulated signal using an interpolated linear fit of values in 10th percentile and moving window size of 30s. The photometry signal was baseline adjusted: dF/F= (demodulated signal-baseline)/baseline.

The averaged dF/F at the start and stop of run bouts was generated using runs of at least 3 seconds in length, defined by periods where wheel speed was greater than 10cm/sec. For this analysis, each run was baselined individually. Run starts were normalized to the average fluorescence 0.5s to 1s prior to the run start and run stops to the fluorescence 0.5s to 1s post run stop.

Optogenetic activation and analysis.

For optical stimulation of ChR2, 470nm LED light (M470F3, Thorlabs) was delivered for 5s pulses (25Hz sinusoid). Light intensity was measured using an optical power meter (PM100D, Thorlabs) at the tip of the fiber canula to be 13-15 mW. During each of the 5-minute light periods, mice received 15 5-second light pulses.

Mice were attached to optic fiber and allowed to run freely on wheel for 30min during the light stimulation protocol with 5-minute alternating no light and light periods. Data was collected from each ChR and control mice once a day with 1-2 rest days in between for 3 weeks. For data to be included in analysis, the following criteria were required: 1) at least 5 minutes total of speeds >5cm/sec and 2) reached speeds greater than 10cm/sec during all 3 no light and light periods. These criteria were used to ensure comparable behavior when animals ran to a similar extent throughout the 30 minutes and remove days where mice stopped running for long periods of time. For wheel speed analysis, wheel rotary encoder output was converted to speed using the wheel circumference and maximum encoder output value and downsampled to 1kHz. Run bouts were defined as periods of wheel speed greater than 10cm/sec for at least 1 second.

Electromyographic recordings and analysis.

To observe the relationship between 5-HT neuron activity and muscle activity during locomotor behavior, electromyographic (EMG) electrodes were implanted into limb muscle to record muscle activity. EMG electrodes were fabricated and implanted as previously described[78]. Briefly, EMG electrodes were made using insulated steel wire (793200, A-M Systems) with 1mm exposed regions 7.5cm away from a 12-pin miniature connector (11P3828, Newark). The wire ends were inserted and crimped inside of a 27-gauge needle to use when inserting electrodes into the muscle. EMG electrodes were implanted into mice that had previously been injected with Gcamp6s virus and had headplates attached. An incision was made above the neck, hip and tibialis anterior (TA) muscle in right hindlimb. The needle end of the wire was guided beneath the skin from the neck to the hip and then to TA. The needle was used to insert the wire through the muscle and a knot made at the end to hold the electrode in place. The

connecter was cemented to the rear edge of the headplate. EMG signals were transmitted from the headplate connecter by an Omnetics connector and amplified using an amplifier chip (RHD2216, Intan Technologies). Amplified EMG signals were recorded using the Intan USB interface board and Intan software. EMG recordings were downsampled to 1kHz, high-pass filtered at 40Hz, and rectified, as described in Miri et al.[78].

 Figure Legends

Figure 1. Spinal motor circuits receive biased 5-HT input form medullary raphe nuclei

a-e. 5-HT immunostaining in adult spinal cord at cervical (C), thoracic (T), lumbar (L) levels. **d**.
ChAT$^+$ motor neurons (MN) with 5-HT immunostaining. Zoom-in of white box in a. **e**. Genetically
labeled Chx10$^+$ ventral interneuron (IN) with 5-HT immunostaining. **f-i**. Genetic labeling of *Egr2-*
Pet1 neurons with synaptophysinGFP (cyan) in cervical spinal cord. **g**. Immunostaining for ChAT
(magenta) to visualize MN. **h**. Distribution of *Egr2-Pet1* puncta (red) and total 5-HT puncta
(black). **i**. Relative density plot of *Egr2-Pet1* puncta. **j-m**. Genetic labeling of *Tac1-Pet1* neurons
with synaptophysinGFP in cervical spinal cord. **k**. Immunostaining for ChAT to visualize MN. **l**.
Distribution of *Tac1-Pet1* puncta (red) and total 5-HT puncta (black). **m**. Relative density plot of
Tac1-Pet1 puncta. **n**. Schematic summary of *Tac1* vs. *Egr2 Pet1* neuron populations and target
innervation of spinal cord. **o-r**. Monosynaptic retrograde rabies tracing to identify 5-HT input to
spinal MNs. **o**. Experimental procedure. EnvA-RVΔG-GFP injection into cervical spinal cord to
infect spinal MN expressing TVA and G. **p**. Cervical spinal cord MNs infected by AAV and RV.
Zoom-in of white-dotted region are below. **q**. Retrogradely labeled neurons expressing
tryptophan hydroxylase 2 (TPH2) within ROb and RPa. **r**. Quantification of RV-labeled TPH2$^+$
neurons. Percentage of total RV$^+$/TPH2$^+$ cells within raphe and LPGi. n=3, each red dot is
average of a single animal.

Figure 2. Activity of ventrally-projecting serotonin neurons increases during locomotion

a. Behavioral assay. Mice freely running on low-profile running wheel with attached fiber for
photometry recordings. **b**. Example photometry traces dF/F (470nm in pink, 405nm in grey) from
ROb/Pa *Pet1* neurons with wheel speed (black) during single session. **c**. Injection of AAV-FLEX-
Gcamp6s into ventral medulla of *Pet1-Cre* mice to infect ROb/Pa *Pet1* neurons. Histology
showing Gcamp6s expression and TPH2 immunostaining in ROb/Pa. **d,h,l**. Example trace
during 60sec of free running with dF/F and wheel speed for ROb/Pa *Pet1* neurons (**d**), RMg
Pet1 neurons (**h**), DRN *Pet1* neurons (**l**). **e,i,m**. Averaged dF/F and wheel speed aligned to start
and stop of running. **f,j,n**. Plot of the average dF/F during the rest (prior to run start) on the x-
axis and during the run on the y-axis. Each dot is an individual run. Triangles show average
dF/F of all runs at 470nm (yellow) and 405nm (black). **o,r,u**. Example photometry traces during
treadmill assay showing 470nm (pink) and 405nm (grey) dF/F with treadmill setting in black.
ROb/Pa *Pet1* neurons (**o**), RMg *Pet1* neurons (**r**), DRN *Pet1* neurons (**u**). **p,s,v**, Averaged
signal across all treadmill run bouts with 10sec run and 10sec rest. ROb/Pa (**p**), RMg (**s**), DRN
(**v**). **q,t,w**, Average 470nm dF/F (pink) and wheel speed (black) at run start on wheel. ROb/Pa
(**q**), RMg (**t**), DRN (**w**).

Figure 3. Locomotor state is a strong predictor of ROb/Pa activity

a,c,e. Linear filters for ROb/Pa *Pet1* neurons (**a**), RMg *Pet1* neurons (**c**), DRN *Pet1* neurons (**e**).
Black line is average (n=3 mice) with grey SEM. **b,d,f**, Actual dF/F trace (pink) overlayed with
model's predicted activity (grey) and wheel speed (black) for ROb/Pa (**b**), RMg (**d**), DRN (**f**).

Figure 4. Brain-wide inputs to raphe *Pet1* neurons targeting ventral spinal cord

a. Identifying inputs to the ventral spinal cord-projecting 5-HT pathway. **b**. Experimental procedure. Injection of EnvA-RVΔG-tdTomato into ROb/Pa of *Tac1-Cre:Pet1-Flpe:N2cG-TVA* mice to identify monosynaptic inputs to *Tac1-Pet1* neurons. **c**. Primary infection of *Tac1-Pet1* neurons in ROb/Pa with EnvA-RVΔG-tdTomato. **d-l**. Representative images of EnvA-RVΔG-tdTomato (tdT) infected neurons LPGi (**d**, zoom-in white box in **f**), cuneiform (CnF, **e**), intermediate reticular nucleus (IRt, **j**), periaqueductal grey (PAG, **k**), hypothalamus (Hyp, **l**). **g-i**. *Vglut2* mRNA expression with EnvA-RVΔG-tdTomato infection in LPGi neurons. **m**. Quantification of rabies-infected neurons. Percentage of total rabies-labeled cells (n=2). **n**. Model suggested by rabies tracing. Locomotor command neurons within MLR (CnF) and LPGi send projections (cyan arrows) to brainstem 5-HT neurons that target ventral spinal cord to facilitate modulation of spinal MNs and INs during locomotor behavior. (lateral paragigantocellularis, LPGi; intermediate reticular nucleus, IRt; periaqueductal grey, PAG; hypothalamus, Hyp; medullary reticular formation, MRF; cuneiform, CnF; superior colliculus, SC; raphe pallidus, RPa; raphe obscurus, ROb; raphe magnus, RMg, mesencephalic locomotor region, MLR)

Figure 5. Activation of ROb/Pa potentiates ongoing locomotor behavior

a-b. Experimental procedure. **a**. Injection of AAV2-DIO-ChR2(H134R)-EYFP into ROb/Pa of *Pet1-Cre* mice. ChR-expressing TPH2⁺ neurons in ROb/Pa with ChR⁺ axon terminals in lumbar spinal cord. **b**. Light delivery protocol during 30min of wheel running with single example wheel speed trace from one ChR animal (orange). Stimulation during 5-minute light periods: 5s 20Hz pulses 470nm light repeating every 15s. **c**. Total time at rest vs. locomoting in control (grey) and ChR (black) animals. n=22-30 trials from 5-6 animals. **d**. Total number of locomotor bouts for ChR and control mice (equal to total number of run starts and run stops) n=22-30 trials from 5-6 animals. **p<0.01 unpaired t-test. **e**. Number of locomotor bouts during each 5-minute light or no light period for control and ChR animals. n=22-30 trials from 5-6 animals. *p<0.05 unpaired t-test. ns=not significant. **f**. Average length of locomotor bouts during each 5-minute light or no light period for control and ChR animals. n=22-30 trials from 5-6 animals. **p<0.01 unpaired t-test. **g**. Fraction of total time control or ChR animals spend within various locomotor speed intervals. n=22-30 trials from 5-6 animals. **p<0.01, ***p<0.001 Bonferroni-corrected t-test. **h**. Distribution of time (ms) across wheel speed during No-light (grey) or Light (blue) periods. Combined data from 5-6 animals across all No-light or Light periods. **i**. Difference in average speed between Light and No-light periods. n=5-6 animals. *p<0.05 unpaired t-test. **j**. Average speed during each 5-minute light or no light period for control and ChR animals. n=22-30 trials from 5-6 animals. *p<0.05 unpaired t-test. **k**. Fraction of time mice spent locomoting greater than speed of 35cm/s during each 5-minute light or no light period for control and ChR animals. n=22-30 trials from 5-6 animals. **p<0.01 unpaired t-test. ns=not significant. **l**. Average maximum speed during each 5-minute light or no light period for control and ChR animals. n=22-30 trials from 5-6 animals. *p<0.05 unpaired t-test. ns=not significant. All Error bars are SEM.

Extended Data Figure 1. Genetic labeling of 5-HT sub-populations

a. Intersectional synaptophysin-GFP allele [37,42]. **b**. Breeding scheme to generate *Egr2-Pet1-synGFP* or *Tac1-Pet1-synGFP* mice. **c**. Distribution of synaptophysinGFP (red) and 5-HT immunostaining (black) puncta in thoracic and lumbar spinal segments for *Egr2-Pet1* and *Tac1-Pet1* neuron populations. Relative density of synGFP puncta (blue). **d**. Distribution of *Tac1-Pet1* and *Egr2-Pet1* neurons in brainstem and midbrain with TPH2 immunostaining. Raphe pallidus, RPa; raphe obscurus, ROb; raphe magnus, RMg; dorsal raphe nucleus, DRN; median raphe nucleus, MRN.

Extended Data Figure 2. Monosynaptic rabies tracing from spinal MNs and Chx10$^+$ INs

a. Rabies virus (RV) labeling in brainstem and midbrain with TPH2 immunostaining showing serotonergic nuclei. **b**. Summary of RV$^+$TPH$^+$ neurons in single *ChAT-Cre* animal with total TPH$^+$ cells. **c**. Experimental strategy for tracing monosynaptic inputs to Chx10$^+$ INs. Injection of *Chx10-cre* mouse cervical spinal cord with AAV-FLEX-TVA-HA-G followed by EnvA-RVΔG-GFP. **d**. Infection of Chx10$^+$ neurons in cervical spinal cord. Zoom-in below showing RV-infected HA$^+$ neurons and 5-HT puncta. **e**. RV labeling within brainstem reticular formation. **f**. Summary of RV$^+$TPH$^+$ neurons in single *Chx10-cre* animal with total TPH$^+$ cells. (Raphe pallidus, RPa; raphe obscurus, ROb; lateral paragigantocellularis, LPGi; raphe magnus, RMg; dorsal raphe nucleus, DRN; median raphe nucleus, MRN.)

Extended Data Figure 3. Retrograde labeling of spinal-projecting *Pet1* neurons

a. Experimental procedure. Injection of AAV2r-FLEX-tdTomato into C5-C8 of *Pet1-Cre* mice. **b**. Expression of tdTomato in brainstem with TPH2 expression. tdTomato$^+$ fibers at nucleus ambiguous (NA). **c**. tdTomato expression within fibers of thoracic (Th) and lumbar (L) spinal cord with ChAT immunostaining. Right most images are zoom-in of black box showing tdTomato$^+$ fibers around MNs. **d**. Schematic depicting highly collateralized ROb/Pa *Pet1* neurons targeting multiple spinal levels and caudal medulla.

Extended Data Figure 4. Neural activity during individual runs on wheel

a-c. Heatmap showing dF/F (left) for all individual runs during wheel assay. *Pet1* neuron activity within ROb/Pa, RMg, or DRN. Each horizontal line is a single run. Runs ordered by length from shortest to longest. Traces include 2 seconds prior to run start. Heatmap showing corresponding wheel speed (right) for all runs. **a**. ROb/Pa (954 runs, 3 animals). **b**. RMg (991 runs, 3 animals). **c**. DRN (916 runs, 3 animals).

Extended Data Figure 5. Additional photometry data and histology quantification

a,c,e. Averaged 470nm (pink) and 405nm (grey) dF/F during treadmill locomotor assay with treadmill setting (black). **a**. ROb/Pa (20 trials from 2 animals). **c**. RMg (10 trials from 2 animals). **e**. DRN (12 trials from 2 animals). **b,d,f**. Post-hoc quantification of histology for animals used in

689 fiber photometry experiments. Gcamp6s$^+$TPH2$^+$ cell counts and fiber position for each animal. **b**.
690 ROb/Pa. **d**. RMg. **f**. DRN. **g**. EMG recordings from tibialis anterior (TA) with Gcamp imaging
691 during wheel running. Raw EMG trace from TA during locomotion (black). Average dF/F (pink)
692 from ROb/Pa overlaying rectified EMG traces aligned to first peak of muscle activity at start of
693 run bout (multicolor). Arrow shows alignment of first muscle burst.

Extended Data Figure 6. Retrograde rabies tracing from raphe neurons targeting ventral spinal cord.

a. Intersectional *N2cG-TVA* allele. **b**. Generation of *Tac1:Pet1:N2cG-TVA* mice. Expression of
HA-tagged N2cG within ROb/Pa and LPGi neurons. **c**. HA and TPH2 immunostaining in *N2cG-*
TVA mouse (no Cre/Flpe) injected with EnvA-RVΔG-tdTomato. **d**. Cre-dependent *N2cG-TVA*
allele. **e**. Generation of *Pet1:N2cG-TVA* mice. Expression of HA-tagged N2cG within ROb/Pa,
RMg, LPGi neurons. **f**. HA and TPH2 immunostaining in *N2cG-TVA* mouse (no Cre) injected
with EnvA-RVΔG-tdTomato. **g**. Rabies infection in *Pet1:N2cG-TVA* caudal brainstem with
labeled presynaptic cells in LPGi, cuneiform (CnF), periaqueductal grey (PAG), and
hypothalamus (Hyp).

Extended Data Figure 7. Optogenetic activation of ROb/Pa *Pet1* neurons during locomotor behavior

a. ChR-infected cells and fiber position for each ChR animal. **b**. ChR expression in lumbar
spinal cord with ChAT immunostaining. **c**. Representative control animal with no ChR
expression. White dotted line denotes position of fiber. **d**. Wheel speed for trials when animal
was still at time of light onset. Averaged wheel speed from the first light pulse of each light
period for control and ChR animals when wheel was 0cm/s at time of light onset. **e**. Distribution
of time (ms) across wheel speeds during each No-light (grey) or Light (blue) periods (1-3).
Combined data from 5-6 animals. **f**. Wheel speed when animal was running at time of light
onset. Averaged wheel speed during first light pulse of each period when wheel speed was
greater than 20cm/s at light onset. **g**. Data in f displayed by control or ChR animals with light
periods overlayed. **h**. Individual wheel traces from light period 3 in f. **i**. Fraction of trials when
animal is running at time of light onset and where running is maintained for 10 seconds following
light onset. n=113-159 runs, from 5-6 animals. Error bars are SEM. *p<0.05 unpaired t-test. **j**.
Averaged wheel speed across all light pulses of each period when wheel speed was greater
than 20cm/s at light onset. **k**. Data in j displayed by control or ChR animals with overlayed light
periods.

References

1. Katz, P.S. (1995). Intrinsic and extrinsic neuromodulation of motor circuits. Curr Opin Neurobiol *5*, 799-808. 10.1016/0959-4388(95)80109-x.
2. Marder, E. (2012). Neuromodulation of neuronal circuits: back to the future. Neuron *76*, 1-11. 10.1016/j.neuron.2012.09.010.
3. Marder, E., and Bucher, D. (2001). Central pattern generators and the control of rhythmic movements. Curr Biol *11*, R986-996. 10.1016/s0960-9822(01)00581-4.
4. Harris-Warrick, R.M. (2011). Neuromodulation and flexibility in Central Pattern Generator networks. Curr Opin Neurobiol *21*, 685-692. 10.1016/j.conb.2011.05.011.
5. Huang, Y.C., Luo, J., Huang, W., Baker, C.M., Gomes, M.A., Meng, B., Byrne, A.B., and Flavell, S.W. (2023). A single neuron in C. elegans orchestrates multiple motor outputs through parallel modes of transmission. Curr Biol *33*, 4430-4445 e4436. 10.1016/j.cub.2023.08.088.
6. Howard, C.E., Chen, C.-L., Tabachnik, T., Hormigo, R., Ramdya, P., and Mann, R.S. (2019). Serotonergic Modulation of Walking in Drosophila. Current Biology *29*, 4218-- 4230.e4218. 10.1016/j.cub.2019.10.042 , pmid = 31786064 , pmcid = PMC6935052.
7. Cabaj, A.M., Majczynski, H., Couto, E., Gardiner, P.F., Stecina, K., Slawinska, U., and Jordan, L.M. (2017). Serotonin controls initiation of locomotion and afferent modulation of coordination via 5-HT7 receptors in adult rats. J Physiol *595*, 301--320. 10.1113/jp272271 , pmid = 27393215.
8. Wei, K., Glaser, J.I., Deng, L., Thompson, C.K., Stevenson, I.H., Wang, Q., Hornby, T.G., Heckman, C.J., and Kording, K.P. (2014). Serotonin Affects Movement Gain Control in the Spinal Cord. The Journal of Neuroscience *34*, 12690--12700. 10.1523/jneurosci.1855-14.2014 , pmid = 25232107 , pmcid = PMC4166156.
9. Lillvis, J.L., and Katz, P.S. (2013). Parallel evolution of serotonergic neuromodulation underlies independent evolution of rhythmic motor behavior. J Neurosci *33*, 2709-2717. 10.1523/JNEUROSCI.4196-12.2013.
10. Dunbar, M.J., Tran, M.A., and Whelan, P.J. (2010). Endogenous extracellular serotonin modulates the spinal locomotor network of the neonatal mouse. J Physiol *588*, 139-156. 10.1113/jphysiol.2009.177378.
11. Perrier, J.F., and Delgado-Lezama, R. (2005). Synaptic release of serotonin induced by stimulation of the raphe nucleus promotes plateau potentials in spinal motoneurons of the adult turtle. J Neurosci *25*, 7993--7999. 10.1523/jneurosci.1957-05.2005 , pmid = 16135756.
12. Clemens, S., and Katz, P.S. (2001). Identified serotonergic neurons in the Tritonia swim CPG activate both ionotropic and metabotropic receptors. J Neurophysiol *85*, 476-479. 10.1152/jn.2001.85.1.476.
13. Zhang, W., and Grillner, S. (2000). The spinal 5-HT system contributes to the generation of fictive locomotion in lamprey. Brain Res *879*, 188-192. 10.1016/s0006-8993(00)02747-5.
14. Kiehn, O., and Kjaerulff, O. (1996). Spatiotemporal characteristics of 5-HT and dopamine-induced rhythmic hindlimb activity in the in vitro neonatal rat. J Neurophysiol *75*, 1472-1482. 10.1152/jn.1996.75.4.1472.
15. Barbeau, H., and Rossignol, S. (1991). Initiation and modulation of the locomotor pattern in the adult chronic spinal cat by noradrenergic, serotonergic and dopaminergic drugs. Brain Res *546*, 250--260. 10.1016/0006-8993(91)91489-n , pmid = 2070262.
16. Katz, P.S., and Harris-Warrick, R.M. (1990). Neuromodulation of the crab pyloric central pattern generator by serotonergic/cholinergic proprioceptive afferents. J Neurosci *10*, 1495-1512. 10.1523/JNEUROSCI.10-05-01495.1990.

17. Wallen, P., Christenson, J., Brodin, L., Hill, R., Lansner, A., and Grillner, S. (1989). Mechanisms underlying the serotonergic modulation of the spinal circuitry for locomotion in lamprey. Prog Brain Res *80*, 321-327; discussion 315-329. 10.1016/s0079-6123(08)62227-x.

18. Hounsgaard, J., and Kiehn, O. (1989). Serotonin-induced bistability of turtle motoneurones caused by a nifedipine-sensitive calcium plateau potential. J Physiol *414*, 265-282. 10.1113/jphysiol.1989.sp017687.

19. Beltz, B., Eisen, J.S., Flamm, R., Harris-Warrick, R.M., Hooper, S.L., and Marder, E. (1984). Serotonergic innervation and modulation of the stomatogastric ganglion of three decapod crustaceans (Panulirus interruptus, Homarus americanus and Cancer irroratus). J Exp Biol *109*, 35-54. 10.1242/jeb.109.1.35.

20. Livingstone, M.S., Harris-Warrick, R.M., and Kravitz, E.A. (1980). Serotonin and octopamine produce opposite postures in lobsters. Science *208*, 76-79. 10.1126/science.208.4439.76.

21. Perrier, J.F., and Cotel, F. (2015). Serotonergic modulation of spinal motor control. Curr Opin Neurobiol *33*, 1--7. 10.1016/j.conb.2014.12.008 , pmid = 25553359.

22. Cazalets, J.R., Sqalli-Houssaini, Y., and Clarac, F. (1992). Activation of the central pattern generators for locomotion by serotonin and excitatory amino acids in neonatal rat. J Physiol *455*, 187-204. 10.1113/jphysiol.1992.sp019296.

23. Barbeau, H., and Rossignol, S. (1990). The effects of serotonergic drugs on the locomotor pattern and on cutaneous reflexes of the adult chronic spinal cat. Brain Research *514*, 55--67. 10.1016/0006-8993(90)90435-e , pmid = 2357531.

24. Johnson, M.D., and Heckman, C.J. (2014). Gain control mechanisms in spinal motoneurons. Front Neural Circuits *8*, 81. 10.3389/fncir.2014.00081 , pmid = 25120435 , pmcid = PMC4114207.

25. Hounsgaard, J., and Kiehn, O. (1985). Ca++ dependent bistability induced by serotonin in spinal motoneurons. Exp Brain Res *57*, 422--425. 10.1007/bf00236551 , pmid = 2578974.

26. Husch, A., Dietz, S.B., Hong, D.N., and Harris-Warrick, R.M. (2015). Adult spinal V2a interneurons show increased excitability and serotonin-dependent bistability. J Neurophysiol *113*, 1124-1134. 10.1152/jn.00741.2014.

27. Perrier, J.F., and Hounsgaard, J. (2003). 5-HT2 receptors promote plateau potentials in turtle spinal motoneurons by facilitating an L-type calcium current. J Neurophysiol *89*, 954--959. 10.1152/jn.00753.2002 , pmid = 12574471.

28. Steinbusch, H.W.M. (1981). Distribution of serotonin-immunoreactivity in the central nervous system of the rat-cell bodies and terminals. Neuroscience *6*, 557--618. 10.1016/0306-4522(81)90146-9 , pmid = 7017455.

29. Dahlstroem, A., and Fuxe, K. (1964). Evidence for the Existence of Monoamine-Containing Neurons in the Central Nervous System. I. Demonstration of Monoamines in the Cell Bodies of Brain Stem Neurons. Acta Physiol Scand Suppl, SUPPL 232:231-255.

30. Hornung, J.P. (2003). The human raphe nuclei and the serotonergic system. J Chem Neuroanat *26*, 331-343. 10.1016/j.jchemneu.2003.10.002.

31. Basbaum, A.I., Clanton, C.H., and Fields, H.L. (1978). Three bulbospinal pathways from the rostral medulla of the cat: An autoradiographic study of pain modulating systems. Journal of Comparative Neurology *178* 209--224. 10.1002/cne.901780203 , pmid = 627624.

32. Holstege, G., and Kuypers, H.G.J.M. (1982). The Anatomy of Brain Stem Pathways to the Spinal Cord in Cat. A Labeled Amino Acid Tracing Study. Progress in Brain Research *57*, 145--175. 10.1016/s0079-6123(08)64128-x , pmid = 7156396.

33. Martin, R.F., Jordan, L.M., and Willis, W.D. (1978). Differential projections of cat medullary raphe neurons demonstrated by retrograde labelling following spinal cord

lesions. Journal of Comparative Neurology *182*, 77--88. 10.1002/cne.901820106 , pmid = 701490.

34. Loewy, A.D. (1981). Raphe pallidus and raphe obscurus projections to the intermediolateral cell column in the rat. Brain Res *222*, 129-133. 10.1016/0006-8993(81)90946-x.

35. Holstege, J.C., and Kuypers, H.G.J.M. (1987). Brainstem projections to spinal motoneurons: An update. Neuroscience *23*, 809--821. 10.1016/0306-4522(87)90160-6 , pmid = 2893995.

36. Brust, R., Corcoran, A., Richerson, G., Nattie, E., and Dymecki, S. (2014). Functional and developmental identification of a molecular subtype of brain serotonergic neuron specialized to regulate breathing dynamics. Cell Rep *9*, 2152--2165. 10.1016/j.celrep.2014.11.027 , pmid = 25497093 , pmcid = PMC4351711.

37. Hennessy, M., Corcoran, A., Brust, R., Chang, Y., Nattie, E., and Dymecki, S. (2017). Activity of Tachykinin1-Expressing Pet1 Raphe Neurons Modulates the Respiratory Chemoreflex. J Neurosci *37*, 1807--1819. 10.1523/jneurosci.2316-16.2016 , pmid = 28073937.

38. Okaty, B., Freret, M., Rood, B., Brust, R., Hennessy, M., deBairos, D., Kim, J., Cook, M., and Dymecki, S. (2015). Multi-Scale Molecular Deconstruction of the Serotonin Neuron System. Neuron *88*, 774--791. 10.1016/j.neuron.2015.10.007 , pmid = 26549332 , pmcid = PMC4809055.

39. Okaty, B., Commons, K., and Dymecki, S. (2019). Embracing diversity in the 5-HT neuronal system. Nat Rev Neurosci *20*, 397--424. 10.1038/s41583-019-0151-3 , pmid = 30948838.

40. Okaty, B.W., Sturrock, N., Escobedo Lozoya, Y., Chang, Y., Senft, R.A., Lyon, K.A., Alekseyenko, O.V., and Dymecki, S.M. (2020). A single-cell transcriptomic and anatomic atlas of mouse dorsal raphe Pet1 neurons. Elife *9*. 10.7554/eLife.55523.

41. Veasey, S.C., Fornal, C.A., Metzler, C.W., and Jacobs, B.L. (1995). Response of serotonergic caudal raphe neurons in relation to specific motor activities in freely moving cats. The Journal of Neuroscience *15*, 5346--5359. 10.1523/jneurosci.15-07-05346.1995 , pmid = 7623157 , pmcid = PMC6577863.

42. Niederkofler, V., Asher, T.E., Okaty, B.W., Rood, B.D., Narayan, A., Hwa, L.S., Beck, S.G., Miczek, K.A., and Dymecki, S.M. (2016). Identification of Serotonergic Neuronal Modules that Affect Aggressive Behavior. Cell Rep *17*, 1934-1949. 10.1016/j.celrep.2016.10.063.

43. Alvarez, F.J., Pearson, J.C., Harrington, D., Dewey, D., Torbeck, L., and Fyffe, R.E. (1998). Distribution of 5-hydroxytryptamine-immunoreactive boutons on alpha-motoneurons in the lumbar spinal cord of adult cats. J Comp Neurol *393*, 69-83.

44. Reardon, T.R., Murray, A.J., Turi, G.F., Wirblich, C., Croce, K.R., Schnell, M.J., Jessell, T.M., and Losonczy, A. (2016). Rabies Virus CVS-N2c(DeltaG) Strain Enhances Retrograde Synaptic Transfer and Neuronal Viability. Neuron *89*, 711-724. 10.1016/j.neuron.2016.01.004.

45. Simon, C.M., Dai, Y., Van Alstyne, M., Koutsioumpa, C., Pagiazitis, J.G., Chalif, J.I., Wang, X., Rabinowitz, J.E., Henderson, C.E., Pellizzoni, L., and Mentis, G.Z. (2017). Converging Mechanisms of p53 Activation Drive Motor Neuron Degeneration in Spinal Muscular Atrophy. Cell Rep *21*, 3767-3780. 10.1016/j.celrep.2017.12.003.

46. Seo, C., Guru, A., Jin, M., Ito, B., Sleezer, B.J., Ho, Y.Y., Wang, E., Boada, C., Krupa, N.A., Kullakanda, D.S., et al. (2019). Intense threat switches dorsal raphe serotonin neurons to a paradoxical operational mode. Science *363*, 538-542. 10.1126/science.aau8722.

47. Veasey, S.C., Fornal, C.A., Metzler, C.W., and Jacobs, B.L. (1997). Single-unit responses of serotonergic dorsal raphe neurons to specific motor challenges in freely moving cats. Neuroscience *79*, 161-169. 10.1016/s0306-4522(96)00673-2.

48. Nagel, K.I., and Doupe, A.J. (2006). Temporal processing and adaptation in the songbird auditory forebrain. Neuron *51*, 845-859. 10.1016/j.neuron.2006.08.030.

49. Dayan, P.A., L.F. (2001). Theoretical Neuroscience (The MIT Press).

50. Capelli, P., Pivetta, C., Esposito, M.S., and Arber, S. (2017). Locomotor speed control circuits in the caudal brainstem. Nature *551*, 373--377. 10.1038/nature24064 , pmid = 29059682.

51. Caggiano, V., Leiras, R., Goñi-Erro, H., Masini, D., Bellardita, C., Bouvier, J., Caldeira, V., Fisone, G., and Kiehn, O. (2018). Midbrain circuits that set locomotor speed and gait selection. Nature *553*, 455--460. 10.1038/nature25448 , pmid = 29342142 , pmcid = PMC5937258.

52. Ferreira-Pinto, M.J., Ruder, L., Capelli, P., and Arber, S. (2018). Connecting Circuits for Supraspinal Control of Locomotion. Neuron *100*, 361--374. 10.1016/j.neuron.2018.09.015 , pmid = 30359602.

53. Berg, E.M., Mrowka, L., Bertuzzi, M., Madrid, D., Picton, L.D., and El Manira, A. (2023). Brainstem circuits encoding start, speed, and duration of swimming in adult zebrafish. Neuron *111*, 372-386 e374. 10.1016/j.neuron.2022.10.034.

54. Tovote, P., Esposito, M.S., Botta, P., Chaudun, F., Fadok, J.P., Markovic, M., Wolff, S.B., Ramakrishnan, C., Fenno, L., Deisseroth, K., et al. (2016). Midbrain circuits for defensive behaviour. Nature *534*, 206-212. 10.1038/nature17996.

55. Leiras, R., Cregg, J.M., and Kiehn, O. (2022). Brainstem Circuits for Locomotion. Annual Review of Neuroscience *45*, 1--23. 10.1146/annurev-neuro-082321-025137 , pmid = 34985919.

56. Hsu, L.-J., Bertho, M., and Kiehn, O. (2023). Deconstructing the modular organization and real-time dynamics of mammalian spinal locomotor networks. Nature Communications *14*, 873. 10.1038/s41467-023-36587-w , pmid = 36797254 , pmcid = PMC9935527.

57. DePuy, S.D., Kanbar, R., Coates, M.B., Stornetta, R.L., and Guyenet, P.G. (2011). Control of Breathing by Raphe Obscurus Serotonergic Neurons in Mice. The Journal of Neuroscience *31*, 1981--1990. 10.1523/jneurosci.4639-10.2011 , pmid = 21307236 , pmcid = PMC3071248.

58. Bacon, S.J., Zagon, A., and Smith, A.D. (1990). Electron microscopic evidence of a monosynaptic pathway between cells in the caudal raphe nuclei and sympathetic preganglionic neurons in the rat spinal cord. Exp Brain Res *79*, 589-602. 10.1007/BF00229327.

59. Lindsay, A.D., and Feldman, J.L. (1993). Modulation of respiratory activity of neonatal rat phrenic motoneurones by serotonin. J Physiol *461*, 213-233. 10.1113/jphysiol.1993.sp019510.

60. Noga, B.R., Turkson, R.P., Xie, S., Taberner, A., Pinzon, A., and Hentall, I.D. (2017). Monoamine Release in the Cat Lumbar Spinal Cord during Fictive Locomotion Evoked by the Mesencephalic Locomotor Region. Front Neural Circuits *11*, 59. 10.3389/fncir.2017.00059.

61. Carrive, P. (1993). The periaqueductal gray and defensive behavior: functional representation and neuronal organization. Behav Brain Res *58*, 27-47. 10.1016/0166-4328(93)90088-8.

62. Watson, T.C., Cerminara, N.L., Lumb, B.M., and Apps, R. (2016). Neural Correlates of Fear in the Periaqueductal Gray. J Neurosci *36*, 12707-12719. 10.1523/JNEUROSCI.1100-16.2016.

63. Fonseca, M.I., Ni, Y.G., Dunning, D.D., and Miledi, R. (2001). Distribution of serotonin 2A, 2C and 3 receptor mRNA in spinal cord and medulla oblongata. Brain Res Mol Brain Res *89*, 11--19. 10.1016/s0169-328x(01)00049-3 , pmid = 11311971.

64. Koyama, Y., Kondo, M., and Shimada, S. (2017). Building a 5-HT3A Receptor Expression Map in the Mouse Brain. Sci Rep *7*, 42884. 10.1038/srep42884 , pmid = 28276429.

65. L, J.B., and A, F.C. (1993). 5-HT and motor control: a hypothesis. Trends Neurosci *16*, 346--352. 10.1016/0166-2236(93)90090-9 , pmid = 7694403.

66. Basbaum, A.I., Clanton, C.H., and Fields, H.L. (1976). Opiate and stimulus-produced analgesia: functional anatomy of a medullospinal pathway. Proc Natl Acad Sci U S A *73*, 4685-4688. 10.1073/pnas.73.12.4685.

67. Cai, Y.Q., Wang, W., Hou, Y.Y., and Pan, Z.Z. (2014). Optogenetic activation of brainstem serotonergic neurons induces persistent pain sensitization. Mol Pain *10*, 70. 10.1186/1744-8069-10-70.

68. Ganley, R.P., de Sousa, M.M., Werder, K., Ozturk, T., Mendes, R., Ranucci, M., Wildner, H., and Zeilhofer, H.U. (2023). Targeted anatomical and functional identification of antinociceptive and pronociceptive serotonergic neurons that project to the spinal dorsal horn. Elife *12*. 10.7554/eLife.78689.

69. Bouvier, J., Caggiano, V., Leiras, R., Caldeira, V., Bellardita, C., Balueva, K., Fuchs, A., and Kiehn, O. (2015). Descending Command Neurons in the Brainstem that Halt Locomotion. Cell *163*, 1191--1203. 10.1016/j.cell.2015.10.074 , pmid = 26590422.

70. Callaway, E.M., and Luo, L. (2015). Monosynaptic Circuit Tracing with Glycoprotein-Deleted Rabies Viruses. The Journal of neuroscience : the official journal of the Society for Neuroscience *35*, 8979--8985. 10.1523/jneurosci.0409-15.2015 , pmid = 26085623 , pmcid = PMC4469731.

71. Miyamichi, K., Shlomai-Fuchs, Y., Shu, M., Weissbourd, B.C., Luo, L., and Mizrahi, A. (2013). Dissecting local circuits: parvalbumin interneurons underlie broad feedback control of olfactory bulb output. Neuron *80*, 1232-1245. 10.1016/j.neuron.2013.08.027.

72. Balaskas, N., Abbott, L.F., Jessell, T.M., and Ng, D. (2019). Positional Strategies for Connection Specificity and Synaptic Organization in Spinal Sensory-Motor Circuits. Neuron *102*, 1143-1156 e1144. 10.1016/j.neuron.2019.04.008.

73. Azim, E., Jiang, J., Alstermark, B., and Jessell, T.M. (2014). Skilled reaching relies on a V2a propriospinal internal copy circuit. Nature *508*, 357-363. 10.1038/nature13021.

74. Bikoff, J.B., Gabitto, M.I., Rivard, A.F., Drobac, E., Machado, T.A., Miri, A., Brenner-Morton, S., Famojure, E., Diaz, C., Alvarez, F.J., et al. (2016). Spinal Inhibitory Interneuron Diversity Delineates Variant Motor Microcircuits. Cell *165*, 207-219. 10.1016/j.cell.2016.01.027.

75. Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. Development *145*. 10.1242/dev.165753.

76. Kuehn, E., Clausen, D.S., Null, R.W., Metzger, B.M., Willis, A.D., and Ozpolat, B.D. (2022). Segment number threshold determines juvenile onset of germline cluster expansion in Platynereis dumerilii. J Exp Zool B Mol Dev Evol *338*, 225-240. 10.1002/jez.b.23100.

77. D'Elia, K.P., Hameedy, H., Goldblatt, D., Frazel, P., Kriese, M., Zhu, Y., Hamling, K.R., Kawakami, K., Liddelow, S.A., Schoppik, D., and Dasen, J.S. (2023). Determinants of motor neuron functional subtypes important for locomotor speed. Cell Rep *42*, 113049. 10.1016/j.celrep.2023.113049.

971   78.   Miri, A., Warriner, C.L., Seely, J.S., Elsayed, G.F., Cunningham, J.P., Churchland, M.M.,
972   and Jessell, T.M. (2017). Behaviorally Selective Engagement of Short-Latency Effector
973   Pathways by Motor Cortex. Neuron *95*, 683-696 e611. 10.1016/j.neuron.2017.06.042.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Extended Data Figure 1

a Rosa26 CAG FRT STOP FRT loxP STOP loxP syn-GFP

Niederkofler, 2016
Hennessy, 2017

b Breeding

Pet1-Flpe X Tac1-Cre or Egr2-Cre X synGFP

c

| Thoracic | Lumbar |

Egr2-Pet1

5-HT
synGFP

Tac1-Pet1

d

Tac1-Pet1   *Egr2-Pet1*

Tac1-Pet1 — Pet1:Tac1 TPH2 — Egr2-Pet1 — Pet1:Tac1 TPH2

Bregma −6.84 — ROb RPa

Bregma −5.68 — RMg

Bregma −4.84 — DRN MRN

Extended Data Figure 2

Extended Data Figure 3

Extended Data Figure 4

Extended Data Figure 5

Extended Data Figure 6

Ai65 (Cre & Flpe-depdendent)

a Rosa26 —CAG—FRT—STOP—FRT—loxP—STOP—loxP— Hist2B-HA — N2cG(O) — TVA66T —

b *Tac1:Pet1:N2cG-TVA*

Tac1-Cre X Pet1-Flpe X N2cG-TVA

ROb/Pa | RMg | LPGi

HA
TPH2

c EnvA-RVΔG-tdTomato

No Cre/Flpe control

N2cG-TVA

TPH2
HA
RV-tdT

ROb/Pa

LPGi

Ai9 (Cre-depdendent)

d Rosa26 —CAG—loxP—STOP—loxP— Hist2B-HA — N2cG(O) — TVA66T —

e *Pet1:N2cG-TVA*

Pet1-Cre X N2cG-TVA

ROb/Pa | RMg | LPGi

HA
TPH2

f EnvA-RVΔG-tdTomato

No Cre control

N2cG-TVA

TPH2
HA
RV-tdT

ROb/Pa

LPGi

g

TPH2
HA
RV-tdT

ROb/Pa

LPGi

RV-tdT | CnF

RV-tdT | PAG

RV-tdT | Hyp

Extended Data Figure 7