Interleukin 6 drives durable T cell-mediated immunity to pancreatic cancer

Short title: IL-6 drives durable T cell immunity in PDAC

Paige C. Arneson-Wissink[1], Alexandra Q. Bartlett[2], Heike Mendez[1], Xinxia Zhu[1], Peter R.

Levasseur[1], Parham Diba[1, 3], Namratha Turuvekere Vittala Murthy[4], Jessica Dickie[1], Matthew

McWhorter[2], Antony Jozic[4], Yulia Eygeris[4], Gaurav Sahay[4, 5], Katelyn T. Byrne[6, 7], Gregory D.

Scott[8], Robert Eil[2], Aaron J. Grossberg[1, 6, 7, 9]*

[1] Department of Radiation Medicine, Oregon Health & Science University, Portland, OR, USA

[2] Division of Surgical Oncology, Department of Surgery, Knight Cancer Institute, Oregon

Health & Science University, Portland, OR, USA

[3] Medical Scientist Training Program, Oregon Health and Science University, Portland, OR,

USA

[4] Department of Pharmaceutical Sciences, College of Pharmacy, Oregon State University,

Portland, OR, USA

[5] Department of Biomedical Engineering, Oregon Health & Science University, Portland, OR,

USA

[6] Department of Cell, Development, and Cancer Biology, Oregon Health & Science University,

Portland, OR, USA

[7] Brenden Colson Center for Pancreatic Care, Oregon Health & Science University, Portland,

OR, USA

[8] Department of Pathology, Oregon Health & Science University, Portland, OR, 97239, USA

[9] Cancer Early Detection Advanced Research Center, Oregon Health & Science University,

Portland, OR, USA

*Corresponding Author: Aaron J. Grossberg, 2730 S Moody Ave, Oregon Health and

Science University, Portland, OR, 97201, 503-494-9945, grossber@ohsu.edu

Preprint Server: https://doi.org/10.1101/2024.09.26.615308

Word Count: 3583

Grant Support: This work was supported by the National Cancer Institute (PCAW:

K99CA286709; GS: R01CA270783; AJG: K08CA245188, R37CA280692, R01CA264133; RE:

K08CA256179), the Brenden Colson Center for Pancreatic Care, the Oregon Pancreas Tissue

Registry, the Histopathology Shared Resource for pathology studies (University Shared

Resource Program at Oregon Health and Sciences University and the Knight Cancer Institute

(P30 CA069533 and P30 CA069533 13S5)), the OHSU Flow Cytometry Shared Resource

(OHSU Knight Cancer Institute NCI Cancer Center Support Grant P30CA069533), and the

Advanced Multiscale Microscopy Shared Resource (OHSU Knight Cancer Institute, NIH P30

CA069533). RE was also supported by grants from the American Association of Cancer

Research, ASCO, and Pancreatic Cancer Action Network.

Data Transparency: All data will be available on the public repository, Mendeley Data, upon

publication. Further information and resources, including plasmid sequences, engineered KPC

cells, and raw data will be shared upon reasonable request to Aaron J. Grossberg

(grossber@ohsu.edu).

Abbreviations:

BSR	Blasticidin Resistance	
CAR	Chimeric Antigen Receptor	
CD	Cell Differentiation	
ELISA	Enzyme-Linked Immunosorbent Assay	
Fbxo32	F-Box Protein 32	
Foxp3	Forkhead Box Protein P3	
H&E	Hematoxylin and Eosin	
IL-6	Interleukin 6	
KPC	$Kras^{G12D/+}$, $Tp53^{R172H/+}$, Pdx1-Cre	
NK	Natural Killer	
OT-PDAC	Orthotopic Pancreatic Ductal Adenocarcinoma	
PDAC	Pancreatic Ductal Adenocarcinoma	
Qpcr	Quantitative Polymerase Chain Reaction	
Trim63	Tripartite Motif Containing 63	

SYNOPSIS

Interleukin-6 induces rapid and durable T cell-driven immune clearance of pancreatic ductal adenocarcinoma. The anti-tumor immune microenvironment is hallmarked by increased lymphoid aggregate formation, increased CD4 T cell abundance, and decreased Treg abundance.

ABSTRACT

Background & Aims: Tumor immune resistance is recognized as a contributor to low survivorship in pancreatic ductal adenocarcinoma (PDAC). The inflammatory cytokine interleukin-6 (IL-6) promotes polarization of CD4 T cell populations away from immune tolerance, and induces differentiation of cytotoxic CD8 T cells. This work aims to test whether IL-6 could stimulate an anti-tumor response in PDAC

Methods: We overexpressed IL-6 in multiple $Kras^{G12D/+}$, $Tp53^{R172H/+}$, Pdx1-Cre (KPC) cell lines, which were orthotopically implanted in mice (OT-PDAC^{IL6}). We followed mouse survival and measured tumor growth, tumor histology, and plasma IL-6 at 5 and 10 days after tumor implantation. We measured tumor immune cell infiltration via flow cytometry and histology. We used antibody-based T cell depletion and secondary tumor implantation rechallenge to test the dependency of the durable immune reaction on T cells. We use lipid nanoparticle (LNP)-based delivery of IL-6 mRNA to the pancreas as an orthogonal approach for testing the effect of elevated IL-6 in the tumor microenvironment on anti-tumor T cell invasion.

Results: Improved survival occurred in all instances of OT-PDAC^{IL6}, with one cell line

(KxPxCx) reproducibly resulting in long-term recurrence-free survival. With KxPxCx cells,

circulating IL-6 was 100-fold higher in OT-KxPxCx^{IL6} than in OT-KxPxCx^{parental} mice. Flow

cytometry revealed increased T cells and NK cells, and decreased T regulatory cells, and we

observed significantly increased lymphoid aggregates in OT-KXPXCX^{IL6} as compared to OT--

KxPxCx^{parental} tumors. Antibody-based CD4$^+$ and CD8$^+$ T cell depletion prevented tumor

clearance and completely abolished the survival advantage in OT-KxPxCx^{IL6} mice. The anti-

tumor immune response to OT-KxPxCx^{IL6} rendered mice immune to re-challenge with OT-

KxPxCx^{parental} tumors. LNP delivery of IL-6 to the pancreas elevated systemic IL-6 levels ~50

fold, lowered tumor burden, and increased anti-tumor T cell phenotypes.

Conclusions: Locally high IL-6 concentrations potently enhance the T cell-mediated anti-

tumor response to PDAC.

Key words: Interleukin 6; Pancreatic Ductal Adenocarcinoma; T regulatory cells

INTRODUCTION

Pancreatic ductal adenocarcinoma (PDAC) is associated with an immunosuppressive microenvironment, characterized by low numbers of infiltrating T cells and relatively higher numbers of immunosuppressive T regulatory cells (Tregs)[1, 2]. Despite clinical trials that have established the safety of immunotherapies in PDAC, the immune profile of PDAC limits the clinical efficacy of these interventions[3]. Conversely, increased T cell infiltration with low Treg infiltration is associated with improved patient survival [5]. Strategies that promote T cell activation, such as agonizing CD40 antibodies, increase tertiary immune structure formation and benefit mouse survival in pre-clinical models[4]. A small but growing body of literature suggests that tumor-localized expression of immunomodulatory cytokines, such as IL-6, IL-12, CXCL13, and CCL21 can also improve immune reactivity to tumors by driving localization and differentiation of tumor-infiltrating lymphocytes, which accumulate in tertiary lymphoid structures or lymphoid aggregates[5-7].

Interleukin-6 (IL-6) signaling is associated with reduced survival in patients with PDAC attributed to the promotion of tumorigenesis and metastasis [8-11]. Elevated circulating IL-6 also associates with sarcopenia in PDAC patients, and we previously demonstrated that host intrinsic IL-6 signaling is necessary for mice to develop PDAC cachexia[12]. Published literature shows that loss of IL-6 attenuates PDAC development, metastasis, and cachexia onset [12-14]. However, IL-6 is also well-established as an immunomodulatory cytokine and contributes to polarization of CD4 T cell populations, differentiation of cytotoxic CD8 T cells, inhibition of inducible Treg development, and expansion and anti-tumor activity in chimeric antigen receptor (CAR) T cell therapies[15-17]. Leveraging the immunomodulatory properties of IL-6 may promote

anti-tumor immunity in PDAC.

In the current study, we sought to investigate the effect of IL-6 overexpression on immune

clearance of PDAC and cachexia development. We used two independent modalities to

elevate IL-6 expression in the pancreas: stable IL-6 overexpressing PDAC cells (PDAC^{IL6})

from PDAC^{parental} (KPC) cell lines[18], and lipid nanoparticles (LNPs) containing IL-6 mRNA. We

found that PDAC^{IL6} cells produced extremely high levels of IL-6 and induced severe cachexia

within days of orthotopic implantation (OT-PDAC^{IL6}). We also observed dramatic changes in

tumor growth dynamics, with one cell line eliciting complete tumor clearance and long-term,

recurrence-free mouse survival. IL-6 LNPs targeted the pancreas, caused high levels of

circulating IL-6, and limited PDAC tumor growth. We then pursued a series of studies to

understand the immune response to high intra-tumoral IL-6. The work shown here presents IL-

6 overexpression as a driver of T cell infiltration in PDAC, which poises mice for a favorable

anti-tumor immune response.

RESULTS

Tumor-specific IL-6 overexpression induces spontaneous tumor clearance and

cachexia recovery. We developed PDAC^{IL6} cells from the KPC PDAC cell line KxPxCx, using

ecotropic retroviral transduction. We first observed that mice implanted with OT-KxPxCx^{IL6}

tumors experienced nearly 100% survival without reaching terminal endpoints (**Figure 1A**).

OT-KxPxCx^{IL6} mice euthanized approximately 10 days after all OT-KxPxCx^{parental} had

significantly lower tumor/pancreas mass, and no detectable IL-6 in the plasma (**Figure 1B-C**).

Deeper histological assessment of tumors showed that both OT-KxPxCx^{IL6} and OT-

KxPxCx^{parental} developed poorly differentiated, infiltrative carcinoma by five days (**Figure 1D-F**).

OT-KxPxCx^{IL6} tumors were slightly smaller in mass and histological area than OT-

KxPxCx^{parental} tumors (**Figure 1F-G**). By 12 days, OT-KxPxCx^{parental} mice had reached humane

euthanasia endpoint, with carcinoma covering approximately 80% of total tissue area, in

contrast, OT-KxPxCx^{IL6} tissue was completely devoid of tumor (**Figure 1E,G, Table S1**). We

confirmed this result molecularly using qPCR for the codon-optimized *Il6* transgene, which was

not detected in OT-KxPxCx^{IL6} whole pancreas tissue after five days (**Figure 1H, Figure S1**).

While lung and liver metastases were detected by *Il6* transgene qPCR at five days, there were

no metastases detected at 12 days (**Figure S2**). Spontaneous tumor clearance also occurred

when KxPxCx^{IL6} cells were implanted subcutaneously, indicating that the phenomenon is not

specific to intrapancreatic administration (**Figure S3**).

OT-KxPxCx^{parental} plasma IL-6 levels were in line with previously published values, while OT-

KxPxCx^{IL6} levels reached 100-fold higher levels at day 5, followed by undetectable levels at

later time points (**Figure 1I**) [10, 12]. Because high circulating IL-6 is associated with cachexia

development in PDAC, we investigated total body and muscle mass[10]. High plasma IL-6 and

peak tumor burden associated with decreased body mass, which recovered with the

normalization of IL-6 levels after tumor clearance in the OT-KxPxCx^{IL6} model (**Figure 1J**).

Gross muscle mass trended downward at five days and was significantly decreased at 10

days, while muscle atrophy gene (*Trim63* and *Fbxo32*) expression was elevated at 5 days and

reduced to baseline at 10 days. These data reflect a delay in tissue recovery as the tumor

resolves and body mass returns to baseline (**Figure 2K-L**). Collectively, these data show that

IL-6 overexpression in PDAC cells leads to severe wasting, followed by body mass recovery

upon spontaneous tumor resolution.

Multiple OT-PDAC[IL6] cell lines, but not CD90.1 transduction control lines, display improved survival outcomes. We then tested whether the impact of IL-6 overexpression on tumor clearance and mouse survival applied to other murine PDAC cell lines: KPC-1199[19], and KPC-1242[20]. While both PDAC[IL6] cell lines showed significantly improved survival, only OT-1242[IL6] tumors remained smaller at humane endpoint, when compared to the respective parental control line (**Figure 2A-B, D-E**). High levels of IL-6 overexpression, quantified as plasma IL-6 levels at endpoint, correlated with less tumor burden: 1242[IL6] caused ~250 fold elevation of IL-6, and 1199[IL6] did not show a significant overexpression or reduction in tumor mass (**Figure 2C,F**). We validated the clearance and survival phenotypes using an independently generated KxPxCx[IL6] (KX2[IL6]) (**Figure 2G-I**). This led us to hypothesize that reduction in tumor mass was driven by IL-6 overexpression. We tested this by using a KxPxCx transduction control cell line, where we expressed CD90.1 alone without IL-6 (OT-KxPxCx[CD90.1]). OT-KxPxCx[CD90.1] mice replicated our findings with non-transduced OT-KxPxCx[parental] mice. Both groups had reached humane euthanasia endpoint by 12 days, carcinoma covering approximately 80% of total tissue area, circulating IL-6 elevated to the expected 200-300 pg/mL range, and body mass loss not exceeding 10% (**Figure 2J-M, Table S1**). Finally, we tested the tumor response to IL-6 overexpression when fewer cells are implanted, thus prolonging the period of host exposure to tumor before humane euthanasia criteria are met. We tested 5,000 KxPxCx cells, 100,000 KPC-1199 cells, and 100,00 KPC-1242 cells, and found that survival, tumor mass, and plasma IL-6 replicated our findings with 1 million cell implantation doses (**Figure 3A-I**). We therefore concluded that the tumor clearance we observed was specific to IL-6 overexpression and most robust in the KxPxCx cell line, which we selected for continued testing. Because we did not observe a dose-dependent effect,

we proceeded to exclusively test the implantation dose of 1 million KxPxCx[IL6] cells, which

aligns with our previously published work.[12]

Tumor IL-6 overexpression induces lymphocytic anti-tumor immune response. Spleens

from OT-KxPxCx[IL6] mice were significantly enlarged, consistent with IL-6-induced peripheral T

cell expansion[15] (**Figure 4A**). We therefore hypothesized that the extreme levels of IL-6

generated in the pancreas of OT-KxPxCx[IL6] mice facilitated immune cell expansion and

immune-mediated tumor clearance. We used flow cytometry to evaluate the immune profile of

OT-KxPxCx[IL6] and OT-KxPxCx[parental] tumors and discovered that pancreata bearing OT-

KxPxCx[IL6] tumors were enriched with CD4$^+$ T cells and Natural Killer (NK) cells, while Foxp3$^+$

T-regulatory cells were decreased (**Figure 4B-I, S4-5**). We did not identify increases in Th17

cells, or Granzyme B$^+$ CD8 T cells, which are two phenotypes associated with increased anti-

tumor immunity and IL-6 signaling (**Figure S6-7**). We did, however, see a decreased ratio of

Foxp3$^+$ CD4 T cells to CD8 T cells, indicating a shift towards a less immunosuppressive tumor

microenvironment (**Figure 4G**). Because increased tumor-infiltrating lymphocytes are

associated with improved outcomes and therapeutic efficacy, we next assessed the

localization of T cells relative to tumor cells[21]. By immunofluorescent staining, we found that

OT-KxPxCx[parental] and OT-KxPxCx[IL6] mice had equal densities of CD3$^+$ T cells in both tumor-

associated stroma and tumor nests (**Figure 4J-L**). These data initially contradicted the flow

cytometry data. However, upon histological assessment, we found that OT-KxPxCx[IL6]

pancreata exhibited a significantly greater number of tumor-associated lymphoid aggregates

(**Figure 4M-N**). In PDAC, lymphoid aggregate formation is associated with improved

survival[22, 23]. In OT-KxPxCx[IL6] mice, the lymphoid aggregates were often near, but not

necessarily located within the tumor stroma, accounting for the differences between our flow

cytometry and immunofluorescence data. In summary, locally high IL-6 induces accumulation

of lymphoid aggregates, increased CD4$^+$ T cells, and decreased Foxp3$^+$ T regulatory cells.

These conditions favor the hypothesis that tumor clearance occurs via an anti-tumor T cell

response.

T cells are necessary for OT-KxPxCx^{IL6} tumor clearance. We hypothesized that T cells are

necessary for OT-KxPxCx^{IL6} tumor clearance, which we tested via antibody-based CD4 and

CD8 depletion in tumor-bearing mice, compared to IgG control treatment. Mice received

antibody injections two days before tumor implantation and every four days following. We

validated T cell depletion via flow cytometry for total T cells (CD90.2), and CD4$^+$ and CD8$^+$

cells (**Figure 5A-C**). We followed humane endpoints for the CD4/CD8-depleted group, which

reached euthanasia criteria at 8-11 days post-implantation (**Figure 5D**). We euthanized all

sham and IgG-treated mice when all CD4/CD8-depleted mice were euthanized, although they

were healthy at the time. We used a second engineered cell line, KxPxCx$^{IL6-LUC}$, which co-

expressed IL-6 and luciferase, to measure tumor burden over time. Both OT-KxPxCx$^{IL6-LUC}$

longitudinal data, and OT-KxPxCx^{IL6} endpoint tumor mass data show that CD4/CD8-depletion

prevents tumor clearance (**Figure 5E-G**). Histological evaluation by a board-certified

pathologist found tumor present in 10/10 CD4/CD8-depleted mice, and in 1/8 IgG control mice

(**Figure 5H**). This was supported by *Il6*-transgene qPCR data and plasma IL-6 levels, which

showed no elevation of IL-6 in 6/8 IgG-treated PDAC mice (**Figure 5I-J**). We conclude that

CD4 and CD8 T cells are necessary for OT-KxPxCx^{IL6} tumor clearance.

Because we previously saw recovery of wasting phenotypes in OT-KxPxCx^{IL6} mice as the

tumor resolved, we also investigated cachexia resolution in T cell-depleted mice. CD4/CD8

depletion prevented body mass recovery and led to sustained muscle wasting as evidenced by

decreased muscle mass and increased atrophy-related gene expression (*Trim63, Fbxo32*)

(**Figure 5K-M**). The recovery in wasting is therefore directly associated with immune-mediated

tumor clearance.

OT-KxPxCx^{IL6} induces a durable T cell response to OT-KxPxCx^{Parental} tumors. Given the

dependency of the anti-OT-KxPxCx^{IL6} response on T cells, which are known to elicit

immunologic memory, we hypothesized that OT-KxPxCx^{IL6} tumors would generate a durable

immune response that would protect mice in the case of a second tumor exposure. We first

tested this hypothesis in a cohort of mice that had recovered from sham surgery or OT-

KxPxCx^{IL6} for over two months, and were then rechallenged with OT-KxPxCx$^{parental-LUC}$

(luciferase-expressing) at 76 days after the initial surgery (**Figure 6A**). Sham-recovered mice

implanted with KxPxCx$^{parental-LUC}$ reached euthanasia criteria 13-14 days after rechallenge

implantation. In contrast, there were no deaths following rechallenge in the OT-KxPxCx^{IL6}-

recovered group (**Figure 6B**). Tumor burden measured longitudinally by IVIS and terminally (in

sham-recovered mice only) revealed significant tumor growth only in sham-recovered mice

(**Figure 6C-D**). Sham-recovered mice also lost more body mass than OT-KxPxCx^{IL6}-recovered

mice following rechallenge (**Figure 6E**). These data indicate that the potent anti-tumor immune

response elicited by high concentrations of IL-6 is durable and, once established, not

dependent on coincident supraphysiologic IL-6 levels. We then used CD4/CD8 antibody

depletion to determine whether the tumor clearance during rechallenge was indeed T cell

mediated. In this study, all mice recovered from OT-KxPxCx^{IL6} for 26 days before starting

antibody treatment and were given sham surgery or implanted with OT-KxPxCx^{parental} tumors

on day 28. We monitored mice for 12 days, until CD4/CD8-depleted mice reached humane

euthanasia criteria (**Figure 6F**). We confirmed that clearance of the rechallenge OT-

KxPxCx[parental] tumor is dependent on T cells, which were effectively depleted by the antibody

treatment (**Figure 6G-I**). Furthermore, we confirmed that CD4/CD8-depletion does not result in

the outgrowth of potentially covert KxPxCx[IL6] cells, as no tumor was present in any of the OT-

KxPxCx[IL6]-recovered, sham-rechallenged, CD4/CD8-depleted mice (**Figure 6G**). These data

show that the T cell response induced by OT-KxPxCx[IL6] provides durable protection against

molecularly similar PDAC tumor growth.

Lipid nanoparticle delivery of IL-6 mRNA replicates the anti-tumor effects observed in

OT-PDAC[IL6]. To test whether the anti-tumor effects of elevated IL-6 in the tumor

microenvironment could be derived through a therapeutically-relevant method, we engineered

lipid nanoparticles (LNPs) that deliver IL-6 mRNA, are administered by IP injection, and target

the pancreas. We used SM-102 LNPs containing luciferase mRNA to validate the particle

delivery 6 hours after IP injection in healthy mice (**Figure 7A**). We found that luciferase

expression was highest in the pancreas, but was also expressed in the liver, and was nearly

undetectable in the spleen (**Figure 7A-B**). Following validation of dosing, we designed an

experimental setup with escalating doses of LNPs, which were administered 3 hours prior to

OT-PDAC[parental] (no IL-6 overexpression by the tumor) implantation and every other day after

(**Figure 7C**). We included an empty LNP group that was dose-matched for LNP cholesterol

content to control for potentially immunogenic effects of the LNPs alone. We found that OT-

PDAC[parental] mice injected with IL-6 LNPs lost significant body mass and had approximately 50-

fold higher circulating IL-6 than empty LNP or no LNP control OT-PDAC[parental] mice (**Figure 7F-**

G). IL-6 LNP OT-PDAC[parental] mice euthanized 8 days post-implantation had significantly lower

tumor burden and larger spleens (**Figure 7G-H**). We used flow cytometry to evaluate the tumor

immune profile and found that tumors from OT-PDAC[parental] mice injected with IL-6 LNPs did

not have significantly more CD4$^+$ or CD8$^+$ T cells (**Figure 7I-J, S9**). Despite this, there were

significantly fewer T-regulatory cells within the CD4 population, and a decreased ratio of

Foxp3$^+$ CD4 T cells to CD8 T cells in IL-6 LNP-treated OT-PDAC[parental] mice, replicating our

findings in OT-PDAC[IL6] (**Figure 7K-L**). Finally, we saw significant enrichment of NK and NK T

cells in the pancreas of IL-6 LNP-treated OT-PDAC[parental] mice (**Figure 7M-N**). There were no

differences in immune populations between the empty LNP and no LNP groups, suggesting

that, in the pancreas with this dosing paradigm, empty SM-102 LNPs did not induce a T cell

response. Together, these data show that IL-6 mRNA delivery to the pancreas results in an

anti-tumor immune landscape and lower tumor burden in OT-PDAC[parental] mice.

DISCUSSION

Our model of PDAC IL-6 overexpression induced a robust, rapid, and durable anti-tumor T cell

response, accompanied by rapid and severe wasting that recovered as the tumor was cleared.

Although IL-6 is traditionally viewed as a negative factor in pancreatic cancer [8-11], we provide

evidence that supraphysiologic levels of IL-6, achieved through tumor cell overexpression of

IL-6 protein or LNP delivery of IL-6 mRNA, were sufficient to induce an anti-tumor immune

landscape in the pancreas. The OT-PDAC[IL6] tumor microenvironment was characterized by

increased lymphoid aggregate formation, elevated CD4$^+$ T cells, decreased Foxp3$^+$ Treg cells,

and elevated NK cells. LNP delivery of IL-6 also drove decreased Treg cell and increased NK

cell infiltration. While the KxPxCx tumor line implanted orthotopically exhibited complete,

recurrence-free survival, subcutaneous tumors of the same line displayed less complete and

slower regression in tumor size. IL-6 overexpression in other KPC cell lines (1199 and 1242)

resulted in significant improvements in mouse survival time, although tumors were never fully

cleared, and mice progressed to humane euthanasia endpoints. These cases highlighted that

the magnitude of anti-tumor immunity due to IL-6 overexpression was partially dependent on

cell-line intrinsic properties.

Unlike cancer types that are now successfully treated with immunotherapy, survival rates for

PDAC patients have increased very slowly over the past decade[24]. PDAC is highly

immunosuppressive, causing immunotherapies, such as checkpoint blockade, to be ineffective

clinically [25]. Previous work indicated that enhancing T cell activation using exogenous agents,

such as agonistic anti-CD40 antibody, improved response to checkpoint blockade therapies

and PDAC tumor regression[1]. The Lesinski Group also used IL-6 blockade to enhance the

effects of PD-L1 or CTLA4 blockade. Combining checkpoint inhibitors with anti-IL-6 reduced

tumor growth and improved T-cell infiltration in implanted murine PDAC models[26, 27]. However,

anti-IL-6 was only effective in bolstering the anti-tumor effects of immune checkpoint blockade;

the IL-6 antibody did not exert anti-tumor effects on its own. Our work raises a fundamental

contradiction regarding the role of IL-6 in PDAC and provides a basis for enhanced IL-6

expression as an alternative method to improve T cell response.

We propose that the effect of IL-6 on PDAC growth is pleiotropic and concentration-dependent.

At low concentrations, IL-6 aids tumor development via signaling directly on neoplastic cells to

drive transformation and growth. This is evidenced by the reduced frequency of high-grade

pancreatic intraepithelial neoplasia observed in an aggressive genetic model of murine PDAC,

when mice were treated with combination PD-L1 and IL-6 blockade[26]. Although some murine

PDAC cells produce IL-6, prior work showed that the source of IL-6 in human PDAC tumors

and orthotopic mouse models is primarily fibroblasts and immune cells[26, 28]. Combination anti-

IL-6 and anti-PD-L1 therapy reduced the number of αSMA+ stromal cells, the loss of which

potentially contributed to limiting PDAC progression in that model[26]. Conversely, at

supraphysiologic concentrations seen in two independent methods of concentrated IL-6

expression in the pancreas (OT-KxPxCx^{IL6} and IL-6 LNP treatment), IL-6 stimulated an anti-

tumor immune response. Circulating levels of IL-6 were approximately 100 times higher in OT-

KxPxCx^{IL6} and 50 times higher after IL-6 LNP treatment than what we detected in OT-

KxPxCx^{parental} mice. We presume that local concentrations in the pancreas were even higher in

both systems. We can infer that pancreatic IL-6 is the initiating signal for T cell accumulation

and eventual tumor clearance; however, the precise manner in which IL-6 mediates this

remains unknown. In addition to increased numbers of tumor-infiltrating T cells, we detected

increased NK cells, and decreased Treg cells intratumorally. It is likely that IL-6, which is a

known immunomodulatory cytokine, impacted multiple cell populations simultaneously to

orchestrate an anti-tumor immune microenvironment[29]. Conversely, it is also likely that lower

levels of IL-6 promote expansion of stromal cells, which can contribute to an immune-deplete

tumor microenvironment. Future work might investigate how high levels of IL-6, as seen in our

OT-KxPxCx^{IL6} model, influence stromal development in PDAC tumors.

IL-6 was also established as a driver of acute-phase protein response in the liver, which

primed the liver for increased metastasis formation[30]. The KxPxCx cell line used most

extensively in this study is not highly metastatic when implanted orthotopically. Liver

metastasis was only detected in one mouse at 5 days post-implantation in the OT-KxPxCx^{IL6}.

At 12 days post-implantation, zero mice had liver metastases detectable by qPCR (**Figure S2**).

Given the high circulating levels of IL-6 in our models, it is likely that STAT3 was activated in

hepatocytes, which could contribute to the rapid onset of cachexia in OT-PDAC[IL6] mice[12].

However, we did not directly assess the activation of acute-phase protein response in our

model. The rapid resolution of the OT-PDAC[IL6] primary tumor and micro-metastatic disease

made this particular model incompatible with rigorously assessing how supraphysiologic IL-6

affected the hepatic metastatic niche. It would be necessary to further investigate the impact

of IL-6 on hepatic inflammation if IL-6 elevation were pursued therapeutically in the future, to

avoid potentially unexpected negative consequences for metastatic disease.

In this work, we confirmed the anti-tumor effects of IL-6 overexpression in PDAC cells using IL-

6 LNPs that targeted the pancreas. LNPs are rising in interest as therapeutic vectors for mRNA

delivery due to their clinical safety and ability to customize particle content and delivery method

to improve tissue-specific targeting[31-34]. Tissue specificity is especially attractive for anti-tumor

therapeutic delivery that minimizes off-target effects[35, 36]. Recent publications show that next-

generation ionizable lipids can improve pancreas specificity of LNPs[37, 38]. LNP-delivered IL-12

was sufficient to prevent PDAC growth[37, 38]. This, and our own work with IL-6, adds to a

growing body of literature showing that local expression of immunomodulatory cytokines leads

to an anti-tumor T cell response[5-7]. Further work is needed to understand how the T cell

response to cytokine overexpression might be conserved and how it could be best translated

to clinical use.

Our data support the widely-accepted notion that IL-6 induces acutely negative effects, as

evidenced by rapid body mass loss of 10-15% in OT-KxPxCx[IL6] mice and mice that received

IL-6 LNPs (**Figure 2H**). In OT-KxPxCx[IL6] mice, which derived IL-6 expression primarily from

the tumor cells, tumor clearance led to rapid recovery in body mass. For patients with PDAC,

who are already vulnerable to weight loss, the pro-cachectic effects of IL-6 would likely limit the

clinical utility of therapies that elevate IL-6 systemically. Because IL-6 is a secreted protein, it is

likely that even pancreas-directed delivery, for example targeted LNP delivery of mRNA, would

pose a high risk of side effects for patients. Instead, future work will focus on identifying the

immune subpopulations that integrate IL-6 signaling into a T cell response, with the goal of

identifying targetable drivers of the anti-tumor response. Using IL-6 overexpression as a model

of effective PDAC clearance will open doors to understanding IL-6-mediated T cell activation in

PDAC and the intricacies of cachexia resolution after PDAC tumor clearance.

ACKNOWLEDGEMENTS

We thank all members of the Aaron Grossberg, Robert Eil, and Katelyn Byrne, and Gaurav

Sahay labs for their helpful discussion and suggestions. We also acknowledge the expert

technical assistance by staff in the Advanced Multiscale Microscopy Shared Resource and

Histopathology Shared Resource. Author contributions are: Conceptualization, PCAW, AQB,

RE, AJG. Methodology, PCAW, AQB, NTVM, AJ, YE, GS, KTB, RE, AJG. Validation, PCAW,

AQB, HM, XZ, JD, MM, PRL, PD. Formal Analysis, PCAW, AQB, HM, XZ, JD, PD, GDS, AJG.

Investigation, PCAW, AQB, HM, XZ, JD, MM, PRL, PD, GDS. Writing—Original Draft, PCAW.

Writing – Review and Editing, PCAW, AQB, KTB, RE, AJG. Visualization, PCAW, AQB, PD.

Supervision, RE, AJG. Project Administration, RE, AJG. Funding Acquisition, PCAW, RE, AJG.

All authors approved this manuscript.

METHODS

Mouse Studies

Husbandry

C57BL/6J (WT, JAX 000664) mice were purchased from The Jackson Laboratory (Bar Harbor, ME) and maintained in our animal facility. All mice were housed and bred in a dedicated mouse room maintained at 26 °C, 40% humidity, and 12-h light/dark cycle. Mice were provided *ad libitum* access to food and water (5L0D, PicoLab) unless otherwise stated. All mice were 12 weeks of age at experiment start. Sex in each experiment is defined in the figure legends. When single housed, mice were allowed a 7-day acclimation period prior to procedure/study start. All tumor studies followed humane endpoints. All mice were humanely euthanized via cardiac puncture or cervical dislocation under deep isoflurane anesthesia. Mouse studies were conducted in accordance with the NIH Guide for the Care and Use of Laboratory animals, and approved by the Oregon Health & Science University IACUC.

Orthotopic Tumor Implantation

A vial of frozen KPC cells was thawed prior to each implantation, and 1 million cells were implanted in 23 uL of PBS per mouse. All mice were anesthetized with isoflurane, scrubbed with betadine, and a para-midline incision was made in the abdomen to expose pancreas. KPC cells or vehicle (PBS) were injected directly into the pancreatic parenchyma. Pancreas was placed back into position and incision was closed using two sutures (4-0 Polysorb) and two skin staples.

Subcutaneous Tumor Implantation

A vial of frozen KPC cells was thawed prior to each implantation, and 1 or 2 million cells were

implanted in 100 uL of PBS per mouse. The mouse's lower right abdomen was shaved, then

the needle was inserted near the right 4th mammary gland.

Antibody-based depletion

Mice were dosed intraperitoneally (IP) with either a combination of CD4 (BioXcell, #BE0003-1)

and CD8a (BioXcell, #BE0061) depletion antibodies, or Rat IgG2b isotype control antibody

(BioXcell, #BE0090), which were resuspended in InVivoPure ph7.0 Dilution Buffer (BioXcell,

#IP0070) per the manufacturer's instructions. First dose was 0.2 mg each antibody per mouse,

given IP two days prior to tumor implantation. Following doses were 0.1 mg each antibody per

mouse, given IP every 4 days after initial dose.

Lipid nanoparticle administration

LNPs were injected with a 30g insulin syringe intraperitoneally, following the dosing scheme

outlined in Figure 7A and D. LNPs were dosed based off of RNA content and were diluted with

calcium- and magnesium-free PBS. Empty LNP dose was calculated by matching the

cholesterol content in the empty and IL-6-containing LNPs.

Saphenous blood collection

We collected blood via saphenous bleeds during the LNP study. Mice were restrained by

scruffing while occluding the vein, and the hindlimb was wiped with petroleum jelly to expose

the vein. A 23G needle was used to gently prick the saphenous vein. Blood was collected in

EDTA-treated tubes, which were then placed on ice. Bleeding was stopped with pressure and

mice were returned to the cage. Each bleed alternated between left and right hindlimb.

Approximately 40 uL of blood was collected at each timepoint. Samples were spun down for 20

minutes at 2,000xg and plasma was stored at -80C.

Cell lines

Growth Conditions and validation

All cells were maintained at 37°C and 5% CO_2 in a humidified incubator, and tested negative in

house for mycoplasma using Universal Mycoplasma Detection Kit (30-1012K). *Kras*$^{G12D/+}$*,*

Tp53$^{R172H/+,}$ *Pdx1-Cre* (KPC) cell lines was generously shared by Dr. Elizabeth Jaffee

(KxPxCx) and Dr. David Tuveson (1199 and 1242)[18-20, 39]. KPC cells were grown on tissue

culture-treated dishes in growth media consisting of RPMI (Gibco) with 10% FBS (Corning)

and 1% penicillin/streptomycin (Gibco).

Engineered KPC

KPC cells expressing the surface marker Thy1.1 (CD90.1) with blasticidin resistance (BSR),

IL-6 with puromycin resistance, and luciferase with hygromycin resistance, were generated

from our stock of KPC cells (female) described above. IL-6 sequence was codon-optimized for

efficient expression (**Figure S9**). Platinum-E ecotropic packaging cells were transfected with

plasmid DNA encoding MSGV-Thy1.1, MSGV-IL6-Puro, or MSGV-Luciferase as described

previously[40]. Retroviral supernatants were spiked with 2ug/mL polybrene and were mixed 1:1

with fresh media before adding to 6-well tissue culture treated plates. Cells were spun at

2000g for 90min, 32C, no brake. Cells were then incubated at 37C for 48 hours before

washing off the viral supernatant and adding DMEM media (Gibco) supplemented with 10%

FBS (Corning). Two days later, KPC cells were placed in complete DMEM media containing

puromycin (5ug/mL) and/or blasticidin (5ug/mL) and/or hygromycin (500 ug/mL) to select for

transduced cells. Following antibiotic selection, successful transduction was confirmed via flow

cytometry staining for Thy1.1.   KxPxCx$^{CD90.1}$, KxPxCx^{IL6}, and KxPxCx^{LUC} cells were implanted

for OT-PDAC as described for parental KPC cells. Continued culturing of engineered cells was

done in selection media described.

Flow cytometry

Brefeldin A injections: For intracellular cytokine staining for flow cytometry, we followed

previously published protocols for golgi transport blockade[41]. Briefly, each mouse received

100ug Brefeldin A (Selleckchem) injected retro-orbitally 5 hours prior to tissue collection.

Sample preparation

We collected tumors from mice 5 days post implantation, and tumors were weighed, then

placed in PBS on ice. Tumor tissue was minced and digested (7). After dissociation, we

strained tumor suspension through at 100 um filter, and performed ACK lysis. We collected

spleens from mice at the endpoint specified. Spleens were pressed through a 70 um filter,

rinsed with PBS, pelleted at 1500 RPM for 5 minutes, and lysed with ACK lysis.

Staining

We stained samples with live/dead stain (1:2000) and surface protein antibodies (1:200 each),

and incubated for 20 minutes room temperature (**Table S3**). After staining, we washed

samples with FACS buffer and pelleted. For intracellular staining (Foxp3), we fixed and

permeabilized cells with 4% paraformaldehyde (BD Cytofix/Cytoperm), washed cells, and then

resuspended cells in antibody diluted 1:200 in permeabilization buffer (BD Perm/Wash) with

overnight incubation at 4C . The next day we washed and resuspended cells with FACS buffer

prior to analysis.

Instrumentation and analysis

All samples were analyzed in the OHSU Flow Cytometry Shared Resource using the Cytek

Aurora flow cytometer (Cytek Biosystems), data was analyzed in FlowJoTM v10.8.1. Samples

were first gated according to size and single cells, then all live cells were captured. The live

population was gated on CD45+ cells to capture all leukocytes. To identify T cells, leukocytes

were gated on CD90.2 then sub-gated for CD4+ or CD8+ cells. The CD4+ T cell population

was gated on Foxp3 to assess T regulatory cells. Natural killer (NK) cells were sub-gated from

the parental CD45+ gate. All NK1.1+ cells were captured, and then divided by expression of

CD90.2 to classify as NK (conventional, CD90.2-) or NK T cells (CD90.2+).

Histology

Pancreas/tumor tissue was fixed overnight in 4% PFA, then stored in 70% ethanol. Tissues

were paraffin embedded, sectioned, and hematoxylin and eosin (H&E) stained by the OHSU

Histopathology Shared Resource. Tumor tissue was sectioned in 5um slices at levels 50 um

apart. H&E stained tumor cross-sections were evaluated by a board-certified gastrointestinal

pathologist. All samples were blinded during sectioning/staining and during evaluation. 3-5

depths of tissue were qualitatively assessed per mouse. For day 5 tumor samples, 2 depths of

tissue were quantified per mouse and averaged together. For day 12 tumor samples, 1 section

was quantified.

Immunofluorescent staining and quantification

Pancreas/tumor tissue was dissected, transferred to BD Cytofix/Cytoperm diluted to 1% PFA,

and kept at 4C overnight. Tissue was then transferred to 30% sucrose in PBS and kept at 4C

overnight. Tissue was then washed twice in PBS before embedding in OCT media (Sakura).

Tissue was cut at 8 um onto superfrost plus slides and stored at -80C until staining. To stain,

slides were washed with PBS, blocked with 2.5% BSA 0.3% TritonX in PBS for one hour,

stained with pre-conjugated antibodies (**Table S3**) for 1h, washed with PBS, quenched with

TrueView Autofluorescence Quench kit (Vector, SP-8400) 2-5 min, stained with DAPI diluted

1:1000 for 10 min, and mounted with Vectashield Vibrance mounting medium (Vector, H-

1700). Whole tissue sections were imaged on a Zeiss Axio Scan 7 at 20x magnification. Two

tissue sections at least 344 um apart were assessed per sample. All images were blinded prior

to annotation and analysis. In QuPath (V0.5.1), areas of tumor, as defined as PanCK positive,

and stroma, as defined as PanCK negative abnormal tissue, were annotated. Adjacent tissue

sections were stained with H&E and used as reference for areas of stroma. Annotations were

made using only the DAPI and PanCK stains. After annotation, $CD3^+$ cells ($CD3^+$, $DAPI^+$,

PanCK-) were manually counted in stroma and tumor areas. Final counts were normalized to

total area of stroma or tumor annotation.

IVIS imaging

In vivo

Mice were injected with 100 uL of 15 mg/mL D-luciferin potassium salt in DPBS (no Ca, no Mg)

(GoldBio, #LUCK-100), then anesthetized with isoflurane. 10 minutes later, a luminescent

image and photo were captured. Longitudinal data was analyzed in one batch by normalizing

tumor ROI luminescence total counts to average background ROI luminescence total counts.

Ex vivo

Organs for analysis were placed in PBS on ice immediately following dissection. When all

organs were collected, PBS was replaced with 1 mL of 15 mg/mL D-Luciferin in DPBS (no Ca, no

Mg) (GoldBio, #LUCK-100). Organs were incubated for 5 minutes before removing from

solution, patting dry, and arranging on the IVIS imaging tray. Luminescent image and photo

were captured. Total luminescence counts were normalized to background luminescence

counts.

Lipid nanoparticle (LNP) preparation

LNPs were prepared using microfluidic mixing of ethanol (lipid) and aqueous (mRNA) phases

at a 3:1 volumetric ratio, with a total flow rate of 12 mL per minute using the Ignite system

(Precision Nanosystems, BC, Canada). The lipid phase contained ionizable lipid SM102,

cholesterol, DSPC, and DMG-PEG2000 at 50/10/38.5/1.5 molar ratios, respectively, diluted in

100% ethanol. mRNA was diluted in sterile 50 mM citrate buffer to achieve an N/P ratio of

5.67. For empty LNPs, an equivalent amount of citrate buffer was used. After aqueous and

lipid solutions were mixed, the LNPs were dialyzed for 4 hours at room temperature, followed

by an overnight dialysis against sterile PBS at 4C in 10 kDa Slide-a-Lyzer G3 cassettes

(ThermoFisher, MA, USA). Then, LNPs were concentrated by centrifugation using 100k

MWCO Amicon ultracentrifugal filters (Millipore Sigma, Burlington, MA) at 3000g and 4C. LNPs

were stored at 4C before administration.

LNP Characterization

Hydrodynamic size and polydispersity of the LNPs were determined with DLS using Stunner

(Unchained Labs, Pleasanton, CA). Encapsulation efficiency and concentration of mRNA were

determined using a modified Quant-iT RiboGreen RNA assay (ThermoFisher, MA, USA) and a

multimode microplate reader. Cholesterol content was determined with the Invitrogen™

Amplex™ Red Cholesterol Assay Kit (ThermoFisher, MA, USA).

Plasma analytes

Plasma was collected, snap frozen in liquid nitrogen, and stored at -80ºC. Plasma

concentrations of IL-6 (Biolegend) were measured using ELISA, and read on a plate reader

(BioTek).

Quantitative real-time polymerase chain reaction (qPCR)

We isolated RNA from cell pellets or tissue samples using the E.Z.N.A. Total RNA Kit I

(Omega BioTek) and we prepared cDNA using high-capacity cDNA reverse transcription kit

(Applied Biosystems). qPCR was run on the ABI 7300 (Applied Biosystems), using TaqMan

Fast Advanced PCR master mix (Applied Biosystems) or SYBR Green master mix (Applied

Biosystems). Relative expression was calculated using the $\Delta\Delta C_t$ method. To confirm the

presence/absence of *Il6 transgene* in tumor-implanted mice, we performed 30 cycles of qPCR

on the ABI7300, followed by running the PCR product on a 3% ethidium bromide gel to

determine the presence of a band at the expected size of 131 bp. Primers/probes are listed in

Table S2.

Statistical Analysis

Specific statistical tests and sample size for each study is indicated in the figure legends. Error

bars in figures show SEM. Statistical analyses were performed using GraphPad Prism (version

9; GraphPad Software Inc) or JMP Pro (version 16; SAS Institute Inc), and graphs were built

using GraphPad Prism (GraphPad Software Inc) statistical analysis software. P values are 2

sided with values less than 0.05 regarded as statistically significant.

Data Availability

All authors had access to the study data and had reviewed and approved the final manuscript.

All data will be available on the public repository, Mendeley Data, upon publication. Further

information and resources, including plasmid sequences, engineered KPC cells, and raw data

will be shared upon reasonable request to Aaron J. Grossberg (grossber@ohsu.edu).

Figure 1

Figure 1: OT-KxPxCx^{IL6} results in complete tumor clearance and body mass recovery.

(A) Survival comparison of OT-KxPxCx^{Parental} and OT-KxPxCx^{IL6}. N = 8 female, 8 male Sham and OT-KxPxCx^{IL6}, 1 female, 4 male OT-KxPxCx^{parental} mice. Statistically tested with Log-rank (Mantel-Cox) test. (B) Tumor mass at humane endpoint (OT-KxPxCx^{Parental}). Pancreas mass of OT-KxPxCx^{IL6} euthanized at a pre-determined endpoint of 24 days (did not meet humane euthanasia criteria); tumor tissue was not present. Statistically tested with 2-way ANOVA main effects only with Tukey correction for multiple comparisons. (C) Plasma IL-6 levels at the endpoints indicated in (H), Il-6 was below the assay detection limit for OT-KxPxCx^{IL6} mice. Statistically tested with 2-way ANOVA main effects only with Tukey correction for multiple comparisons. (A-C) N = 2 female, 3 male KxPxCx^{IL6}, 1 female, 4 male KxPxCx^{parental} mice. (D) Representative pancreas cross sections at 5 (left) and 10 days (right) from OT-KxPxCx^{parental} (top) and OT-KxPxCx^{IL6} (bottom) tumor implantation. Scale bars represent 50 um (top) and 100 um (bottom). (E) Tumor area as a percent of total tissue area at 5 and 12 days, quantified by board-certified pathologist. (F) Percentage of tumor area classified as poorly-differentiated, infiltrative carcinoma at 5 days by board-certified pathologist. (G) Tumor mass at 5 and 12 days. (B-D) N = 3 male, 3 female mice per group. (H) Expression of *Il6 transgene* expression, measured by qPCR at 5 (N = 3 male, 3 female), 10 (N = 6 male), 17 (N = 3 male, 2 female), 24 (N = 3 male, 2 female), and 40 (N = 16 male, 16 female) days. Statistically tested with One-way ANOVA with Tukey correction for multiple comparisons. (I) Plasma IL-6 measured by ELISA at 5 (N = 10 male KxPxCx^{Parental}, 11 male KxPxCx^{IL6}), 12(N = 3 male, 3 female KxPxCx^{Parental}, 3 male, 2 female KxPxCx^{IL6}), and 17 (N = 2 male, 3 female KxPxCx^{IL6}) days. Statistically tested with 2-way ANOVA main effects only with Tukey correction for multiple comparisons. (J) Body mass change as a percentage of initial body mass over time. N = 8

male, 8 female mice per group. Statistically tested with 3-way ANOVA. P<0.0001 for Time,

Sex, Tumor Status, Time x Sex, Time x Tumor Status. p=0.0002 for Sex x Tumor Status. p=

0.2900 for Time x Sex x Tumor Status. (K) Gastrocnemius muscle mass normalized to initial

body mass. (L) Atrophy-related gene expression (*Trim63* and *Fbxo32*) in gastrocnemius

muscle, measured by qPCR. (J-L) 5d N = 1 female, 3 male mice per group. 10d N = 6 male

PDAC, 5 male sham mice. Error bars represent SEM. Unless otherwise noted, 2x2 studies

were statistically tested with a full effects model 2-way ANOVA and Sidak multiple

comparisons test. 2-group analysis tested with unpaired t-test. **** P<0.0001, ***P<0.001,

**P<0.01, *P<0.05.

Figure 2

Figure 2: Tumor-cell IL-6 overexpression transduction controls. A) Survival comparison of

Sham, OT-1242[Parental] and OT-1242[IL6]. Statistically tested with Log-rank (Mantel-Cox) test on

pairwise comparisons. (B) Pancreas and tumor mass at humane endpoint. (C) Plasma IL-6

levels at humane endpoint. (D) Survival comparison of Sham, OT-1199 [Parental] and OT-1199[IL6].

Statistically tested with Log-rank (Mantel-Cox) test on pairwise comparisons. (E) Pancreas and

tumor mass at humane endpoint. (F) Plasma IL-6 levels at humane endpoint. (A-F) N= 5 male

mice per group. (G) Survival comparison of Sham, OT-KX2[Parental] and OT-KX2[IL6]. Statistically

tested with Log-rank (Mantel-Cox) test on pairwise comparisons. (H) Pancreas and tumor

mass at humane endpoint (OT-KX2[Parental]); 5 and 28 days (OT-KX2[IL6]). Statistically tested with

one-way ANOVA. (I) Plasma IL-6 levels at humane endpoint. (G-I) N = 3 male mice OT-KX2[IL6]

d5, 6 male mice OT-KX2[IL6] d28 and OT-KX2[parental]. (J-M) Control-transduced PDAC cells (OT-

KxPxCx[CD90.1]) do not spontaneously clear. (J) Tumor mass at 12 days post implantation. (K)

Tumor area as a percent of total tissue area at 12 days post implantation, quantified by board-

certified pathologist. (L) Plasma IL-6 measured by ELISA at 12 days post implantation. (M)

Body mass change as a percentage of initial body mass over time. Statistically tested with

one-way ANOVA analysis with repeated measures. $P<0.0001$ for Time and Time x Tumor

Status. $p=0.0002$ for Tumor Status. (J-M) N = 3 male, 3 female mice per group. Error bars

represent SEM. Unless otherwise noted, 2-group studies were statistically tested with unpaired

t-test. 3-group studies were statistically tested with 1-way ANOVA and Sidak multiple

comparisons test. **** $P<0.0001$, ***$P<0.001$, **$P<0.01$, *$P<0.05$.

Figure 3

Figure 3: Effects of tumor-specific IL-6 overexpression are maintained with lower tumor inoculation. (A) Survival comparison of OT-KxPxCx^{Parental} and OT-KxPxCx^{IL6} mice implanted with 5,000 cells. Statistically tested with Log-rank (Mantel-Cox) test on pairwise comparisons. (B) Pancreas and tumor mass at humane endpoint (OT-KxPxCx^{Parental}); 42 days (OT-KxPxCx^{IL6}). (C) Plasma IL-6 levels at endpoint. (A-C) N = 11 male mice (OT-KxPxCx^{IL6}), 8 male mice (OT-KxPxCx^{Parental}), and 5 male mice (sham). (D) Survival comparison of Sham, OT-1242^{Parental} and OT-1242^{IL6}. Statistically tested with Log-rank (Mantel-Cox) test on pairwise comparisons. (E) Pancreas and tumor mass at humane endpoint. (F) Plasma IL-6 levels at humane endpoint. (G) Survival comparison of Sham, OT-1199 ^{Parental} and OT-1199^{IL6}. Statistically tested with Log-rank (Mantel-Cox) test on pairwise comparisons. (H) Pancreas and tumor mass at humane endpoint. (I) Plasma IL-6 levels at humane endpoint. Error bars represent SEM. Unless otherwise noted, 2-group studies were statistically tested with unpaired t-test. 3-group studies were statistically tested with 1-way ANOVA and Sidak multiple comparisons test. **** $P<0.0001$, ***$P<0.001$, **$P<0.01$, *$P<0.05$.

Figure 4

Figure 4: OT-KxPxCx^{IL6} induces lymphocytic anti-tumor immune response. (A) Spleen

mass at 5 days. N = 6 male mice per group. (B-I) Intra-tumoral immune cell populations from

pancreas at 5 days: (B) Representative flow gating strategy for T cells. (C) CD8$^+$ T cells, (D)

CD4$^+$ T cells, (E) Foxp3$^+$ T regulatory cells, (F) CD62L$^+$ Naive T cells. (G) Ratio of Foxp3$^+$CD4$^+$

to CD8$^+$ cells, representative of a the degree of anti-tumor activity in the tumor

microenvironment. (H) NK1.1$^+$ Natural Killer cells, (I) NK1.1$^+$/CD90.2$^+$ NK T cells. (B-F) N = 6

male mice per group. (G-I) N = 4 female, 3 male mice per group. (J) representative images of

CD3$^+$ T cell infiltration in tumor and associated stroma of OT-KxPxCx^{parental} (top) and OT-

KxPxCx^{IL6} (bottom) mice. Tissues were stained with DAPI (blue), CD3 (magenta), and

Pancytokeratin (PanCK, green). (scale bars = 100 um). (K-L) CD3$^+$ cells counted per mm^2 of

stroma (K) or tumor nests (L) in immunofluorescence stained pancreas tissues. N = 3 male, 3

female mice per group. (M) Representative H&E image of lymphoid aggregates adjacent to

tumor and stroma in OT-KxPxCx^{IL6} pancreas (left) and magnified image of lymphoid aggregate

(right). Area of magnification is denoted on left image with orange box. (N) Pathologist-

evaluated number of tumor-associated lymphoid aggregates in H&E stained pancreas cross

sections from tissue collected at 5 days. N = 3 male, 3 female mice per group. Error bars

represent SEM. 2-group analysis tested with unpaired t-test. **** P<0.0001, ***P<0.001,

**P<0.01, *P<0.05.

Figure 5

Figure 5: CD4$^+$ and CD8$^+$ T cells are necessary for OT-KxPxCx^{IL6} tumor clearance. (A-C)

Confirmation of CD4/CD8a T cell depletion via flow cytometry on spleen 12 days post

implantation for (A)CD90.2+ total T cells, (B) CD8+ T cells, (C) CD4+ T cells. (D) Survival

comparison of OT-KxPxCx^{IL6} mice given CD4/CD8 depletion or IgG control antibodies.

Statistically tested with Log-rank (Mantel-Cox) test. (E) OT-KxPxCx$^{IL6-LUC}$ tumor growth,

measured by IVIS imaging. Statistically tested with 2-way ANOVA, for time points with

representation from both groups, with Šídák multiple testing correction. (F) Representative

images from IVIS imaging of OT-KxPxCx$^{IL6-LUC}$ 10 minutes after luciferin injection. Images are

of the same animals longitudinally. CD4/CD8 depleted mice reached humane endpoint prior to

D10 scan. (G) Pancreas and tumor mass at endpoint. (H) Representative pancreas cross

sections at humane euthanasia endpoint from IgG control (top) and CD4/CD8 depleted

(bottom) mice. Scale bars represent 100 um. (I) expression of *Il6 transgene* expression,

measured by qPCR. (J) Plasma IL-6 measured by ELISA. (K) Body mass change as a

percentage of initial body mass over time. Statistically tested with Mixed-effects analysis with

repeated measures. P<0.0001 for Time and Time x Tumor Status. p=0.0048 for Tumor Status.

(L) Gastrocnemius muscle mass at humane euthanasia endpoint normalized to initial body

mass. (M) Atrophy-related gene expression (*Trim63* and *Fbxo32*) in gastrocnemius muscle,

measured by qPCR. All male mice, N = 5 sham, 10 KxPxCx^{IL6} per antibody treatment group.

Error bars represent SEM. 3 group studies were statistically tested with a 1-way ANOVA and

Tukey correction for multiple comparisons. **** P<0.0001, ***P<0.001, **P<0.01, *P<0.05.

Figure 6

A

D0 — D76 — Monitor Survival

Implantation
OT-PDAC^{IL6}
or Sham

Implantation
OT-PDAC^{Luc}

B

Survival Post OT-PDAC^{IL6} Implantation

OT-KxPxCx^{Luc}
Implantation

p<0.0001

OT-KxPxCx^{IL6}
Sham

Probability of Survival

Days Post Implantation

C

Tumor Burden

Sham-recovered   KxPxCx^{IL6}-recovered

Luminescence Counts
(background subtracted)

Days Post Implantation OT-PDAC^{Luc}

*   ***   ****

D

**Sham-recovered, OT-PDAC^{Luc}
Terminal Tumor Mass**

Tumor mass (g)

E

Delta Body Mass

Days Post Implantation OT-KxPxCx^{Luc}

Delta Body Mass (% D0)

Sham-recovered   OT-KxPxCx^{IL6} Recovered

Sham recovered   OT-KxPxCx^{IL6} recovered

D2

D5

D7

D12

D21   N.D.

F

D0 — D28 — D40
D26
Antibody Depletion Start

Implantation
OT-PDAC^{IL6}

Rechallenge
Sham or
OT-PDAC^{Parental}

G

Tumor burden 12d post rechallenge

IgG Control   CD4/CD8 Depleted

Pancreas + Tumor Mass (g)

****   ****   ****

Sham   OT-KxPxCx
Rechallenge Status

H

Spleen: CD4 T Cells

IgG Control   CD4/CD8 Depleted

CD4+ (%CD45)

****   ****

Sham   OT-KxPxCx
Rechallenge Status

I

Spleen: CD8 T Cells

IgG Control   CD4/CD8 Depleted

CD8+ (%CD45)

****   ****

Sham   OT-KxPxCx
Rechallenge Status

Figure 6: OT-KxPxCx^{IL6} induces a durable T cell response to OT-KxPxCx^{Parental} tumors.

(A) Schematic timeline for B-E. Mice were implanted with OT-KxPxCx^{IL6} or given sham

surgery, then all mice were rechallenged with OT-KxPxCx$^{Parental-LUC}$ after 76 days. (B) Survival

comparison of OT-KxPxCx^{IL6}-recovered and sham-recovered mice. Statistically tested with

Log-rank (Mantel-Cox) test. (C) OT-KxPxCx$^{Parental-LUC}$ tumor growth, measured by IVIS

imaging. Statistically tested with mixed effects model and Šídák multiple testing correction,

using imputation to match KxPxCx^{IL6} endpoint with the sham endpoint. Corresponding IVIS

images below. (D) Tumor mass at humane euthanasia endpoint for sham-recovered OT-

KxPxCx$^{Parental-LUC}$ tumors. (E) Body mass change as a percentage of initial body mass over

time. Statistically tested with mixed effects model and Šídák multiple testing correction at

timepoints where both groups were represented. p=0.0435 for Time, p=0.4903 for Tumor

Status, p=0.0002 for Time x Tumor Status. (A-E) N = 7 male KxPxCx^{IL6}, 8 male sham, 7

female KxPxCx^{IL6}, 6 female sham. (F) Schematic timeline for G-I. Mice were implanted with

OT-KxPxCx^{IL6} or given Sham surgery. 28 days later, mice were rechallenged with OT-

KxPxCx^{Parental}, and given CD4/CD8 depletion or IgG control antibodies beginning on D26 and

every 4 days thereafter (denoted with double green lines). (G) Pancreas/tumor mass at 12

days. (H) Intra-tumoral CD4$^+$ T cells at 12 days. (I) Intra-tumoral CD8$^+$ T cells at 12 days. (F-I)

All male mice N = 6 per group. Error bars represent SEM. 2x2 studies were statistically tested

with a full effects model 2-way ANOVA and Sidak multiple comparisons test. **** P<0.0001,

***P<0.001, **P<0.01, *P<0.05.

Figure 7

Figure 7: Lipid nanoparticle delivery of IL-6 mRNA replicates the anti-tumor effects observed in OT-PDAC[IL6]. (A) Schematic of luciferase LNP validation. Mice were injected with LNPs containing a total of 2.5 ug luciferase mRNA 6 hours prior to IVIS imaging, euthanasia, tissue collection, and tissue IVIS imaging. N = 3 male mice per group. (B) left- IVIS images of mice injected intraperitoneally with LNPs containing luciferase. Right – IVIS image of the spleen, liver, and pancreas, incubated ex vivo with luciferin prior to imaging. (C) Comparison of total luminescence counts in the pancreas and liver. Statistically tested with one-way ANOVA and Tukey multiple testing correction. (D) Schematic timeline for study presented in D-N. Mice were injected with LNP dose 1, 3 hours prior to tumor implantation. Mice were then injected with subsequent doses on 2, 4, and 6 days post-implantation. Blood was collected by saphenous bleed 1, 3, 5, and 7 days post-implantation. Mice were euthanized on day 8. N = 7 (No LNP), 10 (empty LNP), 8 (IL6 LNP). All mice were male. Error bars represent SEM. (E) Body mass change as a percentage of initial body mass over time. Statistically tested with two-way ANOVA P<0.0001 for Time, Treatment, and Time x Treatment. (F) Plasma IL-6 measured by ELISA at 1, 3, 5, and 7 days post-implantation, and at euthanasia (D8). Statistically tested with repeated measures mixed-effects model with Tukey correction for multiple testing. (G) Tumor and pancreas mass. (H) Spleen mass. (I-N) Intra-tumoral immune cell populations from pancreas: (I) CD8+ T cells, (J) CD4+ T cells, (K) Foxp3+ T regulatory cells, (L) ratio of Foxp3+ to CD8+ cells, representative of a the degree of anti-tumor activity in the tumor microenvironment, (M) NK1.1+ NK cells, (N) NK1.1+/CD90.2+ NK T cells. 3-group analysis tested with one-way ANOVA with Tukey correction for multiple testing. **** P<0.0001, ***P<0.001, **P<0.01, *P<0.05.

SUPPLEMENTAL INFORMATION

Table S1: Histological assessment of transgenic tumors.

		Tumor Area (% total area)	Carcinoma Area (% tumor area)	Lymphoid Aggregates	Neutrophil Presence (Incidence)		
					<5% Tumor	5-50% Tumor	>50% Tumor
5 days	KxPxCx[Parental]	18.75 ± 8.18	45.00 ± 13.78	2.75 ± 1.73	-	10	2
	KxPxCx[IL6]	12.58 ± 6.79	**19.83 ± 9.96**	**12.92 ± 5.19**	-	-	12
12 days	KxPxCx[Parental]	82.83 ± 14.43	95.83 ± 3.06	6.50 ± 3.45	4	2	-
	KxPxCx[IL6]	**0**	NA	NA	NA		
	KxPxCx[CD90.1]	75.00 ± 17.32	91.67 ± 5.85	3.50 ± 3.89	2	4	-

Values represented as mean ± SD. Bold indicates $p<0.05$ when compared to KxPxCx[Parental] .
Day 5: Two sections per sample evaluated for neutrophil presence.

Figure S1: OT-KxPxCx^{IL6} clearance detected by *Il6-transgene* qPCR product. qPCR products from pancreas/tumor tissue, run for 30 PCR cycles, and then run out on a 3% agarose gel. Expected product is 131 bp. Ladder is Promega 100 bp ladder (G210A). (A) 5 days post implantation. (B) 17 days post implantation (samples 7, 9, 11, 16) and 24 days post implantation (samples 3, 4, 8, 10, 14). (C)10 days post implantation. (D) 40 days post implantation.

Figure S2: KxPxCx^{IL6} metastatic cell presence detected by *Il6-transgene* qPCR product. qPCR products, run for 30 PCR cycles, and then run out on a 3% agarose gel. Expected product is 131 bp. Ladder is Promega 100 bp ladder (G210A). (A) Whole liver at 5 days post implantation. (B) Whole lung at 5 days post implantation. (C) Whole lung and liver at 12 days post implantation. (D) Quantification.

Figure S3: KxPxCx[IL6] cells induce anti-tumor immune response when implanted subcutaneously. (A) Tumor volume measured by calipers daily. (B) Injection site/tumor tissue collected at 24 days post implantation qPCR products, run for 30 PCR cycles, and then run out on a 3% agarose gel. Expected product is 131 bp. Ladder is Promega 100 bp ladder (G210A). (C) Plasma IL-6 measured by ELISA at 24 days post implantation.

Figure S4: Gating strategy for flow cytometry presented in Figure 4B-G, and Figure S6 B-E, G (representative sample).

Figure S5: Gating strategies for flow cytometry presented in Figure 4H-I, and Figure S6A, F (representative sample).

Figure S6: Supporting flow cytometry data for Th17 and Granzyme B+ T cell phenotypes. Intra-tumoral immune cell populations from pancreas at 5 days: (A) Granzyme B+ cells as a percent of CD8+ cells (left) and CD8+ Granzyme B MFI (right). (B) Th17 Cells (RORγT+) as a percent of CD4+ cells (left) and CD4+ RORγT MFI (right). Immune cell populations from spleen at 5 days: (C) CD4+ T cells, (D) CD8+ T cells, (E) Ratio of Foxp3+ to CD8+ cells, (F) Granzyme B+ cells as a percent of CD8+ cells (left) and CD8+ Granzyme B MFI (right). (G) Th17 Cells (RORγT+) as a percent of CD4+ cells (left) and CD4+ RORγT MFI (right). (B-E, G) N = 6 male mice per group. (A, F) N = 4 female, 3 male mice per group. Error bars represent SEM. 2-group analysis tested with unpaired t-test. **** p<0.0001, ***p<0.001, **p<0.01, *p<0.05.

CD8+ Cells

Figure S7: Granzyme B gating strategy for flow cytometry presented in Figure S6 (representative samples).

Figure S8: Gating strategies for flow cytometry presented in Figure 7 (representative sample).

Figure S9: Linear sequence map with amino acid annotation for codon-optimized IL6 overexpression cassette transduced into PDAC cells.

Table S2: qPCR primer and probe information

Gene Name	Supplier	Cat. Number
18S	Thermo	4333760F
Fbxo32	IDT	Mm.PT.58.7025875
Trim63	IDT	Mm.PT.58.32840172
Il6-transgene	IDT	F: CGTGGAGATGAGGAAGGAGC R: GGTTGTAGCCGGTCTGGTAG

Table S3: Flow cytometry antibody information

Protein (clone)	Host	Target	Supplier	Cat. Number	Use
CD3 (17A2)	Rat	Mouse	Biolegend	100209	Tissue staining
CD4 (RM4-5)	Rat	Mouse	BD Pharmingen	561115	Flow
CD45.2 (104)	Mouse	Mouse	BD Horizon	563686	Flow
CD62L (MEL-14)	Rat	Mouse	BD Pharmingen	5**605**14	Flow
CD8 (53-6.7)	Rat	Mouse	BD Pharmingen	557959	Flow
CD90.2 (53-2.1)	Mouse	Mouse	BD Pharmingen	553004	Flow
FoxP3 (FJK-16s)	Rat	Mouse	eBioscience (ThermoFisher)	12-5773-82	Flow
Granzyme B (GB11)	Mouse	Mouse	Biolegend	515403	Flow
NK1.1 (PK136)	Mouse	Mouse	BD Pharmingen	552878	Flow
PanCK (AE1/AE3)	Mouse	Mouse	eBioscience (ThermoFisher)	53-9003-82	Tissue staining
RORγT (AFKJS-9)	Rat	Mouse	eBioscience (ThermoFisher)	17-6988-82	Flow

REFERENCES

1.   Winograd R, Byrne KT, Evans RA, et al. Induction of T-cell immunity overcomes

complete resistance to PD-1 and CTLA-4 blockade and improves survival in pancreatic

carcinoma. Cancer immunology research 2015;3:399-411.

2.   Beatty GL, Eghbali S, Kim R. Deploying immunotherapy in pancreatic cancer: defining

mechanisms of response and resistance. American Society of Clinical Oncology

Educational Book 2017;37:267-278.

3.   Aznar MA, Good CR, Barber-Rotenberg JS, et al. Clinical and molecular dissection of

CAR T cell resistance in pancreatic cancer. Cell Reports Medicine 2025.

4.   Byrne KT, Vonderheide RH. CD40 stimulation obviates innate sensors and drives T cell

immunity in cancer. Cell reports 2016;15:2719-2732.

5.   Gnerlich JL, Mitchem JB, Weir JS, et al. Induction of Th17 cells in the tumor

microenvironment improves survival in a murine model of pancreatic cancer. The

Journal of Immunology 2010;185:4063-4071.

6.   Delvecchio FR, Fincham RE, Spear S, et al. Pancreatic cancer chemotherapy is

potentiated by induction of tertiary lymphoid structures in mice. Cellular and Molecular

Gastroenterology and Hepatology 2021;12:1543-1565.

7.   Hotz C, Wagenaar TR, Gieseke F, et al. Local delivery of mRNA-encoded cytokines

promotes antitumor immunity and tumor eradication across multiple preclinical tumor

models. Science translational medicine 2021;13:eabc7804.

8.   Corcoran RB, Contino G, Deshpande V, et al. STAT3 plays a critical role in KRAS-

induced pancreatic tumorigenesis. Cancer research 2011;71:5020-5029.

9.   Suh S-Y, Choi YS, Yeom CH, et al. Interleukin-6 but not tumour necrosis factor-alpha

predicts survival in patients with advanced cancer. Supportive Care in Cancer

2013;21:3071-3077.

10.   Rupert JE, Narasimhan A, Jengelley DH, et al. Tumor-derived IL-6 and trans-signaling

among tumor, fat, and muscle mediate pancreatic cancer cachexia. Journal of

Experimental Medicine 2021;218:e20190450.

11.   Razidlo GL, Burton KM, McNiven MA. Interleukin-6 promotes pancreatic cancer cell

migration by rapidly activating the small GTPase CDC42. Journal of Biological

Chemistry 2018;293:11143-11153.

12.   Arneson-Wissink PC, Mendez H, Pelz K, et al. Hepatic signal transducer and activator

of transcription-3 signalling drives early-stage pancreatic cancer cachexia via

suppressed ketogenesis. Journal of Cachexia, Sarcopenia and Muscle 2024.

13.   Flint TR, Janowitz T, Connell CM, et al. Tumor-induced IL-6 reprograms host

metabolism to suppress anti-tumor immunity. Cell metabolism 2016;24:672-684.

14.   Zhang Y, Yan W, Collins MA, et al. Interleukin-6 is required for pancreatic cancer

progression by promoting MAPK signaling activation and oxidative stress resistance.

Cancer research 2013;73:6359-6374.

15.   Fujimoto M, Nakano M, Terabe F, et al. The influence of excessive IL-6 production in

vivo on the development and function of Foxp3+ regulatory T cells. The Journal of

Immunology 2011;186:32-40.

16.   Jiang Z, Liao R, Lv J, et al. IL-6 trans-signaling promotes the expansion and anti-tumor

activity of CAR T cells. Leukemia 2021;35:1380-1391.

17.   Nejad EB, Labrie C, van Elsas MJ, et al. IL-6 signaling in macrophages is required for

immunotherapy-driven regression of tumors. Journal for ImmunoTherapy of Cancer

2021;9:e002460.

18.   Foley K, Rucki AA, Xiao Q, et al. Semaphorin 3D autocrine signaling mediates the

metastatic role of annexin A2 in pancreatic cancer. Science signaling 2015;8:ra77-ra77.

19.   Quaranta V, Rainer C, Nielsen SR, et al. Macrophage-derived granulin drives resistance

to immune checkpoint inhibition in metastatic pancreatic cancer. Cancer research

2018;78:4253-4269.

20.   Ireland L, Santos A, Ahmed MS, et al. Chemoresistance in pancreatic cancer is driven

by stroma-derived insulin-like growth factors. Cancer research 2016;76:6851-6863.

21.   Ino Y, Yamazaki-Itoh R, Shimada K, et al. Immune cell infiltration as an indicator of the

immune microenvironment of pancreatic cancer. British journal of cancer 2013;108:914-

923.

22.   Hiraoka N, Ino Y, Yamazaki-Itoh R, et al. Intratumoral tertiary lymphoid organ is a

favourable prognosticator in patients with pancreatic cancer. British journal of cancer

2015;112:1782-1790.

23.   Sautès-Fridman C, Petitprez F, Calderaro J, et al. Tertiary lymphoid structures in the

era of cancer immunotherapy. Nature Reviews Cancer 2019;19:307-325.

24.   Siegel RL, Miller KD, Wagle NS, et al. Cancer statistics, 2023. CA: a cancer journal for

clinicians 2023;73:17-48.

25.   Hilmi M, Delaye M, Muzzolini M, et al. The immunological landscape in pancreatic

ductal adenocarcinoma and overcoming resistance to immunotherapy. The Lancet

Gastroenterology & Hepatology 2023.

26.   Mace TA, Shakya R, Pitarresi JR, et al. IL-6 and PD-L1 antibody blockade combination

therapy reduces tumour progression in murine models of pancreatic cancer. Gut

2018;67:320-332.

27.   Ware MB, Phillips M, McQuinn C, et al. Dual IL-6 and CTLA-4 blockade regresses

pancreatic tumors in a T cell–and CXCR3-dependent manner. JCI insight

2023;8:e155006.

28.   Arneson-Wissink PC, Mendez H, Pelz K, et al. Hepatic signal transducer and activator

of transcription-3 signalling drives early-stage pancreatic cancer cachexia via

suppressed ketogenesis. J Cachexia Sarcopenia Muscle 2024;15:975-988.

29.   Soler MF, Abaurrea A, Azcoaga P, et al. New perspectives in cancer immunotherapy:

targeting IL-6 cytokine family. Journal for ImmunoTherapy of Cancer 2023;11.

30.   Lee JW, Stone ML, Porrett PM, et al. Hepatocytes direct the formation of a pro-

metastatic niche in the liver. Nature 2019;567:249-252.

31.   Robinson E, MacDonald KD, Slaughter K, et al. Lipid nanoparticle-delivered chemically

modified mRNA restores chloride secretion in cystic fibrosis. Molecular Therapy

2018;26:2034-2046.

32.   Lokugamage MP, Vanover D, Beyersdorf J, et al. Optimization of lipid nanoparticles for

the delivery of nebulized therapeutic mRNA to the lungs. Nature biomedical engineering

2021;5:1059-1068.

33.   Flamini V, Ghadiali RS, Antczak P, et al. The satellite cell niche regulates the balance

between myoblast differentiation and self-renewal via p53. Stem Cell Reports

2018;10:970-983.

34.   Melamed JR, Yerneni SS, Arral ML, et al. Ionizable lipid nanoparticles deliver mRNA to

830   pancreatic β cells via macrophage-mediated gene transfer. Science Advances

831   2023;9:eade1444.

832   35.   Zhu Q, Yu C, Chen Y, et al. Dual mRNA nanoparticles strategy for enhanced pancreatic

833   cancer treatment and β-elemene combination therapy. Proceedings of the National

834   Academy of Sciences 2025;122:e2418306122.

835   36.   Sahel DK, Renner J, Vlasova KY, et al. Improved localized mRNA delivery using lipid

836   nanoparticles with a novel synthetic cholesterol derivative. RSC pharmaceutics 2026.

837   37.   Qiu X, Yang Y, Ma Q, et al. Thioketal-Incorporated Biodegradable Lipid Nanoparticles

838   Deliver mRNA to Ductal Epithelial Cells for Improved Pancreatic Cancer Treatment.

839   Journal of the American Chemical Society 2025.

840   38.   Shen Q, Liu J, Zeng L, et al. Pancreas-targeted lipid nanoparticles for relatively non-

841   invasive interleukin-12 mRNA therapy in orthotopic pancreatic ductal adenocarcinoma.

842   Journal of Controlled Release 2025;381:113588.

843   39.   Michaelis KA, Zhu X, Burfeind KG, et al. Establishment and characterization of a novel

844   murine model of pancreatic cancer cachexia. Journal of cachexia, sarcopenia and

845   muscle 2017;8:824-838.

846   40.   Eil R, Vodnala SK, Clever D, et al. Ionic immune suppression within the tumour

847   microenvironment limits T cell effector function. Nature 2016;537:539-543.

848   41.   Liu F, Whitton JL. Cutting edge: re-evaluating the in vivo cytokine responses of CD8+ T

849   cells during primary and secondary viral infections. The Journal of Immunology

850   2005;174:5936-5940.

Figure 1

A Survival Post OT-PDAC Implantation

Probability of Survival / Days Post Implantation

OT-KxPxCx^Parental — OT-KxPxCx^IL6 — Sham

B Tumor Mass

Pancreas + Tumor Mass (g) / ****

C Plasma IL-6 ELISA

Plasma [IL-6] pg/mL / *

OT-PDAC^Parental   OT-PDAC^IL6

D 5d   12d

OT-KxPxCx^parental

OT-KxPxCx^IL6

E H&E Tumor Area

Tumor Area (% total area) / Days Post Implantation

F Poorly Differentiated Infiltrative Carcinoma (D5)

Carcinoma Area (% tumor area) / **

G Tumor mass

Resected Tumor Mass (g) / Days Post Implantation

H Pancreas IL6 Transgene

2^-dCT / Days Post Implantation

I Plasma IL-6 ELISA

Plasma [IL-6] pg/mL / Days Post Implantation

J Body mass change (% initial) / Days Post Implantation

Sham (female) — Sham (male) — OT-KxPxCx^IL6 (female) — OT-KxPxCx^IL6 (male)

K Gastrocnemius Mass

Gastroc Mass (mg/g BW, D0) / Days Post Implantation

L Gastrocnemius Gene Expression

mRNA expression (fold change) / Trim63   Fbxo32 / Days Post Implantation

Sham   OT-KxPxCx^IL6

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7