**Abundance measurements reveal the balance between lysis and lysogeny in the human gut microbiome**

Jaime Lopez[1,2], Saria McKeithen-Mead[1,3], Handuo Shi[3], Taylor H. Nguyen[1], Kerwyn Casey Huang[1,3,4,†], Benjamin H. Good[2,4,5,†]

[1]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[2]Department of Applied Physics, Stanford University, Stanford, CA 94305, USA

[3]Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA

[4]Chan Zuckerberg Biohub, San Francisco, CA 94158, USA

[5]Department of Biology, Stanford University School of Medicine, Stanford, CA 94305, USA

[†]Correspondence: kchuang@stanford.edu (K.C.H.) and bhgood@stanford.edu (B.H.G.).

**Abstract**

The human gut contains diverse communities of bacteriophage, whose interactions with the broader microbiome and potential roles in human health are only beginning to be uncovered. Here, we combine multiple types of data to quantitatively estimate gut phage population dynamics and lifestyle characteristics in human subjects. Unifying results from previous studies, we show that an average human gut contains a low ratio of phage particles to bacterial cells (~1:100), but a much larger ratio of phage genomes to bacterial genomes (~4:1), implying that most gut phage are effectively temperate (e.g., integrated prophage, phage-plasmids, etc.). By integrating imaging and sequencing data with a generalized model of temperate phage dynamics, we estimate that phage induction and lysis occurs at a low average rate (~0.001-0.01 per bacterium per day), imposing only a modest fitness burden on their bacterial hosts. Consistent with these estimates, we find that the phage composition of a diverse synthetic community in gnotobiotic mice can be quantitatively predicted from bacterial abundances alone, while still exhibiting phage diversity comparable to native human microbiomes. These results provide a foundation for interpreting existing and future studies on links between the gut virome and human health.

**Introduction**

The human gut harbors a complex community of bacteria, viruses, and microbial eukaryotes that plays important roles in human health (1–3). Previous studies have largely focused on the bacterial portion of this community, but in recent years the bacteriophage ("phage") that infect these bacteria have started to draw more attention. Advances in DNA sequencing and anaerobic culturing have led to extensive databases of gut phage genomes (4,5), as well as increasing numbers of phage isolates that can be propagated in the lab for mechanistic investigation (6,7). Phage can influence the microbiome in multiple ways. They can directly kill their bacterial hosts through lytic infection (7,8) or by inducing lysis from a temperate state (8,9). Temperate phage can also serve as important vectors of horizontal gene transfer (10), carrying cargo genes that enhance the metabolic or defense capabilities of their bacterial hosts (11). These interactions with gut bacterial ecology and evolution have been hypothesized to impact human health. Cohort studies have revealed numerous associations between the composition of the gut virome and various health-related states, including cancer treatment efficacy (2) and lifespan (12). Transplants of sterile phage-containing fecal filtrates from healthy donors can help resolve and protect against infections (13,14) or exacerbate disease phenotypes (15), phenomena potentially mediated by bacteria-phage interactions. Phage particles can also interact directly with the human immune system (16). These results suggest that quantitative characterization of gut phage communities is likely critical for understanding and engineering the gut microbiome.

However, while the individual members of the gut virome are becoming increasingly well

characterized, much less is known about their ecological dynamics within a typical

human and the effects they exert on the surrounding microbial community. In marine

ecosystems, phage particles outnumber bacteria ~10:1 (17,18) and are estimated to kill

~20% of the bacterial population each day (18). Such high rates of lysis generate strong

selection pressures for both bacteria and phage, leading to antagonistic co-evolution

(19) and "kill-the-winner" dynamics of strain turnover (20–22). By contrast, estimates of

the virus-to-microbe ratio (VMR) in the human gut vary widely across studies, from

greater than 1:1 (23,24) to less than 1:10 (25,26). Furthermore, while some studies

have suggested that the gut microbiome is dominated by temperate phage (8,27), little

is known about rates of induction and lysis, and other studies have suggested that

evasion of phage-mediated lysis is a major driver of bacterial evolution within human

hosts (11,28). Inferring these ecological parameters is particularly challenging in the

complex setting of the human gut, as it requires linking existing measurement

approaches with quantitative models of phage population dynamics.

Here, we address this gap by combining mathematical modeling and publicly available

data to obtain quantitative baseline estimates of gut viral populations sizes and

induction rates in human hosts. Using a meta-analysis of gut viral population size

measurements, we show that existing data can be unified into a coherent quantitative

picture in which the gut microbiome has more phage genomes than bacterial genomes,

but many fewer phage particles than bacterial cells. This suggests that the gut is

dominated by temperate phage (here, "temperate" refers to all phage that reproduce

with their host genome, including classical lysogens, phage-plasmids, etc.). Building on

this quantification, we develop a modeling framework that enables inference of mean

gut phage induction rates from microscopy and metagenomic measurements. Our

findings suggest that, in typical adults, gut phage are rarely induced and place a low

mean fitness burden on their bacterial hosts. Finally, we show that similar ecological

dynamics arise in gnotobiotic mice colonized with a synthetic community of >100 human

gut bacterial isolates. As expected for a microbiome dominated by temperate phage, we

find that the virome composition of these mice can be quantitatively predicted from the

bacterial composition alone, while still exhibiting viral diversity comparable to a typical

human stool microbiome. These results suggest that existing methods for predicting gut

phage lifestyles drastically overestimate the fraction of lytic phage, indicating that many

gut phage contain yet uncharacterized host-association genes

**Results**

**The typical human gut microbiome contains fewer phage particles than bacterial cells**

To determine the range of phage population sizes and virus-to-microbe ratios (VMRs) in the gut, we compiled measurements across multiple methodologies and studies (**Table 1**). Although the VMRs, initially appeared to vary across studies, we found that they could be unified into a coherent quantitative picture by employing a consistent calculation approach that accounts for key differences among existing phage quantification techniques (**Fig. 1A**, **Methods**).

Many approaches for estimating phage abundance involve the isolation of virus-like particles (VLPs) as representatives of the free phage particle population within stool samples. In the most common method, isolated VLPs are enumerated via epifluorescence microscopy using a DNA-binding dye (23,29). These microscopy-based methods estimate the concentration of free phage particles in the stool, although their accuracy is constrained by VLP isolation efficiency (23,30) and the presence of non-phage particles (31). Aggregating VLP enumeration data from multiple studies and age groups (**Fig. 1B**), we found that, apart from newborns in which VLP densities are often below the limit of detection (23), stool VLP density stabilizes after >1 month of age to a population average of $\sim 2\times10^9$ VLPs/g stool, which is maintained throughout adulthood. Combining these data with existing estimates of the density of microbial cells in stool in humans older than >1 month ($\sim 10^{11}$ cells/g stool; (32)) yields a VLP-to-microbe ratio

$\sim 10^{-2}$. This estimate is three orders of magnitude lower than the VMRs commonly

reported for surface seawater systems (20), hinting qualitatively different viral ecological

dynamics that we will explore in more detail below. We also find that the inter-individual

variation in VLP counts is similar in magnitude to that of bacterial counts, with post-

infancy VLP measurements exhibiting a population coefficient of variation (CV) of 0.61

versus 0.46 for bacterial counts (32). This suggests that the total gut phage population

does not undergo dramatic abundance fluctuations across hosts.

An alternative form of phage particle measurement utilizes a spike-in approach,

involving shotgun sequencing of amplified DNA from the VLP pool after adding a known

amount of a non-gut reference phage (25). The fraction of sequencing reads mapping to

reference versus non-reference phage can then be used to obtain an independent

estimate of absolute phage particle density (**Fig. 1A**). A recent application of this

approach to longitudinal samples from ~10 healthy adults yielded a ~5-fold higher

concentration than microscopy-based studies (mean of ~1 × $10^{10}$ VLPs/g, inter-

individual CV of 0.9, **Fig. 1B**). These data also provided an estimate of the temporal

variation, with monthly VLP estimates within individuals having a mean CV of 0.78,

suggesting that the total phage load in individual hosts does not undergo dramatic

fluctuations. The differences between this study and microscopy-based quantifications

may be due to underestimation of viral counts by imaging-based approaches relative to

sequencing/qPCR-based approaches (33). However, we found that the two

measurements are largely consistent after exclusion of reads mapping to the

*Microviridae* family of phage (**Fig. 1B**), which are thought to be disproportionately

enriched by the multiple displacement amplification (MDA) protocol commonly

employed in VLP sequencing (34). Regardless, even the larger VMR estimates resulting

from sequencing-based approaches including *Microviridae* reads ($\sim 10^{-1}$) are still far

lower than the ~10:1 ratios reported for surface seawater (20).

**The typical human gut microbiome contains more phage genomes than phage**

**particles**

A third class of quantification methods estimates VMRs directly from metagenomic

sequencing of stool samples (24). This approach has been enabled by the recent

assembly of large databases of viral and prokaryotic genomes from the human gut

(4,5,35), from which >98% of reads from a typical stool sample can be classified using

taxonomic profilers like Phanta (24). By normalizing the ratio of phage to bacterial reads

with corresponding phage and bacterial genome lengths, one can obtain an

independent estimate of the VMR. Applying this approach to a collection of 255

previously sequenced adult gut metagenomes yields an average VMR of ~4:1 (inter-

individual CV = 0.38), corresponding to an absolute density of $\sim 4 \times 10^{11}$ phage

genomes/g after multiplying by the typical bacterial density in Ref. (32). These values

are two orders of magnitude higher than the VLP-based estimates above.

The discrepancy between these estimates can be reconciled by the observation that

bulk stool metagenomics measures the total number of viral genomes in a stool sample,

including those encapsulated in bacterial cells (e.g., as prophage), while VLP-based

methods only measure free viral particles. Hence, it is useful to distinguish between two

distinct abundance measures: the genomic VMR (gVMR), estimated from bulk stool

sequencing, and the particle VMR (pVMR), estimated from VLP-based approaches. The

two measures are roughly equivalent in environments like surface seawater where the

pVMR is much larger than one (and therefore particles dominate the gVMR). However,

they can dramatically diverge in ecosystems like the gut where the pVMR is much less

than one. In this case, the ~100-fold difference between the number of phage particles

and phage genomes in the gut suggests that the vast majority of gut phage are

temperate or otherwise attached to their bacterial hosts. These temperate phage may

not be traditional prophage that are integrated into their host's genome; many gut phage

do not contain recognizable lysogeny-associated genes and thus may utilize other host-

associated lifestyles, such as those of phage-plasmids (36).

Consistent with this temperate-dominated picture, we found that the ratio of phage

genomes to phage particles is also large for many individual viral species. While MDA

amplification biases make precise quantification difficult (34), comparisons between

matched VLP and bulk sequencing in infants and adults revealed three broad classes of

behavior. Some viral species are observed in the bulk metagenome but not in the

associated VLP pool (**Fig. 1C, bottom**). These species account for about half of the

total phage abundance in bulk stool samples (**Fig. 1D**) and could represent cryptic (37)

or inactive (27) prophage, as well as phage that are poorly amplified by MDA (34). A

second set of viral species are observed in VLP sequencing but not in the associated

bulk metagenome (**Fig. 1C, left**). These species account for about half of the total

phage abundance in the VLP pool, and could reflect both MDA amplification biases (34)

as well as phage that are poorly captured by bulk metagenomics. Finally, a third class of

viral species is present in both the VLP and bulk metagenomes. Their abundances are

broadly consistent with the aggregate particle to genome ratio above, with ~80% of

phage-sample pairs having a ratio below 1:100 (**Fig. 1C, Fig. S1**), and contain a mixture

of phage species classified as temperate and purely lytic (**Fig. S2**). These species

account for the other half of the VLP and bulk phage populations (**Fig. 1D**). These data

suggest that even with limitations imposed by MDA biases, a large fraction of gut phage

exhibit a generalized form of temperance, with a small population of viral particles

maintained by a much larger number of host-associated viral genomes.

**The phage particle to phage genome ratio provides a lower bound on the rate of**

**phage induction**

While population sizes and lifestyles are important aspects of gut ecology, they provide

only a static picture of the gut virome and its potential interactions with gut bacteria. To

interpret these data and estimate the rates of phage induction and lysis in the human

gut, we utilized mechanistic models of phage population dynamics over time (38–40).

We begin by considering a simplified model of phage ecology, which approximates each

host gut as a well-mixed ecosystem with mass-action kinetics (**Fig. 2A**, **Methods**). For a

single pair of bacteria and phage, this model can be described by a system of three

differential equations for the concentrations of uninfected susceptible bacteria ($S$),

infected bacteria or prophage ($P$), and free phage particles ($V$):

$$\frac{dS}{dt} = \underbrace{\mu_S(S,P) \cdot S}_{\text{Growth}} - \underbrace{\kappa SV}_{\text{Infection}} - \underbrace{\delta S}_{\text{Dilution}} \, , \tag{1}$$

$$207 \qquad \frac{dP}{dt} = \underbrace{\mu_P(S,P) \cdot P}_{\text{Growth}} + \underbrace{\kappa f_L SV}_{\text{Infection}} - \underbrace{\xi P}_{\text{Induction}} - \underbrace{\delta P}_{\text{Dilution}} , \qquad (2)$$

$$208 \qquad \frac{dV}{dt} = \underbrace{B\xi P}_{\text{Induction}} + \underbrace{B(1-f_L)\kappa SV}_{\text{Direct lysis}} - \underbrace{\kappa SV}_{\text{Infection}} - \underbrace{r_e \kappa PV}_{\text{Failed infection}} - \underbrace{\delta V}_{\text{Dilution}} . \qquad (3)$$

Here, $\mu_S(S,P)$ and $\mu_P(S,P)$ are the growth rates of susceptible and infected bacteria, $\delta$

is the overall dilution rate, and $\kappa$ is the infection rate. We assume that a fraction $f_L$ of

infections result in the formation of prophage, while the remaining infections result in

direct lysis of the cell with burst size $B$. Phage particles are also produced by induction

of prophage at rate $\xi$. We assume that infected cells are immune to further infection by

phage particles, with these failed infections resulting in loss of the infecting phage

particle (e.g., to superinfection inhibition mechanisms (41)) with rate $r_e \kappa$. We also

consider extensions of this model that account for dead cells, dead phage, and actively

lysing cells (**Methods**).

Depending on the induction rate and lysogeny fraction, this minimal model can

interpolate between a classic lytic lifestyle and a purely temperate phase in which phage

primarily reproduce via lysogeny (38,42). In the latter case, the spontaneous induction

of prophage can maintain a small population of phage particles (pVMR $\ll$ 1) while the

ratio of phage to microbial genomes (gVMR) remains near one, similar to the

distributions seen in **Fig. 1**. This prophage-dominated regime emerges for a broad

range of model parameters, particularly when the cost of prophage carriage is low

(**Methods**).

We can extend this basic model to larger numbers of phage and bacterial species,

except that we must now allow for multiple prophage states in each bacterium

(representing simultaneous infection by different combinations of prophage). By

summing **Eq. 3** over phage species and integrating over time, one can derive an

approximate equation relating the aggregate prophage and phage particle

concentrations:

$$0 \approx \bar{\xi}^* \bar{B}^* \left( \frac{P^*}{V^*} \right) - \bar{\psi}^* - \delta^*, \tag{4}$$

where $\bar{x}$ and $x^*$ denote community- and time-weighted averages of the quantity $x$,

respectively (**Methods**), and $\psi$ is the residual phage adsorption rate (e.g., due to failed

infections of lysogens). **Eq. 4** assumes that over sufficiently long timescales, the fluxes

controlling phage population sizes within an individual (induction, degradation, infection,

etc.) are approximately balanced, even though day-to-day fluctuations could still be

substantial (we discuss further details of our calculation assumptions in **Methods**).

Based on the stability and moderate variance of the distribution of phage population

densities (**Fig. 1B**), this assumption appears to hold in healthy humans >1 month of

age.

Rearranging **Eq. 4** yields a relation for the average induction rate as a function of the

other key model parameters:

$$\bar{\xi}^* = \left( \frac{1}{\bar{B}^*} \right) \left( \frac{V^*}{P^*} \right) (\bar{\psi}^* + \delta^*). \tag{5}$$

Consistent with intuition, **Eq. 5** predicts that the average induction rate is linearly

proportional to the phage particle to genome ratio, $V^*/P^*$. It also increases with the

combined rate of particle removal (dilution rate $\delta^*$ and adsorption rate $\bar{\psi}^*$, **Fig. 2C**), and

decreases with average burst size $\bar{B}^*$, since a smaller number of induction events are

required to maintain the same density of phage particles. By combining **Eq. 5** with

order-of-magnitude estimates of the other parameters, we can estimate the underlying

induction rate. The ratio of phage particle and phage genome densities can be

estimated from the population distribution in **Fig. 1** as $(V^*/P^*) \approx 10^{-2}$. The mean

dilution rate $\delta^*$ is determined by the inverse of gut transit time, which can vary across

humans but is approximately 1 day$^{-1}$ (43). The burst size $B$ can vary substantially

across phage, with the model *Escherichia coli* phage $\lambda$ having a burst size $B \approx 100$ (44),

ΦCrAss001 having $B \approx 2.5$ (45), and crAssBcn isolates having $B \approx 50$ (46). Thus, we

estimate the order of magnitude of $\bar{B}^* \approx 10$. The residual adsorption rate $\bar{\psi}^*$ is more

difficult to estimate due to our limited understanding of infection rates and host ranges

of gut phage *in vivo*. Nonetheless, setting this quantity to zero yields a lower bound on

the induction rate,

264
$$\bar{\xi}^* \geq \left(\frac{1}{\bar{B}^*}\right)\left(\frac{V^*}{P^*}\right)\delta^* \approx 10^{-3} \text{ day}^{-1}. \tag{6}$$

This lower bound increases to $\bar{\xi}^* \geq 10^{-2}$ day$^{-1}$ when using the estimate of $V^*/P^* = 10^{-1}$

from VLP spike-in sequencing but is still two orders of magnitude lower than the dilution

rate $\delta^*$. The bound is also relatively tight, with substantial deviations only possible if $\bar{\psi}^*$

is larger than $\delta^*$ (**Fig. 2C**). These results suggest that the gut phage particle pool,

despite having ~1,000-fold higher density than the highly lytic surface seawater virome

(47), can be maintained by a very low rate of induction per infected bacterium.

**The relative coverage of integrated prophage provides an upper bound on the rate of phage induction**

Prophage induction can be identified from metagenomic data by comparing the relative coverage of an integrated prophage genome and nearby regions of its bacterial host genome (48). Such methods have thus far been used to make binary determinations of prophage induction for individual phage-bacteria pairs (48), but they also provide information about the underlying induction rate. To extract this information, we use a generalized version of our model in **Eq. 1** to explicitly model activated prophage, representing the state between the start of induction and lysis (**Methods**). These activated lysogens contain $B_a \approx B$ additional copies of the phage genome that correspond to nascent phage particles. The relative coverage $R$ of the prophage and host genomes in metagenomic sequencing data is given by

$$R = 1 + B_a f_a, \tag{7}$$

where $f_a$ is the fraction of currently activated cells. If activated cells are produced from lysogens at rate $\xi$ and have a mean lysis time of $1/\gamma$, then the ratio of activated to non-activated cells will approach a steady-state value of $\sim \xi / (\gamma + \delta)$ (**Methods**). This result can be combined with **Eq. 7** to relate the mean induction rate to the mean relative coverage:

$$\xi^* \leq \frac{(\gamma^* + \delta^*)(R^* - 1)}{(B^* - 1) - (R^* - 1)}. \tag{8}$$

This estimate is robust to the confounding impact of dead cells and viruses contributing to $R$ (**Methods**). The relationship between the induction rate and the relative coverage critically depends on the characteristic lysis time of activated infected cell, $1/\gamma$. Prior studies suggest that lysis time scales with the bacterial host division time (49–51). This

scaling is consistent with estimates of phage burst energetics: a phage burst consumes

a large fraction of the host bacterial energy budget (52), implying that production of a

phage burst is limited by similar factors as host replication. The mean growth rate

roughly matches the dilution rate $\delta^*$ in the parameter regime implied by **Fig. 2C**. Thus,

for the following calculations we assume that $\gamma^*$ is of the same order of magnitude as

$\delta^*$.

**Eq. 8** applies to the subset of phage that are detected within a contig of an assembled

bacterial genome. While it in principle enables measurements of arbitrarily low induction

rates (**Fig. 2D**), but in practice it is difficult to distinguish small values of $R$ from 1 due to

noise and biases in sequencing. Indeed, in a previously published analysis of positive

and negative controls, induction of individual prophage could only be reliably

determined for $R > 2$, and the median number of such events across fecal

metagenomes was zero (49). To establish a tighter upper bound of the induction rate,

we take the "clipped" average of $R$ (i.e., setting values of $R < 1$ to 1) across all adult

samples analyzed in (48), yielding $R^* - 1 \approx 10^{-1}$. Substituting this value into **Eq. 8** with

$B^* = 10$, and $\gamma^* \approx \delta^* = 1\ \mathrm{day}^{-1}$ yields an upper bound on the induction rate of

$$\xi^* \leq 10^{-2}\ \mathrm{day}^{-1}. \tag{9}$$

The clipped mean is larger in infants ($R^* - 1 \approx 0.4$), even after excluding infants

exposed to antibiotics (**Table 2**), suggesting that the induction rate may be higher in

infants. Combined with our other estimates, we can thus bound the average induction

rate within the range $10^{-3} - 10^{-2}\ \mathrm{day}^{-1}$ for adults (**Fig. 2B**), with a somewhat higher

upper bound for infants**.** Importantly, both estimates are substantially lower than the rate

of microbial growth and dilution from the gut, suggesting that gut phage impose a low

mean fitness burden on their bacterial hosts.

**Similar virome properties arise in gnotobiotic mice colonized with a diverse**

**synthetic community of human gut bacterial isolates**

We next examined the implications of our results for a synthetic gut community

designed to mimic the complexity of a native human microbiome (53). This synthetic

community is composed of 119 bacterial isolates from 48 prevalent genera and stably

colonizes gnotobiotic mice for ≥2 months. We reasoned that the virome of hCom2 would

be exclusively composed of temperate phage (at least initially), since it was constructed

from axenic bacterial cultures (54). Our finding that the human gut is dominated by

rarely inducing temperate phage makes two major predictions about the properties of

the hCom2 virome and its relation to the human data above.

First, if the induction rates in hCom2 are as low as our model predicts (**Fig. 2**), we

expect its viral composition in bulk metagenomic sequencing to be entirely predictable

from the abundances of its bacterial members (since $R \approx 1$). It is usually difficult to test

such a prediction in natural communities like the human gut, in which only a subset of

phage can be directly linked to their bacterial hosts (48). Synthetic communities like

hCom2 provide a unique opportunity to test this prediction, since their initial phage-

bacteria associations can be inferred from the sequenced bacterial isolate genomes. To

carry out this test, we generated *in silico* hCom2 metagenomes based on data from a

recent experiment (53) using the sequenced genome of each bacterial strain in

proportion to their measured abundance in each sample (**Methods**). By construction,

these *in silico* datasets only contain phage sequences that were present within the

original bacterial genomes. We then compared the taxonomic composition of these *in*

*silico* datasets with their corresponding mouse metagenomes using the same pipeline

as above (**Fig. 3A,B**, **Methods**).

Consistent with previous observations in a smaller 15-member community (8), we found

that the abundances of individual phage species were highly correlated across the *in*

*silico* and *in vivo* datasets, with the representative sample in **Fig. 3A,B** having a

Spearman correlation of $\rho = 0.9$ for mutually detected phage, compared to $\rho = 0.97$ for

bacteria (as expected by construction). Similar results were obtained for other

compositional similarity metrics, like the Jaccard index or the total abundance of shared

species (**Fig. S3**). The similarity between the *in vivo* and *in silico* metagenomes was

maintained over time, and even after challenge with an undefined fecal sample (**Fig.**

**3C,D, Fig. S3**). These strong correlations confirm that the hCom2 virome is dominated

by temperate phage, and that the induction rates are consistent with our inferences from

the human data above.

A second – and much stronger – prediction of our prophage-dominated human gut

model is that the hCom2 stool virome should qualitatively resemble the stool virome of a

typical human. We tested this prediction by comparing the taxonomic composition of

hCom2-colonized mouse fecal samples (53) with that of a cohort of 245 healthy human

stool metagenomes (55). We reasoned that if hCom2, a synthetic community composed

of axenic bacterial cultures, was missing a large portion of the normal gut virome, then

feces from hCom2-colonized mice would have substantially lower virome diversity than

a typical human stool sample. However, we found that hCom2-colonized mouse feces

exhibited similar phage Shannon diversity as human stool samples, with the hCom2

samples falling between the 13th and 53rd percentiles of the observed human distribution

(**Fig. 4A**). We obtained a similar correspondence between hCom2 and human stool

using a metric of species richness (**Fig. 4B**, **Fig. S4**), as well as the overall ratio of

phage-to-bacterial genomes (**Fig. 4C**, **Fig. S4**). This similarity between hCom2 and

human stool viromes also holds at finer taxonomic levels, with 16 of the 20 most

prevalent phage genera within the human cohort found at >0.1% abundance in hCom2

samples (**Methods**). Thus, consistent with our estimates above, we find that human-like

levels of viral diversity can be achieved by a synthetic community of exclusively

prophage.

The striking similarities between the viromes of hCom2-colonized mouse feces and

human stool can shed light on other coarse-grained features of the human gut virome.

For example, computational tools have been developed to predict the lifestyles of phage

species from their genomes (56,57), enabling estimation of the ratio of virulent phage

(those that cannot stably replicate within their hosts) to temperate phage (24,57). We

used predictions from widely used tools to estimate the virulent to temperate ratio (VTR)

in hCom2-colonized mouse fecal samples (**Methods**). Since hCom2 was constructed

entirely from axenic bacterial cultures, it might be expected to provide a negative control

with a VTR of ~0. However, hCom2-colonized mouse feces metagenomes yielded a

VTR of ~0.5, similar to the typical values observed in human stool samples (**Fig. 4D**,

**Fig. S4, Fig. S5**). This result suggests that existing methods of phage lifestyle prediction

methods underestimate the number of phage that are capable of stable replication

within their bacterial hosts, consistent with previous observations from human stool

samples (**Fig. S2**) (25).

**Discussion**

Our results complement existing surveys of gut phage diversity (4,23,25,58) by

providing a quantitative assessment of phage population dynamics in typical human

hosts. Our updated estimates of the virus-to-microbe ratio show that the small number

of gut phage particles (pVMR ~ $10^{-2}$–$10^{-1}$) is accompanied by a much larger number of

phage genomes (gVMR ~ 4), implying that the vast majority of gut phage genomes are

replicating within their bacterial hosts. These results support the emerging view that

temperate phage lifestyles play a dominant role in the human gut (8,25,36,59,60), even

if they do not contain recognizable integrase genes (e.g. owing to utilization of novel

integrases or having non-integrative lifestyles) (25,36) (**Fig. 4D**). Our quantitative

framework extends this picture by providing new insights into the corresponding phage

induction rates. By integrating imaging and sequencing measurements with a

generalized model of temperate phage dynamics, we estimated that the average

induction rate in adults lies in the relatively low range of $10^{-3}$–$10^{-2}$ per bacterium per day,

imposing only a modest fitness burden on gut bacteria.

These results starkly contrast with well-studied examples like surface seawater, which

possesses a larger ratio of phage particles (pVMR ≈ gVMR~10) and a higher average

lysis rate (20). The reasons for this difference remain uncertain, but they may partially

stem from the distinct physical structures of the two ecosystems. Previous theoretical

and experimental studies have shown that increased spatial structure can select for

lower virulence and increased lysogeny (61,62), consistent with the dominance of

temperate phage in the more spatially structured gut ecosystem, although more work is

needed to quantify the strength of this effect in the gut. Regardless, our results establish

baseline expectations for the co-variation between phage and bacterial abundances

within the microbiome of a typical human. They imply that tight associations between

the bacterial and phage communities may not be driven by active predator-prey

interactions, but may instead be a simple consequence of their synchronized replication

within the same cells, in line with the "piggyback-the-winner" model (60,63,64). This

latter scenario suggests that phage may impact the gut microbiome primarily by acting

as genetic cargo, altering the behavior of their bacterial hosts in certain conditions

(11,65).

These results have substantial implications for future studies of the gut virome's role in

human health. Many studies have sought to identify biomarkers and characterize

possible mechanistic links between gut virome composition and health states such as

lifespan (12), cancer treatment response (2), diabetes (66), metabolic syndrome (67),

and alcoholic hepatitis (68). Importantly, our results highlight confounding factors that

complicate such analyses of virome-health associations, particularly for studies focused

on bulk stool sequencing in which a high abundance of prophage will likely result in

strong statistical links between phage and bacterial composition if the number of VLPs

is low. In studies focused on VLP sequencing, similar correlations could emerge if the

VLP pool is largely a product of relatively uniformly induced prophage, a scenario

hypothesized by prior work (23) and supported by the substantial overlap between bulk

and VLP virome compositions (**Fig. 1C**) (24). These results suggest that methods

similar to phylogenetic regression (69) may be useful for dealing with these confounding

factors.

The quantities in our modeling framework represent averages over time, space, and

hosts that may mask important behaviors that are transient or localized to a particular

host microniche. In particular, averages over longer timescales may not capture shorter-

term variation in induction rates. While the VLP population appears to be broadly stable

over long timescales, the monthly CV of VLP abundances within individuals has a mean

of 0.77 (25). One explanation for such variation is phage induction driven by

environmental changes within the host, a hypothesis consistent with prior studies

showing increased lytic activity in response to perturbations such as bacterial/phage

invasion (8,11), inflammation (70), or exposure to certain dietary or pharmaceutical

compounds (71,72). In addition to such temporal and host variation, phage population

sizes and induction rates may also vary spatially within an individual gut (60,73–75), as

environmental conditions and bacterial densities change substantially along the

gastrointestinal tract (76). It remains possible that the low pVMRs observed in stool

could be produced by a very high induction rate in a smaller population of bacteria in the

proximal colon or small intestine. In the future, spatial variation could be investigated

using recently developed methods for spatially resolved sampling of the microbiome

(77) to measure population sizes and prophage copy numbers across the intestines.

Our modeling framework can be readily applied to such data to estimate local virome

induction rates.

Variation in lifestyle characteristics across phage is also expected, with some phage

effectively existing as mobile genetic elements that rarely lyse their host and others

being primarily lytic. While our current estimates average over multiple phage taxa, our

modeling framework can also be applied to measurements of individual phage species

to estimate species-specific properties. For example, if the particle-to-genome ratios of

an individual phage species can be more accurately measured (**Fig. 1C**), a species-

specific estimate of the induction rate can be obtained from **Eq. 5**. Applications of this

approach are currently limited by the known amplification biases of existing VLP

sequencing methods (34), but the adoption of sequencing protocols that do not involve

MDA (78) may enable such species-specific resolution in the future.

Beyond our modeling assumptions, there are also limitations in the phage quantification

methods used for experimental measurements. The process of VLP isolation may lead

to substantial loss of phage particles, particularly given the spatially structured nature of

stool and the potential for phage particles to adhere to large particulates. Additionally,

imaging-based quantification methods can both underestimate phage densities due to

loss of material during preparation (33), and overestimate due to the presence of cell

debris or other DNA-containing particles (31). Similarly, RNA phage cannot be

visualized using DNA-staining-based microscopy (23,29). Underestimation of phage

particle densities would imply a higher true pVMR, which would increase the

corresponding induction rate estimate from **Eq. 5.** Note, however, that for our pVMR

estimates to be comparable to that of surface seawater would require very large

differential loss rates (>99%), which could potentially be measured with appropriate

spike-ins. Our analysis framework can easily be applied to updated density estimates as

they become available.

Overall, our work motivates future experimental directions for the gut virome field. While

informative, our estimates of the mean induction rate still encompass 1-2 orders of

magnitude owing to limitations of current data. Given the noise intrinsic to metagenomic

sequencing, we expect that deeper bulk sequencing will have limited benefits for

estimating of the mean induction rate in the parameter regimes suggested by our

analysis. More accurate and direct estimation will likely be dependent on measurement

of rare induced cells. Single-cell bacterial sequencing (79) is a promising avenue to

achieve the needed detection power. Alternatively, measurement of *in vivo* phage

adsorption rates or the degradation rates of lysed cells (**Methods**) would enable

improved estimation of induction rates that we derived in **Fig. 2**. Our results also provide

guidance for the design of virome perturbation experiments, which should focus on

measuring increases in induction and horizontal gene transfer – a major avenue through

which prophage influence their hosts. Finally, the similarities between the estimated

VTR in hCom2-colonized mouse feces (**Fig. 4D**) and human stool metagenomes

highlights the current lack of knowledge regarding the genetic mechanisms enabling

bacterial host-association of gut phage. These results imply that many gut phage

currently computationally identified as virulent in fact contain unidentified and

uncharacterized host-association genes. This pool of genes represents a rich ground for

future phage molecular biology work.

**Acknowledgements**

The authors thank the Huang and Good labs, Ami Bhatt, Danica Schmidtke, Gabriel

Birzu, Colin Hill, and Andrey Shkoporov for helpful discussions. The authors

acknowledge support from NIH RM1 Award GM135102 and R01 AI147023 (to K.C.H.),

NSF Awards EF-2125383 and IOS-2032985 (to K.C.H.), NIH R35 GM146949 (to

B.H.G.), Alfred P. Sloan Foundation grant FG-2021-15708 (to B.H.G.), Human Frontier

Science Program grant RGEC33/2023 (to B.H.G.), and a Friedrich Wilhelm Bessel

Award from the Humboldt Foundation (to K.C.H.). B.H.G. and K.C.H. are Chan

Zuckerberg Biohub Investigators. J.L. was supported by a Stanford PRISM Baker

Fellowship. This work was also supported in part by the National Science Foundation

under Grant PHYS-1066293 and the hospitality of the Aspen Center for Physics. We

thank the Stanford Research Computing Center for use of computational resources on

the Sherlock cluster.

**Figures**

**Figure 1: Comparisons of absolute population densities suggest that the human gut virome is numerically dominated by prophage.** (A) Schematic overview of common gut viral population quantification methods. Metagenomic classification of bulk fecal samples yields phage-to-bacteria genome ratios, which can be combined with absolute bacterial densities to estimate the absolute density of phage genomes. Alternatively, virus-like particles (VLPs) can be extracted from the stool and quantified by epifluorescence microscopy or spike-in sequencing. (B) Gut virome population densities are approximately maintained across human life stages. Each violin plot represents quantification of one population using one measurement method in one

study (**Table 1**), with individual dots representing subjects. The gray line denotes the

average stool bacterial density reported in (32) ($0.92 \times 10^{11}$ bacteria/g stool). (C)

Species-level absolute abundance analysis of the overlap of phage communities found

using VLP- and stool-based quantification approaches. Data represent one healthy

adult population (25) ($n = 10$, VLP WGS absolute quantification) and one population of

4-month-old infants (23) ($n = 19$, VLP microscopy absolute quantification). Each point is

the absolute abundance of one phage species in a matched pair of VLP and stool

samples from one subject (**Methods**). Triangle markers denote species classified as

virulent and circle markers denote species classified as temperate (**Methods**).

Histograms show the distribution of absolute abundances of phage species found

exclusively in either the VLP or stool samples. (D) Relative distribution of phage

genomes or particles between VLP and stool WGS. Each violin plot represents the total

absolute abundances of phage genomes or particles found within only VLP samples,

only stool samples, or shared between VLP and stool. Individual points correspond to a

single subject. Underlying data are the same as in (C).

**Figure 2: Mathematical modeling of phage population dynamics enables**

**estimation of the average phage induction rate in the human gut.** (A) Schematic of

the minimal model of temperate phage dynamics represented by **Eq. 1-3**. Infection of

susceptible bacteria produces prophage, which induce at rate $\xi$, lysing their host and

producing a burst of $B$ free phage particles. (B) Schematic representation of induction

rate estimates. We combine measurements of phage particle to genome ratio and

relative prophage copy number with the model in (A) to estimate upper and lower

bounds on the phage induction rate. Note that given the uncertainty in parameter

values, these estimates are only reported as approximate orders-of-magnitude, with the

combined bound illustrated in grey. (C) Estimated induction rate as a function of total

phage adsorption rate $\psi$ (i.e., all non-dilution phage particle removal mechanisms). The

solid line corresponds to **Eq. 5**, using the phage genome to particle ratio inferred from

**Fig. 1**. (D) Estimated induction rate as a function of the relative coverage of prophage,

$R$. The solid line corresponds to **Eq. 8** with $\gamma \approx \delta$. The solid circle is the mean relative

coverage in adults ($R - 1 \approx 10^{-1}$), using measurements from (48).

**Figure 3: Phage abundance dynamics in a diverse synthetic gut community can be predicted from bacterial abundances alone.** (A,B) Relative read abundances of bacterial (A) and phage species (B) in fecal samples from hCom2-colonized gnotobiotic mice, compared to *in silico* metagenomes ("hMock") generated from their corresponding bacterial genomes weighted according to the fecal bacterial microbiota composition (**Methods**). The example shown is for a single representative sample (mouse 3, week 1). JSS is the Jensen-Shannon similarity and shared $\rho$ denotes the Spearman

correlation computed from species observed in both samples. (C,D) Bacterial and

phage JSS between *in vivo* and *in silico* metagenomes over time and in response to

human stool challenge perturbation. Lines show mean JSS in either unchallenged mice

($n$ = 5) or mice challenged with a human stool perturbation after week 4 ($n$ = 15) over

time. Shaded areas represent 1 standard deviation computed across mice at each time

point, and the dashed line denotes the time of fecal challenge.

**Figure 4: Large-scale features of human stool viromes are recapitulated in a community constructed only of bacterial isolates.** (A-D) Comparison of Shannon diversity (A), weighted species richness (B), virus to microbe ratio (C), and virulent to temperate ratio (D) in fecal samples from hCom2-colonized gnotobiotic mice (53) compared to human stool samples. Violin plots labelled "stool" represent distributions of microbiome properties across $n$ = 245 healthy adults studied in (55). Violin plots labelled "hCom2" represent samples from 20 gnotobiotic mice colonized with the synthetic community hCom2 ($n$ = 77 total samples, all from unchallenged mice or pre-challenge).

583    Virulent to temperate ratios (VTRs) were estimated using the UHGV database phage

584    species virulence predictions (**Methods**). Dashed black line denotes the null

585    expectation of VTR = 0 for a community constructed from axenic bacterial cultures.

 **Methods**

 **Meta-analysis of gut phage quantifications**

 Table 1 summarizes the studies used in our meta-analysis of gut phage abundances.

 Each row corresponds to a single violin plot in **Fig. 1A**, with the order of the table rows

 matching the order in which the datasets appear in the figure.

| Data origin | Secondary quantification | Method | Population |
|---|---|---|---|
| Liang *et al.* 2020 **(23)** | N/A | VLP EFM | Newborns |
| Liang *et al.* 2020 **(23)** | N/A | VLP EFM | 1-month-old infants |
| Liang *et al.* 2020 **(23)** | This study | Stool WGS | 1-month-old infants |
| Liang *et al.* 2020 **(23)** | N/A | VLP EFM | 4-month-old infants |
| Liang *et al.* 2020 **(23)** | This study | Stool WGS | 4-month-old infants |
| Bikel *et al.* 2021 **(80)** | N/A | VLP EFM | 7–10-year-old children |
| Kim *et al.* 2011 **(26)** | N/A | VLP EFM | Healthy adults |
| Hoyles *et al.* 2014 **(29)** | N/A | VLP EFM | Healthy adults |
| Shkoporov *et al.* 2019 **(25)** | N/A | VLP WGS | Healthy adults |
| Shkoporov *et al.* 2019 **(25)** | This study | Stool WGS | Healthy adults |
| Yachida *et al.* 2019 **(55)** | This study | Stool WGS | Healthy adults |

 **Table 1: Studies represented in the quantification meta-analysis in Fig. 1A.** "Data

 origin" column indicates the study that produced the original data, and "Secondary

 analysis" denotes studies that performed additional bioinformatic analyses represented

 in **Fig. 1A**. For all stool WGS quantifications, Phanta was used to estimate the genomic

virus-to-microbe ratio (gVMR) which was then multiplied by an estimate of gut microbial

abundance (32). For studies in which a table of quantifications was not explicitly

provided, counts were digitally extracted from figures using WebPlotDigitizer.

We now briefly describe the different measurement methodologies applied by the

studies analyzed here. The measurement methodologies for gut phage abundance fall

into three classes (**Fig. 1A**), which quantify different subsets of the gut phage

population. One method (labeled "Stool WGS" in **Table 1**) is based on metagenomic

sequencing of whole stool samples, from which the ratio of the abundance of phage

DNA to that of bacteria DNA can be computed (24). By normalizing this ratio by typical

phage and bacterial genome lengths, the ratio of phage to bacterial genome copies is

obtained (24), which combined with quantification of absolute bacterial density

generates an estimate of absolute phage genome density. This method captures both

prophage (e.g., lysogens, phage-plasmids, etc.) and the fraction of phage particles that

lyse during DNA extraction. The other two methods involve isolation of virus-like

particles (VLPs) as representatives of the phage particle population present within stool.

Isolation typically involves 0.2- or 0.45-μm filtration and DNAse/RNAse treatment,

among other steps. In one method, VLPs are stained with a DNA-binding dye and

enumerated via epifluorescence microscopy (23) (labeled "VLP EFM" in **Table 1**), while

in the other method the VLPs are mixed with a known quantity of a non-gut reference

phage and metagenomically sequenced, with the reference phage enabling absolute

quantification (25) (labeled "VLP WGS" in **Table 1**). In contrast to the method based on

bulk stool metagenomics, these VLP-based methods do not capture prophage by

design. Given the drastic methodological differences between bulk and VLP-based

approaches, we define two separate VMRs: the genomic VMR (gVMR), based on bulk

stool sequencing, and the particle VMR (pVMR), based on VLP-approaches.

In our calculations of pVMR and of phage absolute abundance from gVMR, we require

an estimate of the microbial density of the gut microbiome. Note here that we use the

term "microbe" to denote all microorganisms (including archaea, bacteria, and

unicellular eukaryotes); in practice, the vast majority of gut microbes are bacteria (32)

and this is reflected in our taxonomic estimations from Phanta. For all such calculations,

we used a standardized value of $0.92 \times 10^{11}$ microbes/g stool obtained from a

comprehensive meta-analysis of stool microbe abundance quantifications from humans

>1 month of age (32). Using a single standardized value is justified by the minimal

variation of total gut microbial density across human populations >1 month of age (32).

Doing so also eliminates the confounding effect of inter-study variability gut microbial

density. Indeed, a few gut virome studies reported bacterial density estimates of $\sim 10^9$ or

$\sim 10^{10}$ microbes/g stool (23,26), orders of magnitude below well-established values,

which led to inflated values of pVMR. We do not know the origin of these discrepancies,

but we assume based on the weight of evidence that the microbial density is closer to

$0.92 \times 10^{11}$ microbes/g stool.

For the direct VLP-stool comparisons in **Fig. 1C,D**, we used only subjects for which VLP

metagenomes, stool metagenomes, and VLP absolute quantification were available. For

the dataset in (25), matching metagenomes were available for the subjects only at the

8-month timepoint, and VLP quantification was performed only during months 9-12.

Thus, we used the month 8 metagenomes in combination with the average of month 9-

12 measurements of each subject. We found that in both infant and adults these VLP

metagenomic samples had a high gVMR, median ~$10^3$ compared a median gVMR of

~3-4 found in the corresponding stool samples (as measured by Phanta). This large

gVMR difference persists even if *Microviridae* species or all species found only in VLP

sample are removed, indicating that the VLP-stool overlap in **Fig. 1C,D** is likely not due

to bacterial contamination of the VLP pool.

**Analysis of prophage copy number data**

For our estimation of induction rate from prophage copy number, we use results from

Kieft et al. (48), which developed a computational tool, PropagAtE, for estimating

whether an integrated prophage is active. They applied their tool to several

metagenomic sequencing studies and we use the values of $R$ estimated by their tool

(available in Table S3B of their manuscript). We use only prophage-sample

combinations detected as present by their tool and perform additional quality filtering

requiring minimum median host and prophage coverage >1, and prophage coverage

breadth >0.5. We show the resulting summary statistics across cohorts in **Table 2**.

| Sample set | $n$ samples | $n$ pairs | Mean $R$ | Median $R$ | Clipped mean $R$ |
|---|---|---|---|---|---|
| All adults | 123 | 3,459 | 1.04 | 1.01 | 1.09 |
| CRC | 15 | 474 | 1.02 | 0.99 | 1.07 |
| HeQ | 96 | 2842 | 1.05 | 1.02 | 1.09 |
| IjazUZ | 12 | 143 | 1.04 | 1.04 | 1.08 |
| All infants | 79 | 702 | 1.27 | 1.1 | 1.40 |
| Infant (non-abx) | 22 | 254 | 1.33 | 1.22 | 1.43 |
| Infant (abx) | 57 | 448 | 1.24 | 1.02 | 1.39 |

**Table 2: Summary statistics of prophage copy number $R$ across different metagenomic sequencing cohorts, computed based on results from Kieft et al. (48).** "$n$ samples" denotes the number of metagenomic sequencing samples in the cohort, "$n$ pairs" denotes the total number of prophage-bacterial host pairs identified as present and passing the coverage/breadth requirements in those samples. The clipped mean is the mean computed with values of $R < 1$ set to $R = 1$. The CRC dataset is composed of adults with colorectal adenoma and healthy adults, HeQ is composed of adults with Crohn's disease and healthy adults, IjazUZ is composed of adults with Crohn's disease, Infant (non-abx) is composed infants that were not exposed to antibiotics, Infant (abx) is composed of infants that were exposed to antibiotics.

**Processing and analysis of metagenomic datasets**

All metagenomic datasets analyzed in this manuscript were first subjected to quality

control/filtering and adapter removal using the BBDuk decontamination tool in BBTools

(81). Settings used were kmer length "k = 23", "hdist = 1", trim direction "qtrim = rl" (trim

both ends), minimum entropy "entropy = 0.5", sliding window for entropy calculation

"entropywindow = 50", kmer length for entropy calculation "entropyk = 5", minimum

quality "trimq = 25", and minimum read length "minlen = 50". Samples were then

deduplicated using the clumplify tool in BBTools. The maximum number of substitutions

between duplicate reads was zero ("subs = 0"). We found that deduplication minimally

influenced the estimated community compositions.

For taxonomic quantification of samples, we used Phanta, a kmer-based method that

simultaneously profiles phage and bacteria (24). Phanta was run using default settings:

confidence threshold "confidence_threshold 0.1", viral genome coverage requirement

"cov_thresh_viral 0.1", viral unique minimizer threshold "minimizer_thresh_viral 0",

bacterial genome coverage requirement "cov_thresh_bacterial 0.01", and bacterial

unique minimizer threshold "minimizer_thresh_bacterial 0". The

"uhggv2_uhgv_mqplus_v1" database was used, which is based on the prokaryotic

UHGG database and viral UHGV database. For taxonomy-based analyses, the

provided UHGV taxonomy was used, except for quantification of *Microviridae*

abundance, for which we used the provided ICTV taxonomy.

**hCom2 metagenome reconstruction from bacterial genomes**

To generate mock versions of hCom2-colonized mouse fecal metagenomes, we

generated synthetic short-read sequencing datasets using the set of isolate genomes

(53). To determine the relative abundances of genome reads within each sample, we

used the bacterial compositions estimated by NinjaMap (53). NinjaMap is designed to

quantify the composition of synthetic communities in which sequenced genomes are

available for all member strains. For each sample, we specified the relative fraction of

reads from each genome based on that strain's relative abundance and normalized by

its genome length. For genomes that are not assembled into a single contig, the read

abundance was split among the contigs weighted by the length of each contig. To

generate synthetic shotgun samples, we used Grinder (82) with the following settings:

quality levels "-qual levels 33 31", insert distance "-insert_dist 800", read length "-

read_dist 140", forward-reverse mate orientation "-mate_orientation FR", characters

deleted from reference sequences "-delete_chars '-~*NX'", and distribution of mutations

"-mutation_dist uniform 0". The total number of reads generated for each sample was

equal to the post-QC read number of the corresponding original mouse fecal sample.

Fecal samples with <$10^5$ reads were excluded from the analysis. The resulting samples

were subjected to the standard pre-processing pipeline applied to all other

metagenomic sequencing data in the manuscript. For the comparison of hCom2 stool to

human stool, we exclude human stool samples with <$10^5$ reads.

**Computation of community summary statistics in metagenomic samples**

For all analyses except the hCom-hMock comparison in **Fig. 3** and **Fig. S3**, we used

relative taxonomic abundances (which are computed in Phanta by normalizing relative

read abundances to the phage or bacterial genome size). For the hCom-hMock

comparison, relative read abundances were used to better compare reconstruction

fidelity between phage and bacterial communities. To obtain the genomic virus to

microbe ratio (gVMR) of a sample, we calculated the ratio of the total taxonomic

abundance of members within the viral superkingdom to the total taxonomic abundance

of members of the archaeal, bacterial, and non-human eukaryotic superkingdoms. In

practice, the denominator of the gVMR is vastly dominated by the bacterial taxonomic

abundance. To obtain the virulent to temperate ratio (VTR), we calculated the ratio of

total taxonomic abundance of phage classified as virulent to the total taxonomic

abundance of phage classified as temperate. In the main figures, we used the virulence

predictions from the Phanta UHGV database (24), which utilizes a combination of

scores from BACPHLIP (56) along with information from the PHROG database (83) and

geNomad (84). An alternative VTR estimate was performed with scores from PhaTYP

(57) (**Fig. S5**). PhaTYP was run on the UHGV genomes using default settings.

Shannon diversity was computed at the species level as $H = -\sum_i x_i \log_2(x_i)$, where $x_i$

is the relative taxonomic abundance of species $i$ within the bacterial or phage

community. Weighted richness was computed such that the richness contribution of

each species is weighted by $1 - \exp(-x_i/x_0)$, where $x_0 = 10^{-3}$. For the hCom2

reconstruction analysis, the Jensen-Shannon similarity was computed as $\text{JSS} = 1 -$

$\left(\frac{1}{2}\right)\sum_i p_i \log_2\left(\frac{p_i}{m_i}\right) - \left(\frac{1}{2}\right)\sum_i q_i \log_2\left(\frac{q_i}{m_i}\right)$, where $p_i$ and $q_i$ are the relative read abundances

of the communities being compared, normalized to sum to 1 within a given taxonomic

grouping (e.g., phage at the species level), and $m_i = \frac{p_i + q_i}{2}$.

**Overview of phage mathematical model**

We begin with a mathematical model of a single phage and bacterial species in a well-

mixed environment, similar to that of (38). We show here how this model can be

reduced to the model presented in the main text. This model involves the concentration

of a nutrient ($C$), susceptible cells ($S$), cells containing quiescent prophage ($P$), cells in

which the prophage has been activated ($P_a$), viral particles ($V$), and dead cells/viruses of

various kinds ($D_i$). All populations are diluted at rate $\delta$. All populations also experience

non-dilution mortality/degradation at rate $\omega_i$. All bacterial cells experience the same

non-dilution mortality rate $\omega_B$. Susceptible cells and cells containing quiescent prophage

grow by consuming the resource. Resource consumption occurs with uptake rate $\mu(C)$

for susceptible cells and $\mu(C)(1 + s)$ for prophage-containing cells, where $\mu(\cdot)$ is the

growth function and $s$ is the fitness benefit/cost of carrying a quiescent prophage.

Resources are supplied at a constant rate $\Gamma$. Susceptible cells are exposed to viral

particle infection by mass-action kinetics at rate $\kappa$, with a fraction $f_L$ becoming quiescent

prophage-containing cells and a fraction $1 - f_L$ shifting to the activated cell class ($f_L$

models the lysis-lysogeny decision upon initial infection). Prophage-containing cells are

induced at rate $\xi$, shifting to the activated class. Cells in the activated class are

assumed not to grow and lyse at rate $\gamma$, producing a burst of $B$ viral particles. Viral

particles are lost by infecting susceptible cells, failed infection of prophage-containing

cells (e.g., to superinfection inhibition mechanisms (41)), and non-dilution mortality.

Failed infection occurs at rate $r_e\kappa$, where $r_e$ is the ratio of infection coefficients of

prophage-containing and susceptible cells. Dead susceptible, prophage-containing, and

activated cells, with concentrations $D_S$, $D_P$, and $D_a$, respectively, and dead viruses with

concentration $D_V$, are produced by non-dilution mortality. Cells that die by phage lysis

are tracked separately with concentration $D_L$. Non-lysed dead cells are degraded at rate

$\omega_D$, while lysed cells are degraded at rate $\omega_{DL}$, and dead viruses are degraded at rate

$\omega_{VD}$. The dynamics governing this model are thus:

770
$$\frac{dC}{dt} = \Gamma - \mu(C)S - (1+s)\mu(C)P - \delta C, \qquad (S1a)$$

771
$$\frac{dS}{dt} = \mu(C)S - \kappa SV - (\delta + \omega_B)S, \qquad (S1b)$$

772
$$\frac{dP}{dt} = f_L \kappa SV + (1+s)\mu(C)P - \xi P - (\delta + \omega_B)P, \qquad (S1c)$$

773
$$\frac{dP_a}{dt} = (1 - f_L)\kappa SV + \xi P - \gamma P_a - (\delta + \omega_B)P_a, \qquad (S1d)$$

774
$$\frac{dV}{dt} = \gamma B P_a - \kappa SV - r_e \kappa PV - (\delta + \omega_V)V, \qquad (S1e)$$

775
$$\frac{dD_S}{dt} = \omega_B S + (\delta + \omega_D)D_S, \qquad (S1f)$$

776
$$\frac{dD_P}{dt} = \omega_B P + (\delta + \omega_D)D_P, \qquad (S1g)$$

777
$$\frac{dD_a}{dt} = \omega_B P_a + (\delta + \omega_D)D_a, \qquad (S1h)$$

778
$$\frac{dD_L}{dt} = \gamma P_a + (\delta + \omega_{DL})D_L, \qquad (S1i)$$

779
$$\frac{dD_V}{dt} = \omega_V V + (\delta + \omega_{DV})D_V, \qquad (S1j)$$

780    To recover the model discussed in the main text (**Eq. 1-3**), we make a separation of

781    timescales assumption to reduce the number of state variables in the model, assuming

782    that the nutrient, activated cells, and dead cells are in pseudo-steady-state with the

783 remaining state variables (i.e., $\frac{dC}{dt} = \frac{dP_a}{dt} = \frac{dD_i}{dt} = 0$). This assumption yields the following

784 expressions:

$$P_a^* = \frac{(1 - f_L)\kappa SV + \xi P}{\gamma + \delta + \omega_B}, \tag{S2a}$$

$$D_S^* = \frac{\omega_B S}{\delta + \omega_D}, \tag{S2b}$$

$$D_P^* = \frac{\omega_B P}{\delta + \omega_D}, \tag{S2c}$$

$$D_a^* = \frac{\omega_B P_a^*}{\delta + \omega_D}, \tag{S2d}$$

$$D_L^* = \frac{\gamma P_a^*}{\delta + \omega_{DL}}, \tag{S2e}$$

$$D_V^* = \frac{\omega_V V}{\delta + \omega_{DV}}, \tag{S2f}$$

791 These equations can be used to define a simplified set of equations with only the

792 sensitive, prophage, and phage particle abundances

$$\frac{dS}{dt} = \mu(C^*)S - \kappa SV - (\delta + \omega_B)S, \tag{S3a}$$

$$\frac{dP}{dt} = f_L \kappa SV + (1 + s)\mu(C^*)P - \xi P - (\delta + \omega_B)P, \tag{S3b}$$

$$\frac{dV}{dt} = \tilde{B}\xi P + \tilde{B}(1 - f_L)\kappa SV - \kappa SV - r_e \kappa PV - (\delta + \omega_V)V, \tag{S3c}$$

796 where $C^*(S, P)$ is defined implicitly by $0 = \Gamma - \mu(C^*)S - (1 + s)\mu(C^*)P - \delta C^*$ and $\tilde{B} =$

797 $Bf_\gamma = B\left(\frac{\gamma}{\gamma + \delta + \omega_B}\right)$. $f_\gamma$ can be interpreted as the fraction of activated cells that are not

798 diluted or die before lysis occurs and thus $\tilde{B}$ can be interpreted as an effective burst

799 size.

**Conditions for robust phage invasion**

In the following two sections, we will assess invasion and stability of phage populations in the model defined by **Eq. S3**. We assume a linear growth function $\mu(C) = \alpha C$ for these derivations, leading to $C^* = \frac{\Gamma}{\alpha S + (1+s)\alpha P + \delta}$. In the absence of bacteria, the resource concentration will saturate at a steady-state value of $C_0^* = \Gamma/\delta$. Bacteria will be able to invade this ecosystem when their initial growth rate exceeds the death and dilution rate: $\frac{\alpha \Gamma}{\delta} > \delta + \omega_B$. Given the stable bacterial colonization seen in the human gut, we assume this condition to be satisfied. More strongly, given that $\Gamma$ and $\delta$ likely vary substantially over time even within a single host (corresponding to variation in food intake and passage time), robust colonization requires $\frac{\alpha \Gamma}{\delta} \gg \delta + \omega_B$.

In the absence of virus, susceptible bacteria will saturate at an equilibrium abundance

$$S_0^* = \frac{\Gamma}{\delta + \omega_B}\left(1 - \frac{\delta(\delta + \omega_B)}{\Gamma \alpha}\right) \approx \frac{\Gamma}{\delta + \omega_B}, \tag{S4}$$

where the approximation follows from the robust bacterial colonization assumption ($\alpha \Gamma \gg \delta(\delta + \omega_B)$). In the absence of lysogeny ($f_L = 0$), viruses will be able to invade this susceptible population if the initial phage replication is greater than death: $(\tilde{B} - 1)\kappa S_0^* - (\delta + \omega_V) > 0$. Equivalently, the (lytic) basic reproductive number of the virus must be greater than one:

$$R_0 \equiv \frac{(\tilde{B} - 1)\kappa S_0^*}{\delta + \omega_V} > 1. \tag{S5}$$

As above, since $\Gamma$ and $\delta$ will vary (and thus $S_0^*$ will vary), robust phage invasion will require that $R_0 \gg 1$, and thus this is the regime we are primarily interested in.

## Stability of the prophage-dominated steady state

Given the estimated abundance of prophage in the gut (**Fig. 1B**), we are particularly

interested in the properties of the prophage-only steady state of the model. We will

show that under reasonable assumptions this steady state is likely stable in the gut and

thus can be invoked in interpretating our induction rate estimates. The prophage-only

steady state has $S^* = 0$, $P^* \approx \left(\frac{\delta + \omega_B}{\delta + \omega_B + \xi}\right) S_0^*$, and $V^* = \frac{\tilde{B}\xi P^*}{r_e \kappa P^* + \delta + \omega_V}$, and $C^* \approx \frac{\delta + \omega_B + \xi}{\alpha(1+s)}$. As we

are in the robust bacterial colonization regime, we neglect the contribution of dilution to

nutrient elimination. This steady state is robust to small perturbations of $P$, $V$, and $C$.

From an invasion analysis, it will be robust to small invasion of susceptible bacteria if

the net growth rate of these susceptible bacteria is negative:

$$\mu(C^*(0, P^*)) - \kappa V^* - (\delta + \omega_B) < 0. \tag{S6}$$

834    Substituting in the definition of $C^*$, dividing by $\delta + \omega_B$, and rearranging yields

$$\frac{\frac{\xi}{\delta + \omega_B} - s}{1 + s} < \frac{\kappa V^*}{\delta + \omega_B}. \tag{S7}$$

836    We can express $V^*$ in terms of $R_0$ as

$$V^* = \frac{\frac{\tilde{B}R_0}{(\tilde{B} - 1)\kappa}(\delta + \omega_B)\xi}{\left(\frac{r_e R_0}{\tilde{B} - 1}\right)(\delta + \omega_B) + \delta + \omega_B + \xi}, \tag{S8}$$

838    and substituting this equation into the invasion condition yields

$$\frac{\frac{\xi}{\delta + \omega_B} - s}{1 + s} < \frac{\frac{\tilde{B}R_0}{(\tilde{B} - 1)}\left(\frac{\xi}{\delta + \omega_B}\right)}{\frac{r_e R_0}{\tilde{B} - 1} + 1 + \frac{\xi}{\delta + \omega_B}}. \tag{S9}$$

We are particularly interested in the regime where the direct cost (if negative) of the

prophage $s$ is small relative to one, but still potentially large relative to other small

parameters in the system. This limit is consistent with the modest energetic cost of

replicating a phage genome (52) and that non-lytic mobile genetic elements have been

observed to rapidly undergo compensatory adaptation to reach very low fitness costs

(85). Expanding to lowest order in $s$ leads to

$$
\frac{\xi}{\delta + \omega_B} - s < \frac{\frac{\tilde{B}R_0}{(\tilde{B}-1)}\left(\frac{\xi}{\delta + \omega_B}\right)}{\frac{r_e R_0}{\tilde{B}-1} + 1 + \frac{\xi}{\delta + \omega_B}}.
\tag{S10}
$$

847    This condition is violated at very high induction rate ($\xi \gtrsim R_0(\delta + \omega_B)$) and at low

848    induction rate when

$$
\frac{\xi}{\delta + \omega_B} < (-s)\left(\frac{r_e}{\tilde{B}} + \frac{1 - \tilde{B}^{-1}}{R_0}\right).
\tag{S11}
$$

The term on the right-hand-side is much smaller than $-s$ in the empirically relevant

regime where $\tilde{B} \gg 1$ and $R_0 \gg 1$. Thus, as long as the dominant cost of gut prophage is

induction, i.e., $\xi > -s$, as has been experimentally observed for some phage (86), then

the gut ecosystem likely exists within a regime where the prophage-only state is stable.

**Overview of induction rate estimation approach**

In the following sections, we show detailed derivations of the induction rate estimates

presented in the main text, starting from the single phage-bacteria model in **Eq. S3**. In

addition to the estimations based on phage particle to prophage ratio and prophage

copy number, we also show an estimation based on cell viability. With currently

available data, this estimator is poorly constrained and thus not included in the main

text, but some results from this derivation are used in the derivation of the estimate

based on prophage copy number.

In deriving these estimates, we begin with a general form of the calculation that makes

no assumptions about the relative abundance of prophage. This approach leads to

estimates of the total lysis rate, which includes both induction of prophage and lysis of

sensitive cells via non-lysogenic infection. We then simplify these estimates by

assuming that the gut is prophage-dominated, leading to the expressions for the

average induction rate in the main text. This simplification only affects the interpretation

of the resulting estimate: if the prophage-dominated simplification is incorrect and a

substantial amount of phage particle production occurs from sensitive cells, then the

estimates are still valid as total lysis rate estimates. In the final model section, we show

how our framework can be extended to communities with multiple species of bacteria

and phage with explicitly time-varying parameters.

**Total lysis rate and induction rate estimate from phage particle to prophage ratio**

Here, we estimate the average lysis rate (including both induction of lysogens and lysis

after non-lysogenic infection) from the phage particle to prophage ratio. We begin with

the viral dynamics from the timescale-separated prophage model (**Eq. S3**). We define

the population-weighted total lysis rate $\eta$ such that $\eta(S + P) = \xi P + (1 - f_L)\kappa SV$. We can

also rewrite this as $\eta = \xi x_P + (1 - f_L)\kappa V x_S$ where $x_i$ are the population relative

abundance within the $S + P$ pool of cells. By rearranging **Eq. S3c** when $\frac{dV}{dt}$ is on average

zero (i.e. $\frac{1}{\Delta t}\int_0^{\Delta t} \frac{dV}{dt}\, dt \approx 0$), we can obtain an expression for $\eta^*$:

884
$$\eta^* = \frac{1}{\tilde{B}}\left(\frac{V^*}{S^* + P^*}\right)(\kappa S^* + r_e \kappa P^* + \omega_V + \delta), \qquad \text{(S12)}$$

where asterisks denote the time-averaged value $x^* = \frac{1}{\Delta t}\int_0^{\Delta t} x(t)\, dt$. This approximation

assumes that the microbiome is in a statistical steady state (no net trend in $V$) and that

$\Delta t$ is long enough that time averages have converged to their ensemble-averaged

values. At a minimum, this assumption requires that $\Delta t \gtrsim 1/\delta$. The assumption of a

statistical steady state is supported by the results of our absolute abundance meta-

analysis (**Fig. 1**). Given the limited knowledge of $\kappa$, $r_e$, and $\omega_V$ in the gut, we use **Eq.**

**S12** to construct a lower bound on $\eta^*$:

892
$$\eta^* \geq \frac{1}{\tilde{B}}\left(\frac{V^*}{S^* + P^*}\right)\delta. \qquad \text{(S13)}$$

893 In the prophage-dominated regime (i.e., $S^* = 0$) we recover $\eta^* = \xi^* \geq \left(1/\tilde{B}\right)(V^*/P^*)\delta$, a

894 bound on the induction rate.

896 In practice, measured pVMR may not be $\frac{V^*}{S^* + P^*}$ due to the contribution of dead cells and

897 dead viruses. If all populations are represented in the measurement, the pVMR will

898 instead be

899
$$\text{pVMR} = \frac{V^* + D_V^*}{S^* + P^* + P_a^* + D_S^* + D_P^* + D_a^* + D_L^*}. \qquad \text{(S14)}$$

900 This is related to $\frac{V^*}{S^* + P^*}$ by:

901
$$\frac{V^*}{S^* + P^*} = \left(\frac{\nu_V}{\nu_B}\right)\left(\frac{S^* + P^* + P_a^*}{S^* + P^*}\right) \cdot \text{pVMR}, \qquad \text{(S15)}$$

902     where $\nu_i$ is the cell or viral viability fraction (the fraction of cells or viruses that are

903     viable). We now substitute this expression into **Eq. S12** and use the fact that $\frac{\nu_V}{1-\nu_V} =$

904     $\frac{V^*}{D_V^*} = \frac{\delta + \omega_{DV}}{\omega_V}$ to yield

$$\eta^* = \frac{\mathrm{pVMR}}{\tilde{B}} \left( \frac{S^* + P^* + P_a^*}{S^* + P^*} \right) \left( \frac{1}{\nu_B} \right) (\nu_V \kappa S^* + \nu_V r_e \kappa P^* + (1 - \nu_V)\omega_{DV} + \delta). \tag{S16}$$

906     Thus, utilizing a pVMR including the dead populations still functions as a lower bound

907     estimate of $\eta^*$:

$$\eta^* \geq \left( \frac{\mathrm{pVMR}}{\tilde{B}} \right) \delta. \tag{S17}$$

**Cell death and total lysis rate estimates from live cell fraction**

911     Here, we derive estimates of both the non-lysis cell mortality rate $\omega_B$ and the total lysis

912     rate $\eta$ based on the fraction of cells that are living/viable within the microbiome, defined

913     in our model as $\nu_B^* = \frac{S^* + P^* + P_a^*}{S^* + P^* + P_a^* + D_S^* + D_P^* + D_a^* + D_L^*}$. We begin by substituting in the steady-state

914     population abundances to the expression $\frac{\nu_B^*}{1 - \nu_B^*} = \frac{S^* + P^* + P_a^*}{D_S^* + D_P^* + D_a^* + D_L^*}$,

$$\frac{\nu_B^*}{1 - \nu_B^*} = \frac{S^* + P^* + P_a^*}{(S^* + P^* + P_a^*)\left( \frac{\omega_B}{\delta + \omega_D} \right) + \frac{\gamma P_a^*}{\delta + \omega_{DL}}}, \tag{S18}$$

916     which when solved for $\omega_B$ yields

$$\omega_B = (\delta + \omega_D) \left( \frac{1 - \nu_B^*}{\nu_B^*} - \frac{\gamma P_a^*}{(\delta + \omega_{DL})(S^* + P^* + P_a^*)} \right). \tag{S19}$$

918     This equation provides an upper bound estimate for $\omega_B$:

$$\omega_B \leq (\delta + \omega_D) \left( \frac{1 - \nu_B^*}{\nu_B^*} \right), \tag{S20}$$

which will be utilized later in deriving the estimate of total lysis rate from prophage copy

number. The value of $v_B^*$ in stool has been estimated at $\sim 0.5 - 0.8$ based on cell

permeability measurements, leading to $\frac{1-v_B^*}{v_B^*} \approx 1$ (87,88). Thus, the bacterial death rate

is at most similar in magnitude to the sum of dilution and cell degradation rate.

From **Eq. S18**, we can also derive an estimate for the total lysis rate using the steady-

state expression $P_a^* = \frac{\eta(S^*+P^*)}{(\gamma+\delta+\omega_D)}$. Solving **Eq. S18** for $\eta$ yields:

$$\frac{\eta}{\gamma + \delta + \omega_D} = \frac{\left(\dfrac{1 - v_B^*}{v_B^*} - \dfrac{\omega_B}{\delta + \omega_D}\right)}{\dfrac{\gamma}{\delta + \omega_{DL}} - \left(\dfrac{1 - v_B^*}{v_B^*} - \dfrac{\omega_B}{\delta + \omega_D}\right)}. \tag{S21}$$

This equation is potentially usable to provide another lysis rate estimate, and in the

prophage-dominated regime becomes an induction rate estimate. However, the values

of $\omega_D$, $\omega_B$, and $\omega_{DL}$ are currently poorly constrained. For example, one cell viability

measurement method is based on comparing the fraction of 16S rDNA found inside and

outside of intact cells (87) and it is not known how rapidly extracellular DNA is degraded

inside the gut. There are also potential technical issues in the measurement of $v_B$, as it

is unclear to what extent cells lysed by phage are detected by current cell viability

measurements. If the lysis process degrades the host genome or leads to total

destruction of the cellular structure, the population of cells dying due to lysis would be

underestimated by methods relying on extracellular genomic DNA or permeable cell

remains.

**Total lysis rate and induction rate estimates from integrated prophage copy**

**number**

Here, we estimate the average total lysis rate using lysogeny copy number $R$. We

assume that $R$ includes the contribution of viral particles, dead viral particles, and all

dead cells, and we show that the inclusion of these classes does not substantially alter

our induction rate estimation. Each activated cell contributes $B_a$ prophage copies, each

lysogen cell contributes one prophage copy, and lysed cells contribute no prophage

copies. All cells contribute a single bacteria genome copy. We first define $R$ in terms of

our steady-state model populations:

$$R^* \equiv \text{gVMR} = \frac{V^* + D_V^* + P^* + B_a P_a^* + D_P^* + B_a D_a^*}{S^* + P^* + P_a^* + D_S^* + D_P^* + D_a^* + D_L^*}. \qquad \text{(S22)}$$

950 As all cells have the same non-lysis mortality rate, **Eq. S22** can be rearranged to

$$R^* = \text{pVMR} + \frac{P^* + B_a P_a^*}{S^* + P^* + P_a^*}\left(1 + \frac{D_P^*}{P^*}\right)v_B^*, \qquad \text{(S23)}$$

$$R^* - \text{pVMR} = \frac{P^* + B_a P_a^*}{S^* + P^* + P_a^*}\left(1 - \frac{D_L^*}{S^* + P^* + P_a^* + D_S^* + D_P^* + D_a^* + D_L^*}\right). \qquad \text{(S24)}$$

953 We can then substitute in the steady-state population values to express all dead cell

954 populations in terms of living cells populations:

$$R^* - \text{pVMR} = \frac{P^* + B_a P_a^*}{S^* + P^* + P_a^*}\left(1 - \frac{\dfrac{\gamma P_a^*}{\delta + \omega_{DL}}}{(S^* + P^* + P_a^*)\left(1 + \dfrac{\omega_B}{\delta + \omega_D}\right) + \dfrac{\gamma P_a^*}{\delta + \omega_{DL}}}\right). \qquad \text{(S25)}$$

956 Rearranging and using the fact that $\gamma = \left(\frac{f_\gamma}{1-f_\gamma}\right)(\delta + \omega_B)$ yields

$$R^* - \text{pVMR} = \left(\frac{P^* + B_a P_a^*}{S^* + P^* + P_a^*\left[1 + \left(\dfrac{\delta + \omega_D}{\delta + \omega_{DL}}\right)\left(\dfrac{f_\gamma}{1 - f_\gamma}\right)\left(\dfrac{\delta + \omega_B}{\delta + \omega_D + \omega_B}\right)\right]}\right), \qquad \text{(S26)}$$

958    which when solved for $P_a^*$ yields

$$P_a^* = \frac{(S^* + P^*)(R^* - \text{pVMR}) - P^*}{B_a - (R^* - \text{pVMR})Q}, \tag{S27}$$

960    where $Q = 1 + \left(\frac{\delta + \omega_D}{\delta + \omega_{DL}}\right)\left(\frac{f_\gamma}{1 - f_\gamma}\right)\left(\frac{\delta + \omega_B}{\delta + \omega_D + \omega_B}\right)$. From our steady-state solution for $P_a^*$ we

961    have that $\eta^* = P_a^* \left(\frac{\gamma + \omega_B + \delta}{S^* + P^*}\right)$, providing an estimate of $\eta^*$:

$$\eta^* = (\gamma + \omega_B + \delta)\frac{(R^* - \text{pVMR}) - x_P^*}{B_a - (R^* - \text{pVMR})Q}. \tag{S28}$$

963    The effect of dead cells and viruses enters the expression via the factor $Q$, which will

964    inflate the lysis rate. However, this term cannot be greater than $O(1)$, and thus if $B_a$ is

965    large, the impact of dead material is minimal. Empirically, the value of $\omega_B$ is poorly

966    constrained, but we can use results from the cell viability derivation above (**Eq. S20**) to

967    relate this rate to the cell viability fraction $\nu_B$ and the degradation rate of dead cells $\omega_D$:

$$\eta^* \leq \left[\gamma + (\delta + \omega_D)\left(\frac{1 - \nu_B^*}{\nu_B^*}\right) + \delta\right]\frac{(R^* - \text{pVMR}) - x_P^*}{B_a - (R^* - \text{pVMR})Q}. \tag{S29}$$

969    To reach the order of magnitude bound shown in the main text, we assume prophage

970    dominance $x_P^* \approx 1$, that the number of prophage copies in activated cells is similar to the

971    burst size $B_a \approx \tilde{B}$, and that dead cells are primarily removed by dilution $\omega_D \ll \delta$. Based

972    on empirical measurements, we also assume that $R^* \gg \text{pVMR}$, $\tilde{B} \gg 1$, and $\frac{1 - \nu_B^*}{\nu_B^*} \approx 1$,

973    yielding

$$\xi^* \lesssim \frac{(\gamma + \delta)(R^* - 1)}{(\tilde{B} - 1) - (R^* - 1)}. \tag{S30}$$

We now briefly discuss potential bioinformatic/sequencing technical artifacts that could

influence the measurement of $R$. One potential factor that could systematically skew the

above induction rate estimate is sequencing bias between prophage and host (e.g., due

to GC content differences between host bacteria and prophage (89)). However, based

on the negative control analyses performed by (48), these biases do not appear

significant, as non-induced prophage had $R \approx 1$. If large biases existed, $R$ in non-

induced phage would differ significantly from $1$. Another possible confounding factor in

estimating $R$ is the presence of the prophage within only a subpopulation of the bacterial

host, leading to a lower value of $R$. However, this is unlikely to affect our current

analyses, as the $R$ values we analyze were computed based on metagenomically

assembled contigs containing both prophage and bacterial host sequence. Assembly of

such mixed contigs from a mixed lysogen/sensitive population is highly unlikely due to

degeneracies in the possible assembly paths. In the case of both possible biases, our

framework can readily accommodate improved estimates of $R$ as sequencing and

bioinformatic approaches improve.

**Extension of the model to multiple phage and bacterial species in time-varying**

**environments**

We now generalize our model to complex communities in time-varying environments.

For simplicity, we begin with the timescale-separated version of the model and focus on

the prophage-dominated case in which most lysis is due to induction, as for the single

bacteria-phage regime studied above.

We now track the dynamics of multiple types of phage (indexed by $i$) and multiple types

of bacteria (indexed by $j$), such that the total number of phage particles is $V(t) \equiv$

$\sum_i V_i(t)$ and the total number of bacteria is $N(t) \equiv \sum_j N_j(t)$. The bacterial "type" $j$

encompasses both the taxonomic identity of a bacteria and its infection status (i.e., the

$N_j$ also include bacteria infected by a prophage). To keep track of infection status and

the phage-bacteria interaction network, we introduce bookkeeping parameters $I_{ij}$ and

$A_{ijk}$, respectively. The parameter $I_{ij}$ is 1 if bacteria $j$ is infected with a prophage of

phage $i$ and zero otherwise. Thus, the total number of prophage in this system is $P(t) \equiv$

$\sum_i P_i(t) = \sum_{ij} I_{ij} N_j(t)$. Note that generally $P(t) \neq N(t)$, as multiple phage can infect a

single bacteria. The second parameter, $A_{ijk}$, is 1 if an infection of bacteria of type $j$ by

phage $i$ produces an infected bacterium of type $k$, and 0 otherwise. Using this notation,

we now define the multispecies generalization of **Eq. S3**:

1011   
$$\frac{dN_j}{dt} = \mu_j(t)N_j - \sum_i \kappa_{ij}(t)N_jV_i + \sum_{i,k} A_{ikj}\kappa_{ij}(t)N_kV_i - \sum_i I_{ij}\xi_{ij}(t)N_j - \delta(t)N_j - \omega_{B,j}(t)N_j, \text{(S31a)}$$

1012   
$$\frac{dV_i}{dt} = \sum_j I_{ij}B_{ij}(t)\xi_{ij}(t)N_j - \sum_j \kappa_{ij}(t)N_jV_i - \sum_j r_{ij}(t)\kappa_{ij}(t)N_jV_i - \delta(t)V_i - \omega_i(t)V_{V,i}, \text{(S31b)}$$

1013    where we have also allowed the rate parameters to explicitly depend on time. We have

1014    also approximated $f_L = 1$ for simplicity.

1015

1016    To relate these dynamics to the total pVMR, we now sum **Eq. S13b** over the viral index $i$

1017    and substitute $\frac{dx}{dt}\left(\frac{1}{x}\right) = \frac{d\log(x)}{dt}$, yielding

1018   
$$\frac{d\log V}{dt} = \frac{\sum_{ij} I_{ij}B_{ij}(t)\xi_{ij}(t)N_j}{\sum_{ij} I_{ij}N_j} \cdot \left(\frac{P}{V}\right) - \frac{\sum_{ij}\kappa_{ij}(t)N_jV_i}{\sum_i V_i} - \frac{\sum_{ij} r_{ij}(t)\kappa_{ij}(t)N_jV_i}{\sum_i V_i} - \delta(t) - \frac{\sum_{ij}\omega_{V,i}(t)V_i}{\sum_i V_i}. \text{(S32)}$$

1019    **Eq. S14** can be rewritten in a more compact form as

1020   
$$\frac{d\log V}{dt} = \overline{B}(t)\overline{\xi}(t)\left(\frac{P}{V}\right) - \overline{\psi}_I(t) - \overline{\psi}_N(t) - \delta(t) - \overline{\omega}_V(t), \qquad \text{(S33)}$$

1021 where we have defined the microbiome averages

$$\overline{B}(t) \equiv \frac{\sum_{ij} I_{ij} B_{ij}(t) \xi_{ij}(t) N_j}{\sum_{ij} I_{ij} \xi_{ij}(t) N_j}, \tag{S34a}$$

$$\overline{\xi}(t) \equiv \frac{\sum_{i,j} I_{ij} \xi_{ij}(t) N_j}{\sum_{ij} I_{ij} N_j}, \tag{S34b}$$

$$\overline{\psi}_I(t) \equiv \frac{\sum_{ij} \kappa_{ij}(t) N_j V_i}{\sum_i V_i}, \tag{S34c}$$

$$\overline{\psi}_N(t) \equiv \frac{\sum_{ij} r_{ij}(t) \kappa_{ij}(t) N_j V_i}{\sum_i V_i}. \tag{S34d}$$

$$\overline{\omega}_V(t) \equiv \frac{\sum_i \omega_{V,i}(t) V_i}{\sum_i V_i} \tag{S34e}$$

1027 Integrating **Eq. S15** over long times yields

$$0 \approx \overline{B}^* \overline{\xi}^* \left(\frac{P}{V}\right)^* - \overline{\psi}_I^* - \overline{\psi}_N^* - \delta^* - \overline{\omega}_V^*, \tag{S35}$$

where the asterisks again denote the time-averaged value $x^* = \frac{1}{\Delta t} \int_0^{\Delta t} x(t) \, dt$. Since the

microbiome is in a statistical steady state over long times (**Fig. 1B**), we can estimate the

averages over time by taking an average over independent hosts. This procedure yields

a connection between the rate parameters and the VLP-to-prophage ratio from **Fig. 1**

and thus a lower bound similar to the one estimated from the single phage-bacteria

model:

$$\overline{\xi}^* \gtrsim \frac{1}{\overline{B}^*} \left(\frac{V^*}{P^*}\right) \delta^*. \tag{S36}$$

1036 This bound assumes that the burst size, induction rate, and particle-to-prophage ratio

1037 are largely uncorrelated in time. If this assumption is violated, the estimate represents a

1038 particular weighted average of the induction rate bound:

$$1039 \qquad \frac{1}{\overline{B}^*}\left(\frac{V^*}{P^*}\right)\delta^* = \left(\frac{1}{\Delta t}\right)\frac{\int_0^{\Delta t} \overline{\overline{\xi}}\,\overline{B}\left(\frac{P}{V}\right)dt}{\overline{B}^*\left(\frac{V^*}{P^*}\right)}. \qquad (S37)$$

1040 To generalize this bound to the case of pVMR including dead material, we begin with

1041 the multispecies version of the dead virus dynamics:

$$1042 \qquad \frac{dD_{V,i}}{dt} = \omega_{V,i}(t)V_i - \delta(t)D_{V,i} - \omega_{DV,i}(t)D_{V,i}, \qquad (S38)$$

1043 which then yields an expression for the dynamics of the total dead virus population:

$$1044 \qquad \frac{d\log D_{V,i}}{dt} = \overline{\omega}_V(t)\left(\frac{V}{D_V}\right) - \delta(t) - \overline{\omega}_{DV,i}(t), \qquad (S39)$$

1045 where $\overline{\omega}_{DV,i}(t) \equiv \frac{\sum_i \omega_{DV,i}(t)D_{V,i}}{\sum_i D_{V,i}}$. At statistical steady state, this leads to $\frac{v_V^*}{1-v_V^*} = \frac{V^*}{D_V^*} \approx \frac{\delta^* + \overline{\omega}_{DV}^*}{\overline{\omega}_V^*}$,

1046 which when combined with **Eq. S35** shows the lower bound is preserved when the

1047 pVMR accounts for dead populations, as in the earlier single species derivation.

1048

1049 Note that in this section we have only analyzed a simple case of this community model,

1050 and further analysis, such as exploring the role of temporal correlations and the relative

1051 contribution of induction and direct lysis, is a promising direction for future theoretical

1052 phage ecology work.

1053

Here, we have shown the multispecies generalization of the induction rate estimate from

VLP-to-prophage ratio. The rate estimates computed from cell viability will similarly

extend to the multispecies context, as we model all sources of death in aggregate,

independent of which phage causes lysis. The induction rate estimate from the lysogen

copy number is performed on a prophage-by-prophage basis, hence it is not affected in

a multispecies context. However, this context will lead to a difference in the kind of

average used in the estimate: unlike the average computed from the VLP-to-prophage

ratio, the average from copy number average is not abundance weighted and includes

only lysogens captured with their host contig.

1063

1064 **Data Availability**

1065 All code used in this manuscript is available at

1066 https://github.com/jaimegelopez/gut_phage_quantification. All data analyzed in this

1067 manuscript is publicly available. Processed final versions of the datasets (e.g. estimated

1068 taxonomic compositions) are available in the GitHub repository.

 **Supplementary Figures**

1070

**Figure S1: Species-level relative abundance analysis of the overlap between**

**phage community quantifications using VLP- and stool-based approaches.** Data

and plotting methods are the same as **Fig. 1C**, except that relative abundance was

used instead of absolute genome/particle density. This figure includes only phage

shared between the VLP fraction and stool, corresponding to the central scatter plot of

**Fig. 1C**. Relative abundance was defined relative to total taxonomic abundance of

phage within VLPs/stool. Note that without the absolute abundance normalization

employed in **Fig. 1C**, adult VLP abundances appear to be substantially lower than infant

abundances.

1080

**Figure S2: Species-level absolute abundance analysis of the overlap between VLP- and stool-based phage community quantifications, colored by predicted virulence.** This figure is the same as **Fig. 1C**, except points are colored by predicted virulence rather than their dataset of origin. Diamond markers represent points from adult samples, while squares represent points from infant samples.

1086

**Figure S3: Time series of various reconstruction evaluation metrics in hCom2-colonized mice.** Each row is a version of **Fig. 3C,D** but instead showing Spearman correlation of taxa read abundances (A,B), Spearman correlation of read abundances of species shared between hCom2/hMock sample pairs (C,D), Jaccard index (E,F), fraction of read abundances in hCom2-colonized mouse feces that are nonzero in hMock (G,H), mean squared error (MSE) of $\log_{10}$(read abundance) of species shared

between hCom2/hMock sample pairs (I,J), and MSE of $\log_{10}$(read abundance)

computed with a relative abundance pseudocount of $10^{-7}$ (K,L). All metrics were

computed using species-level relative read abundances. Jaccard index is the number of

shared species between an hCom2/hMock sample pair divided by the total number of

species with nonzero abundance in at least one of the two samples. For (G,H), shared

read abundances were normalized to the total bacterial or phage abundance in the

hCom2-colonized mouse fecal sample.

1100

**Figure S4: Comparison of phage and bacterial community properties in the feces of hCom2-colonized mice across conditions.** Community properties and plotting methods are the same as in **Fig. 4.** 'Unchallenged' corresponds to *in vivo* samples from hCom2-colonized gnotobiotic mice that have not been exposed to a human stool sample, while 'Challenged' denotes samples from hCom2-colonized mice that have been exposed to a human stool challenge. '*in vitro*' corresponds to hCom2 communities grown *in vitro*.

1108

**Figure S5: Comparison of the virulent-to-temperate ratio (VTR) between stool**

**from humans and feces from hCom2-colonized mice, as estimated by PhaTYP.**

Equivalent to **Fig. 4D**, except the virulence prediction of phage genomes was performed

using PhaTYP (57) instead of using the Phanta UHGV database (24).

## References

1. Routy B, Le Chatelier E, Derosa L, Duong CP, Alou MT, Daillère R, et al. Gut microbiome influences efficacy of PD-1–based immunotherapy against epithelial tumors. Science. 2018;359(6371):91–7.

2. Thiele Orberg E, Meedt E, Hiergeist A, Xue J, Heinrich P, Ru J, et al. Bacteria and bacteriophage consortia are associated with protective intestinal metabolites in patients receiving stem cell transplantation. Nat Cancer. 2024 Jan 3;5(1):187–208.

3. Zhang F, Aschenbrenner D, Yoo JY, Zuo T. The gut mycobiome in health, disease, and clinical applications in association with the gut bacterial microbiome assembly. Lancet Microbe. 2022 Dec;3(12):e969–83.

4. Nayfach S, Páez-Espino D, Call L, Low SJ, Sberro H, Ivanova NN, et al. Metagenomic compendium of 189,680 DNA viruses from the human gut microbiome. Nat Microbiol. 2021;6(7):960–70.

5. Camarillo-Guerrero LF, Almeida A, Rangel-Pineros G, Finn RD, Lawley TD. Massive expansion of human gut bacteriophage diversity. Cell. 2021;184(4):1098–109.

6. Guerin E, Shkoporov AN, Stockdale SR, Comas JC, Khokhlova EV, Clooney AG, et al. Isolation and characterisation of ΦcrAss002, a crAss-like phage from the human gut that infects Bacteroides xylanisolvens. Microbiome. 2021;9(1):1–21.

7. Shen J, Zhang J, Mo L, Li Y, Li Y, Li C, et al. Large-scale phage cultivation for commensal human gut bacteria. Cell Host Microbe. 2023;31(4):665–77.

8. Reyes A, Wu M, McNulty NP, Rohwer FL, Gordon JI. Gnotobiotic mouse model of phage–bacterial host dynamics in the human gut. Proc Natl Acad Sci. 2013;110(50):20236–41.

9. Sutcliffe SG, Shamash M, Hynes AP, Maurice CF. Common oral medications lead to prophage induction in bacterial isolates from the human gut. Viruses. 2021;13(3):455.

10. Borodovich T, Shkoporov AN, Ross RP, Hill C. Phage-mediated horizontal gene transfer and its implications for the human gut microbiome. Gastroenterol Rep. 2022;10:goac012.

11. Frazão N, Sousa A, Lässig M, Gordo I. Horizontal gene transfer overrides mutation in *Escherichia coli* colonizing the mammalian gut. Proc Natl Acad Sci. 2019 Sep 3;116(36):17906–15.

12. Johansen J, Atarashi K, Arai Y, Hirose N, Sørensen SJ, Vatanen T, et al. Centenarians have a diverse gut virome with the potential to modulate metabolism and promote healthy lifespan. Nat Microbiol. 2023 May 15;8(6):1064–78.

13. Brunse A, Deng L, Pan X, Hui Y, Castro-Mejía JL, Kot W, et al. Fecal filtrate transplantation protects against necrotizing enterocolitis. ISME J. 2022;16(3):686–94.

14. Ott SJ, Waetzig GH, Rehman A, Moltzau-Anderson J, Bharti R, Grasis JA, et al. Efficacy of sterile fecal filtrate transfer for treating patients with Clostridium difficile infection. Gastroenterology. 2017;152(4):799–811.

15. Sinha A, Li Y, Mirzaei MK, Shamash M, Samadfam R, King IL, et al. Transplantation of bacteriophages from ulcerative colitis patients shifts the gut bacteriome and exacerbates the severity of DSS colitis. Microbiome. 2022;10(1):1–23.

16. Adiliaghdam F, Amatullah H, Digumarthi S, Saunders TL, Rahman RU, Wong LP, et al. Human enteric viruses autonomously shape inflammatory bowel disease phenotype through divergent innate immunomodulation. Sci Immunol. 2022;7(70):eabn6660.

17. Fuhrman JA. Marine viruses and their biogeochemical and ecological effects. Nature. 1999 Jun;399(6736):541–8.

18. Suttle CA. Marine viruses — major players in the global ecosystem. Nat Rev Microbiol. 2007 Oct;5(10):801–12.

19. Hussain FA, Dubert J, Elsherbini J, Murphy M, VanInsberghe D, Arevalo P, et al. Rapid evolutionary turnover of mobile genetic elements drives bacterial resistance to phages. Science. 2021 Oct 22;374(6566):488–92.

20. Breitbart M, Bonnain C, Malki K, Sawaya NA. Phage puppet masters of the marine microbial realm. Nat Microbiol. 2018 Jun 4;3(7):754–66.

21. Thingstad TF. Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. Limnol Oceanogr. 2000 Sep;45(6):1320–8.

22. Maslov S, Sneppen K. Population cycles and species diversity in dynamic Kill-the-Winner model of microbial ecosystems. Sci Rep. 2017 Jan 4;7(1):39642.

23. Liang G, Zhao C, Zhang H, Mattei L, Sherrill-Mix S, Bittinger K, et al. The stepwise assembly of the neonatal virome is modulated by breastfeeding. Nature. 2020 May 28;581(7809):470–4.

24. Pinto Y, Chakraborty M, Jain N, Bhatt AS. Phage-inclusive profiling of human gut microbiomes with Phanta. Nat Biotechnol. 2023;1–12.

25. Shkoporov AN, Clooney AG, Sutton TD, Ryan FJ, Daly KM, Nolan JA, et al. The human gut virome is highly diverse, stable, and individual specific. Cell Host Microbe. 2019;26(4):527–41.

26. Kim MS, Park EJ, Roh SW, Bae JW. Diversity and abundance of single-stranded
DNA viruses in human feces. Appl Environ Microbiol. 2011;77(22):8062–70.

27. Dahlman S, Avellaneda-Franco L, Kett C, Subedi D, Young RB, Gould JA, et al.
Temperate gut phages are prevalent, diverse, and predominantly inactive. bioRxiv
[Internet]. 2023; Available from:
https://www.biorxiv.org/content/early/2023/08/18/2023.08.17.553642

28. Shkoporov AN, Khokhlova EV, Stephens N, Hueston C, Seymour S, Hryckowian AJ,
et al. Long-term persistence of crAss-like phage crAss001 is associated with phase
variation in Bacteroides intestinalis. BMC Biol. 2021 Dec;19(1):163.

29. Hoyles L, McCartney AL, Neve H, Gibson GR, Sanderson JD, Heller KJ, et al.
Characterization of virus-like particles associated with the human faecal and caecal
microbiota. Res Microbiol. 2014 Dec;165(10):803–12.

30. Parras-Moltó M, Rodríguez-Galet A, Suárez-Rodríguez P, López-Bueno A.
Evaluation of bias induced by viral enrichment and random amplification protocols in
metagenomic surveys of saliva DNA viruses. Microbiome. 2018;6(1):1–18.

31. Forterre P, Soler N, Krupovic M, Marguet E, Ackermann HW. Fake virus particles
generated by fluorescence microscopy. Trends Microbiol. 2013 Jan;21(1):1–5.

32. Sender R, Fuchs S, Milo R. Revised estimates for the number of human and
bacteria cells in the body. PLoS Biol. 2016;14(8):e1002533.

33. Kaletta J, Pickl C, Griebler C, Klingl A, Kurmayer R, Deng L. A rigorous assessment
and comparison of enumeration methods for environmental viruses. Sci Rep. 2020
Oct 29;10(1):18625.

34. Shkoporov AN, Ryan FJ, Draper LA, Forde A, Stockdale SR, Daly KM, et al.
Reproducible protocols for metagenomic analysis of human faecal phageomes.
Microbiome. 2018 Dec;6(1):68.

35. Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, et al. A unified
catalog of 204,938 reference genomes from the human gut microbiome. Nat
Biotechnol. 2021 Jan;39(1):105–14.

36. Schmidtke DT, Hickey AS, Liachko I, Sherlock GJ, Bhatt AS. Analysis and culturing
of the prototypic crAssphage reveals a phage-plasmid lifestyle. bioRxiv. 2024;2024–
03.

37. Wang X, Kim Y, Ma Q, Hong SH, Pokusaeva K, Sturino JM, et al. Cryptic prophages
help bacteria cope with adverse environments. Nat Commun. 2010 Dec 21;1(1):147.

38. Li G, Cortez MH, Dushoff J, Weitz JS. When to be temperate: on the fitness benefits
of lysis vs. lysogeny. Virus Evol. 2020 Jul 1;6(2):veaa042.

39. Stewart FM, Levin BR. The population biology of bacterial viruses: why be
temperate. Theor Popul Biol. 1984;26(1):93–117.

40. Geng Y, Nguyen TVP, Homaee E, Golding I. Using bacterial population dynamics to
count phages and their lysogens. Nat Commun. 2024 Sep 6;15(1):7814.

41. Bondy-Denomy J, Qian J, Westra ER, Buckling A, Guttman DS, Davidson AR, et al.
Prophages mediate defense against phage infection through diverse mechanisms.
ISME J. 2016 Dec 1;10(12):2854–66.

42. Wang Z, Goldenfeld N. Fixed points and limit cycles in the population dynamics of
lysogenic viruses and their hosts. Phys Rev E. 2010 Jul 22;82(1):011918.

43. Lee YY, Erdogan A, Rao SS. How to assess regional and whole gut transit time with
wireless motility capsule. J Neurogastroenterol Motil. 2014;20(2):265.

44. Shao Y, Wang IN. Effect of late promoter activity on bacteriophage λ fitness.
Genetics. 2009;181(4):1467–75.

45. Shkoporov AN, Khokhlova EV, Fitzgerald CB, Stockdale SR, Draper LA, Ross RP, et
al. ΦCrAss001 represents the most abundant bacteriophage family in the human gut
and infects Bacteroides intestinalis. Nat Commun. 2018;9(1):1–8.

46. Ramos-Barbero MD, Gómez-Gómez C, Sala-Comorera L, Rodríguez-Rubio L,
Morales-Cortes S, Mendoza-Barberá E, et al. Characterization of crAss-like phage
isolates highlights Crassvirales genetic heterogeneity and worldwide distribution.
Nat Commun. 2023;14(1):4295.

47. Parsons RJ, Breitbart M, Lomas MW, Carlson CA. Ocean time-series reveals
recurring seasonal patterns of virioplankton dynamics in the northwestern Sargasso
Sea. ISME J. 2012;6(2):273–84.

48. Kieft K, Anantharaman K. Deciphering active prophages from metagenomes.
mSystems. 2022;7(2):e00084-22.

49. Ellis EL, Delbruck M. The growth of bacteriophage. J Gen Physiol. 1939;22(3):365–
84.

50. Hadas H, Einav M, Fishov I, Zaritsky A. Bacteriophage T4 development depends on
the physiology of its host Escherichia coli. Microbiology. 1997;143(1):179–85.

51. Dennehy JJ, Wang IN. Factors influencing lysis time stochasticity in bacteriophage
λ. BMC Microbiol. 2011;11:1–12.

52. Mahmoudabadi G, Milo R, Phillips R. Energetic cost of building a virus. Proc Natl
Acad Sci. 2017;114(22):E4324–33.

53. Cheng AG, Ho PY, Aranda-Díaz A, Jain S, Feiqiao BY, Meng X, et al. Design, construction, and in vivo augmentation of a complex gut microbiome. Cell. 2022;185(19):3617–36.

54. Wilde J, Boyes R, Robinson AV, Daisley BA, Botschner AJ, Brettingham DJL, et al. Assessing phage-host population dynamics by reintroducing virulent viruses to synthetic microbiomes. Cell Host Microbe. 2024 May;32(5):768-778.e9.

55. Yachida S, Mizutani S, Shiroma H, Shiba S, Nakajima T, Sakamoto T, et al. Metagenomic and metabolomic analyses reveal distinct stage-specific phenotypes of the gut microbiota in colorectal cancer. Nat Med. 2019 Jun;25(6):968–76.

56. Hockenberry AJ, Wilke CO. BACPHLIP: predicting bacteriophage lifestyle from conserved protein domains. PeerJ. 2021 May 6;9:e11396.

57. Shang J, Tang X, Sun Y. PhaTYP: predicting the lifestyle for bacteriophages using BERT. Brief Bioinform. 2023 Jan 19;24(1):bbac487.

58. Reyes A, Haynes M, Hanson N, Angly FE, Heath AC, Rohwer F, et al. Viruses in the faecal microbiota of monozygotic twins and their mothers. Nature. 2010;466(7304):334–8.

59. Sutcliffe SG, Reyes A, Maurice CF. Bacteriophages playing nice: Lysogenic bacteriophage replication stable in the human gut microbiota. iScience. 2023 Feb;26(2):106007.

60. Silveira CB, Rohwer FL. Piggyback-the-Winner in host-associated microbial communities. Npj Biofilms Microbiomes. 2016 Jul 6;2(1):16010.

61. Berngruber TW, Lion S, Gandon S. Spatial structure, transmission modes and the evolution of viral exploitation strategies. PLoS Pathog. 2015;11(4):e1004810.

62. Messinger SM, Ostling A. The consequences of spatial structure for the evolution of pathogen transmission rate and virulence. Am Nat. 2009;174(4):441–54.

63. Knowles B, Silveira C, Bailey B, Barott K, Cantu V, Cobián-Güemes A, et al. Lytic to temperate switching of viral communities. Nature. 2016;531(7595):466–70.

64. Weitz JS, Beckett SJ, Brum JR, Cael BB, Dushoff J. Lysis, lysogeny and virus–microbe ratios. Nature. 2017 Sep 21;549(7672):E1–3.

65. Oh JH, Lin XB, Zhang S, Tollenaar SL, Özçam M, Dunphy C, et al. Prophages in Lactobacillus reuteri are associated with fitness trade-offs but can increase competitiveness in the gut ecosystem. Appl Environ Microbiol. 2019;86(1):e01922-19.

66. Yang K, Niu J, Zuo T, Sun Y, Xu Z, Tang W, et al. Alterations in the Gut Virome in Obesity and Type 2 Diabetes Mellitus. Gastroenterology. 2021 Oct;161(4):1257-1269.e13.

67. de Jonge PA, Wortelboer K, Scheithauer TP, van den Born BJH, Zwinderman AH, Nobrega FL, et al. Gut virome profiling identifies a widespread bacteriophage family associated with metabolic syndrome. Nat Commun. 2022;13(1):3594.

68. Jiang L, Lang S, Duan Y, Zhang X, Gao B, Chopyk J, et al. Intestinal Virome in Patients With Alcoholic Hepatitis. Hepatology. 2020 Dec;72(6):2182–96.

69. Grafen A. The phylogenetic regression. Philos Trans R Soc B. 1989;326(1233):119–57.

70. Diard M, Bakkeren E, Cornuault JK, Moor K, Hausmann A, Sellin ME, et al. Inflammation boosts bacteriophage transfer between *Salmonella* spp. Science. 2017 Mar 17;355(6330):1211–5.

71. Boling L, Cuevas DA, Grasis JA, Kang HS, Knowles B, Levi K, et al. Dietary prophage inducers and antimicrobials: toward landscaping the human gut microbiome. Gut Microbes. 2020;11(4):721–34.

72. Hu J, Ye H, Wang S, Wang J, Han D. Prophage activation in the intestine: insights into functions and possible applications. Front Microbiol. 2021;3930.

73. Shkoporov AN, Stockdale SR, Lavelle A, Kondova I, Heuston C, Upadrasta A, et al. Viral biogeography of the mammalian gut and parenchymal organs. Nat Microbiol. 2022 Aug 2;7(8):1301–11.

74. Wu J, Fu K, Hou C, Wang Y, Ji C, Xue F, et al. Bacteriophage defends murine gut from Escherichia coli invasion via mucosal adherence. Nat Commun. 2024 Jun 4;15(1):4764.

75. Barr JJ, Auro R, Furlan M, Whiteson KL, Erb ML, Pogliano J, et al. Bacteriophage adhering to mucus provide a non–host-derived immunity. Proc Natl Acad Sci. 2013;110(26):10771–6.

76. Donaldson GP, Lee SM, Mazmanian SK. Gut biogeography of the bacterial microbiota. Nat Rev Microbiol. 2016 Jan;14(1):20–32.

77. Shalon D, Culver RN, Grembi JA, Folz J, Treit PV, Shi H, et al. Profiling the human intestinal environment under physiological conditions. Nature. 2023 May 18;617(7961):581–91.

78. Garmaeva S, Sinha T, Gulyaeva A, Kuzub N, Spreckels JE, Andreu-Sánchez S, et al. Transmission and dynamics of mother-infant gut viruses during pregnancy and early life. Nat Commun. 2024 Mar 2;15(1):1945.

79. McNulty R, Sritharan D, Pahng SH, Meisch JP, Liu S, Brennan MA, et al. Probe-
based bacterial single-cell RNA sequencing predicts toxin regulation. Nat Microbiol.
2023 Apr 3;8(5):934–45.

80. Bikel S, López-Leal G, Cornejo-Granados F, Gallardo-Becerra L, García-López R,
Sánchez F, et al. Gut dsDNA virome shows diversity and richness alterations
associated with childhood obesity and metabolic syndrome. iScience.
2021;24(8):102900.

81. Bushnell B. BBTools [Internet]. Available from: sourceforge.net/projects/bbmap/

82. Angly FE, Willner D, Rohwer F, Hugenholtz P, Tyson GW. Grinder: a versatile
amplicon and shotgun sequence simulator. Nucleic Acids Res. 2012 Jul
1;40(12):e94–e94.

83. Terzian P, Olo Ndela E, Galiez C, Lossouarn J, Pérez Bucio RE, Mom R, et al.
PHROG: families of prokaryotic virus proteins clustered using remote homology.
NAR Genomics Bioinforma. 2021 Jun 23;3(3):lqab067.

84. Camargo AP, Roux S, Schulz F, Babinski M, Xu Y, Hu B, et al. Identification of
mobile genetic elements with geNomad. Nat Biotechnol. 2023 Sep 21;

85. Millan AS, Peña-Miller R, Toll-Riera M, Halbert ZV, McLean AR, Cooper BS, et al.
Positive selection and compensatory adaptation interact to stabilize non-
transmissible plasmids. Nat Commun. 2014 Oct 10;5(1):5208.

86. De Paepe M, Tournier L, Moncaut E, Son O, Langella P, Petit MA. Carriage of λ
Latent Virus Is Costly for Its Bacterial Host due to Frequent Reactivation in
Monoxenic Mouse Intestine. Hughes D, editor. PLOS Genet. 2016 Feb
12;12(2):e1005861.

87. Acosta EM, Little KA, Bratton BP, Lopez JG, Mao X, Payne AS, et al. Bacterial DNA
on the skin surface overrepresents the viable skin microbiome. eLife.
2023;12:RP87192.

88. Maurice CF, Haiser HJ, Turnbaugh PJ. Xenobiotics Shape the Physiology and Gene
Expression of the Active Human Gut Microbiome. Cell. 2013 Jan;152(1–2):39–50.

89. Browne PD, Nielsen TK, Kot W, Aggerholm A, Gilbert MTP, Puetz L, et al. GC bias
affects genomic and metagenomic reconstructions, underrepresenting GC-poor
organisms. GigaScience. 2020 Feb 1;9(2):giaa008.

1350