Recurrent introgression and geographical stratification shape *Saccharomyces cerevisiae* in the Neotropics

J Abraham Avelar-Rivas[1], Iván Sedeño[2,3], Luis Fernando García-Ortega[4], Jose A Urban Aragon[2], Eugenio Mancera[4,*], Alexander DeLuna[1,*], Lucía Morales[2,*]

[1]*Unidad de Genómica Avanzada, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Irapuato, Mexico.*

[2] *Laboratorio Internacional de Investigación sobre el Genoma Humano (LIIGH), Universidad Nacional Autónoma de México, 76230 Juriquilla, Mexico*

[3] *Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Coyoacán, 04510 CDMX, Mexico*

[4] *Departamento de Ingeniería Genética, Unidad Irapuato, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Irapuato, Mexico.*

*Correspondence to: eugenio.mancera@cinvestav.mx, alexander.deluna@cinvestav.mx, lmorales@liigh.unam.mx

Running title: Introgressions and population structure in *S. cerevisiae* from the Neotropics

Key words: Evolutionary genomics, Genetic introgression, Gene flow, Geographic population structure, *Saccharomyces cerevisiae*, Agave fermentation

Highlights:

1. Phylogenetic analyses of *S. cerevisiae* isolates from agave fermentation reveal novel clades within the Neotropical cluster.
2. The Neotropical cluster shows geographical population structure.
3. Multiple pulses of interspecific introgression are part of the evolutionary history of *S. cerevisiae* in the Neotropics

ABSTRACT

From yeasts to humans, introgressive hybridization significantly influences the evolutionary history of living organisms by introducing new genetic diversity. Strains of *Saccharomyces cerevisiae* worldwide exhibit introgressions from the sister species *S. paradoxus*, despite the average sequence identity between these species being lower than 90%. While *S. cerevisiae* isolates from the Neotropics are known for their high levels of introgression, the evolutionary events leading to the unusually high prevalence of them remain unclear. Here, we sequenced 216 *S. cerevisiae* isolates living in open agave fermentation across Mexico, a habitat at the interface of natural and industrial environments. The genomes of these strains revealed considerable genetic diversity and population structure linked to geographic distribution, which had been overlooked due to undersampling of this megadiverse region. These strains, along with those from French Guiana, Ecuador, and Brazil, form a broader Neotropical phylogenetic cluster that is notably enriched with introgressions. Surprisingly, the origins and conservation patterns of introgressions indicate multiple hybridization events, suggesting an unprecedented scenario of flexible species barriers in this region. Our findings underscore concurrent evolutionary processes—geographical stratification and multiple introgressions—that shape the genomes of a diverse lineage of *S. cerevisiae*. Neotropical yeasts thus provide a natural laboratory for exploring the mechanisms and adaptive significance of introgressive hybridization in eukaryotic genome evolution.

INTRODUCTION

Beyond its long-established status as the most extensively studied eukaryote in the laboratory, the budding yeast *S. cerevisiae* has emerged as an outstanding model for ecology, population genomics, and evolution (Liti, 2015; Marcet-Houben and Gabaldón, 2015; Alsammar and Delneri, 2020; Bendixsen et al., 2021; Wang et al., 2024). Genomic analyses of thousands of isolates from natural and human-related habitats around the world reveal that the species is structured in clades that broadly correlate with geography and habitat of isolation (Peter et al., 2018; Pontes et al., 2020; Loegler et al., 2024). These global genomic studies have also shown that most strains harbor genetic material from other *Saccharomyces* species, a phenomenon referred to as genetic introgression. For instance, most *S. cerevisiae* isolates from around the world harbor genes introgressed from its sister species *Saccharomyces paradoxus* (Duan et al., 2018; Strope et al., 2015; Peter et al., 2018; Clark et al., 2022; Tellini et al., 2024)*,* even though these species diverged 4-6 MYA and have an average nucleotide divergence of ~14% (Boynton and Greig, 2014; Shen et al., 2018).

The Neotropical region, stretching from northern Mexico to South America, is biogeographically significant due to its high biodiversity (Morrone, 2014). Some of the *S. cerevisiae* strains from this area isolated from anthropogenic and natural environments group in a long-monophyletic branch. This diverged branch, referred to as a complex clade by Pontes et al. (2020), aggregates three groups of strains, namely Mexican Agave (MA), French Guiana (FG), and South American Mix 2 (SAM2). Notoriously, the handful of sequenced genomes available to date indicate that this Neotropical cluster harbors an

unusually high and heterogeneous number of introgressions from *S. paradoxus* (Peter et al.,

2018; Pontes et al., 2020; De Chiara et al., 2022; Tellini et al., 2024). Considering the vast

area encompassed by this region, the few available sequenced isolates likely provide an

underestimate of genetic diversity, thus limiting our understanding of the species

evolutionary history in this part of the world.

The elevated numbers of introgressions observed in *S. cerevisiae* strains from the

Neotropical cluster could explain its high divergence when compared to other clades.

Although a fraction of the introgressions from the Neotropical cluster has been associated

with populations of *S. paradoxus* from the Americas (Barbosa et al., 2016; Peter et al., 2018;

Tellini et al., 2024), their precise origins remain to be determined. In Mexico, there are

reports of *S. paradoxus* in fermentations (Gallegos-Casillas et al., 2024; Peris et al., 2023),

including a lineage that coexists with *S. cerevisiae* in the agave distillery environment and

that has not been considered by previous studies (Gallegos-Casillas et al., 2024). Therefore,

this *S. paradoxus* lineage could be a potential donor for introgressions in the tropical

Americas. Comprehensive understanding of both the vertical evolution of *S. cerevisiae* and

the origins of its introgressions is essential to accurately determine the dynamics of

interspecies gene flow in the region.

To explore the evolutionary history and genomic diversity of *S. cerevisiae* in the tropical

Americas in the context of their strong signatures of introgression, we sequenced 216 strains

primarily isolated from open agave fermentations in Mexico. Previous studies have

highlighted the megadiversity in this region (Mittermeier and Mittermeier, 2005) as well as

high diversity of *S. cerevisiae* in agave fermentation (Lachance, 1995; Verdugo Valdez et

al., 2011; Álvarez-Ainza et al., 2015; Kirchmayr et al., 2017), but whole-genome sequencing efforts have only included nine strains, mostly from a single agave fermenting region (Peter et al., 2018). We show that most of the newly sequenced isolates belong to the Neotropical cluster, contributing new clades that group according to the geographic origin of the strains–i.e. with a geographical population structure. Moreover, by analyzing the patterns and origins of the introgressions in the context of the relatedness of these lineages, we identified multiple introgression events in their evolutionary history. Notably, some of these introgression pulses were unique to strains isolated from agave fermentations. Our refined phylogeny allowed us to illustrate how recurrent episodes of gene flow between two sister *Saccharomyces* species have unfolded in a megadiverse region of the world.

RESULTS

To provide a comprehensive view of *S. cerevisiae* diversity in one of the most biodiverse regions in the world (Mittermeier and Mittermeier, 2005; Morrone, 2014), we sequenced the genomes of 216 isolates across Mexico (Supplementary Figure S1, Supplementary Table S1). Strains were primarily isolated from traditional agave distilleries. This fermentation environment provides a good starting point to study yeasts diversity, as spontaneous fermentations have taken place for over 3,500 years in a wide variety of ecosystems, latitudes, and cultural contexts (Bruman, 2000; Arellano-Plaza et al., 2022; Colón-González et al., 2024). Specifically, 211 of the sequenced strains were isolated from agave fermentations collected through various sampling efforts conducted across the country between the years 1988 and 2021 (Padilla-Camberos et al., 1994; Lachance, 1995; Kirchmayr et al., 2014, 2017; Quezada et al., 2016; Verdugo Valdez et al., 2011; Lopez-Gallegos, 2023; Gallegos-Casillas

et al., 2024). Four of the five remaining isolates were from fermentations of other substrates

— pulque (fermented raw agave sap), apple cider, cactus pears, and cooked stems of sotol

plants (*Dasylirion* spp.)— while the fifth was an uncharacterized commercial yeast strain

used in one of the sampled distilleries; all of them were isolated in Mexico (Supplementary

Table S1). With the 216 genomes (median depth coverage = 163X), we assessed the

phylogenetic position and population structure of the strains in the context of the known

diversity of the species and we also analyzed the dynamics and origins of the prevalent *S.*

paradoxus introgressions within these populations.

Agave fermentation strains belong to the Neotropical cluster

To assess the relatedness of the agave *S. cerevisiae* strains with other isolates of the species,

we first performed multidimensional scaling analysis (MDS) with 1,264 genomes, including

the 216 newly sequenced strains. Most isolates from Mexico clustered next to other strains

from the tropical Americas, such as those from the French Guiana (FG), the South American

Mix 2 (SAM2), and the Mexican Agave 1 (MA1, formerly named "Mexican Agave") clades

(Figure 1A). Only five strains grouped with other clades, namely the Mixed-origin, Wild

North American, or Wine clades. A simple population-structure model (ADMIXTURE,

K=3) showed that ~97% of the agave fermentation strains had a Neotropical ancestral genetic

component (Figure 1B), distinguishing strains from the Neotropics (blue) from wild (green)

and domesticated (red) isolates from other regions of the world. The ADMIXTURE analysis

also indicated that strains from the tequila industry were highly admixed, bearing a

combination of markers from the three ancestral components defined at this specific level

(Figure 1B). All five isolates from non-agave substrates that we sequenced grouped close to

the agave-fermentation strains from the Neotropics.

Figure 1. A novel, diverse yeast lineage encompasses most of the diversity in open agave fermentations. (A) MDS analysis of 1,264 genomes (173,331 informative sites). Large dots are strains from the Tropical Americas Neotropics, colored according to their clade: "Mexican Agave 1" (purple), "Mexican Agave 2" (blue), "French Guiana" (orange), "Tequila Distillery" (yellow), and "South American Mix 2" (SAM2, pink). Small gray dots indicate isolates from other parts of the world, with the Wine and Wild strains shown as references. Black solid outlines are strains sequenced in this study. Gray circles with a solid outline indicate the few isolates from agave fermentation that did not group with other Neotropical strains. **(B)** ADMIXTURE population structure at K=3. At the bottom, strains from the Neotropical cluster, with their phylogenetic clades indicated in the second column and map at the right. **(C)** Maximum likelihood phylogenetic tree constructed using 954,449 in 464 strains including a subset of representative reference genomes and excluding the two admixed clades from the tropical Americas with poor branch support, Tequila distillery, and SAM2.

To establish the phylogenetic relationships of the newly sequenced isolates, we compared

them to a reference panel of strains (see Supplementary Table S2). Consistent with the MDS

analysis, the phylogeny showed that 205 of the 216 newly sequenced strains (94.9%) were

part of a monophyletic group that included the clades MA1, a newly defined clade Mexican

Agave 2 (MA2) and FG, plus the two admixed clades Tequila Distillery and SAM2

(Supplementary Figure S2A-B). We refer to this higher hierarchy clade as the Neotropical

cluster of *S. cerevisiae*, as nearly all the strains in this group come from the tropical Americas

(Figure 1B). Of the remaining sequenced isolates, five strains grouped with the wild North

America Oak clade; other five—including the commercial strain—were closely related to the

Mixed Origin clade and only one clustered with Wine strains.

The mosaic strains of Tequila Distillery and SAM2 branched with poor support and their

presence in the phylogeny changed the relative placement of MA1, MA2 and FG

(Supplementary Figure S2). For that reason, these admixed strains were excluded from the

main phylogenetic analysis (Figure 1C). In contrast, when SNPS from introgressed segments

were removed from the phylogeny, the clades of the Neotropical cluster remained the same.

The presence of introgressed regions had a less significant impact on the branch lengths of

the diverse Neotropical cluster compared to those of the Alpechin clade (Supplementary

Figure S2). Furthermore, the difference in the number of single nucleotide variants was more

pronounced for Alpechin than for the Neotropical cluster after the removal of introgression

(Supplementary Figure S3).

Three clades could be distinguished among the newly sequenced isolates from agave

fermentations in Neotropical *S. cerevisiae*. First, we identified a group of 20 isolates from

northeastern Mexico, which included the seven strains previously reported as "Mexican
Agave" (Peter et al., 2018) herein referred to as the MA1 clade (Figure 1C). It is worth
mentioning that the monophyletic MA1 clade comprised only strains isolated northeast of
the Sierra Madre Oriental, a mountain range which runs parallel to the coast of the Gulf of
Mexico and acts as a natural barrier, exerting strong influence on the climate and biodiversity
on either side of the range (Rzedowski, 1978; Mastretta-Yanes et al., 2015). As a result of
the comprehensive sampling of the isolates reported here, we identified a novel MA2 clade
closely related to MA1 (Figure 1C). Most of the newly sequenced strains in this study
(81.9%) belonged to this clade. Two sequenced strains isolated in Mexico (Peter et al., 2018)
were previously assigned to the SAM2 clade (Tellini et al., 2024) and after these analyses
were reclassified within the new MA2 clade. The third clade of agave isolates was composed
of 15 strains with heterogenous and admixed genotypes, all sequenced in this study (Figure
1B; Supplementary Figure S2). We refer to this as the "Tequila Distillery" clade, since eleven
were isolated from the tequila industry.

Our results show that most strains from open agave fermentations are part of the Neotropical
cluster, revealing that what was previously described as "Mexican Agave" represents only a
minor fraction of the great genetic diversity found in agave fermentations. Furthermore, the
clades within this cluster, including the newly identified groups, rank among the highest in
terms of single nucleotide variants when compared to the reference genome, surpassed only
by the wild Asian clades (**Supplementary Figure S4**). Together, our phylogenomic and
population analyses of *S. cerevisiae* from the tropical Americas bridge the gap of genetic
knowledge for the species in this megadiverse region of the world.

Neotropical yeasts belong to diverse populations stratified by geography

Yeasts from the tropical Americas come from a diverse variety of environments across a broad range of geographical distances. We thus asked to what extent geographic parameters influence the population structure of *S. cerevisiae* in the Neotropical cluster. To this end, we carried out an ADMIXTURE analysis assuming the number of populations that better explained the genotypes (Figure 2A; Supplementary Figure S5). The population structure analysis revealed eleven ancestral components in the Neotropical cluster. Specifically, strains from the MA1, Tequila Distillery, FG, and SAM2 clades each clustered in unique populations, while the remaining seven populations were all from the newly reported MA2 phylogenetic clade.

Figure 2. Yeasts from the Neotropics show population structure correlated with geography.
(A) Population structure of strains from the Neotropics and other regions of the world as reference at
K=24 made with ADMIXTURE (n=466); strains are sorted according to phylogenetic clades (first
track). To the right, strains location is shown in the maps of Mexico (top) and part of South America
encompassing Brazil, Ecuador and French Guiana. The colors of the lines connecting strains with
geographical location are the same as the most prevalent genetic component of the strain of the
Admixture plot. **(B)** Genetic diversity, π (top), heterozygosity (middle) and Tajima's D (bottom) of
clades within the Neotropical cluster.

Given the stratification observed in the Neotropical cluster, we analyzed the geographical

distribution of these isolates according to their population structure. The eleven ancestral

populations of the lineage showed good agreement with their geographic distribution (Figure

2A). This geographic stratification was observed at two levels of resolution. First, each of

the five clades of the Neotropical cluster are constrained to specific regions in the map.

Second, the seven ancestral populations revealed by ADMIXTURE within the MA2 clade

were also stratified by geography. In general, strains from the same agave-distillate

producing region grouped together in the phylogeny and had the same ancestral population

component. This was also shown by a correlation between geographic and genetic distance

in this set of strains (statistic $r=0.28$, $p<0.05$, Pearson Mantel test including MA1 and MA2

strains) and increasing genetic diversity from north to south in the MA2 clade

(Supplementary Figure S6). Together, these findings suggest that the genomic diversity in

the Neotropical cluster is influenced by geography.

Despite the overall agreement between geography and population structure in the strains from

the tropical Americas, there were exceptions suggesting gene flow within and even beyond

the lineage. For instance, a strain isolated from an open agave fermentation shared close

ancestry with wild isolates of the Wild North American clade, while other four isolates

closely related to the Wild North American clade showed signs of admixture with different

ancestral populations of the Neotropical cluster (Figure 2A). Likewise, there were SAM2

strains showing genetic components of the MA2 and French Guiana clades. Moreover, the

Tequila Distillery showed admixture, mostly between different Neotropical populations, but

also with other clades such as Wine. Together, these findings suggest that *S. cerevisiae*

populations in the tropical Americas have diverged to some extent independently from one

another by means of geographic isolation, while other ecological processes or human-related

factors have led to gene flow, which also contributes to their genome diversity.

We used the genomic data to further explore the patterns of genetic diversity in the

Neotropical strains. The strains from each clade varied significantly from each other in

nucleotide diversity (π), Tajima's D, and heterozygosity (Figure 2B; Supplementary Figure

S7), with some of them displaying remarkably high genetic-diversity. For instance, the MA1,

MA2, and Tequila Distillery clades all show higher genetic diversity than yeasts from the

Wine clade (Supplementary Table 2). This held true when comparing with the highly

introgressed Alpechin strains, suggesting that differences in genetic diversity cannot be

solely accounted for by their introgressions. In contrast, the FG clade showed a homogenous

degree of heterozygosity and low genetic diversity. The Tequila clade had the highest

heterozygosity and nucleotide diversity, in agreement with trends observed in domesticated

beer isolates (Peter et al., 2018), while SAM2 also showed high genetic diversity, but lower

heterozygosity, as reported for wild yeast strains (Han et al., 2021). High diversity in these

clades is in part due to their admixed genotypes (Figure 2A), while high heterozygosity

specifically in the Tequila Distillery strains may be associated to their adaptation to the semi-

industrial context where they were isolated. Taken together, our population-structure and

genetic-diversity analyses indicate that yeasts in the tropical Americas constitute highly

diverse, structured populations stratified by geography.

Different distributions of introgressions show recurrent interspecies gene flow

The genomes of *S. cerevisiae* strains within the Neotropical cluster are characterized by an

atypically high number of introgressed regions from *S. paradoxus* (Barbosa et al., 2016;

Tellini et al., 2024). To gain insight into gene flow dynamics, we identified the set of

introgressed genes (Materials and methods), their location, and length patterns in each clade

of the Neotropical cluster (Supplementary Figure S8A). We found that yeasts from MA1

typically exhibited not only more introgressed genes than all other clades, but also the highest

variability in this number, ranging from 119 to 364 introgressed genes per strain. Collectively

the introgressions in MA1 covered 7.1% of the *S. cerevisiae* reference genome (Figure 3A).

We note that the number of introgressed genes in most MA1 strains was within the range

observed for the MA2 strains which featured 96 to 160 introgressed genes. Together, all the

introgressed genes in the MA2 clade covered 3.2% of the nuclear genome. However, seven

strains from the MA1 clade exhibited longer and more heterozygous introgressions than most

MA1 strains (Figure 3A; Supplementary Figure S9). Here, we call an introgression as

heterozygous when both *S. cerevisiae* and *S. paradoxus* alleles have been retained in a single

genome, suggesting a more recent introgression event. The presence of both alleles in several

introgressions of the Tequila Distillery clade suggests admixture with a non-introgressed

population, which may also explain the lower number of introgressed genes (Figure 3A).

Figure 3. Different patterns of introgressions in the clades from the Neotropics. A) Distributions of the number of introgressed genes per strain in each phylogenetic clade. The Alpechin strains are included as a reference of a highly introgressed population. The color of each dot indicates the fraction of introgressed genes that are heterozygous in each strain, meaning that both the *S. cerevisiae* and *S. paradoxus* alleles are present. **B)** Clustering of the introgressed genes found across the Neotropical cluster (n = 1,113). Colors indicate the fraction of strains from each clade that had the introgressed gene. **C)** Distribution of the frequency of introgressions in each clade. **D)** Fraction of the maximum number of possible shared introgressed genes in each comparison. **E)** "Private" introgressed genes, meaning that they are present only in one of the clades.

All Neotropical clades displayed more introgressed genes than the observed average in other groups from around the world, except for the Alpechin clade. The two clades from South

America, SAM2 and FG, displayed fewer and shorter introgressions with a lower degree of

heterozygosity, compared to the two Mexican agave clades (Figure 3A; Supplementary

Figure S9). This trend suggests that the last introgression pulse of SAM2 and FG likely

occurred in the common ancestor of all strains from the Neotropical cluster. These

characteristics in the introgression patterns agree with the trends that we observed in terms

of the introgressions that are shared, whereby a significant fraction of genes was common

between the Mexican Agave clades and, to a lesser extent, with the French Guiana clade

(Figure 3B). Importantly, these patterns of shared introgressions were confirmed using an

alternative strategy that considered *bona fide* species-specific SNP markers (Tellini et al.,

2024) for detecting and scoring introgressed regions (Materials and Methods; Supplementary

Figure S8).

To further understand how introgressions are shared among clades, we quantified the

proportion of strains harboring each introgressed gene within the different clades (Figure

3C). Introgressed genes present at high frequency—at least in 90% of the strains within a/the

clade—were common in French Guiana (43.6%) and Alpechin (22.5%). The high proportion

of shared introgressed genes within these populations suggests either a strong bottleneck or

positive selection acting on introgressed regions. Conversely, introgressed genes present at

10% or lower frequency were more common within the Mexican Agave clades (36% and

45.7%; Figure 3C), which suggests either divergent resolutions of the same hybridization

events, recent independent hybridizations in specific strains or a combination of both.

We found that MA1 and MA2 shared the highest fraction of introgressed genes in paired

comparisons among clades, suggesting that they were present in their common ancestor

(Figure 3D). Moreover, the highest fraction in the triple intersection was that of the MA1,

MA2 and FG (Figure 3C). These results suggest that a fraction of the introgressions present

in the three clades was present in their common ancestor. Additionally, we observed that 14

genes were shared among all four clades, pointing to potential strong selective pressures or

artifacts arising from the use of a reference genome lacking those specific genes. In terms of

"private" introgressions—those present in a clade but absent in all others—we scored 18.9%

in FG, 32.2% in MA2, and 58.6% in MA1 (Figure 3E), while most introgressed genes in the

Alpechin reference clade (76.3%) were exclusively found in this clade (Figure 3E). The high

percentage of private introgressed genes in Alpechin is due to hybridization events not shared

with Neotropical strains (Peter et al., 2018; Tellini et al., 2024). Likewise, MA1 and MA2

isolates exhibited a high fraction of private introgressions, potentially resulting from

introgression pulses that are not shared with the related FG clade. Together, the patterns of

disparate introgression distribution suggest recurrent interspecies gene flow episodes

between *S. cerevisiae* and *S. paradoxus* from the tropical Americas.

Introgressions in the Neotropical cluster trace to different lineages of *S. paradoxus*

To track the history of the observed interspecies gene-flow episodes, we determined the most

likely *S. paradoxus* lineage from which each introgressed gene derived. To this end, we

scored sequence similarity of each introgressed block against a panel of *S. paradoxus*

genomes including representative strains from the Americas (Figure 4A). Our results suggest

that most introgressed genes in the Mexican Agave clades derived from a lineage of *S.*

paradoxus that has only been isolated in Mexico and that cohabits in agave fermentations,

SpB_Mx (Lopez-Gallegos, 2023). In contrast, the majority of the introgressed genes in the

FG strains tracked to *SpB,* a different lineage of *S. paradoxus,* which is not commonly found

in agave distillery contexts (Lopez-Gallegos, 2023). As expected, the introgressed genes in

the Alpechin group most closely resemble the European *SpA* lineage, previously identified

as the parental population of these introgressions (D'Angiolo et al., 2020; Tellini et al., 2024).

These observations suggest that at least two lineages of *S. paradoxus* have hybridized with

ancestors of the extant Neotropical strains, underscoring the close relationship between these

two species in the evolution of *S. cerevisiae* in the region.

A)

B)

Figure 4. Multiple introgression pulses from two *S. paradoxus* lineages in the Neotropical *S.*
***cerevisiae* cluster.** (**A**) The average number of introgressed genes from each lineage of *S.*
paradoxus in each clade of Neotropical *S. cerevisiae*. Error bars are the standard deviation. The
Alpechin clade is shown as a reference. (**B**) Scheme of the model showing multiple introgression
pulses as the most plausible evolutionary scenario for the observed patterns of number and origins
of introgressed genes along with the levels of heterozygosity and length of the introgressed blocks.
335

DISCUSSION

Comprehensive sequencing of *S. cerevisiae* genomes in megadiverse regions, harboring a

substantial portion of Earth's biodiversity, is essential to understand the evolutionary history

and genomic diversity of this prominent model organism. In this study, we sequenced 216

strains from open agave fermentations, providing an extensive population genomics dataset

for the Neotropics. Our phylogenetic and population structure analyses revealed multiple

interspecies gene-flow events resulting in introgressions within this geographically

structured complex clade.

Introgressions impact evolutionary trajectories across the tree of life, from bacteria to humans

(Diop et al., 2022; Edelman and Mallet, 2021). These events are also central to the

evolutionary history of *S. cerevisiae*, as evidenced by their presence in isolates across the

globe (Clark et al., 2022; Peter et al., 2018; Strope et al., 2015; Tellini et al., 2024). In the

Americas, several *S. cerevisiae* lineages feature high number of introgressions from *S.*

paradoxus, both in wild and anthropogenic environments (Barbosa et al., 2016; Pontes et al.,

2020; Jacobus et al., 2021; Tellini et al., 2024), presumably as a consequence of the co-

existence of the two species in the region. Here, we provide a broad picture of the

evolutionary dynamics of introgression happening in the tropical Americas. In the most

plausible scenario, an early introgression pulse occurred in the most recent common ancestor

of the Neotropical cluster before its diversification, where genes of the American *SpB* lineage

of *S. paradoxus* introgressed into *S. cerevisiae* (Figure 4B). This first pulse was presumably

followed by a subsequent introgression event from another lineage, *SpB_Mx*, in the common

ancestor of MA1 and MA2. At least one additional, more recent pulse of introgression from

the *SpB_Mx* lineage took place in the MA1 clade, which is supported by high numbers of

large and heterozygous introgressions in some strains of this clade. It remains to be

determined whether these introgression pulses are simply stochastic byproducts of the

coexistence of *S. cerevisiae* and *S. paradoxus* in the region, or if they underlie adaptation

processes. Wider sampling efforts in other areas, particularly in South America, will likely

expand our knowledge of the interspecies gene-flow dynamics in these yeasts.

Our sequencing effort expanded knowledge of the *S. cerevisiae* diversity and biogeography.

We uncovered two clades within the Neotropical cluster: MA2 and the admixed Tequila

Distillery. Together with MA1, FG, and SAM2, these five clades constitute the Neotropical

cluster. Phylogenetic analysis indicated that the MA1—previously designated as Mexican

Agave—and MA2 are sister clades, which in turn are the closest relatives of the FG strains,

in agreement with the previously reported phylogeny (O'Donnell et al., 2022). The high

variability observed in the remaining SAM2 strains indicates that there probably exist other

undersampled lineages in the tropical Americas, a diverse and biogeographically complex

region (Morrone, 2014). Additional sampling and sequencing in this region would further

elucidate key evolutionary events that have shaped the natural history of yeasts, as discussed

herein.

Importantly, we found that the novel genetic diversity in the Neotropical cluster is structured

and associated with the geographic origins of the yeast isolates. Geographical influence that

shaped the genetic diversity was evident across several dimensions: First, a clear correlation

was observed between each of the eleven ancestral components of the strains from the

Neotropics and their region of isolation. Second, strains from the MA2 clade exhibit

increased nucleotide diversity, in a north-to-south gradient. Finally, the genetic divergence

between the MA1 and MA2 clades is associated with the presence of the Sierra Madre

Oriental mountain range, which is found at the edge where the Neotropical and Nearctic

biogeographic regions coincide. This range is known to act as a geographical barrier

impacting the distributions of multiple species (Mastretta-Yanes et al., 2015; Morrone, 2014;

Rzedowski, 1978).

Beyond biogeographical factors, human activities may have played a role in shaping these

yeast populations, as evidenced by admixture events within and beyond the Neotropical

cluster. It remains to be addressed to what extent the observed yeast population structure is

driven by natural factors such as the ecology and biogeography of plant reservoirs and insect

vectors or by the human dynamics and practices. For instance, the specific local production

practices or the remarkable diversity of fermentation substrates, with over 50 species of

Agave species used for producing spirits (Colunga García Marín et al., 2017), could be

driving population differentiation. The agave fermentation system is not only a reservoir of

remarkable yeast genetic diversity, but also a unique natural laboratory to gain insights into

the interplay of genetic, ecological and evolutionary factors shaping population

differentiation, and interspecies introgression.

MATERIALS AND METHODS

Genome sequencing and variant calling. DNA was purified with the MasterPure DNA purification kit as recommended by the manufacturer, and it was sequenced using DNBSeq (2x150 bp; BGI, China). Raw reads were quality assessed with fastp V0.20.0 (Chen et al., 2018). For each isolate, the filtered reads were aligned with BWA-MEM v0.7.4 (Li, 2013) to a concatenated species reference genome that includes *S. cerevisiae* S288C, *S. paradoxus* YPS138, *S. mikatae* IFO1815, *S. kudriavzevii* Cr85, *S. jurei* M1, *S. arboricola* H6, *S. uvarum* CBS7001, *S. eubayanus* FM1318, *K. marxianus* DMKU31042, and *P. kudriavzevii* CBS573. Duplicated reads were marked with Piccard 2.6.0 (Picard, n.d.), and local realignment around indels and variant calling were performed with GATK v4.1.1.0 (McKenna et al., 2010). Allele balance information was incorporated in the vcf files with GATK Variant annotator. The called genotypes were filtered to keep only biallelic SNPs with a minimum depth of 5, a QUAL score of at least 30 and SNPs presents in at least 90% of the isolates. Additionally, SNPs were also obtained by mapping the filtered reads only to the *S. cerevisiae* genome to obtain variants that include introgressions (Supplementary Figure S3, Supplementary Figure S4).

Multidimension scaling analysis. To analyze population structure, the generated genome-wide SNPs data were used for multidimension scaling (MDS) analysis with plink v1.9 (Purcell et al., 2007). The data set consisted of 1,264 isolates, including 1,006 from Peter et al. (2018) and 258 from the Americas including the 216 sequenced by us. Only nuclear genome SNPs were considered, and low-frequency or rare variants were excluded to this analysis using a threshold of missing call rate > 1 %. After the application of this filter,

isolates with missing genotype > 10% were discarded. The quantitative indices (components)

of the genetic variation for each isolate were calculated on the genome-wide average

proportion of alleles shared between any pair of individuals within the sample.

Phylogenetic analysis. Maximum likelihood trees were built with genomes sequenced here

and references from all over the world with emphasis in American clades. Haplotype

sequences were obtained with vcf2phylip v2.3 (Ortiz, n.d.) from the nuclear genome

variants derived from the concatenated species genome that mapped to the *S. cerevisiae* sub

genome (or directly to *S. cerevisiae* reference to include variation from introgressions for

Supplementary Figure 2). The phylogeny trees were inferred by raxmlHPC-PTHREADS-

AVX2 (Stamatakis, 2014) with the GTR+GAMMA parameter and 100 bootstrap replicates.

The visualization of the tree was performed using Microreact (Argimón et al., 2016). The

first two phylogenetic trees were built for 487 isolates including the 216 sequenced here

(Supplementary Figure S2). As admixed Tequila Distillery and SAM2 strains changed the

topology within the Neotropical cluster, we removed admixed strains from the phylogeny

presented in Figure 1 with 464 genomes (Supplementary Table S2).

ADMIXTURE. To run ADMIXTURE 1.3.0 (Alexander et al., 2009) the VCF with 466

genomes was curated using Plink 1.9 (Purcell and Chang, n.d.) to remove linkage

disequilibrium with parameters 50 window size sliding every 5 nucleotides and correlation

of 0.5 as previously run in *S. cerevisiae* population genetics (Gallone et al., 2016) which led

to 618,253 SNPs. Ten seeds were used to run Admixture. The Cross Validation (CV) error

was obtained for models with k=2 to K=35 populations. The lowest median CV error K was

24 and therefore the iteration with K=24 and the lowest CV error was used for visualization.

Visualization was made in Pong (Behr et al., 2016).

Estimation of genetic diversity. To calculate genetic diversity (π) and Tajima's D, we used

vcftools 0.1.14 (Danecek et al., 2011) and the VCF with variants that mapped to *S. cerevisiae*

after mapping to the concatenated reference and filtering high quality biallelic SNPs with

less than 10% of missing data. To estimate the heterozygosity, bcftools 1.9 (Danecek et al.,

2021) was used as the number of heterozygous variants divided by the total number of

variants in the individual strain VCFs.

Analysis of Introgressions. To identify introgressed genes, we compared the *S. paradoxus*

section of the alignment to the concatenated reference. A gene was considered introgressed

if over half of it had a coverage depth exceeding 25% of the median depth of the entire

reference. Additionally, it had to be the ortholog with the highest depth or, if not, its depth

had to be at least 25% of its ortholog in the *S. cerevisiae* sub genome. We also required a

minimum 5% nucleotide-level difference between the orthologs of *S. cerevisiae* and *S.*

paradoxus. An introgressed gene was considered heterozygous if the ratio of the depth

coverage between the two sub genomes in the competitive mapping was between 0.25 and 4.

Nevertheless, we alternatively used a second strategy with the same VCF file to get the

positions of a set of *bonna fide* introgression markers and retained introgression blocks of at

least five consecutive markers (Tellini et al., 2024) to contrast with our mapping-based

definition (Supplementary Figure S8). Visualization was made using ChromoMap (Anand

and Rodriguez Lopez, 2022).

Once the introgressed genes were determined, consecutive introgressed genes according to

the annotation (Yue et al., 2017) were considered part of a block. To determine the origin of

the blocks, we aligned the *S. paradoxus* from Lopez-Gallegos, 2023 to the concatenated multi

species reference described above and kept only the regions annotated as introgressions in

any of the *S. cerevisiae* strains from this work. We computed the paired distances between

all introgressed blocks and the *S. paradoxus* genomes using bcftools 1.9 (Danecek et al.,

2021) and searched for the *S. paradoxus* with the lowest distance to each introgressed block.

If a given introgression was most similar to strains from more than one *S. paradoxus* lineage,

no origin was assigned to it. All the genes within a given block were considered to be from

the same origin.

DATA ACCESSIBILITY

485 **Genome sequencing.** Genome sequencing data were deposited in the SRA under the

486 project PRJNA1138754.

487 **Custom scripts.** Scripts to identify introgressed genes and their origins along with scripts

488 to run the figures of the article are available at: https://github.com/GELab-

489 LIIGH/ScerYeastGenomesMx2024

ACKNOWLEDGEMENTS

This work was funded by Consejo Nacional de Humanidades, Ciencias y Tecnologías de

México (CONAHCYT grants FORDECYT-PRONACES/103000/2020, CB-2016-

01/284992, CF-2023-G-695), Fondo de Investigación y Desarrollo Tecnológico del

Cinvestav (SEP-CINVESTAV/023), Programa de Apoyo a Proyectos de Investigación e

Innovación Tecnológica DGAPA-UNAM (grants IN209021, IN230420 and IN212524), and

the UK BBSRC under the Global Challenges Research Fund (GCRF) Growing Research

Capability call through the CABANA Innovation Fund (BB/P027849/1). LFG-O is a

postdoctoral researcher funded by Conahcyt (4133922); IS is a PhD student at UNAM with

a graduate scholarship from Conahcyt. We thank Bernard Dujon, Xitlali Aguirre-Dugua and

Maria Avila for critical reading of the manuscript. We thank Luis Aguilar (LAVIS, UNAM),

Porfirio Gallegos (Cinvestav), Claudio Lopez (Cinvestav), Aaron de Luna (LIIGH),

Alejandra Castillo (LIIGH), Carina Uribe (LIIGH), Maritrini Colón-González (LIIGH) and

Jair García (LIIGH) for technical assistance. We thank Xitlali Aguirre-Dugua, Luis Delaye,

Diego Ortega, Marcela Sandoval and Alicia Mastretta-Yanes for helpful discussions and

Manuel Kirchmayr (CIATEJ), Anne Gschaedler (CIATEJ), Maritza Álvarez (CIAD) and

Andre Marc Lachance for sharing their strains.

MANUSCRIPT ITEMS

Figures

Figure 1. A novel, diverse yeast lineage encompasses most of the diversity in open agave fermentations.

Figure 2. Yeasts from the Neotropical cluster show population structure correlated with geography.

Figure 3. Different patterns of introgressions in the clades from the tropical Americas.

Figure 4. Multiple introgression pulses from two *S. paradoxus* lineages in the Neotropical *S. cerevisiae* clades.

Supplementary Tables

Table S1. The 216 Strains sequenced in this study and associated metadata.

Table S2. Strains and metadata used in this study.

Supplementary Figures

Supplementary Figure S1. Origin of the sequenced strains.

Supplementary Figure S2. Phylogenies of the Neotropical cluster show the same groups with or without introgressions, but their relative placement is affected by Tequila Distillery strains.

Supplementary Figure S3. Number of single nucleotide variants of strains with or without introgressions.

Supplementary Figure S4. Neotropical strains grouped in clades show a high number of single nucleotide variants compared to the worldwide diversity.

Supplementary Figure S5. Population structure analysis.

Supplementary Figure S6. Biocultural regions of agave fermentation for spirit production show a gradient in genetic diversity.

Supplementary Figure S7. Genetic diversity parameters in clades of yeasts from the Neotropics.

Supplementary Figure S8. Genomic map of introgressions shows similar distributions regardless of the specific method to define them.

540 **Supplementary Figure S9.** Introgression block size per clade of the Tropical Americas

541 linage Neotropical cluster.

REFERENCES

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast model-based estimation of ancestry in unrelated individuals. Genome Res. 19, 1655–1664. https://doi.org/10.1101/gr.094052.109

Alsammar, H., Delneri, D., 2020. An update on the diversity, ecology and biogeography of the Saccharomyces genus. FEMS Yeast Res. 20, foaa013. https://doi.org/10.1093/femsyr/foaa013

Álvarez-Ainza, M.L., Zamora-Quiñonez, K.A., Moreno-Ibarra, G.M., Acedo-Félix, E., 2015. Genomic Diversity of Saccharomyces cerevisiae Yeasts Associated with Alcoholic Fermentation of Bacanora Produced by Artisanal Methods. Appl. Biochem. Biotechnol. 175, 2668–2676. https://doi.org/10.1007/s12010-014-1469-y

Anand, L., Rodriguez Lopez, C.M., 2022. ChromoMap: an R package for interactive visualization of multi-omics data and annotation of chromosomes. BMC Bioinformatics 23, 33. https://doi.org/10.1186/s12859-021-04556-z

Arellano-Plaza, M., Paez-Lerma, J.B., Soto-Cruz, N.O., Kirchmayr, M.R., Gschaedler Mathis, A., 2022. Mezcal Production in Mexico: Between Tradition and Commercial Exploitation. Front. Sustain. Food Syst. 6.

Argimón, S., Abudahab, K., Goater, R.J.E., Fedosejev, A., Bhai, J., Glasner, C., Feil, E.J., Holden, M.T.G., Yeats, C.A., Grundmann, H., Spratt, B.G., Aanensen, D.M., 2016. Microreact: visualizing and sharing data for genomic epidemiology and phylogeography. Microb. Genomics 2, e000093. https://doi.org/10.1099/mgen.0.000093

Barbosa, R., Almeida, P., Safar, S.V.B., Santos, R.O., Morais, P.B., Nielly-Thibault, L., Leducq, J.-B., Landry, C.R., Gonçalves, P., Rosa, C.A., Sampaio, J.P., 2016. Evidence of Natural Hybridization in Brazilian Wild Lineages of Saccharomyces cerevisiae. Genome Biol. Evol. 8, 317–329. https://doi.org/10.1093/gbe/evv263

Behr, A.A., Liu, K.Z., Liu-Fang, G., Nakka, P., Ramachandran, S., 2016. pong: fast analysis and visualization of latent clusters in population genetic data. Bioinformatics 32, 2817–2823. https://doi.org/10.1093/bioinformatics/btw327

Bendixsen, D.P., Gettle, N., Gilchrist, C., Zhang, Z., Stelkens, R., 2021. Genomic Evidence of an Ancient East Asian Divergence Event in Wild Saccharomyces cerevisiae. Genome Biol. Evol. 13, evab001. https://doi.org/10.1093/gbe/evab001

Boynton, P.J., Greig, D., 2014. The ecology and evolution of non-domesticated Saccharomyces species. Yeast Chichester Engl. 31, 449–462. https://doi.org/10.1002/yea.3040

Bruman, H.J., 2000. Alcohol in Ancient Mexico. University of Utah Press.

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Clark, A., Dunham, M.J., Akey, J.M., 2022. The genomic landscape of Saccharomyces paradoxus introgression in geographically diverse Saccharomyces cerevisiae strains. https://doi.org/10.1101/2022.08.01.502362

Colón-González, M., Aguirre-Dugua, X., Guerrero-Osornio, M.G., Avelar-Rivas, J.A., DeLuna, A., Mancera, E., Morales, L., 2024. Thriving Amidst Sweetness and Harshness: The Life of Yeasts in the Agave Fermentation Environment.

Colunga García Marín, S.P., Torres García, I., Casas Fernández, A., Figueredo Urbina, C.J., Rangel Landa, S., Delgado Lemus, A., Vargas Ponce, O., Cabrera Toledo, D., Zizumbo Villareal, D., Aguirre-Dugua, X., Eguiarte Fruns, L.E., Carrillo Galván, G., 2017. Los agaves y las prácticas mesoamericanas de aprovechamiento, manejo y domesticación.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., 1000 Genomes Project Analysis Group, 2011. The variant call format and VCFtools. Bioinformatics 27, 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and BCFtools. GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008

D'Angiolo, M., De Chiara, M., Yue, J.-X., Irizar, A., Stenberg, S., Persson, K., Llored, A., Barré, B.,
Schacherer, J., Marangoni, R., Gilson, E., Warringer, J., Liti, G., 2020. A yeast living ancestor
reveals the origin of genomic introgressions. Nature 587, 420–425.
https://doi.org/10.1038/s41586-020-2889-1
De Chiara, M., Barré, B.P., Persson, K., Irizar, A., Vischioni, C., Khaiwal, S., Stenberg, S., Amadi,
O.C., Žun, G., Doberšek, K., Taccioli, C., Schacherer, J., Petrovič, U., Warringer, J., Liti, G.,
2022. Domestication reprogrammed the budding yeast life cycle. Nat. Ecol. Evol. 6, 448–460.
https://doi.org/10.1038/s41559-022-01671-9
Diop, A., Torrance, E.L., Stott, C.M., Bobay, L.-M., 2022. Gene flow and introgression are pervasive
forces shaping the evolution of bacterial species. Genome Biol. 23, 239.
https://doi.org/10.1186/s13059-022-02809-5
Duan, S.-F., Han, P.-J., Wang, Q.-M., Liu, W.-Q., Shi, J.-Y., Li, K., Zhang, X.-L., Bai, F.-Y., 2018. The
origin and adaptive evolution of domesticated populations of yeast from Far East Asia. Nat.
Commun. 9, 2690. https://doi.org/10.1038/s41467-018-05106-7
Edelman, N.B., Mallet, J., 2021. Prevalence and Adaptive Impact of Introgression. Annu. Rev. Genet.
55, 265–283. https://doi.org/10.1146/annurev-genet-021821-020805
Gallegos-Casillas, P., García-Ortega, L.F., Espinosa-Cantú, A., Avelar-Rivas, J.A., Torres-Lagunes,
C.G., Cano-Ricardez, A., García-Acero, Á.M., Ruiz-Castro, S., Flores-Barraza, M., Castillo, A.,
González-Zozaya, F., Delgado-Lemus, A., Molina-Freaner, F., Jacques-Hernández, C.,
Hernández-López, A., Delaye, L., Aguirre-Dugua, X., Kirchmayr, M.R., Morales, L., Mancera,
E., DeLuna, A., 2024. Yeast diversity in open agave fermentations across Mexico. Yeast 41,
35–51. https://doi.org/10.1002/yea.3913
Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., Merlevede, A.,
Roncoroni, M., Voordeckers, K., Miraglia, L., Teiling, C., Steffy, B., Taylor, M., Schwartz, A.,
Richardson, T., White, C., Baele, G., Maere, S., Verstrepen, K.J., 2016. Domestication and
Divergence of Saccharomyces cerevisiae Beer Yeasts. Cell 166, 1397-1410.e16.
https://doi.org/10.1016/j.cell.2016.08.020
Han, D.-Y., Han, P.-J., Rumbold, K., Koricha, A.D., Duan, S.-F., Song, L., Shi, J.-Y., Li, K., Wang, Q.-
M., Bai, F.-Y., 2021. Adaptive Gene Content and Allele Distribution Variations in the Wild and
Domesticated Populations of Saccharomyces cerevisiae. Front. Microbiol. 12.
https://doi.org/10.3389/fmicb.2021.631250
Jacobus, A.P., Stephens, T.G., Youssef, P., González-Pech, R., Ciccotosto-Camp, M.M., Dougan, K.E.,
Chen, Y., Basso, L.C., Frazzon, J., Chan, C.X., Gross, J., 2021. Comparative Genomics
Supports That Brazilian Bioethanol Saccharomyces cerevisiae Comprise a Unified Group of
Domesticated Strains Related to Cachaça Spirit Yeasts. Front. Microbiol. 12.
https://doi.org/10.3389/fmicb.2021.644089
Kirchmayr, M.R., Arellano-Plaza, M., Estarron Espinosa, M., Gallardo Valdez, J., Gschaedler Mathis,
A., Lopez Ramirez, J.E., Navarro Hurtado, A., Prado Ramirez, R., Ramirez Romo, E., 2014.
Manual para la estandarización de los procesos de producción del mezcal guerrerense, 1st ed.
Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco A.C.,
Guadalajara, Jalisco. México.
Kirchmayr, M.R., Segura-García, L.E., Lappe-Oliveras, P., Moreno-Terrazas, R., de la Rosa, M.,
Gschaedler Mathis, A., 2017. Impact of environmental conditions and process modifications on
microbial diversity, fermentation efficiency and chemical profile during the fermentation of
Mezcal in Oaxaca. LWT - Food Sci. Technol. 79, 160–169.
https://doi.org/10.1016/j.lwt.2016.12.052
Lachance, M.-A., 1995. Yeast communities in a natural tequila fermentation. Antonie Van Leeuwenhoek
68, 151–160. https://doi.org/10.1007/BF00873100
Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
https://doi.org/10.48550/arXiv.1303.3997
Liti, G., 2015. The Natural History of Model Organisms: The fascinating and secret wild life of the
budding yeast S. cerevisiae [WWW Document]. eLife. https://doi.org/10.7554/eLife.05835

Loegler, V., Friedrich, A., Schacherer, J., 2024. Overview of the Saccharomyces cerevisiae population structure through the lens of 3,034 genomes. https://doi.org/10.1101/2024.09.16.613241

Lopez-Gallegos, C.E., 2023. Population genomics and ecology of yeast associated with traditional Agave fermentation in Mexico (MsC.). CINVESTAV del IPN, https://liigh.unam.mx/yeastgenomesmx/wp-content/uploads/2024/09/TesisDeMaestria_ClaudioLopez_Marzo2023-1.pdf , Irapuato, Guanajuato, Mexico.

Marcet-Houben, M., Gabaldón, T., 2015. Beyond the Whole-Genome Duplication: Phylogenetic Evidence for an Ancient Interspecies Hybridization in the Baker's Yeast Lineage. PLOS Biol. 13, e1002220. https://doi.org/10.1371/journal.pbio.1002220

Mastretta-Yanes, A., Moreno-Letelier, A., Piñero, D., Jorgensen, T.H., Emerson, B.C., 2015. Biodiversity in the Mexican highlands and the interaction of geology, geography and climate within the Trans-Mexican Volcanic Belt. https://doi.org/10.1111/jbi.12546

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., DePristo, M.A., 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res. 20, 1297–1303. https://doi.org/10.1101/gr.107524.110

Mittermeier, R.A., Mittermeier, C.G., 2005. Megadiversity: Earths Biologically Wealthiest Nations. CEMEX.

Morrone, J.J., 2014. Biogeographical regionalisation of the Neotropical region. Zootaxa 3782, 1–110. https://doi.org/10.11646/zootaxa.3782.1.1

O'Donnell, S., Yue, J.-X., Saada, O.A., Agier, N., Caradec, C., Cokelaer, T., Chiara, M.D., Delmas, S., Dutreux, F., Fournier, T., Friedrich, A., Kornobis, E., Li, J., Miao, Z., Tattini, L., Schacherer, J., Liti, G., Fischer, G., 2022. 142 telomere-to-telomere assemblies reveal the genome structural landscape in Saccharomyces cerevisiae. https://doi.org/10.1101/2022.10.04.510633

Ortiz, E.M., n.d. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for phylogenetic analysis. https://doi.org/DOI:10.5281/zenodo.2540861

Padilla-Camberos, E., Pinal-Zuazo, L., Alvarez de la Cuadra Jacob, J., 1994. Catálogo de la colección de cultivos microbianos.

Peris, D., Ubbelohde, E.J., Kuang, M.C., Kominek, J., Langdon, Q.K., Adams, M., Koshalek, J.A., Hulfachor, A.B., Opulente, D.A., Hall, D.J., Hyma, K., Fay, J.C., Leducq, J.-B., Charron, G., Landry, C.R., Libkind, D., Gonçalves, C., Gonçalves, P., Sampaio, J.P., Wang, Q.-M., Bai, F.-Y., Wrobel, R.L., Hittinger, C.T., 2023. Macroevolutionary diversity of traits and genomes in the model yeast genus Saccharomyces. Nat. Commun. 14, 690. https://doi.org/10.1038/s41467-023-36139-2

Peter, J., De Chiara, M., Friedrich, A., Yue, J.-X., Pflieger, D., Bergström, A., Sigwalt, A., Barre, B., Freel, K., Llored, A., Cruaud, C., Labadie, K., Aury, J.-M., Istace, B., Lebrigand, K., Barbry, P., Engelen, S., Lemainque, A., Wincker, P., Liti, G., Schacherer, J., 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates. Nature 556, 339–344. https://doi.org/10.1038/s41586-018-0030-5

Picard, toolkit, n.d. Picard toolkit [WWW Document]. Picard Tools. URL https://broadinstitute.github.io/picard/

Pontes, A., Hutzler, M., Brito, P.H., Sampaio, J.P., 2020. Revisiting the Taxonomic Synonyms and Populations of Saccharomyces cerevisiae—Phylogeny, Phenotypes, Ecology and Domestication. Microorganisms 8, 903. https://doi.org/10.3390/microorganisms8060903

Purcell, S., Chang, C., n.d. PLINK 1.9 [WWW Document]. URL www.cog-genomics.org/plink/1.9/

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P., Bakker, P.I.W. de, Daly, M.J., Sham, P.C., 2007. PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. Am. J. Hum. Genet. 81, 559–575. https://doi.org/10.1086/519795

Quezada, R., Gschaedler, A., Kirchmayr, M., 2016. Characterization of microbial population dynamics associated with different juices of Agave tequilana, in: Sustainable and Integrated Use of Agave, III International Symposium on Agave. 2016, Zapopan, Jalisco, Mexico, p. 2016.

Rzedowski, J., 1978. Vegetación de México, Editorial Limusa S. A. ed.

Shen, X.-X., Opulente, D.A., Kominek, J., Zhou, X., Steenwyk, J.L., Buh, K.V., Haase, M.A.B.,
Wisecaver, J.H., Wang, M., Doering, D.T., Boudouris, J.T., Schneider, R.M., Langdon, Q.K.,
Ohkuma, M., Endoh, R., Takashima, M., Manabe, R., Čadež, N., Libkind, D., Rosa, C.A.,
DeVirgilio, J., Hulfachor, A.B., Groenewald, M., Kurtzman, C.P., Hittinger, C.T., Rokas, A.,
2018. Tempo and Mode of Genome Evolution in the Budding Yeast Subphylum. Cell 175,
1533-1545.e20. https://doi.org/10.1016/j.cell.2018.10.023

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30, 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Strope, P.K., Skelly, D.A., Kozmin, S.G., Mahadevan, G., Stone, E.A., Magwene, P.M., Dietrich, F.S.,
McCusker, J.H., 2015. The 100-genomes strains, an S. cerevisiae resource that illuminates its
natural phenotypic and genotypic variation and emergence as an opportunistic pathogen.
Genome Res. 25, 762–774. https://doi.org/10.1101/gr.185538.114

Tellini, N., De Chiara, M., Mozzachiodi, S., Tattini, L., Vischioni, C., Naumova, E.S., Warringer, J.,
Bergström, A., Liti, G., 2024. Ancient and recent origins of shared polymorphisms in yeast. Nat.
Ecol. Evol. 8, 761–776. https://doi.org/10.1038/s41559-024-02352-5

Verdugo Valdez, A., Segura Garcia, L., Kirchmayr, M., Ramírez Rodríguez, P., González Esquinca, A.,
Coria, R., Gschaedler Mathis, A., 2011. Yeast communities associated with artisanal mezcal
fermentations from Agave salmiana. Antonie Van Leeuwenhoek 100, 497–506.
https://doi.org/10.1007/s10482-011-9605-y

Wang, M., Li, X., Liu, X., Hou, X., He, Y., Yu, J.-H., Hu, S., Yin, H., Xie, B.-B., 2024. Annotation of
2,507 Saccharomyces cerevisiae genomes. Microbiol. Spectr. 12, e03582-23.
https://doi.org/10.1128/spectrum.03582-23

Yue, J.-X., Li, J., Aigrain, L., Hallin, J., Persson, K., Oliver, K., Bergström, A., Coupland, P., Warringer,
J., Lagomarsino, M.C., Fischer, G., Durbin, R., Liti, G., 2017. Contrasting evolutionary genome
dynamics between domesticated and wild yeasts. Nat. Genet. 49, 913–924.
https://doi.org/10.1038/ng.3847

SUPPLEMENTARY FIGURES CAPTIONS

Supplementary Figure S1. Origin of the 216 sequenced strains.

Supplementary Figure S2. Phylogenies of the Neotropical cluster show the same groups with or without introgressions, but their relative placement is affected by mosaic Tequila Distillery and SAM2 strains. Maximum likelihood phylogenetic trees constructed including a subset of representative reference genomes. (A) Maximum likelihood phylogeny with a complete set of 487 strains and 1,002,285 SNPs excluding introgressions. (B) Maximum likelihood phylogeny of the same set of genomes, including SNPs within introgressions (1,177,709 SNPs). (C) Maximum likelihood of 464 strains as described in the main text. The bootstrap values of relevant branching points show how the Tequila Distillery mosaics hamper reconstruction of the branching order of the major phylogenetic clades. Colored dots at the tip of the branches show the origins of *S. cerevisiae* strains from the Neotropical cluster: Mexican Agave 1 (purple), Mexican Agave 2 (blue), French Guiana (orange), Tequila Distillery (yellow), South American Mix 2 (pink); Alpechin strains are also highlighted (salmon).

Supplementary Figure S3. Number of single nucleotide variants of strains with or without introgressions. The numbers are grouped per clade as detected after mapping to the S288c reference genome alone (Including introgressions, blue) or to the multi species concatenated reference as described in Methods (Excluding introgressions, red). Median is shown in red. Wine, Mixed Origin and Taiwanese are shown as reference.

Supplementary Figure S4. Number of single nucleotide variants of strains representing the world-wide diversity arranged by clade. Neotropical clades are at the beginning and the rest are sorted by the median shown in red. The variants from the newly sequenced strains called as described in Methods were merged with the vcf from Peter *et al.*, 2018, and the number of variants was calculated using bcftools 1.9.

Supplementary Figure S5. Population structure analysis with ADMIXTURE. (A) Cross-validation error analysis, where each dot is one of ten replicates at different K values. Cyan and magenta lines are the mean and the median, respectively. (B) Admixture population structure analysis of the simulation with the lowest cross-validation errors (K=3 to K=26). South American Mix clades (Tellini 2024) include the following: SAM1 is formed by strains of B1 lineage of Barbosa (2016) and Ecuadorean strains from Peter (2018), SAM2 is formed by B3 lineage of Barbosa (2016) and different unassigned strains from the Americas from Peter (2018), and SAM3 includes lineage B4 of Barbosa (2016), Cachaca strains of Barbosa (2021) and a Brazil spirit strain from Gallone (2016).

Supplementary Figure S6. Biocultural regions of agave fermentation for spirit production show a gradient in genetic diversity. Regions are as in Gallegos-Casillas *et al.*, 2024.

Supplementary Figure S7. Genetic diversity parameters in clades of yeasts from the Neotropical cluster. Distribution of Heterozygosity (top), Genetic diversity, π (middle), and Tajima's D (bottom). Each matrix at the right shows the *p*-value of a Mann-Witney paired test.

Supplementary Figure S8. Genomic map of introgressions shows similar distributions regardless of the specific method to define them. For each locus, the color scale indicates the percentage of strains in each phylogenetic clade with introgression, (A) using the coverage of depth after competitive mapping to the concatenated multispecies reference and (B) using the *bona fide* markers from Tellini et al. 2024.

Supplementary Figure S9. Introgression block size per clade of the Neotropical cluster. Strains with more genes per block tend to also show a higher number of introgression blocks.

783
784
785

Figure 1

Figure 2

Figure 3

Figure 4

A)

B)

Supplementary Figure S1

Collection Year
- ○ 1988
- ○ 1992
- ○ 2008-2009
- ○ 2013-2016
- ○ 2018-2021
- ○ NA

Reference
- △ Gallegos−Casillas et al., Yeast, 2024
- ▽ Lopez Gallegos, 2023
- ✳ Provided by Manuel Kirchmayr and Anne Gschaedler
- ◇ Lachance, A v L, 1995
- ○ Alvarez−Ainza ML et al., App Biochem and Biotech, 2015
- + Kirchmayr et al, CIATEJ, 2014
- ✕ Kirchmayr et al., LWT− Food Science and Technology, 2017
- ⊠ Quezada−Romero et al., ISA, 2016
- ⊕ Verdugo−Valdez et al., A v L, 2011

Supplementary Figure S2

A)

Neotropical cluster

58 97
65 97
80

Wine

0.040

Mosaic
Region 3

Wild

Mexican Agave 1 ●
Mexican Agave 2 ●
French Guiana ●
South American Mix 2 ●
Tequila Distillery ●
Alpechin ●

B)

Neotropical cluster

Wine

0.040

Mosaic
Region 3

Wild

C)

Neotropical cluster

100
100

Wine

Mosaic
Region 3

Wild

0.040

Supplementary Figure S3

Supplementary Figure S4

Supplementary Figure S5

Supplementary Figure S6

Supplementary Figure S7

Supplementary Figure S8

Supplementary Figure S9