Recurrent introgression and geographical stratification shape *Saccharomyces cerevisiae* in the Neotropics

J Abraham Avelar-Rivas[1], Iván Sedeño[2,3], Luis Fernando García-Ortega[4], Jose A Urban Aragon[2], Eugenio Mancera[4,*], Alexander DeLuna[1,*], Lucía Morales[2,*]

[1]*Unidad de Genómica Avanzada, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Irapuato, Mexico.*

[2] *Laboratorio Internacional de Investigación sobre el Genoma Humano (LIIGH), Universidad Nacional Autónoma de México, 76230 Juriquilla, Mexico*

[3] *Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Coyoacán, 04510 CDMX, Mexico*

[4] *Departamento de Ingeniería Genética, Unidad Irapuato, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Irapuato, Mexico.*

Correspondence to: eugenio.mancera@cinvestav.mx, alexander.deluna@cinvestav.mx, lmorales@liigh.unam.mx

Running title: Introgressions and population structure in *S. cerevisiae* from the Neotropics

Key words: Evolutionary genomics, Genetic introgression, Gene flow, Geographic population structure, *Saccharomyces cerevisiae*, Agave fermentation

Highlights:

1. Phylogenetic analyses of *S. cerevisiae* isolates from agave fermentation reveal novel clades within the Neotropical cluster.
2. The Neotropical cluster shows geographical population structure.
3. Multiple pulses of interspecific introgression are part of the evolutionary history of *S. cerevisiae* in the Neotropics

ABSTRACT

From yeasts to humans, introgressive hybridization significantly influences the evolutionary history of living organisms by introducing new genetic diversity. Strains of *Saccharomyces cerevisiae* worldwide exhibit introgressions from the sister species *S. paradoxus*, despite the average sequence identity between these species being lower than 90%. While *S. cerevisiae* isolates from the Neotropics are known for their high levels of introgression, the evolutionary events leading to the unusually high prevalence of them remain unclear. Here, we sequenced 216 *S. cerevisiae* isolates living in open agave fermentation across Mexico, a habitat at the interface of natural and industrial environments. The genomes of these strains revealed considerable genetic diversity and population structure linked to geographic distribution, which had been overlooked due to undersampling of this megadiverse region. These strains, along with those from French Guiana, Ecuador, and Brazil, form a broader Neotropical phylogenetic cluster that is notably enriched in introgressed DNA. Surprisingly, the origins and conservation patterns of introgressions indicate multiple hybridization events, suggesting an unprecedented scenario of flexible species barriers in this region. Our findings underscore concurrent evolutionary processes—geographical stratification and multiple introgressions—that shape the genomes of a diverse lineage of *S. cerevisiae*. Neotropical yeasts thus provide a natural laboratory for exploring the mechanisms and adaptive significance of introgressive hybridization in eukaryotic genome evolution.

INTRODUCTION

Beyond its long-established status as the most extensively studied eukaryote in the laboratory, the budding yeast *S. cerevisiae* has emerged as an outstanding model for ecology, population genomics, and evolution [1–5]. Genomic analyses of thousands of isolates from natural and human-related habitats around the world reveal that the species is structured in clades that broadly correlate with geography and habitat of isolation [6–8]. These global genomic studies have also shown that most strains harbor genetic material from other *Saccharomyces* species, a phenomenon referred to as genetic introgression. For instance, most *S. cerevisiae* isolates from around the world harbor genes introgressed from its sister species *Saccharomyces paradoxus* [9,10,6,11,12], even though these species diverged 4-6 MYA and have an average nucleotide divergence of ~14% [13,14].

The Neotropical region, which extend from northern Mexico to South America, is biogeographically relevant due to its rich biodiversity [15]. Some of the *S. cerevisiae* strains from this area isolated from anthropogenic and natural environments group in a long-monophyletic branch. This diverged branch, referred to as a complex clade by Pontes et al. (2020), aggregates three groups of strains, namely Mexican Agave (MA), French Guiana (FG), and South American Mix 2 (SAM2). Notoriously, the handful of sequenced genomes available to date indicate that this Neotropical cluster harbors an unusually high and heterogeneous number of introgressions from *S. paradoxus* [6,7,16,12]. Considering the vast area encompassed by this region, the few available sequenced isolates likely provide an underestimate of genetic diversity, thus limiting our understanding of the species evolutionary history in this part of the world.

The elevated numbers of introgressions observed in *S. cerevisiae* strains from the

Neotropical cluster could explain its high divergence when compared to other clades.

Although a fraction of the introgressions from the Neotropical cluster has been associated

with populations of *S. paradoxus* from the Americas [6,12,17], their precise origins remain to

be determined. In Mexico, there are reports of *S. paradoxus* in fermentations [18,19],

including a lineage that coexists with *S. cerevisiae* in the agave distillery environment and

that has not been considered by previous studies [18]. Therefore, this *S. paradoxus* lineage

could be a potential donor for introgressions in the tropical Americas. Comprehensive

understanding of both the vertical evolution of *S. cerevisiae* and the origins of its

introgressions is essential to accurately determine the dynamics of interspecies gene flow

in the region.

To explore the evolutionary history and genomic diversity of *S. cerevisiae* in the tropical

Americas in the context of their strong signatures of introgression, we sequenced 216

strains primarily isolated from open agave fermentations in Mexico. Previous studies have

highlighted the megadiversity in this region [20] as well as high diversity of *S. cerevisiae* in

agave fermentation [21–24], but whole-genome sequencing efforts have only included nine

strains, mostly from a single agave fermenting region [6]. We show that most of the newly

sequenced isolates belong to the Neotropical cluster, including new clades where diversity

correlates with the geographic origin of the strains. Moreover, by analyzing the patterns

and origins of the introgressions in the context of the relatedness of these lineages, we

identified multiple introgression events in their evolutionary history. Notably, some of

these introgression pulses were unique to strains isolated from agave fermentations. Our

refined phylogeny allowed us to illustrate how recurrent episodes of gene flow between

two sister *Saccharomyces* species have unfolded in a megadiverse region of the world.

RESULTS

To provide a comprehensive view of *S. cerevisiae* diversity in one of the most biodiverse

regions in the world [15,20], we sequenced the genomes of 216 isolates across Mexico

(Supplementary Figure S1, Supplementary Table S1). Strains were primarily isolated from

traditional agave distilleries. This fermentation environment provides a good starting point

to study yeasts diversity, as spontaneous fermentations have taken place for over 3,500

years in a wide variety of ecosystems, latitudes, and cultural contexts [25–27]. Specifically,

211 of the sequenced strains were isolated from agave fermentations collected through

various sampling efforts conducted across the country between the years 1988 and 2021

[28,21,29,24,30,22,31,18]. Four of the five remaining isolates were from fermentations of other

substrates — pulque (fermented raw agave sap), apple cider, cactus pears, and cooked

stems of sotol plants (*Dasylirion* spp.)— while the fifth was an uncharacterized commercial

yeast strain used in one of the sampled distilleries; all of them were isolated in Mexico

(Supplementary Table S1). With the 216 genomes (median depth coverage = 163X), we

assessed the phylogenetic position and population structure of the strains in the context of

the known diversity of the species and we also analyzed the dynamics and origins of the

prevalent *S. paradoxus* introgressions within these populations.

Agave fermentation strains belong to the Neotropical cluster

To assess the relatedness of the agave *S. cerevisiae* strains with other isolates of the species,

we first performed multidimensional scaling analysis (MDS) with 1,264 genomes,

including the 216 newly sequenced strains. Most isolates from Mexico clustered next to

other strains from the tropical Americas, such as those from the French Guiana (FG), the

South American Mix 2 (SAM2), and the Mexican Agave 1 (MA1, formerly named

"Mexican Agave") clades (Figure 1A). The rich diversity of the newly sequenced strains is

evident, as strains from the tropical Americas encompass nearly half of the global diversity

shown in MDS component 1, depicted by gray points, and nearly the entire diversity in

component 2 (Figure 1A). Only eleven strains grouped with other clades, namely the

Mixed-origin, Wild North American, or Wine clades. The population-structure

(ADMIXTURE, K=3) showed that ~97% of the agave fermentation strains had a

Neotropical ancestral genetic component (Figure 1B), distinguishing strains from the

Neotropics (blue) from wild (green) and domesticated (red) isolates from other regions of

the world. The ADMIXTURE analysis also indicated that strains from the tequila industry

were highly admixed, bearing a combination of the three ancestral components (Figure 1B).

All five isolates from non-agave substrates that we sequenced grouped close to the agave-

fermentation strains from the Neotropics.

Figure 1. A novel, diverse yeast lineage encompasses most of the diversity in open agave fermentations. (A) MDS analysis of 1,264 genomes (173,331 informative sites). Large dots are strains from the Tropical Americas Neotropics, colored according to their clade: "Mexican Agave 1" (purple), "Mexican Agave 2" (blue), "French Guiana" (orange), "Tequila Distillery" (yellow), and "South American Mix 2" (SAM2, pink). Small gray dots indicate isolates from other parts of the world, with the Wine and Wild strains shown as references. Black solid outlines are strains sequenced in this study (n=216). Gray circles with a solid outline indicate the few isolates from agave fermentation that did not group with other Neotropical strains. **(B)** ADMIXTURE population structure at K=3. At the bottom, strains from the Neotropical cluster, with their phylogenetic clades indicated in the first column and map at the left. **(C)** Maximum likelihood phylogenetic tree constructed using 954,449 SNPs in 464 strains including a subset of representative reference

genomes and excluding the two admixed clades from the tropical Americas with poor branch
support, Tequila distillery, and SAM2.

To establish the phylogenetic relationships of the newly sequenced isolates, we compared
them to a reference panel of strains (see Supplementary Table S2). Consistent with the
MDS analysis, the phylogeny showed that 205 of the 216 newly sequenced strains (94.9%)
were part of a monophyletic group that included the clades MA1, a newly defined clade
Mexican Agave 2 (MA2) and FG, plus the two admixed clades Tequila Distillery and
SAM2 (Supplementary Figure S2A-B). We refer to this higher hierarchy clade as the
Neotropical cluster of *S. cerevisiae*, as nearly all the strains in this group come from the
tropical Americas (Figure 1B). Of the remaining sequenced isolates, five strains grouped
with the wild North America Oak clade; other five—including the commercial strain—
were closely related to the Mixed Origin clade and only one clustered with Wine strains.

The mosaic strains of Tequila Distillery and SAM2 branched with poor support and their
presence in the phylogeny changed the relative placement of MA1, MA2 and FG
(Supplementary Figure S2). For that reason, these admixed strains were excluded from the
main phylogenetic analysis (Figure 1C). In contrast, when SNPS from introgressed
segments were removed from the phylogeny, the clades of the Neotropical cluster remained
the same. The presence of introgressed regions had a less significant impact on the branch
lengths of the diverse Neotropical cluster compared to those of the Alpechin clade
(Supplementary Figure S2). Furthermore, the difference in the number of single nucleotide
variants was more pronounced for Alpechin than for the Neotropical cluster after the
removal of introgression (Supplementary Figure S3).

Three clades could be distinguished among the newly sequenced isolates from agave fermentations in Neotropical *S. cerevisiae*. First, we identified a group of 20 isolates from northeastern Mexico, which included the seven strains previously reported as "Mexican Agave" [6] herein referred to as the MA1 clade (Figure 1C). It is worth mentioning that the monophyletic MA1 clade comprised only strains isolated northeast of the Sierra Madre Oriental, a mountain range which runs parallel to the coast of the Gulf of Mexico and acts as a natural barrier, exerting strong influence on the climate and biodiversity on either side of the range [32,33]. As a result of the comprehensive sampling of the isolates reported here, we identified a novel MA2 clade closely related to MA1 (Figure 1C). Most of the newly sequenced strains in this study (81.9%) belonged to this clade. Two sequenced strains isolated in Mexico [6] were previously assigned to the SAM2 clade [12] and after these analyses were reclassified within the new MA2 clade. The third clade of agave isolates was composed of 15 strains with heterogenous and admixed genotypes, all sequenced in this study (Figure 1B; Supplementary Figure S2). We refer to this as the "Tequila Distillery" clade, since eleven were isolated from the tequila industry.

Our results show that most strains from open agave fermentations are part of the Neotropical cluster, revealing that what was previously described as "Mexican Agave" represents only a minor fraction of the great genetic diversity found in agave fermentations. Furthermore, the clades within this cluster, including the newly identified groups, rank among the highest in terms of single nucleotide variants when compared to the reference genome, surpassed only by the wild Asian clades (**Supplementary Figure S4**). Together, our phylogenomic and population analyses of *S. cerevisiae* from the tropical Americas bridge the gap of genetic knowledge for the species in this megadiverse region of the world.

Neotropical yeasts belong to diverse populations stratified by geography

Yeasts from the tropical Americas come from a diverse variety of environments across a broad range of geographical distances. We thus asked to what extent geographic parameters influence the population structure of *S. cerevisiae* in the Neotropical cluster. To this end, we carried out an ADMIXTURE analysis assuming the number of populations that better explained the genotypes according to the cross-validation error (Figure 2A; Supplementary Figure S5). The population structure analysis revealed eleven ancestral components in the Neotropical cluster. Specifically, strains from the MA1, Tequila Distillery, FG, and SAM2 clades each clustered in unique populations, while the remaining seven populations were all from the newly reported MA2 phylogenetic clade.

Figure 2. Yeasts from the Neotropics show population structure correlated with geography.
(A) Population structure of strains from the Neotropics and other regions of the world as reference
at K=24 made with ADMIXTURE (n=466); strains are sorted according to phylogenetic clades (first
track). To the right, strain's location is shown in the maps of Mexico (top) and part of South America
encompassing Brazil, Ecuador and French Guiana. The colors of the lines connecting strains with
geographical location are the same as the most prevalent genetic component of the strain of the
Admixture plot. **(B)** Genetic diversity, π (top), heterozygosity (middle) and Tajima's D (bottom) of
clades within the Neotropical cluster.

Given the stratification observed in the Neotropical cluster, we analyzed the geographical
distribution of these isolates according to their population structure. The eleven colors in
the ADMIXTURE plot representing ancestral components within the Neotropical cluster
showed good agreement with their geographic distribution (Figure 2A). This geographic

stratification was observed at two levels of resolution. First, each of the five clades of the

Neotropical cluster are constrained to specific regions in the map. Second, at the resolution

within the MA2 clade, the seven colors in the ADMIXTURE were also stratified by

geography. In general, strains from the same agave-distillate producing region grouped

together in the phylogeny and had the same ancestral population component. This was also

shown by a correlation between geographic and genetic distance in this set of strains

(statistic $r=0.28$, $p<0.05$, Pearson Mantel test including MA1 and MA2 strains) and

increasing genetic diversity from north to south in the MA2 clade (Supplementary Figure

S6). Together, these findings suggest that the genomic diversity in the Neotropical cluster is

influenced by geography.

Despite the overall agreement between geography and population structure in the strains

from the tropical Americas, there were exceptions suggesting gene flow within and even

beyond the Neotropical lineage. For instance, a strain isolated from an open agave

fermentation shared close ancestry with wild isolates of the Wild North American clade,

while other four isolates closely related to the Wild North American clade showed signs of

admixture with different ancestral populations of the Neotropical cluster (Figure 2A).

Likewise, there were SAM2 strains showing genetic components of the MA2 and French

Guiana clades. Moreover, the Tequila Distillery showed admixture, mostly between

different Neotropical populations, but also with other clades such as Wine. Together, these

findings suggest that *S. cerevisiae* populations in the tropical Americas have diverged to

some extent independently from one another by means of geographic isolation, while other

ecological processes or human-related factors have led to gene flow, which also contributes

to their genome diversity.

We used the genomic data to further explore the patterns of genetic diversity in the

Neotropical strains. The strains from each clade varied significantly from each other in

nucleotide diversity (π), Tajima's D, and heterozygosity (Figure 2B; Supplementary Figure

S7), with some of them displaying remarkably high genetic diversity. For instance, the

MA1, MA2, and Tequila Distillery clades all show higher genetic diversity than yeasts

from the Wine clade (Supplementary Table 2). This held true when comparing with the

highly introgressed Alpechin strains, suggesting that differences in genetic diversity cannot

be solely accounted for by their introgressions. In contrast, the FG clade showed a

homogenous degree of heterozygosity and low genetic diversity. The Tequila clade had the

highest heterozygosity and nucleotide diversity, in agreement with trends observed in

domesticated beer isolates [6], while SAM2 also showed high genetic diversity, but lower

heterozygosity, as reported for wild yeast strains [34]. High diversity in these clades is in part

due to their admixed genotypes (Figure 2A), while high heterozygosity specifically in the

Tequila Distillery strains may be associated to their adaptation to the semi-industrial

context where they were isolated. Taken together, our population-structure and genetic-

diversity analyses indicate that yeasts in the tropical Americas constitute highly diverse,

structured populations stratified by geography.

Different distributions of introgressions show recurrent interspecies gene flow

The genomes of *S. cerevisiae* strains within the Neotropical cluster are characterized by an

atypically high number of introgressed regions from *S. paradoxus* [12,17]. To gain insight into

gene flow dynamics, we identified the set of introgressed genes (Materials and methods),

their location, and length patterns in each clade of the Neotropical cluster (Supplementary

Figure S8A). We found that yeasts from MA1 typically exhibited not only more

introgressed genes than all other clades, but also the highest variability in this number,

ranging from 119 to 364 introgressed genes per strain. Collectively the introgressions in

MA1 covered 7.1% of the *S. cerevisiae* reference genome (Figure 3A). We note that the

number of introgressed genes in most MA1 strains was within the range observed for the

MA2 strains which featured 96 to 160 introgressed genes. Together, all the introgressed

genes in the MA2 clade covered 3.2% of the nuclear genome. However, seven strains from

the MA1 clade exhibited longer and more heterozygous introgressions than most MA1

strains (Figure 3A; Supplementary Figure S9). Here, we call an introgression as

heterozygous when both *S. cerevisiae* and *S. paradoxus* alleles have been retained in a

single genome, suggesting a more recent introgression event. The presence of both alleles

in several introgressions of the Tequila Distillery clade suggests admixture with a non-

introgressed population, which may also explain the lower number of introgressed genes

(Figure 3A).

Figure 3. Different patterns of introgressions in the clades from the Neotropics. A) Distributions of the number of introgressed genes per strain in each phylogenetic clade. The Alpechin strains are included as a reference of a highly introgressed population. The color of each dot indicates the fraction of introgressed genes that are heterozygous in each strain, meaning that both the *S. cerevisiae* and *S. paradoxus* alleles are present. **B)** Clustering of the introgressed genes found across the Neotropical cluster (n = 1,113). Colors indicate the fraction of strains from each clade that had the introgressed gene. **C)** Distribution of the frequency of introgressions in each clade. **D)** Fraction of the maximum number of possible shared introgressed genes in each comparison. For complete comparisons see Supplementary Figure S10. **E)** "Private" introgressed genes, meaning that they are present only in one of the clades.

All Neotropical clades displayed more introgressed genes than the observed average in other groups from around the world, except for the Alpechin clade. The two clades from

South America, SAM2 and FG, displayed fewer and shorter introgressions with a lower degree of heterozygosity, compared to the two Mexican agave clades (Figure 3A; Supplementary Figure S9). This trend suggests that the last introgression pulse of SAM2 and FG likely occurred in the common ancestor of all strains from the Neotropical cluster. These characteristics in the introgression patterns agree with the trends that we observed in terms of the introgressions that are shared, whereby a significant fraction of genes was common between the Mexican Agave clades and, to a lesser extent, with the French Guiana clade (Figure 3B). Importantly, these patterns of shared introgressions were confirmed using an alternative strategy that considered *bona fide* species-specific SNP markers [12] for detecting and scoring introgressed regions (Materials and Methods; Supplementary Figure S8).

To further understand how introgressions are shared among clades, we quantified the proportion of strains harboring each introgressed gene within the different clades (Figure 3C). Introgressed genes present at high frequency—at least in 90% of the strains within a/the clade—were common in French Guiana (43.6%) and Alpechin (22.5%). The high proportion of shared introgressed genes within these populations suggests either a strong bottleneck or positive selection acting on introgressed regions. Conversely, introgressed genes present at 10% or lower frequency were more common within the Mexican Agave clades (36% and 45.7%; Figure 3C), which suggests either divergent resolutions of the same hybridization events, recent independent hybridizations in specific strains or a combination of both.

We found that MA1 and MA2 shared the highest fraction of introgressed genes in paired comparisons among clades, suggesting that they were present in their common ancestor

(Figure 3D, Supplementary Figure S10). Moreover, the highest fraction in the triple

intersection was that of the MA1, MA2 and FG (Figure 3C). These results suggest that a

fraction of the introgressions present in the three clades was present in their common

ancestor. Additionally, we observed that 14 genes were shared among all four clades,

pointing to potential strong selective pressures or artifacts arising from the use of a

reference genome lacking those specific genes. In terms of "private" introgressions—those

present in a clade but absent in all others—we scored 18.9% in FG, 32.2% in MA2, and

58.6% in MA1 (Figure 3E), while most introgressed genes in the

Alpechin reference clade (76.3%) were exclusively found in this clade (Figure 3E). The

high percentage of private introgressed genes in Alpechin is due to hybridization events not

shared with Neotropical strains [6,12]. Likewise, MA1 and MA2 isolates exhibited a high

fraction of private introgressions, potentially resulting from introgression pulses that are not

shared with the related FG clade. Together, the patterns of disparate introgression

distribution suggest recurrent interspecies gene flow episodes between *S. cerevisiae* and *S.*

paradoxus from the tropical Americas.

Introgressions in the Neotropical cluster trace to different lineages of *S. paradoxus*

To track the history of the observed interspecies gene-flow episodes, we determined the

most likely *S. paradoxus* lineage from which each introgressed gene derived. To this end,

we scored sequence similarity of each introgressed block against a panel of *S. paradoxus*

genomes including representative strains from the Americas (Figure 4A). Our results

suggest that most introgressed genes in the Mexican Agave clades derived from a lineage of

S. paradoxus that has only been isolated in Mexico and that cohabits in agave

fermentations, *SpB_Mx* [31]. In contrast, the majority of the introgressed genes in the FG

strains tracked to *SpB,* a different lineage of *S. paradoxus,* which is not commonly found in

agave distillery contexts [31]. As expected, the introgressed genes in the Alpechin group most

closely resemble the European *SpA* lineage, previously identified as the parental population

of these introgressions [35,36,12]. These observations suggest that at least two lineages of *S.*

paradoxus have hybridized with ancestors of the extant Neotropical strains, underscoring

the close relationship between these two species in the evolution of *S. cerevisiae* in the

region.

Figure 4. Multiple introgression pulses from two *S. paradoxus* lineages in the Neotropical *S. cerevisiae* cluster. (**A**) The average number of introgressed genes from each lineage of *S. paradoxus* in each clade of Neotropical *S. cerevisiae*. Error bars are the standard deviation. The

313   Alpechin clade is shown as a reference. (**B**) Scheme of the model showing multiple introgression

314   pulses as the most plausible evolutionary scenario for the observed patterns of number and origins

315   of introgressed genes along with the levels of heterozygosity and length of the introgressed blocks.

DISCUSSION

Comprehensive sequencing of *S. cerevisiae* genomes in megadiverse regions, harboring a

substantial portion of Earth's biodiversity, is essential to understand the evolutionary

history and genomic diversity of this prominent model organism. In this study, we

sequenced 216 strains from open agave fermentations, providing an extensive population

genomics dataset for the Neotropics. Our phylogenetic and population structure analyses

revealed multiple interspecies gene-flow events resulting in introgressions within this

geographically structured complex clade.

Introgressions impact evolutionary trajectories across the tree of life, from bacteria to

humans [37,38]. These events are also central to the evolutionary history of *S. cerevisiae*, as

evidenced by their presence in isolates across the globe [6,10–12]. In the Americas, several *S.*

cerevisiae lineages feature high number of introgressions from *S. paradoxus*, both in wild

and anthropogenic environments [17,7,39,12], presumably as a consequence of the co-existence

of the two species in the region. Here, we provide a broad picture of the evolutionary

dynamics of introgression happening in the tropical Americas. In the most plausible

scenario, an early introgression pulse occurred in the most recent common ancestor of the

Neotropical cluster before its diversification, where genes of the American *SpB* lineage of

S. paradoxus introgressed into *S. cerevisiae* (Figure 4B). This first pulse was presumably

followed by a subsequent introgression event from another lineage, *SpB_Mx*, in the

common ancestor of MA1 and MA2. At least one additional, more recent pulse of

introgression from the *SpB_Mx* lineage took place in the MA1 clade, which is supported by

high numbers of large and heterozygous introgressions in some strains of this clade. It

remains to be determined whether these introgression pulses are simply stochastic

byproducts of the coexistence of *S. cerevisiae* and *S. paradoxus* in the region, or if they

underlie adaptation processes. Wider sampling efforts in other areas, particularly in South

America, will likely expand our knowledge of the interspecies gene-flow dynamics in these

yeasts.

Our sequencing effort expanded knowledge of the *S. cerevisiae* diversity and

biogeography. We uncovered two clades within the Neotropical cluster: MA2 and the

admixed Tequila Distillery. Together with MA1, FG, and SAM2, these five clades

constitute the Neotropical cluster. Phylogenetic analysis indicated that the MA1—

previously designated as Mexican Agave—and MA2 are sister clades, which in turn are

the closest relatives of the FG strains, in agreement with the previously reported

phylogeny [40]. The high variability observed in the remaining SAM2 strains indicates that

there probably exist other undersampled lineages in the tropical Americas, a diverse and

biogeographically complex region (Morrone, 2014). Additional sampling and sequencing

in this region would further elucidate key evolutionary events that have shaped the natural

history of yeasts, as discussed herein.

Importantly, we found that the novel genetic diversity in the Neotropical cluster is

structured and associated with the geographic origins of the yeast isolates. Geographical

influence that shaped the genetic diversity was evident across several dimensions: First, a

clear correlation was observed between each of the eleven ancestral components of the

strains from the Neotropics and their region of isolation. Second, strains from the MA2

clade exhibit increased nucleotide diversity, in a north-to-south gradient. Finally, the

genetic divergence between the MA1 and MA2 clades is associated with the presence of the

Sierra Madre Oriental mountain range, which is found at the edge where the Neotropical

and Nearctic biogeographic regions coincide. This range is known to act as a geographical

barrier impacting the distributions of multiple species [15,32,33].

Beyond biogeographical factors, human activities may have played a role in shaping these

yeast populations, as evidenced by admixture events within and beyond the Neotropical

cluster. It remains to be addressed to what extent the observed yeast population structure is

driven by natural factors such as the ecology and biogeography of plant reservoirs and

insect vectors or by the human dynamics and practices. For instance, the specific local

production practices or the remarkable diversity of fermentation substrates, with over 50

species of *Agave* species used for producing spirits [41], could be driving population

differentiation. With the rising global demand for agave spirits there is a growing risk that

producers may abandon traditional fermentation practices. This shift could lead to the loss

of the microbial diversity herein described. This work provides a critical framework for

developing conservation and management strategies for these fungi, highlighting their

value beyond basic research. The agave fermentation system is not only a reservoir of

remarkable yeast genetic diversity, but also a unique natural laboratory to gain insights into

the interplay of genetic, ecological and evolutionary factors shaping population

differentiation, and interspecies introgression.

MATERIALS AND METHODS

Genome sequencing and variant calling. DNA was purified with the MasterPure DNA

purification kit as recommended by the manufacturer, and it was sequenced using DNBSeq

(2x150 bp; BGI, China). Raw reads were quality assessed with fastp V0.20.0 [42]. For each

isolate, the filtered reads were aligned with BWA-MEM v0.7.4 [43] to a concatenated species

reference genome that includes *S. cerevisiae* S288C, *S. paradoxus* YPS138, *S. mikatae*

IFO1815, *S. kudriavzevii* Cr85, *S. jurei* M1, *S. arboricola* H6, *S. uvarum* CBS7001, *S.*

eubayanus FM1318, *K. marxianus* DMKU31042, and *P. kudriavzevii* CBS573. Duplicated

reads were marked with Piccard 2.6.0 [44], and local realignment around indels and variant

calling were performed with GATK v4.1.1.0 [45]. Allele balance information was

incorporated in the vcf files with GATK Variant annotator. The called genotypes were

filtered to keep only biallelic SNPs with a minimum depth of 5, a QUAL score of at least

30 and SNPs presents in at least 90% of the isolates. Additionally, SNPs were also obtained

by mapping the filtered reads only to the *S. cerevisiae* genome to obtain variants that

include introgressions (Supplementary Figure S3, Supplementary Figure S4).

Multidimension scaling analysis. To analyze population structure, the generated genome-

wide SNPs data were used for multidimension scaling (MDS) analysis with plink v1.9 [46].

The data set consisted of 1,264 isolates, including 1,006 from Peter et al. (2018) and 258

from the Americas including the 216 sequenced by us. Only nuclear genome SNPs were

considered, and low-frequency or rare variants were excluded to this analysis using a

threshold of missing call rate > 1 %. After the application of this filter, isolates with

missing genotype $> 10\%$ were discarded. The quantitative indices (components) of the

genetic variation for each isolate were calculated on the genome☐wide average proportion

of alleles shared between any pair of individuals within the sample.

Phylogenetic analysis. Maximum likelihood trees were built with genomes sequenced

here and references from all over the world with emphasis in American clades. Haplotype

sequences were obtained with vcf2phylip v2.3 [47] from the nuclear genome variants derived

from the concatenated species genome that mapped to the *S. cerevisiae* sub genome (or

directly to *S. cerevisiae* reference to include variation from introgressions for

Supplementary Figure 2). The phylogeny trees were inferred by raxmlHPC-PTHREADS-

AVX2 [48] with the GTR+GAMMA parameter and 100 bootstrap replicates. The

visualization of the tree was performed using Microreact [49]. The first two phylogenetic

trees were built for 487 isolates including the 216 sequenced here (Supplementary Figure

S2). As admixed Tequila Distillery and SAM2 strains changed the topology within the

Neotropical cluster, we removed them from the phylogeny presented in Figure 1 with 464

genomes (Supplementary Table S2).

ADMIXTURE. To run ADMIXTURE 1.3.0 [50] the VCF with 466 genomes was curated

using Plink 1.9 [51] to remove linkage disequilibrium with parameters 50 window size sliding

every 5 nucleotides and correlation of 0.5 as previously run in *S. cerevisiae* population

genetics [52] which led to 618,253 SNPs. Ten seeds were used to run Admixture. The Cross

Validation (CV) error was obtained for models with k=2 to K=35 populations. The lowest

median CV error K was 24 and therefore the iteration with K=24 and the lowest CV error

was used for visualization. Visualization was made in Pong [53].

Estimation of genetic diversity. To calculate genetic diversity (π) and Tajima's D, we

used vcftools 0.1.14 [54] and the VCF with variants that mapped to *S. cerevisiae* after

mapping to the concatenated reference and filtering high quality biallelic SNPs with less

than 10% of missing data. To estimate the per variant heterozygosity, bcftools 1.9 [55] was

used on individual strain VCFs to calculate the number of heterozygous variants divided by

the total number of variants.

Analysis of Introgressions. To identify introgressed genes, we compared the *S. paradoxus*

section of the alignment to the concatenated reference. A gene was considered introgressed

if over half of it had a coverage depth exceeding 25% of the median depth of the entire

reference. Additionally, it had to be the ortholog with the highest depth or, if not, its depth

had to be at least 25% of its ortholog in the *S. cerevisiae* sub genome. We also required a

minimum 5% nucleotide-level difference between the orthologs of *S. cerevisiae* and *S.*

paradoxus. An introgressed gene was considered heterozygous if the ratio of the depth

coverage between the two sub genomes in the competitive mapping was between 0.25 and

4. Nevertheless, we alternatively used a second strategy with the same VCF file to get the

positions of a set of *bonna fide* introgression markers and retained introgression blocks of

at least five consecutive markers [12] to contrast with our mapping-based definition

(Supplementary Figure S8). Visualization was made using ChromoMap [56].

Once the introgressed genes were determined, consecutive introgressed genes according to

the annotation [57] were considered part of a block. To determine the origin of the blocks, we

aligned the *S. paradoxus* from Lopez-Gallegos, 2023 to the concatenated multi species

reference described above and kept only the regions annotated as introgressions in any of

the *S. cerevisiae* strains from this work. We computed the paired distances between all

introgressed blocks and the *S. paradoxus* genomes using bcftools 1.9 [55] and searched for

the *S. paradoxus* with the lowest distance to each introgressed block. If a given

introgression was most similar to strains from more than one *S. paradoxus* lineage, no

origin was assigned to it. All the genes within a given block were considered to be from the

same origin.

DATA ACCESSIBILITY

Genome sequencing. Genome sequencing data were deposited in the SRA under the

project PRJNA1138754. {Reviewer Link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1138754?reviewer=304775eb7b2b0uud9o

0b1aqb6e }

Custom scripts. Scripts to identify introgressed genes and their origins along with scripts

to run the figures of the article are available at: https://github.com/GELab-

LIIGH/ScerYeastGenomesMx2024

ACKNOWLEDGEMENTS

This work was funded by Consejo Nacional de Humanidades, Ciencias y Tecnologías de

México (CONAHCYT grants FORDECYT-PRONACES/103000/2020, CB-2016-

01/284992, CF-2023-G-695), Fondo de Investigación y Desarrollo Tecnológico del

Cinvestav (SEP-CINVESTAV/023), Programa de Apoyo a Proyectos de Investigación e

Innovación Tecnológica DGAPA-UNAM (grants IN209021, IN230420 and IN212524),

and the UK BBSRC under the Global Challenges Research Fund (GCRF) Growing

Research Capability call through the CABANA Innovation Fund (BB/P027849/1). EM was

funded by CONAHCYT for a sabbatical stay (I0200/111/2024). LFG-O is a postdoctoral

researcher funded by Conahcyt (4133922); IS is a PhD student at UNAM with a graduate

scholarship from Conahcyt. We thank Bernard Dujon, Xitlali Aguirre-Dugua and Maria

Avila for critical reading of the manuscript. We thank Luis Aguilar (LAVIS, UNAM),

Porfirio Gallegos (Cinvestav), Claudio Lopez (Cinvestav), Aaron de Luna (LIIGH),

Alejandra Castillo (LIIGH), Carina Uribe (LIIGH), Maritrini Colón-González (LIIGH) and

Jair García (LIIGH) for technical assistance. We thank Xitlali Aguirre-Dugua, Luis Delaye,

Diego Ortega, Marcela Sandoval and Alicia Mastretta-Yanes for helpful discussions and

Manuel Kirchmayr (CIATEJ), Anne Gschaedler (CIATEJ), Maritza Álvarez (CIAD) and

Marc Andre Lachance for sharing their strains.

MANUSCRIPT ITEMS

Figures

Figure 1. A novel, diverse yeast lineage encompasses most of the diversity in open agave fermentations.

Figure 2. Yeasts from the Neotropical cluster show population structure correlated with geography.

Figure 3. Different patterns of introgressions in the clades from the tropical Americas.

Figure 4. Multiple introgression pulses from two *S. paradoxus* lineages in the Neotropical *S. cerevisiae* clades.

Supplementary Tables

Table S1. The 216 Strains sequenced in this study and associated metadata.

Table S2. Strains and metadata used in this study.

Supplementary Figures

Supplementary Figure S1. Origin of the sequenced strains.

Supplementary Figure S2. Phylogenies of the Neotropical cluster show the same groups with or without introgressions, but their relative placement is affected by Tequila Distillery strains.

Supplementary Figure S3. Number of single nucleotide variants of strains with or without introgressions.

Supplementary Figure S4. Neotropical strains grouped in clades show a high number of single nucleotide variants compared to the worldwide diversity.

Supplementary Figure S5. Population structure analysis.

Supplementary Figure S6. Biocultural regions of agave fermentation for spirit production show a gradient in genetic diversity.

Supplementary Figure S7. Genetic diversity parameters in clades of yeasts from the Neotropics.

Supplementary Figure S8. Genomic map of introgressions shows similar distributions regardless of the specific method to define them.

505   **Supplementary Figure S9.** Introgression block size per clade of the Tropical Americas

506   linage Neotropical cluster.

507   **Supplementary Figure S10.** Fraction of shared introgressed genes between highly

508   introgressed clades.

REFERENCES

REFERENCES

1. Liti, G. The Natural History of Model Organisms: The fascinating and secret wild life of the

budding yeast S. cerevisiae. *eLife* https://elifesciences.org/articles/05835 (2015)

doi:10.7554/eLife.05835.

2. Marcet-Houben, M. & Gabaldón, T. Beyond the Whole-Genome Duplication: Phylogenetic

Evidence for an Ancient Interspecies Hybridization in the Baker's Yeast Lineage. *PLOS Biol.* **13**,

e1002220 (2015).

3. Alsammar, H. & Delneri, D. An update on the diversity, ecology and biogeography of the

Saccharomyces genus. *FEMS Yeast Res.* **20**, foaa013 (2020).

4. Bendixsen, D. P., Gettle, N., Gilchrist, C., Zhang, Z. & Stelkens, R. Genomic Evidence of an

Ancient East Asian Divergence Event in Wild Saccharomyces cerevisiae. *Genome Biol. Evol.* **13**,

evab001 (2021).

5. Wang, M. *et al.* Annotation of 2,507 Saccharomyces cerevisiae genomes. *Microbiol. Spectr.* **12**,

e03582-23 (2024).

6. Peter, J. *et al.* Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature* **556**, 339–

344 (2018).

7. Pontes, A., Hutzler, M., Brito, P. H. & Sampaio, J. P. Revisiting the Taxonomic Synonyms and

Populations of Saccharomyces cerevisiae—Phylogeny, Phenotypes, Ecology and Domestication.

Microorganisms **8**, 903 (2020).

8. Loegler, V., Friedrich, A. & Schacherer, J. Overview of the Saccharomyces cerevisiae population

structure through the lens of 3,034 genomes. 2024.09.16.613241 Preprint at

https://doi.org/10.1101/2024.09.16.613241 (2024).

9. Duan, S.-F. *et al.* The origin and adaptive evolution of domesticated populations of yeast from Far

East Asia. *Nat. Commun.* **9**, 2690 (2018).

10. Strope, P. K. *et al.* The 100-genomes strains, an S. cerevisiae resource that illuminates its natural

phenotypic and genotypic variation and emergence as an opportunistic pathogen. *Genome Res.* **25**,

762–774 (2015).

11. Clark, A., Dunham, M. J. & Akey, J. M. The genomic landscape of Saccharomyces paradoxus

introgression in geographically diverse Saccharomyces cerevisiae strains. 2022.08.01.502362

Preprint at https://doi.org/10.1101/2022.08.01.502362 (2022).

12. Tellini, N. *et al.* Ancient and recent origins of shared polymorphisms in yeast. *Nat. Ecol. Evol.* **8**,

761–776 (2024).

13. Boynton, P. J. & Greig, D. The ecology and evolution of non-domesticated Saccharomyces species.

Yeast Chichester Engl. **31**, 449–462 (2014).

14. Shen, X.-X. *et al.* Tempo and Mode of Genome Evolution in the Budding Yeast Subphylum. *Cell*

175, 1533-1545.e20 (2018).

15. Morrone, J. J. Biogeographical regionalisation of the Neotropical region. *Zootaxa*

3782, 1–110 (2014).

16. De Chiara, M. *et al.* Domestication reprogrammed the budding yeast life cycle. *Nat. Ecol. Evol.* **6**,

448–460 (2022).

17. Barbosa, R. *et al.* Evidence of Natural Hybridization in Brazilian Wild Lineages of Saccharomyces

cerevisiae. *Genome Biol. Evol.* **8**, 317–329 (2016).

18. Gallegos-Casillas, P. *et al.* Yeast diversity in open agave fermentations across Mexico. *Yeast* **41**,

35–51 (2024).

19. Peris, D. *et al.* Macroevolutionary diversity of traits and genomes in the model yeast genus

Saccharomyces. *Nat. Commun.* **14**, 690 (2023).

20. Mittermeier, R. A. & Mittermeier, C. G. *Megadiversity: Earths Biologically Wealthiest Nations*.

(CEMEX, 2005).

21. Lachance, M.-A. Yeast communities in a natural tequila fermentation. *Antonie Van Leeuwenhoek*

68, 151–160 (1995).

22. Verdugo Valdez, A. *et al.* Yeast communities associated with artisanal mezcal fermentations from

Agave salmiana. *Antonie Van Leeuwenhoek* **100**, 497–506 (2011).

23. Álvarez-Ainza, M. L., Zamora-Quiñonez, K. A., Moreno-Ibarra, G. M. & Acedo-Félix, E. Genomic

Diversity of Saccharomyces cerevisiae Yeasts Associated with Alcoholic Fermentation of Bacanora

Produced by Artisanal Methods. *Appl. Biochem. Biotechnol.* **175**, 2668–2676 (2015).

24. Kirchmayr, M. R. *et al.* Impact of environmental conditions and process modifications on microbial

diversity, fermentation efficiency and chemical profile during the fermentation of *Mezcal* in

Oaxaca. *LWT - Food Sci. Technol.* **79**, 160–169 (2017).

25. Bruman, H. J. *Alcohol in Ancient Mexico*. (University of Utah Press, 2000).

26. Arellano-Plaza, M., Paez-Lerma, J. B., Soto-Cruz, N. O., Kirchmayr, M. R. & Gschaedler Mathis,

A. Mezcal Production in Mexico: Between Tradition and Commercial Exploitation. *Front. Sustain.*

Food Syst. **6**, (2022).

27. Colón-González, M. *et al.* Thriving in Adversity: Yeasts in the Agave Fermentation Environment.

Authorea https://www.authorea.com/users/838127/articles/1229396-thriving-in-adversity-yeasts-in-

the-agave-fermentation-environment (2024) doi:10.22541/au.172782726.68006909/v1.

28. Padilla-Camberos, E., Pinal-Zuazo, L. & Alvarez de la Cuadra Jacob, J. Catálogo de la colección de

cultivos microbianos. (1994).

29. Kirchmayr, M. R. *et al. Manual para la estandarización de los procesos de producción del mezcal*

guerrerense. vol. 1 (Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de

Jalisco A.C., Guadalajara, Jalisco. México, 2014).

30. Quezada, R., Gschaedler, A. & Kirchmayr, M. Characterization of microbial population dynamics

associated with different juices of Agave tequilana. in *Sustainable and Integrated use of Agave* vol.

2016 2016 (2016, Zapopan, Jalisco, Mexico, 2016).

31. Lopez-Gallegos, C. E. Population genomics and ecology of yeast associated with traditional Agave

fermentation in Mexico. (CINVESTAV del IPN, https://liigh.unam.mx/yeastgenomesmx/wp-

content/uploads/2024/09/TesisDeMaestria_ClaudioLopez_Marzo2023-1.pdf , Irapuato,

Guanajuato, Mexico, 2023).

586 32. Rzedowski, J. *Vegetación de México.* (1978).

587 33. Mastretta-Yanes, A., Moreno-Letelier, A., Piñero, D., Jorgensen, T. H. & Emerson, B. C.

588   Biodiversity in the Mexican highlands and the interaction of geology, geography and climate within

589   the Trans☐Mexican Volcanic Belt. (2015) doi:10.1111/jbi.12546.

590 34. Han, D.-Y. *et al.* Adaptive Gene Content and Allele Distribution Variations in the Wild and

591   Domesticated Populations of Saccharomyces cerevisiae. *Front. Microbiol.* **12**, (2021).

592 35. Pontes, A., Čadež, N., Gonçalves, P. & Sampaio, J. P. A Quasi-Domesticate Relic Hybrid

593   Population of Saccharomyces cerevisiae × S. paradoxus Adapted to Olive Brine. *Front. Genet.* **10**,

594   (2019).

595 36. D'Angiolo, M. *et al.* A yeast living ancestor reveals the origin of genomic introgressions. *Nature*

596   **587**, 420–425 (2020).

597 37. Diop, A., Torrance, E. L., Stott, C. M. & Bobay, L.-M. Gene flow and introgression are pervasive

598   forces shaping the evolution of bacterial species. *Genome Biol.* **23**, 239 (2022).

599 38. Edelman, N. B. & Mallet, J. Prevalence and Adaptive Impact of Introgression. *Annu. Rev. Genet.*

600   **55**, 265–283 (2021).

601 39. Jacobus, A. P. *et al.* Comparative Genomics Supports That Brazilian Bioethanol Saccharomyces

602   cerevisiae Comprise a Unified Group of Domesticated Strains Related to Cachaça Spirit Yeasts.

603   *Front. Microbiol.* **12**, (2021).

604 40. O'Donnell, S. *et al.* 142 telomere-to-telomere assemblies reveal the genome structural landscape in

605   Saccharomyces cerevisiae. 2022.10.04.510633 Preprint at

606   https://doi.org/10.1101/2022.10.04.510633 (2022).

607 41. Colunga García Marín, S. P. *et al. Los agaves y las prácticas mesoamericanas de aprovechamiento,*

608   *manejo y domesticación.* (2017).

609 42. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

610   *Bioinformatics* **34**, i884–i890 (2018).

611 43. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. Preprint at

612   https://doi.org/10.48550/arXiv.1303.3997 (2013).

44. Picard, toolkit. Picard toolkit. *Picard Tools* https://broadinstitute.github.io/picard/.

45. McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

46. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage

Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

47. Ortiz, E. M. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for phylogenetic

analysis. zenodo https://doi.org/DOI:10.5281/zenodo.2540861.

48. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

49. Argimón, S. *et al.* Microreact: visualizing and sharing data for genomic epidemiology and

phylogeography. *Microb. Genomics* **2**, e000093 (2016).

50. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated

individuals. *Genome Res.* **19**, 1655–1664 (2009).

51. Purcell, S. & Chang, C. PLINK 1.9. www.cog-genomics.org/plink/1.9/.

52. Gallone, B. *et al.* Domestication and Divergence of Saccharomyces cerevisiae Beer Yeasts. *Cell*

166, 1397-1410.e16 (2016).

53. Behr, A. A., Liu, K. Z., Liu-Fang, G., Nakka, P. & Ramachandran, S. pong: fast analysis and

visualization of latent clusters in population genetic data. *Bioinformatics* **32**, 2817–2823 (2016).

54. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

55. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008 (2021).

56. Anand, L. & Rodriguez Lopez, C. M. ChromoMap: an R package for interactive visualization of

multi-omics data and annotation of chromosomes. *BMC Bioinformatics* **23**, 33 (2022).

57. Yue, J.-X. *et al.* Contrasting evolutionary genome dynamics between domesticated and wild yeasts.

Nat. Genet. **49**, 913–924 (2017).

SUPPLEMENTARY FIGURES CAPTIONS

Supplementary Figure S1. Origin of the 216 sequenced strains.

 Supplementary Figure S2. Phylogenies of the Neotropical cluster show the same groups
with or without introgressions, but their relative placement is affected by mosaic Tequila
Distillery and SAM2 strains. Maximum likelihood phylogenetic trees constructed including
a subset of representative reference genomes. (A) Maximum likelihood phylogeny with a
complete set of 487 strains and 1,002,285 SNPs excluding introgressions. (B) Maximum
likelihood phylogeny of the same set of genomes, including SNPs within introgressions
(1,177,709 SNPs). (C) Maximum likelihood of 464 strains as described in the main text.
The bootstrap values of relevant branching points show how the Tequila Distillery mosaics
hamper reconstruction of the branching order of the major phylogenetic clades. Colored
dots at the tip of the branches show the origins of *S. cerevisiae* strains from the Neotropical
cluster: Mexican Agave 1 (purple), Mexican Agave 2 (blue), French Guiana (orange),
Tequila Distillery (yellow), South American Mix 2 (pink); Alpechin strains are also
highlighted (salmon).

656 **Supplementary Figure S3.** Number of single nucleotide variants of strains with or without

657 introgressions. The numbers are grouped per clade as detected after mapping to the S288c

658 reference genome alone (Including introgressions, blue) or to the multi species

659 concatenated reference as described in Methods (Excluding introgressions, red). Median is

660 shown in red. Wine, Mixed Origin and Taiwanese are shown as reference.

661
662 **Supplementary Figure S4.** Number of single nucleotide variants of strains representing
663 the world-wide diversity arranged by clade. Neotropical clades are at the beginning and the
664 rest are sorted by the median shown in red. The variants from the newly sequenced strains
665 called as described in Methods were merged with the vcf from Peter *et al.*, 2018, and the
666 number of variants was calculated using bcftools 1.9.

A)

B)

K=2
K=3
K=4
K=5
K=6
K=7
K=8
K=9
K=10
K=11
K=12
K=13
K=14
K=15
K=16
K=17
K=18
K=19
K=20
K=21
K=22
K=23
K=24
K=25
K=26

Wild Asian
Wild North Am
SAM1
Mixed Origin
Wine2
Apechin
Wine1
SAM3
Tequila Distillery
Mexican Agave 1
SAM2
French Guiana
Mexican Agave 2

Supplementary Figure S5. Population structure analysis with ADMIXTURE. (A) Cross-

validation error analysis, where each dot is one of ten replicates at different K values. Cyan

and magenta lines are the mean and the median, respectively. (B) Admixture population

structure analysis of the simulation with the lowest cross-validation errors (K=3 to K=26).

South American Mix clades (Tellini 2024) include the following: SAM1 is formed by

strains of B1 lineage of Barbosa (2016) and Ecuadorean strains from Peter (2018), SAM2 is

formed by B3 lineage of Barbosa (2016) and different unassigned strains from the

Americas from Peter (2018), and SAM3 includes lineage B4 of Barbosa (2016), Cachaca

strains of Barbosa (2021) and a Brazil spirit strain from Gallone (2016).

Supplementary Figure S6. Biocultural regions of agave fermentation for spirit production

show a gradient in genetic diversity. Regions are as in Gallegos-Casillas *et al.*, 2024.

Supplementary Figure S7. Genetic diversity parameters in clades of yeasts from the Neotropical cluster. Distribution of Heterozygosity (top), Genetic diversity, π (middle), and Tajima's D (bottom). Each matrix at the right shows the *p*-value of a Mann-Witney paired test.

Supplementary Figure S8. Genomic map of introgressions shows similar distributions
regardless of the specific method to define them. For each locus, the color scale indicates
the percentage of strains in each phylogenetic clade with introgression, (A) using the
coverage of depth after competitive mapping to the concatenated multispecies reference
and (B) using the *bona fide* markers from Tellini et al. 2024.

Supplementary Figure S9. Introgression block size per clade of the Neotropical cluster. Strains with more genes per block tend to also show a higher number of introgression blocks.

697 **Supplementary Figure S10.** Fraction of shared introgressed genes between highly introgressed clades.
698 The dark gray bars are the same shown in Figures 3D-E. They highlight cases where the fraction is divided by
699 the maximum number of possible shared introgressed genes in each comparison. The maximum number of
700 possible shared introgressed genes is the number of introgressions present in the least introgressed clade.