The long-term evolutionary potential of four yeast species and their hybrids in extreme temperature conditions

Javier Pinto[1*], Rike Stelkens[1]

[1]Department of Zoology, Stockholm University, Svante Arrheniusväg 18 B, 106 91 Stockholm, Sweden

[*]Corresponding author: Javier Pinto, Department of Zoology Stockholm University, telephone: +46 (0) 816 4223; email: javier.pinto@zoologi.su.se

Abstract
Accelerating climate change and extreme temperature events urge us to better understand
the potential of populations to tolerate and adapt to thermal challenges. Interspecific
hybridization can facilitate adaptation to novel or extreme environments. However,
predicting the long-term fitness effects of hybridization remains a major challenge in
evolutionary and conservation biology. Experimental evolution with microbes provides a
powerful tool for tracking adaptive processes across hundreds of generations in real time.
Here, we investigated thermal adaptation dynamics of four species of budding yeast
(*Saccharomyces*) and their interspecific F2 hybrids, for 200 generations under extremely cold
(5°C) and warm (31°C) conditions. We found significant variation in the evolutionary potential
of species and hybrids. Cold-tolerant species showed larger fitness increases in warm
temperature, whereas warm-tolerant species showed larger fitness gains in cold
temperature. By far the largest fitness improvements occurred in hybrids, with some
populations nearly quadrupling in fitness in the cold environment over the course of
experimental evolution. Some hybrids exceeded both their parents in thermal adaptive
potential. Reciprocal transplanting of evolved populations from the endpoint of evolution into
opposite temperatures revealed that hybrids generally have greater resilience than their
parents when challenged with sudden temperature shifts. Our results highlight that
hybridization alters the fitness outcomes of long-term adaptation to extreme environments
and may render populations more resilient to sudden environmental change, presenting both
opportunities and challenges for conservation and sustainable agriculture.
31
Keywords: climate change, experimental evolution, fitness, hybridization, *Saccharomyces,*
thermal adaptation

Lay summary
Understanding how populations adapt to extreme temperatures is crucial in times of
accelerating climate change. Hybridization and genetic exchange between species can help
populations adapt to new or extreme environments, but can also have negative effects on
fitness. Predicting the long-term effects of hybridization remains challenging for evolutionary
biologists. We usually do not know how hybrid offspring respond to future environmental
change. Experimental evolution with microbes is a powerful tool to provide insight here,
because we can observe fitness changes in real-time, across many generations. We tracked
the fitness of six species of budding yeast *Saccharomyces* and their hybrids, over 200
generations in extreme cold (5°C) and warm (31°C) conditions. We found that species and
hybrids have significantly different abilities to adapt. The largest fitness improvements were
observed in hybrid populations, particularly in the cold, where some nearly quadrupled in
fitness and exceeded both their parents. When exposing evolved populations to a sudden
shift into opposite temperatures, hybrids were more resilient than their parent species. Even
after 200 generations of evolution under opposite temperature conditions, hybrids still
showed a similar performance to their ancestors. Our results suggest that hybridization can
enhance the ability of populations to adapt to extreme environments, offering potential
benefits for conservation and agriculture.
52

53 **Introduction**
54 Human actions in the Anthropocene have led to global temperature increases and more
55 frequent extreme climatic events (Perkins-Kirkpatrick & Gibson, 2017), destabilizing
56 ecosystems and driving declines in biodiversity (Hoegh-Guldberg et al., 2018). This has

intensified the need to understand and forecast the evolutionary responses of populations to temperature changes. However, most experimental studies on the effects of climate change rely on short-term exposure of organisms to higher temperatures, often measured within a single generation only. These brief exposures do not reflect the long-term changes occurring in populations over evolutionary time.

Hybridization, the exchange of genetic material between divergent populations and species, can facilitate adaptation, increase diversity, and rescue populations from extinction because it instantly introduces new genetic variation throughout the genome (Abbott et al., 2013; Grant & Grant, 2019; Seehausen, 2004; Stelkens et al., 2014a). Hybridization rates may rise as a consequence of accelerating climate change and globalization (Chunco, 2014; King et al., 2015; McFarlane & Pemberton, 2019), posing both risks and opportunities for adaptive evolution and conservation (Brauer et al., 2023). However, our knowledge about the long-term fitness consequences of hybridization is limited. This is in part because the focus has traditionally been on long-lived species, often with small population sizes, that are of conservation concern (e.g. birds and mammals). Making hybrid crosses in these systems and using hybrids in experiments spanning multiple generations is often difficult, if not impossible. Yet, finding good conservation practices tailored to populations exposed to hybridization and introgression critically depends on a sound understanding of their long-term evolutionary potential. In an applied context, how do we assess whether supportive breeding is appropriate, if we do not know how outbred offspring respond to future environmental change? To forecast the impact of global warming on the long-term fitness and adaptation of recombined hybrids, it would be helpful to understand how the parental species' ecological and genomic backgrounds affect the hybrids' capacity to thermally adapt, across evolutionary time scales and ecologically relevant conditions.

Another question of major interest in evolutionary biology is how repeatable evolutionary responses to changing environments are. Experimental evolution studies have revealed parallelism at the gene and pathway level in bacteria and yeast (Ament-Velásquez et al., 2022; Harris et al., 2021; Kryazhimskiy et al., 2014.; Tenaillon 2012., 2014) but can we predict fitness outcomes of hybrid crosses as well? The response of hybrid populations to environmental stress can be extremely variable (Brice et al., 2021.; Gaworecki & Klaine, 2008; Lopandic, 2018; Oberhofer et al., 2014; Seehausen, 2004; Stelkens et al., 2014a; Stelkens et al., 2014b), suggesting that complex adaptations are based on different genes in different species, leading to additive, dominant and epistatic genetic interactions in hybrid genomes (Zhang et al., 2020). This highlights the importance of testing a large diversity of parental genomic and ecological backgrounds when generating predictions for the thermal adaptation of hybrids in response to climate change.

We used the microbial system budding yeast (*Saccharomyces* spp.) and experimental evolution across 200 (asexual) generations, to assess the long-term fitness evolution of interspecific hybrids in cold and hot temperatures. Yeast is well suited for this work for two main reasons. First, *Saccharomyces* yeast species vary widely in temperature preferences ranging from 15° to 35°C (Salvadó et al., 2011). Heat resistance and optimal growth temperatures are well known across the genus with some cryotolerant strains able to grow at 0°C, while some heat-resistant strains can grow up to 42°C (Belloch et al., 2008; Pérez-Torrado et al., 2015). Given that *Saccharomyces* species diverged 4-20M years ago (Borneman & Pretorius, 2014; Kellis et al., 2003; Shen et al., 2018) this suggests that thermal adaptation is a deeply diverged trait (Abrams et al., 2021; Weiss et al., 2018), providing interesting starting points for experimental evolution and a rich resource to investigate the long-term

fitness consequences of hybridization at different temperatures. Second, yeast can reproduce
asexually by clonal 'budding', allowing for experiments with large population sizes (up to
200M cells), but also sexually by mating, followed by meiosis and recombination, which allows
for controlled hybrid crosses between all species pairs despite their large evolutionary
divergence (Bendixsen et al., 2022; R. Stelkens & Bendixsen, 2022). Hybridization is known to
have substantially contributed to the origin and evolutionary history of *Saccharomyces*
(Marcet-Houben & Gabaldón, 2015; Peris et al., 2017; Steensels et al., 2021) and interspecific
yeast hybrids have been isolated from natural (D'Angiolo et al., 2020; Eberlein et al., 2019;
Leducq et al., 2016) and industrial environments, e.g. from lager beer fermentation (Gallone
et al., 2019; Gibson & Liti, 2015; Hutzler et al., 2023).
Here, we explore the fitness dynamics and evolutionary potential for thermal
adaptation across four yeast species, which cover the entire range of temperature
preferences and represent all major clades found in the *Saccharomyces* phylogeny (Bendixsen
et al., 2021), and their recombined interspecific hybrid crosses at two extreme temperatures,
5±1°C and 31±1°C. The species *S. kudriavzevii*, and *S. uvarum* are considered cold-tolerant
(Libkind et al., 2011; Naseeb et al., 2017; Salvadó et al., 2011). The Baker's yeast *S.*
cerevisiae represents the other end of the temperature spectrum and is considered warm-
tolerant, with an optimum growth temperature in the range of 30-39°C. The species *S.*
paradoxus is considered a temperature generalist that grows well in a wide range of
temperatures.
124
Materials and methods
Parental strains
We used two strains each of *S. cerevisiae* (XLY011 and Y55)*, S. paradoxus* (YPS744 and Q89.8)*,*
S. kudriavzevii (ZP591 and FM1110), and *S. uvarum* (CBS7001 and JRY9189). These strains are
representative of all major clades in the *Saccharomyces* phylogeny (Bendixsen et al., 2021),
and have a wide range of different ecologies and temperature profiles (more details on strains
and hybrids in Supplementary Table 1). All strains used in this study are derived from diploid,
non-domesticated isolates from the wild with the exception of *S. cerevisiae* XLY011, which is
a homothallic diploid derived from BY4742 and Y55 strain backgrounds commonly used in the
laboratory. All strains are HO-/- (either ho:KanMX, ho:HYGMX, or ho:NATMX), i.e. they cannot
switch mating type.
136
Construction of F1 and F2 hybrids
Haploid parents were grown overnight in liquid YEPD (1% yeast extract, 2% peptone, 2%
dextrose) in a shaking incubator at 30°C and then mixed at equal volumes. Mating was
induced by spinning down to concentrate cells and plating drops of 20µL on YEPD plates (2.5%
agar), kept at room temperature for 12h. We then streaked out for single colonies on a fresh
YEPD plate and incubated for 24 h at 30°C. Colonies were replica-plated to a sporulation plate
(KAC; 2% potassium acetate, 2% agar) and incubated at room temperature for 5 days to
induce sporulation. Diploid hybrids were identified by inspecting individual colonies for
tetrads under the microscope. A pure F1 diploid hybrid colony from each interspecies pair
was picked from the YEPD plate, previously used to streak out single colonies, then
propagated clonally in liquid YEPD overnight, and frozen in 50% glycerol for later use. To
obtain a genetically diverse, recombined F2 hybrid population, F1 hybrids were sporulated as
follows. F1 hybrids were grown in liquid YEPD overnight, cells were pelleted by centrifugation
and transferred to a 500 mL Erlenmeyer flask containing 50 mL of liquid KAC, which was

shaken at room temperature for 7 days to induce meiosis. After microscopic verification of sporulation, we centrifuged and washed cells, and resuspended them in liquid YEPD to initiate germination and population growth. These large, recombinant F2 hybrid populations (which may still contain some unsporulated F1 hybrids) were frozen in 50% glycerol solution for later use.

We used two strains of *S. cerevisiae*, a well-characterized model organism, and crossed them with two strains of three other yeast species: *S. paradoxus*, the most genetically similar; *S. uvarum*, one of the most distant; and *S. kudriavzevii*, representing an intermediate genetic distance. Thus, the two *S. cerevisiae* strains were used as the reference with which all other strains were crossed.

Experimental evolution and fitness assays

We used experimental evolution to mitotically propagate large diploid populations of all four species (eight strains in total, in three or four independent replicate evolution lines each) and six different interspecific F2 hybrids (in three independent replicate evolution lines each) at two temperatures; 5±1°C and 31±1°C (Figure 1). We chose 31°C for the hot evolution regime because this temperature still allows the two cold-tolerant species (*S. kudriavzevii*, and *S. uvarum*) to grow. Note that replicate evolution lines of hybrid crosses are 'biological replicates' in the sense that each population contains genetically different individuals resulting from independent meiosis and recombination, while replicates of pure-species lines are technical, i.e. they were isogenic at the start of evolution.

To initiate experimental evolution, we inoculated 100μL of overnight cell culture (from frozen stock populations) into glass tubes containing 5 mL of liquid YPD. This corresponds to a starting population size of approximately 3×10^7 cells. All populations evolved in the same incubator-shaker (one for each temperature) at the same time, i.e. they experienced the same slight temperature fluctuations if present. Continuous population growth was achieved by serially transferring 50μL of cell culture into 5mL fresh YPD to provide nutrients after 24 hours at 31±1°C, and 96 hours at 5±1°C. This corresponds to approximately 20 generations in each temperature regime, between serial transfers. Generation time was calculated using the formula G= t / (3.3 log b/B) (G = generation time, B = number of cells at the beginning of time interval, b = number of cells at the end of time interval). We used a total of ten serial transfers to reach approximately 200 generations. This transfer protocol leads to periodic reductions in population size but avoids bottlenecks to minimize the loss of beneficial mutations, i.e., selection is stronger than drift in these systems (Wein & Dagan, 2019).

To track population fitness across generations, we measured each population's optical density (OD at 600nm, Biotek Epoch2), which quantifies the maximum biomass before each transfer. OD_{600} was measured ten times for each evolving population across 200 generations (i.e. every 20 generations), in three technical replicates. The dilutions were standardized to approximately 0.5 OD_{600} at each serial transfer. We performed an unpaired t-test with Bonferroni-Dunn correction (at 0.05 alpha-level), to test for differences between ancestors and evolved populations. Relative fitness was calculated by dividing the fitness of evolved populations at any generation by the initial fitness of their respective ancestor (i.e. at generation 0).

In addition, we measured the fitness of populations from the endpoints of evolution (after 200 generations) after reciprocally transplanting them into the opposite temperature, i.e. populations evolved at 31°C were placed in the 5°C environment, and populations evolved at 5°C were placed in the 31°C environment. The relative fitness of these reciprocal transplant

198 populations was assessed as described above, by comparing to the fitness of the un-evolved
199 ancestors.
200

201
Figure 1. A) *Crosses between two S. cerevisiae strains and strains of S. paradoxus, S.*
*kudriavzevii and S. uvarum to obtain six different hybrids. **B)** Experimental evolution protocol*
with hypothetical fitness dynamics. Maximum biomass was measured as optical density
(OD$_{600}$) every 20 generations as a proxy of fitness before each serial transfer (red dot). Every
population was evolved in warm (31°C) and cold (5°C) conditions, in three or 4 independent
evolution lines (replicates), for 200 mitotic generations. At the endpoint of evolution, a
reciprocal transplant experiment was performed (purple square) and the fitness of cold-
evolved populations was measured in warm conditions and vice versa after the sudden
temperature shift.

211
Results
Thermal adaptation dynamics of *Saccharomyces* species
We tracked the fitness dynamics and adaptation of four *Saccharomyces* species across 200
generations in cold (5±1°C) and warm environments (31±1°C). We found significant
differences in fitness gains between species at both temperatures (Figure 2). The largest
improvement in relative fitness (compared to the ancestor's fitness at generation 0) was
found in *S. cerevisiae* at cold temperature (Figure 2A) and *S. kudriavzevii* at warm temperature
(Figure 2B). *S. cerevisiae* XLY011 produced ~120% more biomass (averaged across replicates)

than the ancestor at the end of evolution in the cold environment (t = 13.24, p < 0.01). *S. kudriavzevii ZP591* produced ~100% more biomass than the ancestor at the end of evolution in the warm environment (t = 6.77, p < 0.01).

Figure 2. *Relative fitness dynamics of eight strains of four Saccharomyces yeast species and six hybrid populations evolved for 200 generations in **A)** a cold (5±1°C) and **B)** a warm (31±1°C) environment. Lines show mean relative fitness across three or four independently evolved replicate populations per strain background. Grey-shaded areas show the standard deviations of replicates. Values above 1 represent fitness gains, values below 1 represent fitness losses, relative to the ancestral strain's fitness. Asterisks indicate significant differences between the ancestral and the evolved strains after 200 generations.*

Both of the cold-tolerant species, *S. kudriavzevii*, and *S. uvarum,* showed only a small or no fitness gain in the cold environment, but significant fitness improvements in the warm environment. The warm-tolerant *S. cerevisiae* also improved in fitness in the warm environment, but not to the same extent as in the cold environment. The thermo-generalist

S. paradoxus showed mixed outcomes, with both *S. paradoxus* strains increasing in fitness in
the cold, but only one strain improving in the warm environment.
This suggests that the long-term evolutionary potential to thermally adapt differs
between temperature conditions and among species backgrounds. Comparing warm-tolerant
with cold-tolerant species showed that warm-tolerant species generally showed higher
evolutionary potential at low temperatures (t-test with Welch correction: $t_{13,83}$= 5.76; p <
0.001), whereas cold-tolerant species showed higher evolutionary potential at warm
temperatures (t-test with Welch correction: $t_{16,45}$ = 3.77, p= 0.0065; Figure 3). Note, that as
expected, cold-tolerant species (*S. kudriavzevii*, and *S. uvarum*) had higher absolute starting
fitness than warm-tolerant species at cold temperature (Supplementary Figure 1A, test
results in Supplementary Table S2), whereas at warm temperature, warm-tolerant species (*S.
cerevisiae* and *S. paradoxus)* showed higher absolute starting fitness (Supplementary Figure
1B).

250
Figure 3 *Comparing the relative fitness gains after evolution in cold (5±1°C) and warm*
(31±1°C) environments between warm-tolerant or generalist (S. cerevisiae and S. paradoxus)
and cold-tolerant (S. kudriavzevii and S. uvarum) species. Whiskers show the smallest and
largest values, lines in the middle of the boxes represent the median and + represents the
mean. A t-test with Welch correction shows significant differences in cold and warm
*environments (p< 0.01**).*

257
258

Thermal adaptation dynamics of interspecific hybrids

We also tracked the fitness evolution of six interspecific F2 hybrid populations, each
containing large amounts of genetic variation, across 200 mitotic generations at 5±1°C and
31±1°C. In the cold environment, three hybrid crosses significantly improved in relative fitness
(Figure 2A). As opposed to their parents, the largest improvements in hybrid fitness were seen
in the first 100 generations of experimental evolution, where both *S. cerevisiae* x *S. paradoxus*
hybrid crosses, plus the *S. cerevisiae Y55* x *S. uvarum JRY9189* hybrid, vastly exceeded both
their parents in relative fitness. In the warm environment, hybrids changed overall less in
fitness (Figure 2B). *S. cerevisiae* x *S. paradoxus* and *S. cerevisiae* x *S. kudriavzevii* hybrid crosses
did not change significantly. One of the *S. cerevisiae* x *S. uvarum* hybrids tended to lose fitness
at warm temperature (*S. cerevisiae XL011* x *S. uvarum CBS7001*).

Hybrids had low absolute starting fitness in the cold (often lower than both parents
individually) but improved over the course of evolution, often doubling or tripling in absolute
fitness (Supplementary Figure 1). This was less pronounced in the warm environment, where
the starting fitness of the hybrids was not as low as in the cold, but intermediate or similar to
that of their parents. Absolute fitness values and results of unpaired t-tests comparing
evolved lines with their respective ancestral strains are shown in Supplementary Figure 1 and
Table S2.

Comparing the thermal evolutionary potential of parents and hybrids

To directly compare the fitness of parents and hybrids after experimental evolution in the
warm and cold environment, we plotted the relative fitness gains at 31±1°C against the
relative fitness gains at 5±1°C across all populations. We found that by far the largest
improvements in fitness occurred in hybrids in the cold environment, with some populations
nearly quadrupling in fitness (Figure 4). The majority of the parental populations (grey dots)
gained fitness at both temperatures, but not to the same extent as the hybrids. Confirming
the patterns found above, hybrids showed the highest improvements at cold, particularly
both *S. cerevisiae* and *S. paradoxus* hybrid crosses (Figure 5, orange dots) and *S. cerevisiae*
Y55 x *S. uvarum JRY9189* (purple dots). Hybrid crosses between *S. cerevisiae* and *S.*
kudriavzevii fell in the centre, showing no fitness gains in either the warm or cold environment
(blue dots).

Figure 4*. Relative fitness of parental strains and their six interspecific F2 hybrids after 200*
generations of experimental evolution at 31±1°C, plotted against relative fitness after
evolution at 5±1°C. Each data point is an independently evolved line.

Effects of sudden temperature shifts on the fitness of evolved populations

296   After 200 generations of experimental evolution, we measured the fitness of all warm-
297   evolved populations (at 31±1°C) in the cold environment (5±1°C), and the fitness of cold-

evolved populations in the warm environment, akin to a reciprocal transplant experiment. The transition from warm to cold caused significant fitness decreases in six of the eight evolved parent populations (Supplementary Figure 2). The two cold-tolerant species (*S. kudriavzevii* and *S. uvarum*) were most affected, with significantly lower fitness than the warm-tolerant parents after the sudden temperature shift (ANOVA: $F_{2,11}$ = 4.80; Tukey's test: p= 0.025; Figure 5A left). One strain of the thermo-generalist *S. paradoxus* was unaffected while the other strain (*YPS744*) showed a significant fitness increase. The transition from warm to cold also significantly decreased the fitness of most hybrid crosses (Supplementary Figure 2). Overall, hybrid fitness after the temperature shift to cold was intermediate between that of the parents, with no significant differences to either cold- or warm-tolerant parents (Tukey's test vs. cold-tolerant: p= 0.23; vs. warm-tolerant p= 0.26).

The shift from cold to warm only caused significant fitness decreases in three of the eight evolved parent populations and only the two warm-tolerant species were affected (*S. cerevisiae* and *S. paradoxus*; Supplementary Figure 2). The growth of the two cold-tolerant species (*S. uvarum* and *S. kudriavzevii*) remained mostly unaffected by the sudden temperature transition to warm. A comparison between the fitness of warm-tolerant and cold-tolerant species after the shift to warm was non-significant (ANOVA: $F_{2,11}$= 3.48; Tukey's test: p= 0.06; Figure 5A right). The transition from cold to warm caused a significant fitness decrease in five of the six hybrid populations. Again, hybrid fitness was overall intermediate between that of the parents, with no significant differences to either cold- or warm-tolerant parents (Tukey's test vs. cold-tolerant: p= 0.19; vs. warm-tolerant p= 0.62).

We then tested if the original thermal performance of the strains before experimental evolution predicts how much their fitness is impacted by sudden temperature shifts. We found that the absolute fitness of ancestral hybrid strains before evolution predicted their fitness after the temperature shift when transplanting populations from warm to cold (linear regression; R^2= 0.94, p <0.01 Figure 6B left), but not when transplanting from cold to warm conditions (R^2= 0.41, p= 0.17; Figure 6B right). This was not the case for the parent populations, which showed no correlation between the ancestor's fitness and the temperature-shifted populations in either direction (R^2=0.08, p = 0.49 in cold, R^2=0.16, p = 0.32 in warm). This suggests that hybrids may have greater resilience when faced with sudden temperature shifts from warm to cold, and remain similar in fitness to their hybrid ancestor.

329
Figure 5 A) *Left: Comparing the relative fitness of warm-tolerant species, cold-tolerant species,*
and their hybrids, after evolution in the warm environment (31±1°C), when suddenly exposed
to the cold (5±1°C). Right: Comparing the relative fitness of these three groups after evolution
in the cold environment (5±1°C), when suddenly exposed to the warm (31±1°C). Whiskers show
the smallest and largest values, lines in the middle of the boxes represent the median and +
*represents the mean. Asterisk indicates significant difference in ANOVA. **B)** Linear regression*
of the absolute fitness of the evolved populations after sudden temperature shifts plotted
against the fitness of their ancestors. Asterisks indicate significant correlation coefficient.
338

Discussion
In times of global warming and extreme climate events (Perkins-Kirkpatrick & Gibson, 2017),
the effect of outbreeding on thermal adaptation is of particular interest to evolutionary
biologists, conservation managers, and agriculturalists alike. Genetic exchange in the form of
gene flow, introgression and hybridization occurs in the majority of eukaryotic taxa and may
become intensified through climate change and globalization (Chunco, 2014; McFarlane &
Pemberton, 2019). Outbreeding across large genetic distances is a conservation concern
because it can lead to offspring with poor fitness (Stelkens et al., 2015), maladaptation (Coyne
& Orr, 2004), or produce more potent pathogens (King et al., 2015). But genetic exchange

with divergent populations or species can also be a 'saving grace', particularly for endangered,
genetically eroded populations that face stressful environments and require fast rates of
adaptation to avoid extinction (Abbott et al., 2013; Brauer et al., 2023). However, predicting
the long-term fitness effects of hybridization remains a major challenge in evolutionary and
conservation biology to date.
Here, we tracked the fitness dynamics and thermal adaptation of four *Saccharomyces*
yeast species and their genetically recombined F2 hybrids across 200 generations in a cold
(5±1°C) and a warm environment (31±1°C). We found that species significantly vary in their
long-term evolutionary potential for thermal adaptation. Warm-tolerant species showed
larger fitness improvements at cold temperature (Figure 2A, Figure 3) and cold-tolerant
species showed larger fitness improvements at warm temperature (Figure 2B, Figure 3).
When comparing the adaptive potential of parent species against that of their hybrids, we
found that by far the largest relative fitness improvements occurred in hybrids, with some
populations nearly quadrupling in fitness in the cold environment, especially in the first 100
generations of experimental evolution (Figure 2A). Notably, three of the six hybrids
significantly exceeded both of their parents in their potential to adapt to growing in the cold.
Interestingly, some hybrids made from parent species with opposite thermal
preferences showed no fitness gains in either warm or cold evolution conditions (Figure 4) .
This is consistent with theory (Barton, 2001; Chevin et al., 2014; Rieseberg et al., 1999) and
empiricism (Chhina et al., 2022; Stelkens & Seehausen, 2009) suggesting that F2 hybrids with
parents with a history of divergent selection for ecologically important traits (like
thermotolerance) often display intermediate (or 'mismatched') phenotypes, because there
are fewer options for complementation or epistasis of parental alleles with opposite signs at
quantitative trait loci (which can otherwise result in transgressive traits).
Our reciprocal transplant experiment revealed that sudden transitions from warm to
cold temperatures have larger fitness costs than *vice versa*, especially for cold-tolerant
species that had been forced to adapt to warm conditions (Figure 5A). Shifts from cold to
warm only negatively affected warm-tolerant species and thermo-generalists, which had
been forced to adapt to cold conditions (Figure 5A). These results suggest that beneficial
mutations and molecular mechanisms leading to high-temperature adaptation may have
deleterious fitness effects at cold temperatures (consistent with antagonistic pleiotropy).
Alternatively, when evolving in warm environments, cold-tolerant species may accumulate
loss-of-function mutations in genes normally needed for growth at cold temperatures,
rendering them ill-adapted to their cold 'home' environment. The fact that the sudden shift
from cold to warm left the two cold-tolerant species unaffected (*S. kudriavzevii*, and *S.
uvarum*) suggests that the genes and mechanisms underlying cold adaptation do not have
detrimental effects on fitness at high temperatures.
Interestingly, hybrids showed an overall greater resilience to sudden temperature
changes than the parent species, especially from warm to cold conditions (Figure 5B). Even
after 200 generations of evolution under opposite temperature conditions, hybrids still
showed a similar performance as their ancestors: hybrids initially growing well at 5°C still grew
well at 5°C, even after long-term evolution at 31°C, and hybrids initially growing well at 31°C
still had high fitness at 31°, even after evolution at 5°C. Recombined hybrid genetic
backgrounds may allow for better (or faster) compensation of sudden environmental shifts
than parental genomes, for instance, due to the altered expression of parental genes
(Runemark et al., 2024). For instance, gene expression analysis of ortholog genes in a
domesticated hybrid yeast strain that is used for brewing lager beer (*S. pastorianus*) showed

that alleles inherited from the cold-tolerant *S. eubayanus* parent were significantly overrepresented compared to the warm-tolerant *S. cerevisiae* alleles at cold temperatures (Timouma et al., 2021). Alternatively, hybrid genomes may have had lower access to adaptive mutations over the course of experimental evolution, due to lower rates of loss of heterozygosity (LOH), which can result from gene conversion or mitotic crossover in yeast (Smukowski Heil, 2023). LOH can allow beneficial mutations, even when recessive or only partially dominant, to sweep to fixation quickly in homozygous parental genomes, but may be masked by heterozygosity in the hybrids, as recently demonstrated in yeast hybrids challenged to evolve in an environment containing a UV mimetic drug (Bautista et al., 2023). This is consistent with our finding that hybrids adapted more slowly (or to a lesser extent) than the parents to the warm environment (Figure 2B). Thus, upon temperature reversal and exposure to the cold, hybrids may be better equipped to still grow well in the cold 'ancestral' environment as their cold-adapted alleles were preserved.

Recent work on natural hybrids between climate generalists and species with more narrow climatic ranges has shown that hybrids have reduced vulnerability to environmental change due to the fitness benefits of adaptive introgression (Brauer et al., 2023). Our findings confirm that hybridization alters the fitness outcomes of long-term adaptation to extreme environments and may contribute to population resilience when challenged by sudden temperature changes. More experimental studies on hybrids with different genetic and ecological parental backgrounds are needed if we want to accurately predict the long-term adaptive outcomes of hybridization, which is equally important for the breeding of climate-resilient hybrid crops with improved (a)biotic stress resistance (Cooper & Messina, 2023; Ter Steeg et al., 2022) and the conservation of endangered, climate-sensitive populations.

Author contributions
J.P. conducted the experimental and analytical work and drafted the figures. J.P. and R.S. jointly wrote the manuscript.

Acknowledgements
We thank Chloé Haberkorn and Jennifer Molinet for their constructive comments on an earlier version of the manuscript.

Funding
This work was supported by a Swedish Research Council Project Grant (2022-03427 to R.S.) and a Knut and Alice Wallenberg Foundation Grant (2017.0163 to R.S.).

Conflict of interest statement
The authors declare no conflicts of interest.

References

Abbott, R., Albach, D., Ansell, S., Arntzen, J. W., Baird, S. J. E., Bierne, N., Boughman, J., Brelsford, A., Buerkle, C. A., Buggs, R., Butlin, R. K., Dieckmann, U., Eroukhmanoff, F., Grill, A., Cahan, S. H., Hermansen, J. S., Hewitt, G., Hudson, A. G., Jiggins, C., … Zinner, D. (2013). Hybridization and speciation. *Journal of Evolutionary Biology, 26*(2): 229-246. https://doi.org/10.1111/j.1420-9101.2012.02599.x

441 Abrams, M. B., Dubin, C. A., AlZaben, F., Bravo, J., Joubert, P. M., Weiss, C. V., & Brem, R. B.
442   (2021). Population and comparative genetics of thermotolerance divergence between yeast
443   species. *G3: Genes, Genomes, Genetics*, *11*(7). https://doi.org/10.1093/g3journal/jkab139
444

445 Ament-Velásquez, S. L., Gilchrist, C., Rêgo, A., Bendixsen, D. P., Brice, C., Grosse-Sommer, J. M.,
446   Rafati, N., & Stelkens, R. (2022). The dynamics of adaptation to stress from standing genetic
447   variation and de novo mutations. *Molecular Biology and Evolution*, *39*(11).
448   https://doi.org/10.1093/molbev/msac242
449

450 Barton, N. H. (2001). The role of hybridization in evolution. *Molecular Ecology*, *10*(3), 551–568.
451   https://doi.org/10.1046/j.1365-294X.2001.01216.x
452

453 Bautista, C., Gagnon-Arsenault, I., Utrobina, M., Fijarczyk, A., Bendixsen, D. P., Stelkens, R., &
454   Landry, C. R. (2023). Hybrid adaptation is hampered by Haldane's sieve. *bioRxiv*, 2023-12.
455   https://doi.org/10.1101/2023.12.15.571924
456

457 Belloch, C., Orlic, S., Barrio, E., & Querol, A. (2008). Fermentative stress adaptation of hybrids
458   within the Saccharomyces sensu stricto complex. *International Journal of Food*
459   *Microbiology*, *122*(1–2), 188–195. https://doi.org/10.1016/j.ijfoodmicro.2007.11.083
460

461 Bendixsen, D. P., Frazão, J. G., & Stelkens, R. (2022). Saccharomyces yeast hybrids on the rise.
462   *Yeast 39*(1-2): 40-54. https://doi.org/10.1002/yea.3684
463

464 Bendixsen, D. P., Gettle, N., Gilchrist, C., Zhang, Z., & Stelkens, R. (2021). Genomic evidence of an
465   ancient East Asian divergence event in wild Saccharomyces cerevisiae. *Genome Biology and*
466   *Evolution*, *13*(2), evab001
467

468 Borneman, A. R., & Pretorius, I. S. (2015). Genomic insights into the Saccharomyces sensu stricto
469   complex. *Genetics*, *199*(2), 281-291. https://doi.org/10.1534/genetics.114.173633
470

471 Brauer, C. J., Sandoval-Castillo, J., Gates, K., Hammer, M. P., Unmack, P. J., Bernatchez, L., &
472   Beheregaray, L. B. (2023). Natural hybridization reduces vulnerability to climate change.
473   *Nature Climate Change*, *13*(3), 282–289. https://doi.org/10.1038/s41558-022-01585-1
474

475 Brice, C., Z. Zhang, D. Bendixsen and R. Stelkens (2021). Hybridization outcomes have strong
476   genomic and environmental contingencies. *The American Naturalist 198*(3): E53-
477   E67https://doi.org/10.5061/dryad.66t1g1k21
478

479 Chevin, L. M., Decorzent, G., & Lenormand, T. (2014). Niche dimensionality and the genetics of
480   ecological speciation. *Evolution*, *68*(5), 1244–1256. https://doi.org/10.1111/evo.12346
481

482 Chhina, A. K., Thompson, K. A., & Schluter, D. (2022). Adaptive divergence and the evolution of
483   hybrid trait mismatch in threespine stickleback. *Evolution Letters*, *6*(1), 34-45.
484   https://doi.org/10.1002/evl3.264
485

486 Chunco, A. J. (2014). Hybridization in a warmer world. *Ecology and Evolution, 4*(10): 2019-2031.
487   https://doi.org/10.1002/ece3.1052

Cooper, M., & Messina, C. D. (2023). Breeding crops for drought-affected environments and
improved climate resilience. *The Plant Cell*, *35*(1), 162-186.
https://doi.org/10.1093/plcell/koac321

Coyne, J. A., & Orr, H. A. (2004). Speciation: a catalogue and critique of species concepts.
Philosophy of Biology: An Anthology, 272–292.

D'Angiolo, M., De Chiara, M., Yue, J. X., Irizar, A., Stenberg, S., Persson, K., Llored, A., Barré, B.,
Schacherer, J., Marangoni, R., Gilson, E., Warringer, J., & Liti, G. (2020). A yeast living
ancestor reveals the origin of genomic introgressions. *Nature*, *587*(7834), 420–425.
https://doi.org/10.1038/s41586-020-2889-1

Eberlein, C., Hénault, M., Fijarczyk, A., Charron, G., Bouvier, M., Kohn, L. M., Anderson, J. B., &
Landry, C. R. (2019). Hybridization is a recurrent evolutionary stimulus in wild yeast
speciation. *Nature Communications*, *10*(1), 923. https://doi.org/10.1038/s41467-019-08809-
7

Gallone, B., Steensels, J., Mertens, S., Dzialo, M. C., Gordon, J. L., Wauters, R., Theßeling, F. A.,
Bellinazzo, F., Saels, V., Herrera-Malaver, B., Prahl, T., White, C., Hutzler, M., Meußdoerffer,
F., Malcorps, P., Souffriau, B., Daenen, L., Baele, G., Maere, S., & Verstrepen, K. J. (2019).
Interspecific hybridization facilitates niche adaptation in beer yeast. *Nature Ecology and
Evolution*, *3*(11), 1562–1575. https://doi.org/10.1038/s41559-019-0997-9

Gaworecki, K. M., & Klaine, S. J. (2008). Behavioral and biochemical responses of hybrid striped
bass during and after fluoxetine exposure. *Aquatic Toxicology*, *88*(4), 207–213.
https://doi.org/10.1016/j.aquatox.2008.04.011

Gibson, B., & Liti, G. (2015). Saccharomyces pastorianus: genomic insights inspiring innovation for
industry. *Yeast*, *32*(1), 17–27. https://doi.org/10.1002/yea.3033

Grant, P. R., & Grant, B. R. (2019). Hybridization increases population variation during adaptive
radiation. *Proceedings of the National Academy of Sciences of the United States of America*,
116(46), 23216–23224. https://doi.org/10.1073/pnas.1913534116

Harris, K. B., Flynn, K. M., & Cooper, V. S. (2021). Polygenic Adaptation and Clonal Interference
Enable Sustained Diversity in Experimental Pseudomonas aeruginosa Populations. *Molecular
Biology and Evolution*, *38*(12), 5359–5375. https://doi.org/10.1093/molbev/msab248

Hoegh-Guldberg, O., & Bindi, M. (2018). Impacts of 1.5 ºC global warming on natural and human
systems 2. *Notes*, *41.*

Hutzler, M., Morrissey, J. P., Laus, A., Meussdoerffer, F., & Zarnkow, M. (2023). A new hypothesis
for the origin of the lager yeast Saccharomyces pastorianus. *FEMS Yeast Research*, *23*,
foad023. https://doi.org/10.1093/femsyr/foad023

Kellis, M., Patterson, N., Endrizzi, M., Birren, B., & Lander, E. S. (2003). Sequencing and
comparison of yeast species to identify genes and regulatory elements. *Nature*, *423*(6937),
241-254. https://doi.org/10.1038/nature01644

King, K. C., Stelkens, R. B., Webster, J. P., Smith, D. F., & Brockhurst, M. A. (2015). Hybridization in
parasites: consequences for adaptive evolution, pathogenesis, and public health in a
changing world. *PLoS pathogens*, *11*(9), e1005098.
https://doi.org/10.1371/journal.ppat.1005098

Kryazhimskiy, S., Rice, D. P., Jerison, E. R., & Desai, M. M. (2014). Global epistasis makes
adaptation predictable despite sequence-level stochasticity. *Science*, *344*(6191), 1519-1522.
DOI:10.1126/science.1250939

Leducq, J. B., Nielly-Thibault, L., Charron, G., Eberlein, C., Verta, J. P., Samani, P., ... & Landry, C. R.
(2016). Speciation driven by hybridization and chromosomal plasticity in a wild
yeast. *Nature Microbiology*, *1*(1), 1-10. https://doi.org/10.1038/nmicrobiol.2015.3

Libkind, D., Hittinger, C. T., Valério, E., Gonçalves, C., Dover, J., Johnston, M., Gonçalves, P., &
Sampaio, J. P. (2011). Microbe domestication and the identification of the wild genetic stock
of lager-brewing yeast. *Proceedings of the National Academy of Sciences of the United*
States of America, *108*(35), 14539–14544. https://doi.org/10.1073/pnas.1105430108

Lopandic, K. (2018). Saccharomyces interspecies hybrids as model organisms for studying yeast
adaptation to stressful environments. *Yeast*, *35*(1), 21–38.
https://doi.org/10.1002/yea.3294

Marcet-Houben, M., & Gabaldón, T. (2015). Beyond the whole-genome duplication: phylogenetic
evidence for an ancient interspecies hybridization in the baker's yeast lineage. *PLoS*
biology, *13*(8), e1002220. https://doi.org/10.1371/journal.pbio.1002220

McFarlane, S. E., & Pemberton, J. M. (2019). Detecting the true extent of introgression during
anthropogenic hybridization. *Trends in Ecology & Evolution*, *34*(4), 315-326.
https://doi.org/10.1016/j.tree.2018.12.013

Naseeb, S., James, S. A., Alsammar, H., Michaels, C. J., Gini, B., Nueno-Palop, C., Bond, C. J.,
McGhie, H., Roberts, I. N., & Delneri, D. (2017). Saccharomyces jurei sp. Nov., isolation and
genetic identification of a novel yeast species from Quercus robur. *International Journal of*
Systematic and Evolutionary Microbiology, *67*(6), 2046–2052.
https://doi.org/10.1099/ijsem.0.002013

Oberhofer, M., Güsewell, S., & Leuchtmann, A. (2014). Effects of natural hybrid and non-hybrid
Epichloë endophytes on the response of Hordelymus europaeus to drought stress. *New*
Phytologist, *201*(1), 242–253. https://doi.org/10.1111/nph.12496

Pérez-Torrado, R., González, S. S., Combina, M., Barrio, E., & Querol, A. (2015). Molecular and
enological characterization of a natural Saccharomyces uvarum and Saccharomyces

cerevisiae hybrid. *International Journal of Food Microbiology*, *204*, 101–110.
https://doi.org/10.1016/j.ijfoodmicro.2015.03.012
582

583 Peris, D., Arias, A., Orlić, S., Belloch, C., Pérez-Través, L., Querol, A., & Barrio, E. (2017).
584   Mitochondrial introgression suggests extensive ancestral hybridization events among
585   Saccharomyces species. *Molecular Phylogenetics and Evolution*, *108*, 49–60.
586   https://doi.org/10.1016/j.ympev.2017.02.008
587

588 Perkins-Kirkpatrick, S. E., & Gibson, P. B. (2017). Changes in regional heatwave characteristics as
589   a function of increasing global temperature. *Scientific Reports*, *7*(1), 12256.
590   https://doi.org/10.1038/s41598-017-12520-2
591

592 Rieseberg, L. H., Archer, M. A., & Wayne, R. K. (1999). Transgressive segregation, adaptation and
593   speciation. *Heredity*, *83*(4), 363-372. https://doi.org/10.1038/sj.hdy.6886170
594

595 Salvadó, Z., Arroyo-López, F. N., Guillamón, J. M., Salazar, G., Querol, A., & Barrio, E. (2011).
596   Temperature adaptation markedly determines evolution within the genus Saccharomyces.
597   *Applied and Environmental Microbiology*, *77*(7), 2292–2302.
598   https://doi.org/10.1128/AEM.01861-10
599

600 Seehausen, O. (2004). Hybridization and adaptive radiation. *Trends in ecology & evolution*, *19*(4),
601   198-207. https://doi.org/10.1016/j.tree.2004.01.003
602

603 Shen, X. X., Opulente, D. A., Kominek, J., Zhou, X., Steenwyk, J. L., Buh, K. V., ... & Rokas, A.
604   (2018). Tempo and mode of genome evolution in the budding yeast subphylum. *Cell*, *175*(6),
605   1533-1545. https://doi.org/10.1016/j.cell.2018.10.023
606

607 Smukowski Heil, C. (2023). Loss of heterozygosity and its importance in evolution. *Journal of*
608   *Molecular Evolution*, *91*(3), 369-377. https://doi.org/10.1007/s00239-022-10088-8
609

610 Steensels, J., Gallone, B., & Verstrepen, K. J. (2021). Interspecific hybridization as a driver of
611   fungal evolution and adaptation. *Nature Reviews Microbiology*, *19*(8), 485-500.
612   https://doi.org/10.1038/s41579-021-00537-4
613

614 Stelkens, R. B., Brockhurst, M. A., Hurst, G. D. D., & Greig, D. (2014). Hybridization facilitates
615   evolutionary rescue. *Evolutionary Applications*, *7*(10), 1209–1217.
616   https://doi.org/10.1111/eva.12214
617

618 Stelkens, R. B., Brockhurst, M. A., Hurst, G. D. D., Miller, E. L., & Greig, D. (2014). The effect of
619   hybrid transgression on environmental tolerance in experimental yeast crosses. *Journal of*
620   *Evolutionary Biology*, *27*(11), 2507–2519. https://doi.org/10.1111/jeb.12494
621

622 Stelkens, R. B., Schmid, C., & Seehausen, O. (2015). Hybrid breakdown in cichlid fish. *PLoS ONE*,
623   *10*(5). https://doi.org/10.1371/journal.pone.0127207
624

625 Stelkens, R., & Bendixsen, D. P. (2022). The evolutionary and ecological potential of yeast
626   hybrids. *Current Opinion in Genetics & Development*, *76*, 101958.
627   https://doi.org/10.1016/j.gde.2022.101958
628

629 Stelkens, R., & Seehausen, O. (2009). Genetic distance between species predicts novel trait
630   expression in their hybrids. *Evolution*, *63*(4), 884–897. https://doi.org/10.1111/j.1558-
631   5646.2008.00599.x
632

633 Tenaillon, O., Rodríguez-Verdugo, A., Gaut, R. L., McDonald, P., Bennett, A. F., Long, A. D., &
634   Gaut, B. S. (2012). The molecular diversity of adaptive convergence. *Science*, *335*(6067),
635   457-461. DOI:10.1126/science.1212986
636

637 Ter Steeg, E. M., Struik, P. C., Visser, R. G., & Lindhout, P. (2022). Crucial factors for the feasibility
638   of commercial hybrid breeding in food crops. *Nature Plants*, *8*(5), 463-473.
639   https://doi.org/10.1038/s41477-022-01142-w
640

641 Timouma, S., Balarezo-Cisneros, L. N., Pinto, J., De La Cerda, R., Bond, U., Schwartz, J. M., &
642   Delneri, D. (2021). Transcriptional Profile of the Industrial Hybrid Saccharomyces
643   pastorianus Reveals Temperature-Dependent Allele Expression Bias and Preferential
644   Orthologous Protein Assemblies. *Molecular Biology and Evolution*, *38*(12), 5437–5452.
645   https://doi.org/10.1093/molbev/msab282
646

647 Wein, T., & Dagan, T. (2019). The Effect of Population Bottleneck Size and Selective Regime on
648   Genetic Diversity and Evolvability in Bacteria. *Genome Biology and Evolution*, *11*(11), 3283–
649   3290. https://doi.org/10.1093/gbe/evz243
650

651 Weiss, C. V., Roop, J. I., Hackley, R. K., Chuong, J. N., Grigoriev, I. V., Arkin, A. P., ... & Brem, R. B.
652   (2018). Genetic dissection of interspecific differences in yeast thermotolerance. *Nature*
653   *genetics*, *50*(11), 1501-1504. https://doi.org/10.1038/s41588-018-0243-4
654

655 Zhang, Z., Bendixsen, D. P., Janzen, T., Nolte, A. W., Greig, D., & Stelkens, R. (2020). Recombining
656   Your Way out of Trouble: The Genetic Architecture of Hybrid Fitness under Environmental
657   Stress. *Molecular Biology and Evolution*, *37*(1), 167–182.
658   https://doi.org/10.1093/molbev/msz211
659