stFormer: a foundation model for spatial transcriptomics

Shenghao Cao[1,*], Kaiyuan Yang[1], Jiabei Cheng[1], Hong-Bin Shen[1], Xiaoyong Pan[1], Ye Yuan[1,2,3,*]

Affiliations:

[1] Institute of Image Processing and Pattern Recognition, Shanghai Jiao Tong University, and Key Laboratory of System Control and Information Processing, Ministry of Education of China, Shanghai 200240, China.

[2] Key Laboratory of Biopharmaceutical Preparation and Delivery, Chinese Academy of Sciences, Beijing 100190, China.

[3] State Key Laboratory of Biochemical Engineering, Institute of Process Engineering, Chinese Academy of Sciences, Beijing 100190, China.

[*] Corresponding author,

e-mail: cao.shenghao@sjtu.edu.cn; yyuan@ipe.ac.cn; yuanye_auto@sjtu.edu.cn

Abstract

Recent foundation models for single-cell transcriptomics data generate informative, context-aware representations for genes and cells. The spatially resolved transcriptomics (ST) data offer extra positional insights, yet not considered by these single-cell models. Here, we introduce stFormer, a transformer model tailored for ST data. It employs the cross-attention module to incorporate spatial ligand genes into the transformer encoder of single-cell transcriptomics. To address the trade-off between resolution and gene coverage across ST technologies, we propose a biased cross-attention method that enables the model to do learning with single-cell resolution on whole-transcriptome but low-resolution Visium data. We collected human Visium datasets from a public ST database and performed cell type deconvolution, generating ~4.1 million pretraining samples. After pretraining, stFormer achieved generalization capabilities across diverse tasks including masked gene imputation, cell clustering, and batch effect correction. After fine-tuning, stFormer predicted cell types, gene set partition, and ligand-receptor interactions more accurately than the single-cell foundation model, scFoundation. stFormer also revealed ligand-receptor signaling responses via in silico perturbation.

Introduction

The large language model based on the generative pretrained transformer (GPT) architecture has made groundbreaking progress in the field of natural language processing[1,2]. Recently, the transformer pretraining method was applied to building foundation models for single-cell transcriptomics data[3-5]. The foundation models generated contextual gene and cell embeddings for single cells. These embeddings could serve as input for diverse downstream applications and achieved state-of-the-art performances.

Notably, the intracellular transcription is not only regulated by gene networks, but also mediated by extracellular signals. However, the single-cell transcriptomics data do not provide information in the extracellular microenvironment. Fortunately, the spatially resolved transcriptomics (ST) technologies enable gene expression profiling in native tissue locations, while the extra positional information is not utilized by the single-cell transformer models.

Here, we developed stFormer, a transformer model adapted for ST data to integrate both intracellular and spatial contexts. There are two main innovations in the architecture design. One is adoption of the cross-attention mechanism, previously unused in single-cell models, for modeling ligand-mediated gene interactions; the other is proposal of the biased cross-attention method which enables single-cell resolution learning from cell-type deconvolved visium data, a widely available spatial resource with whole-transcriptome gene coverage.

We assembled a pretraining corpus comprising ~4.1 million spatial samples from public human Visium datasets, spanning diverse tissues, development stages, and disease states. After pretraining, we demonstrated that stFormer gained generalization capabilities, and its derived spatially aware embeddings outperformed single-cell models in both cell-level and gene-level predictions. Moreover, stFormer supports in silico perturbation of spatial ligands and receptors.

Results

stFormer architecture and pretraining

stFormer harnesses the transformer architecture to capture gene interactions for optimizing predictive accuracy of self-supervised masked learning objective. The key improvement over the previous single-cell models is that we insert a cross-attention module after the self-attention module in the transformer block to compute attentions with ligand genes from niche cells (**Fig. 1a and Methods**). In this way, stFormer integrates cell-cell communication effects via ligand secretion and reception into intracellular gene embeddings.

A variety of ST technologies are available with a trade-off between resolution and gene coverage. One of the most commonly used platforms is the 10x Genomics Visium[6]. It captures whole transcriptomes at the arranged spots in the tissue section. However, the spot diameter, i.e. the resolution, is 55 μm, which is several-fold larger than a typical cell size, so each spot may contain multiple cells. Deconvolution methods such as cell2location[7] can estimate the cell-type proportions and cell-type specific expressions at each spot, but the cell number and cell specific expressions remain unknown. To address this problem, we propose the cell-type-wise biased cross-attention method, which enables the model to do single-cell resolution learning on spot-resolution Visium data (**Fig. 1b**). The theoretical details and derivations are provided in the **Methods** section.

We collected 75 human Visium datasets comprised of 432 tissue sections from CROST[8], a comprehensive repository of ST data, and complemented them with paired single-cell transcriptomics data. Then we deconvolved these Visium datasets with their paired single-cell reference, generating ~4.1 million spatial samples at cell-type resolution. These samples represented heterogeneous tissues, development stages, and disease states (**Fig. 1c**). Finally, we pretrained stFormer on these samples through self-supervised learning, which predicted masked gene expressions based on visible genes and spatial ligand genes (**Methods**).

stFormer achieved generalization capabilities after pretraining

For model evaluation, 0.5% of pretraining samples were randomly held out as test data. Both training and test loss for the masked learning objective followed power-law decays with increasing sample size (**Fig. 2a**), indicating progressively enhanced gene imputation capability of stFormer.

We further evaluated stFormer's cell clustering performance using its cell embeddings on a myocardial Visium section[9], which was unseen in the pretraining data. The cell embeddings were produced by max-pooling all gene embeddings output from stFormer in individial cells. We found that as pretraining data scaled up, UMAP visualizations[10] of stFormer-derived cell embeddings showed progressively clearer separation, and both NMI (normalized mutual information) and ARI (adjusted Rand index) demonstrated improved alignment with cell type annotations (**Fig. 2b**). At the end of pretraining (two epochs), the resulting clusters closely matched ground-truth annotations, as evidenced by high NMI (0.941) and ARI (0.961) scores. Furthermore, we pretrained an alternative stFormer instance on a subset of Visium datasets (55 out of 75), yielding ~2.4 million spatial samples. The subset data represented a reduced biological diversity. As a result, although NMI also increased with sample size, it consistently underperformed stFormer-4.1M and exhibited inferior asymptotic performance (**Fig. 2b**).

We also evaluated the batch effect on cell embeddings produced by stFormer using a public benchmarking dataset[11,12], a single-cell RNA-seq project including 8 patients and 3 control subjects. The single-cell transcriptomics data were imported into stFormer at the setting of single-cell mode, where intercellular cross-attention module was disabled (**Methods**). UMAP visualizations revealed that compared to PCA, stFormer-derived cell embeddings improved batch mixing, as quantified by a higher ASW_{batch} score (a batch correction metric), while maintained comparable cell type separation (**Fig. 2c**).

Spatially aware cell embeddings improved cell type predictions

We appended a three-layer perceptron classifier to stFormer for cell type prediction based on cell embeddings (**Methods**). Since cell type annotations were already available through deconvolution for Visium data, there was no need to predict cell types for them. Therefore, we tested on a single-cell ST data, a CosMx SMI dataset from human pancreas tissue[13]. This dataset covered over 18,000 genes using a whole-transcriptome panel. stFormer offers an option—max number of niche cells— to define the radius of spatial niche where ligand genes are taken as input (**Methods**).

To avoid class imbalance effects, we downsampled all cell types to equal sizes. Then, the dataset was split into training (60%), validation (20%), and test (20%) sets. We fine-tuned the last two transformer layers along with the classifier on the training set, and selected the best model based on validation performance. Finally, we compared stFormer at five different settings of niche radius and scFoundation, a single-cell foundation model, on the test set.

As can be seen in **Fig. 3a**, stFormer consistently outperformed scFoundation at all settings of niche radius. The predictive accuracy of stFormer was initially lower than its single-cell mode. As niche radius increased, stFormer's accuracy improved first, peaking at max number of niche cells = 15 and surpassing the single-cell mode. As niche radius further expanded, stFormer's accuracy degraded, while stayed above the single-cell mode. Confusion matrices in **Fig. 3b** compare the predictive accuracy for each cell type. stFormer demonstrated superior predictive accuracy for more cell types compared to both its single-cell mode and scFoundation when using the optimal setting of niche radius (max number of niche cells = 15).

To visualize the spatial distribution of predictions, we excluded one field-of-view (FOV) from the dataset, and downsampled remaining data to balanced class distribution. Then, we split training (80%) and validation (20%) sets, and performed model training and selection on them respectively. Finally, we predicted cell types on the held-out FOV. stFormer (max number of niche cells = 15) outperformed scFoundation with an overall accuracy of 88.4% compared to 85.2%. Acinar.1 cells accounted for the largest proportion (63.6%) of cells in the FOV. We compared the spatial distributions of acinar.1 cell predictions from scFoundation and stFormer in **Fig. 3c**. The circled regions highlighted the acinar.1 cells correctly predicted by stFormer while misclassified by scFoundation. These areas exhibited similar spatial patterns of cell type distributions (**Fig. 3c**).

Spatially aware gene embeddings improved gene function predictions

We appended a three-layer perceptron classifier to stFormer for gene set partition based on gene embeddings (**Methods**). We took the TGF-beta signaling pathway as an example, which exhibited increased activity in the fibrotic myocardium[9]. The TGF-beta signaling pathway is initiated by the binding of TGF-beta ligands to their receptors. We adopted the GSEA[14] gene set "signaling by TGFB

family members" with ligand genes excluded, and randomly sampled an equal number of genes outside this set as negative controls. We evaluated via fivefold cross-validation on a Visium section of fibrotic myocardium[9]. The last two transformer layers and the classifier were fine-tuned on the training sets for predicting gene inclusion within the gene set, with performance assessed on the validation sets. As shown in **Fig. 4a and b**, the average ROC (receiver operating characteristic) curve and PR (precision recall) curve of stFormer both outperformed those of scFoundation, with average AUROC (area under the ROC curve) of 0.77 compared to 0.70 and average AUPRC (area under the PR curve) of 0.70 compared to 0.59.

We next evaluated stFormer on the task of predicting ligand-receptor interactions. Specifically, we concatenated the gene embeddings of ligand-receptor pairs and fed into a three-layer perceptron to classify the interacting versus non-interacting pairs (**Methods**). We adopted the ligand-receptor interaction pairs in the CellTalkDB database[15] as positive samples, and randomly selected ligand-receptor pairs not within this database as negative samples (on average each ligand only interacts with very few receptors so the vast majority of random pairs are expected to be negative). We evaluated via fivefold cross-validation on a Visium section of normal myocardium[9]. The training and validation sets were completely separated to avoid any information leakage (**Methods**). As shown in **Fig. 4c and d**, the average and median ROC (receiver operating characteristic) curves and PR (precision recall) curves of stFormer all outperformed those of scFoundation, with average/median AUROC (area under the ROC curve) of 0.68/0.83 compared to 0.59/0.58 and average/median AUPRC (area under the PR curve) of 0.80/0.79 compared to 0.76/0.75.

In silico perturbation revealed ligand-receptor signaling responses

Finally, we tested whether stFormer-derived gene embeddings captured response of cell-cell signaling via ligand-receptor interactions. It was reported that the WNT signaling pathway plays a pivotal role in cardiac remodeling[16]. Thus, we chose a Visium section from ischemic myocardium[9] and in silico downregulated the gene expressions of WNT ligands and receptors to one-third of baseline levels specifically in cardiomyocytes (**Methods**). The deleterious effects were predicted by alterations of gene embeddings as shown in **Fig. 5**. In silico downregulation of WNT ligands and receptors, whether individual or combined, had a significantly stronger effect on WNT targets than

on randomly selected genes, whereas housekeeping genes remained largely unaffected (**Fig. 5a-c**). Combined downregulation of WNT ligands and receptors demonstrated significantly enhanced effects on WNT targets compared to individual downregulation of either ligands or receptors (**Fig. 5d**).

Discussion

Advances in large language models have inspired the development of foundation models for single-cell transcriptomics data[3-5]. The spatially resolved transcriptomics (ST) technologies, as newly emerging tools for transcriptome quantification, profile gene expressions while preserving spatial position information. However, the extra positional information is not considered in existing single-cell foundation models. Building foundation models for ST data presents two main challenges. One is how to incorporate spatial information into gene expression sequence model; the other is how to accommodate diverse ST technologies, which often involve trade-offs between resolution and gene coverage.

In this study, we introduce stFormer as a solution to the aforementioned challenges. stFormer harnesses the transformer architecture[1]. In addition to the self-attention module, which is widely adopted by the existing single-cell models to encode intracellular gene interactions, stFormer innovatively employs the cross-attention module to encode ligand-mediated intercellular gene interactions. In this way, stFormer incorporates ligand gene expressions in the spatially neighboring cells into single-cell transcriptomics. Furthermore, we proposed the biased cross-attention method which enables single-cell resolution learning from cell-type resolution visium data, a widely available spatial resource with whole-transcriptome gene coverage. We assembled ~4.1 million spatial samples from public human Visium datasets. After pretrained on these samples, stFormer is compatible with both single-cell and spot resolution ST data.

We demonstrated that after pretraining stFormer achieved generalization capabilities across diverse tasks. The masked gene imputation task showcased stFormer's fundamental understanding of associations between genes. The cell clustering task revealed that pooling gene embeddings within

single cells could generate cell embeddings that encode the characteristics of cell states. We further showed that stFormer removed batch effects in the cell embeddings through an scRNA-seq benchmarking dataset. We also found that higher biological diversity in pretraining data accelerated model enhancement and raised its performance limit.

Although stFormer was pretrained on spot-resolution Visium data, it is generalizable to ST data with single-cell resolution, such as CosMx SMI data. Our results indicated that spatial ligand expression information aided in cell type prediction with an appropriately defined niche radius. We provided an example to illustrate how cell type prediction benefited from a stable spatial niche. Using the TGF-beta signaling pathway as an example, we showed that our spatially aware gene embeddings more accurately partitioned the receptor-dependent gene set. Moreover, stFormer improved the predictive accuracy of ligand-receptor interactions. It also uncovered ligand-receptor signaling responses through in silico perturbation, exemplified by the WNT signaling pathway in cardiac remodeling.

In summary, stFormer offers a flexible approach to spatial transcriptomics analysis. It learned general knowledge from pretraining ST data and transferred it to specific ST data for biological exploration. Although the current pretraining dataset was built on Visium data and remained relatively limited in scale, our model architecture was designed to be compatible with different types of ST data. As the volume of ST data in public databases continues to grow—particularly with the advancement of ST technologies that offer both single-cell resolution and high gene coverage—a larger-scale pretrained model can be developed in the future.

Fig.1 | Model schematic. a, For the spatial transcriptomics (ST) data with single-cell resolution, we
regard each cell as a centric node and the neighborhood within a preset radius as its niche. An input
sample comprises gene symbols and values of the center cell as well as ligand gene symbols and
values of the niche cells. The samples are fed into a stack of $L = 6$ identical transformer block
layers and output the embeddings of genes in the center cell. The model parameters are pretrained
by minimizing the reconstruction loss of the masked gene values. Then, the gene embeddings can
serve as input for multiple downstream tasks. **b,** For the ST data based on spot RNA sequencing, we
deconvolve the cell-type composition at each spot first. Then, adding a bias of logarithmic cell-type
proportion to each attention score in cross-attention can unify the case within the single-cell
framework in **a**. See **Methods** for details. **c,** Summary of pretraining data, totally about 4.1 million
samples at cell-type resolution. Pie charts display the proportional data distribution, categorized by
tissue origin, development stage, and disease state, respectively.

Fig.2 | Generalization performance of the pretrained model. a, Train and test loss of the

pretraining objective as the number of pretraining samples increased. MSE, mean squared error. **b,**

Evaluating clustering performance on a Visium test data using cell embeddings from models

pretrained on different data sizes and diversities. The left four plots display UMAP visualizations

of the pretrained cell embeddings, with points colored by cell types. The pretraining data sizes are

indicated above the UMAP plots, and two clustering evaluation metrics (NMI and ARI) are shown

below. The right plot displays the NMI variation with pretraining data size at two diversity levels.

c, UMAP visualizations of PCA and stFormer-derived cell embeddings for an scRNA-seq

benchmarking dataset, with points colored by cell types. stFormer removes batch effects in the data

while preserving biological variance. The batch correction metric ASW_{batch} is indicated below the

UMAP plots.

Fig.3 | Cell type prediction performance fine-tuned on a pancreas single-cell ST data. a, Predictive accuracy of stFormer at different settings of max number of niche cells, compared to its single-cell mode and scFoundation. Max number of niche cells is an option in stFormer determining the spatial niche radius. **b,** Confusion matrices comparing the predictive results from stFormer at the optimal setting of max number of niche cells, its single-cell mode, and scFoundation. **c,** The left two plots compare spatial distributions of correctly and incorrectly classified acinar.1 cells between scFoundation and stFormer. Regions with improved predictions are circled. The right plot displays cell type distributions in the circled regions, which exhibit similar spatial patterns.

Fig.4 | Gene function prediction performance fine-tuned on a myocardium Visium data. a,b,
Average ROC curves (**a**) and average PR curves (**b**) for predicting genes within gene set "signaling by TGFB family members", tested on a fibrotic myocardial tissue section. **c,d,** Average and median ROC curves (**c**), average and median PR curves (**d**) for predicting interacting ligand-receptor pairs, tested on a normal myocardial tissue section.

1
2
Fig.5 | In silico perturbation revealed ligand-receptor signaling responses. a,b,c Effect of in
silico downregulation of WNT receptors (**a**), WNT ligands (**b**), or both WNT ligands and receptors
(**c**) on gene embeddings generated from pretrained stFormer. Target genes of the WNT signaling
pathway were impacted significantly more than randomly selected genes, while housekeeping genes
were significantly less impacted. (**d**) In silico downregulation of WNT ligands and receptors
together impacted on WNT targets significantly more than separate downregulation of ligands or
receptors. These tests were performed on an ischemic myocardium Visium section. FDR-corrected
Wilcoxon p-values are shown beside.
11

Methods

Model architecture

Assumptions. Suppose that the ST data we are analyzing are resolved at the single-cell level. Otherwise, if the ST data are generated through spot RNA sequencing, such as with the 10x Genomics Visium platform, we can follow the methods described in the next subsection to convert them to single-cell scenarios. Centered on each cell in ST data, the neighborhood within a preset radius is defined as the cell's niche. For Visium platform, the niche radius is set to the spot radius. For single-cell ST data, such as CosMx SMI data, stFormer offers an option—'max number of niche cells' (m)—to define the niche radius as the maximum radius ensuring all cells have no more than m niche cells.

Layers. Schematic overview of the model architecture is illustrated in Fig. 1a. It is composed of six identical transformer blocks, each composed of a self-attention layer, cross-attention layer, and feed-forward neural network layer with the following parameters: 768 embedding dimensions, 12 attention heads per layer, and feed-forward size of 3072. Each layer includes attention computation or feed-forward neural network, residue connection, and layer normalization. stFormer can operate in single-cell mode, skipping cross-attention computation while retaining layer normalization in the cross-attention layer.

Input. The input sample consists of the gene symbols and values of the center cell, as well as the ligand gene symbols and values of the niche cells. We constructed a ligand gene library by integrating seven knowledge databases[15,17-22] and only used the ligand genes included in at least two databases. We limit the input to genes with non-zero expressions. For each gene, the symbol embedding is retrieved from a learnable lookup table with a vocabulary size of 19,264. The embeddings in the lookup table are initialized by those from the scFoundation[5] model. The gene value is transformed into a learnable value embedding through a scalar encoding module borrowed from scFoundation[5]. The final input embedding for each gene is the sum of its symbol embedding and value embedding.

Output. The transformer block first calculates self-attention among all gene embeddings within center cell, then computes cross-attention between all gene embeddings of center cell and ligand gene embeddings of niche cells (this step is skipped in single-cell mode), and finally propagates the gene embeddings through a two-layer feed-forward neural network. The model ultimately outputs gene embeddings specific to both the center cell's intracellular context and spatial niche.

Cell-type-wise biased cross-attention method on Visium data

Visium data consists of spot-wise gene expression profiles, with each spot covering multiple cells. We initially used the deconvolution method cell2location[7] with default settings to disentangle the mixed RNA measurements at each spot. Cell2location outputs the proportions of different cell types and cell-type specific expression profiles at each spot. To utilize the model architecture described in the previous subsection, we developed the cell-type-wise biased cross-attention method based on the following two assumptions.

Assumption 1. Cells of the same type within a spot approximately have the same expression profiles.

Assumption 2. The niche of each cell in a spot approximates to the cellular composition of that spot. The assumptions are reasonable since the spot size 55μm is relatively small, several-fold larger than a typical cell size, and *Assumption 1* is also adopted by most cell-type deconvolution methods. Under these two assumptions, calculating cross-attention using cell-wise ligand gene embeddings is equivalent to using cell-type-wise ligand gene embeddings with biases of logarithmic cell-type proportions, so that our model can do learning with single-cell resolution on the cell-type resolution Visium data. The details are described as follows.

Suppose that a total of n cells from t different cell types are located in a given spot, and the cell-type proportions estimated by cell2location are $p_1, p_2, \cdots, p_t$ respectively. For the sake of simplicity and without loss of generality, we suppose only one ligand gene l is detected. According to *Assumption 1*, the np_i cells of the *i*th cell type approximately have identical ligand gene expression, which is estimated by cell2location, resulting in the same ligand gene embedding, denoted as $E_i^l \in \mathbb{R}^{d_e}$ ($i = 1, 2, \cdots, t$).

For any one cell **c** in the spot, according to *Assumption 2*, the niche cells centered on cell **c** are

approximately the n cells in the spot. Denote an arbitrary gene embedding of cell $\mathbf{c}$ as $\boldsymbol{E}^c \in \mathbb{R}^{d_e}$.

Then, the cross-attention between $\boldsymbol{E}^c$ and the ligand gene embeddings of its niche cells $\boldsymbol{E}_i^l$ ($i = 1,2,\cdots,t$) can be calculated as follows.

First, the gene embeddings are projected to the query, key, and value spaces respectively using the learnable parameter matrices $W^Q \in \mathbb{R}^{d_k \times d_e}$, $W^K \in \mathbb{R}^{d_k \times d_e}$, and $W^V \in \mathbb{R}^{d_v \times d_e}$.

$$\boldsymbol{q} = W^Q \boldsymbol{E}^c$$

$$\boldsymbol{k}_i = W^K \boldsymbol{E}_i^l \quad (i = 1,2,\cdots,t)$$

$$\boldsymbol{v}_i = W^V \boldsymbol{E}_i^l \quad (i = 1,2,\cdots,t)$$

Then, the keys and values are packed together into matrices K and V, where $\mathbf{1}_{np_i}$ is an np_i-dimensional vector with all elements being 1 and the prime symbol denotes matrix transpose.

$$K = \left(\boldsymbol{k}_1 \mathbf{1}'_{np_1}, \cdots, \boldsymbol{k}_t \mathbf{1}'_{np_t}\right)$$

$$V = \left(\boldsymbol{v}_1 \mathbf{1}'_{np_1}, \cdots, \boldsymbol{v}_t \mathbf{1}'_{np_t}\right)$$

Finally, the attention function between $\boldsymbol{q}$ and K, V are computed as follows.

$$\text{Attention}(\boldsymbol{q}, K, V) = \text{softmax}\left(\frac{\boldsymbol{q}'K}{\sqrt{d_k}}\right) V'$$

$$= \text{softmax}\left(\frac{\boldsymbol{q}'\boldsymbol{k}_1 \mathbf{1}'_{np_1}}{\sqrt{d_k}}, \cdots, \frac{\boldsymbol{q}'\boldsymbol{k}_t \mathbf{1}'_{np_t}}{\sqrt{d_k}}\right) \cdot \left(\boldsymbol{v}_1 \mathbf{1}'_{np_1}, \cdots, \boldsymbol{v}_t \mathbf{1}'_{np_t}\right)'$$

$$= \text{softmax}\left(\frac{\boldsymbol{q}'\boldsymbol{k}_1}{\sqrt{d_k}} + \log(p_1), \cdots, \frac{\boldsymbol{q}'\boldsymbol{k}_t}{\sqrt{d_k}} + \log(p_t)\right) \cdot (\boldsymbol{v}_1, \cdots, \boldsymbol{v}_t)'$$

$$= \text{softmax}\left(\frac{\boldsymbol{q}'\tilde{K}}{\sqrt{d_k}} + bias\right) \tilde{V}'$$

where $\tilde{K} = (\boldsymbol{k}_1, \cdots, \boldsymbol{k}_t)$, $\tilde{V} = (\boldsymbol{v}_1, \cdots, \boldsymbol{v}_t)$ are the cell-type-wise keys and values; $bias = (\log(p_1), \cdots, \log(p_t))$ is a bias term composed of logarithmic cell-type proportions. The detailed derivatives of the above equations can be found in **Supplementary Note S1**.

Pretraining

The pretraining was accomplished using a masked learning objective. Specifically, 15% of the genes in the center cells were masked. Their value embeddings were replaced with mask embeddings and then added with the symbol embeddings to form the input embeddings. A three-layer perceptron block was added after the model architecture to predict the expression values of the masked genes according to their output gene embeddings. The pretraining objective was to minimize the mean squared error (MSE) loss between the predicted and true values of the masked genes.

We assembled a pretraining corpus by deconvoluting 432 tissue sections from 75 human Visium datasets in CROST[8], a public ST data repository. The corpus encompassed totally ~4.1 million samples across diverse tissues, development stages, and disease states. The model was pretrained with the following hyperparameters: epochs, 2; batch size, 4; gradient accumulation step, 2; learning rate 1×10^{-4}; optimizer, Adam. Pretraining lasted ~10 days on four Nvidia RTX 3090 24GB GPUs.

Downstream tasks

Clustering of cell embeddings. The cell embedding for each cell was generated by max-pooling all of its output gene embeddings. To evaluate the consistency between clustering results and annotated cell types, two metrics NMI (normalized mutual information) and ARI (adjusted Rand index) were calculated using the Python implementation of scib.metrics[23]. The evaluations were conducted on a normal myocardium tissue section (ACH003) from a myocardial infarction Visium dataset[9]. We evaluated clustering results using models pretrained on different data sizes. We also constructed an alternative pretraining corpus by deconvoluting a subset of Visium datasets (55 out of 75), resulting in reduced biological diversity and yielding ~2.4 million spatial samples. We further evaluated clustering results using models pretrained on different data sizes from this reduced corpus. We evaluated the batch effect on cell embeddings using a public benchmarking scRNA-seq dataset[11,12], including 8 patients and 3 controls. In this case, stFormer was run in single-cell mode, and batch correction was quantified by $\text{ASW}_{\text{batch}}$ score, calculated as inverse of the average silhouette width for batch clustering[23]

Cell type prediction. We attached a three-layer perceptron classifier to stFormer to perform the predictive task based on cell embeddings. The objective was to minimize the cross-entropy loss

between predicted cell type probabilities and ground truth labels. We evaluated on a single-cell resolution CosMx SMI dataset from human pancreas tissue[13]. To avoid class imbalance effects, we downsampled all cell types to equal sizes. Then, the dataset was split into training (60%), validation (20%), and test (20%) sets. We fine-tuned the last two transformer layers together with the classifier, and compared stFormer at five different niche radius settings, where max number of niche cells = 5, 10, 15, 20, and 25 respectively. For comparison, we also evaluated the predictive performance of stFormer at single-cell mode and a single-cell transcriptomics foundation model, scFoundation[5]. To visualize the spatial distribution of predictive results, we excluded one field-of-view (FOV=52) from the dataset, and downsampled remaining data to balanced class distribution. Then, we split training (80%) and validation (20%) sets, and performed model training and selection on them respectively. Finally, we predicted cell types on the held-out FOV, and compared the spatial distributions of acinar.1 cell predictions from stFormer and scFoundation.

Gene set partition for receptor-dependent gene network. We attached a three-layer perceptron classifier to stFormer to predict gene inclusion within gene set based on gene embeddings. The objective was to minimize the cross-entropy loss between predicted gene inclusion probabilities and ground truth labels. The last two transformer layers were fine-tuned together with the classifier. We evaluated on the TGF-beta receptor-dependent gene network using fivefold cross-validation, and the results were reported as AUROC and AUPRC±standard deviation. Specifically, we adopted the GSEA[14] gene set "signaling by TGFB family members" with ligand genes excluded, and randomly sampled an equal number of genes outside this set as negative controls. The evaluation samples were obtained from a fibrotic myocardium tissue section (ACH005) in the myocardial infarction Visium dataset[9]. For comparison, we also evaluated the predictive performance of scFoundation.

Ligand-receptor interaction pairs prediction. We attached a three-layer perceptron classifier to stFormer to predict the interacting ligand-receptor pairs. For a specific ligand-receptor pair, the ligand gene embedding from a single-cell sample was derived by averaging the ligand gene embeddings across all niche cells. For Visium data, under the two assumptions we propose in the subsection "Cell-type-wise biased cross-attention method on Visium data", it is equivalent to calculating a weighted average of ligand gene embeddings across cell types, using the cell-type

proportions as weights (see derivatives in **Supplementary Note S2**). We concatenated the ligand gene embedding with the receptor gene embedding output through the single-cell mode and fed into the perceptron classifier. The objective was to minimize the cross-entropy loss between the predicted interacting probabilities and ground truth labels.

We adopted the ligand-receptor interaction pairs in the CellTalkDB database[15] as positive samples (totally 3398 pairs), and randomly selected ligand-receptor pairs not within this database as negative samples. We evaluated the predictive performance using fivefold cross-validation. The training and validation sets were completely separated to avoid any information leakage. Specifically, we respectively split the ligand genes and receptor genes of the interaction pairs into five equal parts. In each of the five iterations, one fold of ligand genes and one fold of receptor genes were left out for validation, with the interaction pairs between them labeled as positive pairs. An equal number of negative pairs between them were randomly selected. The remaining four folds of ligand and receptor genes served as training data. Corresponding positive and negative pairs were generated in the same way. As a result, we ensured a complete separation of training and validation data, with no intersection of ligand genes or receptor genes between them.

The evaluation samples were obtained from a normal myocardium tissue section (ACH003) in the myocardial infarction Visium dataset[9]. For comparison, we also evaluated the performance of scFoundation. Since the input samples used by scFoundation did not include ligand genes of niche cells, the classifier input used intracellular ligand and receptor gene embeddings generated by the model.

In silico perturbation analysis of ligand-receptor signaling responses. We employed stFormer to reveal the signaling responses of ligand-receptor interaction on downstream genes via in silico perturbation. We took the WNT signaling pathway in cardiac remodeling as an example. Specifically, we reduced the expression values of WNT ligand and receptor genes to one-third of baseline levels in cardiomyocytes on an ischemic tissue section (ACH0011) in the myocardial infarction Visium dataset[9]. Since we restricted the input to genes with non-zero expressions, the downregulated ligand genes were WNT2B, WNT5B and the downregulated receptor genes were FZD4, FZD5, FZD6. We

analyzed the effects on various genes, including direct targets of the WNT signaling pathway, housekeeping genes, and 100 randomly selected genes, by calculating the cosine similarity of their output gene embeddings before and after in silico perturbation. Statistical comparisons of differential effects between gene groups were performed using the Wilcoxon rank-sum test, and the reported p-values were adjusted for multiple testing via false discovery rate (FDR) correction. The in silico perturbation was performed on ligand genes and receptor genes either individually or simultaneously.

Acknowledgments

This work was supported by the National Key R&D Program of China (2023YFF1204500), National Natural Science Foundation of China (No. 62103262).

Declaration of interests

The authors declare no competing interests.

References

Vaswani, A. Attention is all you need. *Advances in Neural Information Processing Systems* (2017).

Brown, T. B. Language models are few-shot learners. *arXiv preprint arXiv:2005.14165* (2020).

Theodoris, C. V. *et al.* Transfer learning enables predictions in network biology. *Nature* **618**, 616-624 (2023). https://doi.org/10.1038/s41586-023-06139-9

Cui, H. *et al.* scGPT: toward building a foundation model for single-cell multi-omics using generative AI. *Nat Methods* **21**, 1470-1480 (2024). https://doi.org/10.1038/s41592-024-02201-0

Hao, M. *et al.* Large-scale foundation model on single-cell transcriptomics. *Nat Methods* **21**, 1481-1491 (2024). https://doi.org/10.1038/s41592-024-02305-7

Du, J. *et al.* Advances in spatial transcriptomics and related data analysis strategies. *J Transl Med* **21**, 330 (2023). https://doi.org/10.1186/s12967-023-04150-2

Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial transcriptomics. *Nat Biotechnol* **40**, 661-671 (2022). https://doi.org/10.1038/s41587-021-01139-4

Wang, G. *et al.* CROST: a comprehensive repository of spatial transcriptomics. *Nucleic Acids Res* **52**, D882-d890 (2024). https://doi.org/10.1093/nar/gkad782

Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *Nature* **608**, 766-777 (2022). https://doi.org/10.1038/s41586-022-05060-x

McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

Li, Y. *et al.* Single-Cell Transcriptome Analysis Reveals Dynamic Cell Populations and Differential

Gene Expression Patterns in Control and Aneurysmal Human Aortic Tissue. *Circulation* **142**, 1374-1388 (2020). https://doi.org/10.1161/circulationaha.120.046528

Wen, H. *et al.* in *The Twelfth International Conference on Learning Representations*   (2024).

nanoString.

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences* **102**, 15545-15550 (2005).

Shao, X. *et al.* CellTalkDB: a manually curated database of ligand-receptor interactions in humans and mice. *Brief Bioinform* **22** (2021). https://doi.org:10.1093/bib/bbaa269

Bergmann, M. W. WNT signaling in adult cardiac hypertrophy and remodeling: lessons learned from cardiac development. *Circulation research* **107**, 1198-1208 (2010).

Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature Communications* **12**, 1088 (2021). https://doi.org:10.1038/s41467-021-21246-9

Hou, R., Denisenko, E., Ong, H. T., Ramilowski, J. A. & Forrest, A. R. R. Predicting cell-to-cell communication networks using NATMI. *Nature Communications* **11**, 5011 (2020). https://doi.org:10.1038/s41467-020-18873-z

Zhang, Y. *et al.* Cellinker: a platform of ligand-receptor interactions for intercellular communication analysis. *Bioinformatics* (2021). https://doi.org:10.1093/bioinformatics/btab036

Massenet-Regad, L. & Soumelis, V. ICELLNET v2: a versatile method for cell-cell communication analysis from human transcriptomic data. *Bioinformatics* **40** (2024). https://doi.org:10.1093/bioinformatics/btae089

Shao, X. *et al.* Knowledge-graph-based cell-cell communication inference for spatially resolved transcriptomic data with SpaTalk. *Nature Communications* **13**, 4429 (2022). https://doi.org:10.1038/s41467-022-32111-8

Troulé, K. *et al.* CellPhoneDB v5: inferring cell-cell communication from single-cell multiomics data. *arXiv preprint arXiv:2311.04567* (2023).

Luecken, M. D. *et al.* Benchmarking atlas-level data integration in single-cell genomics. *Nat Methods* **19**, 41-50 (2022). https://doi.org:10.1038/s41592-021-01336-8