1

Plant Mobile Domain protein-DNA motif modules counteract

Polycomb silencing to stabilize gene expression

Thierry Pélissier[1][#], Lucas Jarry[2][#], Margaux Olivier[1], Gabin Dajoux[1], Marie-Noëlle Pouch-Pélissier[1], Charles Courtois[1], Julie Descombin[2], Nathalie Picault[2], Guillaume Moissiard[2][*], and Olivier Mathieu[1][*]

[1] Institute of Genetics Reproduction and Development (iGReD), Université Clermont Auvergne, CNRS-UMR6293, Inserm-U1103, F-63000 Clermont-Ferrand, France.

[2] LGDP-UMR5096, CNRS, Université de Perpignan Via Domitia, F-66000 Perpignan, France.

[#] These authors contributed equally to this work

[*] Corresponding authors: E-mail: guillaume.moissiard@univ-perp.fr (G.M.); olivier.mathieu@uca.fr (O.M.)

Abstract

In plants and animals, Polycomb group (PcG) proteins are crucial for development, regulating gene expression through H3K27me3 deposition and subsequent gene silencing. While Polycomb silencing target specification is increasingly understood, it remains unclear how certain genes with apparent silencing-attracting features escape this process. Here, we show that the plant mobile domain C (PMD-C) containing proteins MAINTENANCE OF MERISTEMS (MAIN), MAIN-LIKE 1 (MAIL1) and MAIL2 oppose Polycomb silencing at numerous actively transcribed genes in Arabidopsis. Mutations in *MAIN*, *MAIL1* or *MAIL2* result in PcG-dependent ectopic H3K27me3 deposition, often associated with transcriptional repression. We show that MAIL1, which functions in concert with MAIN, and MAIL2 target distinct gene sets and associate with chromatin at specific DNA sequence motifs. We demonstrate that the integrity of these motif sequences is essential for promoting expression and antagonizing H3K27me3 deposition. Our results unveil a novel system opposing Polycomb silencing involving PMD-C protein-DNA motif modules, expanding our understanding of eukaryotic gene regulation mechanisms.

Main

Chromatin-based modulation of gene expression is mediated by enzymatic complexes that catalyze post-translational histone modifications and alter chromatin structure. The evolutionarily conserved Polycomb group (PcG) proteins act as a cellular memory system, mediating epigenetic gene repression. PcG proteins form multi-subunit complexes, primarily Polycomb Repressive Complex 1 (PRC1) and Polycomb Repressive Complex 2 (PRC2), which often work cooperatively to establish and maintain gene repression. PRC2 catalyzes the trimethylation of lysine 27 on histone H3 (H3K27me3), a hallmark generally associated with repressed chromatin, while PRC1 mediates ubiquitination on lysine 121 of H2A (H2AK121ub) in plants (or lysine 119 in animals). In the *Arabidopsis thaliana* genome, these two histone marks frequently colocalize, with PRC1-mediated H2AK121ub being required for the incorporation of H3K27me3 at most genes targeted by PcG-mediated silencing[1].

The dynamic regulation of gene expression necessitates mechanisms to counteract PcG-mediated regulation. In *Arabidopsis*, several factors have been identified that oppose PcG activity. These include several histone methyltransferases of the ARABIDOPSIS TRITHORAX (ATX) and ARABIDOPSIS TRITHORAX RELATED (ATXR) protein families, which catalyze H3K4 methylation, thereby promoting active chromatin states[2–5]. Another important group of proteins counteracting PcG-mediated silencing are H3K27me3 demethylases. In Arabidopsis, these include the Jumonji C domain-containing histone demethylases EARLY FLOWERING 6 (ELF6), RELATIVE OF EARLY FLOWERING 6 (REF6), and JMJ13[6–9]. Additionally, ANTAGONIST OF LIKE HETEROCHROMATIN PROTEIN 1 (ALP1) and ALP2 have been proposed to interfere with PcG silencing at a small subset of PcG targets by associating with PRC2 and potentially inhibiting its activity[10,11], although mutations in either *ALP* genes do not alter H3K27me3 levels.

We previously reported that two plant-specific proteins, MAINTENANCE OF MERISTEMS (MAIN) and MAIN-LIKE 1 (MAIL1), are important factors for plant development[12–14]. These proteins are characterized by a Plant Mobile Domain of type C (PMD-C) and have been implicated in the regulation of transposable elements and gene expression. However, their precise molecular functions and potential roles in epigenetic regulation remain poorly understood. Here we demonstrate that MAIN, MAIL1 and the homologous protein MAIL2

62 counteract PcG-mediated silencing at specific genomic loci by preventing ectopic H3K27me3

63 incorporation.

Results

Mutations in *NDX* and PRC core components alleviate *mail1*-associated phenotypes

Mutations in components of the MAIL1/MAIN/PP7L complex result in severe developmental

defects and transcriptional abnormalities characterized by loss of silencing at several TEs and

mis-expression of numerous protein-coding genes[12–15]. To uncover the molecular basis of

these defects, we conducted a genetic screen for suppressors of the *mail1* mutant

phenotypes. Screening individuals from an ethyl methanesulfonate (EMS) mutagenized

population of *mail1* containing the *L5-GUS* transgene[16], we identified a mutant line exhibiting

both improved plant development and suppression of L5 silencing release (Fig. 1a,b).

Compared to *mail1* plants, the suppressed plants showed faster growth and greener leaves.

The traits followed a 1/3 ratio in the F2 progeny from a backcross, indicating a single-locus

nuclear recessive causative mutation. Mapping-by-sequencing using an outcross F2

population identified a knock-out allele of *AT4G03090* (*NODULIN HOMEOBOX; NDX*) that we

named *ndx-7*, associated with a C to T mutation leading to a premature stop codon

(Supplementary Fig. 1a). Crossing *mail1* plants with an independent *ndx-4* T-DNA insertion

mutant line and introduction of an *NDX* genomic copy in the *mail1 ndx-7* background

confirmed that mutation of *NDX* alleviated *mail1*-associated developmental and silencing

defects (Supplementary Fig. 1b-d). Beyond the *L5-GUS* transgene, RT-qPCR analysis revealed

strong suppression of *mail1*-transcriptional activation of several endogenous targets in *mail1*

ndx-7 plants (Fig. 1c). Consistent with MAIL1's role in a complex with MAIN, introducing *ndx-*

7 in the *main* background also mitigated *main*-associated developmental defects and GUS

upregulation (Supplementary Fig. 1b,d).

NDX is a transcription factor recently found to directly interact with PRC1 core

components RING1A and RING1B, regulating a set of ABA-responsive genes[17,18]. Introgression

of the *mail1* mutation into a *ring1a ring1b* (*ring1a/1b*) double mutant backgound[18] resulted

in a similar reversal of *mail1*-associated phenotypic and transcription defects (Fig. 1b,c). NDX

can also indirectly interact with the transcription factor VIVIPAROUS1/ABI3-LIKE (VAL1),

which is involved in PRC1 recruitment at the *FLC* gene[19–21], through its association with

RING1A and RING1B. Crossing *val1* plants, which appear indistinguishable from WT, with

mail1 or *main* mutant lines resulted in double mutants with very severe developmental

phenotype characterized by the apparent absence of functional shoot apical meristem, and

unable to initiate post-embryonic differentiation (Supplementary Fig. 1e). This indicates that

ndx and *ring1a/1b* suppression of *mail1* is independent of *VAL1*.

To assess the genome-wide importance of *NDX* and *RING1A/1B* on *mail1*-associated

transcriptional defects, we performed RNA-seq experiments using WT, *ndx*, *ring1a/1b*, *mail1*,

mail1 ndx and *mail1 ring1a/1b* lines. To account for the variability we observed in *mail1* plant

phenotypic severity -possibly due to subtle differences in growth conditions- we used four

RNA-seq replicates to define a robust set of *mail1*-specific misregulated targets, including

differentially expressed genes and a subset of upregulated TEs (Fig. 2a and Supplementary

Fig. 2a). Gene ontology (GO) analysis revealed that 29% of *mail1*-upregulated genes (79 out

of 273) are linked to hormone and stress pathways (Supplementary Fig. 2b), consistent with

mail1 enhancing plant sensitivity to environmental cues. The expression of most misregulated

loci was unaffected in *ndx* or *ring1a/1b* mutants (Fig. 2a). Combining either *ndx* or *ring1a/1b*

mutations with *mail1* resulted in an overall suppression of TE upregulation (Fig. 2a,b).

Similarly, expression levels of *mail1*-upregulated genes tend to significantly return towards

WT levels in *mail1 ndx* and *mail1 ring1a/1b* (Fig. 2a,b). Additionally, *mail1*-associated

downregulation was suppressed for about 65% of the genes (62 to 68/95) in both mutant

combinations (Fig. 2a,b and Supplementary Fig. 2c). These results indicate that *NDX* and

RING1A/1B are required for the transcriptional misregulation induced in the absence of

MAIL1.

To determine the impact of mutations in the PRC2 core component CLF or SWN, we

generated *mail1 clf*-29 and *mail1 swn*-7 double mutants and performed RNA-seq. CLF and

SWN act largely redundantly to catalyze H3K27me3 deposition in vegetative tissues, although

certain loci appear to be preferentially targeted by one or other H3K27 methyltransferases[22].

We found that *mail1*-induced transcriptional misregulation at genes was significantly

suppressed in *mail1 clf* but not in *mail1 swn* (Supplementary Fig. 2d), although both double

mutants showed developmental defects mostly similar to those of *mail1* (Supplementary Fig.

2e). This indicates that mutation of *CLF* tends to suppress *mail1*-induced transcriptional

defects, albeit to an insufficient extent to result in an improved phenotype, most likely

because of the functional redundancy with SWN[22].

Our results show that *NDX, RING1A/1B,* and to a lesser extent *CLF*, are required for the

transcriptional misregulation induced in the absence of *MAIL1,* suggesting a potential link

between MAIL1 and PcG-mediated silencing.

MORC1 and MORC2 dampen TE activation in *mail1* suppressor backgrounds

MICRORCHIDIA (MORC) proteins are ATPases required for TE silencing and heterochromatin

condensation[23]. We recently linked silencing defects at a subset of TEs to downregulation of

MORC1 in *mail1* plants[15]. In both suppressor backgrounds, loss of TE activation correlated

with increased *MORC1* transcript accumulation relative to *mail1*, albeit limited in *mail1 ndx*

plants (Supplementary Fig. 3a). Interestingly, expression of the closely related *MORC2* gene,

also involved in TE silencing[24], showed a four to six fold induction in the presence of *ndx* or

ring1a/1b mutations (Supplementary Fig. 3a). We tested whether *MORC2* could contribute to

the reestablishment of TE silencing by introgressing the *morc2-1* mutation into the *mail1 ndx*

background and performing RNA-seq to monitor TE expression. Of the 25 *mail1*-upregulated

TEs, at least six showed dependency towards MORC2 for their repression in *mail1 ndx* plants

(Supplementary Fig. 3b,c), and we previously showed that overexpressing *MORC1* in the *mail1*

background also repressed some *mail1*-activated TEs[15] (Supplementary Fig. 3d). This is

consistent with MORC1 and MORC2 acting redundantly in TE silencing[24] and suggests that

upregulation of both *MORC1* and *MORC2* contributes to dampening TE activation in *mail1 ndx*

and *mail1 ring1a/1b*.

Gene silencing in *mail1* correlates with ectopic H3K27me3 incorporation

In light of the suppressor effect of *ndx* and *ring1a/1b* mutations and the known interaction of

NDX and RING1A/1B in PcG-mediated silencing[17,19], we explored a possible link between

mail1-associated gene misregulation and the PcG-mediated silencing diagnostic mark

H3K27me3. Chromatin immunoprecipitation sequencing (ChIP-seq) analyses revealed no

overall change in H3K27me3 levels in *mail1* when considering all Arabidopsis genes (Fig. 2c).

Likewise, the *ndx* mutation did not significantly alter H3K27me3 distribution over genes or

TEs (Fig. 2c and Supplementary Fig. 4). In *mail1,* upregulated genes showed slight loss of

H3K27me3. The addition of the *ndx* mutation had no additional effect, while H3K27me3 levels

were further reduced at these genes in *mail1 ring1a/1b* mutants (Fig. 2c and Supplementary

Fig. 5a). Despite this reduction, both *ndx* and *ring1a/1b* mutations tend to suppress *mail1*-

induced transcriptional activation of these genes (Fig. 2a), suggesting that their misregulation

in *mail1* is not directly connected to H3K27me3 variation. Strikingly, gene downregulation in

mail1 correlated with prominent H3K27me3 enrichment, which was largely reversed upon

additional *NDX* or *RING1A/1B* depletions (Fig. 2c and Supplementary Fig. 5a) that also tended

to restore expression of these genes (Fig. 2a). Many genes acquired high levels of H3K27me3

de novo in *mail1* (Fig. 2d)*,* correlating with transcriptional repression (Supplementary Fig. 5b),

and this was largely reversed in the *mail1 ndx* and *mail1 ring1a/1b* suppressed lines (Fig. 2e

and Supplementary Fig. 5c). One such gene was *MORC1*, gaining H3K27me3 in *mail1*, which

was completely lost in *mail1 ring1a/1b* (Supplementary Fig. 5c). We also found that its

neighbor gene *MORC2* was subject to PcG-mediated silencing in the WT, explaining its

upregulation in the presence of *ndx* or *ring1a/1b* mutations (Supplementary Fig. 3a,5c).

Interestingly, H3K27me3-enriched genes in *mail1* included *SPO11-1* (Supplementary Fig. 5c),

which is mostly expressed in flowers and encodes a topoisomerase-like protein essential for

meiotic recombination and fertility[25,26]. Gain of H3K27me3 was maintained in *mail1* immature

flowers, correlating with about a 70% decrease in *SPO11-1* transcript accumulation

(Supplementary Fig. 5d). Such level of *SPO11-1* downregulation was reported to significantly

affect gamete viability[26], and thus likely contributes to *mail1* hypofertility[27].

RING1A and RING1B are largely required for PRC1-mediated H2A ubiquitination[18],

playing a key role in recruiting H3K27me3 at numerous loci[1]. This is likely the case at genes

gaining H3K27me3 in *mail1* (Supplementary Table 1) as these showed concomitant

enrichment in H2AK121ub (Supplementary Fig. 6a,b). Together, these results suggest that loss

of *MAIL1* directly leads to PcG-mediated silencing at many genes.

To search for possible common features associated with the gain of H3K27me3 at

genes in *mail1*, we performed a DNA motif-based sequence analysis on the 1-kb region

surrounding the transcription start site (TSS) of these genes. Four conserved DNA motifs were

detected in 60-70% of the genes (Supplementary Fig. 6c), but only the top-scoring motif was

significantly enriched compared to the control set (Fig. 2f). We named this motif M1M (for

MAIL1/MAIN Motif), which notably extends a previously reported "down" motif partially

enriched among *mail1-* and *main*-downregulated genes[13]. Interestingly, the other motifs

included telobox and GA-repeat elements, two known polycomb response elements (PREs)

bound by transcription factors to tether PRC2 complexes to specific genes[22,28,29]. Using

previously published data[30], we found that the TELOMERE-REPEAT BINDING FACTOR (TRB)1,

TRB2 and TRB3, which bind telobox elements were loaded in wild-type plants close to the TSS

of a significant fraction (31/49) of *mail1* H3K27me3-enriched genes (Fig. 2e and

Supplementary Fig. 6d,e). Together, these results suggest that the MAIL1/MAIN complex

might promote the expression of some genes by preventing, directly or indirectly, PcG-

mediated silencing. Upon MAIL1/MAIN depletion, PRC2 recruitment, through TRB proteins or

other factors, requires PRC1 activity to be robustly established, potentially through a positive

feedback loop[1].

MAIL1/MAIN complexes are specifically targeted to M1M motifs

To explore the genomic localizations of MAIL1 and MAIN *in vivo*, we performed ChIP-seq using

transgenic lines expressing functional FLAG- or MYC-tagged versions of MAIL1 or MAIN under

their native promoters in their respective mutant backgrounds[13]. MAIL1 and MAIN largely

colocalized and were predominantly enriched at proximal promoter regions of genes (Fig. 3a-

d). Notably, they bound approximately 68% of the genes that gain H3K27me3 in *mail1*, but

showed no enrichment at genes with reduced H3K27me3 levels (Fig. 3e and Supplementary

Fig. 7a). These H3K27me3-enriched genes showed similar transcriptional downregulation and

H3K27me3 accumulation in *mail1*, *main*, and *mail1 main* mutants, consistent with MAIL1 and

MAIN acting in the same regulatory complex (Fig. 3f). Importantly, motif analysis revealed

that approximately 86% of MAIL1/MAIN-bound regions harbor the M1M motif

(Supplementary Fig. 7b), strongly suggesting a functional role for M1M in recruiting

MAIL1/MAIN to proximal promoter regions (Fig. 2e, 3e).

NDX and RING1A are enriched at MAIL1/MAIN binding regions

To further explore how mutations of NDX may suppress *mail1*, we sought to determine the

genomic localization of NDX. We generated transgenic lines expressing functional FLAG- or

MYC-tagged NDX in *ndx-7* and *mail1 ndx* mutant backgrounds (Supplementary Fig. 8a,b).

However, ChiP-seq experiments using anti-MYC or anti-FLAG antibodies were unsuccessful,

potentially due to low NDX protein abundance. Quality control analysis indicated that recently

published NDX ChIP-seq data[31] were not suitable for further use. We therefore turned to a

CUT&Tag approach, which provided good quality data only for the *ndx-7* background. Both

NDX-MYC and FLAG-NDX showed similar genomic distributions, with clear enrichment at

genes, particularly in promoter regions upstream of the TSSs, and depletion at TEs, especially

those located in pericentromeric heterochromatin (Supplementary Fig. 8c,d). This distribution is consistent with NDX belonging to a large family of transcription factors[32,33]. Furthermore, re-analysis of published RING1A ChIP-seq data[34] revealed substantial colocalization with NDX, consistent with their previously reported physical interaction[17,19] (Supplementary Fig. 8e,f). Notably, both NDX and RING1A were enriched at MAIL1/MAIN binding sites (Supplementary Fig. 8e), suggesting that MAIL1/MAIN may function to prevent the recruitment of downstream PcG-silencing components at their target genes.

MAIL2 function is essential to plant development and opposes PcG-mediated silencing at distinct loci

We previously reported that *MAIL1* and *MAIN* are Brassicaceae-specific paralogues of *MAIL2*, which is widely distributed among plant species[13]. The *MAIL2* gene appears to be under strong purifying selection[13] suggesting that it may fulfill an important function. In line with this assumption, the *mail2-1* transfer DNA mutant displayed a very severe developmental phenotype. The shoot apical meristem (SAM) was unable to initiate a proper post-embryonic program, with plants rarely developing true leaves and typically arresting growth at the two cotyledons stage (Fig. 4a and Supplementary Fig. 9a,b). Notably, we were unable to obtain homozygous mutants in the progeny segregating from a self-fertilized heterozygous *mail2* CRISPR mutant, suggesting that *mail2-1* may not be a null mutant.

The MAIL1, MAIN, and MAIL2 proteins share a highly homologous PMD-C domain[27] (Supplementary Fig. 10), which is the only identified domain in these proteins. Given the similarity in structure, we hypothesized that MAIL2 might be involved in a protection mechanism against polycomb silencing similar to the one identified for MAIL1/MAIN. To test this hypothesis, we introduced *ndx*, *clf*, or *ring1a/1b* mutations into the *mail2-1* background through crossing. The resulting mutants showed varying degrees of phenotypic suppression: *mail2-1 ndx* and *mail2-1 clf* double mutants developed a first pair of leaves before arresting growth, while *mail2-1 ring1a/1b* mutants displayed more extensive development, progressing through the rosette stage and approaching the floral transition (Fig. 4a and Supplementary Fig. 9a). These observations indicate that *ndx*, *clf*, and *ring1a/1b* mutations partially suppress the *mail2-1* developmental phenotype, mirroring their effects on *mail1* and *main* mutants. This suggests a functional similarity between MAIL2 and MAIL1/MAIN in

counteracting PcG-mediated silencing, implying that this anti-PcG silencing function may be

an ancestral feature of the PMD-C protein family.

We determined the genomic localization of MAIL2 by ChIP-seq using a transgenic line

expressing a functional, tagged version of MAIL2 (Supplementary Fig. 9c). This analysis

identified 1775 MAIL2-enriched regions across the genome, largely distinct from those bound

by MAIL1/MAIN (Supplementary Fig. 11a), yet similarly associated with elevated NDX and

RING1A occupancy (Supplementary Fig. 8e). As in the case of MAIL1 and MAIN, these regions

were predominantly associated with genes, particularly within proximal promoter regions

(1059 genes) (Fig. 4b and Supplementary Fig. 11b). These genes were significantly

overrepresented in the five deciles showing the highest expression level in the WT (Fig. 4c).

We performed DNA motif-based sequence analyses using MAIL2 peaks or the 1kb

region surrounding TSS of MAIL2-associated gene promoters and identified a conserved

sequence element, which we named MAIL2-Motif (M2M), present in nearly 95% of both the

peaks and promoter regions (Fig. 4d and Supplementary Fig. 11c-e). Intriguingly, the M2M

consensus resembles that of the M1M motif but in a palindromic configuration, suggesting

M1M may have derived from M2M (Supplementary Fig. 11f), and mirroring the evolutionary

relationship between the *MAIL1/MAIN* genes and *MAIL2*. As observed for MAIL1/MAIN-

target genes, GAA-repeats and telobox elements were also present in a substantial fraction

of MAIL2-associated gene promoters (Supplementary Fig. 11c).

To facilitate exploration of MAIL2 function, we generated hypomorphic *mail2* mutants

using artificial microRNAs (amiRNAs) targeting *MAIL2* mRNA (amiR-*mail2*) (Fig. 4a and

Supplementary Fig. 9b). While many amiR-*mail2* plants displayed severely compromised SAM

development under *in vitro* selection (Supplementary Fig. 12a) we were able to isolate two

amiR-*mail2* lines (amiR-*mail2_4* and amiR-*mail2_5*) that developed further. Both homozygous

lines showed around 2.5-fold reduction in *MAIL2* mRNA levels (Supplementary Fig. 12b) and

similar developmental phenotypes, including shorter stature, curly leaves, short stems and

hypofertility, which were partially suppressed by the introduction of *ring1a/1b* mutations

(Fig. 4a and Supplementary Fig. 12c). We confirmed that these phenotypes were caused by

MAIL2 downregulation by restoring MAIL2 function in amiR-*mail2_4* using an amiRNA-

insensitive *MAIL2* transgene (Supplementary Fig. 9b, 12d). We evaluated the impact of *MAIL2*

downregulation on transcription by performing RNA-seq of amiR-*mail2_4* and amiR-*mail2_5*

lines. The transcriptomes of the two lines were highly similar, with differentiated expressed

loci almost exclusively comprising genes, either up- (197) or down-regulated (96 genes)

(Supplementary Table 2). Downregulated genes were significantly enriched for MAIL2-

associated genes and were predominantly found in the top five expression quantiles in WT

plants (Fig. 4c,e). These genes showed no enrichment for specific biological pathway and were

distinct from those downregulated in *mail1* mutants (Supplementary Fig. 13a). Introduction

of amiR-*mail2* into the *mail1* background drastically worsened the *mail1* plant phenotype

(Supplementary Fig. 13b), indicating that MAIL1 and MAIL2 function in parallel pathways

regulating distinct gene sets. In amiR-*mail2*, H3K27me3 distribution was also affected, and a

large fraction of protein-coding genes (75 out of 102) gaining this mark were also bound by

MAIL2 (Supplementary Fig. 13c). We further found that genes showing MAIL2 binding to their

promoter and downregulation in amiR-*mail2* were associated with TRB1/2/3 in WT plants

(Fig. 4f and Supplementary Fig. 13d,e), and specifically gained H3K27me3 upon *MAIL2*

downregulation (Fig. 4e,f). We conclude from these results that MAIL2 and MAIL1 are

involved in similar pathways antagonizing PcG-silencing to promote expression of largely

distinct sets of genes

M1M and M2M are required to promote expression and oppose H3K27me3 deposition

To assess the functional relevance of M1M and M2M motifs, we employed two

complementary approaches based on transgenes. First, we used a transgenic construct in

which the promoter of the *SEP3* gene drives *GUS* expression[29]. The *SEP3* promoter contains a

telobox that allows PRC2-mediated H3K27me3 deposition through TRB1 binding, resulting in

low *GUS* expression[29]. Inserting either M1M or M2M in the *SEP3* promoter between the

telobox and the transgene's TSS significantly enhanced *GUS* expression and reduced

H3K27me3 levels (Fig. 5a). In a complementary approach, we generated transgenic lines using

the promoters of two endogenous genes: *KPI-1*, a MAIL1/MAIN target, or *HD2A,* a MAIL2

target, both of which are transcriptionally downregulated and enriched for H3K27me3 in their

respective mutant backgrounds (Fig. 5b). These promoters naturally contain teloboxes and

either M1M (KPI-1 promoter) or M2M (HD2A promoter) motifs. We independently

transformed WT plants with constructs carrying either the native promoter or mutated

version of M1M or M2M at the most conserved positions. We found that mutating either

M1M or M2M motif led to significant downregulation of *GUS* expression and increased

H3K27me3 deposition (Fig. 5c,d). These results demonstrate that the integrity of M1M and

M2M sequences is required to promote expression and antagonize H3K27me3-associated

repression.

Discussion

We previously reported that MAIL1, MAIN, and PP7L form a regulatory complex that primarily

controls gene expression in Arabidopsis[12–15], yet the molecular mechanisms remained

unclear. Here, we show that mutations in NDX or in the PRC1 components RING1A and

RING1B alleviate the developmental and molecular defects associated with MAIL1 depletion,

revealing an unexpected functional antagonism between the MAIL1/MAIN pathway and

Polycomb silencing.

NDX, a homeodomain-containing protein, interacts with RING1A/B and stabilizes

transcriptional states at euchromatic genes[17,19,35]. It was also recently implicated in

influencing siRNA accumulation and DNA methylation, predominantly at pericentromeric

TEs[31]. While altered DNA methylation can lead to ectopic H3K27me3 deposition at TEs [36,37],

we did not detect global changes in H3K27me3 patterns in *ndx* mutants (Supplementary Fig.

4). We observed reciprocal co-enrichment of NDX and RING1A/B at their binding sites

(Supplementary Fig. 8e,f), consistent with their demonstrated physical interaction[17,19].

Notably, NDX enrichment was detected at protein-coding genes rather than TEs

(Supplementary Fig. 8d), contrasting with prior reports of its localization at centromeric and

pericentromeric regions[31]. Moreover, NDX and RING1A are co-enriched at MAIL1/MAIN and

MAIL2-binding sites (Supplementary Fig. 8e), suggesting that these loci are poised to recruit

PRC2. We propose a model whereby MAIL1/MAIN and MAIL2 maintain gene expression by

limiting PcG activity at these loci. Upon depletion of MAIL1 or MAIL2, RING1A and NDX may

promote PRC2 recruitment, leading to H3K27me3 deposition and gene repression. Additional

cofactors such as TRB proteins, known PRC2 recruiters, may further contribute to this process

at specific targets.

Interestingly, although VAL1 also recruits PRC1 and PRC2 and cooperates with NDX at

FLC[19–21], *val1* mutations do not suppress *mail1* defects but instead exacerbate them

(Supplementary Fig. 1e). This suggests that different PRC1/2 recruitment pathways do not

contribute equally to MAIL1/MAIN-dependent gene regulation. The aggravated phenotype of
mail1 val1 double mutants may reflect misregulation of distinct gene subsets affected in each
background (Supplementary Fig. 14a), although genome-wide binding of VAL1[21] includes
many MAIL1/MAIN targets (Supplementary Fig. 14b), indicating possible context-dependent
cooperation.

Our findings establish MAIL1/MAIN and MAIL2 as key regulators that counteract
Polycomb-mediated repression to safeguard gene expression. Despite targeting distinct
genes via specific promoter motifs (M1M and M2M), these PMD-C proteins likely operate
through a shared molecular mechanism. Structural modeling and DNA-binding predictions
suggest that the PMD-C domains of MAIL1, MAIN and MAIL2 share a conserved concave
surface, a structural feature commonly associated with DNA-binding domains[38], that includes
residues predicted to mediate DNA contact (Supplementary Fig. 15). Moreover, the MAIL1
PMD-C domain shows structural similarity to the DUF1985 domain of the transposon-
encoded anti-silencing protein VANC[39], a known DNA-binding factor[40–42]. This raises the
possibility that PMD-C proteins may directly bind DNA, although this interaction remains to
be experimentally demonstrated. Alternatively, additional DNA- or chromatin-binding factors
may be required to guide or stabilize the interaction of PMD-C with DNA.

We previously linked TE reactivation in *mail1* and *main* mutants to reduced expression
of the silencing effector MORC1[15]. Here, we show that *MORC1* is among the genes protected
from PcG repression by MAIL1/MAIN. In *mail1 ndx* and *mail1 ring1a/b* mutants, TEs are
largely re-silenced, likely due to restored expression of MORC1 along with upregulation of its
paralog MORC2 (Fig. 2b and Supplementary Fig. 3). Many genes are upregulated in *mail1*,
possibly due to the downregulation of transcriptional repressors. However, we did not
identify any clear candidate. *MORC2* is unlikely to be involved, as its expression remains very
low in both *mail1* and WT (Supplementary Fig. 3a), and *morc2* mutants show no gene
deregulation[24]. *MORC1* is also most likely not responsible, as the genes upregulated in *mail1*
and *morc1* are largely distinct (Supplementary Fig. 14c), and *MORC1* overexpression in *mail1*
has minimal impact[15]. Alternatively, gene upregulation in *mail1* may result from changes in
higher-order chromatin architecture associated with loss of MAIL1 function[12].

MAIL1 and *MAIN* likely evolved from a duplication of *MAIL2* that occurred specifically

within the Brassicaceae lineage[13]. The sequence similarity between the M1M and M2M

motifs suggests that the palindromic M2M motif may have evolved into M1M at the same

time. *MAIL2* appears conserved at least in all eudicot plants[13], raising the possibility that this

PMD protein-DNA motif regulatory module may be conserved across a broad range of plant

species. Only a subset (about 20-40%) of M1M and M2M motifs are bound by MAIN/MAIL1

or MAIL2 (Supplementary Fig. 13f), suggesting that additional regulatory layers restrict their

binding to specific active genes. Future studies will address this point.

We previously defined four clades of PMD proteins (A1, A2, B and C). PMD-As are

encoded by retrotransposons, while PMD-B are found in protein-coding genes. MAIL1, MAIN

and MAIL2 belong to the PMD-C clade, which comprises protein synthesized from *bona fide*

host genes as well as DNA transposon-encoded proteins from various monocots and eudicots

plant species[12] (Supplementary Fig. 16 and Supplementary doc. 1). The predicted structures

of MAIL1, MAIN and MAIL2 PMD-C domains are highly similar (Supplementary Fig. 10) and, as

mentioned above, share interesting structural homology with the DUF1985 domain of the

DNA transposon-encoded VANC proteins[39], which function as an anti-silencing factor

antagonizing DNA methylation [40–42]. Recent studies indicate that H3K27me3 plays a role in TE

repression in a broad range of eukaryotes, including Arabidopsis, suggesting an ancestral role

of Polycomb silencing in controlling TE activity[43,44]. This raises the intriguing possibility that

TE-encoded PMD-C proteins also function as Polycomb anti-silencing factors, a hypothesis we

are currently testing. Such a connection between transposon biology and Polycomb silencing

regulation could provide valuable insights into the coevolution of genome defense and gene

expression control mechanisms.

Methods

Plant Material. Mutants *mail1-1* (GK-840E05)[27], *main-2* (GK-728H05)[45], *mail2-1*

(SAIL_1153_F07), *ndx-4* (WiscDsLox344A04)[35], *ring1a-2* (SAIL_393_E05)/*1b-3*

(SAIL_520_B04)[18], *clf-29* (SALK_021003)[46], *swn-7* (SALK_109121)[6] and *val1-2*

(SALK_088606C)[21] are all in the Col-0 background. The transgenic L5 line[16] was kindly provided

by H. Vaucheret (IJPB, France).

Design of the amiRNA (amiR) construct directed against *MAIL2* transcript was done using the

WMD3-Web MicroRNA Designer (http://wmd3.weigelworld.org) and the amiR-*MAIL2*

fragment cloned into the BIN-HYG-TX plasmid under the control of p35S promoter.

The *pMAIL1:gMAIL1-FLAG*, *pMAIL1:gMAIL1-MYC*, *pMAIN:gMAIN-FLAG*, *pMAIN:gMAIN-MYC*

and *SEP3pro:GUS* constructs were described previously[13,29]. For the *pMAIL2:gMAIL2-9xMYC*

construct, two genomic fragments encompassing the *MAIL2* gene (-1834 to +2448 relative to

ATG codon; A designated as +1) were PCR-amplified from Col-0 genomic DNA. PCR primers,

with several point mutations to *MAIL2* sequence, allowed to reconstitute a *MAIL2* transgene

bearing silent mutations and insensitive to amiR-*MAIL2* targeted degradation (Supplementary

Fig. 9b); recombination-based technology (NEB, NEBuilder HiFi DNA Assembly Master Mix

#2621) was then used to replace *MAIL1*-specific sequences of *pMAIL1:gMAIL1-MYC* vector by

the *MAIL2* transgene fragment. The *pNDX:gNDX* fragment was PCR amplified from Col-0

genomic DNA and region from -1631 to +5177 relative to ATG codon was cloned into the

PMDC123 vector. This *pNDX:gNDX* construct was used to produce a Nter-FLAG version of

NDX by inserting three repeats of the FLAG tag in frame with the second amino acids of the

NDX coding sequence. A Cter-9xMYC tag version was obtained through a recombination-

based approach similar to the one described for generating the *pMAIL2:gMAIL2-9xMYC*. For

KPI-1:GUS and *HD2A:GUS* transgenes, proximal gene promoter and 5'UTR amplicons of 459bp

and 357bp respectively were fused to *GUS* coding sequence and cloned into the PMDC123

vector. M1M and M2M were mutated using *pKPI-1:GUS* or *pHD2A:GUS* as matrix and

sequence modifications are described on Fig. 5. Insertion of M1M (-321 to -306 relative to

ATG) or M2M (-460 to -423 relative to ATG) in *proSEP3:GUS* was done by inverse PCR.

All transgenic constructs were introduced in plants through agroinfiltration by floral dipping[47]

and T1 transformants were isolated following adequate antibiotic selection.

Plants were grown on soil (60% relative humidity) or *in vitro* under long-day conditions (16 h

light, 8 h dark) at 23 °C.

Mutagenesis and mapping. Ethyl methanesulfonate (EMS) mutagenesis of the L5/*mail1-1*

mutant line was done as previously described[12]. To isolate suppressors of the *mail1* mutation,

a double screen was performed based on: 1/ visual amelioration of plant phenotype 2/ GUS

staining to confirm L5 transgene resilencing ; GUS assay was done by dissecting one leaf per

M2 plant candidate and leaves were incubated at 37 °C for 24 h in 400µg ml^{-1} 5-bromo-4-

chloro-3-indolyl-b-D-glucuronic acid, 10 mM EDTA, 50 mM sodium phosphate buffer pH 7.2,

0.2% triton X-100. The *ndx-7* mutant allele was identified following mapping-by-sequencing

using genomic DNA from a pool of 47 individual F2 mutant plants segregating from a cross

with Ler

RNA isolation and analyses. Total RNA was extracted from aerial tissues of 12-day-old plants

or immature flower buds using RNAzol RT (RN190, Molecular Research Center, Inc). For real

time RT-qPCR, 0.5 µg of total RNA was DNase treated and reverse transcribed using the

PrimeScript RT reagent kit with gDNA eraser (Perfect real time) (TaKaRa) in a final volume of

20 µl and one microlitre of cDNA was used for subsequent amplification using SYBR Premix Ex

Taq II (Tli RnaseH Plus) (TaKaRa) and Real-Time LightCycler 480 system (Roche) in a final

reaction volume of 15 µl. Data were normalized to a reference gene and analyzed according

to the $2^{-\Delta\Delta Ct}$ method. Means and standard errors of the mean were calculated from

independent biological samples. Primers used are listed in Supplementary Table 3.

For RNA-seq, DNase and purification were performed using RNA Clean &

Concentrator-5 kit (Zymo Research). Libraries were generated from about one µg total RNA

and sequenced as PE100 or PE150 reads on a DNBSEQ platform at Beijing Genomics Institute

(Hong Kong).

ChIP-seq. About two grams of aerial two-week-old rosettes were collected and incubated 15

min under vacuum in 25 ml of freshly prepared cross-linking buffer (10 mM phosphate buffer

pH 7, 100 mM NaCl, 0.1 M sucrose, 1% formaldehyde). 2 M glycine was added to stop cross-

link reaction (0.125 M final) and incubated an additional 5 min under vacuum. Tissues were

then washed three times with 10 mM phosphate buffer pH 7, 100 mM NaCl, 0.1 M sucrose,

pooled in 50 ml falcon tube and snap frozen in liquid nitrogen. Cross-linked tissues were

ground into fine powder by adding 7 ceramic balls to falcon tube and vortexing, and

resuspended into 25 ml of extraction buffer 1 (0.4 M Sucrose, 10 mM Tris-HCl pH 8, 10 mM

MgCl2, 5 mM β-mercaptoethanol, 0.1 mM PMSF, 1X complete EDTA-free cocktail inhibitor

(Roche)) at 4°C. Alternatively, collected tissues were ground into fine powder with liquid

nitrogen and samples were resuspended with 25 ml Hepes buffer containing 1%

formaldehyde (50 mM Hepes pH 8, 1 M sucrose, 5 mM KCl, 5 mM MgCl2, 0.6% triton X-100,

0.4 mM PMSF, 1% formaldehyde, 1X complete EDTA-free cocktail inhibitor), and incubated

for 10 min at RT. Crosslinking was stopped with 2 M glycine (0.125 M final), inverting several

times. The lysates were filtered through one layer of Miracloth and the nuclei were collected

by centrifuge at 4°C with 4000 rpm for 20 min. The nuclei were resuspended with extraction

buffer 2 (0.25 M Sucrose, 10 mM Tris-HCl pH 8, 10 mM $MgCl_2$, 1% triton X-100, 5 mM β-

mercaptoethanol, 0.1 mM PMSF, 1X complete EDTA-free cocktail inhibitor), centrifuge with

14000 g at 4°C for 10 min, and then resuspended with 400 µl extraction buffer 3 (1.7 M

Sucrose, 10 mM Tris-HCl pH 8, 0.15% Triton X-100, 2 mM $MgCl_2$, 5 mM β-mercaptoethanol,

0.1 mM PMSF, 1X complete EDTA-free cocktail inhibitor). The nuclei suspension was loaded

over a cushion of 400 µl extraction buffer 3 and microtubes were centrifugated with 14000 g

for 1 h at 4°C. For chromatin shearing using Bioruptor UCD-200 (Diagenode), nuclei were

resuspended in 250 µl of lysis buffer (50 mM Tris-HCl pH 8, 10 mM EDTA pH 8, 1% SDS, 1X

complete EDTA-free cocktail inhibitor) and 25 cycles of 25 s 'ON', 1 min 15 s 'OFF were applied

at 4°C. Following centrifugation, chromatin was split into two fractions of 80 µl and 9 vol. of

ChIP dilution buffer added (1.1% Triton X-100, 16.7 mM Tris-HCl pH 8, 1.2 mM EDTA pH 8, 167

mM NaCl, 1X complete EDTA-free cocktail inhibitor) before proceeding to preclearing and IP

as described hereafter. For Covaris fragmentation, nuclei were resuspended with 1 ml of

covaris buffer (20 mM Tris pH 8, 2 mM EDTA, 0.1% SDS, 1X complete EDTA-free cocktail

inhibitor) for 30 min at 4°C on a turning wheel. The lysate was transferred into Covaris

milliTube Glass 1 ml AFA Fiber and chromatin was mainly sheared to 200-400 bp fragments

using S220 Focused-ultrasonicator with the following conditions: for histone marks analysis,

PIP175, CBP200, DF20, T 5°C, 5 min; for MAIL2-MYCtag, PIP140, CBP200, DF10, T 5°C, 6 min.

Sheared chromatin was centrifuged with 21000 g for 10 min at 4°C and the supernatant was

split into two fractions of 440µl. After adding 0.5 volume of covaris dilution buffer (20 mM

Tris-HCL pH 8, 2 mM EDTA, 0.1% SDS, 450 mM NaCl, 3% Triton X-100, 1X complete EDTA-free

cocktail inhibitor), chromatin was precleared with 30 µl of magnetic Protein A and Protein G

Dynabeads (Invitrogen) for 1 h at 4°C on a turning wheel. Twenty µl was saved as input and

the remaining supernatant was then incubated at 4°C overnight with specific antibodies (5 µg

of H3K27me3 Ab, Diagenode C15410069 or Millipore 07-449; 10 µl H2AUb Ab, Cell Signaling

Technology ref.8240; 10 µl Myc-Tag Ab, Cell Signaling Technology ref.2276 or ref.2278; 3.5 µl

monoclonal ANTI-FLAG M2 Ab, Sigma F1804-1MG). Next, 40 µl of magnetic Protein A and

Protein G Dynabeads were added and the mixture was rotated at 4°C for 3-4h. The beads

were washed for 5 min rotating at 4°C with low salt solution twice (0.1% SDS; 1% Triton X-

100, 2 mM EDTA pH 8, 20 mM Tris-HCl pH 8, 150 mM NaCl), high salt solution (0.1% SDS; 1%

Triton X-100, 2 mM EDTA pH 8, 20 mM Tris-HCl pH8, 500 mM NaCl), LiCl solution (0.25 M LiCl,

1% IGEPAL CA630, 1% deoxycholic acid, 1 mM EDTA, 10 mM Tris-HCl pH8) and TE solution (10

mM Tris-HCl pH 8, 1 mM EDTA pH 8). A second TE wash was performed after transferring

beads to a new DNA low-bind Eppendorf microtube. The chromatin was eluted with 250 µl

elution buffer twice (1% SDS, 0.1 M NaHCO3) at 65°C for 15 min on a thermomixer with

shaking (950 rpm) and subjected to reverse crosslinking by adding 51 µl of reverse crosslink

mix (2 M NaCl, 0.1 M EDTA pH 8, 0.4 M Tris-HCl pH 8, 0.4 mg/ml proteinaseK) for 3 h at 50°C

then overnight at 65°C. Following digestion with 10 µg RNaseA for 30 min at RT°, DNA was

purified using ChIP DNA clean and concentrator kit (Zymo ZD5205). For MAIL2-MYC, five

independent IP experiments (equivalent 10 g tissues) were pooled.

For immature flower buds, three tubes of about 0.15 g material were harvested and snap-

frozen in liquid nitrogen. Following tissue disruption using Qiagen TissueLyser, samples were

resuspended in 1.8 ml Hepes Buffer containing 1% formaldehyde, pooled and further

processed as described above with covaris fragmentation at 5°C, PIP140, CBP200, DF10 for 6

min. IP was then performed using 4 µg of antibody to H3K27me3 (Diagenode, C15410069).

Libraries were generated and sequenced as PE100 reads on a DNBSEQ platform at Beijing

Genomics Institute (Hong Kong).

ChIP-qPCR. Three tubes of about 0.15 g of aerial 12-day-old rosettes were snap frozen in

liquid nitrogen and processed as described above for floral buds, except that nuclei were

resuspended in 130 µl covaris buffer and sonicated using microtube-130 AFA Fiber. Each IP

was performed using 2.1 µg of H3K27me3 antibody. (Diagenode, C15410069) and one of 15

µl purified DNA was used for real-time PCR amplification using SYBR Premix Ex Taq II (Tli

RnaseH Plus) (TaKaRa) and Real-Time LightCycler 480 system (Roche) in a final reaction

volume of 15 µl. Primers used are listed in Supplementary Table 3.

CUT&Tag. About 0.15g of WT, *ndx-7*::gNDX-9xMYC or *ndx-7*::g3xFLAG-NDX immature flower

buds was snap frozen in liquid nitrogen, grounded to a fine powder using Qiagen TissueLyser

followed by a light *in vitro* cross-linking for 2 min at RT with 1.8ml ChIP extraction buffer 1

containing 0.2% formaldehyde. Cross-linking was stopped with 122.5µl of 2M glycine rotating

for 10 min at RT. Lysate was filtrated through two layers of Miracloth and centrifugated for

20 min at 3000 g and 4°C. Nuclei were resuspended into 1ml of ChIP extraction buffer 2 and

centrifugated with 12000 g at 4°C for 10 min; this step was repeated for a total of 3 washes

until the nuclei pellet appears totally white. The resulting pellet was resuspended in 250 µl

ChIP extraction buffer 2, then loaded over a cushion of 250 µl ChIP extraction buffer 3 and

centrifugated with 16000 g for 1 h at 4°C. Nuclei were resuspended in 1 ml Nuclear Extraction

(NE) buffer (20 mM Hepes pH 7.9, 10 mM KCl , 0.1% triton X-100, 20% glycerol, 0.5 mM

Spermidine, 1X complete EDTA-free cocktail inhibitor), incubated for 10 min on ice followed

with 3 min centrifugation for 600 g at 4°C. The extract was resuspended in 500 µl NE buffer

and centrifuged 3 min for 600 g at 4°C. The nuclei fraction was finally resuspended in 500 µl

NE buffer and nuclei concentration was estimated by counting using Neubauer chamber. Each

CUT&Tag reaction was then performed on 400 000 nuclei essentially as described in the

Epicypher CUTANA CUT&Tag protocol (v1.7) and using 0.5µl monoclonal Myc-Tag Ab (Cell

Signaling Technology ref.2276) or 0.5 µl monoclonal ANTI-FLAG M2 Ab (Sigma F1804-1MG).

Libraries were amplified with 18 PCR cycles and > 75 bp DNA fragments were cleaned through

two rounds of purification using 1.3x, then 0.8x AMPure Beads (Beckman Coulter A63880).

Libraries were pooled and sequenced as PE100 reads on a DNBSEQ platform at Beijing

Genomics Institute (Poland).

RNA-seq analysis. Reads were trimmed and filtered for quality and adapter contamination

using Trim Galore (v.0.6.7, Babraham Institute) and aligned to the *Arabidopsis* reference

genome (TAIR10) genome using STAR (v.2.7.9a)[39]. Reads aligning equally well to more than

one position in the reference genome were discarded, and probable PCR duplicates were

removed using MarkDuplicates from the Picard Tools suite[49]. Sample tracks were generated

using deeptools (v.3.5.1)[50] bamCoverage with the option normalizeUsing CPM and binSize 1.

Subsequent read counting for each gene and TE was performed using featureCounts

(v.2.0.3)[51] using the Araport11 genome annotation. Annotated TEs overlapping strongly (>

80%) with an annotated TE gene were considered TE genes, and the TE annotation was

discarded. Differential expression analysis was performed using DESeq2 (v.1.36.0)[52] with

absolute fold-change ≥ 4 and Benjamini-Hochberg adjusted p-value < 0.05 were considered

differentially expressed. For DEseq2 analysis of amiR-*mail2* hypomorphic lines, criteria of

absolute fold change ≥2 and Benjamini-Hochberg adjusted p-value < 0.05 were retained. All

plots were generated using ggplot2 (v.3.4.4).

ChIP-seq analysis. Reads were trimmed and filtered using Trim Galore (v.0.6.7, Babraham Institute) or Trimmomatic (v.0.39)[53], and aligned to the TAIR10 *Arabidopsis* reference genome using bowtie2 (v.2.4.2)[54], allowing only uniquely mapped reads with perfect matches. PCR duplicates were discarded using MarkDuplicates from the Picard Tools suite or markdup from Sambamba (v.0.8.2)[55]. Sample tracks were generated using deeptools (v.3.5.1)[50] bamCoverage with the option normalizeUsing RPGC and binSize 1. Peaks calling was performed with MACS2 (v.2.2.7.1)[56] using corresponding inputs as a background and --nomodel and –extsize 200 options or MACS3 (v.3.0.0b1)[56] using --model and –call-summits, q-value < 0.01, and peaks called in the anti-MYC or anti-FLAG WT controls were removed. MAIL1/MAIN bound regions are designed as the summit of peaks plus 100bp on each side.

CUT&Tag data processing. Raw paired-end reads were trimmed using Trim Galore (v.0.6.7) with default parameters. Trimmed reads were aligned to the TAIR10 reference genome using Bowtie2 (v.2.4.2 in --very-sensitive-local mode, with insert size constraints (-I 10 -X 700), and with --no-mixed and --no-discordant options to restrict alignments to properly paired reads. Reads mapping to organellar genomes (chloroplast or mitochondria) were filtered out, and only uniquely mapping reads with a mapping quality ≥ 10 were retained. BAM files were then sorted and indexed and insert size distributions were computed using Picard's *CollectInsertSizeMetrics*. For sample normalization, exogenous *E. coli* DNA carried by the pA-Tn5 transposase was used as a spike-in control. Clean reads were aligned to the *E. coli* genome using Bowtie2, and a scaling factor was computed for each sample as the ratio between the number of *E. coli* reads in the sample and the smallest *E. coli* read count observed among all samples. These scaling factors were used to normalize read coverage when generating signal tracks (bigWig format) using *bamCoverage* of deepTools (v.3.5.1), with a bin size of 5 bp.

Motif discovery. DNA sequences of MAIL1/MAIN and MAIL2 ChIP-seq peaks and of gene proximal promoter regions (+/- 500 pb surrounding TSS) were extracted using bedtools (v.2.31.1) getfasta. Conserved motifs were identified using STREME v5.5.5[57] and the shuffled input sequences option with the following parameters: streme --verbosity 1 --oc streme_out -dna --minw 6 --maxw 16 --order 2 --bfile ./background --seed 0 --align center --time 4740 --totallength 4000000 --evalue --thresh 0.05, or using MEME v5.5.5[58] with the Classic mode

option and the following parameters: meme -oc meme_out -objfun classic -mod anr -csites

10000 -nmotifs 10 -minw 6 -maxw 15 -markov_order 0 -dna -revcomp -evt 0.05. Differentially

enriched DNA motifs were recovered using the MEME DE mode option and an equivalent

control set; relative enrichment ratio was evaluated with the SEA program of the MEME suite.

Genomic distribution of M1M and M2M identified from ChIP peaks was determined using

FIMO v5.5.5[59] with P value match < 1e-05 and default parameters. Genomic localization of

telobox elements was determined using FIMO v5.5.5[59] with p-value match < 1e-05 and default

parameters and the consensus sequence "AAACCCTAR".

Gene ontology analysis. GO term enrichment analysis was performed using ShinyGO 0.82[60]

and the *Arabidopsis thaliana* Araport11 annotation. *P*-value was derived from

hypergeometric distribution, adjusted using the FDR (false discovery rate) method. Results

with a *P*-value < 1e-05 were considered as significant.

Protein structure and DNA-binding potential prediction. The structures of the PMD-C

domains of MAIL1, MAIN and MAIL2 were modeled using AlphaFold 3 [61]. Potential DNA-

binding residues of MAIN, MAIL1 and MAIL2 were predicted using the geometry-aware

binding site predictor GPSite [62]. For each PMD-C model, prediction scores were mapped onto

the corresponding residues for visualization using UCSF ChimeraX v. 1.9 [63].

Phylogenetic analysis. Candidate MuDR-encoded PMD sequences were identified through

tBlastN analysis of Repbase database[64] using the MAIL2 PMD-C region as query

(Supplementary doc. 1). The "PhyML+SMS/OneClick" pipeline[65] (https://ngphylogeny.fr/)

with default parameters was then used to build a phylogenetic tree based on MuDR-

associated PMD and including Arabidopsis gene-encoded PMD-B and -C sequences. Resulting

tree was exported using the online Interactive Tree Of Life (ITOL) tool. PMD sequences used

to build the tree are reported in Supplementary Table 4.

Data availability. The data supporting the findings of this study are available within the article

and its Supplementary Information. High throughput sequencing data has been deposited in

the Gene Expression Omnibus (GEO) database and can be accessed with the accession

number GSE278560 (accessible using the token *ktghyqoafluttcz* during the review process).

All data are available from the corresponding author upon reasonable request.

References

1.   Zhou, Y., Romero-Campero, F. J., Gómez-Zambrano, Á., Turck, F. & Calonje, M. H2A monoubiquitination in Arabidopsis thaliana is generally independent of LHP1 and PRC2 activity. *Genome Biol.* **18**, 13–69 (2017).

2.   Chen, L.-Q. *et al.* ATX3, ATX4, and ATX5 Encode Putative H3K4 Methyltransferases and Are Critical for Plant Development. *Plant Physiol.* **174**, 1795–1806 (2017).

3.   Alvarez-Venegas, R. *et al.* ATX-1, an Arabidopsis homolog of trithorax, activates flower homeotic genes. *Curr. Biol. CB* **13**, 627–637 (2003).

4.   Tamada, Y., Yun, J.-Y., Woo, S. chul & Amasino, R. M. ARABIDOPSIS TRITHORAX-RELATED7 Is Required for Methylation of Lysine 4 of Histone H3 and for Transcriptional Activation of FLOWERING LOCUS C. *Plant Cell* **21**, 3257–3269 (2009).

5.   Berr, A. *et al.* Arabidopsis SET DOMAIN GROUP2 Is Required for H3K4 Trimethylation and Is Crucial for Both Sporophyte and Gametophyte Development. *Plant Cell* **22**, 3232–3248 (2010).

6.   Lu, F., Cui, X., Zhang, S., Jenuwein, T. & Cao, X. Arabidopsis REF6 is a histone H3 lysine 27 demethylase. *Nat. Publ. Group* **43**, 715–719 (2011).

7.   Cui, X. *et al.* REF6 recognizes a specific DNA sequence to demethylate H3K27me3 and regulate organ boundary formation in Arabidopsis. *Nat. Publ. Group* (2016).

8.   Yan, W. *et al.* Dynamic and spatial restriction of Polycomb activity by plant histone demethylases. *Nat. Plants* **4**, 681–689 (2018).

9.   Antunez-Sanchez, J. *et al.* A new role for histone demethylases in the maintenance of plant genome integrity. *eLife* **9**, (2020).

10.   Liang, S. C. *et al.* Kicking against the PRCs – A Domesticated Transposase Antagonises Silencing Mediated by Polycomb Group Proteins and Is an Accessory Component of Polycomb Repressive Complex 2. *PLoS Genet.* **11**, e1005660-26 (2015).

11.   Velanis, C. N. *et al.* The domesticated transposase ALP2 mediates formation of a novel Polycomb protein complex by direct interaction with MSI1, a core subunit of Polycomb Repressive Complex 2 (PRC2). *PLoS Genet.* **16**, e1008681 (2020).

12.   Ikeda, Y. *et al.* Arabidopsis proteins with a transposon-related domain act in gene silencing. *Nat. Commun.* **8**, 1–10 (2017).

13.   Nicolau, M. *et al.* The plant mobile domain proteins MAIN and MAIL1 interact with the phosphatase PP7L to regulate gene expression and silence transposable elements in Arabidopsis thaliana. *PLoS Genet.* **16**, e1008324 (2020).

14.   Luxán Hernández, C. *et al.* PP7L is essential for MAIL1-mediated transposable element silencing and primary root growth. *Plant J.* **8**, 15115–15122 (2020).

15.   Jarry, L. *et al.* Plant mobile domain proteins ensure Microrchidia 1 expression to fulfill transposon silencing. *Life Sci. Alliance* **6**, e202201539 (2023).

16.   Morel, J. B., Mourrain, P., Béclin, C. & Vaucheret, H. DNA methylation and chromatin structure affect transcriptional and post-transcriptional transgene silencing in Arabidopsis. *Curr. Biol. CB* **10**, 1591–1594 (2000).

17.   Zhu, Y. *et al.* The Arabidopsis noduliN homeobox factor AtNDX interacts with AtRiNG1a/b and negatively regulates abscisic acid signaling[Open]. *Plant Cell* **32**, 703–721 (2020).

18.   Li, J., Wang, Z., Hu, Y., Cao, Y. & Ma, L. Polycomb Group Proteins RING1A and
RING1B Regulate the Vegetative Phase Transition in Arabidopsis. *Front. Plant Sci.* **8**, 867
(2017).
19.   Mikulski, P. *et al.* VAL1 acts as an assembly platform co-ordinating co-
transcriptional repression and chromatin regulation at Arabidopsis FLC. *Nat. Commun.* **13**,
5542 (2022).
20.   Baile, F., Merini, W., Hidalgo, I. & Calonje, M. EAR domain-containing transcription
factors trigger PRC2-mediated chromatin marking in Arabidopsis. *Plant Cell* **33**, 2701–2715
(2021).
21.   Yuan, L. *et al.* The transcriptional repressors VAL1 and VAL2 recruit PRC2 for
genome-wide Polycomb silencing in Arabidopsis. *Nucleic Acids Res.* **49**, 98–113 (2021).
22.   Shu, J. *et al.* Genome-wide occupancy of histone H3K27 methyltransferases CURLY
LEAF and SWINGER in Arabidopsis seedlings. *Plant Direct* **3**, e00100 (2019).
23.   Moissiard, G. *et al.* MORC family ATPases required for heterochromatin
condensation and gene silencing. *Science* **336**, 1448–1451 (2012).
24.   Moissiard, G. *et al.* Transcriptional gene silencing by Arabidopsis microrchidia
homologues involves the formation of heteromers. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 7474–
7479 (2014).
25.   Stacey, N. J. *et al.* Arabidopsis SPO11-2 functions with SPO11-1 in meiotic
recombination. *Plant J.* **48**, 206–216 (2006).
26.   Xue, M. *et al.* The Number of Meiotic Double-Strand Breaks Influences Crossover
Distribution in Arabidopsis. *Plant Cell* **30**, 2628–2638 (2018).
27.   Ühlken, C., Horvath, B., Stadler, R., Sauer, N. & Weingartner, M. MAIN-LIKE1 is a
crucial factor for correct cell division and differentiation in Arabidopsis thaliana. *Plant J.* **78**,
107–120 (2014).
28.   Xiao, J. *et al.* Cis and trans determinants of epigenetic silencing by Polycomb
repressive complex 2 in Arabidopsis. *Nat. Publ. Group* **49**, 1546–1552 (2017).
29.   Zhou, Y. *et al.* Telobox motifs recruit CLF/SWN-PRC2 for H3K27me3 deposition via
TRB factors in Arabidopsis. *Nat. Publ. Group* **50**, 638 (2018).
30.   Wang, M. *et al.* Arabidopsis TRB proteins function in H3K4me3 demethylation by
recruiting JMJ14. *Nat. Commun.* **14**, 1736 (2023).
31.   Karányi, Z. *et al.* NODULIN HOMEOBOX is required for heterochromatin
homeostasis in Arabidopsis. *Nat. Commun.* **13**, 5058 (2022).
32.   Mukherjee, K., Brocchieri, L. & Bürglin, T. R. A comprehensive classification and
evolutionary analysis of plant homeobox genes. *Mol. Biol. Evol.* **26**, 2775–2794 (2009).
33.   Christensen, R. G. *et al.* Recognition models to predict DNA-binding specificities of
homeodomain proteins. *Bioinformatics* **28**, i84–i89 (2012).
34.   Su, X.-M. *et al.* ALFIN-like proteins link histone H3K4me3 to H2A ubiquitination
and coordinate diverse chromatin modifications in Arabidopsis. *Mol. Plant* **18**, 130–150
(2025).
35.   Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N. J. & Dean, C. R-Loop
Stabilization Represses Antisense Transcription at the Arabidopsis FLC Locus. *Science* **340**,
619–621 (2013).
36.   Deleris, A. *et al.* Loss of the DNA Methyltransferase MET1 Induces H3K9
Hypermethylation at PcG Target Genes and Redistribution of H3K27 Trimethylation to
Transposons in Arabidopsis thaliana. *PLoS Genet.* **8**, e1003062-13 (2012).
37.   Rougée, M. *et al.* Polycomb mutant partially suppresses DNA hypomethylation-
associated phenotypes in Arabidopsis. *Life Sci. Alliance* **4**, (2021).
38.   Barbi, M. & Paillusson, F. Chapter Seven - Protein–DNA Electrostatics: Toward a
New Paradigm for Protein Sliding. in *Advances in Protein Chemistry and Structural Biology*

(ed. Karabencheva-Christova, T.) vol. 92 253–297 (Academic Press, 2013).

39.   Borredá, C., Leduque, B., Colot, V. & Quadrana, L. Transposable element products,
functions, and regulatory networks in Arabidopsis. *bioRxiv* 2024.04.02.587720 (2024)
doi:10.1101/2024.04.02.587720.

40.   Hosaka, A. *et al.* Evolution of sequence-specific anti-silencing systems in
Arabidopsis. *Nat. Commun.* **8**, 2161 (2017).

41.   Sasaki, T. *et al.* Fast co-evolution of anti-silencing systems shapes the invasiveness of
Mu-like DNA transposons in eudicots. *EMBO J.* **41**, e110070 (2022).

42.   Sasaki, T. *et al.* Arms race between anti-silencing and RdDM in noncoding regions of
transposable elements. *EMBO Rep.* **n/a**, e56678 (2023).

43.   Hisanaga, T. *et al.* The Polycomb repressive complex 2 deposits H3K27me3 and
represses transposable elements in a broad range of eukaryotes. *Curr. Biol.* **33**, 4367-4380.e9
(2023).

44.   Hure, V. *et al.* Alternative silencing states of transposable elements in Arabidopsis
associated with H3K27me3. *Genome Biol.* **26**, 11 (2025).

45.   Wenig, U. *et al.* Identification of MAIN, a factor involved in genome stability in the
meristems of Arabidopsis thaliana. *Plant J.* **75**, 469–483 (2013).

46.   Xu, L. & Shen, W.-H. Polycomb Silencing of KNOX Genes Confines Shoot Stem
Cell Niches in Arabidopsis. *Curr. Biol.* **18**, 1966–1971 (2008).

47.   Clough, S. J. & Bent, A. F. Floral dip: a simplified method for -mediated
transformation of. *Plant J.* **16**, 735–743 (1998).

48.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinforma. Oxf. Engl.*
29, 15–21 (2013).

49.   Broad Institute. Picard Toolkit. Broad Institute, GitHub repository http://
broadinstitute.github.io/picard/. (2019).

50.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res.* **44**, W160–W165 (2016).

51.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

52.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

53.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

54.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9**, 357–359 (2012).

55.   Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba: fast
processing of NGS alignment formats. *Bioinformatics* **31**, 2032–2034 (2015).

56.   Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, 1–9
(2008).

57.   Bailey, T. L. STREME: accurate and versatile sequence motif discovery.
Bioinformatics **37**, 2834–2840 (2021).

58.   Bailey, T. L. & Elkan, C. Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. *Proc. Int. Conf. Intell. Syst. Mol. Biol.* **2**, 28–36 (1994).

59.   Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given
motif. *Bioinforma. Oxf. Engl.* **27**, 1017–1018 (2011).

60.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for
animals and plants. *Bioinforma. Oxf. Engl.* **36**, 2628–2629 (2020).

61.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500 (2024).

62.   Yuan, Q., Tian, C. & Yang, Y. Genome-scale annotation of protein binding sites via

language model and geometric deep learning. *eLife* **13**, (2024).
63.   Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein*
Sci. Publ. Protein Soc. **32**, e4792 (2023).
64.   Bao, W., Kojima, K. K. & Kohany, O. Repbase Update, a database of repetitive
elements in eukaryotic genomes. *Mob. DNA* **6**, 11 (2015).
65.   Lemoine, F. *et al.* NGPhylogeny.fr: new generation phylogenetic services for non-
specialists. *Nucleic Acids Res.* **47**, W260–W265 (2019).

Acknowledgments

We thank F. Turck (Max Planck Institute for Plant Breeding Research) for kindly providing the *SEP3pro:GUS* construct. We are also grateful to S. Marquardt (Department of Plant and Environmental Sciences, University of Copenhagen) for his help in setting up the CUT&Tag assay. Work in the Mathieu laboratory was supported by CNRS, Inserm, Université Clermont Auvergne core funding, a grant from the iSITE CAP2025 (to M.O.) and grants from the Agence Nationale de la Recherche (ANR-20-CE12-0009 & ANR-23-CE20-0012 to O.M.). Work in the Moissiard laboratory was supported by CNRS and University of Perpignan Via Domitia (UPVD) core funding, and by grants from the Agence Nationale de la Recherche (ANR-23-CE20-0012 to G.M). This study was also supported by "Laboratoires d'Excellences" (LAbEX) AGRO 2011-LABX-002 (under I-Site Muse framework) coordinated by Agropolis Foundation (ID 2101-009 to G.M), region Occitanie PhD grant to L.J, and by the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-10-LABX-0041)" and "École Universitaire de Recherche (EUR)" TULIP-GS (ANR-18-EURE-0019).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

T.P., L.J., G.M. and O.M. conceived the study. T.P., L.J., M.O., G.D., M.-N.P.-P., C.C., J.D., N.P., G.M. and O.M. conducted laboratory experiments. T.P., L.J., G.M. and O.M interpreted the data. T.P. and O.M. drafted the manuscript. T.P., L.J., G.M. and O.M. edited the manuscript. G.M and O.M. coordinated the research. The authors read and approved the final manuscript.

Additional information

Competing interests: The authors declare that they have no conflict of interest.

Fig. 1. NDX and PRC1 RING1A/1B components suppress *mail1*-induced defects. (**a**)

Histochemical staining for GUS activity in leaves from plants of the indicated genotypes. Scale bar

is 500 μm. (**b**) Developmental phenotype of three-week-old WT, *mail1*, *mail1 ndx-7*, *mail1*

ring1a/1b, *ndx-7* and *ring1a/1b* seedlings. Scale bar is 1 cm. (**c**) Expression analysis of *mail1-*

overexpressed loci by quantitative RT–PCR in the indicated genotypes. Transcript levels are

normalized to *ACT2* and further normalized to *mail1*; values represent means from three

biological replicates ± s.e.m. Asterisks mark statistically significant differences from the *mail1*

mutant (unpaired, two-sided Student's t-test, * $P < 0.05$; ** $P < 0.01$).

Fig. 2. *mail1*-associated TE and gene misregulation is dependent on NDX or RING1A/1B

function. (**a**) Heatmaps of RNA-seq expression z-score computed for *mail1* upregulated TEs

and misregulated genes across the indicated genotypes (left). Means of two biological

replicates were computed. Boxplots with the same data expressed in RPKM are shown on the

right. Center lines show the medians; box limits indicate the 25th and 75th percentiles;

whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles.

Asterisks mark statistically significant differences from the *mail1* mutant (Wilcoxon rank sum

test with continuity correction; two-sided, ** $P < 0.01$; ***$P < 0.001$; ****$P < 0.0001$). (**b**)

Genome browser tracks showing mRNA profiles (CPM) in the indicated genotypes at *mail1*-

misregulated loci. (**c**) Metagene plots of H3K27me3 accumulation (RPGC) in the indicated

genotypes at *mail1*-upregulated, *mail1*-downregulated genes compared to all Arabidopsis

genes. One of the two biological replicates is shown (**d**) H3K27me3 level variations in *mail1*

compared to WT at *mail1*-downregulated genes. (**e**) Representative genome browser views

of RNA-seq (CPM) and H3K27me3 ChIP-seq (RPGC) profiles in the indicated genotypes at

H3K27me3-enriched genes in *mail1*. MAIL1-FLAG and MAIN-FLAG associated patterns are

shown together with TRB1-FLAG data (GEO-GSM6895939 retrieved from[30]). Predicted Mail1

(M1M) and telobox motifs are indicated. (**f**) MEME prediction of a differentially enriched DNA

motif within the 1kb region surrounding TSS of H3K27me3-enriched genes in *mail1*. A control

promoter set was randomly selected from genes with similar H3K27me3 and expression

patterns in wild type plants (ranging from 0 to 3 RPKM and 0.1 to 15 RPKM, respectively).

Percentage of regions exhibiting the M1M motif and relative enrichment ratio are indicated.

Fig. 3. MAIL1 and MAIN are mainly co-targeted to M1M at gene proximal promoters. (**a**)

Heatmaps showing MAIN and MAIL1 ChIPseq signals (Log2 FC / input) over MAIL1 (n = 660)

and MAIN (n = 1064) peaks respectively. Peaks were aligned at their center position. (**b**)

Overlap between MAIL1 and MAIN peaks. (**c**) Diagram illustrating the genomic distribution of

MAIL1/MAIN common peaks (n = 626). (**d**) Metagene plot showing MAIL1 and MAIN

enrichment at MAIL1/MAIN co-bound genes (n = 670). Co-bound genes displayed

MAIL1/MAIN peak overlapping over the 200 bp region upstream TSS (proximal promoter

targeted genes) or encompassed at least 50% of the peak within the TSS - TES region (body

targeted genes). (**e**) Metaplots showing MAIL1, MAIN and M1M enrichment at promoter

region of *mail1*-H3K27me3 enriched genes (n=49; left panel). An equivalent random set of

genes is shown as control (right panel). (**f**) transcript and H3K27me3 levels, expressed in

RPKM, at *mail1*-H3K27me3 enriched genes in the indicated genotypes. Center lines show the

medians; box limits indicate the 25th and 75th percentiles; whiskers extend 1.5 times the

interquartile range from the 25th and 75th percentiles. The effect of genotype was verified

with a Wilcoxon rank sum test with continuity correction; two-sided (ns. not significant; * *P* <

0.05; ** *P* < 0.01; *** *P* < 0.001; **** *P* < 0.0001).

Fig. 4. MAIL2 mainly associates to gene promoters and protects numerous actively transcribed genes from H3K27me3-mediated PcG silencing (**a**) Photos of three-week-old seedlings of indicated genotype. Scale bar is 1 cm. (**b**) Genomic features associated with MAIL2 peaks. (**c**) Distribution of MAIL2-associated promoter genes in relation to their expression level in WT plants. A random set of the same number of genes is shown for comparison. Asterisks mark statistically significant differences from an equal representation in each decile (Fisher's exact test, P-value < 1e-04). All Arabidopsis genes were split in ten deciles based on their WT expression level (RPKM; bottom diagram). Center lines show the medians; box limits indicate the 25th and 75th percentiles; whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles. (**d**) Identification of a highly enriched motif within the 1kb region surrounding TSS of genes harboring MAIL2 peak at proximal promoter (n = 1059). Analysis was carried out using MEME in differential enrichment mode and an equivalent control set randomly selected among genes ranging in the five deciles with the highest expression level in the WT. Percentage of regions exhibiting the M2M motif and relative enrichment ratio are indicated. (**e**) Overlap between MAIL2-associated promoter

genes and misregulated genes in amiR-*mail2* lines. Fold change enrichment compared to

random distribution and p-value from hypergeometric tests are indicated. (**f**) Representative

genome browser views of RNA-seq (CPM) and H3K27me3 ChIP-seq (RPGC) profiles in the

indicated genotypes at MAIL2-associated promoter genes downregulated in amiR-*mail2*

background. MAIL2-MYC associated patterns are shown together with TRB1-FLAG data (GEO-

GSM6895939 retrieved from[30]) and predicted MAIL2 (M2M) and telobox motif positions. (**g**)

Box plots showing transcript and H3K27me3 levels (RPKM) in WT and amiR-*mail2* lines at

MAIL2-associated promoter genes downregulated (n = 39) in hypomorphic amiR-*mail2*

background. Center lines show the medians; box limits indicate the 25th and 75th percentiles;

whiskers extend 1.5 times the interquartile range from the 25th and 75th percentiles. The

effect of genotype was verified with a Wilcoxon rank sum test with continuity correction; two-

sided (ns. not significant; *** $P < 0.001$; **** $P < 0.0001$).

Fig. 5. Promoter-associated M1M and M2M are functionally relevant to safeguard genes from PcG-mediated silencing. (a) Schematic representation of the *proSEP3:GUS* original transgene and derivatives following insertion of one M1 or two M2 motifs. Box plots showing relative GUS expression and H3K27me3 accumulation associated to different transgene combinations. For each transgene, data represent two technical replicates on 4 independent transformed lines. Quantitative RT-PCR analysis of *GUS* expression was normalized to *ACT2*. H3K27me3 ChIP-qPCR is reported as % input. Asterisks mark statistically significant differences from *pSEP3:GUS* transgene (unpaired, two-sided Student's t-test, * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$). (**b**) Genome browser views of RNA-seq (CPM) and H3K27me3 ChIP-seq (RPGC) profiles at *KPI-1* and *HD2A* loci in the indicated genotypes. MAIL1 and MAIL2-associated patterns are shown together with TRB1-FLAG data (GEO-GSM6895939

retrieved from[30]) and predicted M1M, M2M and telobox motifs. (**c**) Schematic representation

of *pPKL1:GUS* transgenes. Most conserved M1M motifs were mutated (m1m) and sequence

modifications are shown under the scheme. Telobox motif is also reported in blue. Relative

GUS expression and H3K27me3 accumulation were performed as described in (a) except that

nine independent lines were analysed for each transgene. (**d**) Schematic representation of

pHD2A:GUS transgenes showing localization of intact (M2M) and mutated (m2m) motifs.

Telobox motifs are indicated in blue and sequence modifications are reported under the

scheme. Molecular analyses were performed as described in (a) except that nine independent

lines were analysed for each transgene. In box plots, center lines show the medians; box limits

indicate the 25th and 75th percentiles; whiskers extend 1.5 times the interquartile range from

the 25th and 75th percentiles.

Supplementary Table 1. List of H3K27me3-enriched genes (Log2 FC > 1; *P*-value < 0.01) in

***mail1* compared to WT background.**

Supplementary Table 2. Lists of differentially expressed loci in amiR-*mail2*_4 and _5 mutant

lines.

Supplementary Table 3. List of qPCR primers used in this study.

Supplementary Table 4. List of PMD sequences used for phylogenetic analyses.

Supplementary Table 5. DNA-binding prediction scores for amino acid residues of MAIN,

MAIL1 and MAIL2.

Supplementary doc. 1. Best results of TBLASTN search on Repbase[64] using MAIL2 PMD-C

domain as query. Only results with e-values < 1e-10 were reported. Dicot species: ArHy

Arachis hypogaea, AvMa *Avicennia marina*, Cba *Capsicum baccatum*, Daca *Daucus carota*, Gar

Gossypium arboretum, GR *Gossypium raimondii*, GM *Glycine max*, Ipa *Ilex paraguariensis*, LU

Linum usitatissimum, MT *Medicago truncatula*, NS *Nicotiana sylvestris*, SeTo *Senna tora*, STu

Solanum tuberosum, VV *Vitis Vinifera*. Monocot species: DRot *Dioscorea cayennensis subsp.*

Rotundata, OS *oryza Sativa*, SBi *Sorghum bicolor*, SHS *Saccharum hybrid cultivar*, Sit *Setaria*

italica, Tae *Triticum aestivum*, ZM *Zea mays*. Basal angiosperm species: ATr *Amborella*

trichopoda.