Title: Early adipose tissue wasting in a novel preclinical model of human lung cancer cachexia

Author List: Deena B. Snoke[1,4], Jos L. van der Velden[2,4], Emma R. Bellafleur[1], Jacob S. Dearborn[3,4], Sean M. Lenahan[2,4], Alexandra E. Beal[1], Reem Aboushousha[2], Skyler C. J. Heininger[1], Jennifer L. Ather[1], Madeleine M. Mank[1], Hailey Sarausky[2,4], Daniel Stephenson[5], Julie A. Reisz[5], Angelo D'Alessandro[5], Devdoot Majumdar[3,4], Thomas P. Ahern[3,4], Kaiwen Xu[6], Kim L. Sandler[8], Bennett A. Landman[6,7,8], Yvonne M.W. Janssen-Heininger[2,4], Matthew E. Poynter[1], David J. Seward[2,4], and Michael J. Toth[1,4,9]

Affiliations: Departments of [1]Medicine, [2]Pathology and Laboratory Medicine, and [3]Surgery, and [4]University of Vermont Cancer Center, University of Vermont College of Medicine, Burlington, VT; [5]Department of Biochemistry and Molecular Genetics, University of Colorado Anschutz Medical Campus, Aurora, CO, and Departments of [6]Computer Science, [7]Biomedical Engineering and [8]Radiology, Vanderbilt University and Vanderbilt University Medical Center, Nashville, TN, USA. [9]Lead contact

Corresponding Author: Michael J. Toth, Ph.D.
Health Science Research Facility 126B
149 Beaumont Avenue
Burlington VT, 05405
Email : Michael.Toth@med.uvm.edu
Tele : (802)-656-7989
Fax : (802)-656-0747

Keywords: Lung neoplasms, lipolysis, skeletal muscle, organoids, inflammation

Summary:

Cancer cachexia (CC), a syndrome of skeletal muscle and adipose wasting, reduces

responsiveness to therapies and increases mortality. There are no approved treatments for CC,

which may relate to discordance between pre-clinical models and human CC. To address the

need for clinically relevant models of lung CC, we generated inducible, lung epithelial cell

specific $Kras^{G12D/+}$ ($G12D$) mice. $G12D$ mice develop CC over a protracted time course and

phenocopy tissue and tumor, cellular, mutational, transcriptomic, and metabolic characteristics

of human lung CC. $G12D$ mice demonstrate early loss of adipose, a phenotype that was

apparent across numerous models of CC and translates to patients with lung cancer. Tumor-

released factors promote adipocyte lipolysis, a driver of adipose wasting in CC, and adipose

wasting was inversely related to tumor burden. Thus, $G12D$ mice model key features of human

lung CC and highlight a role for early tumor metabolic reprogramming of adipose tissue in CC.

Manuscript Body

Introduction:

Cancer cachexia (CC) is characterized by unintentional loss of body weight [1] and

contributes to poor response to treatments and related toxicities and increased morbidity and

mortality [1-11]. Animal models have served as the basis for knowledge of the pathophysiology of

CC [12] because the syndrome typically occurs in aggressive cancers and later-stage disease,

where patients are unwilling or unable to participate in research. While these models have

generated a wealth of data [12], there are no effective, approved treatments for CC. This lack of

translating evidence into treatments may be explained, in part, by the fact that current models

do not mimic human CC [13-21].

Lung cancer is the second most common cancer and the leading cause of cancer-related

death [22, 23]. Most patients with lung cancer exhibit signs of CC [24, 25], and CC reduces mean

survival time and responsiveness to chemo- and immunotherapies [11, 26]. Most preclinical data on

lung CC are derived from the murine Lewis lung carcinoma (LLC) model, in which tumor cells

are injected subcutaneously and grow rapidly. Weight loss begins when the tumor reaches ~5%

of body weight, and mice progress to >10% body weight loss in 1-3 weeks [27]. Neither tumor

growth, location, or size, nor weight loss characteristics of this model reflect lung CC in humans

[28]. Thus, the field would benefit from new models of lung CC that better emulate human lung

CC.

To address this need, we developed mice with an activatable lung epithelial cell-specific

$Kras^{G12D}$ allele ($G12D$) to evaluate its utility as a model of lung CC. We report that $G12D$ mice

display classic adipose and skeletal muscle tissue wasting phenotypes of CC over a protracted

time course (~12 weeks) and phenocopy tumor and tissue cellular, mutational, transcriptomic,

and metabolic characteristics relevant to human lung CC. Additionally, adipose tissue wasting

drove CC initiation in both singular $Kras^{G12D}$ mutation and when combined with other driver

mutations (*e.g., Stk11*$^{-/-}$), was inversely related to tumor burden, and could be incited by tumor

released factors. In a large, prospective cohort study, we confirmed the clinical relevance of

early adipose tissue wasting. Thus, early tumor metabolic reprogramming of adipose tissue may

be important for CC initiation and tumor biology.

Results:

Lung epithelial cell (club cell)-specific Kras^{G12D} induction causes CC.

Induction of the *Kras^{G12D}* allele with tamoxifen administration (**Fig. 1A**) promotes synchronous pulmonary lesions originating in the terminal bronchioles, the predominant anatomic location of club cells (**Fig. 1B**). Lesions exist along the histological spectrum of epithelial proliferation, ranging from florid bronchial hyperplasia to adenoma to atypical adenomatous hyperplasia to adenocarcinoma, with no evidence for dissemination to major organs or lymph nodes. Lung adenocarcinomas were confirmed histologically at the initiation of weight loss (6 weeks post-induction; Fig. 1C) within the spectrum of synchronous pulmonary lesions characterized by marked cellular atypia, architecturally distorted glands, the absence of cilia, and enhanced mitotic index (**Fig. S1A**). In fact, the histologic hallmarks of adenocarcinoma were present, but not as pronounced, as early as 3.5 weeks post-induction (**Fig. S1B**). Next, we examined whether CCSP promoter activity was evident only upon injection of tamoxifen, as non-tamoxifen-inducible (*i.e.,* leaky) Cre expression from the CCSP promoter has been observed [29]. In line with these reports, in *G12D* mice not receiving tamoxifen, we found histological evidence for lung lesion development (**Fig. S2**), albeit much less severe than in mice with tamoxifen injection. Because of this, we used wild-type, *Scgb1a1-CreER^{TM}* littermates (WT; *Kras$^{+/+}$*) injected with tamoxifen as controls, which has the advantage of controlling for toxicity and off-target effects of intracellular Cre activation and tamoxifen, respectively. Finally, tumor growth can affect the function of the organ where it resides, promoting weight loss independent of the tumor. In *G12D* mice, while animals demonstrate labored breathing and reduced activity level at advanced stage CC (weeks 10-12), they show no evidence of labored breathing or reduced activity at the initiation of body weight loss at 6 weeks (Fig. 1C). To evaluate whether there is evidence for impaired lung function that may drive CC, we measured blood oxygen saturation at 3 weeks post-induction, when body weight begins to diverge

between groups (Fig. 1C). We found no evidence for hypoxemia (**Fig. S3**), suggesting that

respiratory dysfunction in *G12D* mice is unlikely to contribute to CC.

G12D mice began losing weight 6 weeks post-induction (**Fig. 1C**), independent of

differences in food intake or fecal energy content (**Fig. S4A-B**). By 12 weeks, *G12D* mice

experienced an average body weight loss of 15%, with most animals (83%) displaying CC

(>10% body weight loss). Weight loss in *G12D* mice was explained by loss of skeletal muscle

(gastrocnemius and soleus), white adipose tissue (inguinal (iWAT) and perigonadal (gWAT)),

and organ (spleen and liver) weight (**Fig. 1D**). No group differences were found in serum

inflammatory cytokine concentrations (**Fig. 1E**).

Skeletal muscle and adipose tissue cellular wasting in G12D mice.

To further characterize tissue wasting, myofiber size was assessed at 12 weeks post-

induction. *G12D* mice exhibited reduced gastrocnemius (**Fig. 2A, Fig. S5A**), but not soleus,

muscle fiber cross-sectional area (CSA; **Fig. 2A**, **Fig. S5B**). Gastrocnemius muscle of *G12D*

mice showed greater expression of key genes involved in protein breakdown that are implicated

in CC (*Fbxo32, Becn1*), and *Il6, Tnf,* and *Lif* mRNA compared with WT (**Fig. 2B**), with cytokines

indicating local inflammation that may drive muscle wasting[30]. In soleus, mRNA levels of *Il6*

were greater in *G12D* mice, with no differences in proteolytic or inflammation markers (**Fig.**

S5C).

Subcutaneous iWAT and visceral gWAT adipocytes exhibited markedly lower CSA at 12

weeks post-induction in *G12D* mice compared with WT mice (**Fig. 2C**), and we observed higher

circulating medium and long-chain fatty acids and markers of carnitine/fatty acid metabolism

(**Fig. S6**), as well as a trend (p=0.06) toward greater circulating glycerol levels (**Fig. 2D**). In

G12D iWAT, mRNA of the lipid transporter, *Cd36,* and mitochondrial uncoupling marker, *Ucp1,*

were lower, and cytokines, *Cxcl1* and *Lif,* greater compared with WT (**Fig. 2E**). In gWAT of

G12D mice, we observed greater expression of mRNA for the beiging marker, *Cidea*, and

trends toward greater *Cd36* (p=0.06) and *Lipe* (p=0.07; **Fig. 2F**), the latter being the transcript of

hormone-sensitive lipase. As expected, mRNA levels of *Lep* (leptin) were lower in iWAT and

gWAT of *G12D* mice compared with WT mice (**Fig. 2E-F**), reflecting reduced adipocyte size. Of

note, controlling for sex did not alter group differences in adipose and muscle cell size and

tissue weights. Collectively, these data show that *G12D* mice exhibit tissue, cellular,

transcriptomic, and metabolomic hallmarks of CC muscle and adipose wasting.

Similar driver mutations can elicit different tumor metabolic phenotypes in epithelial cells

in different organs (lung vs. pancreas [31]), which, in turn, may affect the nature of CC. We chose

to express *Kras^{G12D}* in club cells because this is a recognized model of lung adenocarcinoma[32],

but adenocarcinoma can also develop in alveolar type-2 (AT2) cells[33]. Thus, we evaluated mice

expressing *Kras^{G12D}* in AT2 cells following induction via adenovirus that permits AT2-specific

Cre recombinase expression (Ad5-SPC-Cre). At ~12 weeks post-induction, we found no

differences in skeletal muscle, adipose depot, or organ weights compared to animals

administered control adenovirus (**Fig S7A**). We also activated the *Kras^{G12D}* allele using an

adenovirus that permits epithelial cell agnostic Cre recombinase expression (Ad5-CMV-Cre). In

these mice, we found reduced weights of two adipose depots and the spleen and patterns

towards reductions in other adipose depots, albeit non-significant, but no differences in skeletal

muscle weights (**Fig S7A**), suggesting cell type-specific differences in the ability of *Kras^{G12D}* to

incite CC. The general absence of overt CC wasting phenotypes in mice with both IT Ad-Cre

induction approaches likely related to reduced tumor burden, as we found relative tumor

burdens in Ad-Cre mice equivalent to our *G12D* mice at 3 weeks (~5%; **Fig S7B**). Thus, the

induction method (*i.e.,* intraperitoneal tamoxifen vs. oropharyngeal adenovirus expression Cre

recombinase) affects the timing of CC, although the phenotype of early adipose wasting

appears to be similar when considered relative to tumor burden .

Early tissue wasting phenotypes in G12D mice.

Utilizing the protracted time course of wasting in *G12D* mice, we evaluated *G12D* mice at 6 weeks post-induction, at the initiation of weight loss (Fig. 1C), to examine tumor and peripheral tissue morphological, transcriptomic, and metabolic features of CC initiation. Tumor burden averaged 10.5 ± 1.1% (**Fig. 3A-B**) while *G12D* mice lost an average of 5% of their body weight (**Fig. 3C**). Assuming an average lung weight of 200 mg, we estimate tumor burdens of ~0.08% and ~0.16% of body weight at 6- (Fig. 3B) and 12 (Fig 1B) weeks post-induction, respectively, which is closer to human lung cancer than other lung CC models. No differences in skeletal muscle or organ weights were noted between groups 6 weeks post-induction, but *G12D* mice showed markedly lower iWAT and gWAT weights (**Fig. 3D**). In a separate group of animals, we assessed other skeletal muscles (plantaris, tibialis anterior, diaphragm, and extensor digitorum longus) and found no differences in their weights (**Fig S8A**). Finally, no differences were found in a broad range of circulating cytokines (**Fig. 3E; Fig S8B**).

At the cellular level, consistent with skeletal muscle weights (Fig. 3D), neither gastrocnemius nor soleus muscle fiber CSA differed between groups at 6 weeks post-induction (**Fig. 4A**, **Fig. S8C-D**). Moreover, except for a trend (p=0.06) towards statistically significant greater *Fbxo32* mRNA in the gastrocnemius of *G12D* mice compared with WT mice, no differences in gene expression for markers of protein degradation or inflammatory cytokines were observed (**Fig. S7E-F**). Additionally, we found no group differences in tibialis anterior muscle fiber CSA (**Fig S8G**). In contrast, *G12D* mice showed marked atrophy of adipocytes in both depots at 6 weeks post-induction (**Fig. 4B**). In *G12D* mice, gene expression markers of lipid balance (*Plin1, Pnpla2,* and *Lipe*) were elevated in iWAT (**Fig. 4C**), with a similar expression pattern in *Plin1* and *Pnpla2* (both p=0.09) in gWAT compared with WT mice (**Fig. 4D**). In both depots, we found no differences in mRNA levels of inflammatory cytokines. However, as local tissue inflammation may contribute to CC [30], we conducted a more in-depth,

unbiased assessment of inflammatory cytokine expression in gWAT at 3 and 6 weeks post-

induction by bulk RNA-seq (Fig 4E). At 3 weeks post-induction in gWAT, *G12D* gWAT showed

20 differentially expressed genes (DEG; 7 upregulated and 13 downregulated) compared with

WT gWAT (**Fig S9A**). Pathway enrichment analysis showed no evidence for upregulation of

inflammatory pathways/signaling (**Fig S9B**). In fact, inflammatory response and cytokine

signaling were downregulated. By 6 weeks post-induction in gWAT, *G12D* gWAT showed 31

differentially expressed genes (DEG; 17 upregulated and 14 downregulated) compared with WT

gWAT (**Fig S9C**). Pathway analysis showed upregulation of inflammatory response and

numerous cytokine signaling pathways (**Fig S9D**). Thus, adipose tissue inflammation develops

coincident with CC-associated wasting, which suggest the possibility that, like muscle [30] (Fig

2B), adipose wasting may be modulated by local tissue inflammation.

In keeping with adipocyte wasting, we observed greater circulating glycerol levels in

G12D mice compared with WT mice (**Fig. 4E**), a marker of increased adipose tissue lipolysis,

the process driving adipose wasting in human CC[34]. Patients with NSCLC similarly showed

elevated circulating glycerol levels (**Fig. 4F**), suggesting that increased lipolysis in *G12D* mice

translate to human lung cancer. Finally, to further discriminate the timing of adipose wasting, we

evaluated *G12D* and WT mice 3 weeks post-induction. At 3 weeks, we found no differences in

fat pad or organ weight (**Fig S10A**), and no differences in adipocyte or muscle fiber CSA (**Fig**

S10B-D). Examining *G12D* adipocyte size over time relative to WT, we found that the

progressive reduction in adipocyte size (**Fig 4E**) was inversely associated with increasing tumor

burden (**Fig. 4F**). Coupled with the lack of muscle wasting at 3- and 6-weeks, our data suggest

that early adipose tissue wasting is important for CC initiation and potentially tumor growth.

Absent differences in energy intake or fecal energy content (Fig S4), and considering

that physical activity is not elevated in CC, we explored energetically futile cycles that may

underlie elevated energy expenditure. In some models of CC, adipose wasting may be

explained by beiging of white adipose tissue [35, 36]. In *G12D* mice, however, there were no

differences in thermogenic gene expression or markers of beiging at 12 weeks post-induction,

aside from an upregulation of *Cidea* in gWAT (Fig. 2E-F). In fact, *Ucp1* mRNA expression was

lower in iWAT. At 6-weeks post-induction when weight loss begins, there was a modest

elevation in mRNA levels of *Ucp1* in iWAT and a trend towards a reduction in eWAT in *G12D*

mice (Fig. 4F-G) but no difference in other markers of beiging. To further examine *Ucp1*

expression, we measured UCP-1 tissue protein content via immunohistochemical staining [36]

(**Fig. S11**) and found no differences in UCP-1-positive area (**Fig. S12**), in agreement with work

in other mouse models of CC[37]. Finally, whereas UCP-1 expression in other *Kras*-driven lung

CC models was dependent on increased circulating IL-6 [36], *G12D* mice did not show alterations

in circulating IL-6 concentrations (Fig 1E, 3E) or adipose tissue *Il6* mRNA levels (Fig. 2E-F, 4F-

G). Thus, there is no evidence for upregulation UCP-1-associated WAT beiging.

To explore other UCP-1-independent energy consuming processes that may contribute

to adipose wasting, we evaluated bulk RNA-seq data on adipose for alterations in other energy

wasting futile cycles (lipid [37] creatine [38], Ca^{2+}cycling [39]). Using pathways analysis, with genes

involved in these futile cycles included as custom pathways (**Supplemental Table 1**), pathway

enrichment analysis showed upregulation of fatty acid metabolism and lipogenesis, including

our custom pathway of genes for lipid cycling at 3 weeks (**Fig S9**). The expression of genes for

creatine and Ca^{2+} cycling (**Supplemental Table 3**), in contrast, did not differ between groups.

Upregulation of pathways of lipid turnover in adipocytes are congruent with increased serum

glycerol in *G12D* mice (Fig 4E) and implicate increased lipid cycling as a potential source of the

energetic imbalance contributing to CC, in agreement with other models of CC[37]. At 6 weeks

post-induction, pathway enrichment analysis showed that pathways for fatty acid metabolism

and adipogenesis, including our custom pathway for lipid cycling, were among the top down-

regulated genes (**Fig S9**), which may reflect a compensatory adaptation to slow adipose

wasting given that considerable atrophy has occurred by 6 weeks (**Figs 4B, S8A, S9**). At this

time, neither creatine nor Ca^{2+} cycling pathways were upregulated.

Finally, as tumors may cause liver metabolic reprogramming, we characterized liver

phenotypes. Liver weights did not differ at 3 (**Fig S10A**) or 6 weeks (Fig 3D) but were reduced

at 12 weeks (Fig 1D). Additionally, we observed a steady increase in liver triglyceride content in

G12D mice, which became significant by 12 weeks (**Fig S13**), which may result from

upregulated adipose tissue lipolysis (Fig 4E and [40]). Thus, in addition to adipose and skeletal

muscle tissue, our model shows evidence for liver metabolic reprogramming [41-44], underscoring

its utility as a model of multi-organ, systemic metabolic reprogramming in CC [45].

Tissue wasting phenotypes in lung epithelial cell specific Kras^{G12D/+}-Stk11 null mice.

As patients with lung cancer with mutated *KRAS* lose more weight than patients with

wild-type *KRAS*[46], our model is relevant to human lung tumor oncotypes that predispose to CC.

Recently, studies suggest that serine/threonine kinase 11 (*STK11*) mutations are common

among patients with NSCLC who report weight loss at diagnosis[47]. Additionally, mice with

combined *Kras^{G12D/+}-Stk11^{-/-}* demonstrate more rapid weight loss[48] compared with *G12D* mice

(Fig. 1C). We therefore hypothesized that lung epithelial-specific *Kras^{G12D/+}-Stk11^{-/-}* should

present a more aggressive form of CC than *Kras^{G12D}* alone. To examine this hypothesis, we

generated mice with tamoxifen-inducible, club cell specific *Kras^{G12D/+}-Stk11^{-/-}* (**Fig. 5A**).

Following tamoxifen induction, *KS* mice exhibited body weight loss over a period of 3.5 weeks

(**Fig. 5B**), which diverged from *G12D* and WT 1.5 weeks post-induction, independent of

differences in food intake (**Fig. 5C**). More rapid weight loss in *KS* mice was coincident with

greater tumor burden compared with *G12D* mice (**Fig. 5D**), and weight loss was related

primarily to loss of adipose tissue, as no differences in skeletal muscle or organ weights were

observed (**Fig 5E**). *KS* mice had greater circulating levels of IL-6 compared with WT and *G12D*

mice and greater CXCL1 compared with WT mice, but no differences in other inflammatory

cytokines (**Fig. S14)**. Finally, gastrocnemius muscle fiber CSA was not statistically different

among *KS, G12D,* and WT animals (**Fig. 5F**), whereas adipocyte CSA in iWAT was smaller in

KS compared with *G12D* mice and in gWAT was lower in *KS* and *G12D* compared with WT

(**Fig. 5G**). Collectively, these data confirm reports that combined *Kras*$^{G12D/+}$*-Stk11*$^{-/-}$ promotes

rapidly progressing CC, show that increasing the number of driver mutations hastens CC

development, and indicate that early adipose wasting is maintained across *G12D*-driven models

of lung CC with different rates of weight loss.

Tumor-derived factors promote CC and adipose tissue wasting.

CC is thought to be triggered by factors released from tumors[49]. To examine whether

tumor-derived factors initiate adipose tissue wasting, we developed tumor and lung tissue

organoids from WT and *G12D* mice, respectively (**Fig. 6A**), from lungs collected at 6-weeks

post-induction. Using bulk RNA-seq, we confirmed that the *Kras*^{G12D} mutation present *in vivo*

was retained in the cultured *G12D* organoids (**Fig. 6B**). Moreover, *G12D* lung tumor organoids

showed 1,076 differentially expressed genes (DEG; 541 upregulated and 535 downregulated)

compared with WT lung tissue organoids (**Fig. 6C**), with upregulation of numerous oncogenic

pathways (**Fig. 6D**). As studies suggest that tumor-derived cytokines contribute to CC [49-52], we

examined differences between in the expression of inflammatory genes and found 35 DEG, with

23 upregulated and 12 downregulated (**Fig. S15**). None of these DEGs have been implicated in

CC, except *Gdf15* (growth differentiation factor-15) [53], the expression of which was elevated in

G12D tumor organoids, with no elevation of GDF-15 in serum from *G12D* mice (Fig. 3E and

1E). Thus, we found no evidence that cytokines from *G12D* tumor organoids may incite CC by

increasing systemic inflammation.

To determine whether tumor-released factors may promote CC in *G12D* mice, we

examined whether factors secreted from *G12D* lung tumor organoids alter adipocyte

metabolism to promote atrophy by treating murine 3T3L1 adipocytes with conditioned media

(CM) from WT tissue or *G12D* tumor organoids (**Fig. 6E**) and assessed lipolysis by measuring

media glycerol levels. Treatment of adipocytes with *G12D* tumor organoid CM for 24 hours

increased media glycerol levels compared with WT organoid CM (**Fig. 6F**) independent of

glycerol levels in *G12D* CM (**Fig. S16A**).

To define how tumor-derived factors increase glycerol release in adipocytes, we

evaluated potential mediators. *G12D* CM increased media inflammatory cytokine levels

compared with WT CM (**Fig. 6G-I**), independent of cytokine levels in CM (**Fig. S16B-E**). As one

of these cytokines, IL-6, has been implicated as a mediator of CC and stimulates lipolysis[54] , we

applied *G12D* lung tumor organoid CM that was treated with a neutralizing antibody directed

against IL-6 or an isotype control to 3T3-L1 adipocytes. The IL-6 neutralizing antibody reduced

IL-6 in the culture media to undetectable levels (**Fig. S16F**), but glycerol levels remained

elevated (**Fig. S16G**). Moreover, adding IL-6 to WT lung tissue organoid CM did not increase

glycerol release (**Fig. S16F-G**), suggesting that IL-6 is neither necessary nor sufficient to explain

increased adipocyte glycerol release.

GDF-15 is slightly increased in *G12D* CM, was recently shown to be increased in

circulation of patients experiencing weight loss with recurrence of lung cancer[55] and may

increase adipocyte lipolysis [56]. To explore whether elevated GDF-15 mediates the increased

glycerol release, we treated 3T3-L1 adipocytes with GDF-15 at levels found in *G12D* (2,500

pg/mL) and WT (1,500 pg/mL) organoid CM (**Fig. S16E**). While GDF-15 at both levels increased

glycerol above vehicle control, there was no difference in glycerol between the two

concentrations, suggesting that GDF-15 likely does not increase adipocyte lipolysis.

Studies have also implicated TLR4 activation in adipose wasting in the LLC model of

lung CC[57]. Thus, we evaluated whether TLR4 activation via lipopolysaccharide (LPS) increased

glycerol release by 3T3-L1 adipocytes. Activation of TLR4 by LPS was confirmed by robust

cytokine release into the media (**Fig. S16I-K**). However, LPS did not increase glycerol release in

3T3-L1 adipocytes (**Fig S16L**), arguing against TLR4 activation as the mediator of *G12D* CM

effects on 3T3-L1 adipocytes.

As we observed an increase in adipose inflammation coincident with tissue wasting (Fig.

S9) and that macrophages in tissues may drive tissue wasting in CC [58, 59], we examined the

effects of co-culturing bone marrow-derived macrophages (BMDM) with adipocytes on the

cytokine/chemokine response to CM . We found that BMDM enhanced the ability of *G12D* CM

to increase MCP-1 and CXCL1 release (**Fig. S16N-O**), but not IL-6 (**Fig S16M**), suggesting that

macrophages may recruit circulating immune cells in response to tumor-derived factors to drive

local inflammation (Fig S9B and [30]). Thus, *G12D* tumor organoids release factors that increase

adipose tissue lipolysis and may contribute to the development of inflammation in adipose tissue

in vivo (Fig S9D), phenotypes that parallel adipocyte wasting and inflammation in patients with

CC[34, 59, 60].

Finally, we evaluated whether *G12D* CM could promote wasting of skeletal muscle cells

in culture. Treatment of mature, murine C2C12 myotubes [61] with *G12D* CM for 72 hours did not

reduce myotube diameter compared to WT CM (**Fig. S16P**), in keeping with the lack of effects

of CM from 2D cultures of *Kras^{G12D}*-transformed murine lung epithelial cells and human lung

cancer cell lines[61].

Early adipose tissue wasting in G12D mice translates to human patients with lung cancer.

To examine whether early changes in adipose tissue in *G12D* mice translate to patients

with lung cancer, we carried out a secondary analysis using body composition data collected

during the first 3 years of the National Lung Screening Trial (NLST), a longitudinal study

examining the efficacy of yearly, low-dose chest computed tomography (CT) scans for early

lung cancer diagnosis[62] (baseline patient characteristics in **Supplemental Table 2**). Using

multivariable-adjusted models, females diagnosed with stage 3-4 lung cancer exhibited a

reduction in visceral adipose tissue (VAT) cross-sectional area indexed to height (**Fig. 7A**) and

an increase in VAT density (**Fig. 7B**; reduced adipocyte size [63]) compared with controls. In

males with stage 3-4 lung cancer SAT density (**Fig 7C**) and skeletal muscle were decreased

compared with controls (**Fig. 7D**). Data for non-significant results are shown in **Fig. S17-18**. Our

results demonstrate that there is adipose loss prior to diagnosis in patients with lung cancer and

a sex dimorphism in these adaptations. Nonetheless, these data show that early adipose tissue

adaptations observed in *G12D* mice translates to human lung cancer.

Discussion:

We report the development of a new model of lung CC using mice with a lung epithelial cell-specific, tamoxifen-inducible $Kras^{G12D}$ mutation. *G12D* mice exhibit more physiologically relevant tumors with respect to their anatomic location and burden, a slower time course of CC initiation and progression, and peripheral tissue and tumor morphological, mutational, transcriptomic, and metabolic characteristics that parallel CC in patients with lung cancer. Moreover, adipose tissue wasting drove CC initiation, could be incited by *G12D* lung tumor released factors, was inversely related to tumor burden, was apparent across different *Kras*-driven lung CC models with differing mutational burdens and induction approaches, and translated to human patients, highlighting an early role for tumor-induced metabolic reprogramming of adipose tissue wasting in CC.

Tumor location, burden, and mutations

Most studies of CC have employed heterotopic murine models, in which tumor cells are injected into the flank, producing a tumor akin to a metastatic lesion but in a microenvironment (*i.e.*, subcutaneous) that is atypical outside of melanoma. Recent studies have advanced this approach by administering cancer cells with combined *Kras* and *Tp53* mutations into the vasculature or lungs to seed tumors in the lung [64, 65]. In *G12D* mice, the transgene is targeted to lung epithelial cells, and tumors originate in the terminal bronchioles, the anatomic location of most lung tumors in humans [66]. Moreover, at the initiation of weight loss (6 weeks post-induction), tumor burden in *G12D* mice is small (<<1% of body weight) relative to other models, better reflecting tumor burden in patients. An additional advantage of *G12D* mice is that *KRAS* is the most frequently mutated oncogene ortholog in human NSCLC [67], the majority of patients with lung cancer have elevated *RAS* pathway activation [68] and those with mutated *KRAS* experience greater weight loss [46]. Building on recent reports suggesting that *STK11* loss-of-function amplifies lung CC in humans [47], our data demonstrate that mice with combined

Kras^{G12D}/Stk11^{-/-} develop a more rapidly progressive CC than *Kras^{G12D}* alone, and comparable to

the timeframe of recently developed models using cells with *Kras/Tp53* mutations [64, 65]. Thus,

our approach of using inducible, epithelial cell specific expression of transgenes provides a

useful platform to study human lung tumor oncotypes that predispose to CC and offers a model

that approximates tumor characteristics in human lung CC with respect to anatomic location,

burden, and mutations that predispose to CC. Based on our results showing potentially different

tissue wasting phenotypes when *Kras^{G12D}* is expressed in different lung cell types (club vs. AT2

cells), further studies are needed to evaluate cell- and driver mutation-specific variation in CC

and how this relates to CC in patients with different tumor oncotypes [46, 47].

CC initiation and progression

While recent lung CC models have made advances in localizing tumors to the lungs [64,

65], they show rapid onset and progression of weight loss, which does not reflect what occurs in

patients. In contrast, the initiation and progression of weight loss in *G12D* mice is slower, with

an ~4-6-week latency until weight loss begins, followed by progressive weight loss until 12

weeks post-induction. The relatively slow rate of progression of weight loss in *G12D* mice

(~1.9%/week) more closely approximates that in patients with lung cancer [69] and offers a

comparatively longer time to intervene on CC with therapeutics (~6-8 weeks).

Early adipose tissue wasting

Exploiting the protracted time course of the *G12D* model to examine the temporal

context of body composition changes in CC, we report early wasting of adipose tissue in *G12D*

mice. Considering the relative contribution of adipose tissue to body weight in adult C57BL/6J

mice[70] and our observed reductions at 6 weeks post-induction (>50%), loss of adipose could

explain the majority of body weight loss. Reinforcing these data, we found early adipose tissue

wasting in mice with combined *Kras^{G12D/+}-Stk11^{-/-}*, which exhibit more rapidly progressive weight

loss than *G12D* mice, and in *Kras*-driven lung cancer induced via adenoviral Cre recombinase,

suggesting that early adipose tissue wasting remains a defining characteristic of *Kras*-driven

lung CC across models that differ in mutational burden, time course, and induction techniques.

Factors released from *G12D* tumors may drive early adipose tissue wasting, as we

found that cultured adipocytes treated with *G12D* lung tumor organoid CM release greater

amounts of glycerol into the media, a proxy of lipolysis, compared with adipocytes treated with

WT lung tissue organoid CM. Similar effects were evident *in vivo*, as circulating glycerol levels

were increased in *G12D* mice at 6 weeks post-induction, when weight loss begins. In fact,

trends towards elevated circulating glycerol remained at 12 weeks post-induction, along with

increased circulating markers of fat availability/metabolism. Moreover, we found elevated

glycerol levels in patients with NSCLC compared with non-cancer controls. As increased

lipolysis contributes to adipose wasting in other animal models of CC[37] and humans [34], these

data suggest that tumor-released factors may drive elevated adipose tissue lipolysis in *G12D*

mice.

The importance of early adipose tissue wasting is apparent in recent prospective data in

patients with pancreatic cancer, which showed that the relative loss of adipose tissue was twice

that of skeletal muscle prior to diagnosis and was a stronger predictor of body weight loss [71].

Moreover, longitudinal studies show that adipose wasting constitutes the majority of weight loss

in patients with CC [72]. Patients with lung cancer lose body weight prior to diagnosis [69] but,

based on our recent systematic review, there are no longitudinal data examining early body

composition changes[73]. To address this lack of data, using AI-based approaches [74, 75], we

conducted a secondary analysis of data from the NLST [76]. Both male and female patients with

advanced-stage disease experience changes in adipose tissue size and composition prior to

lung cancer diagnosis, with females experiencing a decrease in VAT size and an increase in

density. Sex differences in the anatomic localization of adipose in the thoracic compartment,

where CT scans were conducted, make it challenging to draw conclusions about biological sex-

specific changes[77] and further studies evaluating other adipose depots and/or whole-body

adiposity are needed. Nonetheless, these data suggest that early adipose tissue wasting in

G12D mice translates to human patients.

Early adipose tissue wasting in CC may have pathological consequences, as lipid

uptake is required to support cancer cell proliferation under certain conditions[78] and elevated or

reduced availability of lipid substrates in conditions of high-fat feeding or caloric restriction,

respectively, increases or decreases proliferation[79]. Further suggesting a role for adipose tissue

substrates in cancer progression, we found an inverse association between adipocyte wasting

and tumor burden in *G12D* mice. The modulatory role of endogenous lipid substrates on tumor

growth is an important topic given the current epidemic of obesity, and our results underscore

the utility of the *G12D* model to evaluate a potentially novel role for CC-induced adipose tissue

wasting in cancer progression.

Limitations of the Study

Several limitations to our study deserve discussion. First, tumors in *G12D* mice were

constrained to the lung, limiting the ability of this model to study the impact of tumor metastases

on CC. Second, organoids do not reproduce all cell types and the physiology of tumors in the

lung, nor does use of their CM in cultured adipocytes reproduce the complex array of cells,

organs and tissues *in vivo*. Similarly, because organoids are developed from whole lung

homogenates, we cannot evaluate how lesion heterogeneity contributes to CC. Third, as the

NLST does not contain information on tumor oncotype, it is unclear if body composition changes

reflect the effect of mutant *KRAS*. However, 84% of lung adenocarcinomas have elevated *RAS*

pathway activation[68], suggesting that tumors in most NLST patients likely exhibit increased *RAS*

pathway activation. Finally, we did not identify the molecule or molecules that are necessary

and sufficient for CC in *G12D* mice. Considering the protracted nature of CC and differential

time course of adipose and skeletal muscle wasting, these could vary by tissue type, along the

continuum of CC, and/or involve complex interactions with other organs and tissues. Further

work will need to identify the complex tumor-host interactions that drive lung CC in this and

other clinically representative models to derive therapies that can alleviate CC.

Conclusions

Our data show that the *G12D* mouse lung cancer model phenocopies tissue, cellular,

mutational, transcriptomic, and metabolic characteristics of human CC. Additionally, because of

the protracted time course for initiation and progression of CC relative to other pre-clinical lung

CC models, *G12D* mice are well-suited to identify mechanisms initiating CC, as well as testing

preventative and remediative therapies. Using this model, we identify adipose tissue wasting as

an early event in *G12D* and *Kras^{G12D}-Stk11$^{-/-}$* mouse models of lung CC that may be mediated

by tumor-released factors. The translational relevance of this model is supported by our data

from a large, prospective trial, where we observed that adipose wasting prior to lung cancer

diagnosis. Together, these findings underscore the utility of this model to study the mechanisms

of CC and to the role of early adipose tissue wasting in CC and cancer progression.

Acknowledgements:

This study was funded by NIH R01 AR065826 (MJT), R21 CA283492 (MJT), T32 HL076122 (MEP), R21 CA280266 (DJS), R01 CA273238 (YJ-H), R01 CA253923 (BAL), and the University of Vermont Cancer Center. Imaging work was performed at the Microscopy Imaging Center at the University of Vermont (RRID# SCR_018821) . The Graphical Abstract was created in BioRender. Snoke, D. (2025) https://BioRender.com/zxbk8g3. A UVM College of Medicine Shared Instrumentation Award to D.T. supported the purchase of the Leica-Aperio VERSA whole slide imaging system used for image acquisition. The University of Colorado SOM Metabolomics Core is supported in part by the University of Colorado Cancer Center via NIH P30 CA046934 and the University of Michigan Animal Phenotyping Core via grants U2CDK110768, KD020572, and DK089503. Funding in support of the RNA-seq was provided by a Zymo Impact Initiative Grant (DBS). The authors thank Vikas Anathy and Zoe Mark for the machinery and technical support needed to measure oxygen saturation.

Author Contributions

Conceptualization: MJT, DBS; Methodology: DBS, MEP, DM, DJS, RA, YJH, KX; Validation: DBS ; Formal analysis: DBS, DM, JSD, TPA; Investigation: DBS, JLA, JV, AEB, RA, SML, HS, ERB, SCJH, MMM, DS, JAR; Resources: MJT, MEP, JV, DM, DJS, YJH, AD, BAL, KLS; Data Curation: DBS, HS, ERB, SCJH, SML, DS, KX; Writing- original draft preparation: DBS, MJT; Writing- review and editing: all authors; Visualization: DBS, TPA, JSD; Supervision: MJT, MEP, DJS, DBS; Project administration: DBS and MJT; Funding acquisition: MJT, MEP, YJ-H and DBS.

Declaration of Interests: K.L.S.: Consulting fees from Re-veal Dx; leadership or fiduciary role in the National Lung Cancer Roundtable, National Comprehensive Cancer Network, Eastern Cooperative Oncology Group–American College of Radiology Imaging Network Cancer Research Group, and the American College of Radiology

Main Figure Titles and Legends

Figure 1. Characterization of lung epithelial cell specific, inducible Kras$^{G12D/+}$ (G12D) mice at 12 weeks post-induction.

Kras$^{G12D/+}$ (G12D) mice harbor a tamoxifen (TAM)-inducible lung epithelial cell specific *Kras^{G12D}* mutation under the club cell secretory protein (CCSP) promoter (A). Four- to 5-month-old mice received intraperitoneal TAM, leading to adenocarcinoma development in lung bronchioles by 12 weeks post-induction compared with wild-type (WT) mice and average tumor burden (percent tumor area / total lung area) (B). Change in body weight relative to Day 0 (C), skeletal muscle, fat pad and organ weights (D) and serum inflammatory cytokines (interleukin 6 (IL-6), soluble IL-6 receptor alpha (IL-6Rα), tumor necrosis factor alpha (TNFα), CXC motif ligand 1 (CXCL1/KC), growth differentiation factor 15 (GDF-15), monocyte chemoattractant protein 1/C-C motif ligand 2 (MCP-1/CCL2), CXCL10, and interleukin-1 beta (IL-1β)) in wild-type (WT; n=7; blue bars/lines) and G12D mice (n=6; gold bars/lines)(E). Data were analyzed using repeated measures ANOVA (C) or Student's t-test (D, E), with * $p<0.05$, ** $p<0.01$, and *** $p< 0.001$. Note that n=5-7/group for serum cytokine measures. Image created with Biorender.com (A). See also: supplemental figures S1-4.

Figure 2. Effect of lung epithelial cell specific, inducible Kras$^{G12D/+}$ (G12D) on skeletal muscle and adipose tissue wasting and mRNA expression at 12-weeks post-induction.

Gastrocnemius (gastroc) and soleus fiber cross-sectional area (CSA) 12 weeks post-induction in wild-type (WT; n=7; blue bars) and *Kras$^{G12D/+}$ (G12D)* mice (n=6; gold bars) and representative images (scale bars=100 μm) (A). mRNA levels for markers of protein degradation and inflammatory cytokines in gastrocnemius muscle, expressed relative to the WT (B). Cross-sectional area (CSA) of adipocytes from inguinal (iWAT) and gonadal white adipose tissue (gWAT) depots in WT (n=7) and *G12D* (n=6) mice and representative sections (scale

bars= 100 μm) (C). Serum glycerol levels in WT and *G12D* mice standardized to body mass (D).

Gene expression of iWAT (E) and gWAT (F) depots for markers of lipid balance, inflammation,

and beiging. Data were analyzed by Student's t-test, with * $p<0.05$ and *** $p<0.001$. Significance

values where $p>0.05$. See also: supplemental figures S5-7.

Figure 3. Characterization of lung epithelial cell specific, inducible Kras[G12D/+] *(G12D) mice CC at*

6-weeks post-induction.

Representative lung cross-section of a *Kras*[G12D/+] (*G12D*) mouse (A) and tumor burden (percent

tumor area / total lung area) (B). Change in body weight relative to Day 0 in wild-type (WT; n=7;

blue lines/bars) and *G12D* (n=8; gold lines/bars) mice (C). Muscle, fat pad, and organ weights of

6-week WT and *G12D* mice (D). Serum inflammatory cytokines in 6-week mice (E). Data were

analyzed by repeated measures analysis of variance (C) and Student's t-tests (D, E), with **

$p<0.01$ and *** $p<0.001$. See also: supplemental figure S8.

Figure 4. Effect of lung club epithelial specific, inducible Kras[G12D/+] *(G12D) on skeletal muscles*

and adipose tissues 6 weeks post-induction.

Muscle fiber cross-sectional area (CSA) in gastrocnemius and soleus muscles of wild-type (WT;

n=7; blue bars) and *G12D* (n=8; gold bars) mice were assessed 6 weeks post tamoxifen

injection, with representative images (scale bars=100 μm) (A). Cross-sectional area (CSA) of

adipocytes from inguinal (iWAT) and perigonadal (gWAT) white adipose tissue depots in 6-week

WT and *G12D* mice, with representative images (scale bars=100 μm) (B). Gene expression in

iWAT (C) and gWAT (D) of *G12D* and WT mice for lipolysis, inflammation, and beiging. Glycerol

levels in WT and *G12D* mice (E), and in plasma of patients with late-stage, non-small cell lung

cancer (NSCLC; n=7; purple bar) compared with healthy controls (Healthy; n=11; black bar) (F).

White adipose tissue CSA in *G12D* mice expressed as a percentage of WT CSA at each

timepoint (G). Tumor area expressed as a percentage of the total lung area over the 12-week

study period (H). Data (A-D) were analyzed Student's t-test with asterisks indicating * $p<0.05$

and ** $p<0.01$. See also: supplemental figures S7-13.

Figure 5. Comparison of lung epithelial cell specific, inducible Kras$^{G12D/+}$ (G12D) and

Kras^{G12D}/Stk11$^{-/-}$ (KS) mice.

KS-tm mice harboring tamoxifen (TAM)-inducible lung epithelial cell specific *Kras^{G12D}*, deletion of

Stk11/Lkb1 and a *Rosa26-tdTomato (tm)* reporter allele (A). Percent change in body weight (B),

cumulative food intake (C) and tumor burden (D) of wild-type (WT-tm; n=4; blue

lines/bars/circles), *G12D-tm* (n=5; gold lines/bars/squares) and *KS-tm* (n=5; red

lines/bars/triangles) mice in. Muscle, fat pad and organ weights at necropsy (E). Mean

gastrocnemius (Gastroc) muscle fiber (F) and iWAT and gWAT adipocyte (G) CSA. Data were

analyzed using a repeated measures ANOVA (B), Student's t-test (D), or a one-way ANOVA (C,

E-G) with post hoc comparisons $p<0.05$ group or group X time interaction effects. ** $p <0.01$

different letters by post hoc comparisons. Image created with BioRender.com (A). See also:

supplemental figure S14.

Figure 6. Organoids developed from lung tissue from wild-type (WT) and lung tumors from

Kras$^{G12D/+}$ (G12D) mice and their characterization.

Lung organoid development with representative images of wild-type (WT; blue) and *G12D*

organoids (gold; A; scale bars = 250 µm). RNA-seq was used to validate the *Kras^{G12D}* mutation

(gold) in lung tumor organoids (B). Volcano plot depicting differentially expressed genes, with

red representing upregulated (n=535) and blue representing downregulated (n=541) genes,

respectively, in *G12D* compared with WT organoids, with a *p*-value exclusion of 0.05 (C).

Pathway analysis depicting upregulated signaling pathways in *G12D* lung tumor compared with

WT lung tissue organoids (D). CM was applied (1:3 with adipocyte culture media) to fully

differentiated 3T3L1 adipocytes (A). Glycerol levels of supernatant of 3T3L1 adipocytes treated

with WT (blue circles) or *G12D* (gold squares) organoid CM for 24 h (F). Supernatant levels of

IL-6 (G), MCP-1 (H), CXCL1 (I) and GDF15 (J) from 3T3L1 adipocytes treated for 24 hours with

WT or *G12D* organoid CM. 3T3L1 experiments are representative of 2-3 independent replicates.

Student's t-tests were used, with *** $p<.001$. Images created with BioRender.com (A, E). See

also: supplemental figures S15-16.

Figure 7. Changes in body composition prior to lung cancer diagnoses among a matched

cohort sample of patients in the National Lung Screening Trial (NLST). We conducted

secondary analysis of body composition from computed tomography scans collected during the

first three years of the NLST using multivariable hierarchical linear models. Modeled data were

visualized by plotting marginal predicted values across the range of time values by exposure

group and stratified by sex, and we present 95% confidence intervals as shaded regions. Data

are shown for non-cases (black lines) and patients diagnosed with stage 1-2 (blue) or stage 3-4

lung cancer (A) when stratified by diagnosis and stage or by non-cases (black lines) and lung

cancer cases (green lines) when stratified by diagnosis only. Visceral adipose tissue (VAT) area

indexed to height transformed via square root to meet normality (A) and density in Hounsfield

units (HU), measured by tissue attenuation (B). Subcutaneous adipose tissue (SAT) density,

measured by tissue attenuation (C). Skeletal muscle area indexed to height in non-cases and

lung cancer cases and transformed via square root to meet normality (D). *P*-values and

comparison groups are listed in each respective figure. See also: supplemental figures S17-18.

532 **Methods**

533 *Resources table*

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
UCP-1	Abcam	Abcam ab23841; RRID: AB_2213764
Goat-anti-mouse IgG	Abcam	Abcam ab205718, RRID: AB_2819160
InVivoMAb anti-mouse IL-6	BioXCell	BE0046
InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase	BioXCell	BE0088
Bacterial and virus strains		
Ad-5CMV-Cre	University of Iowa Viral Vector Core	VVC-U of Iowa-5-HT
Ad-5mSPC-Cre	University of Iowa Viral Vector Core	VVC-Berns-1168-HT
Ad-5CMV-empty	University of Iowa Viral Vector Core	VVC-U of Iowa-272-HT
Biological samples		
Plasma samples	Patients with non-small cell lung cancer undergoing standard-of-care treatment	Snoke et al. 2023[80]
Chemicals, peptides, and recombinant proteins		
Tri Reagent	Sigma-Aldrich	93289
Dexamethasone, 10mM	G Biosciences	API-04
Insulin, Bovine	Sigma-Aldrich	I0516
3-Isobutyl-1methylxanthine (IBMX)	Sigma-Aldrich	I5879
Glycogen, RNA grade	Thermo Fisher Scientific	R0551
Cell Recovery Solution	Corning	354253
Collagenase Type II	Stem Cell Tech	07419
Y-27632 dihydrochloride (Rho-associated protein kinase inhibitor)	AbMole	M1817
Matrigel, GFR, phenol-free	Corning	356231
Recombinant IL-6 protein, carrier-free	R&D Systems	406-ML
LPS-EB Ultrapure	InvivoGen	Tlrl-3pelps
Recombinant Mouse GDF-15 (CHO-expressed) Protein, CF	R+D Systems	10596-GD-025
Critical commercial assays		
Zymo-seq RiboFree Total RNA Library Kit	Zymo Research Corporation	Zymo R3000
Glycerol Assay Kit	Sigma-Aldrich	MAK117
RNeasy Lipid Tissue Mini Kit	Qiagen	74804
DuoSet ELISA kits	R+D Systems	See methods
Direct-zol™ RNA MiniPrep kit	Zymo Research Corporation	R2050

qScript DNA SuperMix	QuantaBio	95048
SYBRGreen SuperMix	BioRAD	1725122
RNA Clean & Concentrator Kit	Zymo	R1013
Triglyceride Colorimetric Assay Kit	Cayman Chemical	10010303
Deposited data		
Raw and figurative lung tumor organoid, 3 week and 6 week gWAT RNA sequencing data	This paper	Zenodo [10.1101/2024.09.27.615385]
BAM and Fastq files for RNA sequencing data	This paper	deposited to GEO [pending]
Experimental models: Cell lines		
3T3L1-MBX mouse embryonic fibroblast	ATCC	RRID:CVCL_0A20
C2C12 myoblasts	ATCC	RRID:CVCL_0188
Experimental models: Organisms/strains		
$Kras^{LSL-G12D/+}$	The Jackson Laboratory	RRID:IMSR_JAX:008179
$Scgb1a1-CreER^{TM}$	The Jackson Laboratory	RRID:IMSR_JAX:016225
$Lkb1^{fl}$ (Stk11$^{fl/fl}$)	The Jackson Laboratory	RRID:IMSR_JAX:014143
$Ai14$ ($Rosa26$ $tdTomato$)	The Jackson Laboratory	RRID:IMSR_JAX:007908
C57BL/6J	The Jackson Laboratory	RRID:IMSR_JAX:000664
Oligonucleotides		
Primers for RT-PCR, see supplemental table S1.	Custom primers from IDT	N/A
Recombinant DNA		
Software and algorithms		
Halo	Indica Labs	v3.3.2541.133
SPSS	IBM	RRID: SCR_002865 v28
Image J	National Institutes of Health	RRID:SCR_003070
DESeq2		v1.38.3

ggplot2		v3.4.4
R Studio	Posit	2023.09.1
Visual Studio Code	Microsoft	V 1.93.1
S-EFOV	https://github.com/MASILab/S-EFOV	V 1.0.3
Other		
QuPath		RRID:SCR_018257, v0.3.2
MetaboAnalyst	Wishart Research Group	RRID:SCR_015539, v. 5.0
Graphpad Prism	Dotmatics	RRID:SCR_000306, v.10.1.2
STATA	StataCorp	RRID: SCR_012763 V 18.5
Python		RRID: SCR_008394 V 3.7.12

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Michael Toth (michael.toth@med.uvm.edu).

Materials availability

Data and code availability

• RNA-seq raw read counts will be available via the Zenodo online repository.

• Original code used to generate figurative data will be available in the Zenodo repository

and from the lead contact upon request.

• Any additional information required to reanalyze the data reported in this work is

available from the lead contact upon request.

Experimental model and study participant details

Animals

All animal experiments were approved by the University of Vermont Institutional Animal Care

and Use Committee, in compliance with the National Institutes of Health Guidelines for Use and

Care of Laboratory Animals. Mice were housed with sex-matched litter mates and given *ad*

libitum access to standard chow and water, except for a subset of mice that were housed singly

to monitor food intake as specified. Animals were allocated by genotype.

Club cell specific, inducible Kras$^{G12D/+}$ (G12D) mice.

G12D mice (*Scgb1a1-CreER^{TM}-Kras$^{LSL-G12D/+}$*) were generated by crossing *Scgb1a1-*

CreER^{TM} mice with *Kras$^{LSL-G12D/+}$* mice and were maintained on a mixed background. At 4-5

months of age, *G12D* or wild-type *Scgb1a1-CreER^{TM}* (WT) littermates were administered five

consecutive daily intraperitoneal (IP) injections of 20 mg/ml tamoxifen (Sigma-Aldrich, St. Louis,

MO, USA) dissolved in corn oil at a dose of 1 mg tamoxifen/10 g body weight. Body weight was

monitored 2-3 times weekly and food intake was measured in a sub-group of individually

housed mice for 6 weeks. At 6- or 12-weeks post-tamoxifen injections, mice were euthanized by

cervical dislocation under pentobarbital anesthesia. Blood was collected by cardiac puncture in

serum separator tubes and centrifuged to isolate serum, aliquoted, and stored at -80°C. Skeletal

muscles, adipose tissues, heart, liver, and spleen were weighed, flash-frozen in liquid nitrogen,

and stored at -80°C. Lungs were collected for histology or organoid development. One

gastrocnemius, soleus and tibialis anterior muscle were mounted in 7% gum tragacanth, flash-

frozen in liquid nitrogen-cooled isopentane, and stored at -80°C. One inguinal (iWAT) and one

perigonadal (gWAT) fat pad were fixed in 10% formalin and paraffin-embedded.

Kras$^{G12D/+}$ -Stk11$^{-/-}$ Mice.

G12D mice were crossed with *Lkb1$^{fl/fl}$* (*Stk11$^{fl/fl}$*)[81] and Ai14 (*Rosa26-tdTomato*; TD)

reporter mice[82] to generate *Kras$^{G12D/+}$-TD* (*G12D*), *Kras$^{G12D/+}$-Stk11$^{-/-}$-TD* (*KS)* and wild-type

Scgb1a1-CreER^{TM}-TD (WT; *Kras$^{+/+}$*) littermate control mice. At 8-10 weeks of age, *G12D, KS,*

and WT mice were injected (IP) for five consecutive days with tamoxifen (Sigma-Aldrich, St.

Louis, MO, USA) dissolved in corn oil as detailed for *G12D* mice. Body weight and food intake

were monitored thrice weekly for 3.5 weeks, after which mice were euthanized, and blood and

tissues were collected similar to *G12D* mice.

Adeno-Cre and Adeno-5-CMV-Cre mice

Mice harboring *Kras*$^{LSL-G12D}$ (JAX 008179) and *Rosa26*$^{LSL-tdTomato}$ (Ai14, JAX 007914)

alleles were used. Cre recombinase was introduced via oropharyngeal administration of 5 x 10^7

pfu/mouse of adenoviral Cre (Ad5-CMV-Cre or Ad5mSPC-Cre; University of Iowa Viral Vector

Core) diluted in 1X PBS to activate recombination in lung epithelial cells. Initially, mice were

monitored once a week and weighed once every other week. At 12-16 weeks post-infection,

mice were euthanized by IP injection of 200 mg/kg of sodium pentobarbital and tissues collected

for analyses.

Lung Tumor Organoids

Tumor or lung tissue from *G12D* or *WT* mice, respectively, were excised 6 weeks post-

tamoxifen injection, and organoid cultures were developed, as described [83]. Briefly, lung tumors

or tissue were excised and minced in high glucose GlutaMAX DMEM containing 0.1% BSA and

1% penicillin-streptomycin (D-BSA). Minced tissue was incubated with 500 μL collagenase type

II (20 mg/mL) and 10 μL of Rho-associated protein kinase inhibitor (10 μM) on an orbital shaker

at 140 rpm for 1-2 hours at 37°C. Fetal bovine serum (FBS) was added to halt collagenase

digestion and tissues were further homogenized by vigorous pipetting. The tissue suspension

was strained through a 100 μM filter, centrifuged at 8°C, the supernatant aspirated leaving ~0.5

mL, and the pellets resuspended. Tissue pellets were washed twice in D-BSA and centrifuged

at 8° C, and tissue pellets were resuspended in Matrigel (200 μL for a 50 μL pellet). The tissue

suspension (50 μL) was seeded into each well of a 6-well plate in ~10 μL drops. Maintenance of

the oncogenic phenotype was confirmed by assessment of the single nucleotide variant

(c.35G>A) in the *Kras* allele via RNA-seq. For subsequent *in vitro* studies, conditioned medium

(CM) from *G12D* or WT organoids were collected over a 3-day period after organoid passaging.

CM was frozen at -80°C in aliquots for *in vitro* studies. For cell treatment, CM was thawed on ice

directly prior to 25% dilution (1:3) in cell culture maintenance medium.

Cells

3T3-L1-MBX mouse embryonic fibroblasts were cultured and differentiated, as described

[84]. For some experiments, cells were treated with recombinant LPS, IL-6, or GDF-15 for a period

of 24 hours and supernatants assessed by ELISA.

C2C12 myoblasts were cultured as previously described [61]. On Day 7 post-

differentiation, myotubes were treated with 25% (1:3) WT or *G12D* organoid CM in maintenance

medium, replenished daily, for a period of 72 hours. Myotube diameter was assessed using

digital images taken at 4x magnification. An average of 5 diameters were taken for each

myotube of n=5-25 myotubes per field from n=4 random fields were measured using ImageJ

software by an assessor blinded to treatment status.

All cells were used prior to passage 10 and all experiments were independently

replicated 2-3 times.

Human volunteers

Patients recently diagnosed with non-small cell lung adenocarcinoma (NSCLC; n=11; 6

male, 5 female) and age- and sex-matched healthy controls (HC; n=11; 6 male, 5 female) were

recruited, as described [80]. After an overnight fast, patients with NSCLC receiving evidence-

based anti-cancer therapy (82 ± 4 days) and matched controls underwent a fasting blood draw,

and plasma was isolated for measurement of glycerol levels. The study was approved by the

University of Vermont Committee on Research in the Medical Sciences and written informed

consent was obtained from each volunteer before participation. Other data collected from these

study participants are published in our recent clinical work[80].

To understand early, pre-diagnosis changes in muscle and adipose in patients with lung

cancer, we conducted a secondary analysis using body composition data collected during the

first 3 years of the NLST (ClinicalTrials.gov identifier NCT00047385) –a large, longitudinal study

examining the utility of yearly low-dose, helical computed tomography scans vs. standard chest

x-ray for earl lung cancer diagnosis. Participants in the original study were 55-74 years of age

with a cigarette smoking history of at least 30 pack-years who either smoked currently or had

quit smoking within the past 15 years. The computed tomography (CT) arm of the study

included 26,722 participants who were invited for, and 25,139 participants who completed CT

screening examinations at 1-year intervals over the first 3 years, with baseline screening

examinations performed after randomization[76] . Additional details of inclusion and exclusion

criteria have been previously described[76]. The exposed group for our analyses included n=436

lung cancer patients with ≥2 CT scans before diagnosis. For each exposed participant, we

calculated the days elapsed between their most recent scan and the date of lung cancer

diagnosis. To each exposed participant, we matched one participant who was alive and without

a lung cancer diagnosis (*i.e.*, unexposed) on the corresponding number of days since their last

scan (n=436).Baseline group characteristics for lung cancer cases and controls are listed in

Supplemental Table S2. Body composition was assessed using AI analysis of lung screening

computed tomography scans, as described previously[74, 75]. We fit crude and multivariable-

adjusted hierarchical linear models to estimate changes in adipose tissue and muscle depot

size and density, with adjustment for race/ethnicity, BMI, and age, as described in the Statistics

section of the Star Methods. Because of sex differences in body composition in the anatomical

region examined (thoracic), all analyses were stratified by biological sex

Definitions of Analytic Variables from National Lung Screening Trial Data

Our main exposure definition was lung cancer cases vs. non-cases. We further explored

parsing lung cancer cases by stage at diagnosis, yielding an exposure definition of stage 3-4,

stage 1-2, and non-cases. Following visual evaluation of body composition variable

distributions, we applied either log or square root transformations to skeletal muscle index, SAT

index, VAT index, and VAT area to improve normality before use in regression models. Patient

age was categorized for descriptive purposes but modeled in continuous form. The time scale

for longitudinal measurements was calculated as the difference in date of lung cancer diagnosis

(or corresponding date for matched non-cases) and date of each CT screening. Race and

ethnicity were left in fine categories for descriptive analysis, but sparsity in most categories

required simplification to "White, non-Hispanic or Latino" and "Other" for regression modeling.

We calculated body mass index (BMI) from baseline weight and height and placed into standard

categories for descriptive analysis; regression models used continuous BMI.

Method details

Fecal energy content

During Week 6 post-tamoxifen injection, food intake and fecal samples were collected

over a 72-hour period to calculate total energy intake and fecal energy content, as previously

described. Within this 72-hour period, fecal samples were collected every 24 hours and

immediately frozen at -80C. Samples were dried, powdered, and mixed with wheat flour of pre-

measured energy content (10% fecal sample). Samples were shaped into pellets and in

duplicate subjected to complete oxidation under O_2 in a bomb calorimeter (6200 isoperibol

bomb calorimeter, Parr) and calorie content and digestible energy content were calculated.

Tumor burden

H&E sections from formalin-fixed, paraffin-embedded lung tissues were imaged and

analyzed by a board-certified anatomic pathologist (DJS) blinded to the identity of the

specimens. Qupath was used to manually annotate tumor vs normal lung tissue and assess %

tumor burden in μm^2 and expressed relative to total lung area (%). The presence of

adenocarcinomas was confirmed by a second board-certified anatomic pathologist who was not

a member of the research team.

Enzyme-linked immunoassay (ELISA)

For organoid CM and cell supernatant, levels of IL-6 (DY-406), MCP-1 (DY-479),

CXCL1/KC (DY-453) were quantified using DuoSet ELISA kits, according to the manufacturer's

protocol. DuoSet ELISA Kits were also used to measure TNFα (DY-410) and IL-6 Rα (DY1830)

in serum, according to the manufacturer's protocol. Absorbance was read at 570 nm and

background was removed using 450 nm absorbance on a Synergy HTX plate reader and Gen5

software (BioTek Inc, Winooski, VT, USA). Mouse serum cytokines were analyzed by a custom

Luminex-based multiplex assay (R&D Systems) for mouse CXCL1, GDF-15, TNFα, IL-6,

CXCL10, IL-1β, IL-7, IL-10, TNFR-II, IL-12, IL-17, IL-18, LIF, CXCL10, and FGF21, according

to manufacturer instructions. In some instances, cytokines produced in response to *G12D*

organoid CM treatment were expressed as the fold change compared to WT organoid CM

treatment to account for variability in cytokine production across experimental replicates.

Muscle fiber cross-sectional area (CSA)

Transverse cryosections (10 µm) of gastrocnemius and soleus muscles were stained

with hematoxylin and eosin, whole slides were scanned (Leica-Aperio VERSA 8, Leica

Biosystems, Deerfield, IL), and 2-3 representative 500 µm x 500 µm images from each of two

representative levels along the length of the muscle were chosen to quantify the mean muscle

fiber CSA for each mouse using ImageJ.

Adipocyte CSA

Sections (4 µm) of iWAT and gWAT were cut and stained with hematoxylin and eosin to

assess adipocyte size. Whole slides were scanned and 2-3 representative 500 µm x 500 µm

images from each of two levels of the tissue were used to quantify the mean adipocyte CSA for

each mouse using ImageJ.

RNA isolation and qPCR

Soleus and gastrocnemius muscles, liver, iWAT and gWAT depots, and lung tumor

(*G12D*) or lung tissue organoids (WT;1 well of a 6-well plate) were used for RNA isolation. For

muscle and adipose tissues, RNA extraction was conducted after tissues were mechanically

homogenized (Polytron PT2100, Kinematica, Bohemia, NY, USA) in 1 mL of TRI reagent.

Thereafter, 200 µL of chloroform was added, the sample was vortexed, and centrifuged at 4°C.

For adipose tissues, RNA extraction was completed using the RNeasy Lipid Tissue Mini Kit

(Qiagen, Venlo, Netherlands), according to the manufacturer's protocols, with the addition of the

elective DNAse digestion steps. For muscle tissues, supernatants were transferred to a new

tube and equal amounts of isopropanol and high salt solution (1.2 M sodium chloride, 0.8 M

sodium citrate) were added to each sample. For soleus muscle, 0.5 µL of glycogen

(ThermoFisher Scientific, Waltham, MA, USA) was added to the RNA precipitation step to

increase RNA yield. Samples were incubated and then centrifuged at 4°C. Pellets were washed

in 95% ethanol and RNA was reconstituted in nuclease-free water for cDNA synthesis or RNA-

seq.

For cell samples, media was aspirated, and cells were washed twice with Hanks'

Buffered Saline Solution (HBSS). TRI reagent was added to each well of a 12-well plate and

incubated for 2 minutes at room temperature. Lysates were transferred to a 2-mL tube and

chloroform (1:4 chloroform:TRI reagent) was added. Samples were shaken vigorously and

centrifuged at 4°C. Supernatants were removed, and pellets were resuspended in 500 µL of

75% ethanol, vortexed briefly to wash, and centrifuged at 12,000 rcf for 5 minutes at 4°C.

Samples were reconstituted in nuclease-free water for cDNA analysis.

For tissue and cell samples, RNA quality and quantity were assessed (NanoDrop;

ThermoFisher Scientific, Waltham, MA, USA) and RNA was reverse transcribed using qScript

cDNA Supermix. qPCR was performed according to a standard protocol using iTaq Universal

SYBR Green Supermix and validated primer sequences (**Supplemental Table S1**). Target

gene expression was normalized to the average of two housekeeping genes (adipose: *Ppia,*

B2m; muscle: *Gapdh, Ppia*). Results are expressed as $2^{-\Delta\Delta Ct}$ relative to the average of the

housekeeping genes for each respective tissue and all data expressed relative to WT [85].

For organoid RNA isolation for RNA-seq, media was collected from cultured organoids

and the adherent organoids were rinsed with phosphate-buffered saline (PBS). Over ice, Cell

Recovery Solution (Corning, Corning, NY, USA) was added to the cells, cultures were agitated

by repeated pipetting and incubated at 4°C. Samples were transferred to microcentrifuge tubes

and centrifuged at 4°C. Supernatant was removed and organoid pellet was washed in PBS and

centrifugation step was repeated. RNA isolation was conducted using the Direct-zol™ RNA

MiniPrep kit (Zymo Research Corp., Irvine, CA, USA), according to manufacturer's instructions,

with Trizol (Zymo Research Corp, Irvine, CA, USA) and column-based spin steps to remove the

hydrogel matrix. Following RNA isolation, RNA Clean & Concentrator™ kit (Zymo Research

Corp., Irvine, CA, USA) was used, according to the manufacturer's instructions to concentrate

RNA for library preparation. RNA quality and quantity was confirmed by Qubit (ThermoFisher

Scientific, Waltham, MA, USA).

RNA-seq

RNA libraries from isolated organoids were prepared from 500 ng of total RNA using the

Zymo-Seq RiboFree Total RNA Library Kit according to the manufacturer's instructions. RNA-

Seq libraries were sequenced on an Illumina NovaSeq to a sequencing depth of at least 30

million read pairs (150 bp paired-end sequencing) per sample. Data were analyzed using the

Zymo Research RNA-Seq pipeline, which was adapted from nf-core/rnaseq pipeline v1.4.2 [86]

and built using Nextflow [87]. Briefly, quality control of raw reads was conducted using FastQC

v0.11.9. Adapter and low-quality sequences were trimmed from raw reads using Trim Galore!

V0.6.6. Trimmed reads were aligned to the C57BL6/J mouse reference genome using STAR

v2.6.1d . BAM file filtering and indexing was carried out using SAMtools v1.9 [88]. RNA-seq

library quality control was implemented using RseQC v4.0.0 [89] and QualiMap v2.2.2-dev [90].

Duplicate reads were marked using Picard tools v2.23.9 [91]. Library complexity was estimated

using Preseq v2.0.3 [92]. Duplication rate quality control was performed using dupRadar v1.18.0

[93]. Reads overlapping with exons were assigned to genes using featureCounts v2.0.1[94].

Classification of rRNA genes/exons and their reads were based on annotations and

RepeatMasker rRNA tracks from UCSC genome browser when applicable. Differential gene

expression analysis was completed using DESeq2 v1.28.0 [95]. Functional enrichment analysis

were performed using g:Profiler python API v1.0.0 [96], the Broad's GSEA software [97] and the R

package PathfindR [98]. Quality control and analysis results plots were visualized using MultiQC

v1.9.

To confirm that organoid cultures maintained the Kras^{G12D} mutation, BAM files generated

from organoid RNA-seq analyses were indexed with SAMtools v1.9 *[88]*. The presence and

frequency of Kras allelic variants were then visualized and quantified using the Integrated

Genome Viewer (IGV) [99]. Differentially expressed genes (DEGs) corresponding to WT and

G12D lung organoid RNA-seq libraries were identified from raw read counts using DESeq2 and

plotted with ggplot2. Finally, DEGs identified with DESeq2 were filtered for p-values adjusted for

false discovery rate of <0.05 and absolute value of fold change >2 were subsequently cross-

referenced against DEGs identified in the TRACERx *study [55]* to identify overlaps.

For analysis of adipose tissues, total RNA was isolated from gWAT and BAT from WT

and G12D tissue samples collected at 3 and 6 weeks post-tamoxifen injection. RNA libraries

were prepared using NEBNext Ultra II RNA Library Prep Kit for Illumina, following the

manufacturer's protocol. Libraries were sequenced on an Illumina NovaSeq platform at a depth

of 30 million reads per sample (150 bp, paired-end). Resulting reads were aligned to the

GRCm39 reference genome using STAR (v2.7.3.a) and , and gene-level quantification was

performed with featureCounts (v2.0.0). Differential expression analysis was conducted using

DESeq2 (v1.48.1), and gene set enrichment analysis (GSEA) was carried out using the FGSEA

R package (v1.34.0).

Glycerol levels

Glycerol levels of mouse serum, human plasma, 3T3L1 adipocyte supernatants and lung

tissue/tumor organoid culture conditioned media were measured using the Glycerol Assay Kit

according to the manufacturer's instructions and were expressed relative to body weight to

control for CC-related wasting.

Hepatic triglycerides

Liver tissue (350-400 mg) was minced in substrate assay reagent. Hepatic triglyceride

content was determined according to the kit manufacturer's instructions and standardized to the

starting tissue weight of the sample.

Serum metabolomics

Serum samples were thawed on ice and metabolites were extracted using ice-cold 5:3:2

methanol:acetonitrile:water [100]. Mixtures were vortexed for 30 min at 4°C and centrifuged at 4°C.

Samples were randomized then 10 μL of extracts were injected into a Thermo Vanquish UHPLC

system (San Jose, CA, USA) and resolved on a Kinetex C18 column (150 × 2.1 mm, 1.7 μm,

Phenomenex, Torrance, CA, USA) at 450 μL/min through a 5 min gradient from 0 to 100% B

(mobile phases: A = 95% water, 5% acetonitrile, 1 mM ammonium acetate; B = 95%

acetonitrile, 5% water, 1 mM ammonium acetate) in negative ion mode [101]. Solvent gradient: 0-

0.5 min 0% B, 0.5-1.1 min 0-100% B, 1.1-2.75 min hold at 100% B, 2.75-3 min 100-0% B, 3-5

min hold at 0% B. Injections were then repeated for positive ion mode at 450 μL/min through a 5

min gradient from 5 to 95% B (mobile phases: A = water, 0.1% formic acid; B = acetonitrile,

0.1% formic acid) in positive ion mode. Solvent gradient: 0-0.5 min 5% B, 0.5-1.1 min 5-95% B,

1.1-2.75 min hold at 95% B, 2.75-3 min 95-5% B, 3-5 min hold at 5% B . Eluant was introduced

to the mass spectrometer (Thermo Q Exactive) using electrospray ionization. For both negative

and positive polarities, signals were recorded at a resolution of 70,000 over a scan range of 65-

900 m/z. The maximum injection time was 200 ms, microscans 2, automatic gain control (AGC)

3×10^6 ions, source voltage 4.0 kV, capillary temperature 320°C, and sheath gas 45, auxiliary

gas 15, and sweep gas 0 (all nitrogen). Resulting .raw files were converted to .mzXML format

using RawConverter. Metabolites were assigned and peak areas integrated using Maven

(Princeton University) in conjunction with the KEGG database and an in-house standard library

of >600 compounds [100, 102]. Top metabolites were determined by t-test p-value, and data were

normalized to median and auto-scaled in MetaboAnalyst.

UCP-1 detection in paraffin-embedded white adipose tissue

Paraffin-embedded sections were heated (70°C), deparaffinized in xylenes, washed in 100%

ethanol, 95% ethanol, and PBS, and endogenous peroxidases quenched with 10% hydrogen

peroxide in methanol. After washing in PBS, epitope retrieval was conducted at 95°C in citrate

EDTA buffer (10 mM citric acid, 2 mM EDTA, 0.05% Tween 20, pH 6.2). Slides were rinsed in

distilled water and PBS for 5 minutes and then blocked in 10% normal goat serum (Vector

Laboratories S-1000-20) in PBS. Slides were incubated in primary antibody (1:1000 dilution in

PBS) for 1 h at room temperature, rinsed in PBS, and secondary antibody (1:2000 dilution in

PBS) was added. After washing in PBS, 3,3'-Diaminobenzidine (DAB) chromogen was added,

slides were incubated in the dark and then rinsed in tap water. Slides were counter-stained in

hematoxylin for 20 seconds and rinsed again in tap water until clear. Slides were dehydrated in

serial ethanol solutions, mounted, and cover slipped. Slides were scanned at 40x magnification

on a Leica-Aperio VERSA whole slide imager (Leica Biosystems, Wetzlar, Germany).

Images were analyzed using Halo software. The hue of the brown chromogen stain

indicating positive UCP1 was segmented with the Halo module "Area Quantification version

2.2.1". Stain selection was set to 2, stain 1 (hematoxylin) with stain color 0.239, 0.230, 0.144;

stain 2 (UCP1) with stain color 0.139, 0.251, 0.216. Settings for stain detection were as follows:

Hematoxylin markup color (255, 0, 0), hematoxylin Min OD (0.15, 0.15, 0.15), UCP1 markup

color (0, 255, 0), UCP1 Min OD (0.15, 0.15, 0.15). Advanced settings were utilized to account

for the negative space in the images (Min Tissue OD 0.037). Nuclei were counted using Halo

module "Nuclei Detection – ISH IHC v 3.1.4." General stain settings were utilized to detect the

nuclear stain (hematoxylin – 0.438, 0.342, 0.171) and exclusion stain settings (0.110, 0.219,

0.189) were used to guard against non-specifically counting tissue lifting defects as nuclei.

Typical settings on our samples were: Nuclear Contrast Threshold (0.497), Minimum Nuclear

Optical Density (0.07), and Nuclear Size (1,1000). Default settings were used for Membrane

and Cytoplasm Detection, and ISH cell scoring. Exclusion Marker threshold was set to 2. The

Minimum Tissue OD setting was set to 0.003 to account for negative space in the images.

Statistics

Data were assessed for normality prior to statistical analyses. Graphical data are

represented as the mean ± the standard error of the mean (SEM). ELISA data, if non-normally

distributed, were log_{10} transformed prior to statistical analyses and the assumption of normality

confirmed. Differences between WT and *G12D* groups for singular time point measurements

were determined with a Student's T-test, whereas differences over time were determined by

repeated measures analysis of variance (ANOVA) with individual comparisons at each timepoint

if group X time interaction effects were noted. For comparison of WT, *G12D,* and *KS* groups, a

one-way ANOVA with multiple comparisons was used to determine differences between groups.

Differences between these three groups over time were distinguished using a repeated

measures one-way ANOVA with group X time interaction effects. *SPSS* was used for these

analyses*.* Data and statistical analyses were visualized using Graphpad Prism.

For human NLST data analyses, we tabulated baseline summary statistics according to

key demographics and study measurements for the exposed (lung cancer patients) and

847 unexposed (lung cancer-free) arms of the NLST cohort. We visualized crude time trends in body

848 composition measurements using both box plots and distributional dot plots overlaid with local

849 regression trends. We then fit crude and multivariable hierarchical linear models to compare

850 changes in the sum of skeletal muscle (SM), subcutaneous adipose tissue (SAT) and visceral

851 adipose tissue (VAT) area between lung cancer cases and matched non-cases across the fifth,

852 eight, and tenth thoracic vertebral levels, indexed to patient height and tissue attenuation and

853 stratified by sex. All outcomes were regressed on the 3-way interaction between exposure (lung

854 cancer/stage vs. non-case), measurement time (continuous), and participant sex; multivariable

855 models further adjusted for race and ethnicity (dichotomized as non-Latino White vs. all other

856 races and ethnicities due to sparse data), body mass index (standard categories modeled as a

857 factor variable), and age (modeled as a continuous variable). All models included study center

858 and patient ID as random effects to account for clustering by study site and correlation of within-

859 patient measurements, respectively. We used changes in Akaike's information criterion (AIC) to

860 guide model specification. We specified contrasts to estimate p-values for overall differences in

861 body composition slopes by exposure group within sex strata. Modeled data were visualized by

862 plotting marginal predicted values across the range of time values by exposure group and

863 stratified by sex, with standard errors estimated by the delta method. All analyses were

864 performed with Stata v.18.5 (Stata Corporation, College Station, TX).

865 Additional resources

866 None.

References:

1. Fearon, K., et al., *Definition and classification of cancer cachexia: an international consensus.* Lancet Oncol, 2011. **12**(5): p. 489-95.
2. Prado, C.M.M., et al., *Sarcopenia as a determinant of chemotherapy toxicity and time to tumor progression in metastatic breast cancer patients receiving capecitabine treatment.* Clin Cancer Res, 2009. **15**: p. 2920-2926.
3. Miyawaki, T., et al., *Desensitizing Effect of Cancer Cachexia on Immune Checkpoint Inhibitors in Patients With Advanced NSCLC.* JTO Clin Res Rep, 2020. **1**(2): p. 100020. DOI: 10.1016/j.jtocrr.2020.100020.
4. Kurk, S., et al., *Skeletal muscle mass loss and dose-limiting toxicities in metastatic colorectal cancer patients.* J Cachexia Sarcopenia Muscle, 2019. **10**(4): p. 803-813. DOI: 10.1002/jcsm.12436.
5. Dewys, W.D., et al., *Prognostic effect of weight loss prior to chemotherapy in cancer patients.* Am J Med, 1980. **69**: p. 491-497.
6. Reeve, B.B., et al., *Impact of cancer on health-related quality of life of older Americans.* J Natl Compr Canc Netw, 2009. **101**(12): p. 860-868. DOI: 10.1093/jnci/djp123.
7. Braun, D., D. Gupta, and E. Staren, *Quality of life assessment as a predictor of survival in non-small cell lung cancer.* BMC Cancer, 2011. **11**(1): p. 353.
8. Jatoi, A., et al., *Daily activities: exploring their spectrum and prognostic impact in older, chemotherapy-treated lung cancer patients.* Support Care Cancer, 2003. **11**(7): p. 460-464. DOI: 10.1007/s00520-003-0449-5.
9. Braithwaite, D., et al., *Long-term prognostic role of functional limitations among women with breast cancer.* J Natl Compr Canc Netw, 2010. **102**(19): p. 1468-1477. DOI: 10.1093/jnci/djq344.
10. Jones, L.W., et al., *Prognostic significance of functional capacity and exercise behavior in patients with metastatic non-small cell lung cancer.* Lung Cancer, 2012. **76**(2): p. 248-252. DOI: 10.1016/j.lungcan.2011.10.009.
11. Ross, P.J., et al., *Do patients with weight loss have a worse outcome when undergoing chemotherapy for lung cancers?* Br J Cancer, 2004. **90**(10): p. 1905-11. DOI: 10.1038/sj.bjc.6601781.
12. Fearon, K.C.H., D.J. Glass, and D.C. Guttridge, *Cancer cachexia: mediators, signaling and metabolic pathways.* Cell Metab, 2012. **16**: p. 1-14.
13. Bennani-Baiti, N. and D. Walsh, *Animal models of the cancer anorexia-cachexia syndrome.* Support Care Cancer, 2011. **19**: p. 1451-1463.
14. Strassmann, G., et al., *Evidence for the involvement of interleukin 6 in experimental cancer cachexia.* J Clin Invest, 1992. **89**: p. 1681-1684.
15. Baracos, V.E., *Bridging the gap: are animal models consistent with clinical cancer cachexia?* Nat Rev Clin Oncol, 2018. **15**(4): p. 197-198. DOI: 10.1038/nrclinonc.2018.14.
16. Johns, N., N.A. Stephens, and K.C. Fearon, *Muscle wasting in cancer.* Int J Biochem Cell Biol, 2013. **45**(10): p. 2215-29.
17. Ramsey, M.L., et al., *Circulating interleukin-6 is associated with disease progression, but not cachexia in pancreatic cancer.* Pancreatology, 2019. **19**(1): p. 80-87. DOI: 10.1016/j.pan.2018.11.002.
18. Talbert, E.E., et al., *Circulating monocyte chemoattractant protein-1 (MCP-1) is associated with cachexia in treatment-naïve pancreatic cancer patients.* J Cachexia Sarcopenia Muscle, 2018. **9**(2): p. 358-368. DOI: 10.1002/jcsm.12251.
19. Tomasin, R., A.C.B.M. Martin, and M.R. Cominetti, *Metastasis and cachexia: alongside in clinics, but not so in animal models.* J Cachexia Sarcopenia Muscle, 2019. **10**(6): p. 1183-1194. DOI: 10.1002/jcsm.12475

20.   Mueller, T.C., et al., *Molecular pathways leading to loss of skeletal muscle mass in*
cancer cachexia – can findings from animal models be translated to humans? BMC
Cancer, 2016. **16**(1): p. 75. DOI: 10.1186/s12885-016-2121-8.
21.   Ferrer, M., et al., *Cachexia: A systemic consequence of progressive, unresolved*
disease. Cell, 2023. **186**(9): p. 1824-1845. DOI: 10.1016/j.cell.2023.03.028.
22.   Siegel, R.L., et al., *Cancer statistics, 2023.* Cancer J Clin 2023. **73**(1): p. 17-48. DOI:
10.3322/caac.21763.
23.   Zhou, B., et al., *Worldwide burden and epidemiological trends of tracheal, bronchus, and*
lung cancer: A population-based study. eBioMedicine, 2022. **78**: p. 103951. DOI:
10.1016/j.ebiom.2022.103951.
24.   Bozzetti, F., *Forcing the vicious circle: sarcopenia increases toxicity, decreases*
response to chemotherapy and worsens with chemotherapy. Ann Oncol, 2017. **28**(9): p.
2107-2118. DOI: 10.1093/annonc/mdx271.
25.   Baracos, V.E., et al., *Body composition in patients with non–small cell lung cancer: a*
contemporary view of cancer cachexia with the use of computed tomography image
analysis. Am J Clin Nutr, 2010. **91**(4): p. 1133S-1137S. DOI: 10.3945/ajcn.2010.28608C.
26.   Miyawaki, T., et al., *Predicting the efficacy of first-line immunotherapy by combining*
cancer cachexia and tumor burden in advanced non-small cell lung cancer. Thorac
Cancer, 2022. **13**(14): p. 2064-2074. DOI: doi.org/10.1111/1759-7714.14529.
27.   DeWys, W.D., *Studies correlating the growth rate of a tumor and its metastases and*
providing evidence for tumor-related systemic growth-retarding factors. Cancer Res,
1972. **32**(2): p. 374-9.
28.   Lee, J., et al., *Longitudinal changes in skeletal muscle mass in patients with advanced*
squamous cell lung cancer. Thorac Cancer, 2021. **12**(11): p. 1662-1667. DOI:
10.1111/1759-7714.13958.
29.   Rawlins, E.L. and A.K. Perl, *The a"MAZE"ing world of lung-specific transgenic mice.* Am
J Respir Cell Mol Biol, 2012. **46**(3): p. 269-82. DOI: 10.1165/rcmb.2011-0372PS.
30.   Pryce, B.R., et al., *Muscle inflammation is regulated by NF-κB from multiple cells to*
control distinct states of wasting in cancer cachexia. Cell Rep, 2024. **43**(11): p. 114925.
DOI: 10.1016/j.celrep.2024.114925.
31.   Mayers, J.R., et al., *Tissue of origin dictates branched-chain amino acid metabolism in*
mutant Kras-driven cancers. Science, 2016. **353**(6304): p. 1161-5. DOI:
10.1126/science.aaf5171.
32.   Rosigkeit, S., et al., *Definitive evidence for Club cells as progenitors for mutant*
Kras/Trp53-deficient lung cancer. Int J Cancer, 2021. **149**(9): p. 1670-1682. DOI:
10.1002/ijc.33756.
33.   Sutherland, K.D., et al., *Multiple cells-of-origin of mutant K-Ras-induced mouse lung*
adenocarcinoma. Proc Natl Acad Sci U S A, 2014. **111**(13): p. 4952-7. DOI:
10.1073/pnas.1319963111.
34.   Ryden, M., et al., *Lipolysis--not inflammation, cell death, or lipogenesis--is involved in*
adipose tissue loss in cancer cachexia. Cancer, 2008. **113**(7): p. 1695-704. DOI:
10.1002/cncr.23802.
35.   Kir, S., et al., *Tumour-derived PTH-related protein triggers adipose tissue browning and*
cancer cachexia. Nature, 2014. **513**(7516): p. 100-4. DOI: 10.1038/nature13528.
36.   Petruzzelli, M., et al., *A switch from white to brown fat increases energy expenditure in*
cancer-associated cachexia. Cell Metab, 2014. **20**(3): p. 433-47. DOI:
10.1016/j.cmet.2014.06.011.
37.   Rohm, M., et al., *An AMP-activated protein kinase-stabilizing peptide ameliorates*
adipose tissue wasting in cancer cachexia in mice. Nat Med, 2016. **22**(10): p. 1120-30.

38. Bunk, J., et al., *The Futile Creatine Cycle powers UCP1-independent thermogenesis in classical BAT.* Nat Commun, 2025. **16**(1): p. 3221. DOI: 10.1038/s41467-025-58294-4.

39. Ikeda, K., et al., *UCP1-independent signaling involving SERCA2b-mediated calcium cycling regulates beige fat thermogenesis and systemic glucose homeostasis.* Nat Med, 2017. **23**(12): p. 1454-1465. DOI: 10.1038/nm.4429.

40. Zhang, S., et al., *Acute activation of adipocyte lipolysis reveals dynamic lipid remodeling of the hepatic lipidome.* J Lipid Res, 2024. **65**(2): p. 100434. DOI: 10.1016/j.jlr.2023.100434.

41. Arneson-Wissink, P.C., et al., *Hepatic signal transducer and activator of transcription-3 signalling drives early-stage pancreatic cancer cachexia via suppressed ketogenesis.* J Cachexia Sarcopenia Muscle, 2024. **15**(3): p. 975-988. DOI: 10.1002/jcsm.13466.

42. das Neves, R.X., et al., *Cachexia causes time-dependent activation of the inflammasome in the liver.* J Cachexia Sarcopenia Muscle, 2023. **14**(4): p. 1621-1630. DOI: 10.1002/jcsm.13236.

43. Wang, G., et al., *Tumour extracellular vesicles and particles induce liver metabolic dysfunction.* Nature, 2023. **618**(7964): p. 374-382. DOI: 10.1038/s41586-023-06114-4.

44. Flint, T.R., et al., *Tumor-Induced IL-6 Reprograms Host Metabolism to Suppress Anti-tumor Immunity.* Cell Metab, 2016. **24**(5): p. 672-684. DOI: 10.1016/j.cmet.2016.10.010.

45. Pryce, B.R., et al., *Cancer cachexia: involvement of an expanding macroenvironment.* Cancer Cell, 2023. **41**(3): p. 581-584. DOI: 10.1016/j.ccell.2023.02.007.

46. Shiono, M., et al., *An analysis of the relationship between metastases and cachexia in lung cancer patients.* Cancer Med, 2016. **5**(9): p. 2641-8. DOI: 10.1002/cam4.841.

47. Iyengar, P., et al., *Tumor loss-of-function mutations in STK11/LKB1 induce cachexia.* JCI Insight, 2023. **8**(8). DOI: 10.1172/jci.insight.165419.

48. Goncalves, M.D., et al., *Fenofibrate prevents skeletal muscle loss in mice with lung cancer.* Proc Natl Acad Sci U S A, 2018. **115**(4): p. E743-e752. DOI: 10.1073/pnas.1714703115.

49. Rupert, J.E., et al., *Tumor-derived IL-6 and trans-signaling among tumor, fat, and muscle mediate pancreatic cancer cachexia.* J Exp Med, 2021. **218**(6). DOI: 10.1084/jem.20190450.

50. Baltgalvis, K.A., et al., *Interleukin-6 and cachexia in ApcMin/+ mice.* Am J Physiol Regul Integr Comp Physiol, 2008. **294**(2): p. R393-R401. DOI: 10.1152/ajpregu.00716.2007.

51. Bonetto, A., et al., *JAK/STAT3 pathway inhibition blocks skeletal muscle wasting downstream of IL-6 and in experimental cancer cachexia.* Am J Physiol Endocrinol Metab, 2012. **303**(3): p. E410-21. DOI: 10.1152/ajpendo.00039.2012.

52. Strassmann, G., et al., *Evidence for the involvement of interleukin 6 in experimental cancer cachexia.* J Clin Invest, 1992. **89**(5): p. 1681-4.

53. Suriben, R., et al., *Antibody-mediated inhibition of GDF15-GFRAL activity reverses cancer cachexia in mice.* Nat Med, 2020. **26**(8): p. 1264-1270. DOI: 10.1038/s41591-020-0945-x.

54. van Hall, G., et al., *Interleukin-6 stimulates lipolysis and fat oxidation in humans.* J Clin Endocrinol Metab, 2003. **88**(7): p. 3005-10. DOI: 10.1210/jc.2002-021687.

55. Al-Sawaf, O., et al., *Body composition and lung cancer-associated cachexia in TRACERx.* Nat Med, 2023. **29**(4): p. 846-858. DOI: 10.1038/s41591-023-02232-8.

56. Laurens, C., et al., *Growth and differentiation factor 15 is secreted by skeletal muscle during exercise and promotes lipolysis in humans.* JCI Insight, 2020. **5**(6). DOI: 10.1172/jci.insight.131870.

57. Henriques, F., et al., *Toll-Like Receptor-4 Disruption Suppresses Adipose Tissue*
Remodeling and Increases Survival in Cancer Cachexia Syndrome. Sci Rep, 2018. **8**(1):
p. 18024. DOI: 10.1038/s41598-018-36626-3.
58. Machado, A.P., L.F. Costa Rosa, and M.C. Seelaender, *Adipose tissue in Walker 256*
tumour-induced cachexia: possible association between decreased leptin concentration
and mononuclear cell infiltration. Cell Tissue Res, 2004. **318**(3): p. 503-14. DOI:
10.1007/s00441-004-0987-2.
59. Batista, M.L., Jr., et al., *Cachexia-associated adipose tissue morphological*
rearrangement in gastrointestinal cancer patients. J Cachexia Sarcopenia Muscle, 2016.
7(1): p. 37-47. DOI: 10.1002/jcsm.12037.
60. Molfino, A., et al., *Histomorphological and inflammatory changes of white adipose tissue*
in gastrointestinal cancer patients with and without cachexia. J Cachexia Sarcopenia
Muscle, 2022. **13**(1): p. 333-342. DOI: 10.1002/jcsm.12893.
61. Guigni, B.A., et al., *Electrical stimulation prevents doxorubicin-induced atrophy and*
mitochondrial loss in cultured myotubes. American Journal of Physiology-Cell
Physiology, 2019. **317**(6): p. C1213-C1228. DOI: 10.1152/ajpcell.00148.2019.
62. Aberle, D.R., et al., *Results of the Two Incidence Screenings in the National Lung*
Screening Trial. New England Journal of Medicine, 2013. **369**(10): p. 920-931. DOI:
doi:10.1056/NEJMoa1208962.
63. Lake, J.E., et al., *CT Fat Density Accurately Reflects Histologic Fat Quality in Adults*
With HIV On and Off Antiretroviral Therapy. J Clin Endocrinol Metab, 2019. **104**(10): p.
4857-4864. DOI: 10.1210/jc.2018-02785.
64. Arneson-Wissink, P.C., A.M. Ducharme, and J.D. Doles, *A novel transplantable model of*
lung cancer-associated tissue loss and disrupted muscle regeneration. Skelet Muscle,
2020. **10**(1): p. 6. DOI: 10.1186/s13395-020-00225-6.
65. van de Worp, W., et al., *A novel orthotopic mouse model replicates human lung cancer*
cachexia. J Cachexia Sarcopenia Muscle, 2023. **14**(3): p. 1410-1423. DOI:
10.1002/jcsm.13222.
66. Chen, Z., et al., *Non-small-cell lung cancers: a heterogeneous set of diseases.* Nature
Rev Cancer, 2014. **14**(8): p. 535-546. DOI: 10.1038/nrc3775.
67. Judd, J., et al., *Characterization of KRAS mutation subtypes in non–small cell lung*
cancer. Mol Cancer Ther, 2021. **20**(12): p. 2577-2584. DOI: 10.1158/1535-7163.Mct-21-
0201.
68. East, P., et al., *RAS oncogenic activity predicts response to chemotherapy and outcome*
in lung adenocarcinoma. Nature Communications, 2022. **13**(1): p. 5632. DOI:
10.1038/s41467-022-33290-0.
69. Gannavarapu, B.S., et al., *Prevalence and survival impact of pretreatment cancer-*
associated weight loss: A tool for guiding early palliative care. J Oncol Pract, 2018.
14(4): p. e238-e250. DOI: 10.1200/jop.2017.025221.
70. Reed, D.R., A.A. Bachmanov, and M.G. Tordoff, *Forty mouse strain survey of body*
composition. Physiol Behav, 2007. **91**(5): p. 593-600. DOI:
10.1016/j.physbeh.2007.03.026.
71. Babic, A., et al., *Adipose tissue and skeletal muscle wasting precede clinical diagnosis*
of pancreatic cancer. Nat Commun, 2023. **14**(1): p. 4317. DOI: 10.1038/s41467-023-
40024-3.
72. Fouladiun, M., et al., *Body composition and time course changes in regional distribution*
of fat and lean tissue in unselected cancer patients on palliative care--correlations with
food intake, metabolism, exercise capacity, and hormones. Cancer, 2005. **103**(10): p.
2189-98.

73. Snoke, D.B., et al., *Body composition alterations in patients with lung cancer.* Am J
Physiol Cell Physiol, 2025. **328**(3): p. C872-c886. DOI: 10.1152/ajpcell.01048.2024.
74. Xu, K., et al., *AI Body Composition in Lung Cancer Screening: Added Value Beyond
Lung Cancer Detection.* Radiology, 2023. **308**(1): p. e222937. DOI:
10.1148/radiol.222937.
75. Xu, K., et al., *Body composition assessment with limited field-of-view computed
tomography: A semantic image extension perspective.* Med Image Anal, 2023. **88**: p.
102852. DOI: 10.1016/j.media.2023.102852.
76. Aberle, D.R., et al., *Reduced lung-cancer mortality with low-dose computed tomographic
screening.* N Engl J Med, 2011. **365**(5): p. 395-409. DOI: 10.1056/NEJMoa1102873.
77. Mourtzakis, M., et al., *A practical and precise approach to quantification of body
composition in cancer patients using computed tomography images acquired during
routine care.* Appl Physiol Nutr Metab, 2008. **33**(5): p. 997-1006. DOI: 10.1139/h08-075.
78. Li, Z., et al., *Cancer cells depend on environmental lipids for proliferation when electron
acceptors are limited.* Nat Metab, 2022. **4**(6): p. 711-723. DOI: 10.1038/s42255-022-
00588-8.
79. Lien, E.C., et al., *Low glycaemic diets alter lipid metabolism to influence tumour growth.*
Nature, 2021. **599**(7884): p. 302-307. DOI: 10.1038/s41586-021-04049-2.
80. Snoke, D.B., et al., *Skeletal muscle adaptations in patients with lung cancer:
Longitudinal observations from the whole body to cellular level.* J Cachexia Sarcopenia
Muscle, 2023. DOI: 10.1002/jcsm.13332.
81. Nakada, D., T.L. Saunders, and S.J. Morrison, *Lkb1 regulates cell cycle and energy
metabolism in haematopoietic stem cells.* Nature, 2010. **468**(7324): p. 653-8. DOI:
10.1038/nature09571.
82. Madisen, L., et al., *A robust and high-throughput Cre reporting and characterization
system for the whole mouse brain.* Nat Neurosci, 2010. **13**(1): p. 133-40. DOI:
10.1038/nn.2467.
83. Sachs, N., et al., *Long-term expanding human airway organoids for disease modeling.*
EMBO J, 2019. **38**(4). DOI: 10.15252/embj.2018100300.
84. Ather, J.L., et al., *Obese adipose tissue modulates proinflammatory responses of mouse
airway epithelial cells.* Am J Physiol Regul Integr Comp Physiol, 2021. **321**(1): p. R79-
r90. DOI: 10.1152/ajpregu.00316.2020.
85. Livak, K.J. and T.D. Schmittgen, *Analysis of relative gene expression data using real-
time quantitative PCR and the 2(-Delta Delta C(T)) method.* Methods, 2001. **25**(4): p.
402-8.
86. Patel, H., et al., *nf-core/rnaseq: nf-core/rnaseq v3.0 - Silver Shark (Version 3.0).* 2020, Zenodo.
87. Di Tommaso, P., et al., *Nextflow enables reproducible computational workflows.* Nat
Biotechnol, 2017. **35**(4): p. 316-319. DOI: 10.1038/nbt.3820.
88. Danecek, P., et al., *Twelve years of SAMtools and BCFtools.* Gigascience, 2021. **10**(2).
DOI: 10.1093/gigascience/giab008.
89. Wang, L., S. Wang, and W. Li, *RSeQC: quality control of RNA-seq experiments.*
Bioinformatics, 2012. **28**(16): p. 2184-5. DOI: 10.1093/bioinformatics/bts356.
90. García-Alcalde, F., et al., *Qualimap: evaluating next-generation sequencing alignment
data.* Bioinformatics, 2012. **28**(20): p. 2678-9. DOI: 10.1093/bioinformatics/bts503.
91. Institute, B., *Picard Toolkit*. 2019, GitHub Repository.
92. Daley, T. and A.D. Smith, *Predicting the molecular complexity of sequencing libraries.*
Nat Methods, 2013. **10**(4): p. 325-7. DOI: 10.1038/nmeth.2375.

93.   Sayols, S., D. Scherzinger, and H. Klein, *dupRadar: a Bioconductor package for the*
1109   *assessment of PCR artifacts in RNA-Seq data.* BMC Bioinformatics, 2016. **17**(1): p. 428.
1110   DOI: 10.1186/s12859-016-1276-2.
1111   94.   Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program*
1112   *for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-
1113   30. DOI: 10.1093/bioinformatics/btt656.
1114   95.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
1115   *dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550. DOI:
1116   10.1186/s13059-014-0550-8.
1117   96.   Raudvere, U., et al., *g:Profiler: a web server for functional enrichment analysis and*
1118   *conversions of gene lists (2019 update).* Nucleic Acids Res, 2019. **47**(W1): p. W191-
1119   W198. DOI: 10.1093/nar/gkz369.
1120   97.   Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*
1121   *interpreting genome-wide expression profiles.* Proc Natl Acad Sci U S A, 2005. **102**(43):
1122   p. 15545-50. DOI: 10.1073/pnas.0506580102.
1123   98.   Ulgen, E., O. Ozisik, and O.U. Sezerman, *pathfindR: An R Package for Comprehensive*
1124   *Identification of Enriched Pathways in Omics Data Through Active Subnetworks.* Front
1125   Genet, 2019. **10**: p. 858. DOI: 10.3389/fgene.2019.00858.
1126   99.   Robinson, J.T., et al., *Integrative genomics viewer.* Nature Biotechnology, 2011. **29**(1):
1127   p. 24-26. DOI: 10.1038/nbt.1754.
1128   100.   Nemkov, T., K.C. Hansen, and A. D'Alessandro, *A three-minute method for high-*
1129   *throughput quantitative metabolomics and quantitative tracing experiments of central*
1130   *carbon and nitrogen pathways.* Rapid Commun Mass Spectrom, 2017. **31**(8): p. 663-
1131   673. DOI: 10.1002/rcm.7834.
1132   101.   Nemkov, T., et al., *High-Throughput Metabolomics: Isocratic and Gradient Mass*
1133   *Spectrometry-Based Methods.* Methods Mol Biol, 2019. **1978**: p. 13-26. DOI:
1134   10.1007/978-1-4939-9236-2_2.
1135   102.   Nemkov, T., A. D'Alessandro, and K.C. Hansen, *Three-minute method for amino acid*
1136   *analysis by UHPLC and high-resolution quadrupole orbitrap mass spectrometry.* Amino
1137   Acids, 2015. **47**(11): p. 2345-57. DOI: 10.1007/s00726-015-2019-9.

1138

Graphical Abstract

Fig. 1. Characterization of lung club cell-specific, inducible Kras$^{G12D/+}$ (G12D) mice at 12 weeks post-induction.

Figure 2. Effect of lung club cell-specific, inducible Kras$^{G12D/+}$ (G12D) on skeletal muscle and adipose tissue wasting and mRNA expression at 12-weeks post-induction.

Figure 3. Characterization of lung club cell-specific, inducible Kras$^{G12D/+}$ (G12D) murine model of CC at 6-weeks post-induction

Figure 4. *Effect of lung club cell-specific, inducible Kras^{G12D/+} (G12D) on skeletal muscles and adipose tissues 6 weeks post-induction*

Figure 5. *Comparison of lung club cell-specific, inducible Kras$^{G12D/+}$ (G12D-tm) and Kras$^{G12D/+}$-Stk11$^{-/-}$ (KS-tm) murine models of CC.*

Figure 6. *Organoids developed from lung tissue from wild type (WT) and lung tumors from Kras$^{G12D/+}$ (G12D) mice and their characterization.*

Figure 7. Changes in body composition prior to lung cancer diagnoses among patients in the National Lung Screening Trial