Full Title: The p24-family member, TMED9, clears misfolded GPI-anchored proteins from the ER to the Golgi via the Rapid ER Stress-Induced Export pathway

Short Title: TMED9 is required for ER-to-Golgi export of misfolded GPI-APs

Authors: Elsa Ronzier[1] & Prasanna Satpute-Krishnan[1]
[1]Department of Biochemistry and Molecular Biology, Uniformed Services University of the Health Sciences, Bethesda, MD, USA

Abstract:

The p24-family member TMED9 has recently emerged as a player in secretory pathway protein quality control (PQC) that influences the trafficking and degradation of misfolded proteins. However, TMED9's mechanism of action is not well understood. Here we show that TMED9 plays a central role in the PQC of GPI-anchored proteins (GPI-APs) and provide mechanistic insight. Typically, misfolded GPI-APs dissociate from the endoplasmic reticulum (ER)-resident chaperone calnexin and move to the Golgi by the Rapid ER stress-induced Export (RESET) pathway, and are subsequently trafficked to lysosomes via the plasma membrane. We used biochemical and imaging approaches in cultured cells to demonstrate that misfolded GPI-APs dissociate from calnexin and associate with TMED9 during RESET. Depletion of TMED9 with siRNA or its chemical inhibitor, BRD4780, blocked ER-export of misfolded GPI-APs. By contrast, TMED9-depletion did not prevent ER-export of wild type GPI-APs for secretion, indicating a specific role for TMED9 in GPI-AP PQC. Intriguingly, we discovered that acute treatment with BRD4780 induced a shift in TMED9 localization from the ER to the downstream Golgi cisternae and a block to the RESET pathway. Removal of BRD4780 following acute treatment returned TMED9 to the ER where TMED9 was able to associate with the RESET substrate and restore the RESET pathway. Based on these data, we propose a model in which TMED9 participates in RESET by capturing misfolded GPI-APs that are released by calnexin within the ER and conveying them to the Golgi.

Keywords: GPI-anchored proteins, TMED9, TMED2, TMED10, TMP21, p24-family, RESET, BRD4780, secretory pathway, protein quality control, calnexin.

Abbreviations List:

ER	Endoplasmic Reticulum
ERGIC	ER-Golgi Intermediate Compartment
PQC	Protein Quality Control
NRK	Normal Rat Kidney
GPI	Glycosylphosphatidylinositol
GPI-AP	Glycosylphosphatidylinositol-Anchored Protein
PrP	Prion Protein
YFP-PrP*	YFP-tagged Prion Protein C179A
YFP-PrP* NRK	stable transfectants of NRK expressing YFP-PrP*
YFP-PrP WT NRK	stable transfectants of NRK expressing YFP-PrP WT
GFP-CD59 C94S NRK	stable transfectants of NRK expressing GFP-CD59 C94S
ATZ	Alpha-1-antitrypsin Z variant
MUC1-fs	Mucin-1 frame shift mutant
CNX	Calnexin
RESET	Rapid ER Stress-induced Export
ERAD	ER-Associated Degradation
ERLAD	ER-to-Lysosome-Associated Degradation
siTMED9	TMED9 siRNA
LMW band of TMED9	Lower Molecular Weight band of TMED9
HMW band of TMED9	Higher Molecular Weight band of TMED9
YFP	Yellow Fluorescent Protein
GFP	Green Fluorescent Protein
CER	Cerulean
TG	Thapsigargin
BafA1	Bafilomycin A1
3MA	3-Methyladenine
BFA	Brefeldin A
CHX	Cycloheximide
PNGase F or PNG	Peptide-N-Glycosidase F
Endo H	Endoglycosidase H
NA	Neuraminidase
O-G	O-Glycosidase

Introduction

Protein folding is an error-prone process that often generates misfolded and potentially toxic byproducts [1, 2]. To avoid the accumulation and aggregation of misfolded proteins, cells employ "protein quality control (PQC)" systems. PQC systems comprise elaborate networks of chaperones and enzymes that work together to fold and refold, sequester or destroy misfolded proteins [3, 4].

In the case of secretory pathway proteins, PQC systems monitor folding within the early secretory pathway [i.e. the endoplasmic reticulum (ER), ER-Golgi intermediate compartment (ERGIC) and Golgi] before releasing properly folded proteins to their final destination [5]. To manage the diverse topologies of secretory pathway proteins, eukaryotic cells have evolved numerous distinct PQC pathways to capture and clear misfolded products from the early secretory pathway [6]. Misfolded transmembrane or soluble secretory proteins are typically retained in the early secretory pathway and retrotranslocated to the cytosol for proteasomal degradation by the ER-associated degradation (ERAD) pathways [7, 8] or are directly transferred to lysosomes by the ER-to-lysosome-associated degradation (ERLAD) pathways, including selective ER autophagy [9-11]. By contrast, trafficking studies of misfolded GPI-APs, including naturally occurring mutants of prion protein (PrP) and artificial misfolding mutants PrP (C179A) "PrP*" and CD59 (C94S), revealed that misfolded GPI-APs are poor substrates for ERAD and instead exit the ER and access the cell surface prior to being internalized for lysosomal degradation [12, 13].

During steady-state conditions, cells constitutively release misfolded GPI-APs from the ER to the Golgi for eventual turnover in lysosomes [12, 14]. However, the flux of ER to Golgi clearance is dramatically enhanced by the chemical ER stressors, thapsigargin or dithiothreitol, or overexpression of other ER PQC substrates [12, 14]. Thus, we referred to the ER-clearance pathway as "RESET" for "Rapid ER Stress-induced ExporT" [12]. RESET involves dissociation from the ER-resident chaperone calnexin, and association with p24-family members, including TMP21 [12, 14, 15]. The p24-family is composed of type I transmembrane proteins that are conserved between yeast and mammals, form hetero-oligomers, and associate with COPI and COPII coat proteins to mediate vesicular trafficking between the ER and Golgi [16-19]. SiRNA knockdown of p24-family member TMP21 or TMED2, or treatment with chemical inhibitor, brefeldin A, completely blocks ER to Golgi export of misfolded GPI-APs [12, 20], indicating a role for p24-family hetero-oligomers and the conventional secretory pathway in RESET, respectively.

Several recent findings have provided compelling evidence for the involvement of the p24-family member, TMED9, in regulating diverse aspects of PQC in the early secretory pathway [21]. In particular, Dvela-Levitt *et al.* discovered that TMED9 traps the disease mutant of the secretory protein mucin-1, called MUC1-fs, in TMED9-enriched compartments [22]. Importantly, they identified a small molecule, BRD4780, that induces the acute degradation of TMED9 and concurrently allows for the specific degradation of MUC1-fs but not wild type mucin-1 [22]. Additionally, Roberts *et al.* found TMED9, along with TMP21, preferentially associates with the

misfolding Z mutant of alpha-1-antitrypsin (ATZ) over the wild type variant, and is required for ATZ clearance from the ER [23]. Finally, Zavodszky and Hegde discovered that PrP* co-immunoprecipitates with TMED9, along with TMP21 and TMED2, and that partial knockdown of TMED9 partially reduced the ER-export of PrP* [20].

In this study, we sought to define TMED9's role and its mechanism of action in ER-clearance of misfolded GPI-APs by using a combination of biochemical and imaging approaches in cells. We demonstrated that properly folded GPI-APs are able to traffic from the ER to the cell surface regardless of TMED9 expression or knock down. Conversely, inhibiting TMED9 with siRNA or BRD4780 specifically blocked ER-export of misfolded GPI-APs. Critically, we provide evidence that in order to execute its role in RESET, TMED9 must be localized in the ER where it can bind to the RESET substrates. Complete degradation of TMED9 blocks RESET. Similarly, acute mislocalization of TMED9 from the ER to the Golgi blocks RESET. However, relocalization of TMED9 to the ER restores the RESET pathway. Altogether, the results we present here strongly suggest that TMED9, in association with the known RESET factors TMP21 and TMED2, participates in RESET by capturing misfolded GPI-APs that are released by calnexin in the ER and coordinating their transport to the Golgi.

Results

Rationale behind our model substrate, system and strategy to dissect the role of TMED9 in RESET

To dissect the role of TMED9 in RESET, we used a mutant variant of YFP-PrP WT, called "YFP-PrP*" as our primary misfolding model substrate. YFP-PrP WT includes the N-terminal prolactin signal sequence, which drives efficient translocation into the ER [24], and yields a homogenous population of PrP that efficiently folds and traffics to the cell surface [13, 25]. YFP-PrP* is derived from YFP-PrP WT but contains a C179A mutation that disrupts the single disulfide bond in PrP. YFP-PrP* was stably expressed in an NRK cell line "YFP-PrP* NRK" at levels similar to endogenous PrP in mouse brain lysates [12, 14]. Here, we verified our previous observations that at steady-state YFP-PrP* NRK cells expressed a population of proteins that were strongly localized to the ER and poorly localized on the plasma membrane (Supp Figure S1A-B and [12]).

Additionally, to test for the universality of YFP-PrP* as a model misfolded GPI-AP, we used an alternate, unrelated, misfolded GPI-AP model substrate GFP-CD59 (C94S) [12, 14, 15, 26], as a control. GFP-CD59 (C94S) contains a single cysteine mutation that disrupts 1 of 5 possible disulfide bonds. Here we show that GFP-CD59 (C94S) NRK cells expressed more heterogenous populations of GFP-CD59 (C94S), in comparison to YFP-PrP*, that included clearly detectable ER and plasma membrane-localized populations (Supp Figure S1C-D). Thus, we opted to use YFP-PrP* NRK cells as our primary model RESET substrate.

During steady-state conditions, misfolded GPI-APs are exported out of the ER via RESET for subsequent lysosomal turnover [12]. However, various ER-stressors including the induction of new glycoprotein expression or chemical ER-stressors enhance the flux of misfolded GPI-APs through RESET [12, 14, 20]. Thapsigargin (TG)-treatment, in particular, induces the immediate and near-synchronized release of YFP-PrP* or GFP-CD59 (C94S) out of the ER and through the RESET pathway in YFP-PrP* or GFP-CD59 (C94S) NRK cells [12]. Thus, we used TG as a facile tool and our stable NRK cell lines as a tractable system for the induction and study of RESET, respectively.

During RESET, misfolded GPI-APs release calnexin and increase binding to TMED9 and TMP21

We set out to validate and expand upon the previously reported finding that TMED9 associates with GFP-PrP* at steady-state and may be involved in its clearance from the ER via the RESET pathway in TG-treated cells [20]. We confirmed that during steady-state conditions, YFP-PrP* co-immunoprecipitates with TMED9, along with calnexin, TMP21, and TMED2 (Supp Figure S1E), as previously published [12, 20]. Similarly, we found that the alternate model RESET substrate, GFP-CD59 (C94S), co-immunoprecipitated with TMED9, calnexin, TMP21, and TMED2 at steady-state (Supp Figure S1F).

To gain insights into the dynamics of YFP-PrP* protein-interactions during RESET, we correlated live-cell time-lapse imaging experiments with co-immunoprecipitation time courses. The majority of YFP-PrP* molecules was previously shown to have been cleared from the ER to the Golgi via the RESET pathway within 30 minutes of TG-treatment [12]. Here, we verified this point using YFP-PrP* NRK cells that were either co-transfected with CER-CNX, a Cerulean (CER)-tagged version of the ER-resident chaperone calnexin, to mark the ER or CER-GalT to mark the Golgi. Within the first 20 min of TG-treatment, the majority of YFP-PrP* actively trafficked out of the ER into the Golgi leading to a significant decrease in the colocalization of YFP-PrP* with CER-CNX and significant increase in the colocalization of YFP-PrP* with CER-GalT by 20 min (Figure 1A-D). Additionally, we imaged the trafficking of YFP-PrP* during RESET in cells co-transfected with CER-TMED9 and Golgi marker, FusionRed (FusRed)-SiT. Again, within 20 min of TG-treatment, the majority of YFP-PrP* moved into the FusRed-SiT-marked Golgi (Figure 1E-F, Video 1). Intriguingly, TG-treatment did not significantly alter the distribution of total CER-TMED9 across the 0 to 60 min time points, despite the dramatic shift in YFP-PrP* localization, nor did it significantly alter the Pearson's colocalization coefficient between CER-TMED9 and FusRed-SiT from the 0 to 20 min time points (Figure 1E-F; Video 1).

To examine the dynamics of TMED9 and misfolded GPI-AP protein-protein interactions during RESET, we treated YFP-PrP* NRK cells with TG, co-immunoprecipitated YFP-PrP* and analyzed the changes in YFP-PrP*'s association with calnexin, TMED9 and TMP21 over time. During steady-state conditions, YFP-PrP* associates with calnexin, TMED9 and TMP21. However, within 20 min of TG-treatment densitometry measurements of the western blot bands revealed that YFP-PrP*'s association with calnexin declined by 57±9% (Stdev.p, n=3) and concurrently YFP-PrP*'s association with TMED9 and TMP21 increased by 335±25% (Stdev.p, n=3) and 267±91% (Stdev.p, n=3), respectively (Figure 2A-B). From 20 min to 40 min after addition of TG, YFP-PrP*'s association with TMED9 and Tmp21 declined.

Synthesis of the data from Figures 1, Video 1 and Figure 2 demonstrate that within the first 20 min after TG-treatment during the time that YFP-PrP* was actively clearing out of the ER via RESET, YFP-PrP*'s movement out of the ER and into the Golgi correlated with YFP-PrP*'s dissociation from calnexin and increased association with TMED9 and TMP21. Beyond 20 min after TG-treatment, YFP-PrP*'s association with TMED9 and TMP21 declined during the time when most of the YFP-PrP* was moving through and out of the Golgi.

The association of TMED9 with unrelated misfolded proteins during steady-state conditions, shown here and in other reports (Supp Figure S1E-F)[20, 22, 23], may indicate its role as a possible PQC factor. In particular, the dynamic association between TMED9 and YFP-PrP* that is heightened specifically during the period that YFP-PrP* is actively undergoing RESET (i.e. ER-export to the Golgi) implicates TMED9's involvement in the RESET pathway. Therefore, we set out to test whether TMED9 plays an essential role in coordinating RESET of misfolded GPI-APs.

siRNA depletion of TMED9 blocks ER-export of misfolded but not wild type GPI-APs

To test for a role of TMED9 in RESET, we used siRNA to deplete TMED9 levels in YFP-PrP* NRK cells. Our siRNA knockdown conditions resulted in near complete depletion of TMED9 at the population level, as observed by western blot (Figure 3A, quantified in the next section). However, immunofluorescence of TMED9 revealed that, on an individual cell level, siRNA knockdown produced a heterogenous population of siTMED9-treated cells with varying levels of TMED9-depletion that included complete TMED9 knockdown in approximately 65-70% of the cells, and partial knockdown in the remainder (Supp Figure S2A). We examined the impact of TMED9-depletion on RESET at the population level using western blotting and live-cell imaging techniques, and examined the impact on RESET at the individual cell level using immunofluorescence. For the below siTMED9 experiments, we exploited our previous observation that TG-treatment of YFP-PrP* NRK cells induces the rapid clearance of YFP-PrP* out of the ER via RESET, resulting in almost complete degradation of YFP-PrP* within 90 min [12].

First, we examined the effect of TMED9-depletion on TG-induced clearance of YFP-PrP* using western blot. In control siRNA-treated cells, TG induced a 86.3±6.3% (Stdev.p, n=3) decrease in the levels of YFP-PrP* within 90 min (Figure 3A-B, compare lanes 1 and 2). On the other hand, in the siTMED9-treated cells, TG induced a 34.2±10.52% (Stdev.p, n=3) decrease in the levels of YFP-PrP* (Figure 3A-B, compare lanes 3 and 4), indicating a role for TMED9 in the efficient degradation of YFP-PrP*. Additionally, we observed a slight and insignificant increase to 113.88±17.28% (Stdev.p, n=3) in the steady-state levels of YFP-PrP* in the siTMED9 cells when compared to control cells (Figure 3A-B, compare lanes 1 and 3). This demonstrates that at a population level, TMED9-depletion reduces the efficiency of YFP-PrP* clearance.

Second, we examined the impact of TMED9-depletion on TG-induced ER-export of YFP-PrP* by live-cell time-lapse imaging using the YFP-fluorescent patterns as a readout for the localization of YFP-PrP*. Typically, upon TG-induced release of YFP-PrP* from the ER to the Golgi, YFP-PrP* relocalizes from the ER, which appears as a cytoplasm-filling reticulate organelle that encircles the nucleus, to the Golgi, which appears as perinuclear and crescent shaped organelle by confocal microscopy (Supp Figure S1A and Figure 1 and [12]). In control YFP-PrP* NRK cells, addition of TG resulted in the rapid relocalization of YFP-PrP* from the reticulate ER pattern to the perinuclear Golgi pattern within 30 min in 100% of the cells (n=136, Figure 3C-D). On the other hand, in siTMED9-treated YFP-PrP* NRK cells, addition of TG did not alter the ER-localization of YFP-PrP* in the majority (65.6%, n=180) of the cells. Instead, YFP-PrP* remained in the reticulate ER pattern for at least 90 min after TG-treatment (Figure 3C-D). This demonstrates that at a population level, siTMED9-treatment reduces the number of cells with efficient ER-to-Golgi clearance of YFP-PrP*.

Third, we examined the effect of TMED9-depletion on TG-induced YFP-PrP* clearance from the ER using immunofluorescence against endogenous TMED9 to reveal the efficacy of TMED9 knockdown and the ER-resident chaperone, calnexin, as an ER-marker. In control siRNA-treated YFP-PrP* NRK cells, YFP-PrP* localized to the ER at steady-state, but was released into post-ER compartments after 90 min of TG-treatment (Figure 3E, Supp Figure S2A). The idea that YFP-PrP* underwent RESET in control siRNA-treated cells was supported by

Pearson's colocalization analysis, which demonstrated a strong r value for colocalization of YFP-PrP* with calnexin of 0.834 +/- 0.05 (n=14 cells) at steady-state, but dropped significantly to 0.538 ± 0.067 (n=12 cells) after 90 min of TG-treatment (Figure 3F). On the other hand, in siTMED9-treated YFP-PrP* NRK cells in which TMED9 levels were depleted to background levels, YFP-PrP* appeared to remain localized to the calnexin-marked ER despite 90 min of TG-treatment (Figure 3E, Supp Figure S2A). Moreover, in TMED9-depleted cells, Pearson's r value between YFP-PrP* and calnexin was not significantly altered despite TG-treatment, 0.80 ± 0.11 (n=15 cells) at steady-state (TG 0 min) and to 0.84 ± 0.10 (n=14 cells) in thapsigargin-treated cells (TG 90 min) (Figure 3F).

As with YFP-PrP* (Figure 3 and Supp. Figure S2A), siTMED9-treatment appeared to inhibit RESET of the alternate misfolded GPI-AP, GFP-CD59 (C94S) (Supp. Figure S2B-E). In comparison to control scrambled siRNA-treated cells, siTMED9 depleted GFP-CD59 (C94S) NRK cells exhibited (1) inefficient TG-induced clearance of GFP-CD59 (C94S), as shown through western blot analysis (Supp Figure S2B-C), (2) increased steady-state levels of GFP-CD59 (C94S), as shown by western blot (Supp Figure S2B-C), and (3) a significant decrease in the percentage of cells in which GFP-CD59 (C94S) underwent RESET upon TG treatment, as shown by live-cell imaging (Supp Figure S2D-E). The demonstration that TMED9-depletion inhibited ER-export of two unrelated misfolded GPI-APs suggest that TMED9 could play a general role in the ER-export step of the RESET pathway for misfolded GPI-APs.

As a control to test for general effects of TMED9-depletion on ER-export, we used YFP-PrP WT. YFP-PrP WT folds efficiently and traffics via the secretory pathway from the ER to the plasma membrane where it stably resides and is predominantly localized at steady-state [12]. Between control and siTMED9-treated YFP-PrP WT NRK cells, we observed no discernable difference in YFP-PrP WT localization in the post-ER secretory pathway compartments and the plasma membrane. Specifically, to determine whether YFP-PrP WT requires the expression of TMED9 to access post-ER compartments, we fixed the cells and co-stained them for TMED9 expression and endogenous calnexin and counted the number of cells in which YFP-PrP WT localized to the plasma membrane. In 100% of the TMED9-expressing cells (n=28) and TMED9-depleted cells (n=26), YFP-PrP WT visibly localized to the plasma membrane (Figure 3G), demonstrating that TMED9 is not required for ER-export of the wild type variant. Further support for the idea that TMED9 is not required for ER-export of wild type GPI-APs was provided by Pearson's colocalization analysis of YFP-PrP WT with calnexin, which revealed similar r values of 0.325 ± 0.056 for n=22 control siRNA-treated cells and 0.404 ± 0.100 for n=24 siTMED9-treated cells (Figure 3H). This is in marked contrast to the misfolded variant, YFP-PrP* (Figure 3E-F).

Thus, the results from the siTMED9 knockdown experiments indicate that TMED9-expression is required for the ER-export of misfolded GPI-APs. TMED9-depletion did not prevent the wild type GPI-AP from trafficking to the plasma membrane, suggesting that the mechanisms underlying ER-export and post-ER trafficking remained intact. Thus, TMED9 is a key component of the machinery that is specifically required for misfolded GPI-APS to undergo ER-export for subsequent degradation.

Chemical inhibitor of TMED9, BRD4780, inhibits RESET of misfolded GPI-APs

The siTMED9 depletion studies strongly suggest that TMED9 expression is required for RESET. However, TMED9-depletion by knock down or knock out was reported to be associated with a decrease in the levels of other p24/TMED-family proteins, including TMP21 and TMED2 [27, 28]. We replicated this phenomenon here with siTMED9-treated YFP-PrP* NRK cells (Figure 4A-B). As expected, siTMED9-depletion reduced TMED9 expression levels by 91.34% +/- 1.54% (Stdev.p, n=3). However, siTMED9-depletion also resulted in a 62.18±22.95% (Stdev.p, n=3) reduction in TMP21 and 48.26±16.89% (Stdev.p, n=3) reduction in TMED2 levels. The concomitant reduction of known RESET factors, TMP21 and TMED2, exposes the possibility that TMED9's involvement in RESET pathway may be indirect through the stabilization of TMP21 or TMED2. Thus, to gain insight into whether TMED9 plays a direct role in RESET, we opted to use the recently published chemical inhibitor of TMED9, BRD4780, that has been shown to acutely and specifically destabilize TMED9 [22].

To determine the optimal parameters for acute TMED9 inhibition by BRD4780, we used live cell imaging assays to identify the shortest pretreatment time and lowest concentration for BRD4780 that would inhibit TG-induced RESET in 100% of the YFP-PrP* NRK cells. Using YFP-PrP*NRK cells that were cotransfected with ER marker mCherry-Sec61β, we discovered that a 30 min pretreatment of 100 μM BRD4780 was sufficient to completely block TG-induced RESET in 100% (n=34) of the cells. YFP-PrP* to remain in the ER of the YFP-PrP* NRK cells for at least 90 min after the addition of TG in BRD4780-pretreated cells, instead of relocalizing to downstream secretory pathway compartments (Figure 4C). Pearson's colocalization analysis of YFP-PrP* with the ER marker, mCherry-Sec61β, revealed strong colocalization at steady-state that was reduced upon TG-treatment in control cells, but maintained in BRD4780-pretreated cells (Figure 4D). Similar experiments conducted with GFP-CD59 (C94S) NRK cells revealed that in 100% of the cells that were treated with TG alone, the ER-localized population of GFP-CD59 (C94S) underwent RESET (n=33), while in BRD4780-pretreated cells, the ER-population of GFP-CD59 (C94S) remained trapped in the ER in 100% of the cells (n=32) (Supp Figure S3A). As with YFP-PrP* (Figure 4D), colocalization analysis of GFP-CD59 (C94S) with mCherry-Sec61β supported the observation that GFP-CD59 (C94S) underwent TG-induced ER-export in the control cells, but remained in the ER of the BRD4780-pretreated cells (Supp Figure S3B).

Although BRD4780 pretreatment appeared to completely inhibit RESET, it did not prevent the loss of YFP-PrP*. After BRD4780 pretreatment, addition of TG induced a gradual fading of the YFP-fluorescence over the 90 min time course, without any obvious change in the localization (Figure 4C). This suggested that the YFP-PrP* was degraded from the ER by an alternate mechanism. Western blot analysis produced results that were consistent with the live-cell imaging. While TG alone induced degradation of YFP-PrP* within 90 min, the 30 min BRD4780-pretreatment partially inhibited this TG-induced clearance (Figure 4E-F). Altogether, our results demonstrate that although misfolded GPI-APs appear to get degraded by an alternate pathway, BRD4780 completely blocks the TG-induced clearance of YFP-PrP* via the RESET pathway.

Next, we sought to verify that the BRD4780-block on RESET was due to downregulation of TMED9 in NRK cells. In untreated and TG-treated cells, our western blot results for TMED9 revealed a doublet of TMED9 bands between 20 and 25 KDa (Figure 4E). However, upon the 30 min BRD4780-pretreatment prior to the 90 min TG-treatment (BRD4780 + TG), we observed that only the approximately 23.5 KDa lower molecular weight (LMW) band of TMED9 was lost while part of the approximately 24 KDa higher molecular weight (HMW) band remained (Figure 4G-H). This connected the BRD4780-block on RESET with an alteration in TMED9.

To gain more insight into BRD4780's effect on TMED9, we performed a BRD4780-treatment time course and probed for TMED9 by western blot. We observed the gradual loss of the LMW band from 0 to 30 min. After 30 min of BRD4780-treatment, the LMW band decreased by 75.70±1.24 (Stdev.p, n=3) (Figure 4I-J), which corresponds with a complete inhibition of RESET (Figure 4C). By 60 min there was a complete loss of the LMW band, followed by an eventual loss of the HMW band over 180 min of BRD4780-treatment. Critically, at the 30 min timepoint when the LMW TMED9 band disappeared and RESET was completely blocked, TMP21 and TMED2 demonstrated no significant change in expression levels by western blot (Figure 4K-L). These results demonstrate that BRD4780 provides a useful tool to inhibit TMED9's function as a RESET factor, while leaving the previously established RESET factors, TMP21 and TMED2, intact.

The results from these live-cell imaging and western blot based BRD4780 experiments are consistent with the idea that TMED9 plays an essential role in the clearance of YFP-PrP* via RESET. The rapid 30 min BRD4780-induced blockade of RESET, BRD4780-induced molecular weight shift of TMED9 and unaltered TMP21 and TMED2 bands spotlight TMED9 as RESET factor that plays a unique role beyond simply stabilizing the essential RESET factors, TMP21 and TMED2.

BRD4780 alters TMED9 trafficking and glycosylation

The BRD4780-treatment experiments uncovered a connection between the loss of the LMW band of TMED9 and the loss of the cell's ability to carry out RESET (Figure 4C, G-J). Intriguingly, the LMW band disappeared within 30 min of adding BRD4780, while the HMW band persisted for up to 180 min. We considered two explanations for this observation. First, the LMW band may be relatively unstable in comparison to the HMW band, and thus degrade more quickly upon BRD4780-treatment. Second, the LMW band may be converted to the HMW band as a step in TMED9's path toward its eventual degradation. A major clue was presented in Figure 4I. Between 0 and 15 min of BRD4780-treatment, there was an evident and reproducible shift between the balance of the LMW and HMW TMED9 bands that comprised the total pool of TMED9 (Figure 4I). At time 0, total TMED9 was made up of 64.83±3.92% (Stdev.p, n=3) LMW and 34.17±3.92% (Stdev.p, n=3) HMW pools. However, within 15 min of BRD4780-treatment, the total TMED9 pool was comprised of 28.60±1.13% (Stdev.p, n=3) in the LMW and 71.40±1.13% (Stdev.p, n=3) in the HMW band. Given that the total levels of TMED9 remain steady over the first 15 min, this flip in the relative distribution of the TMED9 LMW and HMW populations negates the idea that the HMW band simply persists longer than the LMW band.

Instead, this flip is consistent with the idea that the LMW population converts to the HMW population as TMED9 traffics along the BRD4780-induced degradation pathway for near complete degradation by 180 min (Figure 4I-J).

To delineate the BRD4780-induced degradation pathway for TMED9, we started by assessing which of the two major cellular degradatory systems, lysosomes and proteasomes, are involved. We treated cells for 180 min with BRD4780 alone, BRD4780 with the lysosomal degradation inhibitor, bafilomycin A1 (BafA1 + BRD4780), or BRD4780 with the proteasomal degradation inhibitor, MG132 (MG132 + BRD4780) (Figure 5A-B, lanes 1-3). Bafilomycin A1 inhibited BRD4780-induced degradation of TMED9 HMW species (Figure 5A-B, compare lanes 2 and 3), while MG132 did not (Figure 5A-B, compare lanes 2 and 3), suggesting that the pathway is lysosomal. We next assessed which of the two major categories of trafficking pathways between ER and lysosomes are involved: (1) one of the LC3-II-dependent ER-to-lysosome associated degradation pathways, including autophagy, that are inhibited by 3-methyladenine (3MA) [29], or (2) one of the ER-export to Golgi-dependent pathways that are inhibited with brefeldin A (BFA) [30-32]. We found that while 3-methyladenine did not affect BRD4780-induced degradation (Figure 5A-B, compare lanes 2 and 5), brefeldin A strongly inhibited BRD4780-induced degradation of both the LMW and HMW bands (Figure 5A-B, compare lanes 2 and 6). Brefeldin A causes the entire Golgi to fuse into the ER within 10 min [33]. Therefore, stabilization of both LMW and HMW bands by co-incubating cells with brefeldin A + BRD4780 indicates that the entire population of TMED9 must traffic through the Golgi prior to degradation.

A possible explanation for the BRD4780-induced shift in TMED9's molecular weight is that it reflects the maturation of glycosylation as TMED9 moves from the ER and ERGIC through the Golgi. In eukaryotic cells, the organization of glycosyltransferases across the ER and Golgi allow for predictable sequential glycosylation reactions to occur as secretory pathway proteins traffic through these compartments. Thus, glycosidase digests are a standard procedure to monitor glycoprotein movement through the secretory pathway [34, 35]. Upon entry into the ER, most secretory pathway proteins are N-linked glycosylated [36]. PNGase F cleaves all N-linked glycosylations. As glycoproteins move from the cis to the medial cisternae, they encounter Golgi-resident glycosyltransferases that modify the existing N-linked glycosylations to Endo H resistant forms [34, 35]. As they move through medial and trans cisternae, they may encounter O-linked glycosyltransferases and sialyltransferases, which conjugate sialic acids to either N- or O-linked glycosylations [34, 37, 38]. With this in mind, we hypothesized that the steady-state pool of LMW and HMW TMED9 bands may be a reflection of different glycosylation-states of TMED9 whose ratios are affected by BRD4780-induced of TMED9 ER-to-Golgi transport. We therefore took an open approach to test our hypothesis with a panel of glycosidase digests.

PNGase F and Endo H digests of the YFP-PrP* NRK lysates confirmed that at steady-state TMED9 is N-linked glycosylated with Endo H resistant and sensitive populations (Figure 5C-D, Supp Figure S4A-B). The Endo H resistant populations aligned with the LMW and HMW bands, and the Endo H sensitive population aligned with the PNGase F digested TMED9 core protein. The neuraminidase + O-glycosidase digest appeared to collapse the HMW and LMW doublet into single band with a similar mobility to the LMW band, indicating that the HMW band

corresponded with the subset of TMED9 that accessed the trans cisternae. After BRD4780-treatment for 2 hrs in cells that were pretreated with bafilomycin to prevent TMED9 degradation, the entire TMED9 population ran as a single HMW band that was entirely Endo H resistant and neuraminidase + O-glycosidase sensitive (Figure 5C-D). The following two results suggest that TMED9 does not get O-linked glycosylated, but instead its N-linked glycosylation gets sialylated: (1) PNGase F and PNGase F + neuraminidase + O-glycosidase digests each produced a single band for TMED9 that ran at the exact same molecular weight as the core TMED9 protein (Figure 5C-D), and (2) neuraminidase and neuraminidase + O-glycosidase each produced a single band for TMED9 that ran at the exact same molecular weight as LMW TMED9 in untreated and BRD4780 (30 min)-treated cells (Supp Figure S4A-B). Our results match previously published reports that at steady-state TMED9 obtains an N-linked glycosylation that can be modified with Endo H resistance and neuraminidase sensitivity [17, 39, 40], but does not get O-linked glycosylated [17]. Taken together, these glycosidase digests demonstrate that TMED9 localization shifts from the ER to the Golgi upon BRD4780-treatment in YFP-PrP* NRK cells.

To test whether BRD4780 induces a shift in localization of TMED9 from the ER to the Golgi, we transiently transfected the YFP-PrP* NRK cells with CER-TMED9 to perform live cell dynamic studies. We co-transfected the cells with either the ER marker, mCherry-Sec61β or the Golgi marker, FusionRed-SiT. At steady-state, the CER-TMED9 appeared to colocalize with the ER marker, mCherry-Sec61β (Figure 5E) and the Golgi marker, FusionRed-SiT (Figure 5F). Upon BRD4780-treatment, the ER-localized population of CER-TMED9 appeared to move out of the ER and through the Golgi (Figure 5G-K). In YFP-PrP* NRK cells that were co-transfected with CER-TMED9 and mCherry-Sec61β, CER-TMED9 visibly moved out of the mCherry-Sec61β marked ER (Figure 5G). We quantified the decrease of cumulative CER-TMED9 fluorescence intensity in the ER by establishing an ER mask by segmentation of the mCherry-Sec61β image (Figure 5H), and measuring the cumulative, background-subtracted pixel intensity of CER-TMED9 within the ER at the 0 and 30 min timepoints (Figure 5I). BRD4780 (30 min)-treatment induced a 55.17±14.26% (Stdev.p, n=5) decrease in the ER pool of CER-TMED9 (Figure 5I). Conversely, within 30 min of BRD4780-treatment, we detected an increase in the Pearson's colocalization coefficient r between CER-TMED9 and GM130-stained Golgi in YFP-PrP* NRK cells that were transfected with CER-TMED9 (Figure 5J-K).

The results in Figure 5 suggest that BRD4780 changes TMED9's distribution within the secretory pathway. The glycosidase digests panel for endogenous TMED9 and live-cell imaging results using exogenously expressed CER-TMED9 each represent TMED9 populations that span across the ER, Golgi and intermediate compartments at steady-state, but shift away from the ER and towards the Golgi and downstream compartments within 30 min of BRD4780-treatment. Altogether, the above results demonstrate that RESET inhibition by BRD4780 is associated with alterations in TMED9 trafficking and glycosylation.

TMED9's glycosylation state does not impact its function in RESET

The BRD4780 (30 min)-induced inhibition of RESET coincided with conversion of TMED9 from the LMW form to the fully Endo H resistant, sialylated HMW form and relocalization of CER-TMED9 from the ER to the Golgi (Figure 5). These observations could be explained by two possible scenarios: (1) Only the LMW population of TMED9 is able to carry out RESET or (2) TMED9 must be localized to the ER and/or immediately ER-accessible compartments to carry out RESET. If the first scenario were correct, then TMED9 that was converted to HMW would not be able to support RESET. If the second scenario were correct, then TMED9 that was relocated into compartments downstream of the cis-Golgi would not be able to carry out RESET.

To test the possible explanations, we performed a series of identical incubations in two separate but identical dishes of YFP-PrP* NRK cells. One dish culminated in the collection of the lysates to examine the LMW/HMW states of TMED9. The other dish was assessed by a wide-field fluorescent microscope to test for the cells' ability to carry out TG-induced RESET of YFP-PrP* based on the relocalization of the YFP-PrP* signal from ER (cytoplasm-filling reticular pattern excluding the nucleus) to Golgi (perinuclear crescents). First, we replicated our previous finding shown in Figures 3G. In untreated cells, the majority of TMED9 was in the LMW state and 100% of the cells were able to undergo RESET (Figure 6A, lane 1 and Figure 6B, panel 1). Upon a BRD4780 (30 min)-treatment, TMED9 converted to the HMW population and all of the cells lost their ability to undergo RESET (Figure 6A, compare lanes 1 and 2, and Figure 6B, panels 1 and 2). However, when we introduced a 30 min wash-out step immediately after the BRD4780 (30 min)-treatment, TMED9 remained in the HMW form but the majority of the cells (72%, n=50) regained competency to carry out RESET (Figure 6A, lane 3 and Figure 6B, panel 3). Thus acute inhibition of RESET with BRD4780-treatment is reversible, despite the glycosylation state of TMED9. As shown previously (Figure 3I-J), long-term treatment with BRD4780 for 180 min induced near total TMED9-degradation (Figure 6A, lane 4) and blocked RESET (Figure 6B, panel 4), and was not reversible upon BRD4780 wash-out (Figure 6A, lane 5, Figure 6B, panel 5).

To eliminate the possibility that newly synthesized TMED9 could be contributing to the pool of TMED9 involved in RESET, we performed the above experiments in the presence of the protein translation inhibitor, cycloheximide. Again, in the presence of cycloheximide, BRD4780 (30 min)-treatment induced the conversion of TMED9 to the HMW population and blocked RESET (Figure 6A, compare lanes 6 and 7, Figure 6B, compare panels 6 and 7). Following the cycloheximide + BRD4780 (30 min)-co-treatment, BRD4780 wash-out using medium containing only cycloheximide restored the cell's ability to carry out RESET in the majority of the cells (67%, n=99), but did not alter the HMW state of TMED9 (Figure 6A, lane 8, Figure 6B, panel 8). The finding that simply washing out BRD4780 restored RESET in the presence of a protein translation inhibitor disconnects the glycosylation state of TMED9 from its ability to support RESET. Finally, near complete depletion of the existing pool of TMED9 with cycloheximide + BRD4780 (180 min) co-treatment did not allow for the restoral of RESET with BRD4780 wash-out (Figure 6A, lanes 9 and 10, Figure 6B, panels 9 and 10). This reiterates the requirement for TMED9's presence to carry out RESET.

TMED9 must have access to misfolded GPI-APs in the ER to coordinate RESET

To gain insight into whether TMED9's localization was involved in its ability to execute RESET, we again used exogenously expressed CER-TMED9 as a proxy for TMED9. We co-transfected NRK cells with YFP-PrP*, CER-TMED9 and mCherry-Sec61β. We co-incubated these cells with cycloheximide + BRD4780 and then performed BRD4780 wash-out using medium with only cycloheximide (Figure 7A). Live cell imaging revealed that cycloheximide + BRD4780 (30min) induced export of CER-TMED9 out of the ER, as demonstrated earlier. However during the cycloheximide-only wash-out step, CER-TMED9 appeared to move back into the ER and the cells regained their ability to carry out RESET (Figure 7A, Video 2).

Based on our combined western blot and live-cell imaging results for siRNA depletion or BRD4780-inhibition and depletion of TMED9 (Figures 3, 4, 6, 7A, Supp Figures S2 and S3), we concluded that TMED9 expression was required for RESET, regardless of whether TMED9 was in the LMW or HMW form. The acute inhibition of RESET by BRD4780-treatment was concomitant with the relocalization of TMED9 out of the ER, while restoration of RESET was concomitant with the return of CER-TMED9 to the ER upon BRD4780 wash-out (Figures 4, 6, 7A). Importantly, we observed that YFP-PrP* did not co-traffic with CER-TMED9 out of the ER upon BRD4780-treatment (Figure 7A, Supp Figure S5). Instead, YFP-PrP* appeared to be left behind.

The above observations are consistent with the idea that in order to coordinate RESET, TMED9 requires access to the ER where RESET substrates are largely localized in association with calnexin (Supp Figure S1 and Figure 2, [12, 14]). To directly test this idea, we performed a co-immunoprecipitation time course, pulling down YFP-PrP* from untreated or BRD4780-treated YFP-PrP* NRK cells. In untreated cells, YFP-PrP* co-purified with calnexin and TMED9 (Figure 7B-C, t=0). However, upon BRD4780-treatment, we observed the rapid and complete dissociation of TMED9 from YFP-PrP* between 5 and 30 min (Figure 7B-C). Additionally, TMP21 and TMED2 appeared to dissociate from YFP-PrP* within the same time-frame (Supp Figure S6). Intriguingly, YFP-PrP* steadily maintained association with calnexin, even while the p24-family members rapidly disengaged (Figure 7B-C), which is consistent with the imaging results showing YFP-PrP* remaining in the ER upon BRD4780-treatment (Figure 7A, Supp Figure S5).

Finally, to test our hypothesis that BRD4780 wash-out restores RESET by allowing TMED9 to return to the ER where it can interact with the RESET substrate, we performed co-immunoprecipitations of YFP-PrP* with anti-GFP antibody conjugated beads under the BRD4780-treatment and wash-out conditions (Figure 7D). We prevented the involvement of newly synthesized TMED9 by including cycloheximide in all of the incubations. As a negative control for the co-immunoprecipitation, we included the parental cells with CHX. We treated YFP-PrP* NRK cells as follows: (1) CHX for 30 min, (2) CHX + BRD4780 for 30 min, and (3) CHX + BRD4780 for 30 min followed by a 30 min wash-out in medium containing only CHX. Again, we observed the TMED9 band shift from LMW to HMW upon the CHX + BRD4780 (30 min)-treatment in the input lanes. In the eluate lanes, the CHX (30 min) treatment allowed YFP-

PrP* to pull down both calnexin and TMED9, while the CHX + BRD4780 (30 min)-treatment allowed YFP-PrP* to pull down calnexin but not TMED9. Excitingly, in the eluate lane for CHX + BRD4780 (30 min) followed by the CHX-only wash-out, pull-down of YFP-PrP* co-eluted calnexin and TMED9. After the wash-out, the TMED9 that co-eluted with YFP-PrP* ran at the higher molecular weight, indicating that it had trafficked to downstream compartments of the Golgi and returned to the ER where it could associate with YFP-PrP*.

Taken together, our results demonstrate that TMED9's location in the ER is essential for it to carry out RESET. We showed that TMED9 relocalizes from the ER to the Golgi upon BRD4780 (30 min)-treatment. Upon relocalization out of the ER, TMED9 loses contact with the RESET substrate and its ability to conduct RESET. However if TMED9 is restored to the ER upon BRD4780 wash-out, then TMED9 is able to associate with the RESET substrate and perform its essential function in the RESET pathway. These results presented here are consistent with a model in which TMED9, in conjunction with other p24-family members, is a key constituent of the essential ER-export machinery that captures misfolded GPI-APs released by calnexin and conveys them from the ER to the Golgi (Supp Figure S7).

Discussion

Our observations that BRD4780-treatment phenocopies siTMED9 in blocking RESET of misfolded GPI-APs (Figures 3, 4 and Supp Figure S2) support the argument presented by Dvela-Levitt *et al.* that BRD4780 is a specific inhibitor of TMED9 function [22, 41]. However, the acute action of BRD4780 allowed for a more nuanced understanding of where within the RESET pathway TMED9 functions. TMED9 and p24-family proteins are known to localize between the ER, ERGIC and Golgi [40, 42, 43]. Here, using a combination of glycosidase digests, live cell imaging and co-immunoprecipitations of the RESET substrate, we showed that BRD4780 induced a shift in TMED9 from the ER to the Golgi. The continued application of BRD4780 induced degradation of TMED9 through a brefeldin A- and bafilomycin-sensitive pathway, suggesting that TMED9 traffics through the Golgi to eventually reach lysosomes. Conversely, wash-out of BRD4780 allowed the Golgi-localized TMED9 to be rapidly retrieved back to the ER. Critically, simply shifting TMED9 out of and back into the ER with BRD4780 and wash-out, respectively, caused TMED9 to dissociate and reassociate with RESET substrates and concomitantly disengage and re-engage in the RESET pathway. These findings reveal the potential of BRD4780 as facile tool to manipulate and dissect the relationship between TMED9, misfolded proteins, and chaperone interactions.

Our results demonstrate that BRD4780-treatment causes misfolded GPI-APs, which are membrane bound, to remain in the ER in association with calnexin, while TMED9 leaves the ER through the Golgi for subsequent bafilomycin-sensitive degradation. By contrast, Dvela-Levitt *et al.* reported that BRD4780-treatment induces the misfolded luminal truncation mutant of mucin 1 (MUC1-fs) to traffic out of the ER through the Golgi for subsequent bafilomycin-sensitive degradation [22]. We speculate that the difference in the fates of the misfolded luminal MUC1-fs and misfolded GPI-APs may be based on their interaction with calnexin. Misfolded GPI-APs are well known to interact with the ER-resident chaperone, calnexin [12, 14, 15, 26, 44, 45], which may anchor them to the ER. On the other hand, MUC1-fs may preferentially interact with luminal chaperones, such as calreticulin, BiP, GRP94 and PDI-family members, which are able to cycle between the ER and Golgi. Alternatively, there may be a "tug of war" between TMED9 and calnexin for these substrates, with different outcomes depending on TMED9 or calnexin substrate binding affinities. These ideas remain to be tested. The findings presented here invoke several additional salient questions for future studies regarding the effects of BRD4780. For example, the very first p24-family member to be discovered, canine TMED11 whose closest human homolog is TMED9, was in a co-immunoprecipitation with calnexin **[45]**. This suggests a strong affinity between calnexin and p24-family members, and evokes the question how does BRD4780 impact TMED9 and ER chaperone interactions? Further questions include (1) how does BRD4780 induce rapid re-localization of TMED9 out of the ER to the Golgi and subsequent degradation? (2) how does BRD4780 wash-out allow for the return of TMED9 from the Golgi to the ER and restoration of TMED9's function in RESET? and (3) how does BRD4780 impact the other p24-family members?

Our study unveils new uncharted territories. In particular, given the advent of BRD4780 and the potential development of clinically beneficial derivatives [22, 41], this study revealed an

important new direction of research: determine the fate of misfolded GPI-APs whose clearance via RESET has been blocked. As with siTMED9-treatment, BRD4780-treatment completely blocked RESET. However, upon addition of TG in either BRD4780-treated and siTMED9-treated cells, imaging and western blot results showed a marked reduction in YFP-PrP* levels, suggesting that YFP-PrP* could be degraded by alternate ER-localized degradation pathways (Figures 3 and 4). Similarly, Dvela-Levitt *et al.* previously demonstrated that BRD4780 induced the degradation of another misfolded GPI-AP protein, uromodulin (C126R), that accumulates intracellularly [22]. Blocking RESET and consequent access to the cell surface may be advantageous because cell surface populations of misfolding proteins have the opportunity to engage in various pathological activities. In the case of misfolded prion protein, they have been shown to bind and internalize infectious prion aggregates into cells [46], aggregate within the axonal endosomes of primary neurons and cause neuronal death [47], and aggregate in extracellular deposits in the brain tissue of patients with prion disease [48, 49]. Preventing the extracellular access of misfolded GPI-APs by exploiting BRD4780 to reroute them for an internal route of degradation presents an exciting possible therapeutic intervention that may help ameliorate diseases, including prion diseases.

Materials and Methods

Cell lines

The isolation of the clonal line of Normal Rat Kidney (NRK) cells that we refer to here as parental NRK cells here was described previously [50]. YFP-PrP* NRK, YFP-PrP WT NRK, and GFP-CD59 (C94S) NRK cells are each derived from the NRK cells and their creation were described previously [12, 14]. In short, each of the cell lines were created by transiently transfecting a 10 cm dish with expression plasmids, expanding the transfected cells and selecting them for the stable integrants by using antibiotic resistance, and sorting them for by their fluorescence intensity. As described previously, YFP-PrP* NRK cells were sorted for expression of YFP-PrP* at similar levels to endogenous PrP in mouse brain lysate and YFP-PrP WT NRK and GFP-CD59 (C94S) NRK each express YFP-PrP WT and GFP-CD59 at similar levels to YFP-PrP* [12, 14].

Plasmids

The construction of YFP-PrP WT, YFP-PrP* and GFP-CD59 (C94S) was described in detail [12]. Briefly, YFP-PrP WT includes the N-terminal prolactin signal sequence, which drives efficient translocation into the ER [24], and YFP-PrP* is derived from YFP-PrP WT, but contains a C179A mutation that disrupts the single disulfide bond in PrP. GFP-CD59 (C94S) includes the N-terminal rabbit lactase phlorizin-hydrolase "lactase" signal sequence, which drives efficient translocation into the ER [51-55].

Cerulean (CER)-TMED9 includes the N-terminal rabbit lactase phlorizin-hydrolase signal sequence and its construction was described in detail [23].

Golgi markers include CER-GalT (gift from Jennifer Lippincott-Schwartz, Addgene; plasmid# 11930 [56]) and FusionRed-SiT-15 (gift from Michael W. Davidson, Addgene; plasmid# 56133 [57]).

ER markers include mCherry (mCh)-Sec61β (gift from Jennifer Lippincott-Schwartz, Addgene; plasmid# 90994 [58]) and Cerulean (CER)-calnexin, which we constructed for this study. For CER-calnexin we used the same vector backbone as the SS(lactase)-CER-TMED9 described previously [23]. However, instead of the mature domain of TMED9, we PCR amplified the mature domain of calnexin (human) from a plasmid template obtained from DNASU (DNASU, HsCD00042644). "SS(lactase)" refers to rabbit lactase-phlorizin hydrolase signal sequence. "Mature domain" refers to the ORF that encodes the processed type I transmembrane protein after the N-terminal signal sequence has been cleaved by the signal peptidase. For assembly we used NEBuilder (NEB E5520S). The final ORF was verified to be the following through sequencing analysis with SS(lactase) in all capital letters, followed by linker and CER in lowercase, followed by calnexin mature domain in capital letters:
ATG GAG CTC TTT TGG AGT ATA GTC TTT ACT GTC CTC CTG AGT TTC TCC TGC CGG
GGG TCA GAC TGG GAA TCT CTG CAG TCG ACG GTA CCG CGG GCC CGG GAT CCA

CCG GTC GCC ACC atg gtg agc aag ggc gag gag ctg ttc acc ggg gtg gtg ccc atc ctg gtc gag
ctg gac ggc gac gta aac ggc cac aag ttc agc gtg tcc ggc gag ggc gag ggc gat gcc acc tac ggc
aag ctg acc ctg aag ttc atc tgc acc acc ggc aag ctg ccc gtg ccc tgg ccc acc ctc gtg acc acc ctg
acc tgg ggc gtg cag tgc ttc gcc cgc tac ccc gac cac atg aag cag cac gac ttc ttc aag tcc gcc atg
ccc gaa ggc tac gtc cag gag cgc acc atc ttc ttc aag gac gac ggc aac tac aag acc cgc gcc gag gtg
aag ttc gag ggc gac acc ctg gtg aac cgc atc gag ctg aag ggc atc gac ttc aag gag gac ggc aac
atc ctg ggg cac aag ctg gag tac aac gcc atc agc gac aac gtc tat atc acc gcc gac aag cag aag
aac ggc atc aag gcc aac ttc aag atc cgc cac aac atc gag gac ggc agc gtg cag ctc gcc gac cac
tac cag cag aac acc ccc atc ggc gac ggc ccc gtg ctg ctg ccc gac aac cac tac ctg agc acc cag tcc
aag ctg agc aaa gac ccc aac gag aag cgc gat cac atg gtc ctg ctg gag ttc gtg acc gcc gcc ggg atc
act ctc ggc atg gac gag ctg tac aag gca gga ggc agc CAT GAT GGA CAT GAT GAT GAT GTG
ATT GAT ATT GAG GAT GAC CTT GAC GAT GTC ATT GAA GAG GTA GAA GAC TCA AAA
CCA GAT ACC ACT GCT CCT CCT TCA TCT CCC AAG GTT ACT TAC AAA GCT CCA GTT
CCA ACA GGG GAA GTA TAT TTT GCT GAT TCT TTT GAC AGA GGA ACT CTG TCA GGG
TGG ATT TTA TCC AAA GCC AAG AAA GAC GAT ACC GAT GAT GAA ATT GCC AAA TAT
GAT GGA AAG TGG GAG GTA GAG GAA ATG AAG GAG TCA AAG CTT CCA GGT GAT
AAA GGA CTT GTG TTG ATG TCT CGG GCC AAG CAT CAT GCC ATC TCT GCT AAA CTG
AAC AAG CCC TTC CTG TTT GAC ACC AAG CCT CTC ATT GTT CAG TAT GAG GTT AAT
TTC CAA AAT GGA ATA GAA TGT GGT GGT GCC TAT GTG AAA CTG CTT TCT AAA ACA
CCA GAA CTC AAC CTG GAT CAG TTC CAT GAC AAG ACC CCT TAT ACG ATT ATG TTT
GGT CCA GAT AAA TGT GGA GAG GAC TAT AAA CTG CAC TTC ATC TTC CGA CAC AAA
AAC CCC AAA ACG GGT ATC TAT GAA GAA AAA CAT GCT AAG AGG CCA GAT GCA GAT
CTG AAG ACC TAT TTT ACT GAT AAG AAA ACA CAT CTT TAC ACA CTA ATC TTG AAT
CCA GAT AAT AGT TTT GAA ATA CTG GTT GAC CAA TCT GTG GTG AAT AGT GGA AAT
CTG CTC AAT GAC ATG ACT CCT CCT GTA AAT CCT TCA CGT GAA ATT GAG GAC CCA
GAA GAC CGG AAG CCC GAG GAT TGG GAT GAA AGA CCA AAA ATC CCA GAT CCA
GAA GCT GTC AAG CCA GAT GAC TGG GAT GAA GAT GCC CCT GCT AAG ATT CCA GAT
GAA GAG GCC ACA AAA CCC GAA GGC TGG TTA GAT GAT GAG CCT GAG TAC GTA
CCT GAT CCA GAC GCA GAG AAA CCT GAG GAT TGG GAT GAA GAC ATG GAT GGA
GAA TGG GAG GCT CCT CAG ATT GCC AAC CCT AGA TGT GAG TCA GCT CCT GGA TGT
GGT GTC TGG CAG CGA CCT GTG ATT GAC AAC CCC AAT TAT AAA GGC AAA TGG AAG
CCT CCT ATG ATT GAC AAT CCC AGT TAC CAG GGA ATC TGG AAA CCC AGG AAA ATA
CCA AAT CCA GAT TTC TTT GAA GAT CTG GAA CCT TTC AGA ATG ACT CCT TTT AGT
GCT ATT GGT TTG GAG CTG TGG TCC ATG ACC TCT GAC ATT TTT TTT GAC AAC TTT
ATC ATT TGT GCT GAT CGA AGA ATA GTT GAT GAT TGG GCC AAT GAT GGA TGG GGC
CTG AAG AAA GCT GCT GAT GGG GCT GCT GAG CCA GGC GTT GTG GGG CAG ATG
ATC GAG GCA GCT GAA GAG CGC CCG TGG CTG TGG GTA GTC TAT ATT CTA ACT GTA
GCC CTT CCT GTG TTC CTG GTT ATC CTC TTC TGC TGT TCT GGA AAG AAA CAG ACC
AGT GGT ATG GAG TAT AAG AAA ACT GAT GCA CCT CAA CCG GAT GTG AAG GAA GAG
GAA GAA GAG AAG GAA GAG GAA AAG GAC AAG GGA GAT GAG GAG GAG GAA GGA
GAA GAG AAA CTT GAA GAG AAA CAG AAA AGT GAT GCT GAA GAA GAT GGT GGC ACT
GTC AGT CAA GAG GAG GAA GAC AGA AAA CCT AAA GCA GAG GAG GAT GAA ATT TTG
AAC AGA TCA CCA AGA AAC AGA AAG CCA CGA AGA GAG TAG.

Cell culture, transfection, siRNA knock down, chemical inhibitors

Cell culture conditions included incubation at 37°C, 5% CO2 and 90% humidity in Dulbecco's Modified Eagle Medium (DMEM, Corning, 17-205-CV) containing 10% fetal bovine serum (FBS, Corning 35-011-CV), 1x L-glutamine (Corning, 25-005-CL). Cells used in experiments were never allowed to grow beyond 80% confluency and regularly split every 2 or 3 d using 0.05% Trypsin/0.53 mM EDTA in HBSS (Corning, 25-051-CL). Cells were typically cultured on tissue culture treated plastic (SCBT, sc-251460) and seeded onto #1.5 coverslip glass-bottom dishes (Cell Vis, D35C4-20-1.5-N) for imaging.

Transfections of YFP-PrP* NRK cells were performed at 70% confluency using PolyJet™ In Vitro DNA Transfection Reagent (SignaGen Laboratories, SL100688). For all experiments involving transient transfections, we allowed time for cells that were overexpressing the plasmid-encoded protein to die. We split transiently transfected cells 24 h after transfection onto #1.5 glass-bottom coverslips (Cell Vis, D35C4-20-1.5-N) for imaging experiments. Imaging experiments were performed 48–72 h after transfection.

siRNA knockdown of TMED9 was carried out by two sequential transfections 48 h apart of the ON-TARGETplus SMARTpool Rat Tmed9 siRNA (Dharmacon, L-092937-01-0005) with Dharmafect 2 transfection reagent (Dharmacon, T-2002-02) exactly as recommended by the manufacturer. For imaging experiments, immediately after the second siRNA transfection, cells were transferred to #1.5 coverslips (Cell Vis, D35C4-20-1.5-N), and 24 h later, the imaging experiments were performed. For western blots, immediately after the second siRNA transfection, cells were transferred to 6-well or 12-well tissue culture-treated dishes (Genesee, 25-105MP or 25-106MP).

Chemical inhibitors were added to cell culture in complete DMEM at the following working concentrations: 0.1 µM thapsigargin (diluted from 10mM stock prepared in DMSO; Calbiochem, 586006), 100 µM BRD4780 (also called AGN 192403 hydrochloride; diluted from 50mM stock prepared in water; Tocris, 1072), 10 µM of MG132 (diluted from 20mM stock prepared in DMSO; Calbiochem, 474790), 250 nM bafilomycin A1 (diluted from 1mM stock prepared in DMSO; SCBT, SC-201550B), 10 mM 3-methyladenine (3MA, diluted directly into cell culture medium at 10mM; Tocris, 3977), 2.5 µM brefeldin A (diluted from 25mM stock prepared in DMSO; LC Laboratories, B-8500) and 50 µg/ml cycloheximide (diluted from 10mg/ml stock prepared in water; Calbiochem, 23-97-64). As a control for either thapsigargin or bafilomycin-treatments, the equivalent concentrations of DMSO was added to cells. For BRD4780, 3MA, and CHX, since they were originally diluted in water, control conditions are simply the cell culture medium without the drug.

Co-immunoprecipitation

Cells were cultured in 10 cm dishes to ~70% confluency and treated as described. The entire procedure prior to elution was performed in a 4°C cold room and on ice to stabilize interactions. First, cells were washed twice with 5 mL of cold Low Salt Buffer (50 mM HEPES, pH7.4, 100

mM NaCl, 2 mM MgCl$_2$). Cells were lysed with 1 mL of IP buffer (1% CHAPS, 50 mM HEPES, pH 7.4, 100 mM NaCl, and 2 mM CaCl2) and transferred to a microfuge tube. The lysate was pipetted up and down 20 times through a 200 µl pipet tip to disrupt the cell membranes. To clear the lysates of debris that may clog the column, lysates were subjected to two rounds of centrifugation for 5 min at 16,000 g at 4°C, and supernatants were transferred to new 1.5 mL microfuge tube after each centrifugation. 90 µL of the cleared lysate were set aside to load as "inputs" and the remaining 900 µL were incubated on ice for 15 min with 60 µL of super-paramagnetic eGFP µMACS MicroBead from the µMACS GFP Isolation Kit (Miltenyi, 130-091-288). In parallel, µColumns (Miltenyi, 130-042-701) were placed in the µMacs Separator (Miltenyi, 130-042-602) and equilibrated with 200 µL of IP buffer. At the end of the incubation period, the bead/lysate mixtures were loaded into the columns. Columns were then rinsed five times with 200 µL of IP buffer and washed once with 20 mM Tris Buffer, pH 7.5. After the washes, the experiment was transferred to room temperature and elution was performed by adding 40 µL of boiling elution buffer made of 2x Laemmli Sample Buffer (Bio-Rad, 1610737) and 20mM TCEP (tris(2-carboxyethyl)phosphine), Sigma, 646547). The input samples were denatured with 35 µL of 4X sample buffer (Bio-Rad, 1610747) with 40mM TCEP. The input and eluate samples were further analyzed by western blot.

Western blots and Antibodies

After treatments, cells were washed with cold 1X phosphate-buffered saline (PBS), lysed in a small volume of cold radioimmunoprecipitation assay buffer (RIPA) buffer (50 mM Tris HCl, pH 7.6, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate and 0.1% SDS). Protein lysates were centrifuged at 4°C for 10 min at 15,000 rpm on a microcentrifuge. Supernatants were stored at -20°C and processed as follows. Protein lysate concentrations were determined with Pierce Micro BCA Protein Assay kit (ThermoFisher, 23235). Equal total quantities of protein were denatured using Laemmli sample buffer supplemented with tris(2-carboxyethyl)phosphine (TCEP) and boiled for 5 min before loading into a 12% or 4-20% acrylamide pre-cast SDS-PAGE gels (Bio-Rad, 4561045 or 4568094) for electrophoresis. Proteins were transferred to a PVDF (polyvinylidene difluoride, Bio-Rad, 1704275) membrane using a Trans-Blot Turbo Transfer System (Bio-Rad, 1704150). Membranes were blocked with 5% non-fat milk in Tris-buffered saline with 0.1% Tween-20 (TBS-T + 5% Milk) for 1 hour at room temperature prior to probing with primary antibodies.

The following primary antibodies were added to the TBS-T + 2.5% milk solution at the following concentrations: mouse monoclonal αGFP (Proteintech, 66002-1-IG) at 1:2000, rabbit polyclonal αTMED9 (Proteintech, 21620-1-AP) at 1:2500, rabbit polyclonal αTMP21 (homemade and described previously [12]) at 1:2000, rabbit polyclonal αTMED2 (Proteintech, 11981-1-AP) at 1:2000, mouse monoclonal αCalnexin (Proteintech, 66903-1) at 1:2500, rabbit polyclonal αPrP (Proteintech, 12555-1-AP) at 1:2500, mouse monoclonal αGAPDH antibody (ThermoFisher, MA5-15738) at 1:4000, for 2 hours at room temperature or overnight at 4°C.

Membranes were washed three times with TBS-T and incubated in TBS-T + 5% milk for 1 hour at room temperature with either of the following secondary antibodies at 1:5000: HRP-

conjugated Donkey anti-Rabbit (Sigma Aldrich, NA934-1ML) or HRP-conjugated Goat anti-Mouse (ThermoFisher, A28177). Membranes were washed three times, and briefly incubated in in Pierce™ ECL Western Blotting Substrate (ThermoFisher, 32209) or SuperSignal™ West Femto Maximum Sensitivity Substrate (ThermoFisher, 34096) prior to imaging the chemiluminescence on the ChemiDoc™ Touch Imaging System (Bio-Rad, 17001401).

ImageJ Software (NIH, 1.53q) was used to measure relative pixel densities of bands on unsaturated images collected as eight-bit .tif files from the Chemi-Doc (Bio-Rad). A rectangular ROI sized to enclose the band with the largest area was created and the same ROI was used to measure the integrated density of the rest of the bands. Integrated density values were background subtracted and normalized against "time 0," "untreated" or "control" bands to obtain relative pixel densities. Relative pixel density values from three independently performed experiments were presented as mean ± SD (n = 3), as indicated in the figure legends. Means and SD were calculated and plotted using Microsoft Excel. Quantification details for specific experiments are provided in the adjoining figure legends.

Glycosidase digests

Cell lysates were divided equally into multiple samples that were treated individually with water (no digested control), or with Endoglycosidase H (Endo H, New England Biolabs, P0702), with a2-3,6,8,9 Neuraminidase A (neuraminidase, New England Biolabs, P0722), with a combination of neuraminidase and O-Glycosidase (NA and O-G, New England Biolabs, P0733), with Peptide-N-glycosidase F (PNGase F, New England Biolabs, P0704) or with a combination of O-Glyc+PNGase (O-G + PNGase F) for 3 hrs at 37°C according to the manufacturer's instructions.

To draw the band density histograms associated with the glycosidase digests, ImageJ Software (NIH, 1.53q) was used. Since the Endo H digest in the untreated lysate produced bands spanned the greatest distance from highest molecular weight band to lowest molecular weight band, a rectangular ROI sized to enclose the Endo H bands was created. This same ROI was dragged from lane to lane and the plot graph was generated by selecting the "Analyze," "Gel" and "Plot lanes" menu items in Image J and graphs representing the densitometry intensity versus pixel distance were generated by selecting the "Analyze," "Tools" menu items in Image J to run the Analyze Line Graph.

Live-cell imaging, Immunofluorescence, Antibodies, Nuclear Stain and the Microscope

Cells were seeded on a #1.5 glass bottom dish (Cell Vis, D35-20-1.5-N or D35CA-20-1.5-N) 24 hrs prior to the imaging experiment and imaged at 60-80% confluency. Treatment conditions (e.g. untreated, BRD4780 or TG-treatment) were specified in the figure legend.

For live-cell experiments and time-lapse imaging, cells were imaged in their normal DMEM medium inside of a pre-equilibrated stage top incubator chamber adjusted to 37°C, 5% CO_2 and 90% humidity. To stain nuclei for live cell imaging, cells were incubated for 10 min with NucBlue™ Live ReadyProbes™ Reagent Hoechst 33342 (Thermofisher, R37605).

For immunofluorescence against calnexin (CNX), TMED9, and GM130, cells were washed once with PBS and then fixed and permeabilized with methanol/acetone (3:1) solution for 10 min at -20°C. For immunofluorescence against cell surface GFP, cells were washed once with PBS and then fixed with 4% paraformaldehyde (PFA) in PBS for 15 min at room temperature, but not permeabilized. After the fixation step, cells were washed three times with PBS and blocked for 30 min at room temperature with UltraCruz blocking reagent (SantaCruz; SC-516214) to stain GFP, calnexin, or GM130, or 5% normal goat serum (ThermoFisher, 31872) diluted into PBS to stain TMED9. Primary antibodies were diluted 1:200 in either UltraCruz blocking reagent or 2.5% normal goat serum in PBS buffer and added to the cells for 2 h at room temperature. The following primary antibodies were used: mouse monoclonal αGFP (Proteintech, 66002-1-IG), rabbit polyclonal αTMED9 (Novus, NBP1-31650, lot# 40121), mouse monoclonal αCalnexin (Proteintech, 66903-1), or rabbit polyclonal αGM130 (Novus Bio, NBP2-53420). Cells were washed 3 times with PBS. Secondary antibodies, Goat anti-Rabbit IgG, Alexa Fluor™ 660 (ThermoFisher Scientific, A21074) or Goat anti-Mouse IgG, Alexa Fluor™ 568 (ThermoFisher Scientific, A1103), were diluted at 1:2500 in either UltraCruz blocking reagent or 2.5% normal goat serum in PBS buffer and added to the cells for 1 h at room temperature. Cells were washed 3 times with PBS and incubated with fresh PBS with NucBlue nuclear stain (ThermoFisher, 33342) diluted at 1:20. Images were captured using the Nikon spinning disk (described below).

Images were acquired using a Nikon inverted spinning disk confocal microscope equipped with a Yokogawa CSU-X1 Spinning Disk, EMCCD camera, 60x Plan Apo 1.40 NA oil / 0.13mm WD, Tokai live-cell incubator, and 445 nm, 514, 561, and 647 nm laser lines, and Perfect Focus system to prevent focus drift.

Image analysis

Pearson R coefficients were obtained by the following steps. CER-TMED9 expressing cells were stained for GM130 to mark the Golgi (immunofluorescence described above). Multichannel images were cropped tightly around a single cell. The cell edge was identified by temporarily increasing the contrast of the image and ROIs were manually drawn around the edges of the cell and saved using ROI manager. CER-TMED9 and AlexaFluor 647-GM130, were cropped identically for each individual cell. Pearson's correlation coefficient value *r* was measured using the Coloc2 plugin by selecting Analyze, Colocalization, and Coloc 2 commands in FIJI (ImageJ 1.52p, NIH USA) for the CFP and AlexaFluor 647 channels, for CER-TMED9 and anti-GM130, respectively.

Measuring fluorescence intensity in an image generated ER mask

To measure the total fluorescence intensity of ER-localized CER-TMED9, cells were co-transfected with mCh-Sec61β to mark the ER and CER-TMED9. Multichannel images were collected before and after treatment in RFP (for mCh-Sec61β) and CFP (for CER-TMED9) channels and cropped tightly around a single cell. For each cell, the cell edge was identified by

temporarily increasing the contrast of the image and ROIs were manually drawn around the edges of the cell and saved using ROI manager. ER masks were created from Sec61-mCherry images using Fiji (NIH Image J, 1.52p) by setting the threshold to cover a maximum of the ER compartment by selecting the following Image, Adjust from the menu, and selecting the Threshold command. The threshold setting was used to create and save the ER-mask by selecting Edit, Selection from the menu and selecting the Create Mask command. ER-localized CER-TMED9 fluorescence intensity was extrapolated using the ER mask by selecting Process from the menu and the Image Calculator command with Image 1 set as "ER mark", operation set as "AND" and Image 2 set as "CER-TMED9". The ER-localized CER-TMED9 fluorescence intensity was then measured using Measurement command.

Statistics

Statistical analyses were performed using Prism 9 (GraphPad). Statistical significance was calculated from unpaired t-test with Welch's correction on the mean value of at least 3 independent experiments. The p values that were equal or less than 0.05, were annotated with one star (*); equal or less than 0.01 were annotated with two stars (**); equal or less than 0.001, were annotated with three stars (***); equal or less than 0.0001, were annotated with three stars (****).

Acknowledgements: We thank Alexandra Graninger for technical advice on the glycosidase digests and scientific discussions. We thank Xin Xiang and Yihong Ye for critical feedback on the study and manuscript. This research was supported by National Institute of General Medical Sciences, National Institutes of Health grant no. R01 GM134327 and Uniformed Services University of the Health Sciences startup funds to P. Satpute-Krishnan.

Figure Legends and Figures listed in the order they appear in the text:

Supporting Figure S1: Misfolded GPI-APs localize to the ER and interact with calnexin, TMP21, TMED2, and TMED9.

(A-B) Confocal images of YFP-PrP* NRK cells at steady state. Scale bar represents 10 µm.

(A) Immunofluorescence image of endogenous calnexin (CNX) in a YFP-PrP* NRK cell. The nucleus was stained with Hoechst.

(B) Immunofluorescence image of cell-surface YFP-PrP* using anti-GFP antibody on cells that were not permeabilized. The nucleus was stained with Hoechst.

(C-D) Confocal images of GFP-CD59 (C94S) NRK cells at steady state. Scale bar represents 10 µm.

(C) Immunofluorescence image against endogenous calnexin (CNX) in GFP-CD59 (C94S). The nucleus was stained with Hoechst.

(D) Immunofluorescence of cell-surface GFP-CD59 (C94S) using anti-GFP antibody on cells that were not permeabilized. The nucleus was stained with Hoechst.

(E) Western blots of GFP-tag co-immunoprecipitates from the parental untransfected NRK cells (P) or stably transfected YFP-PrP* NRK cells at steady state. Blots were probed for GFP for YFP-PrP*, and probed for endogenous calnexin (CNX), TMP21, TMED2 and TMED9.

(F) Western blots of GFP-tag co-immunoprecipitates from parental untransfected NRK cells (P) or stably transfected GFP-CD59 (C94S) NRK cells (n=1). Blots were probed for GFP for GFP-CD59 (C94S), and probed for endogenous calnexin (CNX), TMP21, TMED2 and TMED9.

Supplemental Figure S1

Figure 1: During RESET, misfolded GPI-APs traffic from the ER to the Golgi.
(A) Time-lapse imaging sequence collected immediately after thapsigargin (TG)-treatment of a typical YFP-PrP* NRK cell that was co-expressing ER-marker, Cerulean-calnexin (CER-CNX). Scale bar represents 10 μm.
(B) Plot of the average Pearson's r values between YFP-PrP* and CER-CNX for 9 individual cells across the 0 and 20 min time points of time-lapses collected after addition of TG, as shown in (A). Pearson's colocalization coefficients, r, were measured between YFP-PrP* and CER-CNX within the boundaries of the cell. For each image, the boundary of the cell was revealed by temporarily increasing the gain for CER-CNX. The Pearson's r value for individual cells are color-coded.
(C) Time-lapse imaging sequence collected immediately after TG-treatment of a typical YFP-PrP* NRK cell that was co-expressing the Golgi marker, CER-GalT. Scale bar represents 10 μm.
(D) Plot of the average Pearson's r values between YFP-PrP* and CER-GalT for 10 individual cells across the T0 and T20min time points of time-lapses collected after addition of TG, as shown in (C). Pearson's colocalization coefficients, r, were measured between YFP-PrP* and CER-GalT within the boundaries of the cell. For each image, the boundary of the cell was revealed by temporarily maximizing the gain for YFP-PrP*. The Pearson's r value for individual cells are color-coded.
(E) Time-lapse images collected immediately after TG-treatment of a typical YFP-PrP* NRK cell that was co-expressing CER-TMED9 and FusionRed (FusRed)-SiT as a Golgi-marker. This figure panel is also provided as Video 1. Scale bar represents 10 μm.
(F) Plot of the average Pearson's r values between YFP-PrP* and FusRed-SiT or CER-TMED9 and FusRed-SiT, as indicated, for 6 individual cells across the 0 and 20min time points of time-lapses collected after addition of TG, as shown in (E). Pearson's colocalization coefficients, r, were measured between YFP-PrP* and FusRed-SiT or CER-TMED9 and FusRed-SiT, within the boundaries of the cell. For each data point at 0 and 20 min time points, the boundary of the cell was revealed by temporarily maximizing the gain for YFP-PrP*. The Pearson's r value for individual cells are color-coded.

Figure 1

Figure 2: During RESET, misfolded GPI-APs dissociate from calnexin and increase association with TMED9 and TMP21.

(A) Representative western blots of GFP-tag co-immunoprecipitations from the parental untransfected NRK cells (P) or stably transfected YFP-PrP* NRK cells (n=3 biological replicates). YFP-PrP* NRK cells were treated with thapsigargin (TG) and collected for co-immunoprecipitation at the indicated time points. In addition to using anti-GFP antibody to detect co-immunoprecipitation of YFP-PrP*, blots were probed for calnexin (CNX), TMED9 and TMP21. "L" indicates the lane that the ladder was loaded in.

(B) Bar graph representing the mean band intensity for CNX, TMED9 and TMP21 in the eluates from 3 independently performed experiments as shown in (A). For each time point, CNX, TMED9 and TMP21 eluate band intensities were double normalized. First, they were normalized by the band intensity of eluate YFP-PrP*. Second, they were normalized against eluate time 0 band intensity of each protein probed. Error bars represent standard deviations of the mean of 3 independent experiments. Symbols were coded for each experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated $p<0.05$ and ** indicated $p<0.01$.

Figure 2

Figure 3: siRNA depletion of TMED9 blocks ER-export of misfolded GPI-APs but not wild type GPI-APs.

(A) Representative western blots of control (siTMED9 –) or siTMED9-treated (siTMED9 +) YFP-PrP* NRK cells that were either untreated (TG –) or treated with TG for 90 min (TG +) (n=3 biological replicates). Blots were probed for TMED9, PrP to detect YFP-PrP* and GAPDH. Each lane was numbered to facilitate cross-referencing with the quantification shown in (B).

(B) Bar graph representing the mean band intensity for YFP-PrP* from 3 biological replicates as shown in (A). For each condition, YFP-PrP* band intensities were double normalized. First, they were normalized by the band intensity of GAPDH. Second, they were normalized against the untreated control ("siTMED9 – TG – ") band intensity. Error bars represent standard deviation of the mean. Symbols were coded for each independently performed experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated p<0.05 and ** indicated p<0.01. The bars were numbered 1-4 to facilitate cross-referencing with the representative blot in (A).

(C) Time-lapse images of control (CTL) or TMED9 siRNA (siTMED9)-treated YFP-PrP* NRK cells. Prior to starting the time-lapses, nuclei were stained with Hoechst. Image-collection was started immediately after the addition of thapsigargin (TG). Scale bar represents 20 µm**.**

(D) Percentage of cells as represented in (C) with a perinuclear Golgi-localized pattern of YFP-PrP* after 30 min of TG treatment. The bar graph represents 3 biological replicates. Symbols were coded for each independently executed experiment. The number of cells analyzed for each biological replicate are as follows (CTL: triangle 29, diamond 56, circle 51; siTMED9: triangle 69, diamond 58, circle 61). Error bars represent standard deviation of the mean. Statistics were calculated from unpaired t-test with ** indicated p<0.01.

(E) Immunofluorescence images of endogenous TMED9 in control siRNA or siTMED9-treated YFP-PrP* NRK cells that were depleted for TMED9. For each condition, cells were treated with thapsigargin (TG) for 0 min or 90 min before fixation and staining for TMED9 and ER-resident chaperone, calnexin (CNX). Nuclei were stained with Hoechst. Larger fields of views for these panels and additional data for these experiments are presented in Supp. Figure S2A. Scale bar represents 10 µm.

(F) Plot of the average Pearson's r values between YFP-PrP* and endogenous calnexin (CNX) for control (CTL) siRNA or siTMED-treated cells at 0 min or 90 min after thapsigargin addition, as shown in (E). The number of cells analyzed for each condition are listed as follows: CTL siRNA 0 min post-TG-treatment (n=14), CTL siRNA 90 min post-TG-treatment (n=12), siTMED9 0 min post-TG-treatment (n=15), siTMED9 90 min post-TG-treatment (n=14). Pearson's colocalization coefficients, r, were measured between YFP-PrP* and CNX, within the boundaries of the cell as defined by the outline of the CNX staining.

(G) Immunofluorescence images of endogenous TMED9 in control (CTL) or TMED9 siRNA (siTMED9)-treated YFP-PrP WT NRK cells that were depleted for TMED9. Cells were fixed and stained for TMED9 and ER-resident chaperone, calnexin (CNX) at steady-state conditions. Nuclei were stained with Hoechst. Scale bar represents 10 µm.

(H) Plot of the average Pearson's r values between YFP-PrP WT and endogenous calnexin (CNX) for control (CTL) siRNA or siTMED-treated cells at steady-state, as shown in (G). The number of cells analyzed for each condition are listed as follows: CTL siRNA (n=22) and siTMED9 (n=24). Pearson's colocalization coefficients, r, were measured between YFP-PrP* and CNX, within the boundaries of the cell as defined by the outline of the CNX staining.

Figure 3

Supporting Figure S2: TMED9 is required for RESET of misfolded GPI-APs.

(A) Immunofluorescence image stained for endogenous TMED9 and CNX in control YFP-PrP* NRK (CTL) or siTMED9-treated YFP-PrP* NRK cells (siTMED9 cells) at steady state (TG 0min) after 90 min of TG-treatment (TG 90 min). Nuclei were stained with Hoechst. For the siTMED9-treated cells, large white arrows point to cells with only partial TMED9 knockdown. Scale bar represents 10 μm. Scale bar represents 10 μm.

(B) Representative western blots of either control (CTL) or TMED9 siRNA (siTMED9)-treated GFP-CD59 (C94S) NRK cells that were either untreated (TG –) or treated with 90 min TG (TG +) (n=3 biological replicates).

(C) Bar graph representing the mean band intensity for GFP-CD59 (C94S) from 3 biological replicates as shown in (B). For each condition, GFP-CD59 (C94S) band intensities were double normalized. First, GFP band intensity was normalized against the band intensity of GAPDH. Second, the GFP band intensity was normalized against the untreated control ("siTMED9 – TG – ") band intensity. Error bars represent standard deviation of the mean. Symbols were coded for each independently performed experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated p<0.05 and ** indicated p<0.01. **(D)** Time-lapse images of control (CTL) or TMED9 siRNA (siTMED9) GFP-CD59 (C94S) NRK cells. Image-collection was started immediately after the addition of thapsigargin (TG). Scale bar represents 20 μm.

(E) Percentage of cells showing a perinuclear-pattern for GFP-CD59 (C94S), indicative of Golgi-localization, after 30 min of TG-treatment. These data are derived from 4 independent experiments (n=4). Symbols were coded for each independently performed experiment. The number of cells analyzed for each biological replicate are as follows (CTL: triangle 37, diamond 59, circle 18, square 54; siTMED9: triangle 73, diamond 93, circle 28, square 64). Error bars represent standard deviation of the mean. Statistics were calculated from unpaired t-test with Welch's correction with ** indicated p<0.01.

Supplemental Figure S2

Figure 4: BRD4780 modifies TMED9 and inhibits RESET of misfolded GPI-APs.

(A) Representative western blots of either non-targeting siRNA (CTL) or TMED9 siRNA (siTMED9)-treated YFP-PrP* NRK cells. Membranes were probed for TMP21 and GAPDH or TMED2 and GAPDH. (n=3 biological replicates).

(B) Bar graph representing the mean band intensity for TMED9, TMP21 or TMED2 from 3 independently performed experiments as shown in (A). For each condition, TMED9. TMP21 or TMED2 band intensities were double normalized: first against the band intensity of GAPDH, second against the CTL band intensity. Error bars represent the standard deviation of the mean. Symbols were coded for each independently performed experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated p<0.05.

(C) Time-lapse images of YFP-PrP* NRK cells that were co-transfected with mCh-Sec61β. Cells were either not pretreated (Control) or pretreated with BRD4780 (BRD pretreat) for 30 min. Image collection was started immediately after the addition of thapsigargin (TG). Scale bars represent 10 µm.

(D) Plot of the average Pearson's r values between YFP-PrP* and mCh-Sec61β. For Control vs. BRD4780-pretreated conditions, as described in (C), 10 time-lapses of individual cells were analyzed for the 0 and 30 min time points. Pearson's colocalization coefficients, r, were measured between YFP-PrP* and mCh-Sec61β within the boundaries of the cell. For each data point, the boundaries of each cell at 0 and 30min time points were revealed by temporarily maximizing the gain for YFP-PrP*.

(E) Representative western blots of YFP-PrP* NRK cells that were untreated (CTL), TG-treated for 90 min (TG) or BRD4780 (30 min)-pretreated and harvested after 90 min of TG treatment (BRD+TG) (n=3 biological replicates). Blots were probed for TMED9, PrP (YFP-PrP*) and GAPDH.

(F) Average of normalized YFP-PrP* band intensity from 3 independently performed experiments as shown in (E). For each condition, YFP-PrP* band intensities were double normalized. First, they were normalized against the band intensity of GAPDH. Second, they were normalized against the CTL band intensity. Error bars represent standard deviations of the mean. Statistics were calculated from unpaired t-test with Welch's correction with *** indicated p<0.001.

(G) Representative western blots of YFP-PrP* NRK cells that were either untreated control (CTL) or treated with BRD4780 for 30 min (30' BRD). Blots were probed for TMED9, TMP21, and TMED2 and their respective GAPDH (n=3 biological replicates). TMED9 protein profile shows 2 distinct bands, one lower band (orange arrow, LMW) and a higher band (black arrow, HMW).

(H) Average of normalized High Molecular Weight (HMW) and Low Molecular Weight (LMW) TMED9 band intensity from 3 independently performed experiments as shown in (G). For each condition, TMED9 band intensities were double normalized. First, they were normalized against the band intensity of GAPDH. Second, they were normalized against the CTL intensity of each protein probed. Orange bars represent the quantification of the lower molecular weight (LMW) band intensity and gray bars represent the higher molecular weight (HMW) band intensity. Error bars represent standard deviations of the mean. Symbols were coded for each independently performed experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated p<0.05, and *** p<0.001.

(I) Representative western blots of endogenous TMED9 from YFP-PrP* NRK cells after 180 min of BRD-treatment (n=3 biological replicates). Blots were probed for TMED9 and GAPDH. TMED9 protein profile shows 2 distinct bands, one LMW (orange arrow) and a HMW (black arrow).

(J) Average of normalized HMW and LMW TMED9 band intensity from 3 independently performed experiments as shown in (I). For each time point TMED9 band intensities were double normalized: first by the band intensity of GAPDH and second against the BRD4780 0 min band intensity. Orange bars representing the measurement for the LMW of TMED9 and gray, the HMW. Error bars represent standard deviations of the mean. Symbols were coded for each independently performed experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated p<0.05 and ** p<0.01.

(K) Representative western blots of TMP21, TMED2 and their representatives GAPDH either control (CTL) or BRD4780 (30 min)-treated YFP-PrP* NRK cells (n=3 biological replicates).

(L) Average of normalized TMP21 and TMED2 band intensities from 3 independently performed experiments as shown in (K). For each condition, TMP21 or TMED2 band intensities were double normalized: first against the band intensity of GAPDH, second against the CTL band intensity. Error bar represents standard deviation of the mean. Symbols were coded for each independently performed experiment. Statistics were calculated from unpaired t-test with Welch's correction with * indicated p<0.05.

Figure 4

Supporting Figure S3: Chemical inhibitor of TMED9, BRD4780, inhibits RESET.
(A) Representative time-lapse imaging of stably transfected GFP-CD59 C94S NRK cells in control condition (CTL) or pretreated with BRD4780 for 30 min (pretreated with BRD4780). Image-collection was started immediately after the addition of thapsigargin (TG). Scale bars represent 10 µm.
(B) Plot of the average Pearson's r values between GFP-CD59 (C94S) and mCh-Sec61β for Control vs. BRD4780-pretreated conditions, as described in (A). For Control and BRD4780-pretreated conditions, 10 and 9 time-lapses of individual cells, respectively, were analyzed for the 0 and 30 min time points. Pearson's colocalization coefficients, r, were measured between GFP-CD59 (C94S) and mCh-Sec61β within the boundaries of the cell. For each data point, the boundaries of the cells were revealed by temporarily maximizing the gain for mCh-Sec61β.

Supplemental Figure S3

Figure 5: BRD4780 alters TMED9 trafficking and glycosylation.

(A) Representative western blots of YFP-PrP* NRK cells that were treated as listed below (n=3 biological replicates). Untreated (Control), treated with BRD4780 for 180 min (BRD), pretreated with bafilomycin A1 (BAF) for 180 min and co-treated with BRD for 180 min (BAF+BRD), pretreated with MG132 for 180 min and co-treated with BRD for 180 min (MG132+BRD), pretreated with 3-methyladenine (3MA) for 180 min and co-treated with BRD for 180 min (3MA+BRD) or pretreated with BFA for 180 min and co-treated with BRD for 180 min (BFA+BRD). Blots are probed for endogenous TMED9 and GAPDH.

(B) Bar graph representing the mean band intensity for TMED9 from 3 biological replicates as shown in (A). For each condition, TMED9 band intensities were double normalized first by the band intensity of GAPDH and second within individual experiments using control band intensity. Orange bars represent the measurement for the low molecular weight (LMW) band of TMED9 and gray bars represent the measurement of the high molecular weight (HMW) band of TMED9. Error bars represent standard deviations of the mean. Symbols were coded for individual experiments. Statistics were calculated from unpaired t-test with Welch's correction with * indicated $p<0.05$ and ** $p<0.01$.

(C) Western blots of digested protein lysates of stably transfected YFP-PrP* NRK cells either in control condition (Control) or after a pretreatment of 2 h of bafilomycin followed by co-incubation with 2 h of BRD4780 (BAF+BRD). Protein lysates undigested (water) or digested with Endoglycosidase H (Endo H), Neuraminidase+O-Glycosidase (NA+O-Glyc.), Peptide-N-glycosidase F (PNG) or NA+O-Glyc+PNGase (NA+O-G+PNG) for 3 h at 37°C (n=1 experiment). Blots were probed for TMED9 and calnexin (CNX).

(D) Western blot band graphics obtained by first analyzing the plot lanes and second generating the line graphs of pre-selected regions of interest (ROI) from the western blot for control samples (left) or BAF+BRD samples (right). Black arrow represents the HMW form, orange arrow represents the LMW and purple, the unglycosylated form of TMED9**.**

(E) (left) Fluorescence image of a typical untreated YFP-PrP* NRK cell that was co-transfected with CER-TMED9 and ER-marker, mCherry-Sec61β. Scale bar represents 10 μm. (right) Magnified panel of the stroked region on the right. Scale bar represents 1 μm.

(F) (left) Fluorescence image of a typical untreated YFP-PrP* NRK cell that was co-transfected with CER-TMED9 and Golgi marker, FusionRed-SiT. Scale bar represents 10 μm. (right) Magnified panel of the stroked region on the right. Scale bar represents 1 μm.

(G) Time-lapse imaging sequence of a NRK cell that was co-transfected with CER-TMED9 and ER marker, mCherry-Sec61β. Image-collection was started immediately after the addition of 100 μM BRD4780 treatment. Scale bar is 10 μm.

(H) Example of an ER mask created from the ER-marker, mCherry-Sec61β. The ER mask was used to measure the CER-TMED9 fluorescence intensity within the ER at 0 or 30 min time points after the addition of BRD4780 from 9 individual time-lapses as shown in (G) and quantified in (I).

(I) Bar graph representing the mean of ER-localized CER-TMED9 fluorescence-intensity measurements of 9 individual time-lapses of CER-TMED9 and mCherry-Sec61β-expressing cells, as shown in (G). The ER mask was generated based on and as analyzed in in (H). Statistics were calculated from unpaired t-test with Welch's correction with ** indicated $p<0.01$

(J) Representative immunofluorescence images of YFP-PrP* NRK cells that were transfected with CER-TMED9 under control untreated (CTL) or 100 μM BRD4780 (30min)-treated (BRD) conditions and stained for GM130. Scale bar is 20 μm.

(K) Bar graph representing the mean of Pearson's correlation coefficient r to measure colocalization between CER-TMED9 and GM130 under untreated (CTL) and BRD4780 (30 min)-treated (BRD) conditions, as exemplified in (J). Colocalization for 18 cells were measured for control conditions and 11 cells were measured for BRD4780 (30min)-treated cells. Statistics were calculated from unpaired t-test Welch's correction with *** indicating $p<0.001$.

Figure 5

Supporting Figure S4: BRD4780-treatment for 30 min induces a shift in the TMED9 population towards endo H resistance and neuraminidase-sensitivity. (A) Western blots of digested lysates of YFP-PrP* NRK cells in control and BRD4780-treated conditions. Protein lysates were not digested (water) or digested with Endoglycosidase H (Endo H), Neuraminidase only (NA), Neuraminidase+O-Glycosidase (NA+O-G.), Peptide-N-glycosidase F (PNG) or Neuraminidase+O-Glycosidase + Peptide-N-glycosidase F (NA+O-G+PNG) for 3 h at 37°C (n=1 experiment). Blots were probed for TMED9 and calnexin (CNX). (**B**) Western blot band graphics were obtained by first analyzing the plot lanes and second by generating line graphs of pre-selected regions of interest (ROI) from the western blot for control samples (left) or BRD4780 (30 min)-treated samples (right). Black arrow represents the HMW form, orange arrow represents the LMW and purple, the unglycosylated form of TMED9.

Supplemental Figure S4

Supporting Figure S5: Cerulean-TMED9 relocalizes upon BRD4780-treatment leaving YFP-PrP* behind in the ER. Time-lapse images of a typical YFP-PrP* NRK cell that was transfected with CER-TMED9. Image-collection was started immediately after the addition of 100 μM BRD4780 treatment. Scale bar represents 20 μm.

Supporting Figure S5

Figure 6: TMED9's function in RESET is not related to its glycosylation state.

(**A**) Table presenting the different treatments and their effects on TMED9's molecular weight. As depicted in the treatment time-course, YFP-PrP* NRK cells were pre-treated with BRD4780 (BRD) or cycloheximide + BRD4780 (CHX BRD) for different periods of time with or without a 30 min BRD4780 wash-out (wash). "CHX wash" indicates that after pre-treatment with CHX + BRD, the BRD4780 wash-out included cycloheximide. The numbers below the treatment time-course diagram indicate the percentage of cells that were competent for RESET after each treatment course. The percentages were obtained by dividing the number of cells in which YFP-PrP* relocalized from the ER to the Golgi within 30 min of adding thapsigargin by the total number of cells counted. The n values (i.e. total number of cells counted per treatment course) were included directly under the percentage. We assessed "ER" vs "Golgi" localization based on the pattern of YFP fluorescence. At steady-state YFP-PrP* fluorescence is dispersed throughout the cell in a reticulo-tubular pattern with a sharp outline of nucleus that is consistent with ER and the nuclear envelope subdomain of the ER. During RESET, YFP-PrP* fluorescence changes from disperse a perinuclear localization that is consistent with the Golgi. Aligned directly below the treatment courses and percentages, are the TMED9 and GAPDH western blots. Each lane encompassing the treatment, % RESET competent cells, TMED9 molecular weight, GAPDH loading control, was numbered on the bottom. There were 10 individual treatment conditions.

(**B**) Representative images of YFP-PrP* NRK cells after each indicated treatment conditions (1-10). The outline of each cell has been traced in yellow using the ROI tool in Image J.

Figure 6

A

% RESET competent cells (n)	100% (115)	0% (72)	72% (50)	0% (59)	0% (56)	100% (101)	0% (67)	67% (99)	0% (84)	0% (88)

αTMED9 ... -25

αGAPDH ... -37

1   2   3   4   5   6   7   8   9   10

B

YFP-PrP* (grey) + ROIs delimiting cell edges (yellow)

Figure 7: TMED9 must have access to misfolded GPI-APs in the ER to coordinate RESET.
(A) Still frames of a typical YFP-PrP* NRK cell that was co-transfected with CER-TMED9 and mCherry-Sec61β and treated as follows. The first image is of the cell before treatment began (Untreated). The second image is of the cell after 30 min with 50 µg/ml of CHX + BRD4780 (CHX + BRD4780 30 min). The third image is of the cell 30 min after washing with CHX-only medium (CHX wash 30 min). The fourth image is of the cell 30 min after treatment with 50 µg/ml of CHX + 0.1 µM thapsigargin (CHX + TG 30 min). Juxtaposed with each image is an area zoom of the region within the stroked frame. Scale bar represents 10 µm in the larger field of view and in the magnified panel of the stroked region. These select images are also provided in Video 2.
(B) Representative western blots of GFP-tag co-immunoprecipitations (co-IPs) from the parental untransfected NRK cells (P) or YFP-PrP* NRK cells (n=3 biological replicates). Quantified in (C). Cells were harvested at the indicated time points after BRD4780-treatment. Blots were probed for PrP to detect YFP-PrP*, or for endogenous calnexin (CNX) and TMED9.
(C) Bar graph representing the mean band intensity for CNX and TMED9 in the eluates from 3 independently performed experiments as shown in (B). For each time point, CNX and TMED9 eluate band intensities were double normalized. First, they were normalized by the band intensity of the eluted YFP-PrP* band. Second, they were normalized by the time 0 eluate band intensity for each protein probed. Error bars represent standard deviations of the mean. Symbols were coded for individual experiments. Statistics were calculated from unpaired t-test with Welch's correction with and ** indicated p<0.01.
(D) Western blots of GFP-tag co-immunoprecipitations (co-IPs) from the parental untransfected NRK cells (P) or YFP-PrP* NRK cells (n=1 experiment). Cells were harvested after the indicated treatments. Blots were probed for PrP to detect YFP-PrP*, or for endogenous calnexin (CNX) and TMED9.

Figure 7

Supporting Figure S6: BRD4780 treatment dissociates TMED9, TMP21 and TMED2 from YFP-PrP*.
Western blots of GFP-tag co-immunoprecipitations (co-IPs) from the parental untransfected NRK cells (P) or stably transfected YFP-PrP* NRK cells (n=1 experiment). YFP-PrP* NRK cells were treated with 100 µM BRD4780 (BRD) and collected for co-IP at the indicated time points. Cells were harvested at the indicated time points for co-immunoprecipitation of YFP-PrP* with anti-GFP antibody conjugated beads in addition to GFP to detect co-immunoprecipitation of YFP-PrP* constructs, blots were probed for endogenous calnexin (CNX), TMED2, TMP21, and TMED9.

Supporting Figure S6

Supporting Figure S7: Model depicting the role of TMED9 during the steady-state clearance of misfolded GPI-APs from the ER to the Golgi via the RESET pathway. (A) At steady-state, TMED9 (cyan), in conjunction with TMP21 (green), TMED2 (gold) and likely other export machinery (gray), such as other p24-family members, cargo receptors, COPII and COPI coat proteins (gray), traffic between the ER and Golgi, creating a pre-existing pathway for the ER-to-Golgi transport of select substrates. **(B)** Upon release from calnexin (red) during steady-state or ER-stress conditions, misfolded GPI-anchored proteins (GPI-APs) (depicted as purple circles) piggyback with p24-family members to access these p24-family populated ER-exit sites. **(C)** When TMED9 is depleted from the ER either by siRNA knockdown or by BRD4780-treatment, the TMED9 and p24-family populated ER-exit pathway collapses and misfolded GPI-APs are unable to exit the ER to the Golgi. Instead, they remain in association with calnexin. **(B-C)** Properly folded GPI-APs (depicted as purple triangles with yellow stripe) are able to exit the ER to the Golgi through alternate ER-exit sites for subsequent secretion regardless of the presence or absence of TMED9.

Supporting Figure S7

Video 1: Thapsigargin induces rapid ER-export of YFP-PrP* to the Golgi, but does not alter CER-TMED9 localization at the population level. Time-lapse images of a YFP-PrP* NRK cell that is co-expressing CER-TMED9 and Golgi-marker, FusionRed (FusRed)-SiT immediately after the addition of thapsigargin. Images were collected at 5 min intervals. Scale bar represents 10 µm. This Video is associated with Figure 1E.

Video 2: BRD4780-induced ER-export of TMED9 and block on RESET are reversible within 30 min of treatment. Time-lapse images collected at 5 min intervals of a YFP-PrP* NRK cell that was co-transfected with CER-TMED9 and mCherry-Sec61β. As specified in text in the video, the cell was treated with CHX + BRD4780 for 30min, then washed with medium containing CHX-only for 30 min, and then treated with CHX + thapsigargin (TG). Scale bar represents 10 µm. This Video is associated with Figure 7A.

References

1. Princiotta MF, Finzi D, Qian SB, Gibbs J, Schuchmann S, Buttgereit F, et al. Quantitating protein synthesis, degradation, and endogenous antigen processing. Immunity. 2003;18(3):343-54. doi: 10.1016/s1074-7613(03)00051-7. PubMed PMID: 12648452.

2. Dobson CM. Protein folding and misfolding. Nature. 2003;426(6968):884-90. doi: 10.1038/nature02261. PubMed PMID: 14685248.

3. Hartl FU, Bracher A, Hayer-Hartl M. Molecular chaperones in protein folding and proteostasis. Nature. 2011;475(7356):324-32. Epub 20110720. doi: 10.1038/nature10317. PubMed PMID: 21776078.

4. Braakman I, Hebert DN. Protein folding in the endoplasmic reticulum. Cold Spring Harb Perspect Biol. 2013;5(5):a013201. Epub 20130501. doi: 10.1101/cshperspect.a013201. PubMed PMID: 23637286; PubMed Central PMCID: PMCPMC3632058.

5. Anelli T, Sitia R. Protein quality control in the early secretory pathway. EMBO J. 2008;27(2):315-27. doi: 10.1038/sj.emboj.7601974. PubMed PMID: 18216874; PubMed Central PMCID: PMCPMC2234347.

6. Phillips BP, Gomez-Navarro N, Miller EA. Protein quality control in the endoplasmic reticulum. Curr Opin Cell Biol. 2020;65:96-102. Epub 20200511. doi: 10.1016/j.ceb.2020.04.002. PubMed PMID: 32408120; PubMed Central PMCID: PMCPMC7588826.

7. Ruggiano A, Foresti O, Carvalho P. Quality control: ER-associated degradation: protein quality control and beyond. J Cell Biol. 2014;204(6):869-79. doi: 10.1083/jcb.201312042. PubMed PMID: 24637321; PubMed Central PMCID: PMCPMC3998802.

8. Sun Z, Brodsky JL. Protein quality control in the secretory pathway. J Cell Biol. 2019;218(10):3171-87. Epub 20190919. doi: 10.1083/jcb.201906047. PubMed PMID: 31537714; PubMed Central PMCID: PMCPMC6781448.

9. Molinari M. ER-phagy responses in yeast, plants, and mammalian cells and their crosstalk with UPR and ERAD. Dev Cell. 2021;56(7):949-66. Epub 20210324. doi: 10.1016/j.devcel.2021.03.005. PubMed PMID: 33765438.

10. De Leonibus C, Cinque L, Settembre C. Emerging lysosomal pathways for quality control at the endoplasmic reticulum. FEBS Lett. 2019;593(17):2319-29. Epub 20190813. doi: 10.1002/1873-3468.13571. PubMed PMID: 31388984.

11. Wilkinson S. Emerging Principles of Selective ER Autophagy. J Mol Biol. 2020;432(1):185-205. Epub 20190514. doi: 10.1016/j.jmb.2019.05.012. PubMed PMID: 31100386; PubMed Central PMCID: PMCPMC6971691.

12. Satpute-Krishnan P, Ajinkya M, Bhat S, Itakura E, Hegde RS, Lippincott-Schwartz J. ER stress-induced clearance of misfolded GPI-anchored proteins via the secretory pathway. Cell. 2014;158(3):522-33. doi: 10.1016/j.cell.2014.06.026. PubMed PMID: 25083867; PubMed Central PMCID: PMCPMC4121523.

13. Ashok A, Hegde RS. Selective processing and metabolism of disease-causing mutant prion proteins. PLoS Pathog. 2009;5(6):e1000479. Epub 2009/06/23. doi: 10.1371/journal.ppat.1000479. PubMed PMID: 19543376; PubMed Central PMCID: PMCPMC2691595.

14. Cheatham AM, Sharma NR, Satpute-Krishnan P. Competition for calnexin binding regulates secretion and turnover of misfolded GPI-anchored proteins. J Cell Biol. 2023;222(10). Epub 20230913. doi: 10.1083/jcb.202108160. PubMed PMID: 37702712; PubMed Central PMCID: PMCPMC10499038.

15. Guo XY, Liu YS, Gao XD, Kinoshita T, Fujita M. Calnexin mediates the maturation of GPI-anchors through ER retention. J Biol Chem. 2020;295(48):16393-410. Epub 2020/09/25. doi: 10.1074/jbc.RA120.015577. PubMed PMID: 32967966; PubMed Central PMCID: PMCPMC7705322.

16. Belden WJ, Barlowe C. Erv25p, a component of COPII-coated vesicles, forms a complex with Emp24p that is required for efficient endoplasmic reticulum to Golgi transport. J Biol Chem. 1996;271(43):26939-46. doi: 10.1074/jbc.271.43.26939. PubMed PMID: 8900179.

17. Dominguez M, Dejgaard K, Fullekrug J, Dahan S, Fazel A, Paccaud JP, et al. gp25L/emp24/p24 protein family members of the cis-Golgi network bind both COP I and II coatomer. J Cell Biol.

1998;140(4):751-65. doi: 10.1083/jcb.140.4.751. PubMed PMID: 9472029; PubMed Central PMCID: PMCPMC2141742.

18. Strating JR, van Bakel NH, Leunissen JA, Martens GJ. A comprehensive overview of the vertebrate p24 family: identification of a novel tissue-specifically expressed member. Mol Biol Evol. 2009;26(8):1707-14. Epub 20090508. doi: 10.1093/molbev/msp099. PubMed PMID: 19429673.

19. Schimmoller F, Singer-Kruger B, Schroder S, Kruger U, Barlowe C, Riezman H. The absence of Emp24p, a component of ER-derived COPII-coated vesicles, causes a defect in transport of selected proteins to the Golgi. EMBO J. 1995;14(7):1329-39. doi: 10.1002/j.1460-2075.1995.tb07119.x. PubMed PMID: 7729411; PubMed Central PMCID: PMCPMC398218.

20. Zavodszky E, Hegde RS. Misfolded GPI-anchored proteins are escorted through the secretory pathway by ER-derived factors. Elife. 2019;8. Epub 20190516. doi: 10.7554/eLife.46740. PubMed PMID: 31094677; PubMed Central PMCID: PMCPMC6541436.

21. Roberts BS, Satpute-Krishnan P. The many hats of transmembrane emp24 domain protein TMED9 in secretory pathway homeostasis. Front Cell Dev Biol. 2022;10:1096899. Epub 20230116. doi: 10.3389/fcell.2022.1096899. PubMed PMID: 36733337; PubMed Central PMCID: PMCPMC9888432.

22. Dvela-Levitt M, Kost-Alimova M, Emani M, Kohnert E, Thompson R, Sidhom EH, et al. Small Molecule Targets TMED9 and Promotes Lysosomal Degradation to Reverse Proteinopathy. Cell. 2019;178(3):521-35 e23. doi: 10.1016/j.cell.2019.07.002. PubMed PMID: 31348885.

23. Roberts BS, Mitra D, Abishek S, Beher R, Satpute-Krishnan P. The p24-family and COPII subunit SEC24C facilitate the clearance of alpha1-antitrypsin Z from the endoplasmic reticulum to lysosomes. Mol Biol Cell. 2024;35(3):ar45. Epub 20240131. doi: 10.1091/mbc.E23-06-0257. PubMed PMID: 38294851.

24. Rane NS, Yonkovich JL, Hegde RS. Protection from cytosolic prion protein toxicity by modulation of protein translocation. EMBO J. 2004;23(23):4550-9. Epub 20041104. doi: 10.1038/sj.emboj.7600462. PubMed PMID: 15526034; PubMed Central PMCID: PMCPMC533048.

25. Rane NS, Chakrabarti O, Feigenbaum L, Hegde RS. Signal sequence insufficiency contributes to neurodegeneration caused by transmembrane prion protein. J Cell Biol. 2010;188(4):515-26. Epub 20100215. doi: 10.1083/jcb.200911115. PubMed PMID: 20156965; PubMed Central PMCID: PMCPMC2828915.

26. Liu YS, Guo XY, Hirata T, Rong Y, Motooka D, Kitajima T, et al. N-Glycan-dependent protein folding and endoplasmic reticulum retention regulate GPI-anchor processing. J Cell Biol. 2018;217(2):585-99. Epub 2017/12/20. doi: 10.1083/jcb.201706135. PubMed PMID: 29255114; PubMed Central PMCID: PMCPMC5800811.

27. Tashima Y, Hirata T, Maeda Y, Murakami Y, Kinoshita T. Differential use of p24 family members as cargo receptors for the transport of glycosylphosphatidylinositol-anchored proteins and Wnt1. J Biochem. 2022;171(1):75-83. doi: 10.1093/jb/mvab108. PubMed PMID: 34647572.

28. Zhang M, Liu L, Lin X, Wang Y, Li Y, Guo Q, et al. A Translocation Pathway for Vesicle-Mediated Unconventional Protein Secretion. Cell. 2020;181(3):637-52 e15. Epub 20200408. doi: 10.1016/j.cell.2020.03.031. PubMed PMID: 32272059.

29. Fregno I, Fasana E, Bergmann TJ, Raimondi A, Loi M, Solda T, et al. ER-to-lysosome-associated degradation of proteasome-resistant ATZ polymers occurs via receptor-mediated vesicular transport. EMBO J. 2018;37(17). Epub 20180803. doi: 10.15252/embj.201899259. PubMed PMID: 30076131; PubMed Central PMCID: PMCPMC6120659.

30. Doms RW, Russ G, Yewdell JW. Brefeldin A redistributes resident and itinerant Golgi proteins to the endoplasmic reticulum. J Cell Biol. 1989;109(1):61-72. doi: 10.1083/jcb.109.1.61. PubMed PMID: 2745557; PubMed Central PMCID: PMCPMC2115463.

31. Lippincott-Schwartz J, Yuan LC, Bonifacino JS, Klausner RD. Rapid redistribution of Golgi proteins into the ER in cells treated with brefeldin A: evidence for membrane cycling from Golgi to ER. Cell. 1989;56(5):801-13. doi: 10.1016/0092-8674(89)90685-5. PubMed PMID: 2647301; PubMed Central PMCID: PMCPMC7173269.

32. Fujiwara T, Oda K, Yokota S, Takatsuki A, Ikehara Y. Brefeldin A causes disassembly of the Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. J Biol Chem. 1988;263(34):18545-52. PubMed PMID: 3192548.

33. Sciaky N, Presley J, Smith C, Zaal KJ, Cole N, Moreira JE, et al. Golgi tubule traffic and the effects of brefeldin A visualized in living cells. J Cell Biol. 1997;139(5):1137-55. doi: 10.1083/jcb.139.5.1137. PubMed PMID: 9382862; PubMed Central PMCID: PMCPMC2140213.

34. Freeze HH, Kranz C. Endoglycosidase and glycoamidase release of N-linked glycans. Curr Protoc Mol Biol. 2010;Chapter 17:Unit 17 3A. doi: 10.1002/0471142727.mb1713as89. PubMed PMID: 20069534; PubMed Central PMCID: PMCPMC3869378.

35. Varki A CR, Esko JD, Stanley P, Hart GW, Aebi M, Mohnen D, Kinoshita T, Packer NH, Prestegard JH, Schnaar RL, Seeberger PH Essentials of Glycobiology [Internet]. In: Varki A, Cummings RD, Esko JD, Stanley P, Hart GW, Aebi M, et al., editors. Essentials of Glycobiology. 4th ed. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2022.

36. Hammond C, Braakman I, Helenius A. Role of N-linked oligosaccharide recognition, glucose trimming, and calnexin in glycoprotein folding and quality control. Proc Natl Acad Sci U S A. 1994;91(3):913-7. doi: 10.1073/pnas.91.3.913. PubMed PMID: 8302866; PubMed Central PMCID: PMCPMC521423.

37. Bhide GP, Colley KJ. Sialylation of N-glycans: mechanism, cellular compartmentalization and function. Histochem Cell Biol. 2017;147(2):149-74. Epub 20161214. doi: 10.1007/s00418-016-1520-x. PubMed PMID: 27975143; PubMed Central PMCID: PMCPMC7088086.

38. Rabouille C, Hui N, Hunte F, Kieckbusch R, Berger EG, Warren G, et al. Mapping the distribution of Golgi enzymes involved in the construction of complex oligosaccharides. J Cell Sci. 1995;108 (Pt 4):1617-27. doi: 10.1242/jcs.108.4.1617. PubMed PMID: 7615680.

39. Fullekrug J, Suganuma T, Tang BL, Hong W, Storrie B, Nilsson T. Localization and recycling of gp27 (hp24gamma3): complex formation with other p24 family members. Mol Biol Cell. 1999;10(6):1939-55. doi: 10.1091/mbc.10.6.1939. PubMed PMID: 10359607; PubMed Central PMCID: PMCPMC25391.

40. Lavoie C, Paiement J, Dominguez M, Roy L, Dahan S, Gushue JN, et al. Roles for alpha(2)p24 and COPI in endoplasmic reticulum cargo exit site formation. J Cell Biol. 1999;146(2):285-99. doi: 10.1083/jcb.146.2.285. PubMed PMID: 10427085; PubMed Central PMCID: PMCPMC3206572.

41. Dvela-Levitt M, Shaw JL, Greka A. A Rare Kidney Disease To Cure Them All? Towards Mechanism-Based Therapies for Proteinopathies. Trends Mol Med. 2021;27(4):394-409. Epub 20201216. doi: 10.1016/j.molmed.2020.11.008. PubMed PMID: 33341352.

42. Mitrovic S, Ben-Tekaya H, Koegler E, Gruenberg J, Hauri HP. The cargo receptors Surf4, endoplasmic reticulum-Golgi intermediate compartment (ERGIC)-53, and p25 are required to maintain the architecture of ERGIC and Golgi. Mol Biol Cell. 2008;19(5):1976-90. Epub 20080220. doi: 10.1091/mbc.e07-10-0989. PubMed PMID: 18287528; PubMed Central PMCID: PMCPMC2366877.

43. Strating JR, Martens GJ. The p24 family and selective transport processes at the ER-Golgi interface. Biol Cell. 2009;101(9):495-509. doi: 10.1042/BC20080233. PubMed PMID: 19566487.

44. Rajagopalan S, Xu Y, Brenner MB. Retention of unassembled components of integral membrane proteins by calnexin. Science. 1994;263(5145):387-90. doi: 10.1126/science.8278814. PubMed PMID: 8278814.

45. Wada I, Rindress D, Cameron PH, Ou WJ, Doherty JJ, 2nd, Louvard D, et al. SSR alpha and associated calnexin are major calcium binding proteins of the endoplasmic reticulum membrane. J Biol Chem. 1991;266(29):19599-610. PubMed PMID: 1918067.

46. Fehlinger A, Wolf H, Hossinger A, Duernberger Y, Pleschka C, Riemschoss K, et al. Prion strains depend on different endocytic routes for productive infection. Sci Rep. 2017;7(1):6923. Epub 20170731. doi: 10.1038/s41598-017-07260-2. PubMed PMID: 28761068; PubMed Central PMCID: PMCPMC5537368.

47. Chassefeyre R, Chaiamarit T, Verhelle A, Novak SW, Andrade LR, Leitao ADG, et al. Endosomal sorting drives the formation of axonal prion protein endoggresomes. Sci Adv. 2021;7(52):eabg3693. Epub 20211222. doi: 10.1126/sciadv.abg3693. PubMed PMID: 34936461; PubMed Central PMCID: PMCPMC8694590.

48. Jeffrey M, Goodsir CM, Bruce M, McBride PA, Scott JR, Halliday WG. Correlative light and electron microscopy studies of PrP localisation in 87V scrapie. Brain Res. 1994;656(2):329-43. Epub 1994/09/12. doi: 10.1016/0006-8993(94)91477-x. PubMed PMID: 7820594.

49. Scheckel C, Aguzzi A. Prions, prionoids and protein misfolding disorders. Nat Rev Genet. 2018;19(7):405-18. Epub 2018/05/02. doi: 10.1038/s41576-018-0011-4. PubMed PMID: 29713012.

50. Hailey DW, Rambold AS, Satpute-Krishnan P, Mitra K, Sougrat R, Kim PK, et al. Mitochondria supply membranes for autophagosome biogenesis during starvation. Cell. 2010;141(4):656-67. doi: 10.1016/j.cell.2010.04.009. PubMed PMID: 20478256; PubMed Central PMCID: PMCPMC3059894.

51. Mantei N, Villa M, Enzler T, Wacker H, Boll W, James P, et al. Complete primary structure of human and rabbit lactase-phlorizin hydrolase: implications for biosynthesis, membrane anchoring and evolution of the enzyme. EMBO J. 1988;7(9):2705-13. doi: 10.1002/j.1460-2075.1988.tb03124.x. PubMed PMID: 2460343; PubMed Central PMCID: PMCPMC457059.

52. Hauri HP, Sterchi EE, Bienz D, Fransen JA, Marxer A. Expression and intracellular transport of microvillus membrane hydrolases in human intestinal epithelial cells. J Cell Biol. 1985;101(3):838-51. doi: 10.1083/jcb.101.3.838. PubMed PMID: 3897250; PubMed Central PMCID: PMCPMC2113743.

53. Naim HY, Sterchi EE, Lentze MJ. Biosynthesis and maturation of lactase-phlorizin hydrolase in the human small intestinal epithelial cells. Biochem J. 1987;241(2):427-34. doi: 10.1042/bj2410427. PubMed PMID: 3109375; PubMed Central PMCID: PMCPMC1147578.

54. Ouwendijk J, Peters WJ, van de Vorstenbosch RA, Ginsel LA, Naim HY, Fransen JA. Routing and processing of lactase-phlorizin hydrolase in transfected Caco-2 cells. J Biol Chem. 1998;273(12):6650-5. doi: 10.1074/jbc.273.12.6650. PubMed PMID: 9506961.

55. Keller P, Toomre D, Diaz E, White J, Simons K. Multicolour imaging of post-Golgi sorting and trafficking in live cells. Nat Cell Biol. 2001;3(2):140-9. doi: 10.1038/35055042. PubMed PMID: 11175746.

56. Cole NB, Smith CL, Sciaky N, Terasaki M, Edidin M, Lippincott-Schwartz J. Diffusional mobility of Golgi proteins in membranes of living cells. Science. 1996;273(5276):797-801. doi: 10.1126/science.273.5276.797. PubMed PMID: 8670420.

57. Shemiakina, II, Ermakova GV, Cranfill PJ, Baird MA, Evans RA, Souslova EA, et al. A monomeric red fluorescent protein with low cytotoxicity. Nat Commun. 2012;3:1204. doi: 10.1038/ncomms2208. PubMed PMID: 23149748.

58. Nixon-Abell J, Obara CJ, Weigel AV, Li D, Legant WR, Xu CS, et al. Increased spatiotemporal resolution reveals highly dynamic dense tubular matrices in the peripheral ER. Science. 2016;354(6311). Epub 20161027. doi: 10.1126/science.aaf3928. PubMed PMID: 27789813; PubMed Central PMCID: PMCPMC6528812.