Genome-directed study reveals the diversity of *Salmonella* T6SS effectors and

identifies a novel family of lipid-targeting antibacterial toxins

Gianlucca G. Nicastro[2,3¶], Stephanie Sibinelli-Sousa[1, 2¶], Julia Takuno Hespanhol[1],

Thomas W. C. Santos[2], Joseph P. Munoz[1], Rosangela S. Santos[4], Blanca M. Perez-

Sepulveda[5], Sayuri Miyamoto[4], L. Aravind[3], Robson F. de Souza[2,*], Ethel Bayer-

Santos[1,2,6,*].

[1]Department of Molecular Biosciences, College of Natural Sciences, The University of

Texas at Austin, Austin, USA.

[2]Departamento de Microbiologia, Instituto de Ciências Biomédicas, Universidade de

São Paulo, São Paulo, Brazil.

[3]Computational Biology Branch, National Center for Biotechnology Information,

National Library of Medicine, National Institutes of Health, USA.

[4]Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São

Paulo, Brazil.

[5]Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool,

Liverpool, UK.

[6]LaMontagne Center for Infectious Disease, The University of Texas at Austin, Austin,

Texas, USA.

Short title: Novel toxin domains and lipid-targeting NlpC/P60 toxins

*Corresponding authors: ebayersantos@austin.utexas.edu (EBS); rfsouza@usp.br

(RFS)

¶These authors contributed equally to this work.

Abstract

Bacterial warfare is a common and ancient phenomenon in nature, where bacterial species use strategies to inhibit the growth or kill competitors. This involves the production and deployment of antibacterial toxins that disrupt essential cellular processes in target cells. The continuous arms race in which bacteria acquire new toxin and immunity proteins to promote increased adaptation to their environment is responsible for the diversification of this toxin repertoire. Here, we deployed *in-silico* strategies to analyze 10,000 genomes and identify effectors secreted via the type VI secretion system of *Salmonella*. We identified 128 candidates, which are widespread in a vast array of *Salmonella* serovars and other bacterial species. Tox-Act1 is among the most frequent candidates and was selected for in-depth characterization. Tox-Act1 contains a permuted NlpC/P60 papain-like catalytic core characteristic of lipid-targeting members rather than the typical peptidases or amidases. Evolutionary analysis revealed the relationship of Tox-Act1 with acyltransferases. Biochemical assays with purified toxin and lipidomics of intoxicated cells showed that Tox-Act1 exhibits phospholipase activity, cleaving off acyl groups from phosphatidylglycerol and phosphatidylethanolamine. In addition, we demonstrate that Tox-Act1 is secreted in a T6SS-dependent manner and provide a competitive advantage during colonization of the gut of infected mice. This work broadens our understanding of toxin domains and provides the first direct characterization of a lipid-targeting NlpC/P60 domain in biological conflicts.

Keywords: *Salmonella*, T6SS, effector, toxin, NlpC/P60, phospholipase, acyltransferase.

Introduction

Competition is a fundamental biological process in nature, occurring both within and between species that share a common environment. Bacteria actively participate in these ecological battles employing a potent arsenal of toxins as their weaponry[1]. A prominent mechanism for toxin delivery among Gram-negative bacteria is via the type 6 secretion system (T6SS)[2]. Phylogenetic analysis of T6SS components showed that there are four types of T6SSs (T6SS[i-iv])[3-5], with the canonical T6SS[i] present in *Proteobacteria* being further classified into six subtypes (i1, i2, i3, i4a, i4b and i5)[4, 6, 7]. The T6SS functions in a contact-dependent manner and relies on the biochemical properties of secreted effector toxins for its function[8]. These toxin domains frequently recombine and move via lateral gene transfer, allowing them to be delivered by different secretion systems, which warrants their name as polymorphic toxins[9].

During secretion via the T6SS, toxins are loaded onto a spear-like structure formed by hexameric rings of Hcp proteins capped by a spike comprising a trimer of VgrG sharpened by a PAAR protein[10]. Effectors are translocated into target cells either fused at the C-terminus of Hcp, VgrG, and PAAR proteins (named specialized effectors), or associated with these proteins via adaptors (cargo effectors)[11]. Antibacterial toxins are often paired with immunity proteins that prevent self-intoxication, thus forming gene pairs that are frequently located near structural components of the T6SS[8, 12].

The protective role of the endogenous microbiota against *Salmonella* infection has been recognized for years[13]; however, only recently have studies started to reveal the mechanism by which the microbiota maintains gut homeostasis and promotes colonization resistance[14]. Despite this understanding, there is a relative scarcity of information about the direct antimicrobial mechanisms employed by commensals and

pathogens during these disputes for niche control[15]. T6SSs clusters are conserved

across several *Salmonella* spp., highlighting their importance to the fitness of these

bacteria[16, 17]. *Salmonella* spp. encode T6SSs in different pathogenicity islands

(SPIs), which were acquired by distinct horizontal transfer events[16, 17]. *S.*

Typhimurium encodes a T6SS subtype i3 within SPI-6, which is important for

interbacterial competition and gut colonization in mice[18, 19]. However, there is

limited information about the identity or mechanism of the T6SS effector repertoire that

contributes to these phenotypes.

Here, we set out to identify the repertoire of T6SS effectors in a dataset of

10,000 *Salmonella* isolates utilizing a computational approach. Employing sensitive

sequence and structure searches alongside comparative genomics, we identified 128

candidates that are widespread among several *Salmonella* serovars and additional

bacterial species. This comprehensive analysis indicates that T6SSs subtypes i3 and i1

are associated with antibacterial and anti-eukaryotic effectors, respectively.

Furthermore, our findings reveal that each of the 149 serovars harbors a unique

combination of effectors, likely required for their specific ecological interactions. A

detailed examination of a selected candidate (Tox-Act1) encoding a newly identified

domain showed that it is an antibacterial effector belonging to the NlpC/P60

superfamily with a permuted catalytic core. Tox-Act1 is evolutionarily related to

acyltransferases and displays phospholipase activity to modify target-cell membrane

composition. This study offers a comprehensive characterization of new toxins,

especially the arsenal linked to T6SSs in *Salmonella*, identifying novel effector families

and providing an in-depth analysis of a new protein family that specifically targets

lipids.

Results

Computational pipeline for the automatic identification of T6SS components

T6SS effectors are often encoded in close genomic proximity to structural components of the system [16, 20, 21]. To identify new effectors, we employed a "guilt-by-association" approach, which relies on the conservation of genomic context [22]. We applied this methodology to a curated dataset of 10,000 *Salmonella* genomes (10KSG) [23] (Fig 1A). First, we located and classified the T6SS clusters using hidden Markov models (HMMs) from different sources[9, 24-26]. We collected 10 genes upstream and downstream of each locus encoding T6SS components, referred to as genomic sites (Fig 1A). A total of 42,560 genomic sites housing at least one T6SS component were identified. We then applied a graph theoretic strategy leveraging the Jaccard index for network construction[27], followed by the Louvain algorithm[28] to automatically categorize genomic sites. Analysis resulted in five categories: T6SS subtype i1, i2, i3, i4b, and orphan (Fig 1B, 1C).

We observed mobility of the distinct T6SS subtypes among genomic sites and pathogenicity islands, depending on the strain and/or serovar. Using tRNAs as markers to identify the insertion sites[29], we found that subtype i3 (90%, 4929/5461 genomic sites) is mainly located within SPI-6 and flanked by the aspartate tRNA (tRNA^{Asp})[16], while subtype i1 is not well conserved and found within SPI-19 associated with tRNA^{Ala} (12%, 66/532 genomic sites)[16], and SPI-6 (3%, 18/532 genomes) flanked by tRNA^{Asp} - most subtype i1 could not be assigned to a specific location (82%, 437/532 genomic sites). These results indicate that distinct T6SSs subtypes are inserted into different genomic sites that are hot spots for horizontal transfer events, and the

combination between the insertion site and the introduced subtype varies according to

the *Salmonella* strain/serovar.

Fig 1. Computational pipeline for the identification of T6SS subtypes and effector

repertoire within 10,000 *Salmonella* genomes. (a) Pipeline used for classification of

genomic sites and T6SS subtypes. **(b)** Number of genomes containing the different

T6SS subtypes within the 10KSG dataset. **(c)** Examples of the genomic organization of

T6SS structural clusters from distinct phylogenetic subtypes. Colors denote structural

proteins forming the membrane complex (orange), sheath and inner tube (light blue),

baseplate and spike components (green). Effectors are shown in red and immunities in

dark blue. **(d)** *In silico* strategies used for the identification and classification of T6SS

effectors. **(e)** Comparison of profile Hidden Markov Models (pHMMs) of STox to

published models of related families. Each circle corresponds to a population of proteins

detected by an STox model. The relative frequency of proteins detected by an STox model compared to a reference model is shown on the horizontal axis. The HMM divergence score is shown in the vertical axis. The blue-white-red color scale of each dot represents the values for the Spearman correlation index between STox and reference alignment scores for the same proteins. Yellow circles represent STox models that detect proteins that are not recognized by previously existing models. The radius of each circle is proportional to the total number of proteins detected only by STox models. The regression line does not include the data points for the models represented by yellow circles. **(f)** Schematic representation of the functional classes of T6SS effectors identified in the 10KSG.

Identification of the T6SS effectors repertoire in 10,000 *Salmonella* genomes

Next, we focused on identifying effector toxins using three strategies (Fig 1D). First, we focused on proteins containing N-terminal PAAR, VgrG or Hcp domains and additional C-terminal domains with more than 50 amino acids. These C-terminal regions were isolated and grouped based on similarity (80% coverage and 1e-3 e-value). Second, we used the genomic sites containing T6SS components and analyzed up to five genes upstream and downstream via the software BastionX[30]. Third, we used amino acid sequences, HMMs and PSSMs (position-specific scoring matrix) from SecReT6[31], Pfam[25], and Zhang et al. [9] to search the 10KSG dataset for previously described effectors. For candidates not recognized by previously described models, we collected homologs from NCBI using 4 iterations of PSI-BLAST[32] and generated multiple sequence alignments and HMMs.

We then used a series of sequence and structure-based strategies to classify and

annotate the function of these candidates: i) profile-profile comparison methods such as

HHsearch[33] were used to detect distant homologs; ii) structural models were created

using Alphafold2[34] from multiple sequence alignments to perform searches in

FoldSeek[35] and DALI[36] against PDB[37] and AlphaFoldDB[38]; iii) the structure-

structure comparison algorithm from FoldSeek was used to cluster groups of candidates.

All information collected was manually examined to establish the final domain

annotation (Fig 1D). Candidates that displayed sequence or structural similarity to

known toxin domains or proteins of unknown function that displayed conserved

genomic organization and/or adjacent conserved putative immunity proteins across

several species were maintained.

In total, we identified 128 groups of effector toxins (Table S1 and Fig 1F).

Eighty-thee were already described in public databases (e.g. Ntox47)[9, 25], or

represent individual effectors that were experimentally characterized but for which

HMMs have not been produced and made available (e.g. TreTu)[21, 39-54]. For the

latter, the newly created HMMs were named with a ".st" suffix (e.g. TreTu.st). Within

the 128 candidates, 45 groups comprise new toxin domains or highly divergent

variations that were not detected by previously published HMMs (Fig 1E). This

justified them being distinct groups requiring the design of new HMMs. These groups

of effectors were named STox followed by a number (e.g. STox_1) (Table S1 and Fig

1F). It is worth noting that these effectors are not just present in *Salmonella* and are

detected widespread across several species, comprising polymorphic toxin domains[9].

A few examples of species encoding each STox, together with the amino acid sequence

alignments   and   AlphaFold   prediction   can   be   found   at

https://leepbioinfo.github.io/10ksgt6ss.

Inspection of genomic context across several bacterial species revealed that most

candidates exhibited a conserved adjacent gene coding for a predicted immunity

protein, thus suggesting antibacterial activity (83.6%, 107/128). Some effectors, which

lacked conserved immunity proteins, were predicted to display anti-eukaryotic activity

(12.4%, 14/128) (Table S1 and Fig 1F). The analysis revealed a diverse array of cellular

targets and biochemical activities among the 128 toxin groups (Table S1 and Fig 1F).

Notably, the activities of a few STox effectors could not be predicted confidently and

will require further analysis (Fig 1F). Overall, these findings highlight the significant

diversity of effector toxins encoded by *S. enterica* serovars and reveal an array of novel

proteins used in interbacterial competition and as virulence factors.

Salmonella serovars encode unique subsets of effector toxins

The most frequent effectors detected within the 10KSG dataset were the

peptidoglycan-targeting effectors Tlde1, a L,D-transpeptidase[20]; and Tae4, a papain-

like amidase[39] (Fig 2A). These effector families were followed by Ntox47, a

predicted RNase with the BECR fold[55]; the metallopeptidase Tox-HopH1; a second

more divergent group of Ntox47 (Ntox47.st2); the ADP-ribosyltransferases TreTu[40]

and STox_62; and another peptidoglycan-targeting effector Tae2 [39] (Fig 2A).

Together, these eight effectors were found in most genomes within the 10KSG and

constitute the core T6SS effectors. In addition, by analyzing the combination of

effectors in each genome, we found that 88% of the genomes within the 10KSG dataset

were estimated to encode up to 4 effectors per genome (mode = 3), while 12% encode

combinations that range between 5-18 effectors (mode = 5) (Fig S1).

Fig 2. Unique subsets of effectors are associated with specific *Salmonella* serovars and T6SS subtypes. (a) The most frequent effectors identified in the 10KSG dataset. Each bar represents the number of genomes encoding a specific effector. Colors represent different effector activities, with light colors representing orphan effectors while dark colors represent effectors encoded within the structural cluster. **(b)** Schematic representation of the most common sets of effectors in genomes encoding different T6SS subtypes. The number of genomes is indicated on the right. Colors represent activity as shown in (a). **(c)** The five most frequent effectors encoded in different *Salmonella* serovars. Colors indicate the effector activity as in (a). **(d)** Pie chart illustrating the relative proportions of effectors classified by activity encoded within the T6SSs subtypes i3, i1, and i2. Colors as shown in (a). **(e)** Venn diagram illustrating the proportion of overlap between effectors encoded within each T6SS structural cluster (blue: i1; purple: subtype i2; red: subtype i3; and green: orphan). **(f)** Schematic representation of the genetic organization of T6SSs showing the position of variable regions in which the effector and immunity proteins are encoded. Colors denote structural proteins forming the membrane complex (orange), sheath and inner tube

(light blue), baseplate and spike components (green). Effectors are shown in red, and

immunities in dark blue.

Next, we determined the 5 most frequent effectors detected in each of the 149

Salmonella serovars. Serovars that predominantly encode subtype i3, such as *S.*

Typhimurium. *S.* Panama, *S.* Infantis, and *S.* Typhi, harbor effectors targeting

peptidoglycan (e.g. Tae4, Tlde1, Tae2), nucleases (e.g. Ntox47, Tox-WHH, ColE7,

Tox-AHH) and ART superfamily enzymes (e.g. TreTu, STox_62) (Fig 2B, 2C). *S.*

Dublin, which contains both subtypes i1 and i3, displays a mixture of effectors targeting

the peptidoglycan (Tae2 and Tlde1), a nuclease (Ntox47), and a metallopeptidase (Tox-

HopH1) (Fig 2C). *S.* Agona encodes only a T6SS subtype i1 and contains effectors with

nuclease (ColE7 and Ntox19), pore-forming (STox_47), and undetermined activities

(STox_18 and STox_53) (Fig 2C). The core effectors of each serovar can be found in

Fig S2.

244   Each *Salmonella* T6SS subtype is associated with target-

245   specific effectors

We analyzed the combination of effectors most frequently encoded in genomes

containing phylogenetically distinct T6SSs. Our findings revealed that the antibacterial

subtype i3 predominantly displays a combination of peptidoglycan-targeting effectors

(e.g. Tae4 and Tlde1); nucleases (e.g. Ntox47 and Tox-WHH); and ART family

enzymes (e.g. TreTu and STox_62) (Fig 2A, 2B). Conversely, genomes encoding the

anti-eukaryotic subtype i1 lack peptidoglycan-targeting effectors and show a

combination of nucleases (e.g. ColE7), pore-forming toxins (e.g. Ssp6) and effectors

with undetermined activity (Fig 2B, 2C). For genomes encoding both subtypes i1 and

i3, there was a mix of effectors with antibacterial (e.g. Tae2, Tlde1, Ntox47) and anti-

eukaryotic activity (e.g. Tox-HopH1) (Fig 2B, 2C). This data supports the classification

of *Salmonella* T6SS subtype i3 as antibacterial and subtype i1 as anti-eukaryotic.

Notably, effectors encoded within or close to T6SS structural clusters, show

minimum overlap (Fig 2E), suggesting an evolutionary scenario where T6SS clusters

are acquired alongside with the associated set of effectors and/or that certain subsets of

effectors are preferably exchanged among bacteria harboring similar T6SS subtypes.

Previous analysis of subtype i3 identified the insertion of thee variable regions between

the structural genes (VR1-3) in which effectors are encoded [16] (Fig 2F). Our results

indicate that VR1 and VR2 contain mainly toxins targeting the periplasm (e.g. Tae2,

Tae4, Tlde1, Tox-Act1) whereas VR3 primarily houses toxins targeting the cytoplasm

(e.g. Ntox47, Tox-WHH, TreTu). The latter are typically associated with an N-terminal

PAAR domain and Rhs (Rearrangement hotspot) repeats, which assist in the

translocation across the bacterial inner membrane to reach the cytoplasmic targets[40,

55-57] (Fig 2F). The position of effectors at the edge of the T6SS cluster and the

domain architecture containing Rhs repeats facilitate recombination events[55], possibly

leading to the greater diversity of nuclease effectors observed at the VR3 (Fig S1).

Tox-Act1 is a T6SS effector used for interbacterial

competition in the mouse gut

Among the newly identified effectors, STox_15, which was renamed Tox-Act1

(toxin acyltransferase 1, see details below), emerged as one of the most abundant (Fig

2A). To analyze whether Tox-Act1 and its associated downstream gene (Imm-Act1)

form an effector and immunity pair (Fig 3A), we cloned these genes in compatible

vectors under the control of different promoters and assessed toxicity upon expression

in *Escherichia coli*. Tox-Act1 is toxic in the periplasm (SP-Tox-Act1) but not in the

cytoplasm (Tox-Act1) of *E. coli* and co-expression with Imm-Act1 neutralizes the effect

(Fig 3B).

We performed time-lapse microscopy to evaluate bacterial growth and

morphology at the single cell level. *E. coli* carrying the plasmid with SP-Tox-Act1 grew

normally in d-glucose (repressed conditions) (Movie S1); however, shortly after

induction of SP-Tox-Act1 with l-arabinose, cells began lysing (Movie S2). It was

curious that cells lysed without losing their rod shape, which suggests that the

peptidoglycan was not affected (Fig 3C). After lysing, residual structures resembling the

intact peptidoglycan sacculus remained (Fig 3C), indicating that this is not the target of

Tox-Act1. In addition, we noticed that the cognate immunity protein Imm-Act1 contains

a conserved domain with two transmembrane helices (Fig S3), suggesting that the site

of its neutralizing action occurs at the cell membrane.

Fig 3. Tox-Act1 is a T6SS effector used for bacterial competition in the mouse gut.

(a) Scheme of the genomic region encoding Tox-Act1 and Imm-Act1 effector/immunity

pair (FD01843896). **(b)** *E. coli* toxicity assay. Serial dilutions of *E. coli* carrying pBRA

and pEXT22 constructs. Images are representative of thee independent experiments. **(c)**

Time-lapse microscopy of *E. coli* carrying pBRA SP-Tox-Act1 grown on repressed or

induced conditions. Scale bar: 5 µm. Timestamps in hh:mm. **(d)** C57BL/6 mice were

infected by oral gavage with equal numbers of each strain. Bacteria were recovered

from the cecum 4 days after infection, and CI values calculated. The log10 Cis were

used for statistical analysis. Single sample t-test was used to compare the CI to the

hypothetical value of 0, and *p* value is indicated in brackets. Unpaired t-test (****p <0.01)

was used to compare the two groups.

The SPI-6 T6SS of *Salmonella enterica* serovar Typhimurium is repressed by

the silencer protein H-NS [18]. Bacterial competition assays performed *in vitro* have not

been able to recapitulate the natural conditions found in the mouse gut to fully activate

this system [19]. To analyze whether Tox-Act1 is a T6SS substrate and contributes to

interbacterial competition *in vivo*, we performed competitive index assays using *S.*

Typhimurium strain IC33Q that naturally encodes Tox-Act1 (Fig 3A). Mice were

infected by oral gavage with a 1:1 mixture of wild-type (WT) and the effector-immunity

double mutant Δ*tox-act1/imm-act1*, or a mixture of the T6SS mutant (Δ*tssL*) and Δ*tox-*

act1/imm-act1. Results showed that Δ*tox-act1/imm-act1* has a competitive disadvantage

during colonization of the mouse gut when compared to the WT, and such disadvantage

is not observed when compared to Δ*tssL* (Fig 3D). These results confirm that Tox-Act1

is an antibacterial effector secreted via the SPI-6 T6SS and that it is actively expressed

by this *S. enterica* strain and used during competition in the mouse gut.

Tox-Act1 is evolutionarily related to NlpC/P60 enzymes

targeting lipids

The Tox-Act1 domain was not identified by any of the HMMs deployed in the

initial steps of this study. However, subsequent HHpred analysis revealed a significant

probability of homology with DUF4105 and the effector TseH from *Vibrio*

cholerae[45] (data not shown), both of which are members of the NlpC/P60

superfamily[58, 59]. This superfamily was previously defined as encompassing four

families, which are divided into two higher-order groups (canonical and permuted)[58].

Members of the canonical group (AcmB-like and P60-like) function as peptidases

involved in peptidoglycan hydrolysis[60], while permuted members (YaeF-like and

LRAT-like) exhibit a circular permutation in their catalytic core, creating a hydrophobic

binding pocket that provides specificity for lipids [61, 62]. Closer inspection of the

sequence and structure of Tox-Act1 revealed a circular permutation of the catalytic

domain indicating that it belongs to the permuted NlpC/P60 group

(https://leepbioinfo.github.io/10ksgt6ss/)[58].

To predict the function of Tox-Act1, we sought to understand its evolutionary

relationship by constructing a phylogenetic tree using the sequences of Tox-Act1, TseH

and additional permuted members, such as LRAT and YiiX. The analysis revealed the

formation of five major clades including YiiX, LRAT, TseH, Tox-Act1 and DUF4105

(Fig 4A, Table S2 and Table S3). Genomic context analysis predicted that proteins of

the Tox-Act1 clade are toxins deployed in biological conflicts (Fig 4A and 4B).

Similarly, the TseH clade (Pfam DUF6695) displays genomic contexts indicative of

biological conflicts (Fig 4A and 4B). The clades LRAT and YiiX harbor proteins

known to be involved in lipid metabolism: LRAT (lecithin:retinol acyltransferase) is an

enzyme present in mammals and involved in the transference of acyl groups from

phosphatidylcholine to all-trans retinol to produce all-trans retinyl esters that are storage

forms of vitamin A[63]; H-RAS-like suppressor (HASLS) proteins are a group within

the LRAT family that display both acyltransferase and phospholipase A1/2

activities[64]; and YiiX-like family members from *Bacillus cereus* are active against

lipids[62]. We decided to name Tox-Act1 (toxin acyltransferase 1) due to its

evolutionary relationship with acyltransferases.

Fig 4. Tox-Act1 is evolutionarily related to lipid-targeting enzymes with a permuted NlpC/P60 domain. (a) Maximum-likelihood phylogenetic tree of permuted NlpC/P60 members. Dots represent the number of PSI-BLAST iterations required to collect homologs and the red star marks the query. **(b)** Genomic organization of representatives from clades TseH and Tox-Act1 showing the genes are encoded in the context of conflict systems, and DUF4105 showing context of lipid metabolism. **(c)** Sequence logo from the enzymatic core of permuted NlpC/P60 from all clades shown in (a). The arrows indicate conserved His and Cys residues that were mutated in (d). **(d)** *E. coli* toxicity assay. Serial dilution of *E. coli* containing pBRA and pEXT22 constructs. Images are representative of thee independent experiments.

Interestingly, Tox-Act1 emerged as the sister clade of DUF4105 (Fig 4A and 4B). Our comparative genomics analysis revealed a recurring evolutionary pattern in

which DUF4105 domain-containing proteins are repeatedly displaced by apolipoprotein N-acyltransferases (Lnt) across thee distinct genomic contexts (Fig 4B). Hence, we proposed that DUF4105 could be working as an acyltransferase. Remarkably, DUF4105 was recently identified as the missing lipoprotein N-acyltransferase in *Bacteroides*[65], which was named Lnb (N-acyltransferase in *Bacteroides*). These experimental results confirmed our independent *in-silico* prediction for the function of DUF4105 and provide two different lines of evidence leading to similar conclusions. It is noteworthy that the DUF4105 clade identified in our analysis consists primarily, though not exclusively, of *Bacteroides* species, with a branch enriched in Gram-positives like *Firmicutes* (Fig 3A). The list of homologs containing DUF4105 can be found in Table S3C and S3D.

Multiple sequence alignments of each of the permuted clades including Tox-Act1 revealed the conserved catalytic His and Cys residues characteristic of the NlpC/P60 superfamily (Fig 4C)[58]. Substitution of these residues for alanine (Tox-Act1$_{H43A}$ and Tox-Act1$_{C151A}$) eliminated toxicity in *E. coli* (Fig 4D). These findings collectively confirm that the enzymatic function of the NlpC/P60 papain-like fold domain is crucial for toxicity. Collectively, the periplasmic-acting phenotype of Tox-Act1, the presence of a membrane-associated immunity protein and the fatty acyl linkage targeting activities common in the permuted NlpC/P60 members strongly support a function for Tox-Act1 in targeting membrane lipids.

Tox-Act1 displays phospholipase activity and changes the membrane composition of intoxicated cells

To analyze the enzymatic activity of Tox-Act1, we incubated purified recombinant protein (Fig S4) with either purified phosphatidylglycerol (PG) 16:0-18:1

or phosphatidylethanolamine (PE) 16:0-18:1 and analyzed the reaction product by

HPLC coupled to mass spectrometry (Fig 5A and B). Results showed that Tox-Act1 has

predominantly phospholipase A1 activity and cleaves both PG and PE (a preference for

cleaving off the 16:0 acyl chain) as observed by the accumulation of 18:1

lysophospholipids (Fig 5A and B).

Fig 5. Tox-Act1 has phospholipase A activity and changes the composition of

target cell membranes. (a) *In vitro* enzymatic assay with recombinant Tox-Act1 (red)

or Tox-Act1$_{C151A}$ (blue) incubated with different phospholipids (both 16:0-18:1 PG and

PE). The amount of lysophospholipids produced was analyzed and quantified by HPLC-

MS/MS. **(b)** Quantification of the peak area of lysophospholipids normalized by the

intact substrate. Data corresponds to the mean ± SD. ***p < 0.001 and *p < 0.05, *ns* not

significant (unpaired t-test). **(c)** UHPLC-MS total ion chomatogram showing the profile

of total lipids extracted from *E. coli* expressing Tox-Act1 (red) or Tox-Act1$_{C151A}$ (blue).

(d) Heatmap plot of top 20 altered lipids of intoxicated *E. coli*. Results display four

biological replicates of each condition (WT or C151A) with the quantification of lipids

species (Tukey test; $p < 0.05$ FDR adjusted). Data were expressed in nM mg of protein^{-1}

and normalized by log transformation (base 10) prior to analysis. Red (up) and blue

(down) bars represent changes in lipid species concentration relative to the normalized

mean. Letters differentiate between the isomers.

Next, we set out to determine whether Tox-Act1 could cause changes in the

composition of phospholipids when ectopically expressed by target cells. *E. coli*

harboring the SP-Tox-Act1 plasmid were grown to OD_{600nm} of 1.0 in the presence of d-

glucose (repressed), washed and resuspended in AB medium with l-arabinose to induce

the expression of the toxin. Total lipids were extracted and analyzed by UHPLC-MS.

We observed a general decrease in intact phospholipid forms in Tox-Act1$_{WT}$, especially

phosphatidylglycerol (PG), when compared with the catalytic mutant Tox-Act1$_{C151A}$

(Fig 5A and 5B). In addition, an increase in lysophospholipids forms - either

lysophosphatidylglycerol (LPG) or lysophosphatidylethanolamine (LPE) - and free fatty

acids (FFA) was detected in the wild-type (Fig 5A and 5B). Lysophospholipids possess

amphiphilic properties and have an inverted cone-shaped molecular structure that

interacts with and modifies membrane properties (e.g. curvature) similarly to

detergents[66]. The accumulation of lysophospholipids on the target cell membrane

likely promotes the observed membrane disruption (Fig 3C) due to its detergent-like

properties[67]. Collectively, these results confirm that Tox-Act1 targets phospholipids

similarly to other proteins possessing a permuted NlpC/P60 domain, and that Tox-Act1

displays phospholipid-degrading activity. It remains to be determined whether Tox-

Act1 also possesses acyltransferase activity as reported for other permuted NlpC/P60.

Discussion

Our comprehensive analysis of 10,000 *Salmonella* genomes has significantly expanded the repertoire of T6SS effectors and polymorphic toxins in general, identifying 128 candidates, among which 45 comprise novel protein domains. Our study employed a robust bioinformatic pipeline, integrating classical methods with the latest structural bioinformatics techniques. The combination of sensitive sequence and structure searches with comparative genomics provided a comprehensive understanding of the identified toxin domains. In addition, the manual curation of candidates ensured high confidence in our results, distinguishing our approach from previous large-scale *in-silico* analyses. The identification of these novel toxin domains builds upon the foundational work by Zhang et al. [9] and others [30, 68-76], which characterized the diversity of polymorphic toxin systems across bacterial lineages. The identification of novel toxin domains highlights the constant evolutionary arms race between bacteria, driving the diversification of toxin and immunity proteins.

This study not only broadens our understanding of *Salmonella* T6SS effectors and toxin domains but also provides the first direct characterization of a lipid-targeting NlpC/P60 domain. The phospholipase activity of Tox-Act1, which cleaves acyl groups from phosphatidylglycerol and phosphatidylethanolamine, underscores its role in membrane disruption during bacterial competition. Interestingly, our phylogenetic analysis revealed an evolutionary link between Tox-Act1 and a new family of lipoprotein N-acyltransferases in *Bacteroides*, adding a new dimension to the functional diversity of these toxins. Notably, one of the homologs of Tox-Act1 is TseH [45, 77], which has been proposed to be an endopeptidase due to its NlpC/P60 domain and similarity to the amidase Tse1[78, 79]. However, unlike Tse1, TseH exhibits a permutation in its catalytic core [77]. This permutation, along with its evolutionary

452   relationship to Tox-Act1 and other permuted NlpC/P60, suggest that TseH actual

453   substrate might be an acyl group in phospholipids rather than a peptide/amide bond.

454   Exploring the potential acyltransferase activity of Tox-Act1 and its homologs could

455   reveal further biochemical diversity within the NlpC/P60 superfamily.

456   In the context of *Salmonella* biology, the unprecedented diversity of T6SS

effectors presents numerous opportunities for new studies. Our findings reveal the

existence of individual subsets of T6SS effectors for each serovar, suggesting that

Salmonella acquire and maintain effectors in response to specific environmental

pressures rather than accumulating an increasingly larger array of effectors. Notably, we

observed a higher number of effectors in serovars isolated from environmental sources

compared to those from patients, indicating that the number of effectors increases in

more diverse environments where there are potentially more encounters with a variety

of rival species. In addition, we observed that *Salmonella* serovars encode

phylogenetically distinct T6SS clusters, which are specialized to target either eukaryotic

or bacterial cells. These results further highlight the versatility of the T6SS aiding in

adaptation to many environments and hosts.

468   In conclusion, our comprehensive analysis has greatly enhanced the

469   understanding of toxins involved in bacterial competition and pathogenesis. The

470   identification of previously uncharacterized toxin domains highlights the potential for

471   discovering novel biochemical activities. This study provides a solid foundation for

472   future research into the complex dynamics of conflict systems and their implications for

473   bacterial ecology and pathogenesis.

Materials and Methods

Comparative genomic analysis

The .gff files from the genome assemblies retrieved from the 10KSG project[23]

were organized and stored in tabular format using Python scripts, based on the

Biopython[80] and pandas[81] libraries. Iterative searches were conducted using

Jackhmmer[82] with a 10e-6 e-value cutoff. Protein clustering was performed using

MMseqs[83] to remove redundancy (80% coverage and 70% identity) and form

homologous groups (80% coverage and e-value 10e-3). Multiple sequence alignments

were generated using the local-pair algorithm in MAFFT[84], and phylogenetic trees

were constructed using FastTree[85]. Domain identification and annotation was

performed using HMMsearch and HMMscan[82, 86] and models from the databases

Pfam[25], TXSScan[24], and BastionHub[87]. Remote homology identification was

performed using HHpred[33] and FoldSeek.

Bacterial strains

A list of bacterial strains used in this work can be found in Table S4. Strains

were grown at 37 °C in Lysogeny Broth (10 g/L tryptone, 10 g/L NaCl, 5 g/L yeast

extract) under agitation. AB medium was used for lipidomics: 0.2% (NH4)2SO2, 0.6%

Na2HPO4, 0.3% KH2PO4, 0.3% NaCl, 0.1 mM CaCL2, 1 mM MgCl2, 3 µM FeCL3,

supplemented with 0.2% sucrose, 0.2% casamino acids, 10 µg/mL thiamine, and 25

µg/mL uracil. Cultures were supplemented with antibiotics in the following

concentration when necessary: 50 µg/mL kanamycin, 100 µg/mL ampicillin, and 50

µg/mL streptomycin.

Cloning and mutagenesis

498   All primers are listed in Table S4. Tox-Act1 and Imm-Act1 were amplified by

499   PCR and cloned into pBRA vector under the control of P_{BAD} promoter[88] with or

500   without pelB signal peptide sequence from pET22b (Novagen)[89]. Imm-Act1 was

501   cloned into pEXT22 under the control of P_{TAC} promoter[90]. For protein expression and

purification, Tox-Act1 was cloned into pET28a (Novagen), including a C-terminal Strep II tag. Point mutations (Tox-Act1$_{H43A}$, Tox-Act1$_{C151A}$) were created using QuikChange II XL Site-Directed Mutagenesis Kit (Agilent Technologies) or by splicing by overlap extension (SOE) PCR. All constructs were confirmed by sequencing. *S.* Typhimurium mutant strains used for competition assays were constructed by λ-Red recombination system using a one-step inactivation procedure [91].

E. coli toxicity assay

Overnight cultures of *E. coli* DH5α co-expressing effectors for cytoplasmic (pBRA Tox-Act1) or periplasmic (pBRA SP-Tox-Act1) localization and immunity protein (pEXT22 Imm-Act1) were adjusted to OD$_{600nm}$ 1, serially diluted in LB (1:4) and 5 μL were spotted onto LB agar (1.5%) containing either 0.2% d-glucose or 0.2% l-arabinose plus 200 μM IPTG, supplemented with streptomycin and kanamycin, and incubated at 37°C for 20 h.

Time-lapse microscopy

For time-lapse microscopy, LB agar (1.5%) pads were prepared by cutting a rectangular piece out of a double-sided adhesive tape, which was taped onto a microscopy slide as described previously[89]. *E. coli* DH5α harboring pBRA SP-Tox-Act1 was subcultured in LB (1:50) with 0.2% d-glucose until reaching OD$_{600nm}$ 0.4–0.6 and adjusted to OD$_{600nm}$ 1. Cultures were spotted onto LB agar pads supplemented either with 0.2% d-glucose or 0.2% l-arabinose plus antibiotics. Images were acquired every 15 min for 16 h using a Leica DMi-8 epifluorescent microscope fitted with a DFC365 FX camera (Leica) and Plan-Apochomat ×63 oil objective (HC PL APO ×63/1.4 Oil ph3 objective Leica). Images were analyzed using FIJI software [92].

Competitive index

Female C57BL/6 mice (6-8 weeks old) were purchased from Jackson's Laboratory. Mice were housed in pathogenic free conditions with unlimited access to food and water, except for 4 hours prior oral gavages. Mice were pre-treated with 20 mg of streptomycin 24 hours prior infection with *Salmonella*. Mice were infected by oral gavage with a 1:1 mixture (total of 10^{10} CFUs) of *S*. Typhimurium wild-type (WT) and Δtox-*act1*/*imm-act1* (Cm^R), or a mixture of $\Delta tssL$ and Δtox-*act1*/*imm-act1*. Mice were euthanized four days after infection by CO_2 asphyxiation, and cecum were harvested and the content serial diluted in PBS 1x (Phosphate-buffered saline) and plated in MacConkey Agar to determine the total CFU counts. One hundred colonies from each mouse were patched on chloramphenicol plates to determine the proportion of each strain in the mixture. The same procedure was performed with the initial mixture prior infection. Competitive index was calculated as previously described [93]. The unpaired t-test was used to compare the CI between WT and $\Delta tssL$ groups, while the single sample t-test was used to compare each log10 CI to the hypothetical value of 0 (the value of 0 means that two strains grew equally well *in vivo*).

Protein expression and purification

E. coli BL21(DE3) carrying pET28a Tox-Act1$_{WT}$-Strep or Tox-Act1$_{C151A}$-Strep were grown in 4 L of LB supplemented with kanamycin (37 °C, 180 rpm) until OD$_{600nm}$ 0.7. Expression was induced with 1 mM IPTG for 16 h at 16 °C. Cells were harvested via centrifugation at 5000 *g* for 20 min, and pellets were resuspended in lysis buffer (50 mM Tris HCl pH 8.0, 350 mM NaCl, 45 mM β-mercaptoethanol, 5 mg/mL lysozyme, 10% glycerol) and lysed at 4°C using a sonicator. The lysate was centrifuged at 40,000 *g* for 45 min at 4°C. The supernatant was loaded onto a 1 ml StrepTrap HP column (Cytiva) equilibrated in buffer (50 mM Tris-HCl pH 8.0, 350 mM NaCl, 10% glycerol). The column was washed with 40 column volumes (CV) of wash buffer (50 mM Tris-

HCl pH 8.0, 1 M NaCl, 10% glycerol, 45 mM β-mercaptoethanol), followed by a

second wash with 12 CV (50 mM Tris-HCl pH 8.0, 1.5 M urea) to remove chaperonin

GroEL[94]. The column was subjected to a third round of washes with 40 CV of wash

buffer and eluted with 10 CV of elution buffer (50 mM Tris-HCl pH 8.0; 350 mM

NaCl; 10% glycerol; 50 mM biotin). Fractions were buffer exchanged (25mM Tris-HCl

pH 7.5, 100 mM NaCl, 5% glycerol) and concentrated using an Amicon of 30 kDa

(Sigma). Protein aliquots were snap-frozen until use.

***In vitro* phospholipase assay**

For *in vitro* enzymatic assay, phospholipids 1-palmitoyl-2-oleoyl-sn-glycero-3-

phospho-(1'-rac-glycerol) (PG 16:0-18:1) and 1-palmitoyl-2-oleoyl-sn-glycero-3-

phosphoethanolamine (PE 16:0-18:1) were purchased from Avanti Polar Lipids.

Substrates were resuspended and diluted in methanol to adjust the concentration of

aliquots. The methanol of each aliquot was dried under a nitrogen flow. A total of 1.2

mM of phospholipids (PG or PE) were resuspended in reaction buffer (25 mM Tris-HCl

pH 7.5, 100 mM NaCl, 0.5 mM $CaCl_2$, 0.5 mM $MgCl_2$, 180 mM sodium deoxycholate

and 0.5 mM DTT) and incubated with 0.8 mM of enzyme (Tox-Act1$_{WT}$ or Tox-

Act1$_{C151A}$) in a total volume of 100 µL for 2 h at 37°C under agitation (350 rpm). Lipids

were extracted by adding 830 µL of a mixture of MTBE/methanol/water (10:3:2.5,

v/v/v), followed by incubation under agitation for 1 h at room temperature. Samples

were centrifuged for 2 min at 220 *g* and 350 µL of the top fraction was transferred to a

new tube, dried in a SpeedVac and stored at –80 °C until analysis.

For mass spectrometry analysis, samples were resuspended in 350 µL of

isopropanol and analyzed in a Shimadzu 8060 Triple Quadrupole Liquid Chomatograph

Mass Spectrometer. Samples (0.1-0.5 µL) were loaded into an Agilent column C18

ZORBAX Eclipse Plus (4.6 x 150 mm, 5 µm, 400 bar) with a flow rate of 0.5 mL/min

and an oven temperature of 40 °C. HPLC gradients were as described below for

lipidomic analysis. The phospholipids and lysophospholipids of interest were analyzed

in the multiple reaction monitoring (MRM) mode using m/z transitions, collision

energies (CE) and dwell times as shown in Table S5A. Data was acquired by Shimadzu

LabSolutions and processed in LabSolutions Browser. Graphs were plotted using

GraphPrism 5.

Lipidomics of Tox-Act1-intoxicated *E. coli*

E. coli MG1655 containing the plasmids pBRA SP-Tox-Act1$_{WT}$ or SP-Tox-

Act1$_{C151A}$ were cultured in LB containing 0.2% d-glucose at 37 °C for 14 h. Cells were

subcultured in LB 0.2% d-glucose until an OD$_{600nm}$ of 1 and centrifuged (10 min, 2900

g, 30 °C). Cells were washed with 40 mL of preheated AB medium at 37 °C,

centrifuged (10 min, 2900 *g*, 30 °C), and resuspended in 5 mL of AB supplemented

with 0.2% l-arabinose to induce Tox-Act1 expression. Cells were incubated for 1 h at

37 °C with agitation (100 rpm). The cells were centrifuged (15 min, 2900 *g*, 4 °C) and

washed once with 1 mL of PBS pH 7.4. PBS was removed by centrifugation and the

cell pellet was stored at -80 °C until lipid extraction. A cocktail of class specific internal

standards was added to the cell mass prior to lipid extraction for subsequent

quantification and normalization (Table S5B). Total lipid extraction was performed

using an adapted version of the protocol described by Yoshida, Kodai (95). Briefly, cell

pellets were resuspended with 500 µL of cold methanol and 1 mL of MilliQ water and

transferred to glass tubes. A mixture of chloroform and ethyl acetate (4:1) was added,

followed by agitation for 1 min at 25 °C. Samples were centrifuged (2 min, 1500 *g*, 4

°C) and the lipid-containing phase (lower phase) was extracted and transferred to a new

glass tube that was dried under a nitrogen (N$_2$) flow until all solvent traces were

evaporated. Samples were stored at -80 °C until analysis.

Lipid extracts were diluted in 100 μL of isopropanol and analyzed using ultra-

high performance liquid chomatography (UHPLC Nexera, Shimadzu) coupled with an

ESI-Q-TOF mass spectrometer (Triple TOF 6600, Sciex) (UHPLC-Q-TOF/MS). 2 μL

of each sample were injected into the UHPLC-MS, and molecules were separated using

a CORTECS column (C18, 1.6 μm, 2.1x100 mm, Waters) with a flow rate of 0.2

mL/min and temperature set to 35 °C [96]. The mobile phases consisted of (A)

water/acetonitrile (60:40) and (B) isopropanol/acetonitrile/water (88:10:2). Ammonium

acetate at a final concentration of 10 mM was incorporated in both mobile phases A and

B for the negative ionization acquisition mode. The gradient elution used in the

chomatography was from 40 to 100% (mobile phase) B over the first 10 min; 100% B

from 10-12 min; 100 to 40% B for 12-13 and holding 40% B for 13-20 min. The

negative mode was utilized for the examination of phospholipids and free fatty acids.

MS and MS/MS data acquisition was performed using Analyst 1.7.1 software (Sciex).

Mass spectrometry data was inspected using PeakView 2.0 software (Sciex), and lipid

molecular species were manually identified with the help of an in-house manufactured

Excel-based macro. Lipid species were quantified using MultiQuant software (Sciex),

where the precursor ions areas were normalized by the internal standards for each class

(Table S5B).

Data availability

All data supporting the findings of this study are available within the paper and

its Supplementary Information files or at https://leepbioinfo.github.io/10ksgt6ss/. All

data and code used for sequence and genome context analyses are available on a GitHub

repository at https://github.com/leepbioinfo/10ksgt6ss. The ROTIFER package can be

downloaded from the GitHub repository https://github.com/leepbioinfo/rotifer. The

contents of both repositories are made available under the Creative Commons

Attribution 4.0 International or the GNU Lesser General Public License (LGPL) v. 2.0.

We have also uploaded the GitHub data and code to Zenodo (DOI:

10.5281/zenodo.13845778).

Ethics Statement

The animal experiments were performed with protocols approved by the

University of Texas at Austin, Institutional Animal Care and Use Committee. The

University of Texas at Austin animal management program is accredited by the

Association for the Assessment and Accreditation of Laboratory Animal Care,

International (AAALAC), and meets National Institutes of Health standards as set forth

in the Guide for the Care and Use of Laboratory Animals (DHHS Publication No. (NIH)

85–23 Revised 1996).

Acknowledgments

We are grateful to Jay Hinton for initially sharing the 10KSG dataset with us, to

Ian Riddington for assistance with mass-spectrometry analysis of phospholipids, and

Gerd Prehna and Larissa Diniz for discussions about enzymatic assay. Marcos

Yoshinaga from PinguisLab for lipidomic data processing. Edgar Llontop and Diorge

Paulo Souza for troubleshooting with protein purification. Cristiano G. Moreira for

sharing strains. Alexandre Bruni for access to the microscope. Luize Nobrega Silva and

Gustavo Chagas for technical support. This research was supported by Sao Paulo

Research Foundation grant #2017-02178-2 to E.B-S, #2013/07937-8 to S.M,

#2016/09047-8 and # 2021/10577-0 to R.F.S and #2021/10577-0 to T.W.C.S. Startup

funds from MBS and CNS to E.B-S. This research was supported in part by an

appointment to the National Library of Medicine Research Participation Program

administered by the Oak Ridge Institute for Science and Education (ORISE) though an

interagency agreement between the U.S. Department of Energy (DOE) and the National

Library of Medicine. ORISE is managed by ORAU under DOE contract number DE-

SC0014664. All opinions expressed in this paper are the author's and do not necessarily

reflect the policies and views of NIH, NLM, DOE, or ORAU/ORISE. This work is

supported by the funds of the Intramural Research Program of National Library of

Medicine at the National Institutes of Health, USA (to L.A. and G.G.N). This work

utilized the NIH HPC Biowulf computer cluster.

References

1.   Granato ET, Meiller-Legrand TA, Foster KR. The Evolution and Ecology of
Bacterial Warfare. Curr Biol. 2019;29(11):R521-R37. doi: 10.1016/j.cub.2019.04.024.
PubMed PMID: 31163166.
2.   Hood RD, Peterson SB, Mougous JD. From Striking Out to Striking Gold:
Discovering that Type VI Secretion Targets Bacteria. Cell host & microbe.
2017;21(3):286-9. doi: 10.1016/j.chom.2017.02.001. PubMed PMID: 28279332;
PubMed Central PMCID: PMCPMC6404758.
3.   Russell AB, Wexler AG, Harding BN, Whitney JC, Bohn AJ, Goo YA, et al. A
type VI secretion-related pathway in Bacteroidetes mediates interbacterial antagonism.
Cell   host   &   microbe.   2014;16(2):227-36.   Epub   20140725.   doi:
10.1016/j.chom.2014.07.007. PubMed PMID: 25070807; PubMed Central PMCID:
PMCPMC4136423.
4.   Li J, Yao Y, Xu HH, Hao L, Deng Z, Rajakumar K, et al. SecReT6: a web-based
resource for type VI secretion systems found in bacteria. Environ Microbiol.
2015;17(7):2196-202. doi: 10.1111/1462-2920.12794. PubMed PMID: 25640659.
5.   Bock D, Medeiros JM, Tsao HF, Penz T, Weiss GL, Aistleitner K, et al. In situ
architecture, function, and evolution of a contractile injection system. Science.
2017;357(6352):713-7. doi: 10.1126/science.aan7904. PubMed PMID: 28818949;
PubMed Central PMCID: PMCPMC6485382.
6.   Boyer F, Fichant G, Berthod J, Vandenbrouck Y, Attree I. Dissecting the
bacterial type VI secretion system by a genome wide in silico analysis: what can be
learned from available microbial genomic resources? BMC Genomics. 2009;10:104.
Epub 2009/03/17. doi: 10.1186/1471-2164-10-104. PubMed PMID: 19284603; PubMed
Central PMCID: PMCPMC2660368.
7.   Barret M, Egan F, Fargier E, Morrissey JP, O'Gara F. Genomic analysis of the
type VI secretion systems in Pseudomonas spp.: novel clusters and putative effectors
uncovered. Microbiology (Reading). 2011;157(Pt 6):1726-39. Epub 20110407. doi:
10.1099/mic.0.048645-0. PubMed PMID: 21474537.

8.   Hood RD, Singh P, Hsu F, Guvener T, Carl MA, Trinidad RR, et al. A type VI
secretion system of Pseudomonas aeruginosa targets a toxin to bacteria. Cell host &
microbe. 2010;7(1):25-37. doi: 10.1016/j.chom.2009.12.007. PubMed PMID:
20114026; PubMed Central PMCID: PMCPMC2831478.
9.   Zhang D, de Souza RF, Anantharaman V, Iyer LM, Aravind L. Polymorphic
toxin systems: Comprehensive characterization of trafficking modes, processing,
mechanisms of action, immunity and ecology using comparative genomics. Biol Direct.
2012;7:18. Epub 20120625. doi: 10.1186/1745-6150-7-18. PubMed PMID: 22731697;
PubMed Central PMCID: PMCPMC3482391.
10.   Cherrak Y, Rapisarda C, Pellarin R, Bouvier G, Bardiaux B, Allain F, et al.
Biogenesis and structure of a type VI secretion baseplate. Nat Microbiol.
2018;3(12):1404-16. Epub 2018/10/17. doi: 10.1038/s41564-018-0260-1. PubMed
PMID: 30323254.
11.   Jana B, Salomon D. Type VI secretion system: a modular toolkit for bacterial
dominance. Future Microbiol. 2019;14:1451-63. Epub 20191113. doi: 10.2217/fmb-
2019-0194. PubMed PMID: 31718312.
12.   Allsopp LP, Bernal P. Killing in the name of: T6SS structure and effector
diversity. Microbiology (Reading). 2023;169(7). doi: 10.1099/mic.0.001367. PubMed
PMID: 37490402; PubMed Central PMCID: PMCPMC10433429.
13.   Bohnhoff M, Drake BL, Miller CP. Effect of streptomycin on susceptibility of
intestinal tract to experimental Salmonella infection. Proc Soc Exp Biol Med.
1954;86(1):132-7. doi: 10.3181/00379727-86-21030. PubMed PMID: 13177610.
14.   Rogers AWL, Tsolis RM, Baumler AJ. Salmonella versus the Microbiome.
Microbiol Mol Biol Rev. 2021;85(1). Epub 20201223. doi: 10.1128/MMBR.00027-19.
PubMed PMID: 33361269; PubMed Central PMCID: PMCPMC8549850.
15.   Sibinelli-Sousa S, de Araujo-Silva AL, Hespanhol JT, Bayer-Santos E.
Revisiting the steps of Salmonella gut infection with a focus on antagonistic
interbacterial interactions. FEBS J. 2021. Epub 2021/09/22. doi: 10.1111/febs.16211.
PubMed PMID: 34546626.
16.   Blondel CJ, Jimenez JC, Contreras I, Santiviago CA. Comparative genomic
analysis uncovers 3 novel loci encoding type six secretion systems differentially
distributed in Salmonella serotypes. BMC Genomics. 2009;10:354. Epub 20090804.
doi: 10.1186/1471-2164-10-354. PubMed PMID: 19653904; PubMed Central PMCID:
PMCPMC2907695.
17.   Bao H, Zhao JH, Zhu S, Wang S, Zhang J, Wang XY, et al. Genetic diversity
and evolutionary features of type VI secretion systems in Salmonella. Future Microbiol.
2019;14:139-54. Epub 20190123. doi: 10.2217/fmb-2018-0260. PubMed PMID:
30672329.
18.   Brunet YR, Khodr A, Logger L, Aussel L, Mignot T, Rimsky S, et al. H-NS
Silencing of the Salmonella Pathogenicity Island 6-Encoded Type VI Secretion System
Limits Salmonella enterica Serovar Typhimurium Interbacterial Killing. Infection and
immunity. 2015;83(7):2738-50. Epub 20150427. doi: 10.1128/IAI.00198-15. PubMed
PMID: 25916986; PubMed Central PMCID: PMCPMC4468533.
19.   Sana TG, Flaugnatti N, Lugo KA, Lam LH, Jacobson A, Baylot V, et al.
Salmonella Typhimurium utilizes a T6SS-mediated antibacterial weapon to establish in
the host gut. Proceedings of the National Academy of Sciences of the United States of
America. 2016;113(34):E5044-51. Epub 20160808. doi: 10.1073/pnas.1608858113.
PubMed PMID: 27503894; PubMed Central PMCID: PMCPMC5003274.
20.   Sibinelli-Sousa S, Hespanhol JT, Nicastro GG, Matsuyama BY, Mesnage S,
Patel A, et al. A Family of T6SS Antibacterial Effectors Related to l,d-Transpeptidases

Targets the Peptidoglycan. Cell Rep. 2020;31(12):107813. doi: 10.1016/j.celrep.2020.107813. PubMed PMID: 32579939.

21. Hespanhol JT, Sanchez-Limache DE, Nicastro GG, Mead L, Llontop EE, Chagas-Santos G, et al. Antibacterial T6SS effectors with a VRR-Nuc domain are structure-specific nucleases. Elife. 2022;11. Epub 2022/10/14. doi: 10.7554/eLife.82437. PubMed PMID: 36226828; PubMed Central PMCID: PMCPMC9635880.

22. Aravind L. Guilt by association: contextual information in genome analysis. Genome Res. 2000;10(8):1074-7. doi: 10.1101/gr.10.8.1074. PubMed PMID: 10958625.

23. Perez-Sepulveda BM, Heavens D, Pulford CV, Predeus AV, Low R, Webster H, et al. An accessible, efficient and global approach for the large-scale sequencing of bacterial genomes. Genome Biol. 2021;22(1):349. Epub 20211221. doi: 10.1186/s13059-021-02536-3. PubMed PMID: 34930397; PubMed Central PMCID: PMCPMC8690886.

24. Abby SS, Cury J, Guglielmini J, Neron B, Touchon M, Rocha EP. Identification of protein secretion systems in bacterial genomes. Sci Rep. 2016;6:23080. Epub 20160316. doi: 10.1038/srep23080. PubMed PMID: 26979785; PubMed Central PMCID: PMCPMC4793230.

25. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al. Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49(D1):D412-D9. doi: 10.1093/nar/gkaa913. PubMed PMID: 33125078; PubMed Central PMCID: PMCPMC7779014.

26. Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, et al. The conserved domain database in 2023. Nucleic Acids Res. 2023;51(D1):D384-D8. doi: 10.1093/nar/gkac1096. PubMed PMID: 36477806; PubMed Central PMCID: PMCPMC9825596.

27. Jaccard P. The Distribution of the Flora in the Alpine Zone. New Phytologist. 1912;11:37-50. doi: 10.1111/J.1469-8137.1912.TB05611.X.

28. De Meo PF, E.; Fiumara, G.; Provetti, A. Generalized Louvain method for community detection in large networks. International Conference on Intelligent Systems Design and Applications. 2011. doi: 10.1109/ISDA.2011.6121636.

29. Hansen-Wester I, Hensel M. Salmonella pathogenicity islands encoding type III secretion systems. Microbes and infection / Institut Pasteur. 2001;3(7):549-59. doi: 10.1016/s1286-4579(01)01411-3. PubMed PMID: 11418329.

30. Wang J, Yang B, Leier A, Marquez-Lago TT, Hayashida M, Rocker A, et al. Bastion6: a bioinformatics approach for accurate prediction of type VI secreted effectors. Bioinformatics. 2018;34(15):2546-55. doi: 10.1093/bioinformatics/bty155. PubMed PMID: 29547915; PubMed Central PMCID: PMCPMC6061801.

31. Zhang J, Guan J, Wang M, Li G, Djordjevic M, Tai C, et al. SecReT6 update: a comprehensive resource of bacterial Type VI Secretion Systems. Sci China Life Sci. 2022. Epub 2022/11/09. doi: 10.1007/s11427-022-2172-x. PubMed PMID: 36346548.

32. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res. 1997;25(17):3389-402. doi: 10.1093/nar/25.17.3389. PubMed PMID: 9254694; PubMed Central PMCID: PMCPMC146917.

33. Soding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology detection and structure prediction. Nucleic Acids Res. 2005;33(Web Server issue):W244-8. Epub 2005/06/28. doi: 10.1093/nar/gki408. PubMed PMID: 15980461; PubMed Central PMCID: PMCPMC1160169.

34.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al.
Highly accurate protein structure prediction with AlphaFold. Nature.
2021;596(7873):583-9. Epub 2021/07/16. doi: 10.1038/s41586-021-03819-2. PubMed
PMID: 34265844; PubMed Central PMCID: PMCPMC8371605.
35.   van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchist CLM, et al.
Fast and accurate protein structure search with Foldseek. Nat Biotechnol.
2024;42(2):243-6. Epub 20230508. doi: 10.1038/s41587-023-01773-0. PubMed PMID:
37156916; PubMed Central PMCID: PMCPMC10869269.
36.   Holm L. Using Dali for Protein Structure Comparison. Methods Mol Biol.
2020;2112:29-42. Epub 2020/02/02. doi: 10.1007/978-1-0716-0270-6_3. PubMed
PMID: 32006276.
37.   Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, et al. The
Protein Data Bank. Nucleic Acids Res. 2000;28(1):235-42. doi: 10.1093/nar/28.1.235.
PubMed PMID: 10592235; PubMed Central PMCID: PMCPMC102472.
38.   Varadi M, Bertoni D, Magana P, Paramval U, Pidruchna I, Radhakrishnan M, et
al. AlphaFold Protein Structure Database in 2024: providing structure coverage for over
214 million protein sequences. Nucleic Acids Res. 2024;52(D1):D368-D75. doi:
10.1093/nar/gkad1011. PubMed PMID: 37933859; PubMed Central PMCID:
PMCPMC10767828.
39.   Russell AB, Singh P, Brittnacher M, Bui NK, Hood RD, Carl MA, et al. A
widespread bacterial type VI secretion effector superfamily identified using a heuristic
approach. Cell host & microbe. 2012;11(5):538-49. Epub 2012/05/23. doi:
10.1016/j.chom.2012.04.007. PubMed PMID: 22607806; PubMed Central PMCID:
PMCPMC3358704.
40.   Jurenas D, Rey M, Byrne D, Chamot-Rooke J, Terradot L, Cascales E.
Salmonella antibacterial Rhs polymorphic toxin inhibits translation though ADP-
ribosylation of EF-Tu P-loop. Nucleic Acids Res. 2022;50(22):13114-27. Epub
2022/12/10. doi: 10.1093/nar/gkac1162. PubMed PMID: 36484105; PubMed Central
PMCID: PMCPMC9825190.
41.   Mariano G, Trunk K, Williams DJ, Monlezun L, Strahl H, Pitt SJ, et al. A family
of Type VI secretion system effector proteins that form ion-selective pores. Nat
Commun. 2019;10(1):5484. Epub 2019/12/04. doi: 10.1038/s41467-019-13439-0.
PubMed PMID: 31792213; PubMed Central PMCID: PMCPMC6889166.
42.   Hu H, Zhang H, Gao Z, Wang D, Liu G, Xu J, et al. Structure of the type VI
secretion phospholipase effector Tle1 provides insight into its hydrolysis and membrane
targeting. Acta Crystallogr D Biol Crystallogr. 2014;70(Pt 8):2175-85. Epub 20140725.
doi: 10.1107/S1399004714012899. PubMed PMID: 25084336.
43.   Bernal P, Allsopp LP, Filloux A, Llamas MA. The Pseudomonas putida T6SS is
a plant warden against phytopathogens. ISME J. 2017;11(4):972-87. Epub 20170103.
doi: 10.1038/ismej.2016.169. PubMed PMID: 28045455; PubMed Central PMCID:
PMCPMC5363822.
44.   Miyata ST, Kitaoka M, Brooks TM, McAuley SB, Pukatzki S. Vibrio cholerae
requires the type VI secretion system virulence factor VasX to kill Dictyostelium
discoideum. Infection and immunity. 2011;79(7):2941-9. Epub 20110509. doi:
10.1128/IAI.01266-10. PubMed PMID: 21555399; PubMed Central PMCID:
PMCPMC3191968.
45.   Altindis E, Dong T, Catalano C, Mekalanos J. Secretome analysis of Vibrio
cholerae type VI secretion system reveals a new effector-immunity pair. mBio.
2015;6(2):e00075. Epub 20150310. doi: 10.1128/mBio.00075-15. PubMed PMID:
25759499; PubMed Central PMCID: PMCPMC4453574.

46.　　Whitney JC, Chou S, Russell AB, Biboy J, Gardiner TE, Ferrin MA, et al.
Identification, structure, and function of a novel type VI secretion peptidoglycan
glycoside hydrolase effector-immunity pair. The Journal of biological chemistry.
2013;288(37):26616-24. Epub 20130722. doi: 10.1074/jbc.M113.488320. PubMed
PMID: 23878199; PubMed Central PMCID: PMCPMC3772208.
47.　　Bullen NP, Sychantha D, Thang SS, Culviner PH, Rudzite M, Ahmad S, et al.
An ADP-ribosyltransferase toxin kills bacterial cells by modifying structured non-
coding RNAs. Molecular cell. 2022;82(18):3484-98 e11. Epub 20220906. doi:
10.1016/j.molcel.2022.08.015. PubMed PMID: 36070765.
48.　　Li DY, Liu YL, Liao XJ, He TT, Sun SS, Nie P, et al. Identification and
Characterization of EvpQ, a Novel T6SS Effector Encoded on a Mobile Genetic
Element in Edwardsiella piscicida. Front Microbiol. 2021;12:643498. Epub 20210311.
doi: 10.3389/fmicb.2021.643498. PubMed PMID: 33776977; PubMed Central PMCID:
PMCPMC7991086.
49.　　Berni B, Soscia C, Djermoun S, Ize B, Bleves S. A Type VI Secretion System
Trans-Kingdom Effector Is Required for the Delivery of a Novel Antibacterial Toxin in
Pseudomonas aeruginosa. Front Microbiol. 2019;10:1218. Epub 20190605. doi:
10.3389/fmicb.2019.01218. PubMed PMID: 31231326; PubMed Central PMCID:
PMCPMC6560169.
50.　　Yah TL, Vallis AJ, Hancock MK, Barbieri JT, Frank DW. ExoY, an adenylate
cyclase secreted by the Pseudomonas aeruginosa type III system. Proceedings of the
National Academy of Sciences of the United States of America. 1998;95(23):13899-
904. doi: 10.1073/pnas.95.23.13899. PubMed PMID: 9811898; PubMed Central
PMCID: PMCPMC24955.
51.　　Poole SJ, Diner EJ, Aoki SK, Braaten BA, t'Kint de Roodenbeke C, Low DA, et
al. Identification of functional toxin/immunity genes linked to contact-dependent
growth inhibition (CDI) and rearrangement hotspot (Rhs) systems. PLoS Genet.
2011;7(8):e1002217. Epub 20110804. doi: 10.1371/journal.pgen.1002217. PubMed
PMID: 21829394; PubMed Central PMCID: PMCPMC3150448.
52.　　Russell AB, LeRoux M, Hathazi K, Agnello DM, Ishikawa T, Wiggins PA, et al.
Diverse type VI secretion phospholipases are functionally plastic antibacterial effectors.
Nature. 2013;496(7446):508-12. Epub 20130403. doi: 10.1038/nature12074. PubMed
PMID: 23552891; PubMed Central PMCID: PMCPMC3652678.
53.　　Koskiniemi S, Lamoureux JG, Nikolakakis KC, t'Kint de Roodenbeke C, Kaplan
MD, Low DA, et al. Rhs proteins from diverse bacteria mediate intercellular
competition. Proceedings of the National Academy of Sciences of the United States of
America. 2013;110(17):7032-7. Epub 20130409. doi: 10.1073/pnas.1300627110.
PubMed PMID: 23572593; PubMed Central PMCID: PMCPMC3637788.
54.　　Jiang F, Waterfield NR, Yang J, Yang G, Jin Q. A Pseudomonas aeruginosa type
VI secretion phospholipase D effector targets both prokaryotic and eukaryotic cells.
Cell host & microbe. 2014;15(5):600-10. doi: 10.1016/j.chom.2014.04.010. PubMed
PMID: 24832454.
55.　　Koskiniemi S, Garza-Sanchez F, Sandegren L, Webb JS, Braaten BA, Poole SJ,
et al. Selection of orphan Rhs toxin expression in evolved Salmonella enterica serovar
Typhimurium. PLoS Genet. 2014;10(3):e1004255. Epub 20140327. doi:
10.1371/journal.pgen.1004255. PubMed PMID: 24675981; PubMed Central PMCID:
PMCPMC3967940.
56.　　Quentin D, Ahmad S, Shanthamoorthy P, Mougous JD, Whitney JC, Raunser S.
Mechanism of loading and translocation of type VI secretion system effector Tse6. Nat

Microbiol. 2018;3(10):1142-52. Epub 2018/09/05. doi: 10.1038/s41564-018-0238-z.
PubMed PMID: 30177742; PubMed Central PMCID: PMCPMC6488228.

57. Ahmad S, Tsang KK, Sachar K, Quentin D, Tashin TM, Bullen NP, et al.
Structural basis for effector transmembrane domain recognition by type VI secretion
system chaperones. Elife. 2020;9. Epub 2020/12/16. doi: 10.7554/eLife.62816. PubMed
PMID: 33320089; PubMed Central PMCID: PMCPMC7773334.

58. Anantharaman V, Aravind L. Evolutionary history, structural features and
biochemical diversity of the NlpC/P60 superfamily of enzymes. Genome Biol.
2003;4(2):R11. Epub 2003/03/07. doi: 10.1186/gb-2003-4-2-r11. PubMed PMID:
12620121; PubMed Central PMCID: PMCPMC151301.

59. Ozhelvaci F, Steczkiewicz K. Identification and classification of papain-like
cysteine proteinases. The Journal of biological chemistry. 2023;299(6):104801. Epub
20230508. doi: 10.1016/j.jbc.2023.104801. PubMed PMID: 37164157; PubMed Central
PMCID: PMCPMC10318531.

60. Griffin ME, Klupt S, Espinosa J, Hang HC. Peptidoglycan NlpC/P60 peptidases
in bacterial physiology and host interactions. Cell Chem Biol. 2023;30(5):436-56. Epub
20221122. doi: 10.1016/j.chembiol.2022.11.001. PubMed PMID: 36417916; PubMed
Central PMCID: PMCPMC10192474.

61. Golczak M, Kiser PD, Sears AE, Lodowski DT, Blaner WS, Palczewski K.
Structural basis for the acyltransferase activity of lecithin:retinol acyltransferase-like
proteins. The Journal of biological chemistry. 2012;287(28):23790-807. Epub
20120517. doi: 10.1074/jbc.M112.361550. PubMed PMID: 22605381; PubMed Central
PMCID: PMCPMC3390653.

62. Xu Q, Rawlings ND, Chiu HJ, Jaroszewski L, Klock HE, Knuth MW, et al.
Structural analysis of papain-like NlpC/P60 superfamily enzymes with a circularly
permuted topology reveals potential lipid binding sites. PloS one. 2011;6(7):e22013.
Epub 20110722. doi: 10.1371/journal.pone.0022013. PubMed PMID: 21799766;
PubMed Central PMCID: PMCPMC3142127.

63. Sears AE, Palczewski K. Lecithin:Retinol Acyltransferase: A Key Enzyme
Involved in the Retinoid (visual) Cycle. Biochemistry. 2016;55(22):3082-91. Epub
20160523. doi: 10.1021/acs.biochem.6b00319. PubMed PMID: 27183166; PubMed
Central PMCID: PMCPMC5555363.

64. Mardian EB, Bradley RM, Duncan RE. The HASLS (PLA/AT) subfamily of
enzymes. J Biomed Sci. 2015;22:99. Epub 20151026. doi: 10.1186/s12929-015-0210-7.
PubMed PMID: 26503625; PubMed Central PMCID: PMCPMC4624172.

65. Armbruster KM, Jiang J, Sartorio MG, Scott NE, Peterson JM, Sexton JZ, et al.
Identification and Characterization of the Lipoprotein N-acyltransferase in Bacteroides.
bioRxiv. 2024. Epub 20240601. doi: 10.1101/2024.05.31.596883. PubMed PMID:
38853980; PubMed Central PMCID: PMCPMC11160734.

66. Zheng L, Lin Y, Lu S, Zhang J, Bogdanov M. Biogenesis, transport and
remodeling of lysophospholipids in Gram-negative bacteria. Biochim Biophys Acta
Mol Cell Biol Lipids. 2017;1862(11):1404-13. Epub 20161209. doi:
10.1016/j.bbalip.2016.11.015. PubMed PMID: 27956138; PubMed Central PMCID:
PMCPMC6162059.

67. Henriksen JR, Andresen TL, Feldborg LN, Duelund L, Ipsen JH. Understanding
detergent effects on lipid membranes: a model study of lysolipids. Biophys J.
2010;98(10):2199-205. doi: 10.1016/j.bpj.2010.01.037. PubMed PMID: 20483328;
PubMed Central PMCID: PMCPMC2872274.

68. Salomon D, Kinch LN, Trudgian DC, Guo X, Klimko JA, Grishin NV, et al.
Marker for type VI secretion system effectors. Proceedings of the National Academy of

Sciences of the United States of America. 2014;111(25):9271-6. Epub 20140609. doi: 10.1073/pnas.1406110111. PubMed PMID: 24927539; PubMed Central PMCID: PMCPMC4078801.

69.   Jana B, Fridman CM, Bosis E, Salomon D. A modular effector with a DNase domain and a marker for T6SS substrates. Nat Commun. 2019;10(1):3595. Epub 20190809. doi: 10.1038/s41467-019-11546-6. PubMed PMID: 31399579; PubMed Central PMCID: PMCPMC6688995.

70.   Fridman CM, Keppel K, Gerlic M, Bosis E, Salomon D. A comparative genomics methodology reveals a widespread family of membrane-disrupting T6SS effectors. Nat Commun. 2020;11(1):1085. Epub 20200227. doi: 10.1038/s41467-020-14951-4. PubMed PMID: 32109231; PubMed Central PMCID: PMCPMC7046647.

71.   Blondel CJ, Amaya FA, Bustamante P, Santiviago CA, Pezoa D. Identification and distribution of new candidate T6SS effectors encoded in Salmonella Pathogenicity Island 6. Front Microbiol. 2023;14:1252344. Epub 2023/09/04. doi: 10.3389/fmicb.2023.1252344. PubMed PMID: 37664116; PubMed Central PMCID: PMCPMC10469887.

72.   Kanarek K, Fridman CM, Bosis E, Salomon D. The RIX domain defines a class of polymorphic T6SS effectors and secreted adaptors. Nat Commun. 2023;14(1):4983. Epub 20230817. doi: 10.1038/s41467-023-40659-2. PubMed PMID: 37591831; PubMed Central PMCID: PMCPMC10435454.

73.   Geller AM, Shalom M, Zlotkin D, Blum N, Levy A. Identification of type VI secretion system effector-immunity pairs using structural bioinformatics. Mol Syst Biol. 2024;20(6):702-18. Epub 20240424. doi: 10.1038/s44320-024-00035-8. PubMed PMID: 38658795; PubMed Central PMCID: PMCPMC11148199.

74.   Carobbi A, Leo K, Di Nepi S, Bosis E, Salomon D, Sessa G. PIX is an N-terminal delivery domain that defines a class of polymorphic T6SS effectors in Enterobacterales. Cell Rep. 2024;43(4):114015. Epub 20240402. doi: 10.1016/j.celrep.2024.114015. PubMed PMID: 38568810.

75.   Jana B, Keppel K, Fridman CM, Bosis E, Salomon D. Multiple T6SSs, Mobile Auxiliary Modules, and Effectors Revealed in a Systematic Analysis of the Vibrio parahaemolyticus Pan-Genome. mSystems. 2022;7(6):e0072322. Epub 20221013. doi: 10.1128/msystems.00723-22. PubMed PMID: 36226968; PubMed Central PMCID: PMCPMC9765294.

76.   Tchelet D, Keppel K, Bosis E, Salomon D. Vibrio parahaemolyticus T6SS2 effector repertoires. Gut Microbes. 2023;15(1):2178795. doi: 10.1080/19490976.2023.2178795. PubMed PMID: 36803660; PubMed Central PMCID: PMCPMC9980498.

77.   Hersch SJ, Watanabe N, Stietz MS, Manera K, Kamal F, Burkinshaw B, et al. Envelope stress responses defend against type six secretion system attacks independently of immunity proteins. Nat Microbiol. 2020;5(5):706-14. Epub 2020/02/26. doi: 10.1038/s41564-020-0672-6. PubMed PMID: 32094588; PubMed Central PMCID: PMCPMC7190449.

78.   Russell AB, Hood RD, Bui NK, LeRoux M, Vollmer W, Mougous JD. Type VI secretion delivers bacteriolytic effectors to target cells. Nature. 2011;475(7356):343-7. Epub 20110720. doi: 10.1038/nature10244. PubMed PMID: 21776080; PubMed Central PMCID: PMCPMC3146020.

79.   Chou S, Bui NK, Russell AB, Lexa KW, Gardiner TE, LeRoux M, et al. Structure of a peptidoglycan amidase effector targeted to Gram-negative bacteria by the type VI secretion system. Cell Rep. 2012;1(6):656-64. Epub 20120531. doi:

10.1016/j.celrep.2012.05.016. PubMed PMID: 22813741; PubMed Central PMCID: PMCPMC3401384.

80.   Cock PJ, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely available Python tools for computational molecular biology and bioinformatics. Bioinformatics. 2009;25(11):1422-3. Epub 20090320. doi: 10.1093/bioinformatics/btp163. PubMed PMID: 19304878; PubMed Central PMCID: PMCPMC2682512.

81.   McKinney Wo. Data structures for statistical computing in python. . Proceedings of the 9th Python in Science Conference. 2010;445:51-6.

82.   Johnson LS, Eddy SR, Portugaly E. Hidden Markov model speed heuristic and iterative HMM search procedure. BMC Bioinformatics. 2010;11:431. Epub 20100818. doi: 10.1186/1471-2105-11-431. PubMed PMID: 20718988; PubMed Central PMCID: PMCPMC2931519.

83.   Hauser M, Steinegger M, Soding J. MMseqs software suite for fast and deep clustering and searching of large protein sequence sets. Bioinformatics. 2016;32(9):1323-30. Epub 20160106. doi: 10.1093/bioinformatics/btw006. PubMed PMID: 26743509.

84.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80. Epub 20130116. doi: 10.1093/molbev/mst010. PubMed PMID: 23329690; PubMed Central PMCID: PMCPMC3603318.

85.   Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for large alignments. PloS one. 2010;5(3):e9490. Epub 20100310. doi: 10.1371/journal.pone.0009490. PubMed PMID: 20224823; PubMed Central PMCID: PMCPMC2835736.

86.   Eddy SR. Accelerated Profile HMM Searches. PLoS computational biology. 2011;7(10):e1002195. Epub 20111020. doi: 10.1371/journal.pcbi.1002195. PubMed PMID: 22039361; PubMed Central PMCID: PMCPMC3197634.

87.   Wang J, Li J, Hou Y, Dai W, Xie R, Marquez-Lago TT, et al. BastionHub: a universal platform for integrating and analyzing substrates secreted by Gram-negative bacteria. Nucleic Acids Res. 2021;49(D1):D651-D9. doi: 10.1093/nar/gkaa899. PubMed PMID: 33084862; PubMed Central PMCID: PMCPMC7778982.

88.   Souza DP, Oka GU, Alvarez-Martinez CE, Bisson-Filho AW, Dunger G, Hobeika L, et al. Bacterial killing via a type IV secretion system. Nat Commun. 2015;6:6453. Epub 20150306. doi: 10.1038/ncomms7453. PubMed PMID: 25743609.

89.   Bayer-Santos E, Cenens W, Matsuyama BY, Oka GU, Di Sessa G, Mininel IDV, et al. The opportunistic pathogen Stenotrophomonas maltophilia utilizes a type IV secretion system for interbacterial killing. PLoS pathogens. 2019;15(9):e1007651. Epub 2019/09/13. doi: 10.1371/journal.ppat.1007651. PubMed PMID: 31513674; PubMed Central PMCID: PMCPMC6759196.

90.   Dykxhoorn DM, St Pierre R, Linn T. A set of compatible tac promoter expression vectors. Gene. 1996;177(1-2):133-6. doi: 10.1016/0378-1119(96)00289-2. PubMed PMID: 8921858.

91.   Datsenko KA, Wanner BL. One-step inactivation of chomosomal genes in Escherichia coli K-12 using PCR products. Proceedings of the National Academy of Sciences of the United States of America. 2000;97(12):6640-5. doi: 10.1073/pnas.120163297. PubMed PMID: 10829079; PubMed Central PMCID: PMCPMC18686.

92.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods.

2012;9(7):676-82.   Epub   20120628.   doi: 10.1038/nmeth.2019.   PubMed   PMID:
22743772; PubMed Central PMCID: PMCPMC3855844.
93.   Beuzon CR, Holden DW. Use of mixed infections with Salmonella strains to
study virulence genes and their interactions in vivo. Microbes and infection / Institut
Pasteur. 2001;3(14-15):1345-52. doi: 10.1016/s1286-4579(01)01496-4. PubMed PMID:
11755424.
94.   Belval L, Marquette A, Mestre P, Piron MC, Demangeat G, Merdinoglu D, et al.
A fast and simple method to eliminate Cpn60 from functional recombinant proteins
produced   by   E.   coli   Arctic   Express.   Protein   Expr   Purif.   2015;109:29-34.   Epub
20150203. doi: 10.1016/j.pep.2015.01.009. PubMed PMID: 25655203.
95.   Yoshida Y, Kodai S, Takemura S, Minamiyama Y, Niki E. Simultaneous
measurement of F2-isoprostane, hydroxyoctadecadienoic acid, hydroxyeicosatetraenoic
acid,   and   hydroxycholesterols   from   physiological   samples.   Anal   Biochem.
2008;379(1):105-15. Epub 20080425. doi: 10.1016/j.ab.2008.04.028. PubMed PMID:
18482573.
96.   Chaves-Filho AB, Pinto IFD, Dantas LS, Xavier AM, Inague A, Faria RL, et al.
Alterations in lipid metabolism of spinal cord linked to amyotrophic lateral sclerosis.
Sci Rep. 2019;9(1):11642. Epub 20190812. doi: 10.1038/s41598-019-48059-7. PubMed
PMID: 31406145; PubMed Central PMCID: PMCPMC6691112.
97.   Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam
H, et al. Clustal W and Clustal X version 2.0. Bioinformatics. 2007;23(21):2947-8.
Epub 20070910. doi: 10.1093/bioinformatics/btm404. PubMed PMID: 17846036.
98.   Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P,
Nielsen H, et al. DeepTMHMM predicts alpha and beta transmembrane proteins using
deep   neural   networks.   bioRxiv.   2022:2022.04.08.487609.   doi:
10.1101/2022.04.08.487609.

1062

Supporting information

S1 Fig. T6SS effector repertoire in the 10KSG dataset. Each column indicates the

presence or absence of a toxin as identified by the models developed in this study. Lines

denote unique combinations of effectors. The histogram on the right shows the

frequency of genomes in the 10KSG dataset containing each specific repertoire. The

histogram at the bottom illustrates the frequency of genomes with at least one protein

identified by the above HMM model. Effector activities are color-coded as described in

Fig 2A.

S2 Fig. List of five most frequent T6SS effectors identified in each of the 149 *Salmonella* **serovars contained in the 10KSG dataset.** Colors indicate the effector activity as in Fig 2A.

S3 Fig. Imm-Act1 is a two-transmembrane helices protein. (a) Sequence alignment of Imm-Act1 homologs with orange rectangles indicating predicted transmembrane helices. Sequences are colored according to the Clustal X color scheme[97]. Consensus abbreviations: h, hydrophobic (A, C, F, I, L, M, V, W, Y); l, aliphatic (L, I, V); a, aromatic (F, W, Y); b, large residues (L, I, Y, E, R, F, Q, K, M, W); s, small residues (A, G, S, V, C, D, N); u, tiny residues (G, A, S); p, polar residues (S, T, E, D, K, R, N, Q, H, C); c, charged residues (D, E, H, K, R); +, positively charged residues (H, K, R); –, negatively charged residues (D, E). (B) Transmembrane helix prediction for Imm-Act1 using DeepTMHMM[98].

S4 Fig. Recombinant protein purification used in enzymatic assay. SDS–PAGE of recombinant proteins during purification steps to obtain purified protein for enzymatic assays. Affinity chomatography using Strep-Tactin sepharose to purify Tox-Act1 versions with C-terminus Strep-tag II: WT in (a) or C151A in (b). Recombinant proteins were purified from the soluble fraction. An additional step of washing with 1.5M urea was performed after the traditional washes to remove contamination with GroEL before elution with biotin. Additional bands were identified by mass-spectrometry to confirm identity. MK: marker; INS: insoluble; SOL: soluble; FT: flowthough. GroEL: chaperonin GroEL; Bccp: biotin carboxyl carrier protein; OmpF: outer membrane porin F.

1097   **S1 Table. List of all T6SS toxin domains identified in this study in the 10K**

1098   ***Salmonella* genomes dataset.**

1099

1100   **S2 Table. List of all homologs collected by JackHMMER searches and used to**

1101   **build the phylogenetic tree shown in Fig 4A.**

1102

1103   **S3 Table. Genomic context of members of each permuted NlpC/P60 members**

1104   **shown in Fig 4A.**

1105

1106   **S4 Table. List of strains, plasmids and primers used in the study.**

1107

1108   **S5 Table. (a) MRM transition for LC-MS/MS method of lysophospholipids. (b)**

1109   **Internal standards used for lipidomics analysis in negative mode.**

1110

1111   **S1 Movie. Time-lapse microscopy of *E. coli* harboring pBRA SP-Tox-Act1 growing**

1112   **in media supplemented with 0.2% d-glucose.** Timestamp in hh:mm. Scale bar: 5 μm.

1113

1114   **S2 Movie. Time-lapse microscopy of *E. coli* harboring pBRA SP-Tox-Act1 growing**

1115   **in media supplemented with 0.2% l-arabinose.** Timestamp in hh:mm. Scale bar: 5

1116   μm.

1117

1118