Technical Report

Simultaneous observation of RNAs and their binding proteins in plant cells

Junfei Ma[1†*], Yunhan Wang[2†], Svetlana Y. Folimonova[1,2], Bin Liu[3], Ying Wang[1,2*]

1 Department of Plant Pathology, University of Florida, Gainesville, FL, USA

2 Plant Molecular and Cell Biology Program, University of Florida, Gainesville, FL, USA

3 Food Science and Human Nutrition Department, University of Florida, FL, USA

[†] These authors contributed equally to the work.
*Correspondence: Junfei Ma and Ying Wang

Email: junfeima@ufl.edu (JM) and ying.wang1@ufl.edu (YW)

Keywords

viroid, RNA fluorescence *in situ* hybridization (FISH), immunofluorescence (IF), RNA-protein co-localization, RNA polymerase II (Pol II)

Abstract

RNA is key component of a versatile group of complexes that play diverse roles in a wide range of biological processes, including plant acclimation to ever-changing environments. From synthesis to degradation, RNA molecules interact with cognate proteins that assist in processes such as transcription, splicing, modification, trafficking, and the execution of their functions. While numerous valuable techniques exist to study RNA-protein interactions, visualizing RNAs and their associated proteins simultaneously within cells remains challenging, despite its potential to provide deeper insights into the biology of RNA-protein interactions. In this study, we adapted a modified immunofluorescence (IF) assay in combination with RNA fluorescence in situ hybridization (FISH) to successfully visualize the colocalization of endogenous as well as infectious RNAs with their cognate binding proteins in plant nucleus. This new method that combines IF and FISH will facilitate future studies on RNA and protein colocalization in various plant systems.

Introduction

Mounting evidence has highlighted the essential roles of diverse classes of RNAs across a wide range of biological processes (Sharp, 2009). From transcription to decay, RNAs interact with specific RNA-binding proteins, and these interactions often correlate with distinct biological functions (Street et al., 2024). In certain cases, RNA-protein complexes organize into membrane-less organelles via liquid-liquid phase separation (Miao, Chodasiewicz, & Fang, 2024). Examples include Processing bodies and Stress Granules involved in mRNA turnover or storage (Chantarachot & Bailey-Serres, 2018), SERRATE-driven dicing bodies in microRNA processing (Xie et al., 2021), FLL2-promoted phase separation of polyadenylation complexes for mRNA 3' end processing (Fang et al., 2019), etc. Although tools such as RNA immunoprecipitation and gel electrophoresis mobility shift assays are valuable for studying RNA-protein interactions, observing RNA subcellular localization or simultaneously visualizing specific RNAs with their associated proteins in cells remains challenging, hindering the understanding of RNA-protein dynamics in the context of cell biology.

The MS2-MCP system enables live-cell imaging of RNAs by using the MS2-binding coat protein (MCP) tagged with a green fluorescent protein (GFP) to recognize RNAs fused with MS2 hairpins (Syed & Lim, 2024). By tracking MSP-fused GFP fluorescence, RNA subcellular localization can be observed in living cells with high specificity and temporal resolution. RNA-binding proteins can be co-visualized by tagging them with fluorescent proteins other than GFP. In addition to the MS2-MCP system, RNA fluorescence aptamers also allow visualization of fusion RNAs in living cells (Sachdeva, Garg, Godding, Way, & Silver, 2014). However, both systems pose a risk of disrupting RNA overall structures and functions. For example, any insertion in the viroid RNA genome can lead to a loss of infectivity, limiting the applications of these methods to certain RNAs.

Observing proteins and RNAs in chemically fixed cells can provide insights into their subcellular dynamics as well. Immunofluorescence (IF) detects specific proteins, while RNA fluorescence in situ hybridization (FISH) reveals RNA subcellular localization patterns. The development of single-molecule RNA FISH (smFISH) enables the

quantitative observation of mRNAs, but it requires dozens of short, specific oligos (17-
22 nt) and may not be suitable for short RNAs or those with limited unique sequences
(Duncan, Johansson, & Ding, 2023). Moreover, traditional IF and FISH are often
incompatible, because the formamide used in FISH tend to mask antigens, hindering
protein detection (B. Ma & Tanese, 2013; Meng, Zhao, & Lao, 2018; Meyer, Garzia, &
Tuschl, 2017). Notably, some reports showed that fluorescent proteins (FPs) could
retain fluorescence activity to outline protein subcellular localization after RNA FISH
steps (Duncan et al., 2023; Huang, Guillotin, Rahni, Birnbaum, & Wagner, 2023; Zhao,
Fonseca, Meschichi, Sicard, & Rosa, 2023). However, this approach is mostly limited to
observing proteins colocalized with cytosolic RNAs because RNA-FISH protocols for
observing nuclear RNAs often denature FPs by high temperatures and stringent
chemical treatments (Alkaabi, Yafea, & Ashraf, 2005; Jolly, Mongelard, Robert-Nicoud,
& Vourc'h, 1997; Qi & Ding, 2003; Yokoyama, Ogawa, Nozu, & Hashimoto, 1990).

To address this shortcoming, we developed a new strategy that combines IF and
FISH to simultaneously visualize an RNA and its interacting protein partner in same
cells. This approach is straightforward, does not require extensive cloning or transgenic
plants (if a specific primary antibody is available), and reliably detects RNAs and
proteins in the nucleus. We successfully tested the method by using a classic RNA-
binding protein fibrillarin and one of its RNA partners (U3 small nucleolar RNA/snoRNA),
which often function in biomolecular condensates for processing nascent rRNAs (Yao et
al., 2019).

We also applied this method to examining viroid-host interactions at the cellular and
subcellular levels. Viroids are single-stranded, circular, noncoding RNA pathogens that
primarily infect crops (J. Ma, Dissanayaka Mudiyanselage, Hao, & Wang, 2023; Ortola
& Daros, 2023). There are two viroid families, *Pospiviroidae* and *Avsunviroidae*.
Members of *Pospiviroidae* replicate in the nucleus using host RNA polymerase II (Pol
II)(Wang, 2021). Unlike the well-known 12-subunit Pol II, a modified Pol II with 6-7
subunits transcribes the potato spindle tuber viroid (PSTVd) RNA (Dissanayaka
Mudiyanselage et al., 2022), assisted by an RNA-specific transcription factor (TFIIIA-
7ZF)(Dissanayaka Mudiyanselage et al., 2022; Wang et al., 2016). Despite the progress
achieved in the understanding of the PSTVd replication, the detailed subcellular

localization and the pattern of Pol II transcription on PSTVd RNA remain unclear. Using
our new method, we observed the nuclear distribution patterns of Pol II (represented by
RPB1) and PSTVd sense-strand RNA in protoplasts. Therefore, this method for
simultaneous observation of RNAs and proteins in plant cells is valuable for studying
various nuclear-localized noncoding RNAs and their associated proteins, including
those involved in liquid-liquid phase separation across diverse biological processes.
100
Results
Optimizing RNA FISH protocol for detecting RNA in the nucleus
First, we sought to use PSTVd RNA as an example to optimize the procedures for
detecting RNA in fixed plant cells. Notably, a detailed protocol for this type of
experiments is not currently available. Due to PSTVd's relatively short length (359 nt),
designing dozens of oligo probes, as required in some single-molecule RNA FISH
protocols, is not possible. Instead, we used traditional *in vitro* transcription method to
generate RNA oligos.
The key factor is optimizing probe size. We designed two probes around 100 nt in
length that are short enough to facilitate nuclear entry. Each probe contained a
maximum of 25-30 AlexaFluor488-labeled UTPs. We used both probes, achieving up to
54 labeled UTPs per PSTVd RNA to enhance the signal-to-background ratio. We tested
two general protocols reported in previous RNA FISH experiments that observed
nuclear PSTVd accumulation (Qi & Ding, 2002; Yokoyama et al., 1990). Interestingly,
combining the penetration steps from both protocols yielded better, more consistent
results (Figure 1). PSTVd displayed a unique distribution pattern, with a dominant signal
near the nucleolar region, as indicated by the hollow DAPI staining (Figure 1). This
PSTVd RNA condensate pattern is consistent with previous reports (Qi & Ding, 2002,
2003). Notably, this protocol also worked well for the endogenous U3 snoRNA as we
performed colocalization analysis of U3 snoRNA and fibrillarin (Figure 2A).
121
Modifying the traditional immunofluorescence IF protocol allowing simultaneous
observation of fibrillarin and U3 small nucleolar RNA
Only a limited number of immunofluorescence protocols are available for plant

samples. Based on a previous protocol (Borg, Buendia, & Berger, 2019), we performed

extensive trials by using fibrillarin as a target and established one that works well for this

endogenous protein. Fibrillarin is an RNA-binding protein that associates with many

endogenous as well as infectious RNAs, such as U3, U8, and U13 small nucleolar

RNAs (snoRNAs) (Tyc & Steitz, 1989), ribosomal RNAs (rRNAs) (Rakitina, Taliansky,

Brown, & Kalinina, 2011), ELF18-INDUCED LONG NONCODING RNA 1 (Seo,

Diloknawarit, Park, & Chua, 2019), tobacco mosaic virus (Rakitina et al., 2011), bamboo

mosaic virus satellite RNA (Chang et al., 2016), etc. Fibrillarin is a methyltransferase

enzyme that is critical for rRNAs' processing (Rodriguez-Corona, Sobol, Rodriguez-

Zapata, Hozak, & Castano, 2015), and U3 snoRNA-containing ribonucleoprotein guides

fibrillarin to perform the first step of pre-ribosomal RNA processing (Kass, Tyc, Steitz, &

Sollner-Webb, 1990). To allow simultaneous observation of U3 snoRNA and fibrillarin in

the nucleolus, we slightly modified the IF protocol to crosslink the samples twice, pre-IF

crosslinking and post-IF crosslinking. The purpose is to preserve the antibody-antigen

interaction in harsh RNA-FISH procedures. As shown in Figure 2A, we successfully

observed a distinct signal of fibrillarin in the nucleolus.

Based on our RNA FISH protocol, we designed a full-length U3 snoRNA-specific

probe, which allow maximum of 38 AlexaFluor488-labeled UTPs. As shown in Figure 2A,

U3 snoRNA specific signals were observed in the nucleolus and often co-localized with

fibrillarin when both specific antibodies and probes were applied. Separate color

channels of another example also confirmed the colocalization around the nucleolar

region (Figure 2B). This colocalization pattern resembles previous observations in HeLa

cells (Leary, Terns, & Huang, 2004). We analyzed the RNA-protein colocalization by

using the JACoP plugin of Image J (Bolte & Cordelieres, 2006) and found the Pearson's

co-efficiency values range from 0.735 to 0.949 (Figure S1), an indication of co-

localization.

An alternative immunofluorescence (IF) approach allowing simultaneous

observation of PSTVd and Pol II

With the success of observing the co-localization of U3 snoRNA and fibrillarin, we

aimed to observe the subcellular localization of Pol II and PSTVd, which would deepen

our understanding of plant-viroid interactions. However, our initial attempts using the
above-mentioned IF protocol were unsuccessful mainly due to the failure of antibody
penetration into the fixed nuclei. Inspired by a recent study on RNase L-induced bodies
that directly incubate antibodies with culture cells (Watkins & Burke, 2024), we
reasoned that incubating antibodies directly with protoplasts could be an effective
alternative IF approach. We tested this approach by observing GFP protein localization
in the *Nicotiana benthamiana* 16C line, which expresses a GFP5 variant with an
endoplasmic reticulum localization signal (Philips et al., 2017). Figure 3 shows the GFP
fluorescence signal in the live protoplasts as well as in the cells incubated with the
antibodies. Signals from the Alexa Fluor594-conjugated anti-GFP antibody largely
highlighted GFP localization comparable to the direct fluorescence from GFP,
demonstrating that directly incubating antibodies with protoplasts can serve as an
alternative IF technique.

Using this system, we aimed to understand the subcellular distribution of Pol II and
PSTVd RNA. We used the well-known 8WG16 monoclonal antibody to detect the
localization of the largest subunit (RPB1) of Pol II. As shown in Figure 4A, there is no
fluorescence signal from samples in which we did not add the primary antibody,
demonstrating the specificity in detecting target protein. When both antibodies and
probes were applied, we observed significant co-localization of PSTVd and Pol II in the
nucleoplasm (Figure 4A).

In another representative image (Figure 4B), we showed that the RPB1 and PSTVd
colocalized loci are spread in the nucleoplasm. This pattern demonstrates that Pol II
transcribes the template of PSTVd sense-strand RNA in discrete loci mainly in the
nucleoplasm. The JACoP analysis showed that Pearson's co-efficiency values range
from 0.867 to 0.997 (Figure S1), indicating a positive co-localization.

The overall strategy combining IF and FISH to simultaneously observe RNA and
cognate binding protein

The success in observing the colocalization of U3 snoRNA/fibrillarin and
PSTVd/RPB2 indicates that combined IF-FISH protocol is efficient for observing
proteins and RNA simultaneously in the cells (Figure 5). This strategy involves

generating protoplasts from leaves, followed by IF process. The key is to fix the cells

with antibody before the subsequent RNA FISH steps, which include permeabilization,

blocking, hybridization, and finally, observation. The entire procedure can be completed

in four days.

Discussion

By combining IF and RNA FISH, we successfully achieved the simultaneous

visualization of RNAs and their cognate proteins in the nucleus of plant cells. This

approach eliminates the need of extensive cloning, as required in the MS2–MCP

system (Syed & Lim, 2024), or the generation of transgenic plants when a specific

primary antibody is available. Notably, our protocol is effective for detecting both

endogenous and foreign RNAs in the nucleus, which has previously proven challenging

based on both literature and our experience. Additionally, this method allows the direct

visualization of proteins and RNAs in their native state without ectopic expression,

making it a robust and straightforward approach for plant cell biology studies.

Compared with smFISH (Duncan et al., 2023), our probe design strategy offers an

advantage in detecting relatively short RNAs that are insufficient in length for designing

multiple oligos. Nevertheless, our method could be adapted in combination with smFISH

to detect endogenous mRNAs. Furthermore, multiple proteins and/or RNAs can be

observed within the same cells by utilizing various AlexaFluor dyes, allowing different

targets to be marked with distinct colors. This flexibility enables researchers to adapt the

method to address a range of biological questions.

Notably, we present two distinct IF protocols for different proteins. The conventional

IF protocol is suitable for detecting fibrillarin in our practice. In contrast, the live-cell-

antibody-incubation strategy is more suitable for DNA ligase 1 (Wang, Ma, Hao, Liu, &

Wang, 2024) and RPB1. It is expected that the two available protocols will increase the

chance for successful detection of a wide-spectrum of proteins.

However, we'd like to note that both antibodies and probes penetrate into

protoplasts at a relatively low frequency (~20-30% of cells). Therefore, there are limited

cells with both signals. The reasons behind such low frequency are not immediately

clear. Nevertheless, it is still possible to find dozens of samples with colocalization

signals. Future trials may be needed to further improve the efficiency.

Recent studies have increasingly highlighted the importance of dynamic RNA

organization and their associated factors within the cellular environment, particularly in

the context of phase separation across diverse biological processes (Miao et al., 2024).

However, most knowledge has been derived either from methods lacking cellular

context (e.g., RIP-Seq, iCLIP, etc) (Lewinski et al., 2024) or indirectly by observing

either protein (Citovsky et al., 2006; Cox et al., 2024; Paciorek, Sauer, Balla,

Wisniewska, & Friml, 2006; Sharma, Klosgen, & Bennewitz, 2023) or RNA subcellular

localization alone (Duncan, Olsson, Hartley, Dean, & Rosa, 2017; J. Ma et al., 2022;

Zollner et al., 2021). Our method provides a means to directly investigate the dynamic

interactions between RNAs and proteins in cells, offering a valuable tool for future

research. One recent success is demonstrating the co-localization of DNA ligase1 and

PSTVd in a biomolecular condensate (Wang et al., 2024).

Although it is well-known that PSTVd replication depends on Pol II and TFIIIA-7ZF

activity (Wang et al., 2016), the exact nuclear distribution pattern involved in PSTVd

RNA-templated transcription remain unclear. By using our method, we now understand

that Pol II transcribes sense-strand of PSTVd in multiple discrete loci in the nucleoplasm.

Pol II transcription loci do not all overlap with the major PSTVd biomolecular

condensate, and this biomolecular condensate contains PSTVd and DNA ligase1

(Wang et al., 2024), the latter of which is a critical enzyme for generating progeny

(Nohales, Flores, & Daros, 2012). Therefore, the observation support the model that

there is an active enrichment process of sense-strand PSTVd RNA for enzymatic

processing (Qi & Ding, 2003).

Materials and methods

Plant growth and protoplast preparation

Nicotiana benthamiana (wild type and line 16C) plants were grown in a growth shelf

at 25 °C with a 16-hour light and 8-hour dark cycle. PSTVd-infected plants were verified

by RNA gel blots as described previously (Takeda, Zirbel, Leontis, Wang, & Ding, 2018).

Protoplasts were isolated from the 4[th] to 6[th] leaves of the plants by following a published

protocol with slight modifications (Jiang, Ma, Liu, & Wang, 2019). Briefly, leaves were

digested with 3% cellulose (Onozuka Yakult Pharmaceutical IND., Tokyo, Japan) and

0.8% macerase (MilliporeSigma, Burlington, MA, USA) in protoplast digestion buffer

(0.4 M mannitol, 20 mM KCl, 20 mM MES, 10 mM $CaCl_2$, 0.1% BSA, 5 mM β-

mercaptoethanol, pH 5.7). The lower epidermis layer of the leaves was removed by

tapes before digestion. After 1 hr of digestion, the protoplasts were pelleted by

centrifugation at 150× g for 1 min. The pelleted protoplasts were then washed

sequentially with W5 buffer (5 mM MES, 154 mM NaCl, 125 mM $CaCl_2$, 5 mM KCl, pH

5.7) at room temperature. Finally, the protoplasts were incubated in the IF buffer (0.5 M

mannitol, 20 mM MES, pH 5.7, 20 mM KCl, 1% BSA, 0.1% Tween-20).

Traditional immunofluorescence (IF)

After extensive trials, we found the following procedures can lead to most clear and

repeatable results. Protoplasts were first fixed with 3.7% formaldehyde and 0.2% picric

acid in KE buffer (50 mM KH_2PO_4, 5 mM EGTA, pH 5.0) at room temperature for 12 min.

After fixation, cells were quenched in 1X KE buffer with 300mM glycine for 30 min and

washed in 1X KE buffer twice. The fixed cells were then permeabilized in 1X PBS buffer

containing 2% BSA, 0.1% cellulase, 0.05% macerozyme, 1% Triton X-100 and 0.5%

Nonidet P-40 at room temperature for 30 min, followed by methyl alcohol incubation at

minus 20°C for 30 min. Cell blocking was performed using 1X PBS containing 0.2%

Tween20, 1% BSA, and 10% DMSO for 1 hr in the room temperature. Cells were

incubated with 1:250 diluted fibrillarin primary antibody (Santa Cruz Biotechnology,

Dallas, Texas) at 4°C overnight. After three washes, cells were then incubated with

1:250 diluted AlexaFluor594-conjugated anti-mouse secondary antibody

(ThermoFisherSci) at 37°C for 1 hr. After three washes, cells were fixed for the second

time with 3.7% formaldehyde and 0.2% picric acid in KE buffer at room temperature for

30 min.

Live-cell immunofluorescence

The live cell IF method was adapted from a study on *in vivo* antibody uptake

(Watkins & Burke, 2024). Freshly prepared protoplasts were incubated overnight at 4°C

with primary antibodies diluted 1:500 in IF buffer. After primary antibody incubation,

protoplasts were washed three times with IF buffer with a 10 min interval. Secondary
antibodies were added in IF buffer with a 1:50 dilution and incubated at room
temperature for 3 hr. The protoplasts were washed three times with IF buffer (10 min
each) to remove the secondary antibody. Sample fixation was done by using 3.7%
formaldehyde and 0.2% picric acid in KE at room temperature for 1 hr. Cross-linking
was stopped by using 1X KE buffer containing 0.3M Glycine, followed by two times
washing with 1X KE buffer for 1h each. Primary antibodies are mouse monoclonal
8WG16 (ThermoFisherSci, Waltham, MA) for Pol II RPB1 and anti-GFP polyclonal
antibody raised in rabbit (Genscript, Piscataway, NJ). AlexaFluor594-conjugated anti-
mouse or anti-rabbit antibodies (both from ThermoFisherSci) was used as secondary
antibody.

RNA fluorescence in situ hybridization (FISH)

Following IF treatment and fixation, the protoplasts were stored in 70% ethanol at
4°C overnight before FISH steps. The samples were baked onto Superfrost®
microscope slides (ThermoFisherSci) at -20°C for 30 minutes, followed by baking at
70°C for 2 hours. Next, the samples were washed with 1X PBS containing 2% bovine
serum albumin (BSA), 0.1% cellulase, 0.05% macerozyme, 0.3% Triton X-100, and 0.5%
Nonidet P-40 at room temperature for 30 min. Then, samples were washed three times
with 100 mM Tris (pH 6.8) containing 1% Nonidet P-40 and 0.5% Triton X-100 at 37°C
for 10 minutes each. Pre-chilled methanol with 2% hydrogen peroxide was added to the
samples, which were incubated at -20°C for 30 min. The slides were subjected to three
rounds of liquid nitrogen treatment, each lasting 5 seconds. The samples were then
treated with 0.5% (v/v) acetic anhydride in 0.1 M triethanolamine (TEA) buffer (pH 8.0)
at room temperature for 20 min.

Samples were then blocked with PerfectHyb™ Plus hybridization buffer
(ThermoFisherSci) at 55°C for 30 min. Probes (corresponding to the 98-188 and 270-13
nucleotide positions in the PSTVd genome but in the complementary orientation) were
synthesized using T3 Maxiscript kit (ThermoFisherSci) mixed with AlexaFluor488-
labeled UTP (ThermoFisherSci) (J. Ma et al., 2022). Probes (2.5-5 μg/ml) were
denatured by heating at 100°C for 1 minutes, followed by cooling on ice. The denatured

probes, along with 5 nM ribonucleoside vanadyl complex (Sigma-Aldrich, St. Louis, MO, USA), were added to PerfectHyb™ Plus Hybridization Buffer and applied to samples on slides, followed by incubation at 55°C overnight.

The samples were then rinsed with 2X SSC, washed with 2X SSC containing 50% formamide at 37°C for 30 min, and subsequently washed with 2X SSC and 1X SSC at room temperature for 30 minutes each. 4',6-diamidino-2-phenylindole (DAPI) was added directly to the samples stain (1 µg/ml) for 5 minutes and then rinsed with distilled water. Anti-fade reagent (ThermoFisherSci) was applied to each sample, and the slides were sealed with nail polish.

Microscopy imaging

Fluorescence signals were obtained by using a Discover Echo Revolve microscope with DAPI, FITC, and TRITC cubes and Leica Stellaris 5 confocal microscope.

Acknowledgments

This work is supported by US National Science Foundation (MCB-1906060, MCB-2350392, and IOS-2410009 to YW). Yunhan Wang is supported by CALS Deans' scholarship from the University of Florida. Symbols in Figure 5 are from bioicons.com. We apologize to colleagues whose work was not cited due to the page limits.

References

Alkaabi, K. M., Yafea, A., & Ashraf, S. S. (2005). Effect of pH on thermal- and chemical-induced denaturation of GFP. *Appl Biochem Biotechnol, 126*(2), 149-156. doi:10.1385/abab:126:2:149

Bolte, S., & Cordelieres, F. P. (2006). A guided tour into subcellular colocalization analysis in light microscopy. *J Microsc, 224*(Pt 3), 213-232. doi:10.1111/j.1365-2818.2006.01706.x

Borg, M., Buendia, D., & Berger, F. (2019). A simple and robust protocol for immunostaining Arabidopsis pollen nuclei. *Plant Reprod, 32*(1), 39-43. doi:10.1007/s00497-018-00360-7

Chang, C. H., Hsu, F. C., Lee, S. C., Lo, Y. S., Wang, J. D., Shaw, J., . . . Lin, N. S. (2016). The Nucleolar Fibrillarin Protein Is Required for Helper Virus-Independent Long-Distance Trafficking of a Subviral Satellite RNA in Plants. *Plant Cell, 28*(10), 2586-2602. doi:10.1105/tpc.16.00071

Chantarachot, T., & Bailey-Serres, J. (2018). Polysomes, Stress Granules, and Processing Bodies: A Dynamic Triumvirate Controlling Cytoplasmic mRNA Fate and Function. *Plant Physiol, 176*(1), 254-269. doi:10.1104/pp.17.01468

Citovsky, V., Lee, L. Y., Vyas, S., Glick, E., Chen, M. H., Vainstein, A., . . . Tzfira, T. (2006).
Subcellular localization of interacting proteins by bimolecular fluorescence
complementation in planta. *J Mol Biol, 362*(5), 1120-1131.
doi:10.1016/j.jmb.2006.08.017
Cox, K. L., Pardi, S. A., O'Connor, L., Klebanovych, A., Huss, D., Nusinow, D. A., . . . Czymmek, K. J.
(2024). ExPOSE: A comprehensive toolkit to perform expansion microscopy in plant
protoplast systems. *bioRxiv*, 2024.2007.2012.603300. doi:10.1101/2024.07.12.603300
Dissanayaka Mudiyanselage, S. D., Ma, J., Pechan, T., Pechanova, O., Liu, B., & Wang, Y. (2022).
A remodeled RNA polymerase II complex catalyzing viroid RNA-templated transcription.
PLoS Pathog., 18(9), e1010850. doi:10.1371/journal.ppat.1010850
Duncan, S., Johansson, H. E., & Ding, Y. (2023). Reference genes for quantitative Arabidopsis
single molecule RNA fluorescence in situ hybridization. *J Exp Bot, 74*(7), 2405-2415.
doi:10.1093/jxb/erac521
Duncan, S., Olsson, T. S. G., Hartley, M., Dean, C., & Rosa, S. (2017). Single Molecule RNA FISH in
Arabidopsis Root Cells. *Bio Protoc, 7*(8), e2240. doi:10.21769/BioProtoc.2240
Fang, X., Wang, L., Ishikawa, R., Li, Y., Fiedler, M., Liu, F., . . . Dean, C. (2019). Arabidopsis FLL2
promotes liquid-liquid phase separation of polyadenylation complexes. *Nature,*
569(7755), 265-269. doi:10.1038/s41586-019-1165-8
Huang, T., Guillotin, B., Rahni, R., Birnbaum, K. D., & Wagner, D. (2023). A rapid and sensitive,
multiplex, whole mount RNA fluorescence in situ hybridization and
immunohistochemistry protocol. *Plant Methods, 19*(1), 131. doi:10.1186/s13007-023-
01108-9
Jiang, J., Ma, J., Liu, B., & Wang, Y. (2019). Combining a simple method for DNA/RNA/protein
co- purification and Arabidopsis protoplast assay to facilitate viroid research. *Viruses,*
11(4), 324. doi:10.3390/v11040324
Jolly, C., Mongelard, F., Robert-Nicoud, M., & Vourc'h, C. (1997). Optimization of nuclear
transcript detection by FISH and combination with fluorescence immunocytochemical
detection of transcription factors. *J Histochem Cytochem, 45*(12), 1585-1592.
doi:10.1177/002215549704501201
Kass, S., Tyc, K., Steitz, J. A., & Sollner-Webb, B. (1990). The U3 small nucleolar
ribonucleoprotein functions in the first step of preribosomal RNA processing. *Cell, 60*(6),
897-908. doi:10.1016/0092-8674(90)90338-f
Leary, D. J., Terns, M. P., & Huang, S. (2004). Components of U3 snoRNA-containing complexes
shuttle between nuclei and the cytoplasm and differentially localize in nucleoli:
implications for assembly and function. *Mol Biol Cell, 15*(1), 281-293.
doi:10.1091/mbc.e03-06-0363
Lewinski, M., Bruggemann, M., Koster, T., Reichel, M., Bergelt, T., Meyer, K., . . . Staiger, D.
(2024). Mapping protein-RNA binding in plants with individual-nucleotide-resolution
cross-linking and immunoprecipitation (plant iCLIP2). *Nat Protoc, 19*(4), 1183-1234.
doi:10.1038/s41596-023-00935-3
Ma, B., & Tanese, N. (2013). Combined FISH and immunofluorescent staining methods to co-
localize proteins and mRNA in neurons and brain tissue. *Methods Mol Biol, 1010*, 123-
138. doi:10.1007/978-1-62703-411-1_9

Ma, J., Dissanayaka Mudiyanselage, S. D., Hao, J., & Wang, Y. (2023). Cellular roadmaps of viroid
infection. *Trends Microbiol, 31*(11), 1179-1191. doi:10.1016/j.tim.2023.05.014
Ma, J., Dissanayaka Mudiyanselage, S. D., Park, W. J., Wang, M., Takeda, R., Liu, B., & Wang, Y.
(2022). A nuclear import pathway exploited by pathogenic noncoding RNAs. *Plant Cell,
34*(10), 3543-3556. doi:10.1093/plcell/koac210
Meng, C., Zhao, X., & Lao, J. (2018). A modified immunofluorescence in situ hybridization
method to detect long non-coding RNAs and proteins in frozen spinal cord sections. *Exp
Ther Med, 15*(6), 4623-4628. doi:10.3892/etm.2018.6046
Meyer, C., Garzia, A., & Tuschl, T. (2017). Simultaneous detection of the subcellular localization
of RNAs and proteins in cultured cells by combined multicolor RNA-FISH and IF. *Methods,
118-119*, 101-110. doi:10.1016/j.ymeth.2016.09.010
Miao, Y., Chodasiewicz, M., & Fang, X. (2024). Navigating biomolecular condensates in plants
from patterns to functions. *Mol Plant, 17*(9), 1329-1332.
doi:10.1016/j.molp.2024.08.001
Nohales, M. A., Flores, R., & Daros, J. A. (2012). Viroid RNA redirects host DNA ligase 1 to act as
an RNA ligase. *Proc. Natl. Acad. Sci. U. S. A., 109*(34), 13805-13810.
doi:10.1073/pnas.1206187109
Ortola, B., & Daros, J. A. (2023). Viroids: Non-Coding Circular RNAs Able to Autonomously
Replicate and Infect Higher Plants. *Biology (Basel), 12*(2). doi:10.3390/biology12020172
Paciorek, T., Sauer, M., Balla, J., Wisniewska, J., & Friml, J. (2006). Immunocytochemical
technique for protein localization in sections of plant tissues. *Nat Protoc, 1*(1), 104-107.
doi:10.1038/nprot.2006.16
Philips, J. G., Naim, F., Lorenc, M. T., Dudley, K. J., Hellens, R. P., & Waterhouse, P. M. (2017).
The widely used Nicotiana benthamiana 16c line has an unusual T-DNA integration
pattern including a transposon sequence. *PLoS One, 12*(2), e0171311.
doi:10.1371/journal.pone.0171311
Qi, Y., & Ding, B. (2002). Replication of Potato spindle tuber viroid in cultured cells of tobacco
and Nicotiana benthamiana: the role of specific nucleotides in determining replication
levels for host adaptation. *Virology, 302*(2), 445-456.
Qi, Y., & Ding, B. (2003). Differential subnuclear localization of RNA strands of opposite polarity
derived from an autonomously replicating viroid. *Plant Cell, 15*(11), 2566-2577.
doi:10.1105/tpc.016576
Rakitina, D. V., Taliansky, M., Brown, J. W., & Kalinina, N. O. (2011). Two RNA-binding sites in
plant fibrillarin provide interactions with various RNA substrates. *Nucleic Acids Res,
39*(20), 8869-8880. doi:10.1093/nar/gkr594
Rodriguez-Corona, U., Sobol, M., Rodriguez-Zapata, L. C., Hozak, P., & Castano, E. (2015).
Fibrillarin from Archaea to human. *Biol Cell, 107*(6), 159-174.
doi:10.1111/boc.201400077
Sachdeva, G., Garg, A., Godding, D., Way, J. C., & Silver, P. A. (2014). In vivo co-localization of
enzymes on RNA scaffolds increases metabolic production in a geometrically dependent
manner. *Nucleic Acids Res, 42*(14), 9493-9503. doi:10.1093/nar/gku617
Seo, J. S., Diloknawarit, P., Park, B. S., & Chua, N. H. (2019). ELF18-INDUCED LONG NONCODING
RNA 1 evicts fibrillarin from mediator subunit to enhance PATHOGENESIS-RELATED
GENE 1 (PR1) expression. *New Phytol, 221*(4), 2067-2079. doi:10.1111/nph.15530

Sharma, M., Klosgen, R. B., & Bennewitz, B. (2023). Fluorescent Protein-Based Approaches for
Subcellular Protein Localization in Plants. *Methods Mol Biol, 2564*, 203-211.
doi:10.1007/978-1-0716-2667-2_9

Sharp, P. A. (2009). The centrality of RNA. *Cell, 136*(4), 577-580. doi:10.1016/j.cell.2009.02.007

Street, L. A., Rothamel, K. L., Brannan, K. W., Jin, W., Bokor, B. J., Dong, K., . . . Yeo, G. W. (2024).
Large-scale map of RNA-binding protein interactomes across the mRNA life cycle. *Mol*
Cell. doi:10.1016/j.molcel.2024.08.030

Syed, S., & Lim, B. (2024). Multi-labeling Live Imaging to Quantify Gene Expression Dynamics
During Drosophila Embryonic Development. *Methods Mol Biol, 2805*, 137-151.
doi:10.1007/978-1-0716-3854-5_9

Takeda, R., Zirbel, C. L., Leontis, N. B., Wang, Y., & Ding, B. (2018). Allelic RNA Motifs in
Regulating Systemic Trafficking of Potato Spindle Tuber Viroid. *Viruses, 10*(4).
doi:10.3390/v10040160

Tyc, K., & Steitz, J. A. (1989). U3, U8 and U13 comprise a new class of mammalian snRNPs
localized in the cell nucleolus. *EMBO J, 8*(10), 3113-3119. doi:10.1002/j.1460-
2075.1989.tb08463.x

Wang, Y. (2021). Current view and perspectives in viroid replication. *Curr. Opin. Virol., 47*, 32-37.
doi:10.1016/j.coviro.2020.12.004

Wang, Y., Ma, J., Hao, J., Liu, B., & Wang, Y. (2024). DNA Ligase I Circularises Potato Spindle
Tuber Viroid RNA in a Biomolecular Condensate. *Mol Plant Pathol, 25*(12), e70047.
doi:10.1111/mpp.70047

Wang, Y., Qu, J., Ji, S., Wallace, A. J., Wu, J., Li, Y., . . . Ding, B. (2016). A land plant-specific
transcription factor directly enhances transcription of a pathogenic noncoding RNA
template by DNA-dependent RNA polymerase II. *Plant Cell, 28*(5), 1094-1107.
doi:10.1105/tpc.16.00100

Watkins, J. M., & Burke, J. M. (2024). RNase L-induced bodies sequester subgenomic flavivirus
RNAs to promote viral RNA decay. *Cell Rep, 43*(9), 114694.
doi:10.1016/j.celrep.2024.114694

Xie, D., Chen, M., Niu, J., Wang, L., Li, Y., Fang, X., . . . Qi, Y. (2021). Phase separation of SERRATE
drives dicing body assembly and promotes miRNA processing in Arabidopsis. *Nat Cell*
Biol, 23(1), 32-39. doi:10.1038/s41556-020-00606-5

Yao, R. W., Xu, G., Wang, Y., Shan, L., Luan, P. F., Wang, Y., . . . Chen, L. L. (2019). Nascent Pre-
rRNA Sorting via Phase Separation Drives the Assembly of Dense Fibrillar Components in
the Human Nucleolus. *Mol Cell, 76*(5), 767-783 e711. doi:10.1016/j.molcel.2019.08.014

Yokoyama, M., Ogawa, M., Nozu, Y., & Hashimoto, J. (1990). Detection of Specific RNAs by in
situ Hybridization in Plant Protoplasts. *Plant and Cell Physiology, 31*(3), 403-406.
doi:10.1093/oxfordjournals.pcp.a077923

Zhao, L., Fonseca, A., Meschichi, A., Sicard, A., & Rosa, S. (2023). Whole-mount smFISH allows
combining RNA and protein quantification at cellular and subcellular resolution. *Nat*
Plants, 9(7), 1094-1102. doi:10.1038/s41477-023-01442-9

Zollner, N. R., Bezrutczyk, M., Laureyns, R., Nelissen, H., Simon, R., & Frommer, W. B. (2021). An
RNA in situ hybridization protocol optimized for monocot tissue. *STAR Protoc, 2*(2),
100398. doi:10.1016/j.xpro.2021.100398

Figure Legends

Figure Legends

Figure 1. An optimized RNA FISH protocol for observing RNA in plant nucleus.

Nucleolar-enriched potato spindle tuber viroid (PSTVd) detected by Alexa Fluor488-

labeled riboprobe. Scale bar, 8 µm.

Figure 2. Colocalization of fibrillarin and U3 snoRNA. A) fibrillarin (Fri) is detected by ▢-

Fri1 monoclonal antibody and Alexa Fluor594-labeled anti-mouse secondary antibody

(magenta color). Secondary antibody was applied to all treatments. U3 snoRNA is

detected by Alexa Fluor488-labeled specific riboprobe (green). White arrow heads point

to some colocalized loci. B) Separate color channels of a representative sample

showing the colocalization of U3 snoRNA and fibrillarin. Scale bar, 5 µm.

Figure 3. An alternative immunofluorescence protocol. Protoplasts from *Nicotiana*

benthamiana line 16C were used for testing. GFP5-ER in protoplasts was detected by

▢-GFP polyclonal primary antibody and Alexa Fluor594-labeled anti-Rabbit secondary

antibody. anti-Rabbit secondary antibody was added to all IF samples. Scale bar, 5 µm.

Figure 4. Combining immunofluorescence (IF) and RNA fluorescence in situ

hybridization (FISH) to colocalize PSTVd and RPB1. A) RPB1 was detected by ▢-RPB1

polyclonal antibody and Alexa Fluor594-labeled anti-Rabbit secondary antibody

(magenta color). Secondary antibody was applied to all treatments. PSTVd was

detected by Alexa Fluor488-labeled specific riboprobe (green). B) Separate color

channels of a representative sample showing the colocalization of PSTVd and Pol II

(RPB1). White arrow heads point to some colocalized loci. Scale bar, 5 µm.

Figure 5. Schematic workflow of combined IF and FISH.

A

	No α-RPB1	α-RPB1 added
No Probe		
PSTVd Probe		

B

DAPI   Merge

PSTVd   RPB1

Day 1 Sample preparation and IF

Protoplast isolation

option 1

fixation

option 2

1' antibody incubation → 2' antibody incubation → fixation

Day2 FISH

Bake slides → penetration → pre-hybridization → hybridization

Day 3 Wash and View

Wash → slides mounting → observation