Zebrafish Pol Theta and Human Pol Theta: Orthologues with Homologous Function.

Corey Thomas[1], Sydney Green[1], Lily Kimball[2], Isaiah Schmidtke[2], Makayla Griffin[2], Lauren Rothwell[2], Ivy Par[1], Sophia Schobel[1], Yayleene Palacio[1], Jamie B Towle-Weicksel[1], Steven E Weicksel[2]

[1]Department of Physical Sciences, Rhode Island College, Providence, RI

[2]Department of Biology and Biological Sciences, Bryant University, Smithfield RI

ABSTRACT

DNA polymerase theta (Pol θ) is a conserved an A-family polymerase that plays an essential role in repairing double strand breaks, through micro-homology end joining, and bypassing DNA lesions, through translesion synthesis, to protect genome integrity. Despite its essential role in DNA repair, Pol θ is inherently error prone. Recently, Pol θ has been identified in various cancer types, suggesting Pol θ's promiscuous nature aids in cancer progression.

Here we present a study comparing the structure and function of the polymerase domain of zebrafish and human Pol θ to determine the potential of zebrafish as a model for pol θ function. We show that these two proteins share a large amount of sequence and structural homology, but differ in key loop areas thought to drive defining functions of the enzyme. Despite these differences zebrafish Pol θ still displays characteristics identify in human Pol θ, including DNA template extension, microhomology-mediated

end joining, and translesion synthesis. Additionally, we found several important residues

within loops of the polymerase domain that support function unique to Pol θ. These

results will support future studies with zebrafish as a model to investigate Pol θ function.

INTRODUCTION

Melanoma is the rarest but deadliest form of skin cancer, with an estimate in

2020 of 325,000 new cases worldwide, increase to nearly 510,000 new cases by 2040[1].

Unlike the expansion of melanocytes horizontally along the epidermis as benign nevi,

(commonly referred to a moles) melanoma is characterized by expansion vertically,

down, to and through the basement membrane where cells then begin to spread out

from the original tumor location. Melanoma cells are highly mobile in part due to the

migratory nature of melanocytes[2], making migration at key tropism of the disease.

Surgical treatments prior to migration of melanoma to the lymph nodes have a high

degree of success, with low rates of tumor reoccurrence and favorable patient

outcomes. Patient outcomes, however, become less favorable after migration, as

metastatic tumors often are no longer sensitive to radiation or chemotherapy-based

approaches. Recent advances in new immune-boosting therapies have shown promise

in treating metastatic melanoma, but early detection and tumor resection treatments are

still the most effective[3,4]. This indicates that there is still much to know about melanoma

carcinogenesis. Particularly, what are the genetic markers for disease and what

potential targets exist for novel treatments metastatic melanoma.

One emerging factor identified in a recent study of patient derived melanoma

samples is DNA polymerase-theta (Pol θ or POLQ)[5]. An A-family DNA repair enzyme,

Pol θ is essential for cell function and organismal development[6]. Inherently error prone

Pol θ[7,8] plays a predominant role repairing double strand breaks (DSB) in the DNA

strand through microhomology-mediated end joining (MMEJ, also known as theta-

mediated end joining, (TMEJ)), and translesion nucleotide bypass[9–12]. Unlike

homologous recombination (HR), the favored DSB repair pathway, TMEJ is highly error

prone and is proposed to be activated when HR is overwhelmed (when the genome

occurs many double strand breaks) and/or inactive (such as in cancer states). The

activity of Pol θ in translesion nucleotide bypass plays a critical role in replication, that

while perpetuating genomic mutations, allows replicative DNA polymerases to continue

replication while also avoiding more DNA DSBs and potential mutagenesis through

replication fork collapse[13]. Pol θ is also a key factor in repairing DBSs during early

embryonic development[6]. Together this indicates that Pol θ function is intrinsically

mutagenic yet required for cell function.

This duality, mutagenic enzymatic behavior while also supporting cell

survival[6,14,15], along with aberrant Pol θ activity in cancer cells[16–18] that has led to many

hypothesizing that Pol θ activity drives carcinogenesis. In the case of melanoma, it is

hypothesized carcinogenesis is driven in part by improper DNA repair of UV damage

generated by sun exposure[8,19]. UV radiation has the potential to create thymine dimers

on the DNA strand which either need to be resolved by base excision repair, a

mechanism with low mutagenic potential, or by translesion nucleotide bypass, a

mutagenic mechanism which Pol θ can perform. As mentioned before, bypassing or

repairing these damaged sites in the genome are important for genomic stability as they

will result in replication fork collapse and DSBs during replication. These DSBs could

potentially lead to greater mutagenic events if instead of HR, DSBs are repaired though

the MMEJ/TMEJ pathways by Pol θ. Interestingly, variant forms of Pol θ identified in

patient melanoma samples, display a range of enzymatic properties different from wild-

type (WT)[20]. These data suggest enhanced mutagenic capabilities for these versions of

Pol θ. However, few models for assessing the function of Pol θ as well as the outcome

of Pol θ function in the context of an organism exist.

Zebrafish have long been employed to model melanoma for organismal study[21].

With nearly 70% homology in their genes and 85% in human disease-related genes[22],

zebrafish have a high degree of similarity to humans. Like humans, zebrafish have

melanin producing melanocytes which are displayed most prominently in their horizonal

stripes. Zebrafish melanocytes produce melanin through pathways homologous to those

in mammals and develop from neural crest cells in the embryo[2]. Xenografts of human

melanoma tumors into zebrafish have also shown that these tumors have similar

metastatic tropism in zebrafish to those observed in human malignancies[23,24]. These

studies suggest many of the cellular cues of melanoma spread and disease are similar

between human and zebrafish.

Here we present the first comparative analysis of protein structure and function of

purified zebrafish POLQ (zPol θ or zPOLQ) and human POLQ (hPol θ or hPOLQ)

polymerase domains. Protein alignment indicates that many of the residues present in

the polymerase domain between the two proteins are conserved resulting in similar

folded structures. However, within loop regions (unresolved in the human crystal

structure), specific to PolQ relative to other A-type proteins, there is little conservation.

Despite this lack of conservation, we observe similar zPOLQ behavior compared to

hPOLQ. zPolQ can extend DNA templates, perform TMEJ, and bypass DNA lesions,

hallmarks of PolQ function in the cell.

MATERIALS AND METHODS

Materials

All materials were purchased from Sigma-Aldrich (St. Louis, MO), Bio-rad

Laboratories (Hercules, CA), AmericanBio (Canton, MA), and Research Products

(Mount Prospect, IL). DNA oligonucleotides were purchased from Integrated DNA

Technologies (Newark, NJ) and deoxynucleotides from New England Biolabs (Ipswich,

MA). All DNA oligos were purified via HPLC with standard desalting from the

manufacturer.

Zebrafish Pol θ cloning

Total RNA from 4-hour post fertilization embryos was extracted using TRIzol

(Invitrogen) following manufactures instructions. A library of cDNAs was generated from

the pool of polyA mRNAs using ProtoScript II Reverse Transcriptase (New England

Biolabs, NEB) following manufactures instructions, primed by oligo(dT). The polymerase

domain of zebrafish Pol θ was then amplified for cloning into the POLQM1 vector[12] a

pSUMO3 based expression vector. This was a two-step cloning process as the

POLQM1 vector did not have multiple cloning sites.

First, the polymerase domain of zebrafish Pol θ (residues 1801-2579) was

amplified from the cDNA library with primers contain a 5' KpnI site and a 3' BamHI site:

Pol θ RVS BamHI- TATACTGGATCCTTATATGTCCAGGTCTTGAAGGTTACC

Pol θ FWD KpnI- ATTAGGTACCTCAACATCAGTGTTAGGCGCAC

Second, the 6xHIS and SUMO sequences (HIS-SUMO) of POLQM1 were

amplified off of the plasmid using primers containing a 5' XbaI site and a 3' KpnI site:

HIS-SUMO RVS KpnI- ATTAGGTACCTCCCGTCTGCTGC

HIS-SUMO FWD XbaI- TTCCCCTCTAGAAATAATTTTGTTTAACTTTAAGAAG

PCRs reactions used Phusion High-Fidelity DNA Polymerase (NEB) following

manufacturer's instructions and were run for 30 cycles. PCR Products were gel isolated

from a 1% TAE agarose gel using Freeze 'N Squeeze DNA gel extraction columns (Bio-

Rad), following manufacturer's instructions.

Next, the zPol θ, HIS-SUMO, and POLQM1 DNAs were digested with

appropriate enzymes (NEB) overnight at 37□C:

zPol θ PCR - BamHI and KpnI

HIS-SUMO PCR - KpnI and XbaI

POLQM1 - XbaI and BamHI

Digested samples were gel separated on a 1% TAE agarose gel and fragments were

isolated using Freeze 'N Squeeze DNA gel extraction columns. zPol θ and HIS-SUMO

digested fragments were then ligated using T4 ligase (NEB) incubating at 16□C

overnight and gel isolated from a 1% TAE agarose gel using Freeze 'N Squeeze DNA

gel extraction columns. zPol θ-HIS-SUMO fragment was ligated into linearized

POLQM1 vector using T4 ligase and incubating at 16□C overnight. NEB 5-alpha

competent *E.coli* (NEB) were transformed by ligated productions using manufacturer's

instructions. Bacteria were selected for through ampicillin resistance.

zPol θ polymerase domain modeling and alignments

Amino acid sequence alignments were completed using EMBL-EBI Clustal

Omega MSA[25] on default settings. Structural rendering of the zPol θ polymerase

domain was completed using ColabFold v1.5.5: AlphaFold2 using MMseqs2[26–30] on

default settings. The resulting zPol θ structure was compared to the solved hPol θ

polymerase domain structure (AX0Q)[31] using the pairwise alignment tool in FATCAT[32]

on default setting, and visualized using RCSB PDB visualization tools[33].

Expression and Purification of hPol θ and zPol θ

Recombinant pSUMO3 plasmids containing the truncated polymerase domain of

hPol θ and zPol θ gene were expressed in *E.coli* and purified as previously described in

Thomas et al[34]. Briefly, Rosetta2(DE3) cells containing the zPol θ plasmid were

inoculated into autoinduction Terrific Broth and grown at 20°C for 60 hours. Cells were

harvested through centrifugation and lysed in Lysis buffer. After 6 rounds of sonication,

cell fractions were further separated via centrifugation. The fraction containing soluble

protein were applied to a 5mL His-Trap FF crude Nickel Column (Cytiva) by FPLC using

a high imidazole gradient. Fractions containing Pol θ were separated again on a HiTrap

Heparin HP (Cytiva) column for further purification. Eluate containing Pol θ was

incubated overnight at 4°C with SUMO2 Protease (Fisher Scientific) to remove the

6xHIS-SUMO tag. Untagged Pol θ was separated from the 6X-HIS-SUMO on a Hi-Trap

Chelating HP column, reserving the flowthrough that contained only Pol θ. A final

HiTrap Heparin column removed any remaining non-specific binding proteins and

exchanged the imidazole buffer for a high NaCl buffer. Protein purification was verified

on 10% denaturing SDS visualized on an Odyssey CL-x IR scanner (LiCOR). Purified

protein is highly unstable and was flash-frozen in liquid nitrogen and stored at -80°C for

3 months maximum.

DNA Substrate Generation

Double stranded DNA substrates (dsDNA) were generated using complementary

oligodeoxynucleotides from IDT. Templates representing cyclobutane pyrimidine

Thymine-Thymine dimer (CPD) damaged and undamaged were synthesized in the

Delaney laboratory (Sarah Delaney, Brown University).

The CPD-containing sequence is as follows: 5'-AAG AGT TCG AXX GCC TAC

ACT GGA GTA CCG GAG-3' where XX denotes the CPD lesion. The oligonucleotide

was synthesized on a MerMade 4 (BioAutomation) using standard phosphoramidite

chemistry. All reagents were purchased from Glen Research. The 5'-dimethoxytrityl

group was retained for HPLC purification (Agilent PLRP-S column, 250 mm × 4.6 mm;

mobile phase A = 1% acetonitrile, 10% triethylammonium acetate (TEAA), 89% water;

mobile phase B = 10% TEAA, 90% acetonitrile). The gradient was as follows: 95% A /

5% B to 65% A / 35% B over 35 min at 1 mL/min. Oligonucleotide was subject to

detritylation by incubation for 60 min at room temperature in 20% (v/v) aqueous glacial

acetic acid. The reaction was quenched by precipitation of the oligonucleotides in room-

temperature ethanol. A second HPLC purification was then performed (same column

and mobile phases as above) using the following gradient: 100% A / 0% B to 75% A /

25% B over 40 min at 1 mL/min. The purified oligonucleotide was flash-frozen with liquid

nitrogen and lyophilized.

The 5'6-FAM primers were annealed to complementary DNA templates with

sequence context as previously described [35,36] and are described below:

25/40 undamaged DNA substrate

`5'-/FAM/ TTT GCCT TGA CCA TGT AAC AGA GAG`

`        CGGA ACT GGT ACA TTG TCT CTC GCA CTC ACT CTC TTC`

`TCT`

24/33 CPD damaged DNA substrate

`5'-/FAM/-TTT CTC CGG TAC TCC AGT GTA GGC`

`        GAG GCC ATG AGG TCA CAT CCG `**`TT`**`A GCT TGA GAA`

24/33 undamaged DNA substrate

`5'-/FAM/-TTT CTC CGG TAC TCC AGT GTA GGC`

`        GAG GCC ATG AGG TCA CAT CCG `**`TT`**`A GCT TGA GAA`

Confirmation of annealed substrates was determined 12% Native PAGE and samples

scanned on an RB Amersham Typhoon Fluorescent Imager (Cytiva) with a FAM filter.

Single oligodeoxynucleotides were purchased from IDT for MMEJ with internal

consensus sequence as previously described[37].

`5'-/FAM/ GGT TAG CCC GGG`

Circular Dichroism and Melting Temperature

Secondary protein characteristics of hPol θ to zPol θ were determined on a J-

815-CD Spectropolarimeter (Jasco, Brown University) with a 0.2 cm quartz cuvette at

room temperature (20°C). For each sample, 3 µM of protein in 10 mM K_2HPO_4 buffer

were scanned in triplicate from 190-280 nm. The thermal denaturation profile was

determined by using the same instrument by heating the same sample from 20-90°C

with a 5°C/min temperature rate increase at 222 nm using the same. Data were

analyzed on Prism 10 GraphPad and the melting temperature (Tm) was estimated using

the halfway point of the denaturing curve

Electrophoretic Mobility Shift Assay (EMSA)

The DNA binding affinity constant $K_{D(DNA)}$ was determined as previously described[34].

zPol θ was titrated from 0-1000nM against 10nM 25/40 dsDNA substrate in binding

buffer and incubated for 1 hour at room temperature. Samples were separated on a 6%

Native PAGE and scanned on an RB Typhoon scanner (Cytiva) with the FAM

fluorescence filter.   Separated bound and unbound products were quantified using

ImageQuant. $K_{D(DNA)}$ was determined by equation 1.

$Y = \left[\frac{(mx)}{(x+K_D)}\right] + b$ Equation 1

Four replicates and two protein preparations were used to generate this data.

Rapid Chemical Quench Assay

Biphasic burst kinetics were measured as previously described[34]. Briefly,100 nM

Pol θ was pre-mixed with 300 nM 25/40 dsDNA substrate and rapidly mixed with 100

µM of dCTP (correct nucleotide) with 10 mM $MgCl_2$ using an RQF-3 Rapid Chemical

Quench instrument (KinTek Corporation) at 37°C between 0.004-0.6 seconds.

Reactions were quenched by addition of 0.5M EDTA. Products were separated on a

15% Urea-denaturing polyacrylamide gel and scanned using an Amersham Typhoon RB

Fluorescent imager (Cytiva). Extended product (n+1) was quantified using ImageQuant

software and then plotted to a full biphasic pre-steady state burst equation via non-

linear regression using Prism 9 GraphPad software (equation 2). A minimum of three

replicates were included for each assay on two independent protein preparations.

$$[Product] = [E]_{app}[\frac{k_{obs}^2}{(k_{obs}+k_{ss})^2} \times (1 - e^{-(k_{obs}+k_{ss})t} + \frac{k_{obs}k_{ss}}{(k_{obs}+k_{ss})}t]$$ Equation 2

Primer Extension Assays

Qualitative primer extension assays were performed as previously described[34].

Varying conditions were used to explore the primer extension capabilities between the

hPol θ and zPol θ. Under single-turnover conditions, excess Pol θ (200 nM) was pre-

incubated with 50 nM DNA and incubated for 5 minutes at 37°C. Nucleotide (125 μM) of

either none, all, individual nucleotides were preincubated with 20 mM *x* inorganic salt (x

= MgCl$_2$, CaCl$_2$, MnCl$_2$). Under Michaelis-Menton conditions, DNA (200nM) was in

excess to Pol θ (50 nM). All reactions were carried out in buffer containing 20 mM Tris

HCl, pH 8.0, 25 mM KCl, 4% glycerol, 1 mM βME, and 80 μg/mL BSA. The reaction was

initiated by combining Pol-θ/DNA with dNTP/salt. Reactions were incubated at 37°C for

5 minutes before being stopped by 80% Formamide/EDTA quench. Products were

separated out on a 15% urea-denaturing polyacrylamide gel, and scanned on a RB

Amersham Typhoon fluorescent imager (Cytiva).

MMEJ Assay

Microhomology mediated end-joining assay for both hPol θ and zPol θ were

carried out as previously described[37] on a 12-mer FAM labeled oligodeoxynucleotide.

Pol θ (20 nM) was preincubated with 30 nM 5'-FAM ssDNA in reaction buffer (25mM

Tris-HCl pH8.8, 1mM βME, 0.01% NP-40, 0.1 mg/mL BSA, 10% glycerol, 10 mM MgCl$_2$,

30mM NaCl) for 5 minutes at 37°C. Nucleotides (20 µM) were added and incubated at

37°C for an additional 45 minutes. Reactions were terminated by addition of non-

denaturing stop buffer (100 mM Tris-HCl pH 7.5, 10 mg/mL proteinase K, 80 mM EDTA,

and 0.5% SDS) for an additional 15 minutes. DNA products were separated on a 12%

native polyacrylamide gel and scanned by an RB fluorescent Amersham Typhoon

(Cytiva) with a FAM filter.

RESULTS

Zebrafish and human polymerase domains display high degree of structural similarity

To determine the degree of similarity between the zPol θ and hPol θ PD primary

amino acid sequences were aligned (Table 1, Supplemental figure 1). The alignment of

the full-length Pol θ protein indicates, zebrafish and human Pol θ share 46% identity.

This degree of similarity increases when comparing the predicted polymerase domain

(63%), as well as subdomains containing catalytic activity, fingers (75.3%), thumb

(74.6%), and palm (66.7%). These data suggest a structurally similar molecule.

To assess the extent of structural similarity we generated a predicted structure for

zPol θ PD using ColabFold[26] to compare to the solved crystal structure of hPol θ PD[31].

Upon visual inspection the predicted zPol θ PD displays classical DNA polymerase PD

structures (Figure 1). The three major subunits, the fingers, thumb, and palm are visible,

and when modeled in, a DNA molecule can fit in the presumed catalytic domain. The

model also indicates the presence of unstructured loop domains, that have functional

importance[12], that were not resolved in the hPol θ PD structure. An over lay of the hPol

θ PD and zPol θ PD show that the structures have a high degree of similarity. As

predicted by the amino-acid alignment, these data indicate that much of the structure of

the hPol θ PD is conserved in zPol θ.

Table 1. Sequence alignment analysis

	Polymerase theta comparisons		
	zebrafish	% identity	human
Polymerase Domain	thumb (122 aa)	74.6%	thumb (125 aa)
	fingers (142 aa)	75.3%	fingers (142 aa)
	palm (163 aa)	66.7%	palm (177 aa)
	total (744 aa)	63%	total (799 aa)
Full length	2576 aa	46%	2590 aa

Figure 1. Structural modeling of zPol θ. (A) AlphaFold rendering of zPol θ with subdomains colored, thumb – blue,

fingers – red, palm – green, and exo-nuclease in yellow. DNA is colored in light blue. (B) FATCAT overlay of AlphaFold rending of

zPol θ (bronze) and hPol θ (navy) crystal structure[31] without loop inserts, DNA is colored in light blue.

hPol θ and zPol θ are structural similar

The plasmid containing the c-terminal recombinant zPol θ was expressed and

purified in the same way as hPol θ[34] and as summarized in the Materials and Methods.

Similar to hPol θ, one protein preparation yields approximately 5-10 µM and we

observed similar expression and purification levels as seen with hPol θ (Figure 2A)

To confirm similarity in secondary structure between hPol θ and zPol θ, circular

dichroism spectroscopy (CD) was performed at 20°C. The same sample was heated

from 20-90°C in order to determine the thermal denaturation profile. Both spectra were

overlayed and indicated minimal variance suggesting that both hPol θ and zPol θ have

similar secondary characteristics and thermal stability with a Tm of about 55°C.

298
299

Figure 2. hPol θ and zPol θ display similar secondary characteristics and thermal stability. (A) Expression and purification of

zPol θ was the same as hPol θ as described in the Materials and Methods. For each sample, approximately 56-60 pmol of cleaved,

purified protein were loaded on a 10% SDS PAGE and Coomassie stained. Both hPol θ and zPol θ migrate to approximately 90

kDa as expected. (B) Circular dichroism spectra of 3 µM hPol θ (solid line) and zPol θ (dashed) proteins in 10 mM Potassium

Phosphate buffer. Samples were scanned from 190 to 280 nm. (C)The same samples were heated from 20-90°C and ellipticity

measured at 222 nm.

zPol θ binds to dsDNA substrate

DNA binding by a DNA polymerase is one of the first steps in its catalytic

mechanism. To determine the DNA binding capabilities of zPol θ, we titrated zPol θ from

0-1000 nM protein against 10 nM 25/40 dsDNA. Complexed DNA/protein products

were separated on a denaturing gel to determine a dissociation constant ($K_{D(DNA)}$) for

DNA binding. Similar to hPol θ, zPol θ has a low $K_{D(DNA)}$ value of approximately 19.8 ±

3.1 nM.

Figure 3. zPol θ binds tightly to ds DNA. zPol θ was titrated from 0-1000 nM against 10 nM 25/40 dsDNA. Bound and unbound

products were separated on a 6% non-denaturing gel and quantified using ImageQuant. $K_{D(DNA)}$ was mathematically calculated using

Equation 1 and is the midpoint between bound and unbound fractions.

zPol θ can extend dsDNA similar to hPol θ

The second step in the DNA polymerase catalytic pathway is nucleotide binding

and formation of the phosphodiester bond. To explore this fundamental step of DNA

Polymerase activity, we assayed zPol θ's ability to extend 25/40 dsDNA under varying

conditions. Under standard steady-state conditions, 200 nM of zPol θ or hPol θ was

pre-incubated with 50 nM 25/40 dsDNA. The reaction was initiated by the addition of

125 nM dNTP as described in Figure 4 along with 20 mM $MgCl_2$, the preferred metal for

DNA polymerase [38]. We observed under these conditions both hPol θ and zPol θ were

able to extend the full 18-mer template with all nucleotides present (Figure 4).   Both

enzymes were able to incorporate single nucleotides, correct and incorrect as well.

Notably, zPol θ was able to incorporate incorrect dGTP to full extension (n+1) compared

to only n+6 with hPol θ. DNA polymerases can utilize other metals including Mn^{2+}, and

we observed an increase in mutagenesis through misincorporation for both hPol θ and

zPol θ. When provided with all dNTP, zPol θ can extend past the end of the template

(n+18).   Overall, zPol θ experiences more extension products especially with incorrect

nucleotides dATP, dGTP, and dTTP compared with hPol θ under similar conditions.

Steady-state conditions highlight overall DNA polymerase activity, but because the

dsDNA substrate is in excess, activity highlights multiple turnovers[39].   Although DNA pol

θ has been shown to have robust de novo activity with manganese[40], we wanted to be

sure this over extension observed with zPol θ and Mn^{2+} was the result of extension and

not an artifact.   We changed the ratio of protein to DNA to reflect single-turnover

conditions; excess protein over dsDNA substrate.   Here we are able to observe

polymerization events for theoretically every available DNA substrate. Similar to steady-

state conditions, we observe an even more robust de novo extension with not only all

nucleotides, but also with dATP, suggesting that zPol θ misincorporation with dATP is

preferred (Supplement 2).

5'-/FAM/ TTT GCCT TGA CCA TGT AAC AGA GAG
         CGGA ACT GGT ACA TTG TCT CTC GCA CTC ACT CTC TTC TCT

n+18

n+1
n

None All dATP dCTP dGTP dTTP None All dATP dCTP dGTP dTTP None All dATP dCTP dGTP dTTP None All dATP dCTP dGTP dTTP

hPol θ   zPolθ   hPol θ   zPol θ

Mg²⁺   Mn²⁺

zPol θ catalytic activity similar to other DNA polymerases

To further explore the mechanism of nucleotide incorporation of zebrafish Pol θ,

we assayed zPol θ under presteady-state conditions in which there is an excess DNA

substrate to enzyme with correct nucleotide. This assay focuses on the DNA

polymerase ability to extend DNA by incorporating the correct nucleotide opposite a

templating base. This activity is biphasic in which there is a rapid polymerization step of

nucleotide incorporation at the DNA primer's 3'OH and a slower, rate limiting step of

product release[39].   If biphasic activity is not observed, it suggests a step before

nucleotide incorporation is the rate-limiting step[41].   To ensure that purified c-terminal

zPol θ follows the traditional DNA polymerase mechanism, 100 nM zPol θ was

preincubated with 300 nM 25/40 dsDNA.   The DNA/Pol θ complex was rapidly

combined with 100 μM correct nucleotide and 10 mM $MgCl_2$ from 0.004-0.6 seconds.

DNA products were separated on a denaturing polyacrylamide gel and primer extension

of n+1 was quantified and data fit to a full biphasic burst equation.   zPol θ fit to a

biphasic equation with an observable polymerization rate (k_{obs}) of 15.9 ± 2.5 s^{-1} (Figure

5).

Figure 5. zPol θ experiences biphasic burst activity. Zebrafish Pol θ (100 nM) was preincubated with 300 nM 25/40 dsDNA. The

DNA/Pol θ complex was rapidly combined with 100 μM dCTP (correct nucleotide) and 10 mM $MgCl_2$. Reactions were carried out at

37°C and quenched with 0.5 M EDTA. Products were separated on a denaturing gel and quantified with ImageQuant software.

Data were fit to a biphasic burst equation to obtain observed k_{obs} rates 15.9 ± s^{-1}. The slower rate k_{ss} was calculated to be 3.4 ± 0.46

s^{-1}.

zPol θ performs MMEJ activity

One of the major functions of DNA Pol θ is its ability to repair double-strand

breaks and is the primary DNA polymerase for microhomology-mediated end joining. In

doing so, Pol θ utilizes internal homology within the DNA sequence to act as a template.

Pol θ aligns these complementary pieces and extends in the 5' to 3' direction[37,42,43].

Truncated hPol θ has been shown to able to perform MMEJ activity on short 12-mer

single-stranded DNA, but the full 290 kDa Pol θ with the N-terminal helicase and central

domains are needed to anneal and extend larger segments of DNA[37]. We wanted

ensure that zPol θ could also perform MMEJ in a similar manner to hPol θ on short

fragments of DNA. Figure 6 is a representative gel of hPol θ and zPol θ performing

MMEJ on a ssDNA. As indicated in the schematic above, the CCCGGG are aligned

through Pol θ in the presence of (+) dNTP and subsequently extended in the opposite

direction giving rise to a slower moving double-stranded DNA product. Both hPol θ and

zPol θ are able to perform this activity. We hypothesize the smaller product bands are

indicative of classic snap-back synthesis in which the DNA substrate anneals onto itself

for Pol θ to extend. This behavior has been observed by others on hPol θ and there is

little variation between the two species[37].

zPol θ is able to bypass CPD lesion DNA.

Pol θ is a versatile DNA polymerase in not only can it perform MMEJ, it has also

been shown to bypass cyclobutane pyrimidine dimers (CPD)[36]. By being able to extend

a DNA primer passed a template containing a contorted Thymine-Thymine lesion,

human and mouse Pol θ have been demonstrated to be critical in suppressing DNA

damage and preventing skin lesions. On a molecular level, human Pol θ has

demonstrated that not only can it insert opposite the initial T in the T-T dimer but is able

to mutagenically extend past this lesion for the remaining DNA template. We

hypothesized that zPol θ has the same ability to bypass CPD lesions and we assayed

both Pol θ under single-turnover conditions (4:1 protein to DNA) in the presence of a

24/33 CPD damaged DNA template with both Mg^{2+} and Mn^{2+}. As predicted, there was

little variance in bypass activity of zPol θ compared to hPol θ. Both enzymes were able

to readily insert opposite a T-T dimer as well as extend past this lesion with both all

dNTPs present and dATP and to some extent dGTP. Both Pol θs could not incorporate

dCTP opposite T-T, but we observed only insertion of dTTP opposite and no extension.

In the presence of Mn^{2+}, both Pol θs readily bypassed T-T dimers, again demonstrating

de novo synthesis past the template. zPol θ was more robust in extension with the other

incorrect nucleotides suggesting Mn^{2+} has an increased mutagenic effect.

Figure 7. zPol θ is able to bypass CPD lesions. As described in the Materials and Methods, 200 nM Pol θ (human or zebrafish) was preincubated with 50 nM CPD Damaged DNA substrate. Reactions were initiated by the addition of 125 nM nucleotides as described and either 10 mM $MgCl_2$ or $MnCl_2$. Reactions were carried out at 37°C for 5 minutes and products visualized on a 12% denaturing gel. Higher migrating products are indicative of full extension (n+12) with smeared bands representing de novo synthesis with extension past n+12.

zPol θ experiences unusual extension of DNA substrates in the presence of Ca^{2+}.

To explore the role of divalent metals in DNA polymerase activity for Pol θ, we

performed a DNA polymerase extension assay again with either these assays with their

specific DNA substrates swapping out the active metals for Ca^{2+} which has traditionally

used as an inert control. Unlike the other divalent metals, Ca^{2+} allows for ternary

complex formation, but extension is limited or slow[44,45]. Using 50 nm if the 24/33

undamaged and CPD damaged DNA substrate, we performed a primer extension assay

with 200 nM hPol θ or z WT Pol θ with Mg^{2+} substituted for $CaCl_2$. Extension products

were separated on a denaturing polyacrylamide gel and quantified based on the percent

extension. Figure 8A is a representative gel of extension on 24/33 undamaged DNA

template. We observe that hPol θ could incorporate every nucleotide to some extent,

with an n+3 extension product only observed in the presence of all dNTP or purines.

zPol θ was observed to generate full extension product (94%) on this DNA template

(n+12) with all nucleotides present and, like hPol θ, could extend with purines as well.

Incorporation of dATP led to 92% conversion to product although the enzyme stalled

around n+2. Interestingly, zPol θ appears to skip the first thymine in the undamaged

sequence for both all nucleotides and dATP. The same experiment was carried out with

24/33 CPD damaged DNA. Under these conditions we report that Ca^{2+} reduced DNA

polymerase activity for both hPol θ and zPol θ with incorporation of only one nucleotide

irrespective if that nucleotide was matched or mismatched with the templating base.

449
Figure 8. DNA Pol θ is able incorporate and extend dsDNA in the presence of Ca^{2+}. Under single turnover conditions 200 nM

Pol θ (human or zebrafish) was preincubated with 50 nM 24/33 undamaged (A) or damaged (B) DNA substrate and reacted with 125

nM nucleotides in 10 mM CaCl$_2$. Reactions were carried out at 37°C for 5 minutes and products visualized on a 12% denaturing

gel. Percent extended was calculated using ImageQuant software by quantifying the intensity of the extended products (n+1 and

higher) divided by the intensity of the total amount of DNA.

DISCUSSION

Zebrafish and human Pol θ structures have a high degree of similarity

Comparisons of the amino acid sequence (table 1) and the structures (Figure 1)

of zPol θ and hPol θ reveal that the two proteins share a high degree of similarity.

Importantly, and perhaps not surprisingly, the areas of greatest similarity are around the

catalytic subdomains, the fingers, thumb, and palm, of the polymerase domain. These

sites of activity would be important to the protein function across evolutionary time.

Interestingly, unlike other A-type polymerase family members Pol θ has three loop

structures within the PD which have been identified to be important for function[12].

Comparison of the zPol θ and hPol θ sequences indicate that zPol θ also contains these

inserted loops however, they contain little homology (Supplement 1) to that observed

between human and mouse[31] Pol θ. Despite this difference, expression and purification

yields of zPol θ were similar to that of hPol θ as were the secondary structural

characteristics and thermal stability (Figure 2); an early indication similar protein folding.

We also show here that zPol θ still retains the same activity observed in hPol θ.

Zebrafish Pol θ extends dsDNA

DNA polymerase θ primary role in the human cell is the primary DNA polymerase

repair enzyme in microhomology-mediated end joining and thus, the major goal of this

study was to query if zPol θ retained a similar function. Initially we simplify the activity

by asking 'can zPol θ bind to a primer/template dsDNA substrate and then extend it?'

Our data suggests that yes it can. zPol θ binds tightly to this substrate (Figure 3)[34]

similar to values obtained with hPol θ. This is expected because loop 1 which is located

in the thumb domain or DNA binding domain is thought to be involved with contacts to

DNA[12] and is the only conserved loop region between zPol θ and hPol θ. zPol θ can in

fact extend a DNA substrate, and we show it has robust activity on this particular DNA

substrate (Figure 4) especially in the presence of Mn^{2+}. While most of the data

presented in this work was qualitative, probing how fast a DNA polymerase makes a

phosphodiester bound through biochemical kinetics can provide insight into mechanism

of incorporation[39]. Like most DNA polymerases, zPol θ performs biphasic burst kinetics

which is indicative of a two-step mechanism with a rate limiting step of product release.

zPol θ experiences an observed polymerization rate of around 16 s^{-1} (Figure 5) which is

almost 4 times slower than its human ortholog[34], but not uncommonly slow as a similar

DNA Polymerase β experiences a similar rate[46–48]. Why zPol θ might experience a

slower rate is unknown. It could be due to the lack of conservation within the looping

structures in the palm domain which in human Pol θ may drive substrate alignment for

rapid polymerization.

Zebrafish Pol θ is able to perform microhomology-mediated End Joining

Despite being able to extend DNA, it is important that zPol θ also be able to

perform MMEJ as in humans it is its primary function. Although a majority of human Pol

θ's N-terminal and central domains are critical for this function, studies have shown that

the c-terminal polymerase domain of Pol θ does retain limited function for aligning and

extending short single-stranded DNA[37]. zPol θ was no exception (Figure 6) and was

able to complement two single-strands and extend which is perhaps the most

compelling evidence of homologous function.

zPol θ can bypass CPD lesions

Another function of human Pol θ is its ability to bypass DNA damage and a more

recent study in mice suggest that bypassing UV damage is critical in the prevention of

skin cancer[36]. Our data provides evidence that zPol θ is able to bypass and extend

CPD lesions *in vivo* similar to that of hPol θ (Figure 7). Translesion bypass activity has

been highlighted as a function of loops 2 and 3 in human Pol θ[12,31]. Surprisingly,

zebrafish display very little homology through similar inserts (Supplement 1). However,

our studies might suggest that the only critical residues for this function are isolated to

the c-terminal end of this insert beginning with the sequence GMXFSXSMR. Further

studies exploring this insert in zebrafish are needed to determine if these conserved

amino acids are truly critical and that the function is either dictated through the overall

presence of the loop or that the loop dependent activities require the few conserved

amino acids retain in zPol θ.

zPol θ experiences Ca^{2+} dependent polymerization

Our data shows that zPol θ retains all of the DNA polymerase activities of hPol θ,

with the exception being the robust Ca^{2+} dependency during DNA extension (Figure 8).

While it is unusual to see DNA polymerases extend past the initial insertion event, there

have been instances where high-fidelity *Sulfolobus solfataricus* Dpo4 polymerase uses

calcium [49]. Whether or not swapping Ca^{2+} in zebrafish makes it a faster or mutagenic

polymerase has not been explored but evolutionarily it is possible that Ca^{2+} makes for a

stable ion swap for structural alignment and catalytic activity. Given that in freshwater

contains nearly identical concentrations of the divalent ions[50], its plausible that Ca^{2+}

could readily be a co-factor substitute.

Our data presented is clear evidence that zebrafish Pol θ is a homolog to human

Pol θ and that structurally and enzymatically behave with similar functions. This study is

significant as it highlights the availability of zebrafish as a model organism for studying

Pol θ and its potential function in DNA repair and disease. In particular given the robust

assortment of tools zebrafish offers a powerful, functionally relevant model for human

melanoma. Future studies could introduce patient derived mutations by way of germline

alterations and study the effects over the life time of the animal. Thus, adding new

insights into potential disease markers and mechanisms of disease progression and

treatment.

ACKNOWLEDGEMENTS

We would like to thank Sarah Delaney and Mary Tarantino from Brown University for the generation of the CPD damaged DNA. Thank you to Sylvie Doublié from University of Vermont for the human pol θ plasmid.

FUNDING

Research reported in this publication was supported by the Rhode Island Institutional Development Award (IDeA) Network of Biomedical Research Excellence under P20GM103430 and in part by the National Institute of General Medical Sciences of the National Institutes of Health under grant number R15GM144903-01. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

WORK CITED

1. Arnold, M. *et al.* Global Burden of Cutaneous Melanoma in 2020 and Projections to 2040. *JAMA Dermatol* **158**, 495–503 (2022).

2. Mort, R. L., Jackson, I. J. & Patton, E. E. The melanocyte lineage in development and disease. *Development* **142**, 620–632 (2015).

3. Jilani, S. *et al.* CAR-T cell therapy targeting surface expression of TYRP1 to treat cutaneous and rare melanoma subtypes. *Nat Commun* **15**, 1244 (2024).

4. Larkin, J. *et al.* Combined Nivolumab and Ipilimumab or Monotherapy in Previously Untreated Melanoma. *N Engl J Med* **373**, 23–34 (2015).

5. Farshidfar, F. *et al.* Integrative molecular and clinical profiling of acral melanoma links focal amplification of 22q11.21 to metastasis. *Nat Commun* **13**, 898 (2022).

6. Thyme, S. B. & Schier, A. F. Polq-mediated end joining is essential for surviving DNA double-strand breaks during early zebrafish development. *Cell Rep* **15**, 707–714 (2016).

7. Brambati, A., Barry, R. M. & Sfeir, A. DNA polymerase theta (Polθ) – an error-prone polymerase necessary for genome stability. *Current Opinion in Genetics & Development* **60**, 119–126 (2020).

8. Yoon, J.-H. *et al.* Error-Prone Replication through UV Lesions by DNA Polymerase θ Protects against Skin Cancers. *Cell* **176**, 1295-1309.e15 (2019).

9. Chan, S. H., Yu, A. M. & McVey, M. Dual Roles for DNA Polymerase Theta in Alternative End-Joining Repair of Double-Strand Breaks in Drosophila. *PLOS Genetics* **6**, e1001005 (2010).

10. Kent, T., Chandramouly, G., McDevitt, S. M., Ozdemir, A. Y. & Pomerantz, R. T.

Mechanism of microhomology-mediated end-joining promoted by human DNA

polymerase θ. *Nat Struct Mol Biol* **22**, 230–237 (2015).

11. Yousefzadeh, M. J. *et al.* Mechanism of Suppression of Chromosomal Instability by

DNA Polymerase POLQ. *PLOS Genetics* **10**, e1004654 (2014).

12. Hogg, M., Seki, M., Wood, R. D., Doublié, S. & Wallace, S. S. Lesion Bypass Activity

of DNA Polymerase θ (POLQ) Is an Intrinsic Property of the Pol Domain and

Depends on Unique Sequence Inserts. *Journal of Molecular Biology* **405**, 642–652

(2011).

13. Yu, A. M. & McVey, M. Synthesis-dependent microhomology-mediated end joining

accounts for multiple types of repair junctions. *Nucleic Acids Res* **38**, 5706–5717

(2010).

14. Ceccaldi, R. *et al.* Homologous-recombination-deficient tumours are dependent on

Polθ-mediated repair. *Nature* **518**, 258–262 (2015).

15. Goff, J. P. *et al.* Lack of DNA Polymerase θ (POLQ) Radiosensitizes Bone Marrow

Stromal Cells In Vitro and Increases Reticulocyte Micronuclei after Total-Body

Irradiation. *Radiat Res* **172**, 165–174 (2009).

16. Goullet de Rugy, T. *et al.* Excess Polθ functions in response to replicative stress in

homologous recombination-proficient cancer cells. *Biology Open* **5**, 1485–1492

(2016).

17. Allera-Moreau, C. *et al.* DNA replication stress response involving PLK1, CDC6,

POLQ, RAD51 and CLASPIN upregulation prognoses the outcome of early/mid-

stage non-small cell lung cancer patients. *Oncogenesis* **1**, e30–e30 (2012).

18. Lemée, F. *et al.* DNA polymerase θ up-regulation is associated with poor survival in

breast cancer, perturbs DNA replication, and promotes genetic instability.

Proceedings of the National Academy of Sciences **107**, 13390–13395 (2010).

19. Seki, M. *et al.* High efficiency bypass of DNA damage by human DNA polymerase

Q. *The EMBO Journal* **23**, 4484–4494 (2004).

20. Thomas, C. *et al.* Melanoma-Derived DNA Polymerase Theta Variants Exhibit

Altered DNA Polymerase Activity. *Biochemistry* **63**, 1107–1117 (2024).

21. Patton, E. E. *et al.* BRAF mutations are sufficient to promote nevi formation and

cooperate with p53 in the genesis of melanoma. *Curr Biol* **15**, 249–254 (2005).

22. Howe, K. *et al.* The zebrafish reference genome sequence and its relationship to the

human genome. *Nature* **496**, 498–503 (2013).

23. Groenewoud, A. *et al.* Patient-derived zebrafish xenografts of uveal melanoma

reveal ferroptosis as a drug target. *Cell Death Discov.* **9**, 1–11 (2023).

24. Fazio, M., Ablain, J., Chuan, Y., Langenau, D. M. & Zon, L. I. Zebrafish patient

avatars in cancer biology and precision cancer therapy. *Nat Rev Cancer* **20**, 263–

273 (2020).

25. Madeira, F. *et al.* The EMBL-EBI Job Dispatcher sequence analysis tools framework

in 2024. *Nucleic Acids Res* **52**, W521–W525 (2024).

26. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods*

19, 679–682 (2022).

27. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

(2021) doi:10.1038/s41586-021-03819-2.

28. Mirdita, M., Steinegger, M. & S"oding, J. MMseqs2 desktop and local web server

app for fast, interactive sequence searches. *Bioinformatics* **35**, 2856–2858 (2019).

29. Mirdita, M. *et al.* Uniclust databases of clustered and deeply annotated protein

sequences and alignments. *Nucleic Acids Res.* **45**, D170–D176 (2017).

30. Mitchell, A. L. *et al.* MGnify: the microbiome analysis resource in 2020. *Nucleic Acids*

Res. (2019) doi:10.1093/nar/gkz1035.

31. Zahn, K. E., Averill, A. M., Aller, P., Wood, R. D. & Doublié, S. Human DNA

polymerase θ grasps the primer terminus to mediate DNA repair. *Nat Struct Mol Biol*

22, 304–311 (2015).

32. Li, Z., Jaroszewski, L., Iyer, M., Sedova, M. & Godzik, A. FATCAT 2.0: towards a

better understanding of the structural diversity of proteins. *Nucleic Acids Res* **48**,

W60–W64 (2020).

33. Bittrich, S., Segura, J., Duarte, J. M., Burley, S. K. & Rose, Y. RCSB protein Data

Bank: exploring protein 3D similarities via comprehensive structural alignments.

Bioinformatics **40**, btae370 (2024).

34. Thomas, C. *et al.* Melanoma-Derived DNA Polymerase Theta Variants Exhibit

Altered DNA Polymerase Activity. *Biochemistry* **63**, 1107–1117 (2024).

35. Prasad, R. *et al.* Human DNA polymerase θ possesses 5′-dRP lyase activity and

functions in single-nucleotide base excision repair in vitro. *Nucleic Acids Research*

37, 1868–1877 (2009).

36. Yoon, J. H. *et al.* Error-Prone Replication through UV Lesions by DNA Polymerase θ

Protects against Skin Cancers. *Cell* **176**, 1295-1309.e15 (2019).

37. Black, S. J. *et al.* Molecular basis of microhomology-mediated end-joining by purified

full-length Polθ. *Nature Communications* **10**, (2019).

38. Steitz, T. A. DNA Polymerases: Structural Diversity and Common Mechanisms *.

Journal of Biological Chemistry **274**, 17395–17398 (1999).

39. Johnson, K. A. Transient-State Kinetic Analysis of Enzyme Reaction Pathways.

Enzymes **20**, 1–61 (1992).

40. Kent, T., Mateos-Gomez, P. A., Sfeir, A. & Pomerantz, R. T. Polymerase θ is a robust

terminal transferase that oscillates between three different mechanisms during end-

joining. *eLife* **5**, e13740.

41. Joyce, C. M. Techniques used to study the DNA polymerase reaction pathway.

Biochimica et biophysica acta **1804**, 1032–1040 (2010).

42. Brambati, A., Barry, R. M. & Sfeir, A. DNA polymerase theta (Polθ) – an error-prone

polymerase necessary for genome stability. *Current Opinion in Genetics and*

Development **60**, 119–126 (2020).

43. Li, C. *et al.* Structural basis of DNA polymerase θ mediated DNA end joining. *Nucleic*

Acids Research **51**, 463–474 (2023).

44. Freudenthal, B. D., Beard, W. A., Shock, D. D. & Wilson, S. H. Observing a DNA

Polymerase Choose Right from Wrong. *Cell* **154**, 157–168 (2013).

45. Nakamura, T., Zhao, Y., Yamagata, Y., Hua, Y. J. & Yang, W. Watching DNA

polymerase eta make a phosphodiester bond. *Nature* **487**, 196–201 (2012).

46. Towle-Weicksel, J. B. *et al.* Fluorescence resonance energy transfer studies of DNA

polymerase β the critical role of fingers domain movements and a novel non-

covalent step During nucleotide selection. *Journal of Biological Chemistry* **289**,

16541–16550 (2014).

47. Werneburg, B. G. *et al.* DNA Polymerase β:□ Pre-Steady-State Kinetic Analysis and

Roles of Arginine-283 in Catalysis and Fidelity†. *Biochemistry* **35**, 7041–7050

(1996).

48. Murphy, D. L., Kosa, J., Jaeger, J. & Sweasy, J. B. The Asp285 Variant of DNA

Polymerase Beta Extends Mispaired Primer Termini via Increased Nucleotide

Binding. *Biochemistry* **47**, 8048–8057 (2008).

49. Ralec, C., Henry, E., Lemor, M., Killelea, T. & Henneke, G. Calcium-driven DNA

synthesis by a high-fidelity DNA polymerase. *Nucleic Acids Res* **45**, 12425–12440

(2017).

50. Guerin, A. J. *et al.* High-resolution freshwater dissolved calcium and pH data layers

for Canada and the United States. *Sci Data* **11**, 370 (2024).

674