Zebrafish Pol Theta and Human Pol Theta: Orthologues with Homologous Function.

Cory Thomas[1], Sidney Green[1], Lily Kimball[2], Isaiah Schmidtke[2], Makayla Griffin[2], Lauren Rothwell[2], Ivy Par[1], Sophia Schobel[1], Yayleene Palacio[1], Jamie B Towle-Weicksel[1], Steven E Weicksel[2]

[1]Department of Physical Sciences, Rhode Island College, Providence, RI

[2]Department of Biology and Biological Sciences, Bryant University, Smithfield RI

ABSTRACT

DNA polymerase theta (Pol θ) is a conserved an A-family polymerase that plays an essential role in repairing double strand breaks, through micro-homology end joining, and bypassing DNA lesions, through translesion synthesis, to protect genome integrity. Despite its essential role in DNA repair, Pol θ is inherently error prone. Recently, Pol θ has been identified in various cancer types, suggesting Pol θ's promiscuous nature aids in cancer progression.

Here we present a study comparing the structure and function of the polymerase domain of zebrafish and human Pol θ to determine the potential of zebrafish as a model for pol θ function. We show that these two proteins share a large amount of sequence and structural homology, but differ in key loop areas thought to drive defining functions of the enzyme. Despite these differences zebrafish Pol θ still displays characteristics identify in human Pol θ, including DNA template extension, microhomology-mediated end

joining, and translesion synthesis. Additionally, we found several important residues within

loops of the polymerase domain that support function unique to Pol θ. These results will

support future studies with zebrafish as a model to investigate Pol θ function.

INTRODUCTION

Melanoma is the rarest but deadliest form of skin cancer, with an estimate in 2020

of 325,000 new cases worldwide, increase to nearly 510,000 new cases by 2040[1]. Unlike

the expansion of melanocytes horizontally along the epidermis as benign nevi, (commonly

referred to a moles) melanoma is characterized by expansion vertically, down, to and

through the basement membrane where cells then begin to spread out from the original

tumor location. Melanoma cells are highly mobile in part due to the migratory nature of

melanocytes[2], making migration at key tropism of the disease. Surgical treatments prior

to migration of melanoma to the lymph nodes have a high degree of success, with low

rates of tumor reoccurrence and favorable patient outcomes. Patient outcomes, however,

become less favorable after migration, as metastatic tumors often are no longer sensitive

to radiation or chemotherapy-based approaches. Recent advances in new immune-

boosting therapies have shown promise in treating metastatic melanoma, but early

detection and tumor resection treatments are still the most effective[3,4]. This indicates that

there is still much to know about melanoma carcinogenesis. Particularly, what are the

genetic markers for disease and what potential targets exist for novel treatments

metastatic melanoma.

One emerging factor identified in a recent study of patient derived melanoma

samples is DNA polymerase-theta (Pol θ or POLQ)[5]. An A-family DNA repair enzyme,

Pol θ is essential for cell function and organismal development[6]. Inherently error prone

Pol θ[7,8] plays a predominant role repairing double strand breaks (DSB) in the DNA strand

through microhomology-mediated end joining (MMEJ, also known as theta-mediated end

joining, (TMEJ)), and translesion nucleotide bypass[9–12].   Unlike homologous

recombination (HR), the favored DSB repair pathway, TMEJ is highly error prone and is

proposed to be activated when HR is overwhelmed (when the genome occurs many

double strand breaks) and/or inactive (such as in cancer states). The activity of Pol θ in

translesion nucleotide bypass plays a critical role in replication, that while perpetuating

genomic mutations, allows replicative DNA polymerases to continue replication while also

avoiding more DNA DSBs and potential mutagenesis through replication fork collapse[13].

Pol θ is also a key factor in repairing DBSs during early embryonic development[6].

Together this indicates that Pol θ function is intrinsically mutagenic yet required for cell

function.

This duality, mutagenic enzymatic behavior while also supporting cell survival[6,14,15],

along with aberrant Pol θ activity in cancer cells[16–18] that has led to many hypothesizing

that Pol θ activity drives carcinogenesis. In the case of melanoma, it is hypothesized

carcinogenesis is driven in part by improper DNA repair of UV damage generated by sun

exposure[8,19]. UV radiation has the potential to create thymine dimers on the DNA strand

which either need to be resolved by base excision repair, a mechanism with low

mutagenic potential, or by translesion nucleotide bypass, a mutagenic mechanism which

Pol θ can perform. As mentioned before, bypassing or repairing these damaged sites in

the genome are important for genomic stability as they will result in replication fork

collapse and DSBs during replication. These DSBs could potentially lead to greater

mutagenic events if instead of HR, DSBs are repaired though the MMEJ/TMEJ pathways

by Pol θ. Interestingly, variant forms of Pol θ identified in patient melanoma samples,

display a range of enzymatic properties different from wild-type (WT)[20]. These data

suggest enhanced mutagenic capabilities for these versions of Pol θ. However, few

models for assessing the function of Pol θ as well as the outcome of Pol θ function in the

context of an organism exist.

Zebrafish have long been employed to model melanoma for organismal study[21].

With nearly 70% homology in their genes and 85% in human disease-related genes[22],

zebrafish have a high degree of similarity to humans. Like humans, zebrafish have

melanin producing melanocytes which are displayed most prominently in their horizonal

stripes. Zebrafish melanocytes produce melanin through pathways homologous to those

in mammals and develop from neural crest cells in the embryo[2]. Xenografts of human

melanoma tumors into zebrafish have also shown that these tumors have similar

metastatic tropism in zebrafish to those observed in human malignancies[23,24]. These

studies suggest many of the cellular cues of melanoma spread and disease are similar

between human and zebrafish.

Here we present the first comparative analysis of protein structure and function of

purified zebrafish POLQ (zPol θ or zPOLQ) and human POLQ (hPol θ or hPOLQ)

polymerase domains. Protein alignment indicates that many of the residues present in

the polymerase domain between the two proteins are conserved resulting in similar folded

structures. However, within loop regions (unresolved in the human crystal structure),

specific to PolQ relative to other A-type proteins, there is little conservation. Despite this

lack of conservation, we observe similar zPOLQ behavior compared to hPOLQ. zPolQ

can extend DNA templates, perform TMEJ, and bypass DNA lesions, hallmarks of PolQ

function in the cell.

MATERIALS AND METHODS

Materials

97   All materials were purchased from Sigma-Aldrich (St. Louis, MO), Bio-rad

98 Laboratories (Hercules, CA), AmericanBio (Canton, MA), and Research Products (Mount

99 Prospect, IL). DNA oligonucleotides were purchased from Integrated DNA Technologies

100 (Newark, NJ) and deoxynucleotides from New England Biolabs (Ipswich, MA). All DNA

101 oligos were purified via HPLC with standard desalting from the manufacturer.

Zebrafish Pol θ cloning

Total RNA from 4-hour post fertilization embryos was extracted using TRIzol

(Invitrogen) following manufactures instructions. A library of cDNAs was generated from

the pool of polyA mRNAs using ProtoScript II Reverse Transcriptase (New England

Biolabs, NEB) following manufactures instructions, primed by oligo(dT). The polymerase

domain of zebrafish Pol θ was then amplified for cloning into the POLQM1 vector[12] a

pSUMO3 based expression vector. This was a two-step cloning process as the POLQM1

vector did not have multiple cloning sites.

First, the polymerase domain of zebrafish Pol θ (residues 1801-2579) was

amplified from the cDNA library with primers contain a 5' KpnI site and a 3' BamHI site:

Pol θ RVS BamHI- TATACTGGATCCTTATATGTCCAGGTCTTGAAGGTTACC

Pol θ FWD KpnI- ATTAGGTACCTCAACATCAGTGTTAGGCGCAC

Second, the 6xHIS and SUMO sequences (HIS-SUMO) of POLQM1 were

amplified off of the plasmid using primers containing a 5' XbaI site and a 3' KpnI site:

HIS-SUMO RVS KpnI- ATTAGGTACCTCCCGTCTGCTGC

HIS-SUMO FWD XbaI- TTCCCCTCTAGAAATAATTTTGTTTAACTTTAAGAAG

PCRs reactions used Phusion High-Fidelity DNA Polymerase (NEB) following

manufacturer's instructions and were run for 30 cycles. PCR Products were gel isolated

from a 1% TAE agarose gel using Freeze 'N Squeeze DNA gel extraction columns (Bio-

Rad), following manufacturer's instructions.

Next, the zPol θ, HIS-SUMO, and POLQM1 DNAs were digested with appropriate

enzymes (NEB) overnight at 37ºC:

zPol θ PCR - BamHI and KpnI

HIS-SUMO PCR - KpnI and XbaI

POLQM1 - XbaI and BamHI

Digested samples were gel separated on a 1% TAE agarose gel and fragments were

isolated using Freeze 'N Squeeze DNA gel extraction columns. zPol θ and HIS-SUMO

digested fragments were then ligated using T4 ligase (NEB) incubating at 16ºC overnight

and gel isolated from a 1% TAE agarose gel using Freeze 'N Squeeze DNA gel extraction

columns. zPol θ-HIS-SUMO fragment was ligated into linearized POLQM1 vector using

T4 ligase and incubating at 16ºC overnight. NEB 5-alpha competent *E.coli* (NEB) were

transformed by ligated productions using manufacturer's instructions. Bacteria were

selected for through ampicillin resistance.

zPol θ polymerase domain modeling and alignments

Amino acid sequence alignments were completed using EMBL-EBI Clustal Omega

MSA[25] on default settings. Structural rendering of the zPol θ polymerase domain was

completed using ColabFold v1.5.5: AlphaFold2 using MMseqs2[26–30] on default settings.

The resulting zPol θ structure was compared to the solved hPol θ polymerase domain

structure (AX0Q)[31] using the pairwise alignment tool in FATCAT[32] on default setting, and

visualized using RCSB PDB visualization tools[33].

Expression and Purification of hPol θ and zPol θ

Recombinant pSUMO3 plasmids containing the truncated polymerase domain of

hPol θ and zPol θ gene were expressed in *E.coli* and purified as previously described in

Thomas et al[34]. Briefly, Rosetta2(DE3) cells containing the zPol θ plasmid were inoculated

into autoinduction Terrific Broth and grown at 20°C for 60 hours. Cells were harvested

through centrifugation and lysed in Lysis buffer. After 6 rounds of sonication, cell fractions

were further separated via centrifugation. The fraction containing soluble protein were

applied to a 5mL His-Trap FF crude Nickel Column (Cytiva) by FPLC using a high

imidazole gradient. Fractions containing Pol θ were separated again on a HiTrap Heparin

HP (Cytiva) column for further purification. Eluate containing Pol θ was incubated

overnight at 4°C with SUMO2 Protease (Fisher Scientific) to remove the 6xHIS-SUMO

tag. Untagged Pol θ was separated from the 6X-HIS-SUMO on a Hi-Trap Chelating HP

column, reserving the flowthrough that contained only Pol θ. A final HiTrap Heparin

column removed any remaining non-specific binding proteins and exchanged the

imidazole buffer for a high NaCl buffer. Protein purification was verified on 10% denaturing

SDS visualized on an Odyssey CL-x IR scanner (LiCOR). Purified protein is highly

unstable and was flash-frozen in liquid nitrogen and stored at -80°C for 3 months

maximum.

DNA Substrate Generation

Double stranded DNA substrates (dsDNA) were generated using complementary

oligodeoxynucleotides from IDT.   Templates representing cyclobutane pyrimidine

Thymine-Thymine dimer (CPD) damaged and undamaged were synthesized in the

Delaney laboratory Sarah Delaney (Brown University). The 5'6-FAM primers were

annealed to complementary DNA templates with sequence context as previously

described [35,36] and are described below:

25/40 undamaged DNA substrate

`5'-/FAM/ TTT GCCT TGA CCA TGT AAC AGA GAG`

`                CGGA ACT GGT ACA TTG TCT CTC G̲CA CTC ACT CTC TTC TCT`

24/33 CPD damaged DNA substrate

`5'-/FAM/-TTT CTC CGG TAC TCC AGT GTA GGC`

`                GAG GCC ATG AGG TCA CAT CCG `**`TT`**`A GCT TGA GAA`

24/33 undamaged DNA substrate

`5'-/FAM/-TTT CTC CGG TAC TCC AGT GTA GGC`

`                GAG GCC ATG AGG TCA CAT CCG `**`TT`**`A GCT TGA GAA`

Confirmation of annealed substrates was determined 12% Native PAGE and samples

scanned on an RB Amersham Typhoon Fluorescent Imager (Cytiva) with a FAM filter.

Single oligodeoxynucleotides were purchased from IDT for MMEJ with internal consensus

sequence as previously described[37].

`5'-/FAM/GGTTAGCCCGGG`

Circular Dichroism and Melting Temperature

Secondary protein characteristics of hPol θ to zPol θ were determined on a J-815-CD Spectropolarimeter (Jasco, Brown University) with a 0.2 cm quartz cuvette at room temperature (20°C). For each sample, 3 μM of protein in 10 mM K_2HPO_4 buffer were scanned in triplicate from 190-280 nm. The thermal denaturation profile was determined by using the same instrument by heating the same sample from 20-90°C with a 5°C/min temperature rate increase at 222 nm using the same. Data were analyzed on Prism 10 GraphPad and the melting temperature (Tm) was estimated using the halfway point of the denaturing curve

Electrophoretic Mobility Shift Assay (EMSA)

The DNA binding affinity constant $K_{D(DNA)}$ was determined as previously described[34]. zPol θ was titrated from 0-1000nM against 10nM 25/40 dsDNA substrate in binding buffer and incubated for 1 hour at room temperature. Samples were separated on a 6% Native PAGE and scanned on an RB Typhoon scanner (Cytiva) with the FAM fluorescence filter. Separated bound and unbound products were quantified using ImageQuant. $K_{D(DNA)}$ was determined by equation 1.

$$Y = \left[\frac{(mx)}{(x+K_D)}\right] + b \hspace{3cm} \text{Equation 1}$$

Four replicates and two protein preparations were used to generate this data.

Rapid Chemical Quench Assay

Biphasic burst kinetics were measured as previously described[34]. Briefly,100 nM

Pol θ was pre-mixed with 300 nM 25/40 dsDNA substrate and rapidly mixed with 100 µM

of dCTP (correct nucleotide) with 10 mM MgCl$_2$ using an RQF-3 Rapid Chemical Quench

instrument (KinTek Corporation) at 37°C between 0.004-0.6 seconds. Reactions were

quenched by addition of 0.5M EDTA. Products were separated on a 15% Urea-denaturing

polyacrylamide gel and scanned using an Amersham Typhoon RB Fluorescent imager

(Cytiva). Extended product (n+1) was quantified using ImageQuant software and then

plotted to a full biphasic pre-steady state burst equation via non-linear regression using

Prism 9 GraphPad software (equation 2). A minimum of three replicates were included for

each assay on two independent protein preparations.

$[Product] = [E]_{app}[\frac{k_{obs}{}^2}{(k_{obs}+k_{ss})^2} \times (1 - e^{-(k_{obs}+k_{ss})t} + \frac{k_{obs}k_{ss}}{(k_{obs}+k_{ss})}t]$ Equation 2

Primer Extension Assays

Qualitative primer extension assays were performed as previously described[34].

Varying conditions were used to explore the primer extension capabilities between the

hPol θ and zPol θ. Under single-turnover conditions, excess Pol θ (200 nM) was pre-

incubated with 50 nM DNA and incubated for 5 minutes at 37°C. Nucleotide (125 µM) of

either none, all, individual nucleotides were preincubated with 20 mM *x* inorganic salt (x

= MgCl$_2$, CaCl$_2$, MnCl$_2$). Under Michaelis-Menton conditions, DNA (200nM) was in excess

to Pol θ (50 nM). All reactions were carried out in buffer containing 20 mM Tris HCl, pH

8.0, 25 mM KCl, 4% glycerol, 1 mM βME, and 80 µg/mL BSA. The reaction was initiated

by combining Pol-θ/DNA with dNTP/salt. Reactions were incubated at 37°C for 5 minutes

before being stopped by 80% Formamide/EDTA quench. Products were separated out on

a 15% urea-denaturing polyacrylamide gel, and scanned on a RB Amersham Typhoon

fluorescent imager (Cytiva).

MMEJ Assay

Microhomology mediated end-joining assay for both hPol θ and zPol θ were carried

out as previously described[37] on a 12-mer FAM labeled oligodeoxynucleotide. Pol θ (20

nM) was preincubated with 30 nM 5'-FAM ssDNA in reaction buffer (25mM Tris-HCl

pH8.8, 1mM βME, 0.01% NP-40, 0.1 mg/mL BSA, 10% glycerol, 10 mM $MgCl_2$, 30mM

NaCl) for 5 minutes at 37°C. Nucleotides (20 µM) were added and incubated at 37°C for

an additional 45 minutes. Reactions were terminated by addition of non-denaturing stop

buffer (100 mM Tris-HCl pH 7.5, 10 mg/mL proteinase K, 80 mM EDTA, and 0.5% SDS)

for an additional 15 minutes. DNA products were separated on a 12% native

polyacrylamide gel and scanned by an RB fluorescent Amersham Typhoon (Cytiva) with

a FAM filter.

RESULTS

Zebrafish and human polymerase domains display high degree of structural similarity

To determine the degree of similarity between the zPol θ and hPol θ PD primary

amino acid sequences were aligned (Table 1, Supplemental figure 1). The alignment of

the full-length Pol θ protein indicates, zebrafish and human Pol θ share 46% identity. This

degree of similarity increases when comparing the predicted polymerase domain (63%),

as well as subdomains containing catalytic activity, fingers (75.3%), thumb (74.6%), and

palm (66.7%). These data suggest a structurally similar molecule.

To assess the extent of structural similarity we generated a predicted structure for

zPol θ PD using ColabFold[26] to compare to the solved crystal structure of hPol θ PD[31].

Upon visual inspection the predicted zPol θ PD displays classical DNA polymerase PD

structures (Figure 1). The three major subunits, the fingers, thumb, and palm are visible,

and when modeled in, a DNA molecule can fit in the presumed catalytic domain. The

model also indicates the presence of unstructured loop domains, that have functional

importance[12], that were not resolved in the hPol θ PD structure. An over lay of the hPol

θ PD and zPol θ PD show that the structures have a high degree of similarity. As predicted

by the amino-acid alignment, these data indicate that much of the structure of the hPol θ

PD is conserved in zPol θ.

Table 1. Sequence alignment analysis

| | Polymerase theta comparisons | | |
	zebrafish	% identity	human
Polymerase Domain	thumb (122 aa)	74.6%	thumb (125 aa)
	fingers (142 aa)	75.3%	fingers (142 aa)
	palm (163 aa)	66.7%	palm (177 aa)
	total (744 aa)	63%	total (799 aa)
Full length	2576 aa	46%	2590 aa

Figure 1. Structural modeling of zPol θ. (A) AlphaFold rendering of zPol θ with subdomains colored, thumb – blue, fingers

– red, palm – green, and exo-nuclease in yellow. DNA is colored in light blue. (B) FATCAT overlay of AlphaFold rending of zPol θ

(bronze) and hPol θ (navy) crystal structure[31] without loop inserts, DNA is colored in light blue.

hPol θ and zPol θ are structural similar

The plasmid containing the c-terminal recombinant zPol θ was expressed and

purified in the same way as hPol θ[34] and as summarized in the Materials and Methods.

Similar to hPol θ, one protein preparation yields approximately 5-10 μM and we observed

similar expression and purification levels as seen with hPol θ (Figure 2A)

To confirm similarity in secondary structure between hPol θ and zPol θ, circular

dichroism spectroscopy (CD) was performed at 20°C. The same sample was heated from

20-90°C in order to determine the thermal denaturation profile. Both spectra were

overlayed and indicated minimal variance suggesting that both hPol θ and zPol θ have

similar secondary characteristics and thermal stability with a Tm of about 55°C.

Figure 2. hPol θ and zPol θ display similar secondary characteristics and thermal stability. (A) Expression and purification of

zPol θ was the same as hPol θ as described in the Materials and Methods. For each sample, approximately 56-60 pmol of cleaved,

purified protein were loaded on a 10% SDS PAGE and Coomassie stained. Both hPol θ and zPol θ migrate to approximately 90 kDa

as expected. (B) Circular dichroism spectra of 3 µM hPol θ (solid line) and zPol θ (dashed) proteins in 10 mM Potassium Phosphate

buffer. Samples were scanned from 190 to 280 nm. (C)The same samples were heated from 20-90°C and ellipticity measured at 222

nm.

zPol θ binds to dsDNA substrate

DNA binding by a DNA polymerase is one of the first steps in its catalytic

mechanism. To determine the DNA binding capabilities of zPol θ, we titrated zPol θ from

0-1000 nM protein against 10 nM 25/40 dsDNA. Complexed DNA/protein products were

separated on a denaturing gel to determine a dissociation constant ($K_{D(DNA)}$) for DNA

binding. Similar to hPol θ, zPol θ has a low $K_{D(DNA)}$ value of approximately 19.8 ± 3.1 nM.

Figure 3. zPol θ binds tightly to ds DNA. zPol θ was titrated from 0-1000 nM against 10 nM 25/40 dsDNA. Bound and unbound

products were separated on a 6% non-denaturing gel and quantified using ImageQuant. $K_{D(DNA)}$ was mathematically calculated using

Equation 1 and is the midpoint between bound and unbound fractions.

zPol θ can extend dsDNA similar to hPol θ

The second step in the DNA polymerase catalytic pathway is nucleotide binding

and formation of the phosphodiester bond. To explore this fundamental step of DNA

Polymerase activity, we assayed zPol θ's ability to extend 25/40 dsDNA under varying

conditions. Under standard steady-state conditions, 200 nM of zPol θ or hPol θ was pre-

incubated with 50 nM 25/40 dsDNA. The reaction was initiated by the addition of 125 nM

dNTP as described in Figure 4 along with 20 mM $MgCl_2$, the preferred metal for DNA

polymerase [38]. We observed under these conditions both hPol θ and zPol θ were able to

extend the full 18-mer template with all nucleotides present (Figure 4). Both enzymes

were able to incorporate single nucleotides, correct and incorrect as well. Notably, zPol θ

was able to incorporate incorrect dGTP to full extension (n+1) compared to only n+6 with

hPol θ. DNA polymerases can utilize other metals including Mn^{2+}, and we observed an

increase in mutagenesis through misincorporation for both hPol θ and zPol θ. When

provided with all dNTP, zPol θ can extend past the end of the template (n+18). Overall,

zPol θ experiences more extension products especially with incorrect nucleotides dATP,

dGTP, and dTTP compared with hPol θ under similar conditions. Steady-state conditions

highlight overall DNA polymerase activity, but because the dsDNA substrate is in excess,

activity highlights multiple turnovers[39]. Although DNA pol θ has been shown to have

robust de novo activity with manganese[40], we wanted to be sure this over extension

observed with zPol θ and Mn^{2+} was the result of extension and not an artifact. We

changed the ratio of protein to DNA to reflect single-turnover conditions; excess protein

over dsDNA substrate. Here we are able to observe polymerization events for

theoretically every available DNA substrate. Similar to steady-state conditions, we

observe an even more robust de novo extension with not only all nucleotides, but also

with dATP, suggesting that zPol θ misincorporation with dATP is preferred (Supplement

2).

5'-/FAM/ TTT GCCT TGA CCA TGT AAC AGA GAG
CGGA ACT GGT ACA TTG TCT CTC GCA CTC ACT CTC TTC TCT

Figure 4. zPol θ experiences greater nucleotide extension activity compared to hPol θ. Under steady-state conditions 50 nM hPol θ or zPol θ were preincubated with 200 nM 25/40 dsDNA and combined with either 10 mM $MgCl_2$ or $MnCl_2$ for 5 minutes and 37°C. DNA extension products were separated on a denaturing gel and visualized on a Typhoon scanner. Each n+1 band represents an extension of one nucleotide following the DNA template as described above. N+1 would represent either correct nucleotide incorporation of dCTP opposite a templating G (underlined) or a misincorporation event of dATP, dGTP, or dTTP opposite templating G. Each subsequent band is another nucleotide extension with a maximum of template-dependent extension of n+18. Bands migrating higher than n+18 represent de novo synthesis.

zPol θ catalytic activity similar to other DNA polymerases

To further explore the mechanism of nucleotide incorporation of zebrafish Pol θ, we assayed zPol θ under presteady-state conditions in which there is an excess DNA substrate to enzyme with correct nucleotide. This assay focuses on the DNA polymerase

ability to extend DNA by incorporating the correct nucleotide opposite a templating base.

This activity is biphasic in which there is a rapid polymerization step of nucleotide

incorporation at the DNA primer's 3'OH and a slower, rate limiting step of product

release[39]. If biphasic activity is not observed, it suggests a step before nucleotide

incorporation is the rate-limiting step[41]. To ensure that purified c-terminal zPol θ follows

the traditional DNA polymerase mechanism, 100 nM zPol θ was preincubated with 300

nM 25/40 dsDNA. The DNA/Pol θ complex was rapidly combined with 100 μM correct

nucleotide and 10 mM $MgCl_2$ from 0.004-0.6 seconds. DNA products were separated on

a denaturing polyacrylamide gel and primer extension of n+1 was quantified and data fit

to a full biphasic burst equation. zPol θ fit to a biphasic equation with an observable

polymerization rate (k_{obs}) of 15.9 ± 2.5 s^{-1} (Figure 5).

Figure 5. zPol θ experiences biphasic burst activity. Zebrafish Pol θ (100 nM) was preincubated with 300 nM 25/40 dsDNA. The

DNA/Pol θ complex was rapidly combined with 100 μM dCTP (correct nucleotide) and 10 mM $MgCl_2$. Reactions were carried out at

37°C and quenched with 0.5 M EDTA. Products were separated on a denaturing gel and quantified with ImageQuant software. Data

were fit to a biphasic burst equation to obtain observed k_{obs} rates 15.9 ± s^{-1}. The slower rate k_{ss} was calculated to be 3.4 ± 0.46 s^{-1}.

zPol θ performs MMEJ activity

One of the major functions of DNA Pol θ is its ability to repair double-strand breaks and is the primary DNA polymerase for microhomology-mediated end joining. In doing so, Pol θ utilizes internal homology within the DNA sequence to act as a template. Pol θ aligns these complementary pieces and extends in the 5' to 3' direction[37,42,43]. Truncated hPol θ has been shown to able to perform MMEJ activity on short 12-mer single-stranded DNA, but the full 290 kDa Pol θ with the N-terminal helicase and central domains are needed to anneal and extend larger segments of DNA[37]. We wanted ensure that zPol θ could also perform MMEJ in a similar manner to hPol θ on short fragments of DNA. Figure 6 is a representative gel of hPol θ and zPol θ performing MMEJ on a ssDNA. As indicated in the schematic above, the CCCGGG are aligned through Pol θ in the presence of (+) dNTP and subsequently extended in the opposite direction giving rise to a slower moving double-stranded DNA product. Both hPol θ and zPol θ are able to perform this activity. We hypothesize the smaller product bands are indicative of classic snap-back synthesis in which the DNA substrate anneals onto itself for Pol θ to extend. This behavior has been observed by others on hPol θ and there is little variation between the two species[37].

Figure 6. zPol θ is able to perform MMEJ activity of short DNA fragments. Pol θ (20 nM) was preincubated with 30 nM 5'-FAM ssDNA in reaction buffer. All nucleotides (+ dNTP) were added and the ternary complex was incubated for 45 minutes at 37°C. A no dNTP (-dNTP) control was carried out in the same manner. Reactions were stopped and products separated on a 12% Native PAGE. The gel was visualized on a Typhoon scanner.

zPol θ is able to bypass CPD lesion DNA.

Pol θ is a versatile DNA polymerase in not only can it perform MMEJ, it has also been shown to bypass cyclobutane pyrimidine dimers (CPD)[36]. By being able to extend

a DNA primer passed a template containing a contorted Thymine-Thymine lesion, human

and mouse Pol θ have been demonstrated to be critical in suppressing DNA damage and

preventing skin lesions. On a molecular level, human Pol θ has demonstrated that not

only can it insert opposite the initial T in the T-T dimer but is able to mutagenically extend

past this lesion for the remaining DNA template. We hypothesized that zPol θ has the

same ability to bypass CPD lesions and we assayed both Pol θ under single-turnover

conditions (4:1 protein to DNA) in the presence of a 24/33 CPD damaged DNA template

with both Mg^{2+} and Mn^{2+}. As predicted, there was little variance in bypass activity of zPol

θ compared to hPol θ. Both enzymes were able to readily insert opposite a T-T dimer as

well as extend past this lesion with both all dNTPs present and dATP and to some extent

dGTP. Both Pol θs could not incorporate dCTP opposite T-T, but we observed only

insertion of dTTP opposite and no extension. In the presence of Mn^{2+}, both Pol θs readily

bypassed T-T dimers, again demonstrating de novo synthesis past the template. zPol θ

was more robust in extension with the other incorrect nucleotides suggesting Mn^{2+} has

an increased mutagenic effect.

5'-/FAM/-TTT CTC CGG TAC TCC AGT GTA GGC
            GAG GCC ATG AGG TCA CAT CCG **TT**A GCT TGA GAA

Figure 7. zPol θ is able to bypass CPD lesions. As described in the Materials and Methods, 200 nM Pol θ (human or zebrafish) was

preincubated with 50 nM CPD Damaged DNA substrate. Reactions were initiated by the addition of 125 nM nucleotides as described

and either 10 mM $MgCl_2$ or $MnCl_2$. Reactions were carried out at 37°C for 5 minutes and products visualized on a 12% denaturing

gel. Higher migrating products are indicative of full extension (n+12) with smeared bands representing de novo synthesis with

extension past n+12.

zPol θ experiences unusual extension of DNA substrates in the presence of Ca^{2+}.

To explore the role of divalent metals in DNA polymerase activity for Pol θ, we

performed a DNA polymerase extension assay again with either these assays with their

specific DNA substrates swapping out the active metals for Ca^{2+} which has traditionally

used as an inert control. Unlike the other divalent metals, Ca^{2+} allows for ternary complex

formation, but extension is limited or slow[44,45]. Using 50 nm if the 24/33 undamaged and

CPD damaged DNA substrate, we performed a primer extension assay with 200 nM hPol

θ or z WT Pol θ with Mg^{2+} substituted for $CaCl_2$. Extension products were separated on

a denaturing polyacrylamide gel and quantified based on the percent extension. Figure

8A is a representative gel of extension on 24/33 undamaged DNA template. We observe

that hPol θ could incorporate every nucleotide to some extent, with an n+3 extension

product only observed in the presence of all dNTP or purines. zPol θ was observed to

generate full extension product (94%) on this DNA template (n+12) with all nucleotides

present and, like hPol θ, could extend with purines as well. Incorporation of dATP led to

92% conversion to product although the enzyme stalled around n+2. Interestingly, zPol

θ appears to skip the first thymine in the undamaged sequence for both all nucleotides

and dATP. The same experiment was carried out with 24/33 CPD damaged DNA. Under

these conditions we report that Ca^{2+} reduced DNA polymerase activity for both hPol θ and

zPol θ with incorporation of only one nucleotide irrespective if that nucleotide was

matched or mismatched with the templating base.

Figure 8. DNA Pol θ is able incorporate and extend dsDNA in the presence of Ca²⁺. Under single turnover conditions 200 nM Pol

θ (human or zebrafish) was preincubated with 50 nM 24/33 undamaged (A) or damaged (B) DNA substrate and reacted with 125 nM

nucleotides in 10 mM $CaCl_2$. Reactions were carried out at 37°C for 5 minutes and products visualized on a 12% denaturing gel.

Percent extended was calculated using ImageQuant software by quantifying the intensity of the extended products (n+1 and higher)

divided by the intensity of the total amount of DNA.

DISCUSSION

Zebrafish and human Pol θ structures have a high degree of similarity

Comparisons of the amino acid sequence (table 1) and the structures (Figure 1) of

zPol θ and hPol θ reveal that the two proteins share a high degree of similarity.

Importantly, and perhaps not surprisingly, the areas of greatest similarity are around the

catalytic subdomains, the fingers, thumb, and palm, of the polymerase domain. These

sites of activity would be important to the protein function across evolutionary time.

Interestingly, unlike other A-type polymerase family members Pol θ has three loop

structures within the PD which have been identified to be important for function[12].

Comparison of the zPol θ and hPol θ sequences indicate that zPol θ also contains these

inserted loops however, they contain little homology (Supplement 1) to that observed

between human and mouse[31] Pol θ. Despite this difference, expression and purification

yields of zPol θ were similar to that of hPol θ as were the secondary structural

characteristics and thermal stability (Figure 2); an early indication similar protein folding.

We also show here that zPol θ still retains the same activity observed in hPol θ.

Zebrafish Pol θ extends dsDNA

DNA polymerase θ primary role in the human cell is the primary DNA polymerase

repair enzyme in microhomology-mediated end joining and thus, the major goal of this

study was to query if zPol θ retained a similar function. Initially we simplify the activity by

asking 'can zPol θ bind to a primer/template dsDNA substrate and then extend it?' Our

data suggests that yes it can. zPol θ binds tightly to this substrate (Figure 3)[34] similar to

values obtained with hPol θ. This is expected because loop 1 which is located in the

thumb domain or DNA binding domain is thought to be involved with contacts to DNA[12]

and is the only conserved loop region between zPol θ and hPol θ. zPol θ can in fact

extend a DNA substrate, and we show it has robust activity on this particular DNA

substrate (Figure 4) especially in the presence of Mn^{2+}. While most of the data presented

in this work was qualitative, probing how fast a DNA polymerase makes a phosphodiester

bound through biochemical kinetics can provide insight into mechanism of

incorporation[39]. Like most DNA polymerases, zPol θ performs biphasic burst kinetics

which is indicative of a two-step mechanism with a rate limiting step of product release.

zPol θ experiences an observed polymerization rate of around 16 s^{-1} (Figure 5) which is

almost 4 times slower than its human ortholog[34], but not uncommonly slow as a similar

DNA Polymerase β experiences a similar rate[46–48]. Why zPol θ might experience a slower

rate is unknown. It could be due to the lack of conservation within the looping structures

in the palm domain which in human Pol θ may drive substrate alignment for rapid

polymerization.

Zebrafish Pol θ is able to perform microhomology-mediated End Joining

474   Despite being able to extend DNA, it is important that zPol θ also be able to perform

MMEJ as in humans it is its primary function. Although a majority of human Pol θ's N-

terminal and central domains are critical for this function, studies have shown that the c-

terminal polymerase domain of Pol θ does retain limited function for aligning and

extending short single-stranded DNA[37]. zPol θ was no exception (Figure 6) and was able

to complement two single-strands and extend which is perhaps the most compelling

evidence of homologous function.

zPol θ can bypass CPD lesions

483   Another function of human Pol θ is its ability to bypass DNA damage and a more

recent study in mice suggest that bypassing UV damage is critical in the prevention of

skin cancer[36]. Our data provides evidence that zPol θ is able to bypass and extend CPD

lesions *in vivo* similar to that of hPol θ (Figure 7). Translesion bypass activity has been

highlighted as a function of loops 2 and 3 in human Pol θ[12,31]. Surprisingly, zebrafish

display very little homology through similar inserts (Supplement 1). However, our studies

might suggest that the only critical residues for this function are isolated to the c-terminal

end of this insert beginning with the sequence GMXFSXSMR. Further studies exploring

this insert in zebrafish are needed to determine if these conserved amino acids are truly

critical and that the function is either dictated through the overall presence of the loop or

that the loop dependent activities require the few conserved amino acids retain in zPol θ.

zPol θ experiences Ca^{2+} dependent polymerization

Our data shows that zPol θ retains all of the DNA polymerase activities of hPol θ,

with the exception being the robust Ca^{2+} dependency during DNA extension (Figure 8).

While it is unusual to see DNA polymerases extend past the initial insertion event, there

have been instances where high-fidelity *Sulfolobus solfataricus* Dpo4 polymerase uses

calcium [49]. Whether or not swapping Ca^{2+} in zebrafish makes it a faster or mutagenic

polymerase has not been explored but evolutionarily it is possible that Ca^{2+} makes for a

stable ion swap for structural alignment and catalytic activity. Given that in freshwater

contains nearly identical concentrations of the divalent ions[50], its plausible that Ca^{2+} could

readily be a co-factor substitute.

Our data presented is clear evidence that zebrafish Pol θ is a homolog to human

Pol θ and that structurally and enzymatically behave with similar functions. This study is

significant as it highlights the availability of zebrafish as a model organism for studying

Pol θ and its potential function in DNA repair and disease. In particular given the robust

assortment of tools zebrafish offers a powerful, functionally relevant model for human

melanoma. Future studies could introduce patient derived mutations by way of germline

alterations and study the effects over the life time of the animal. Thus, adding new insights

into potential disease markers and mechanisms of disease progression and treatment.

ACKNOWLEDGEMENTS

We would like to thank Sarah Delaney and Mary Tarantino from Brown University for the

generation of the CPD damaged DNA. Thank you to Sylvie Doublié from University of

Vermont for the human pol θ plasmid.

FUNDING

Research reported in this publication was supported by the Rhode Island Institutional

Development Award (IDeA) Network of Biomedical Research Excellence under

P20GM103430 and in part by the National Institute of General Medical Sciences of the

National Institutes of Health under grant number R15GM144903-01.

The content is solely the responsibility of the authors and does not necessarily represent

the official views of the National Institutes of Health.

WORK CITED

1. Arnold, M. *et al.* Global Burden of Cutaneous Melanoma in 2020 and Projections to 2040. *JAMA Dermatol* **158**, 495–503 (2022).

2. Mort, R. L., Jackson, I. J. & Patton, E. E. The melanocyte lineage in development and disease. *Development* **142**, 620–632 (2015).

3. Jilani, S. *et al.* CAR-T cell therapy targeting surface expression of TYRP1 to treat cutaneous and rare melanoma subtypes. *Nat Commun* **15**, 1244 (2024).

4. Larkin, J. *et al.* Combined Nivolumab and Ipilimumab or Monotherapy in Previously Untreated Melanoma. *N Engl J Med* **373**, 23–34 (2015).

5. Farshidfar, F. *et al.* Integrative molecular and clinical profiling of acral melanoma links focal amplification of 22q11.21 to metastasis. *Nat Commun* **13**, 898 (2022).

6. Thyme, S. B. & Schier, A. F. Polq-mediated end joining is essential for surviving DNA double-strand breaks during early zebrafish development. *Cell Rep* **15**, 707–714 (2016).

7. Brambati, A., Barry, R. M. & Sfeir, A. DNA polymerase theta (Polθ) – an error-prone polymerase necessary for genome stability. *Current Opinion in Genetics & Development* **60**, 119–126 (2020).

8. Yoon, J.-H. *et al.* Error-Prone Replication through UV Lesions by DNA Polymerase θ Protects against Skin Cancers. *Cell* **176**, 1295-1309.e15 (2019).

9. Chan, S. H., Yu, A. M. & McVey, M. Dual Roles for DNA Polymerase Theta in Alternative End-Joining Repair of Double-Strand Breaks in Drosophila. *PLOS Genetics* **6**, e1001005 (2010).

10. Kent, T., Chandramouly, G., McDevitt, S. M., Ozdemir, A. Y. & Pomerantz, R. T.

Mechanism of microhomology-mediated end-joining promoted by human DNA

polymerase θ. *Nat Struct Mol Biol* **22**, 230–237 (2015).

11. Yousefzadeh, M. J. *et al.* Mechanism of Suppression of Chromosomal Instability by

DNA Polymerase POLQ. *PLOS Genetics* **10**, e1004654 (2014).

12. Hogg, M., Seki, M., Wood, R. D., Doublié, S. & Wallace, S. S. Lesion Bypass Activity

of DNA Polymerase θ (POLQ) Is an Intrinsic Property of the Pol Domain and

Depends on Unique Sequence Inserts. *Journal of Molecular Biology* **405**, 642–652

(2011).

13. Yu, A. M. & McVey, M. Synthesis-dependent microhomology-mediated end joining

accounts for multiple types of repair junctions. *Nucleic Acids Res* **38**, 5706–5717

(2010).

14. Ceccaldi, R. *et al.* Homologous-recombination-deficient tumours are dependent on

Polθ-mediated repair. *Nature* **518**, 258–262 (2015).

15. Goff, J. P. *et al.* Lack of DNA Polymerase θ (POLQ) Radiosensitizes Bone Marrow

Stromal Cells In Vitro and Increases Reticulocyte Micronuclei after Total-Body

Irradiation. *Radiat Res* **172**, 165–174 (2009).

16. Goullet de Rugy, T. *et al.* Excess Polθ functions in response to replicative stress in

homologous recombination-proficient cancer cells. *Biology Open* **5**, 1485–1492

(2016).

17. Allera-Moreau, C. *et al.* DNA replication stress response involving PLK1, CDC6,

POLQ, RAD51 and CLASPIN upregulation prognoses the outcome of early/mid-

stage non-small cell lung cancer patients. *Oncogenesis* **1**, e30–e30 (2012).

18. Lemée, F. *et al.* DNA polymerase θ up-regulation is associated with poor survival in

breast cancer, perturbs DNA replication, and promotes genetic instability.

Proceedings of the National Academy of Sciences **107**, 13390–13395 (2010).

19. Seki, M. *et al.* High-efficiency bypass of DNA damage by human DNA polymerase

Q. *The EMBO Journal* **23**, 4484–4494 (2004).

20. Thomas, C. *et al.* Melanoma-Derived DNA Polymerase Theta Variants Exhibit

Altered DNA Polymerase Activity. *Biochemistry* **63**, 1107–1117 (2024).

21. Patton, E. E. *et al.* BRAF mutations are sufficient to promote nevi formation and

cooperate with p53 in the genesis of melanoma. *Curr Biol* **15**, 249–254 (2005).

22. Howe, K. *et al.* The zebrafish reference genome sequence and its relationship to the

human genome. *Nature* **496**, 498–503 (2013).

23. Groenewoud, A. *et al.* Patient-derived zebrafish xenografts of uveal melanoma

reveal ferroptosis as a drug target. *Cell Death Discov.* **9**, 1–11 (2023).

24. Fazio, M., Ablain, J., Chuan, Y., Langenau, D. M. & Zon, L. I. Zebrafish patient

avatars in cancer biology and precision cancer therapy. *Nat Rev Cancer* **20**, 263–

273 (2020).

25. Madeira, F. *et al.* The EMBL-EBI Job Dispatcher sequence analysis tools framework

in 2024. *Nucleic Acids Res* **52**, W521–W525 (2024).

26. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods*

19, 679–682 (2022).

27. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

(2021) doi:10.1038/s41586-021-03819-2.

28. Mirdita, M., Steinegger, M. & S"oding, J. MMseqs2 desktop and local web server

app for fast, interactive sequence searches. *Bioinformatics* **35**, 2856–2858 (2019).

29. Mirdita, M. *et al.* Uniclust databases of clustered and deeply annotated protein

sequences and alignments. *Nucleic Acids Res.* **45**, D170–D176 (2017).

30. Mitchell, A. L. *et al.* MGnify: the microbiome analysis resource in 2020. *Nucleic Acids*

Res. (2019) doi:10.1093/nar/gkz1035.

31. Zahn, K. E., Averill, A. M., Aller, P., Wood, R. D. & Doublié, S. Human DNA

polymerase θ grasps the primer terminus to mediate DNA repair. *Nat Struct Mol Biol*

22, 304–311 (2015).

32. Li, Z., Jaroszewski, L., Iyer, M., Sedova, M. & Godzik, A. FATCAT 2.0: towards a

better understanding of the structural diversity of proteins. *Nucleic Acids Res* **48**,

W60–W64 (2020).

33. Bittrich, S., Segura, J., Duarte, J. M., Burley, S. K. & Rose, Y. RCSB protein Data

Bank: exploring protein 3D similarities via comprehensive structural alignments.

Bioinformatics **40**, btae370 (2024).

34. Thomas, C. *et al.* Melanoma-Derived DNA Polymerase Theta Variants Exhibit

Altered DNA Polymerase Activity. *Biochemistry* **63**, 1107–1117 (2024).

35. Prasad, R. *et al.* Human DNA polymerase θ possesses 5′-dRP lyase activity and

functions in single-nucleotide base excision repair in vitro. *Nucleic Acids Research*

37, 1868–1877 (2009).

36. Yoon, J. H. *et al.* Error-Prone Replication through UV Lesions by DNA Polymerase θ

Protects against Skin Cancers. *Cell* **176**, 1295-1309.e15 (2019).

37. Black, S. J. *et al.* Molecular basis of microhomology-mediated end-joining by purified full-length Polθ. *Nature Communications* **10**, (2019).

38. Steitz, T. A. DNA Polymerases: Structural Diversity and Common Mechanisms *. *Journal of Biological Chemistry* **274**, 17395–17398 (1999).

39. Johnson, K. A. Transient-State Kinetic Analysis of Enzyme Reaction Pathways. *Enzymes* **20**, 1–61 (1992).

40. Kent, T., Mateos-Gomez, P. A., Sfeir, A. & Pomerantz, R. T. Polymerase θ is a robust terminal transferase that oscillates between three different mechanisms during end-joining. *eLife* **5**, e13740.

41. Joyce, C. M. Techniques used to study the DNA polymerase reaction pathway. *Biochimica et biophysica acta* **1804**, 1032–1040 (2010).

42. Brambati, A., Barry, R. M. & Sfeir, A. DNA polymerase theta (Polθ) – an error-prone polymerase necessary for genome stability. *Current Opinion in Genetics and Development* **60**, 119–126 (2020).

43. Li, C. *et al.* Structural basis of DNA polymerase θ mediated DNA end joining. *Nucleic Acids Research* **51**, 463–474 (2023).

44. Freudenthal, B. D., Beard, W. A., Shock, D. D. & Wilson, S. H. Observing a DNA Polymerase Choose Right from Wrong. *Cell* **154**, 157–168 (2013).

45. Nakamura, T., Zhao, Y., Yamagata, Y., Hua, Y. J. & Yang, W. Watching DNA polymerase eta make a phosphodiester bond. *Nature* **487**, 196–201 (2012).

46. Towle-Weicksel, J. B. *et al.* Fluorescence resonance energy transfer studies of DNA polymerase β the critical role of fingers domain movements and a novel non-

covalent step During nucleotide selection. *Journal of Biological Chemistry* **289**,

16541–16550 (2014).

47. Werneburg, B. G. *et al.* DNA Polymerase β: Pre-Steady-State Kinetic Analysis and

Roles of Arginine-283 in Catalysis and Fidelity†. *Biochemistry* **35**, 7041–7050

(1996).

48. Murphy, D. L., Kosa, J., Jaeger, J. & Sweasy, J. B. The Asp285 Variant of DNA

Polymerase Beta Extends Mispaired Primer Termini via Increased Nucleotide

Binding. *Biochemistry* **47**, 8048–8057 (2008).

49. Ralec, C., Henry, E., Lemor, M., Killelea, T. & Henneke, G. Calcium-driven DNA

synthesis by a high-fidelity DNA polymerase. *Nucleic Acids Res* **45**, 12425–12440

(2017).

50. Guerin, A. J. *et al.* High-resolution freshwater dissolved calcium and pH data layers

for Canada and the United States. *Sci Data* **11**, 370 (2024).

651