Target journal: Frontiers in Endocrinology

Special issue: Oxytocin and Metabolic Dysregulation: From Pathophysiology to

Pharmacotherapy

Full title: Effects of systemic oxytocin and beta-3 receptor agonist (CL 316243) treatment

on body weight and adiposity in male diet-induced obese rats

Abbreviated title: Systemic Oxytocin and CL 316243 Treatment Reduces Weight

Jared D. Slattery[1], June R. Rambousek[1], Edison Tsui[1], Mackenzie K. Honeycutt[1], Matvey

Goldberg[1], James L. Graham[5], Tomasz A. Wietecha[2,4], Tami Wolden-Hanson[1], Kevin D.

O'Brien[3,4], Peter J. Havel[5,6], and James E. Blevins[1,2]

[1]VA Puget Sound Health Care System, Office of Research and Development Medical

Research Service, Department of Veterans Affairs Medical Center, Seattle, WA 98108,

USA

[2]Division of Metabolism, Endocrinology and Nutrition, Department of Medicine, University

of Washington School of Medicine, Seattle, WA 98195, USA

[3]Division of Cardiology, Department of Medicine, University of Washington School of

Medicine, Seattle, WA 98195, USA

[4]UW Medicine Diabetes Institute, University of Washington School of Medicine, Seattle,

WA 98109

[5]Department of Nutrition, University of California, Davis, CA 95616, USA

[6]Department of Molecular Biosciences, School of Veterinary Medicine, University of

California, Davis, CA 95616, USA

Corresponding author's address:

J. E. Blevins, Ph.D.

VA Puget Sound Health Care System

Research-151

1660 South Columbian Way

Seattle, WA 98108

Key words: Obesity, brown adipose tissue, white adipose tissue, food intake, oxytocin

This material was based upon work supported by the Office of Research and
Development, Medical Research Service, Department of Veterans Affairs (VA) and the
VA Puget Sound Health Care System Rodent Metabolic Phenotyping Core and the
Cellular and Molecular Imaging Core of the Diabetes Research Center at the University
of Washington and supported by National Institutes of Health (NIH) grant P30DK017047.
This work was also supported by the VA Merit Review Award
5 I01BX004102, from the United States (U.S.) Department of Veterans Affairs Biomedical
Laboratory Research and Development Service and NIH 5R01DK115976 grant to James
Blevins. PJH's research program also received research support during the project period
from NIH grants DK-095980, HL-091333, HL-107256 and a Multi-campus grant from the
University of California Office of the President.
45
46   The contents do not represent the views of the U.S. Department of Veterans Affairs or
47   the United States Government.
48
49   **ACKNOWLEDGMENTS**
50   The authors thank the technical support of Nishi Ivanov.
51

52   Disclosures: JEB had a financial interest in OXT Therapeutics, Inc., a company that
53   developed highly specific and stable analogs of oxytocin to treat obesity and metabolic
54   disease but this is no longer the case. The authors' interests were reviewed and were
55   managed by their local institutions in accordance with their conflict of interest policies.
56   The other authors have nothing to report.
57

Abstract

Previous studies have implicated hindbrain oxytocin (OT) receptors in the control of food intake and brown adipose tissue (BAT) thermogenesis. We recently demonstrated that hindbrain [fourth ventricle (4V)] administration of oxytocin (OT) could be used as an adjunct to drugs that directly target beta-3 adrenergic receptors (β3-AR) to elicit weight loss in diet-induced obese (DIO) rodents. What remains unclear is whether systemic OT can be used as an adjunct with the β3-AR agonist, CL 316243, to increase BAT thermogenesis and elicit weight loss in DIO rats. We hypothesized that systemic OT and β3-AR agonist (CL 316243) treatment would produce an additive effect to reduce body weight and adiposity in DIO rats by decreasing food intake and stimulating BAT thermogenesis. To test this hypothesis, we determined the effects of systemic (subcutaneous) infusions of OT (50 nmol/day) or vehicle (VEH) when combined with daily systemic (intraperitoneal) injections of CL 316243 (0.5 mg/kg) or VEH on body weight, adiposity, food intake and brown adipose tissue temperature (T_{IBAT}). OT and CL 316243 monotherapy decreased body weight by $8.0\pm0.9\%$ ($P<0.05$) and $8.6\pm0.6\%$ ($P<0.05$), respectively, but OT in combination with CL 316243 produced more substantial weight loss ($14.9\pm1.0\%$; $P<0.05$) compared to either treatment alone. These effects were associated with decreased adiposity, energy intake and elevated T_{IBAT} during the treatment period. The findings from the current study suggest that the effects of systemic OT and CL 316243 to elicit weight loss are additive and appear to be driven primarily by OT-elicited changes in food intake and CL 316243-elicited increases in BAT thermogenesis.

Introduction

The obesity epidemic and its associated complications, increase the risk for cardiovascular disease (CVD), hypertension, cancer, type 2 diabetes, and COVID-19 [1; 2]. Many of the monotherapies to treat obesity are of limited effectiveness, associated with adverse and/or unwanted side effects (i.e. diarrhea, nausea, vomiting, sleep disturbance and depression) and/or are poorly tolerated. Improvements have been made in monotherapies to treat obesity, particularly within the family of drugs that target the glucagon-like peptide-1 receptor (GLP-1R). While the FDA recently approved the use of the long-acting and highly effective GLP-1R agonist, semaglutide [3], it can also be associated with mild to moderate gastrointestinal (GI) side effects [3; 4], thus highlighting the need for continued optimization of existing treatments.

Recent studies suggest that combination therapy (co-administration of different compounds) and monomeric therapy (dual or triple agonists in single molecule) are more effective than monotherapy for prolonged weight loss [5; 6]. Marked weight loss has been reported in long-term (20 weeks to ≥ 1 year) clinical studies in humans treated with the amylin analogue, cagrilintide, and semaglutide (≈ 15.6 to 17.1% of initial body weight [7; 8]) and the FDA-approved drug, Qsymia (topiramate + phentermine) (≈ 10.9% of initial body weight; [9]). Alternatively, the monomeric compound, tirzepatide (Zepbound^{TM}), targets both GLP-1R and glucose-dependent insulinotropic polypeptide (GIP), and was recently reported to elicit 20.9% and 25.3% weight loss in humans with obesity over 72- [10] and 88-week trials [11]. In addition, retatrutide, a triple-agonist that targets GIP, glucagon receptors (GCGR) and GLP-1R was reported to reduce body weight by 24.2% over a 48-week trial [12]. Despite the considerable improvements that have been made

with respect to weight loss, these treatments are still associated with adverse GI side

effects [10], leading, in some cases, to the discontinuation of the drug in up to 7.1% of

participants [10].

While the hypothalamic neuropeptide, oxytocin (OT) is largely associated with

reproductive behavior [13], recent studies implicate an important role for OT in the

regulation of body weight [14; 15; 16; 17]. Studies to date indicate that OT elicits weight

loss, in part, by reducing food intake and increasing lipolysis [18; 19; 20] and energy

expenditure [18; 21; 22; 23]. While OT is effective at evoking prolonged weight loss in

DIO rodents [19; 22; 23; 24; 25; 26; 27; 28] and nonhuman primates [18], its overall

effectiveness as a monotherapy to treat obesity is relatively modest following 4-8 week

treatments in DIO mice ($\approx$4.9%) [28], rats ($\approx$8.7%) [28] and rhesus monkeys ($\approx$3.3%) [18]

thus making it more suited as a combination therapy with other drugs that work through

other mechanisms. Head and colleagues recently reported that systemic OT and the

opioid antagonist, naltrexone, resulted in an enhanced reduction of high-fat, high-sugar

meal in rats [29]. Recently, we found that hindbrain (fourth ventricle; 4V) OT treatment in

combination with systemic treatment with CL-316243, a drug that directly targets beta-3

adrenergic receptors (β3-AR) to increase BAT thermogenesis [28; 30; 31; 32; 33],

resulted in greater weight loss (15.5 $\pm$ 1.2% weight loss) than either OT (7.8 $\pm$ 1.3% weight

loss) or CL 316243 (9.1 $\pm$ 2.1% weight loss) alone [34].

The goal of the current study was to test if systemic OT treatment could be used as

an adjunct with the β3-AR agonist, CL 316243, to increase BAT thermogenesis and elicit

weight loss in DIO rats when using a more translational route of administration for OT

delivery. We hypothesized that systemic OT and β3-AR agonist (CL 316243) treatment

would produce an additive effect to reduce body weight and adiposity in DIO rats by

decreasing food intake and stimulating BAT thermogenesis. To test this, we determined

the effects of systemic (subcutaneous) infusions of OT (50 nmol/day) or vehicle (VEH)

when combined with daily systemic (intraperitoneal) injections of CL 316243 (0.5 mg/kg)

or VEH on body weight, adiposity, food intake, brown adipose tissue temperature (T_{IBAT})

and thermogenic gene expression.

Methods

Animals

Adult male Long-Evans rats [~ 8-9 weeks old, 292-349 grams at start of high fat dietary

(HFD) intervention/~ 8-10 months old, 526-929 g body weight at study onset] were initially

obtained from Envigo (Indianapolis, IN) and maintained for at least 4 months on a high

fat diet (HFD) prior to study onset. All animals were housed individually in Plexiglas cages

in a temperature-controlled room (22±2°C) under a 12:12-h light-dark cycle. All rats were

maintained on a 1 a.m./1 p.m. light cycle. Rats had *ad libitum* access to water and a HFD

providing 60% kcal from fat (approximately 6.8% kcal from sucrose and 8.9% of the diet

from sucrose) (Research Diets, D12492, New Brunswick, NJ). The research protocols

were approved both by the Institutional Animal Care and Use Committee of the Veterans

Affairs Puget Sound Health Care System (VAPSHCS) and the University of Washington

in accordance with NIH's *Guide for the Care and Use of Laboratory Animals* (NAS, 2011)

[35] .

Drug Preparation

Fresh solutions of OT acetate salt (Bachem Americas, Inc., Torrance, CA) were

solubilized in sterile water and subsequently primed in sterile 0.9% saline at 37º C for

approximately 40 hours prior to minipump implantation based on manufacturer's

recommended priming instructions for ALZET® model 2004 minipumps. CL 316243

(Tocris/Bio-Techne Corporation, Minneapolis, MN) was solubilized in sterile water each

day of each experiment.

Subcutaneous implantations of osmotic minipumps

Animals were implanted with an osmotic minipump (model 2004, DURECT Corporation

Cupertino, CA) one week prior to CL 316343 treatment as previously described [34].

Implantation of temperature transponders underneath IBAT

Rats were anesthetized with isoflurane prior to having the dorsal surface along the upper

midline of the back shaved and scrubbed with 70% ethanol followed by betadine swabs

as previously described [28]. Following an incision (1 ") along the midline of the

interscapular area a temperature transponder (14 mm long/2 mm wide) (HTEC IPTT-300;

Bio Medic Data Systems, Inc., Seaford, DE) was implanted underneath the left IBAT pad

as previously described [28; 36; 37]. The transponder was subsequently secured in place

by suturing it to the brown fat pad with sterile silk suture. The interscapular incision was

closed with Nylon sutures (5-0), which were removed in awake animals approximately

10-14 days post-surgery. HTEC IPTT-300 transponders were used in place of IPTT-300

transponders to enhance accuracy in our measurements as previously described [28].

Acute IP injections and measurements of T_{IBAT}

CL 316243 (or saline vehicle; 0.1 ml/kg injection volume) was administered immediately

prior to the start of the dark cycle following 4 hours of food deprivation. Animals remained

without access to food for an additional 1 (**Study 2-3**) or 4 h (**Study 1**) during the course

of the T_{IBAT} measurements. A handheld reader (DAS-8007-IUS Reader System; Bio

Medic Data Systems, Inc.) was used to collect measurements of T_{IBAT}.

Body   *Composition*

Determinations of lean body mass and fat mass were made on un-anesthetized rats by

quantitative magnetic resonance using an EchoMRI 4-in-1-700™ instrument (Echo

Medical Systems, Houston, TX) at the VAPSHCS Rodent Metabolic Phenotyping Core.

Measurements were taken prior to 4V cannulations and minipump implantations as well

as at the end of the infusion period.

Study Protocols

Study 1: Determine the dose-response effects of systemic (SC) infusion of OT (16

and 50 nmol/day) on body weight, adiposity and energy intake in DIO rats.

Rats were fed *ad libitum* and maintained on HFD for approximately 5.5 months prior to

prior to being implanted with a temperature transponder underneath the left IBAT depot.

Rats were subsequently maintained on a daily 4-h fast and received minipumps to infuse

vehicle or OT (16 or 50 nmol/day) over 29 days. These doses was selected based on a

dose of OT found to be effective at reducing body weight when administered

subcutaneously [19] or into the 4V [38] of DIO rats. Daily food intake and body weight

were also tracked for 29 days.

Study 2: Effect of chronic 4V OT infusions (16 nmol/day) and systemic beta-3

receptor agonist (CL 31643) administration (0.5 mg/kg) on body weight, body

adiposity, energy intake and T_{IBAT} in male DIO rats.

Rats (~ 10 mo old; 526-929 g at start of study) were fed *ad libitum* and maintained on

HFD for approximately 7.5 months prior to receiving implantations of temperature

transponders underneath the left IBAT pad in addition to 4V cannulas and 28-day

minipumps to infuse vehicle or OT (16 nmol/day) over 28 days, respectively. After having

matched animals for OT-elicited reductions in body weight (infusion day 7), DIO rats

subsequently received 1x daily IP injections of VEH or CL 316243 (0.5 mg/kg). We

selected this dose because it elevated T_{IBAT} at doses that failed to produce elevations in

heart rate in lean rats [39]. In addition, this dose produced comparable weight loss to that

of OT alone in **Study 2**. T_{IBAT} was measured daily at baseline (-4 h; 9:00 a.m.),

immediately prior to IP injections (0 h; 12:45-1:00 p.m.), and at 0.25, 0.5, 0.75, 1, 20 and

24-h post-injection. In addition, daily food intake and body weight were also tracked for

28 days. Data from animals that received the single dose of CL-31643 were analyzed

over the 28-day infusion period.

Adipose tissue processing for adipocyte size and UCP-1 analysis

IWAT and EWAT depots were collected at the end of the infusion period in rats from

Study 2. Rats from each group were euthanized following a 3-h fast. Rats were

euthanized with intraperitoneal injections of ketamine cocktail [ketamine hydrochloride

(214.3 mg/kg), xylazine (10.71 mg/kg) and acepromazine (3.3 mg/kg) in an injection

volume up to 2 mL/rat] and transcardially exsanguinated with PBS followed by perfusion

with 4% paraformaldehyde in 0.1 M PBS. Adipose tissue (IBAT, IWAT, and EWAT) was

dissected and placed in 4% paraformaldehyde-PBS for 24 h and then placed in 70%

ethanol (EtOH) prior to paraffin embedding. Sections (5 μm) sampled were obtained using

a rotary microtome, slide-mounted using a floatation water bath (37°C), and baked for 30

min at 60°C to give approximately 15-16 slides/fat depot with two sections/slide.

Adipocyte size analysis and UCP-1 staining

Adipocyte size analysis was performed on deparaffinized and digitized IWAT and EWAT

sections. The average cell area from two randomized photomicrographs was determined

using the built-in particle counting method of ImageJ software (National Institutes of

Health, Bethesda, MD). Fixed (4% PFA), paraffin-embedded adipose tissue was

sectioned and stained with a–primary rabbit anti-UCP-1 antibody (1:100; Abcam,

Cambridge, MA (#ab10983/RRID: AB_2241462)] as has been previously described in

lean C57BL/6J mice [40] and both lean and DIO C57BL/6 mice after having been

screened in both IBAT and IWAT of Ucp1$^{+/-}$ and Ucp1$^{-/-}$ mice [41]. Immunostaining

specificity controls included omission of the primary antibody and replacement of the

primary antibody with normal rabbit serum at the same dilution as the respective primary

antibody. Area quantification for UCP1 staining was performed on digital images of

immunostained tissue sections using image analysis software (Image Pro Plus software,

Media Cybernetics, Rockville, MD, USA). Slides were visualized using bright field on an

Olympus BX51 microscope (Olympus Corporation of the Americas; Center Valley, PA)

and photographed using a Canon EOS 5D SR DSLR (Canon U.S.A., Inc., Melville, NY)

camera at 100X magnification. Values for each tissue within a treatment were averaged

to obtain the mean of the treatment group.

Blood collection

Blood was collected from 4-h (**Study 1**) or 6-h fasted rats (**Study 2**) within a 2-h window

towards the end of the light cycle (10:00 a.m.-12:00 p.m.) as previously described in DIO

CD® IGS rats and mice [24; 28]. Animals from **Study 2** were euthanized at 2-h post-CL

316243 or VEH treatment. Treatment groups were counterbalanced at time of euthanasia

to avoid time of day bias. Blood samples [up to 3 mL] were collected immediately prior to

transcardial perfusion by cardiac puncture in chilled K2 EDTA Microtainer Tubes (Becton-

Dickinson, Franklin Lakes, NJ). Whole blood was centrifuged at 6,000 rpm for 1.5-min at

4°C; plasma was removed, aliquoted and stored at −80°C for subsequent analysis.

Plasma hormone measurements

Plasma leptin and insulin were measured using electrochemiluminescence detection

[Meso Scale Discovery (MSD®), Rockville, MD] using established procedures [28; 42].

Intra-assay coefficient of variation (CV) for leptin was 2.7% and 3.2% for insulin. The

range of detectability for the leptin assay is 0.137-100 ng/mL and 0.069-50 ng/mL for

insulin. Plasma fibroblast growth factor-21 (FGF-21) (R&D Systems, Minneapolis, MN)

and irisin (AdipoGen, San Diego, CA) levels were determined by ELISA. The intra-assay

CV for FGF-21 and irisin were 4.5% and 8.4%, respectively; the ranges of detectability

were 31.3-2000 pg/mL (FGF-21) and 0.078-5 μg/mL (irisin). Plasma adiponectin was also

measured using electrochemiluminescence detection Meso Scale Discovery (MSD$^®$),

Rockville, MD] using established procedures [28; 42]. Intra-assay CV for adiponectin was

1.1%. The range of detectability for the adiponectin assay is 2.8-178 ng/mL. The data

were normalized to historical values using a pooled plasma quality control sample that

was assayed in each plate.

Blood glucose and lipid measurements

Blood was collected for glucose measurements by tail vein nick following a 4 (**Study 1**)

or 6-h fast (**Study 2**) and measured with a glucometer using the AlphaTRAK 2 blood

glucose monitoring system (Abbott Laboratories, Abbott Park, IL) [43]. Tail vein glucose

was measured at 2-h post-CL 316243 or VEH treatment (**Study 2**). Total cholesterol (TC)

[Fisher Diagnostics (Middletown, VA)] and free fatty acids (FFAs) [Wako Chemicals USA,

Inc., Richmond, VA)] were measured using an enzymatic-based kits. Intra-assay CVs for

TC and FFAs were 1.4 and 2.3%, respectively. These assay procedures have been

validated for rodents [44].

Tissue collection for quantitative real-time PCR (qPCR)

IBAT and IWAT tissue was collected from 4 (**Study 1**) or 6-h fasted rats (**Study 2**). In

addition, animals from **Study 2** were euthanized at 2-h post-CL 316243 (0.5 mg/kg) or

VEH administration. IBAT and IWAT were collected within a 2-h window towards the end

of the light cycle (10:00 a.m.-12:00 p.m.) as previously described in DIO CD$^®$ IGS/Long-

Evans rats and C57BL/6J mice [24; 28; 38]. Tissue was rapidly removed, wrapped in foil

and frozen in liquid N2. Samples were stored frozen at -80°C until analysis.

qPCR

RNA extracted from samples of IBAT and IWAT (Studies 1-2) were analyzed using the

RNeasy Lipid Mini Kit (Qiagen Sciences Inc, Germantown, MD) followed by reverse

transcription into cDNA using a high-capacity cDNA archive kit (Applied Biosystems,

Foster City, CA). Quantitative analysis for relative levels of mRNA in the RNA extracts

was measured in duplicate by qPCR on an Applied Biosystems 7500 Real-Time PCR

system (Thermo Fisher Scientific, Waltham, MA) and normalized to the cycle threshold

value of Nono mRNA in each sample. The TaqMan® probes used in the study were

Thermo Fisher Scientific Gene Expression Assay probes. The probe for rat *Nono*

(Rn01418995_g1), UCP-1 (*Ucp1*; catalog no. Rn00562126_m1), beta 1 adrenergic

receptor (β1-AR) (*Adrb1*; catalog no. Rn00824536_s1), β3-AR (*Adrb3*; catalog no.

Rn01478698_g1), type 2 deiodinase (D2) (*Dio2*; catalog no. Rn00581867_m1), G-protein

coupled receptor 120 (*Gpr120*; catalog no. Rn01759772_m1), cell death-inducing DNA

fragmentation factor α-like effector A (*Cidea*; catalog no. Rn04181355_m1), peroxisome

proliferator-activated receptor gamma coactivator 1α (*Ppargc1a*; catalog no.

Rn00580241_m1) and PR domain containing 16 (*Prdm16*; catalog no. Rn01516224_m1)

were acquired from Thermo Fisher Scientific. Relative amounts of target mRNA were

determined using the Comparative C_T or $2^{-\Delta\Delta C_T}$ method [45] following adjustment for the

housekeeping gene. Specific mRNA levels of all genes of interest were normalized to the

cycle threshold value of Nono mRNA in each sample and expressed as changes

normalized to controls (vehicle/vehicle treatment).

Transponder placement

All temperature transponders were confirmed to have remained underneath the IBAT

depot at the conclusion of the study.

Statistical Analyses

All results are expressed as means ± SE. Comparisons between multiple groups involving

between subjects designs were made using one- or two-way ANOVA as appropriate,

followed by a post-hoc Fisher's least significant difference test. Comparisons involving

within-subjects designs were made using a one-way repeated-measures ANOVA

followed by a post-hoc Bonferroni Test. Analyses were performed using the statistical

program SYSTAT (Systat Software, Point Richmond, CA). Differences were considered

significant at $P<0.05$, 2-tailed.

Results

Study 1: Determine the dose-response effects of systemic (SC) infusion of OT (16

and 50 nmol/day) on body weight, adiposity and energy intake in DIO rats

Prior to treatment, groups were matched for body weight and adiposity

(vehicle: 749±50 grams; OT (16 nmol/day): 763.6±37.6 grams/42±2.6% fat; OT (50

nmol/day): 759±45.4 grams/39.2±1.6%. There was no difference in body weight

$[(F_{(2,17)} = 0.027, P=NS)]$ or percent adiposity $[(F_{(2,17)} = 0.003, P=NS)]$ between groups

prior to treatment onset. As expected, body weight of DIO rats remained stable over the

month of vehicle treatment relative to pre-treatment [($F_{(1,5)}$ = 2.865, P=0.151)] (**Figure**

1A). In contrast to vehicle treatment, systemic OT (16 nmol/day) resulted in a significant

reduction of body weight relative to OT pre-treatment [($F_{(1,6)}$ = 140.799, P<0.01)] (**Figure**

1A; P<0.05). Furthermore, SC OT, at a 3-fold higher dose (50 nmol/day), also resulted in

a significant elevation of body weight relative to pre-treatment [($F_{(1,6)}$ = 47.271, P<0.01)].

In addition, SC OT (16 and 50 nmol/day) was able to reduce weight gain (**Figure 1B**)

relative to vehicle treatment throughout the 28-day infusion period. SC OT (50 nmol/day),

at a dose that was at least 3-fold higher than the centrally effective dose (16 nmol/day),

reduced weight gain throughout the entire 28-day infusion period. SC OT (16 nmol/day)

treated rats had reduced weight gain between days 2-29 (P<0.05) while SC OT (50

nmol/day) reduced weight gain between days 1-29 (P<0.05). There was an overall effect

of OT to reduce relative fat mass (pre- vs post-intervention) [($F_{(2,17)}$ = 6.052, P=0.010)].

SC OT (50 nmol/day) reduced fat mass (P<0.05) and there was also a tendency for the

lower dose (16 nmol/day) to reduce relative fat mass (P=0.066) (**Figure 1C**; P<0.05).

There was also an overall effect of OT to reduce relative lean mass (pre- vs post-

intervention) [($F_{(2,17)}$ = 5.572, P=0.014)]. Specifically, SC OT (16 nmol/day) reduced

relative lean mass at the lower dose (16 nmol/day; P<0.01) while the higher dose (50

nmol/day) tended to reduce relative lean mass (P=0.090). Note that there was no

significant reduction in total fat mass or lean mass (P=NS).

The changes in body weight and relative fat mass were not associated with any changes

in plasma leptin, insulin, glucose or total cholesterol (**Table 1**). These effects that were

mediated, at least in part, by a modest reduction of energy intake that was apparent during

weeks 1 (50 nmol/day) and 2 (16 and 50 nmol/day) of OT treatment (**Figure 1D**; $P<0.05$).

Study 2: Effect of chronic 4V OT infusions (16 nmol/day) and systemic β3-AR

agonist (CL 31643) administration (0.5 mg/kg) on body weight, body adiposity and

energy intake in male DIO rats.

The goal of this study was to determine the effects of chronic OT treatment in combination

with a single dose (identified in **Study 1**) of the β3-AR agonist, CL 316243, on body weight

and adiposity in DIO rats. By design, DIO rats were obese as determined by both body

weight (804±14 g) and adiposity (310±11 g fat mass; 38.3±4.7% adiposity) after

maintenance on the HFD for approximately 7 months months. Prior to the onset of CL

316243 treatment on infusion day 7, both OT treatment groups were matched for OT-

elicited reductions of weight gain.

OT and CL 316243 alone reduced body weight by ≈ 7.8±1.3% ($P<0.05$) and 9.1±2.3%

($P<0.05$), respectively, but the combined treatment produced more pronounced weight

loss (pre- vs post-intervention) (15.5±1.2%; $P<0.05$) (**Figure 2A**) than either treatment

alone ($P<0.05$). OT alone tended to reduce weight gain on days 16-28 ($0.05<P<0.1$) while

CL 316243 alone tended to reduce or reduced weight gain on day 24 ($0.05<P<0.1$), day

25 ($P=0.05$), and days 26-28 ($P<0.05$) (**Figure 2B**). OT and CL 316243 together tended

to reduce weight gain on day 9 ($0.05<P<0.1$) reduced weight gain on days 10-28

(*P*<0.05). The combination treatment appeared to produce a more pronounced reduction

of weight gain relative to OT alone on day 25 (0.05<*P*<0.1) and this reached significance

on days 26-28 (*P*<0.05).

In addition, the combination treatment tended to produce a greater reduction of weight

gain relative to CL 316243 alone on days 21-22 and 26-28 (0.05<*P*<0.1). While OT alone

did not significantly reduce fat mass (*P*=NS), there was a tendency for CL 316243 alone

(0.05<*P*<0.1), and the combination of OT and CL 316243 (*P*<0.05) to reduce fat mass

without impacting lean body mass (**Figure 2C**; *P*=NS). However, the combination

treatment did not result in a significant reduction of fat mass relative to OT alone or CL

316243 alone (*P*=NS). Consistent with these effects, OT alone did not produce a

reduction in relative fat mass (*P*=0.108) while CL 316243 alone reduced relative fat mass

(pre- vs post-intervention; *P*<0.05) without changing lean mass (**Figure 2D**; *P*=NS). The

combination treatment produced a significant reduction of relative fat mass (pre- vs post-

intervention; *P*<0.05) which exceeded that of OT alone (*P*<0.05) but not CL 316243 alone

(*P*=NS). OT, CL 316243 and the combined treatment were all effective at reducing energy

intake at week 2 (**Figure 2E**; *P*<0.05) with no difference between treatments (*P*=NS). All

treatments were ineffective at reducing energy intake over weeks 3 and 4 (*P*=NS).

Two-way ANOVA revealed an overall significant effect of OT [($F_{(1,39)}$ = 63.434, *P*<0.01)],

CL 316243 [($F_{(1,39)}$ = 74.939, *P*<0.01)] but no significant interactive effect between OT

and CL 316243 [($F_{(1,39)}$ = 0.058, *P*=NS)] on weight loss. Consistent with this finding,

two-way ANOVA revealed consistent overall effects of OT and CL-3162343 on reduction

of body weight gain between days 10-29 but no significant overall effect. In addition, two-

way ANOVA revealed an overall significant effect of OT [(F(1,39) = 4.989, P=0.031)], CL

316243 [(F(1,39) = 7.607, P=0.009)] and a near significant interactive effect between OT

and CL 316243 [(F(1,39) = 2.945, P=0.094)] on fat mass. There was no significant overall

effect of OT [(F(1,39) = 1.351, P=NS)], on lean mass but there was an overall effect of

CL 316243 [(F(1,39) = 4.829, P=0.034)], and an interactive effect of OT and CL 316243

[(F(1,39) = 4.905, P=0.033)] on lean mass.

Lastly, two-way ANOVA revealed an overall significant effect of OT [(F(1,39) = 21.464,

P<0.01)], CL 316243 [(F(1,39) = 62.681, P<0.01)] and a near significant interactive effect

between OT and CL 316243 [(F(1,39) = 3.190, P=0.082)] on energy intake (week 2).

Overall, these findings suggest an additive effect of OT and CL 316243 to produce

sustained weight loss in DIO rats. The effects of the combination treatment on adiposity

and energy intake appear to be driven largely by CL 316243 and OT, respectively.

CL 316243 elevated T_{IBAT} on injection day 1 at 0.5, 0.75 and 1-h post-injection (P<0.05;

Table 2A) and tended to elevate T_{IBAT} at 0.25-h post-injection (0.05<P<0.1). Similarly, CL

316243, when given in combination with OT, also increased T_{IBAT} at 0.5, 0.75 and 1-h

post-injection and tended to elevate T_{IBAT} at 0.25-h post-injection on injection day 1

(0.05<P<0.1). Both CL 316243 and CL 316243 + OT treatments elevated T_{IBAT} relative to

vehicle treated animals when the T_{IBAT} data from injection day 1 were averaged over 1-h

post-injection (TIBATP<0.05). There was no significant difference in T_{IBAT} response to CL

316243 CL 316243 + OT treatments when the T_{IBAT} data were averaged over 60 min

(P=NS).

CL 316243 also elevated T_{IBAT} on injection day 22 at 0.25, 0.5, 0.75 and 1-h post-injection

(P<0.05; **Table 2B**). Similarly, CL 316243, when given in combination with OT, also

increased T_{IBAT} at 0.5, 0.75 and 1-h post-injection and tended to elevate T_{IBAT} at 0.25 and

1-h post-injection on injection day 22 (0.05<P<0.1). Both CL 316243 and CL 316243 +

OT treatments elevated T_{IBAT} relative to vehicle treated animals when the T_{IBAT} data from

injection day 22 were averaged over 1-h post-injection (P<0.05). There was no significant

difference in the T_{IBAT} response to CL 316243 and CL 316243 treatments when the T_{IBAT}

data were averaged over 60 min (P=NS).

Adipocyte size

OT and CL 316243 given in combination reduced IWAT adipocyte size whereas there

was no significant effect of OT or CL 316343 on adipocyte size when given alone (P=NS)

(**Figure 3A**-**D**; **Figure 4A**). In contrast, CL 316243 alone (P<0.05) or in combination with

OT (P<0.05) reduced EWAT adipocyte size in DIO rats relative to vehicle

treatment **(Figure 5E**-**H**; **Figure 4B**). There were no significant differences in the ability

of the combined treatment to reduce EWAT adipocyte size relative to OT alone (P=0.119)

or CL 316243 alone (P=0.156).

Plasma hormone concentrations

To characterize the endocrine and metabolic effects of 4V OT (16 nmol/day) and systemic

beta-3 receptor agonist (CL 316243) in DIO rats in a chronic study using as single dose

of CL 316243 (**Study 2; Table 3**), we measured blood glucose levels and plasma

concentrations of leptin, insulin, FGF-21, irisin, adiponectin, TC, triglycerides, and FFAs.

CL 316243 alone or in combination with OT resulted in a reduction of plasma leptin

relative to OT (*P*<0.05) or vehicle alone (*P*<0.05). The combination treatment was also

associated with a reduction of blood glucose and insulin relative to vehicle (*P*<0.05) and

OT treatment (*P*<0.05) but it was not statistically different from CL 316243 (*P*=NS). We

also found that FGF-21 was reduced in response to CL 316243 and CL 316243 in

combination with OT relative to vehicle and OT treatment (*P*<0.05). In addition, the

combination treatment was associated with an elevation of adiponectin relative to vehicle

(*P*<0.05) and OT treatment (*P*<0.05) but it was not statistically different from CL 316243

(*P*=NS).

Gene expression data

We next determined the extent to which CL 316243 (0.5 mg/kg), OT (16 nmol/day), or the

combination treatment increased thermogenic gene expression in IBAT relative to vehicle

at 2-hour post-CL 316243/vehicle injections.

IBAT: Consistent with published findings in mice and rats, chronic CL 316243

administration elevated relative levels of the thermogenic markers *Ucp1* [46; 47; 48], *Dio2*

[34; 49], *Ppargc1a* [47] and *Gpr120* [34; 47] (**Table 4A**; *P*<0.05). CL 316243 in

combination with OT also elevated *Ucp1*, *Dio2, Ppargc1a* and *Gpr120* (*P*<0.05). In

addition, chronic CL 316243 alone and in combination with OT reduced *Adrb3* (β3-AR)

mRNA expression in IBAT (**Table 4A**; *P*<0.05).

IWAT: There were no significant differences in the thermogenic markers *Ucp1*, *Dio2*,

Ppargc1a or *Adrb3* in response to chronic CL 316243 alone or in combination with OT

(**Table 4B**; *P*=NS).

As a functional readout of BAT thermogenesis for the gene expression analyses, we

measured T_{IBAT} in response to CL 316243 alone or CL 316243 + OT during the time

period that preceded tissue collection. CL 316243 alone resulted in an increase in T_{IBAT}

at 0.25, 0.5, 0.75 and 1-h post-injection (**Table 5**; *P*<0.05). Similarly, CL 316243 + OT

resulted in an increase in T_{IBAT} at 0.25, 0.5 and 1-h post-injection (**Table 5**; *P*<0.05) and

it also tended to increase T_{IBAT} at 0.75-h post-injection (**Table 5**; 0.05<*P*<0.1).

Discussion

The goal of the current study was to determine the extent to which systemic OT could be

used as an adjunct with the β3-AR agonist, CL 316243, to increase BAT thermogenesis

and elicit weight loss in DIO rats. We hypothesized that systemic OT and beta-3 agonist

(CL 316243) treatment would produce an additive effect to reduce body weight and

adiposity in DIO rats by decreasing food intake and stimulating BAT thermogenesis. To

test this hypothesis, we determined the effects of systemic (subcutaneous) infusions of

OT (50 nmol/day) or vehicle (VEH) when combined with daily systemic (intraperitoneal)

injections of CL 316243 (0.5 mg/kg) or VEH on body weight, adiposity, food intake and

brown adipose tissue temperature (T_{IBAT}). OT and CL 316243 monotherapy decreased

body weight by 8.0±0.9% (*P*<0.05) and 8.6±0.6% (*P*<0.05), respectively, but OT in

combination with CL 316243 produced more substantial weight loss (14.9±1.0%; *P*<0.05)

compared to either treatment alone. These effects were associated with decreased

adiposity, energy intake and elevated T_{IBAT} during the treatment period. In addition,

systemic OT and CL 316243 combination therapy increased IBAT thermogenic gene

expression suggesting that increased BAT thermogenesis may also contribute to these

effects. The findings from the current study suggest that the effects of systemic OT and

CL 316243 to reduce body weight and adiposity are additive and appear to be driven

primarily by OT-elicited changes in food intake and CL 316243-elicited increases in BAT

thermogenesis.

Recent findings also indicate that other agents, namely the glucagon-like peptide-1

receptor agonist, liraglutide, and the fat signal, oleoylethanolamide (OEA), act in an

additive fashion with CL 316243 to reduce body weight or body weight gain. Oliveira

recently reported that the combination of liraglutide and CL 316243 produce additive

effects to reduce the change in body weight in a mouse model [50]. These effects

appeared to be attributed to additive effects on energy intake and increased oxygen

consumption in IBAT and IWAT. In addition, the combined treatment increased

expression of UCP-1 in IWAT (indicative of browning). Similarly, Suarez and colleagues

demonstrated that the peroxisome proliferator-activating receptor-α (PPARα) agonist and

fat signal, oleoylethanolamide (OEA), act in an additive fashion with CL 316243 produced

to reduce food intake and weight gain in rats [32]. These effects were associated with

pronounced reductions in fat mass and increases in expression of thermogenic genes

(*PPARα* and *Ucp1*) in EWAT [32]. Of interest is the finding that systemic OT and central

OT can increase OEA expression in EWAT [19]. Furthermore, Deblon reported that the

effectiveness of OT to decrease body weight was partially blocked in PPARα null mice

[19], indicating that PPARα may partially mediate OT's thermogenic effects in EWAT.

OEA has also been found to stimulate 1) hypothalamic expression of OT mRNA [51], 2)

PVN OT neurons [52], and 3) OT release within the PVN [52]. In addition, OEA also

decreases food intake, in part, through OT receptor signaling [51]. Thus, it is possible that

high fat diet-elicited stimulation of OEA [53] may reduce food intake, in part, through an

OTR signaling and that the effects of OT to stimulate WAT thermogenesis might occur

through PPARα. Additional studies that utilize adipose depot knockdown of PPARα will

enable us to determine if PPARα in specific adipose depots may contribute to the ability

of OT and CL 316243 to reduce body weight and adiposity.

Our findings and others raise the possibility that systemic OT could be reducing food

intake and adiposity, in part, through a direct effect on peripheral OT receptors. Asker

reported that OT-B12, a BBB-impermeable OT analogue, reduced food intake in rats,

thus providing evidence that peripheral OTR signaling is important in the control of food

intake [54]. Consistent with these findings, Iwasaki found that the ability of peripheral

administration of OT to reduce food intake was attenuated in vagotomized mice [55; 56].

In addition, Brierley extended these findings and found that the effect of systemic

administration of OT to suppress food intake required NTS preproglucagon neurons that

receive direct synaptic input OTR-expressing vagal afferent neurons [57]. We also found

that systemic administration of a non-BBB penetrant OTR antagonist, L-371257,

stimulated food intake and body weight gain in [58]. Several studies have also found that

subcutaneous infusion of OT reduced adipocyte size in 1) visceral fat in female Wistar

rats that were peri- and postmenopausal [59] and 2) visceral fat in a dihydrotestosterone-

elicited model of polycystic ovary syndrome in female Wistar rats [60], 3) subcutaneous

fat in female ovariectomized Wistar rats [61], and 4) EWAT in male Zucker fatty rats [62].

More recent studies have confirmed that peripheral administration of long-acting OT

analogues (including ASK2131 and ASK1476) also reduced both food intake and body

weight [63; 64]. Together, these findings suggest that OT may also act in the periphery to

decrease adipocyte size by a direct effect on OTRs found on adipose tissue [19; 20; 65].

Of translational importance is the finding that subcutaneous [19; 24; 25; 26; 27; 66] or

intraperitoneal [27] administration of OT or long-acting OT analogues can recapitulate the

effects of chronic CNS administration of OT on reductions of food intake and body weight.

The combination treatment and CL 316243 monotherapy reduced body weight and

adiposity, in part, through increased BAT thermogenesis. Both CL 316243 alone and in

combination with OT elevated T_{IBAT} throughout the course of the injection study and

increased IBAT thermogenic genes (UCP-1, DIO2 and Ppargc1a) and UCP-1 content in

IBAT. These findings coincided with CL- 316243-elicited increases in T_{IBAT} from the same

animals during the time that preceded tissue collection. These findings are consistent with

our previously published findings in rats [34] and other studies mice and rats that found

chronic CL 316243 administration to increase the thermogenic markers DIO2 [49] and

Gpr120 [46; 47; 48]. Similar to our findings, others also reported that systemic CL 316243

increased UCP-1 mRNA expression in mice [67]. We also found that the combination

treatment and CL 316243 alone caused a downregulation of the β3-AR @ 2-h post-

injection. This finding is consistent with we have previously reported [34] and others who

have reported that both cold exposure and norepinephrine reduced β3-AR mRNA

expression in mouse IBAT [68; 69] and.mouse brown adipocytes [70].

In summary, our results indicate that systemic administration of OT in combination

with systemic CL 316243 treatment results in more profound reductions of body weight

compared to either OT or CL 316243 alone. Moreover, the combined treatment of OT and

CL 316243 stimulated BAT thermogenesis as determined by increased T_{IBAT} and

thermogenic gene expression in IBAT. Together, our data support the hypothesis that

systemic OT and β3-AR agonist (CL 316243) treatment produce an additive effect to

reduce body weight and adiposity in DIO rats. The effects of the combined treatment on

body weight and adiposity appeared to be additive and driven predominantly by OT-

elicited reductions of food intake and CL-316243-elicited increases in BAT

thermogenesis.

Collectively, these findings suggest that systemic OT treatment could be a viable

adjunct to other anti-obesity treatment strategies. While intranasal OT has been found to

reduce body weight by approximately 9.3% in a small study with limited subjects (≈9.3%)

[71], it was not found, however, to have any effect on body weight in a larger scale clinical

study (N=subjects) in which subjects were matched well for body weight, adiposity and

gender [72]. Importantly, Plessow, Lawson and colleagues did find a significant effect of

intranasal OT to reduce energy intake at 6-weeks post-treatment, which served as an

important control. It is possible that a more extended length of treatment might have been

required to take advantage of the reductions of energy intake that were not observed until

the 6-week post-treatment time point. In addition, changes in dose, dosing frequency, or

co-administration with Mg^{2+} [73; 74] might need to be taken into consideration in order to

maximize the effects of intranasal OT on body weight in humans who are overweight or

obese. Given that OT can be an effective delivery approach to reduce energy intake and

elicit weight loss in several rodent models (see [15; 75; 76] for review) and obese nonhuman

primates [18], it will also be important to determine if chronic systemic OT treatment can

elicit weight loss when given in combination with CL 316243 at doses that are sub-

threshold for producing adverse effects on heart rate or blood pressure [77]. Recent

findings indicate that OT can be effective at reducing food intake and/or body weight in

female rats [78] and DIO male mice [25], respectively. Thus, it will be important to examine

if this combination treatment produces an additive effect to reduce body weight and

adiposity in female DIO rodents and nonhuman primates.

References

[1] S. Smyth, and A. Heron, Diabetes and obesity: the twin epidemics. Nature medicine 12 (2006) 75-80.

[2] G. Targher, A. Mantovani, X.B. Wang, H.D. Yan, Q.F. Sun, K.H. Pan, C.D. Byrne, K.I. Zheng, Y.P. Chen, M. Eslam, J. George, and M.H. Zheng, Patients with diabetes are at higher risk for severe illness from COVID-19. Diabetes & metabolism (2020).

[3] J.P.H. Wilding, R.L. Batterham, S. Calanna, M. Davies, L.F. Van Gaal, I. Lingvay, B.M. McGowan, J. Rosenstock, M.T.D. Tran, T.A. Wadden, S. Wharton, K. Yokote, N. Zeuthen, R.F. Kushner, and S.S. Group, Once-Weekly Semaglutide in Adults with Overweight or Obesity. The New England journal of medicine 384 (2021) 989.

[4] J.P. Frias, M.J. Davies, J. Rosenstock, F.C. Perez Manghi, L. Fernandez Lando, B.K. Bergman, B. Liu, X. Cui, K. Brown, and S.-. Investigators, Tirzepatide versus Semaglutide Once Weekly in Patients with Type 2 Diabetes. The New England journal of medicine 385 (2021) 503-515.

[5] O.G. Chepurny, R.L. Bonaccorso, C.A. Leech, T. Wollert, G.M. Langford, F. Schwede, C.L. Roth, R.P. Doyle, and G.G. Holz, Chimeric peptide EP45 as a dual agonist at GLP-1 and NPY2R receptors. Scientific reports 8 (2018).

[6] J.P. Frias, E.J. Bastyr, L. Vignati, M.H. Tschop, C. Schmitt, K. Owen, R.H. Christensen, and R.D. DiMarchi, The Sustained Effects of a Dual GIP/GLP-1 Receptor Agonist, NNC0090-2746, in Patients with Type 2 Diabetes. Cell Metabolism 26 (2017) 343-+.

[7] L.B. Enebo, K.K. Berthelsen, M. Kankam, M.T. Lund, D.M. Rubino, A. Satylganova, and D.C.W. Lau, Safety, tolerability, pharmacokinetics, and pharmacodynamics of concomitant administration of multiple doses of cagrilintide with semaglutide 2.4 mg for weight management: a randomised, controlled, phase 1b trial. Lancet 397 (2021) 1736-1748.

[8] J.P. Frias, S. Deenadayalan, L. Erichsen, F.K. Knop, I. Lingvay, S. Macura, C. Mathieu, S.D. Pedersen, and M. Davies, Efficacy and safety of co-administered once-weekly cagrilintide 2.4 mg with once-weekly semaglutide 2.4 mg in type 2 diabetes: a multicentre, randomised, double-blind, active-controlled, phase 2 trial. Lancet 402 (2023) 720-730.

[9] D.B. Allison, K.M. Gadde, W.T. Garvey, C.A. Peterson, M.L. Schwiers, T. Najarian, P.Y. Tam, B. Troupin, and W.W. Day, Controlled-release phentermine/topiramate in severely obese adults: a randomized controlled trial (EQUIP). Obesity 20 (2012) 330-42.

[10] A.M. Jastreboff, L.J. Aronne, N.N. Ahmad, S. Wharton, L. Connery, B. Alves, A. Kiyosue, S. Zhang, B. Liu, M.C. Bunck, A. Stefanski, and S.-. Investigators, Tirzepatide Once Weekly for the Treatment of Obesity. The New England journal of medicine 387 (2022) 205-216.

[11] L.J. Aronne, N. Sattar, D.B. Horn, H.E. Bays, S. Wharton, W.Y. Lin, N.N. Ahmad, S. Zhang, R. Liao, M.C. Bunck, I. Jouravskaya, M.A. Murphy, and S.-. Investigators, Continued Treatment With Tirzepatide for Maintenance of Weight Reduction in Adults With Obesity: The SURMOUNT-4 Randomized Clinical Trial. JAMA : the journal of the American Medical Association (2023).

[12] A.M. Jastreboff, L.M. Kaplan, and M.L. Hartman, Triple-Hormone-Receptor Agonist Retatrutide for Obesity. Reply. The New England journal of medicine 389 (2023) 1629-1630.

[13] G. Gimpl, and F. Fahrenholz, The oxytocin receptor system: structure, function, and regulation. Physiol Rev 81 (2001) 629-83.

[14] J.E. Blevins, and D.G. Baskin, Translational and therapeutic potential of oxytocin as an anti-obesity strategy: Insights from rodents, nonhuman primates and humans. Physiol Behav (2015).

[15] E.A. Lawson, The effects of oxytocin on eating behaviour and metabolism in humans. Nature reviews. Endocrinology 13 (2017) 700-709.

[16] E.A. Lawson, P.K. Olszewski, A. Weller, and J.E. Blevins, The role of oxytocin in regulation of appetitive behaviour, body weight and glucose homeostasis. J Neuroendocrinol 32 (2020) e12805.

[17] S.E. McCormack, J.E. Blevins, and E.A. Lawson, Metabolic Effects of Oxytocin. Endocrine reviews 41 (2020).

[18] J.E. Blevins, J.L. Graham, G.J. Morton, K.L. Bales, M.W. Schwartz, D.G. Baskin, and P.J. Havel, Chronic oxytocin administration inhibits food intake, increases energy expenditure, and produces weight loss in fructose-fed obese rhesus monkeys. Am J Physiol Regul Integr Comp Physiol 308 (2015) R431-8.

[19] N. Deblon, C. Veyrat-Durebex, L. Bourgoin, A. Caillon, A.L. Bussier, S. Petrosino, F. Piscitelli, J.J. Legros, V. Geenen, M. Foti, W. Wahli, V. Di Marzo, and F. Rohner-Jeanrenaud, Mechanisms of the anti-obesity effects of oxytocin in diet-induced obese rats. PloS one 6 (2011) e25565.

[20] K.J. Yi, K.H. So, Y. Hata, Y. Suzuki, D. Kato, K. Watanabe, H. Aso, Y. Kasahara, K. Nishimori, C. Chen, K. Katoh, and S.G. Roh, The regulation of oxytocin receptor gene expression during adipogenesis. J Neuroendocrinol 27 (2015) 335-42.

[21] E.E. Noble, C.J. Billington, C.M. Kotz, and C. Wang, Oxytocin in the ventromedial hypothalamic nucleus reduces feeding and acutely increases energy expenditure. Am J Physiol Regul Integr Comp Physiol 307 (2014) R737-45.

[22] G. Zhang, H. Bai, H. Zhang, C. Dean, Q. Wu, J. Li, S. Guariglia, Q. Meng, and D. Cai, Neuropeptide exocytosis involving synaptotagmin-4 and oxytocin in hypothalamic programming of body weight and energy balance. Neuron 69 (2011) 523-35.

[23] G. Zhang, and D. Cai, Circadian intervention of obesity development via resting-stage feeding manipulation or oxytocin treatment. Am J Physiol Endocrinol Metab 301 (2011) E1004-12.

[24] J.E. Blevins, B.W. Thompson, V.T. Anekonda, J.M. Ho, J.L. Graham, Z.S. Roberts, B.H. Hwang, K. Ogimoto, T.H. Wolden-hanson, J.O. Nelson, K.J. Kaiyala, P.J. Havel, K.L. Bales, G.J. Morton, M.W. Schwartz, and D.G. Baskin, Chronic CNS oxytocin signaling preferentially induces fat loss in high fat diet-fed rats by enhancing satiety responses and increasing lipid utilization. Am J Physiol-Reg I (2016).

[25] Y. Maejima, M. Aoyama, K. Sakamoto, T. Jojima, Y. Aso, K. Takasu, S. Takenosihita, and K. Shimomura, Impact of sex, fat distribution and initial body weight on oxytocin's body weight regulation. Scientific reports 7 (2017) 8599.

[26] Y. Maejima, Y. Iwasaki, Y. Yamahara, M. Kodaira, U. Sedbazar, and T. Yada, Peripheral oxytocin treatment ameliorates obesity by reducing food intake and visceral fat mass. Aging 3 (2011) 1169-77.

[27] G.J. Morton, B.S. Thatcher, R.D. Reidelberger, K. Ogimoto, T. Wolden-Hanson, D.G. Baskin, M.W. Schwartz, and J.E. Blevins, Peripheral oxytocin suppresses food intake and causes weight loss in diet-induced obese rats. Am J Physiol-Endoc M 302 (2012) E134-E144.

[28] Z.S. Roberts, T.H. Wolden-Hanson, M.E. Matsen, V. Ryu, C.H. Vaughan, J.L. Graham, P.J. Havel, D.W. Chukri, M.W. Schwartz, G.J. Morton, and J.E. Blevins, Chronic Hindbrain Administration of Oxytocin is Sufficient to Elicit Weight Loss in Diet-Induced Obese Rats. Am J Physiol Regul Integr Comp Physiol (2017) ajpregu 00169 2017.

[29] M.A. Head, A.S. Levine, D.G. Christian, A. Klockars, and P.K. Olszewski, Effect of combination of peripheral oxytocin and naltrexone at subthreshold doses on food intake, body weight and feeding-related brain gene expression in male rats. Physiol Behav 238 (2021) 113464.

[30] P.J. Enriori, P. Sinnayah, S.E. Simonds, C. Garcia Rudaz, and M.A. Cowley, Leptin action in the dorsomedial hypothalamus increases sympathetic tone to brown adipose tissue in spite of systemic leptin resistance. J Neurosci 31 (2011) 12189-97.

[31] M.R. Mirbolooki, S.K. Upadhyay, C.C. Constantinescu, M.L. Pan, and J. Mukherjee, Adrenergic pathway activation enhances brown adipose tissue metabolism: A [F-18]FDG PET/CT study in mice. Nucl Med Biol 41 (2014) 10-16.

[32] J. Suarez, P. Rivera, S. Arrabal, A. Crespillo, A. Serrano, E. Baixeras, F.J. Pavon, M. Cifuentes, R. Nogueiras, J. Ballesteros, C. Dieguez, and F.R. de Fonseca, Oleoylethanolamide enhances beta-adrenergic-mediated thermogenesis and white-to-brown adipocyte phenotype in epididymal white adipose tissue in rat. Disease models & mechanisms 7 (2014) 129-141.

[33] C. Xiao, M. Goldgof, O. Gavrilova, and M.L. Reitman, Anti-obesity and metabolic efficacy of the beta3-adrenergic agonist, CL316243, in mice at thermoneutrality compared to 22 degrees C. Obesity 23 (2015) 1450-9.

[34] M.M. Edwards, H.K. Nguyen, A.D. Dodson, A.J. Herbertson, T.A. Wietecha, T. Wolden-Hanson, J.L. Graham, M.K. Honeycutt, J.D. Slattery, K.D. O'Brien, P.J. Havel, and J.E. Blevins, Effects of combined oxytocin and beta-3 receptor agonist (CL 316243) treatment on body weight and adiposity in male diet-induced obese rats. Frontiers in physiology (2021).

[35] Guide for the Care and Use of Laboratory Animals: Eighth Edition, The National Academies Press, Washington, D.C., 2011.

[36] M.N. Brito, N.A. Brito, D.J. Baro, C.K. Song, and T.J. Bartness, Differential activation of the sympathetic innervation of adipose tissues by melanocortin receptor stimulation. Endocrinology 148 (2007) 5339-47.

[37] C.H. Vaughan, Y.B. Shrestha, and T.J. Bartness, Characterization of a novel melanocortin receptor-containing node in the SNS outflow circuitry to brown adipose tissue involved in thermogenesis. Brain Res 1411 (2011) 17-27.

[38] M.M. Edwards, H.K. Nguyen, A.J. Herbertson, A.D. Dodson, T. Wietecha, T. Wolden-Hanson, J.L. Graham, K.D. O'Brien, P.J. Havel, and J.E. Blevins, Chronic Hindbrain Administration of Oxytocin Elicits Weight Loss in Male Diet-Induced Obese Mice. Am J Physiol Regul Integr Comp Physiol (2021).

[39] B. Malinowska, and E. Schlicker, Further evidence for differences between cardiac atypical beta-adrenoceptors and brown adipose tissue beta3-adrenoceptors in the pithed rat. British journal of pharmacology 122 (1997) 1307-14.

[40] Q. Cao, J. Jing, X. Cui, H. Shi, and B. Xue, Sympathetic nerve innervation is required for beigeing in white fat. Physiological reports 7 (2019) e14031.

[41] M.M. Veniant, G. Sivits, J. Helmering, R. Komorowski, J. Lee, W. Fan, C. Moyer, and D.J. Lloyd, Pharmacologic Effects of FGF21 Are Independent of the "Browning" of White Adipose Tissue. Cell Metab 21 (2015) 731-8.

[42] A.A. Bremer, K.L. Stanhope, J.L. Graham, B.P. Cummings, W. Wang, B.R. Saville, and P.J. Havel, Fructose-fed rhesus monkeys: a nonhuman primate model of insulin resistance, metabolic syndrome, and type 2 diabetes. Clinical and translational science 4 (2011) 243-52.

[43] J.E. Blevins, D.H. Moralejo, T.H. Wolden-Hanson, B.S. Thatcher, J.M. Ho, K.J. Kaiyala, and K. Matsumoto, Alterations in activity and energy expenditure contribute to lean phenotype in Fischer 344 rats lacking the cholecystokinin-1 receptor gene. Am J Physiol Regul Integr Comp Physiol 303 (2012) R1231-40.

[44] B.P. Cummings, E.K. Digitale, K.L. Stanhope, J.L. Graham, D.G. Baskin, B.J. Reed, I.R. Sweet, S.C. Griffen, and P.J. Havel, Development and characterization of a novel rat model of type 2 diabetes mellitus: the UC Davis type 2 diabetes mellitus UCD-T2DM rat. Am J Physiol Regul Integr Comp Physiol 295 (2008) R1782-93.

[45] K.J. Livak, and T.D. Schmittgen, Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25 (2001) 402-8.

[46] E. Gonzalez-Hurtado, J. Lee, J. Choi, and M.J. Wolfgang, Fatty acid oxidation is required for active and quiescent brown adipose tissue maintenance and thermogenic programing. Molecular metabolism 7 (2018) 45-56.

[47] M. Rosell, M. Kaforou, A. Frontini, A. Okolo, Y.W. Chan, E. Nikolopoulou, S. Millership, M.E. Fenech, D. MacIntyre, J.O. Turner, J.D. Moore, E. Blackburn, W.J. Gullick, S. Cinti, G. Montana, M.G. Parker, and M. Christian, Brown and white adipose tissues: intrinsic differences in gene expression and response to cold exposure in mice. Am J Physiol Endocrinol Metab 306 (2014) E945-64.

[48] J. Suarez, P. Rivera, S. Arrabal, A. Crespillo, A. Serrano, E. Baixeras, F.J. Pavon, M. Cifuentes, R. Nogueiras, J. Ballesteros, C. Dieguez, and F. Rodriguez de Fonseca, Oleoylethanolamide enhances beta-adrenergic-mediated thermogenesis and white-to-brown adipocyte phenotype in epididymal white adipose tissue in rat. Disease models & mechanisms 7 (2014) 129-41.

[49] J.M.A. de Jong, R.T.F. Wouters, N. Boulet, B. Cannon, J. Nedergaard, and N. Petrovic, The beta3-adrenergic receptor is dispensable for browning of adipose tissues. Am J Physiol Endocrinol Metab 312 (2017) E508-E518.

[50] F.C.B. Oliveira, E.J. Bauer, C.M. Ribeiro, S.A. Pereira, B.T.S. Beserra, S.M. Wajner, A.L. Maia, F.A.R. Neves, M.S. Coelho, and A.A. Amato, Liraglutide Activates Type 2 Deiodinase and Enhances beta3-Adrenergic-Induced Thermogenesis in Mouse Adipose Tissue. Frontiers in endocrinology 12 (2021) 803363.

[51] S. Gaetani, J. Fu, T. Cassano, P. Dipasquale, A. Romano, L. Righetti, S. Cianci, L. Laconca, E. Giannini, S. Scaccianoce, J. Mairesse, V. Cuomo, and D. Piomelli, The fat-induced satiety factor oleoylethanolamide suppresses feeding through central release of oxytocin. J Neurosci 30 (2010) 8096-101.

[52] A. Romano, T. Cassano, B. Tempesta, S. Cianci, P. Dipasquale, R. Coccurello, V. Cuomo, and S. Gaetani, The satiety signal oleoylethanolamide stimulates oxytocin neurosecretion from rat hypothalamic neurons. Peptides 49 (2013) 21-6.

[53] I. Sospedra, R. Moral, R. Escrich, M. Solanas, E. Vela, and E. Escrich, Effect of High Fat
Diets on Body Mass, Oleylethanolamide Plasma Levels and Oxytocin Expression in
Growing Rats. J Food Sci 80 (2015) H1425-31.
[54] M. Asker, J.P. Krieger, A. Liles, I.C. Tinsley, T. Borner, I. Maric, S. Doebley, C.D. Furst, S.
Borchers, F. Longo, Y.R. Bhat, B.C. De Jonghe, M.R. Hayes, R.P. Doyle, and K.P.
Skibicka, Peripherally restricted oxytocin is sufficient to reduce food intake and
motivation, while CNS entry is required for locomotor and taste avoidance effects.
Diabetes, obesity & metabolism 25 (2023) 856-877.
[55] Y. Iwasaki, P. Kumari, L. Wang, S. Hidema, K. Nishimori, and T. Yada, Relay of peripheral
oxytocin to central oxytocin neurons via vagal afferents for regulating feeding.
Biochemical and biophysical research communications 519 (2019) 553-558.
[56] Y. Iwasaki, Y. Maejima, S. Suyama, M. Yoshida, T. Arai, K. Katsurada, P. Kumari, H.
Nakabayashi, M. Kakei, and T. Yada, Peripheral oxytocin activates vagal afferent neurons
to suppress feeding in normal and leptin-resistant mice: a route for ameliorating
hyperphagia and obesity. Am J Physiol Regul Integr Comp Physiol 308 (2015) R360-9.
[57] D.I. Brierley, M.K. Holt, A. Singh, A. de Araujo, M. McDougle, M. Vergara, M.H. Afaghani,
S.J. Lee, K. Scott, C. Maske, W. Langhans, E. Krause, A. de Kloet, F.M. Gribble, F.
Reimann, L. Rinaman, G. de Lartigue, and S. Trapp, Central and peripheral GLP-1 systems
independently suppress eating. Nat Metab 3 (2021) 258-273.
[58] J.M. Ho, V.T. Anekonda, B.W. Thompson, M. Zhu, R.W. Curry, B.H. Hwang, G.J. Morton,
M.W. Schwartz, D.G. Baskin, S.M. Appleyard, and J.E. Blevins, Hindbrain oxytocin
receptors contribute to the effects of circulating oxytocin on food intake in male rats.
Endocrinology 155 (2014) 2845-57.
[59] O. Erdenebayar, T. Kato, T. Kawakita, K. Kasai, Y. Kadota, K. Yoshida, T. Iwasa, and M.
Irahara, Effects of peripheral oxytocin administration on body weight, food intake,
adipocytes, and biochemical parameters in peri- and postmenopausal female rats.
Endocrine journal (2020).
[60] T. Iwasa, T. Matsuzaki, Y. Mayila, T. Kawakita, R. Yanagihara, and M. Irahara, The effects
of chronic oxytocin administration on body weight and food intake in DHT-induced PCOS
model rats. Gynecological endocrinology : the official journal of the International Society
of Gynecological Endocrinology 36 (2020) 55-60.
[61] T. Iwasa, T. Matsuzaki, Y. Mayila, R. Yanagihara, Y. Yamamoto, T. Kawakita, A. Kuwahara,
and M. Irahara, Oxytocin treatment reduced food intake and body fat and ameliorated
obesity in ovariectomized female rats. Neuropeptides 75 (2019) 49-57.
[62] L. Balazova, K. Krskova, M. Suski, V. Sisovsky, N. Hlavacova, R. Olszanecki, D. Jezova,
and S. Zorad, Metabolic effects of subchronic peripheral oxytocin administration in lean
and obese zucker rats. Journal of physiology and pharmacology : an official journal of the
Polish Physiological Society 67 (2016) 531-541.
[63] C.T. Elfers, J.E. Blevins, T.S. Salameh, E.A. Lawson, D. Silva, A. Kiselyov, and C.L. Roth,
Novel Long-Acting Oxytocin Analog with Increased Efficacy in Reducing Food Intake
and Body Weight. International journal of molecular sciences 23 (2022).
[64] C.T. Elfers, J.E. Blevins, E.A. Lawson, R. Pittner, D. Silva, A. Kiselyov, and C.L. Roth,
Robust Reductions of Body Weight and Food Intake by an Oxytocin Analog in Rats.
Frontiers in physiology 12 (2021) 726411.
[65] A. Schaffler, N. Binart, J. Scholmerich, and C. Buchler, Hypothesis paper Brain talks with
fat--evidence for a hypothalamic-pituitary-adipose axis? Neuropeptides 39 (2005) 363-7.

[66] J. Altirriba, A.L. Poher, A. Caillon, D. Arsenijevic, C. Veyrat-Durebex, J. Lyautey, A. Dulloo,
and F. Rohner-Jeanrenaud, Divergent effects of oxytocin treatment of obese diabetic mice
on adiposity and diabetes. Endocrinology 155 (2014) 4189-201.
[67] H. Yoshitomi, K. Yamazaki, S. Abe, and I. Tanaka, Differential regulation of mouse
uncoupling proteins among brown adipose tissue, white adipose tissue, and skeletal muscle
in chronic beta 3 adrenergic receptor agonist treatment. Biochemical and biophysical
research communications 253 (1998) 85-91.
[68] Y. Kasahara, K. Sato, Y. Takayanagi, H. Mizukami, K. Ozawa, S. Hidema, K.H. So, T.
Kawada, N. Inoue, I. Ikeda, S.G. Roh, K. Itoi, and K. Nishimori, Oxytocin receptor in the
hypothalamus is sufficient to rescue normal thermoregulatory function in male oxytocin
receptor knockout mice. Endocrinology 154 (2013) 4305-15.
[69] Y. Kasahara, Y. Tateishi, Y. Hiraoka, A. Otsuka, H. Mizukami, K. Ozawa, K. Sato, S.
Hidema, and K. Nishimori, Role of the Oxytocin Receptor Expressed in the Rostral
Medullary Raphe in Thermoregulation During Cold Conditions. Frontiers in endocrinology
6 (2015) 180.
[70] T. Bengtsson, B. Cannon, and J. Nedergaard, Differential adrenergic regulation of the gene
expression of the beta-adrenoceptor subtypes beta1, beta2 and beta3 in brown adipocytes.
The Biochemical journal 347 Pt 3 (2000) 643-51.
[71] H. Zhang, C. Wu, Q. Chen, X. Chen, Z. Xu, J. Wu, and D. Cai, Treatment of obesity and
diabetes using oxytocin or analogs in patients and mouse models. PloS one 8 (2013)
e61477.
[72] F. Plessow, L. Kerem, M.L. Wronski, E. Asanza, M.L. O'Donoghue, F.C. Stanford, K.T.
Eddy, T.M. Holmes, M. Misra, J.J. Thomas, F. Galbiati, M. Muhammed, A.C. Sella, K.
Hauser, S.E. Smith, K. Holman, J. Gydus, A. Aulinas, M. Vangel, B. Healy, A. Kheterpal,
M. Torriani, L.M. Holsen, M.A. Bredella, and E.A. Lawson, Intranasal Oxytocin for
Obesity. NEJM Evid 3 (2024) EVIDoa2300349.
[73] V.N. Bharadwaj, J. Meyerowitz, B. Zou, M. Klukinov, N. Yan, K. Sharma, D.J. Clark, X.
Xie, and D.C. Yeomans, Impact of Magnesium on Oxytocin Receptor Function.
Pharmaceutics 14 (2022).
[74] F.A. Antoni, and S.E. Chadio, Essential role of magnesium in oxytocin-receptor affinity and
ligand specificity. The Biochemical journal 257 (1989) 611-4.
[75] J. Niu, J. Tong, and J.E. Blevins, Oxytocin as an Anti-obesity Treatment. Front Neurosci 15
(2021) 743546.
[76] P.K. Olszewski, E.E. Noble, L. Paiva, Y. Ueta, and J.E. Blevins, Oxytocin as a Potential
Pharmacological Tool to Combat Obesity. Journal of Neuroendocrinology In press (2022).
[77] A.M. Cypess, L.S. Weiner, C. Roberts-Toler, E. Franquet Elia, S.H. Kessler, P.A. Kahn, J.
English, K. Chatman, S.A. Trauger, A. Doria, and G.M. Kolodny, Activation of human
brown adipose tissue by a beta3-adrenergic receptor agonist. Cell Metab 21 (2015) 33-8.
[78] C.M. Liu, E.A. Davis, A.N. Suarez, R.I. Wood, E.E. Noble, and S.E. Kanoski, Sex Differences
and Estrous Influences on Oxytocin Control of Food Intake. Neuroscience 447 (2020) 63-
73.
872
873
874
875

Figure legend

Figure 1A-D: Determine the dose-response effects of systemic (SC) infusion of OT (16 and 50 nmol/day) on body weight, adiposity and energy intake in DIO rats. *A*, Rats were maintained on HFD (60% kcal from fat; N=6-7/group) for approximately 5.5 months prior to being implanted with temperature transponders and allowed to recover for 1-2 weeks prior to being implanted with subcutaneous minipumps. *A*, Effect of chronic subcutaneous OT or vehicle on body weight in DIO rats; *B*, Effect of chronic subcutaneous OT or vehicle on body weight change in DIO rats; *C*, Effect of chronic subcutaneous OT or vehicle on adiposity in DIO rats; *D*, Effect of chronic subcutaneous OT or vehicle on adiposity in DIO rats. Data are expressed as mean ± SEM. *$P<0.05$, †$0.05<P<0.1$ OT vs. vehicle.

Figure 2A-E: Effect of chronic 4V OT infusions (16 nmol/day) and systemic beta-3 receptor agonist (CL 31643) administration (0.5 mg/kg) on body weight, body adiposity and energy intake in male DIO rats. Ad libitum fed rats were either maintained on HFD (60% kcal from fat; N=8-10/group) for approximately 8 months prior to receiving continuous infusions of vehicle or OT (16 nmol/day) in combination with a single dose of CL 316243 (0.5 mg/kg). *A*, Change in body weight in HFD-fed DIO rats; *B*, Change in body weight gain in HFD-fed DIO rats; *C*, Change in fat mass and lean mass in HFD-fed DIO rats; *D*, Change in relative fat mass and lean mass in HFD-fed DIO rats; *E*, Weekly energy intake (kcal/week) in HFD-fed DIO rats. ↑ indicate 1x daily injections. Different letters denote significant differences between treatments *B*, Colored bars represent specific group comparisons vs vehicle. Data are expressed as mean ± SEM. *$P<0.05$, **P=0.05, †$0.05<P<0.1$ vs. vehicle or baseline (pre-treatment; **Fig. 3A**).

Figure 3A-H: Effect of chronic 4V OT infusions (16 nmol/day) and systemic beta-3 receptor agonist (CL 31643) administration (0.5 mg/kg) on adipocyte size in IWAT and EWAT in male DIO rats. Adipocyte size was analyzed using ImageJ. Images were taken from fixed (4% PFA) paraffin embedded sections (5 μm) containing IWAT (A-D) or EWAT (E-H) in HFD-fed rats treated with 4V OT (16 nmol/day) or 4V vehicle in combination with IP CL 316243 (0.5 mg/kg) or IP vehicle. A/E, Veh/Veh. B/F, OT/Veh. C/G, Veh/CL 316243. D/H, OT-CL 316243; (A–H) all visualized at 100X magnification.

Figure 4A-B: Effect of chronic 4V OT infusions (16 nmol/day) and systemic beta-3 receptor agonist (CL 31643) administration (0.5 mg/kg) on adipocyte size in IWAT and EWAT in male DIO rats. *A*, Adipocyte size (pixel^2) was measured in IWAT from rats that received chronic 4V infusion of OT (16 nmol/day) or vehicle in combination with daily CL 316243 (0.5 mg/kg) or vehicle treatment (N=4-9/group). *B*, Adipocyte size was measured in EWAT from rats that received chronic 4V infusion of OT (16 nmol/day) or vehicle in combination with daily CL 316243 (0.5 mg/kg) or vehicle treatment (N=5-9/group). Data are expressed as mean ± SEM. *$P<0.05$ OT vs. vehicle.

Table 1. Plasma measurements following systemic (SC) infusions of OT (16 and 50
nmol/day) or vehicle DIO rats. Data are expressed as mean ± SEM. *$P<0.05$ OT vs.
vehicle (N=6-7/group).

Table 2A-C. T$_{IBAT}$ measurements following acute systemic administration of the beta-3
receptor agonist (CL 316243) or vehicle in male DIO rats. *A*, injection day 1 (0.5 mg/kg)
and *B*, injection day 22 (0.5 mg/kg). Data are expressed as mean ± SEM. *P<0.05 vs
VEH; †$0.05<P<0.1$ vs VEH.

Table 3. Plasma measurements following 4V infusions of OT (16 nmol/day), CL 316243
(0.5 mg/kg) or OT (16 nmol/day) + CL 316243 (0.5 mg/kg) in DIO rats. Data are expressed
as mean ± SEM. Different letters denote significant differences between treatments
($P<0.05$).

Table 4. Changes in IBAT mRNA expression following chronic 4V OT (16 nmol/day) and
systemic CL 316243 (0.5 mg/kg) treatment in male DIO rats. Data are expressed as mean
± SEM. Different letters denote significant differences between treatments ($P<0.05$).

Table 5. T$_{IBAT}$ measurements following acute systemic administration of the beta-3
receptor agonist (CL 316243) or vehicle in male DIO rats. *A*, injection day 23 prior to
euthanasia (0.5 mg/kg), euthanasia day. Data are expressed as mean ± SEM. *P<0.05
vs VEH; †$0.05<P<0.1$ vs VEH.

943
944

FIGURE 1A

FIGURE 1B

FIGURE 1C

FIGURE 1D

FIGURE 2A

FIGURE 2B

FIGURE 2C

FIGURE 2D

FIGURE 2E

IWAT

A. Veh/Veh

B. Veh/CL 316243

C. OT/Veh

D. OT/CL 316243

EWAT

E. Veh/Veh

F. Veh/CL 316243

G. OT/Veh

H. OT/CL 316243

FIGURE 3

FIGURE 4A

FIGURE 4B

Table 1. Plasma Measurements Following SC OT (16 and 50 nmol/day)

SC	VEH	OT (16 nmol/day)	OT (50 nmol/day)
Leptin (ng/mL)	65.9 ± 7.8^{a}	71.3 ± 8.2^{a}	59.6 ± 7.7^{a}
Insulin (ng/mL)	4.7 ± 1.7^{a}	3.4 ± 0.9^{a}	4.7 ± 1.3^{a}
Blood Glucose (mg/dL)	137.3 ± 4.5^{a}	160.4 ± 8.3^{b}	157.7 ± 7.3^{ab}
Glucose (mg/dL)	194.6 ± 5.7^{a}	202.2 ± 8.0^{a}	205 ± 12.5^{a}
Total Cholesterol (mg/dL)	108.1 ± 10.4^{a}	118.4 ± 13.4^{a}	105.5 ± 11.5^{a}

Different letters denote significant differences between treatments

Shared letters are not significantly different from one another

N=6-7/group

Table 2A. Changes in T_{IBAT} Following Systemic OT +/- CL 316243 Treatment in Male DIO Rats (Injection Day 1)

SC	IP	0 min	15 min	30 min	45 min	60 min
		Temp (C°)	Temp (C°)	Temp (C°)	Temp (C°)	Temp (C°)
VEH	VEH	37.1±0.4	37.3±0.8	37.7±0.3	37.5±0.3	37.4±0.3
VEH	CL 316243	36.8±0.2	38.3±0.2†	38.6±0.2*	38.5±0.2*	38.5±0.2*
OT	VEH	36.6±0.3	37.7±0.2	37.3±0.2	37.3±0.2	37.0±0.2
OT	CL 316243	36.6±0.2	38.2±0.2†	38.4±0.2*	38.3±0.2*	38.1±0.1*

*P<0.05 vs VEH

N=8-12/group

Table 2B. Changes in T_{IBAT} Following Systemic OT +/- CL 316243 Treatment in Male DIO Rats (Injection Day 22)

SC	IP	0 min	15 min	30 min	45 min	60 min
		Temp (C°)	Temp (C°)	Temp (C°)	Temp (C°)	Temp (C°)
VEH	VEH	36.5±0.3	37.7±0.2	37.1±0.3	37.0±0.2	37.1±0.3
VEH	CL 316243	36.8±0.2	38.5±0.2*	38.3±0.2*	38.2±0.2*	38.1±0.2*
OT	VEH	36.2±0.2	37.4±0.1	37.0±0.2	36.9±0.2	37.0±0.2
OT	CL 316243	36.4±0.1	38.1±0.2†	37.9±0.1*	37.7±0.2*	37.7±0.2†

*P<0.05 vs VEH

N=10-12/group

Table 3. Plasma Measurements Following SC OT +/- CL 316243 in DIO Rats

SC	VEH/VEH	OT/VEH	VEH/CL 316243	OT/CL 316243
Leptin (ng/mL)	47.2 ± 6.0^a	45.9 ± 4.9^a	25.8 ± 3.4^b	22.0 ± 2.4^b
Insulin (ng/mL)	10.9 ± 2.3^{ac}	11.7 ± 1.5^a	6.6 ± 1.2^{bc}	4.3 ± 0.8^b
FGF-21 (pg/mL)	187.3 ± 22.5^a	163.9 ± 9.1^a	87.7 ± 8.0^b	101.5 ± 11.6^c
Irisin (μg/mL)	3.9 ± 0.3^a	4.5 ± 0.6^a	4.6 ± 0.4^a	4.4 ± 0.4^a
Adiponectin (μg/mL)	6.4 ± 0.6^{ac}	6.3 ± 0.6^{ac}	7.3 ± 0.6^{bc}	8.7 ± 0.5^b
Blood Glucose (mg/dL)	179.3 ± 24.5^a	174.9 ± 9.3^a	150 ± 3.7^{ab}	138 ± 4.5^b
Glucose (mg/dL)	206.2 ± 10.7^a	213.1 ± 12.2^a	173.2 ± 7.7^b	180.9 ± 6.3^b
Triglycerides (mg/dL)	81.5 ± 13.7^{ab}	97.5 ± 33.1^a	39.8 ± 3.9^b	47.8 ± 3.3^{ab}
FFA (mEq/L)	0.5 ± 0.1^a	0.5 ± 0.1^a	0.2 ± 0.02^b	0.3 ± 0.03^{ab}
Total Cholesterol (mg/dL)	100.0 ± 6.0^a	108.1 ± 9.7^a	94.5 ± 5.1^a	107.0 ± 10.2^a

Different letters denote significant differences between treatments

Shared letters are not significantly different from one another

N=10-12/group

Table 4A. Changes in IBAT mRNA Expression Following SC OT and CL 316243 Treatment in Male DIO Rats

SC Treatment	VEH/VEH	OT/VEH	VEH/CL 316243	OT/CL 316243
IBAT				
Adrb1	1.0 ± 0.3^a	1.0 ± 0.1^a	0.8 ± 0.1^a	0.9 ± 0.4^a
Adrb3	1.0 ± 0.1^a	0.9 ± 0.2^a	0.5 ± 0.1^b	0.5 ± 0.04^b
Ucp1	1.0 ± 0.1^a	1.0 ± 0.1^a	1.6 ± 0.1^b	1.6 ± 0.1^b
Cidea	1.0 ± 0.1^a	1.0 ± 0.1^a	1.0 ± 0.1^a	0.9 ± 0.1^a
Dio2	1.0 ± 0.1^a	1.5 ± 0.2^a	3.3 ± 0.4^b	3.5 ± 0.3^b
Gpr120	1.0 ± 0.2^a	0.9 ± 0.2^a	19.8 ± 2.4^b	15.9 ± 2.1^b
Prdm16	1.0 ± 0.1^a	0.9 ± 0.1^a	0.8 ± 0.1^a	0.9 ± 0.1^a
Ppargc1a	1.0 ± 0.05^a	1.0 ± 0.2^a	4.8 ± 0.4^b	5.9 ± 0.5^c

Different letters denote significant differences between treatments

Shared letters are not significantly different from one another

N=10-12/group

Table 4B. Changes in IWAT mRNA Expression Following SC OT and CL 316243 Treatment in Male DIO Rats

SC Treatment	VEH/VEH	OT/VEH	VEH/CL 316243	OT/CL 316243
IWAT				
Adrb3	1.0 ± 0.4^a	1.1 ± 0.6^a	1.0 ± 0.3^a	0.5 ± 0.2^a
Ucp1	1.0 ± 0.4^a	0.3 ± 0.1^a	0.5 ± 0.2^a	1.7 ± 0.8^a
Dio2	1.0 ± 0.3^a	1.0 ± 0.2^a	0.6 ± 0.2^a	0.9 ± 0.3^a
Ppargc1a	1.0 ± 0.7^a	0.5 ± 0.4^a	1.5 ± 1.0^a	0.04 ± 0.02^a

Different letters denote significant differences between treatments

Shared letters are not significantly different from one another

N=9-12/group

Table 5. Changes in T_{IBAT} Following Systemic OT +/- CL 316243 Treatment in Male DIO Rats (Euthanasia Day)

SC	IP	0 min Temp (C°)	15 min Temp (C°)	30 min Temp (C°)	45 min Temp (C°)	60 min Temp (C°)
VEH	VEH	36.7±.0.3	37.5±.0.2	37.4±.0.2	37.5±.0.2	37.2±.0.7
VEH	CL 316243	36.7±.0.1	38.3±.0.3*	38.5±.0.2*	38.5±.0.2*	38.3±.0.2*
OT	VEH	36.3±.0.1	37.1±.0.3	37.5±.0.2	37.5±.0.2	37.1±.0.3
OT	CL 316243	36.7±.0.2	37.8±.0.2*	38.1±.0.2*	38.0±.0.2†	37.8±.0.2*

*P<0.05 vs VEH

†0.05<P<0.1

N=9-12/group