1 RESEARCH ARTICLE

2 *Running title:* Tailocins of Soft Rot *Pectobacteriaceae* bacteria

4 Tailocin-mediated interactions among Soft Rot *Pectobacteriaceae*

Marcin Borowicz [a], Dorota M. Krzyżanowska [a], Marta Sobolewska [a], Magdalena Narajczyk [b], Inez

Mruk [c], Paulina Czaplewska [c], Jacques Pédron [d], Marie-Anne Barny [d], Pierre Yves Canto [d], Joanna

Dziadkowiec [e] and Robert Czajkowski [a,*]

[a] Laboratory of Biologically Active Compounds, Intercollegiate Faculty of Biotechnology of UG and

MUG, University of Gdansk, A. Abrahama 58, 80-307 Gdańsk, Poland;

[b] Bioimaging Laboratory, Faculty of Biology, University of Gdansk, Wita Stwosza 59, 80-308

Gdansk, Poland;

[c] Laboratory of Mass Spectrometry-Core Facility Laboratories, Intercollegiate Faculty of

Biotechnology of UG and MUG, University of Gdansk, Antoniego Abrahama 58, 80-307 Gdansk,

Poland;

[d] Institute of Ecology and Environmental Sciences of Paris, Sorbonne Université, INRAE, IRD,

CNRS, UPEC, UMR 7618, Paris, France;

[e] The Njord Centre, Departments of Geosciences and Physics, University of Oslo, Blindern, 0316

Oslo, Norway;

* Correspondence: robert.czajkowski@ug.edu.pl; phone: 0048 58 523 6333

Abstract

Bacteria carry phage-derived elements within their genomes, some of which can produce phage-like particles (tailocins) used as weapons to kill kin strains in response to environmental conditions. This study investigates the production and activity of tailocins by plant pathogenic bacteria: *Pectobacterium*, *Dickeya,* and *Musicola* genera, which compete for niche, providing an attractive model to study the ecological role of tailocins. Microscopy revealed that most analyzed strains (88%) produced tailocins. Tailocin-mediated killing interactions were assessed across 351 strain pairs, showing that *Dickeya* spp. had a higher likelihood of killing neighbors (57.1%) than *Pectobacterium* spp. (21.6%). Additionally, *Dickeya* spp. strains exhibited broader phylogenetic killing, targeting both *Pectobacterium* spp. and *Musicola* sp., while *Pectobacterium* spp. tailocins were genus-specific. Mutual killing was observed in 33.9% of interactions, predominantly within *Dickeya* spp. Although tailocins were morphologically indistinguishable between producers, genomic analyses identified conserved clusters having distinct differences between *Pectobacterium* spp. and *Dickeya* spp tailocins. This suggests different origins of these particles. Induction experiments demonstrated that tailocin production was boosted by hydrogen peroxide, supporting the role of these particles in bacteria-bacteria competition during infection. Tailocins were detectable in infected potato tissue but not in river water, highlighting the particular ecological relevance of tailocins in plant environments.

Keywords:

bacteriophage, phage tail-like particles, hydrogen peroxide, inter-genus, interactions, *Erwinia chrysanthemi*, *Erwinia carotovora*, prophage

Introduction

In highly competitive and resource-limited environments that bacteria occupy, their survival and fitness often depend on their ability to outcompete neighboring rival cells (Bauer et al., 2018;Wagner, 2022). To gain a competitive edge, bacteria have evolved diverse strategies, including producing and utilizing phage tail-like particles (tailocins) that can be lethal to other bacteria. These structures directly resemble the tails of bacteriophages but function independently from viral infection and replication (Booth et al., 2023). These particles are highly specialized killing instruments that, by puncturing the cell envelope, lead to the disruption of the cell membrane and the death of the targeted cell (Scholl, 2017). Until recently, the utilization of phage tail-like particles has been primarily studied in *Pseudomonas* species (Carim et al., 2021). However, there are also reports of tailocins isolated from various other Gram-negative and Gram-positive bacterial species, including human, animal, and plant pathogens, as well as saprophytic bacteria that inhabit a wide range of environments (Backman et al., 2024a).

We previously characterized a new type of phage tail-like particle produced by the plant pathogenic bacterium, *Dickeya dadantii* reference strain 3937 (Borowicz et al., 2023). This tailocin, dickeyocin P2D1, was tolerant of most environmental conditions and exhibited broad killing potential against members of different *Dickeya* species but was harmless to *Pectobacterium* species and non-toxic to *Caenorhabditis elegans.* We postulated that the dickeyocin P2D1 targets were restricted to phylogenetically closely related strains. Only one other phage tail-like particle, carotovoricin Er from *Pectobacterium carotovorum* Er, was described in detail (Yamada et al., 2006). The sequence of Er differs from that of dickeyocin P2D1, indicating a distinct evolutionary origin of these two tailocins (Yamada et al., 2006). Besides these limited studies, little attention has been paid to the presence and ecological role of tailocins in SRP bacteria.

SRP bacteria (consisting of *Pectobacterium* spp., *Dickeya* spp., and *Musicola* spp., previously collectively described as pectinolytic *Erwinia* spp.) are a particularly appropriate group of bacteria for studying the environmental role of phage tail-like particles in the environment. *Pectobacterium* spp., *Dickeya* spp., and *Musicola* spp. belong to the same species complex and

are often found together in infected hosts. Consequently, due to their metabolic similarities and

overlapping niches, they are known to compete against each other (Toth et al., 2021a).

Furthermore, these bacteria occupy various ecological niches, including host and non-host plants,

natural and agricultural soils, rainwater, surface water, sewage, and insects (Van Gijsegem et al.,

2021). In these diverse environments, SRP bacteria often encounter a diverse array of other

species, both closely and distantly related, but their competitive strategies in these alternative

biomes remain unknown (Bellieny-Rabelo et al., 2019).

Our understanding of the mechanisms of competition in SRP bacteria mediated by phage

tail-like particles remains poorly understood (Marquez-Villavicencio et al., 2011;Hugouvieux-Cotte-

Pattat, 2016;Hugouvieux-Cotte-Pattat et al., 2023). Specifically, no systematic studies have been

conducted to assess whether the production of tailocins is common in this group of bacteria and

how it contributes to their overall ecological success (Borowicz et al., 2023). Addressing this gap

remains important for understanding how plant-pathogenic bacteria establish and thrive in diverse

environments, particularly in those niches where microbial competition between kin strains is

expected to be intense.

This study aimed to assess the expression of phage tail-like particles (tailocins) in SRP

strains obtained from one selected environment – the Durance River in France (Ben Moussa et al.,

2022). Although river water is not recognized as a primary environment where SRP bacteria thrive,

this niche links the several distinct ecosystems that support the growth and survival of SRP. The

tested SRP strains could potentially interact in water or be transmitted to a common plant host,

where they might interact during infection (Van Gijsegem et al., 2021). We aimed to characterize

interactions among SRP bacteria guided by phage tail-like particles to understand better how these

pathogens share the environment and whether they compete for niche during infection. For this,

we characterized the SRP tailocins and identified gene clusters associated with their synthesis,

thus determining the prevalence of tailocin production among environmental SRP strains and their

apparent role in intragenus competition among the diverse strains that coexist in this aquatic

habitat.

Materials and Methods

Bacterial strains and culture conditions

Bacterial strains included in this work are listed in Supplementary Table S1. The pool comprised 25 SRP isolates from the waters of Durance River (France), collected between 2015 and 2017 (Ben Moussa et al., 2022), as well as two reference strains: *Dickeya dadantii* 3937 (Pritchard et al., 2013) and *Musicola paradisiaca* NCPPB 2511 (Hugouvieux-Cotte-Pattat et al., 2021). The latter strains have been previously studied (Borowicz et al., 2023). Routine propagation of bacterial cells was performed at 28 ºC in liquid Trypticase Soy Broth (TSB, Oxoid), with agitation (120 rpm), or on solid Trypticase Soy Agar (TSA, Oxoid) under the same growth conditions (Czajkowski et al., 2010).

Induction and purification of tailocins

Tailocin particles were purified as described previously (Borowicz et al., 2023). Briefly, an overnight culture (*ca.* 16 h) of each strain was rejuvenated (1:40) in a fresh aliquot of TSB and cultured for 2.5 hours. Next, tailocin production was induced by the addition of 0.1 µg mL^{-1} of mitomycin C (Abcam, Poland). Twenty-four hours after induction, tailocins were purified using PEG from 10 mL volumes of filtered (0.2 µm, PES membrane, GoogLab™) culture supernatants. Purified particles were stored in phosphate buffer saline (PBS), pH 7.2, at 4 °C.

Imaging

The morphology of purified tailocin particles was investigated by transmission electron microscopy (TEM) and atomic force microscopy (AFM). TEM imaging was performed as described previously (Borowicz et al., 2023) using a Tecnai Spirit BioTWIN microscope (FEI). AFM imaging was performed using JPK NanoWizard® 4 (NanoScience), in non-destructive quantitative imaging (QI™) force spectroscopy mode, employing the SCANASYST-AIR probes (f0 7.0 kHz, diameter <12 nm, k: 0.4 N/m) and the SCANASYST-FLUID+ probes (f0 150 kHz, diameter <12 nm, k: 0.7

N/m). Spring constants and sensitivity of the probes were calibrated using the thermal calibration

method. Before imaging, the samples were deposited on freshly cleaved mica surfaces and air-

dried. Images were processed using JPK Data Processing software. Dimensions of tailocins were

calculated based on TEM images. Tailocins from a given strain were considered for measurement

only if more than 4 individual particle images were available for analysis. Analysis was

accomplished on 13 out of 18 *Pectobacterium* spp. strains, with a median of 10 particles measured

per strain, and 7 *Dickeya* spp., with a median of 17 particles measured per strain.

Killing and sensitivity assays

Twenty-seven SRP strains, including 25 water isolates (Supplementary Table S1), were tested in

pairs to determine the target range (killing spectrum) of the tailocin produced by each group

member. Likewise, the sensitivity of strains to the tailocins produced by all other strains in the

analyzed pool was assessed. Both the killing spectrum and sensitivity were tested using the spot

assay as previously described (Hockett and Baltrus, 2017;Yao et al., 2017;Borowicz et al., 2023).

Each combination was tested in duplicate, and the entire experiment was performed twice using

independently-obtained batches of tailocins.

SDS-PAGE, ESI LC-MS/MS, and mapping of identified proteins to genomic *loci*

Proteins in PEG-purified tailocin preparations were separated by SDS-PAGE, fragmented, and the

resulting peptides sequenced using mass spectrometry, as described in detail previously

(Golebiowski et al., 2022;Borowicz et al., 2023). Briefly, bands were excised from the gel and

subjected to in-gel trypsin digestion, and the peptides were eluted, cleaned up, and concentrated

(Schmidt and Sinz, 2017;Goldman et al., 2019). The peptides were then analyzed by ESI LC-

MS/MS. This included separating peptides on an Eksigent microLC column ChromXP C18CL (3

μm, 120 Å, 150 × 0.3 mm). The samples were placed onto the column with a CTC Pal Autosampler

(CTC Analytics AG, Zwinger, Switzerland), and the injection volume was 5 μl. Solvents A and B

consisted of 0.1% (v/v) formic acid (FA) in water and acetonitrile, respectively. LC gradient

parameters: 18–50% solvent B in 30 min (Fiołka et al., 2019). The mass spectra with Triple Tof

5600+ mass spectrometer with DuoSpray Ion Source (AB SCIEX, Framingham, MA, USA) enabled

the identification of proteins based on the obtained fragmentation spectra using the Peaks Studio

11 (Bioinformatic Solutions) and the appropriate protein database (Supplementary Table S1) (1%

FDR). The annotations of the identified proteins were searched for phage-associated functions,

and the encoding genes were mapped to the genomes of the tested strains. Tailocin-encoding

regions were identified based on the clustering of the identified phage/tailocin-related genes within

the genomes of the analyzed strains and previous studies concerning the structure of the P2D1

tailocin cluster (Borowicz et al., 2023). As the full annotated sequence of the carotovoricin Er

cluster is unavailable, the originally uploaded sequence (accession: AB017338.2) (Yamada et al.,

2006) annotated by us using RAST (https://rast.nmpdr.org/rast.cgi) (Brettin et al., 2015) was used

for cluster synteny analysis.

Phylogenetic and bioinformatic analyses

Synteny of tailocin clusters

Gene clusters encoding tailocins were extracted from the genomes of *Pectobacterium*, *Dickeya,*

and *Musicola* species except for *Pectobacterium odoriferum* A122-S21-F16, where the putative

cluster was split across multiple contigs. For *Pectobacterium* spp. the genomic region containing

the tailocin cluster was defined as the region between the genes annotated as *tolC* (locus tag in *P.*

versatile A69-S13-O15: EG331_00440) and *ybiB* (locus tag in *P. versatile* A69-S13-O15:

EG331_00565). In contrast, for *Dickeya* species and a single *Musicola* sp. strain, the genomic

region containing the tailocin cluster was identified as the region between the genes annotated as

methyl-accepting chemotaxis protein (locus tag in *D. dadantii* 3937: DDA3937_RS1200) and *guaD*

(locus tag in *D. dadantii* 3937: DDA3937_RS12145).

Accession numbers of genomes from which the tailocin regions were extracted are listed in

Supplementary Table 1. The extracted sequences were then subjected to synteny analysis and

visualization using the Clinker tool (Gilchrist and Chooi, 2021). The analysis was performed

separately for *Pectobacterium* and *Dickeya* species, along with a single strain of *Musicola* sp.

Genes within the clusters were color-coded based on their putative roles in tailocin production. In

the visualization, links between genes indicate sequence identity percentages greater than 50%.

The order of strains in the visualization was determined according to their phylogenetic

relationships. Synteny analysis was based on *Pectobacterium versatile* A73-S18-O15 for

Pectobacterium spp. and *Dickeya dadantii* 3937 for *Dickeya* spp.

Multilocus sequence analysis

A subset of strains was used for phylogenetic analysis. The selected strains possessed the

following features: (1) the tailocin cluster had to be present on a single genomic contig; (2) genes

annotated as fiber and sheath had to be present within the region identified as the tailocin cluster;

and (3) the strain had to demonstrate the ability to target at least one other analyzed strain through

tailocin interaction. Orthologous sequences were clustered into homologous families using the

SiLix software package v1.2.9 (Miele et al., 2011) with an 80% identity threshold and at least 80%

overlap. Orthologous sequences (658 coding sequences) were concatenated, aligned using

MUSCLE (Edgar, 2004) software v5.1, and filtered using the GBLOCK tool (Castresana, 2000).

The alignments were used to build a phylogenetic tree with the BioNJ algorithm with SeaView

software v5.0.5 (Gouy et al., 2010), with 200 bootstrap replications.

Phenotype-based dendrogram

Agglomerative clustering (Nielsen, 2016) was applied to prepare a dendrogram for tested strains

based on the similarity of their killing spectra. The ability to kill a given strain was assigned the

value of 1, and the lack of this ability was noted as 0. The scikit-learn (Virtanen et al., 2020) and

SciPy (Virtanen et al., 2020) libraries were used for clustering and dendrogram plotting,

respectively. A Phyton script for implementing the above libraries was written with the assistance of

ChatGPT-4o (OpenAI) (Supplementary Dataset 3).

Prevalence of carotovoricin and dickeyocin clusters in SRP complete genomes

The carotovorocin cluster was blasted against 116 *Pectobacterium* spp. complete, high-quality

genomes available on the NCBI database in June 2024 using the NCBI Megablast algorithm with

default parameters (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The presence of the cluster was

defined according to the 70% sequence coverage and identity thresholds. The available sequence

of the originally described carotovoricin Er cluster was incomplete (accession: AB017338.2).

Therefore, as a reference carotovoricin sequence for *Pectobacterium* spp. we used the full

nucleotide sequence of the carotovoricin cluster from *P. versatile* A73-S18-O15 (19,528 bp.). For

Dickeya spp. and the single *Musicola* sp. strain, the full nucleotide sequence of the dickeyocin

cluster of *D. dadantii* 3937 strain (26,580 bp) was blasted against a single *Musicola* sp. and 74

Dickeya spp. complete genomes available on the NCBI database in June 2024. Results were

interpreted based on analogous thresholds as described for P2D1 tailocin.

Detection of tailocins in river water and potato tubers

To determine whether SRPs produce detectable levels of tailocins in river water and potato tuber

tissue, we inoculated these environments with bacteria. After incubation, the tailocins were purified

and spot tests were conducted as outlined below. Each experiment was performed twice, and

detection was considered positive when all replicates gave positive results. To prepare the

inoculation suspensions, the six strains *P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-

F17, *P. aquaticum* A212-S19-A16, *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, and *D. zeae*

A586-S18-A17, were grown in TSB (Oxoid) for 24 h at 28 °C. The cells were harvested by

centrifugation (4200 RCF, 5 min) and resuspended in PBS buffer. Turbidity of the suspension was

adjusted to 0.06 McFarland (approx. 10^6 CFU mL^{-1}). Five ml of filter-sterilized (0.2 μm, PES

membrane) and autoclaved Durance River water was inoculated with 50 μl of bacterial suspension,

three replicates per strain. Uninoculated water was used as a control. Samples were incubated for

72 h at 15 °C with shaking (120 rpm) to mimic natural conditions. Following incubation, 2 ml of

water from each sample was centrifuged to remove bacterial cells (8000 RCF, 10 min), and the

supernatant was processed for purification and detection of tailocins. Tubers cv. Gala were surface

sterilized by immersion for 20 min in 5% commercial bleach (ACE, Procter and Gamble), followed

by a double rinse in distilled water and air drying under laminar flow. Each tuber was inoculated by

inserting a pipette tip containing 50 µl of the test suspension (Krzyzanowska et al., 2019). Tubers

inoculated with PBS buffer alone were used as a negative control. Three potato tubers were tested

per treatment. Inoculated tubers were placed in humid boxes (85 to 90% relative humidity).

Samples were incubated at 28°C for 72 h to enable the development of soft rot symptoms. The

tubers were then cut at the inoculation site. Tissue macerated by bacteria was excised, weighed,

and placed in homogenization bags (Bioreba). PBS buffer supplemented with 0.02%

diethyldithiocarbamic acid (DIECA) (Sigma-Aldrich) was added to the samples (1:2, w/v)

(Perombelon and van Der Wolf, 2002). The samples were homogenized, and 1 mL aliquots were

diluted 1:1 with PBS 0.02% DIECA in new tubes. Bacteria and plant debris were pelleted by

centrifugation (8000 RCF, 10 min, RT), and the supernatant was processed for purification and

detection of tailocins. The supernatants from both experimental setups were filtered (0.2 µm, PES

membrane) and transferred to new tubes containing PEG-8000 (final concentration 10 % m/v).

Samples were incubated overnight at 4°C with shaking. The putative tailocins were collected by

centrifugation (1 h, 16 000 RCF, 4°C) and resuspended in 1/20 volume of the initial sample in PBS.

Tailocins were detected using a spot test (Hockett and Baltrus, 2017;Yao et al., 2017). Five µL

aliquots of purified tailocins were spotted on soft top agar plates, each inoculated with a strain

sensitive to tailocins of the six investigated SRPs (Supplementary Table S2).

Semiquantitative assessment of tailocin production depending on the type of inductive

factor

We investigated whether hydrogen peroxide can induce the production of tailocins in SRPs in a

way similar to mitomycin C. The experiment was conducted on the same group of six strains tested

for tailocin production in river water and potato tuber tissue. Tailocin production was induced in

rejuvenated cells by the addition of 0.1 µg mL^{-1} mitomycin C (Abcam, Poland) or 0.003 % (0.88

mM) hydrogen peroxide (Laboratorium Galenowe Olsztyn Sp. z o. o., Poland) and cultures with no

inductive factor were used as controls. Twenty-four hours post-induction, tailocins were PEG-

purified from 2 mL volumes of filtered (0.2 µm, PES membrane) culture supernatants and

resuspended in 1/10 of the initial volume in PBS (Borowicz et al., 2023). The concentration of

tailocins was estimated by a semiquantitative spot test (Yao et al., 2017;Borowicz et al., 2023)

using tailocin-susceptible strains (Supplementary Table S2). The reciprocal of the highest dilution

causing a visible plaque was defined as the relative activity in arbitrary units (=1 AU), and the AU

per mL of culture was calculated. The experiment was conducted twice, with tree replicates per

combination.

Competition assay *in vitro*

Competition assays were performed using a spontaneous rifampicin-resistant mutant of *D. solani*

A623-S20-A17 selected on TSA with 50 µg mL^{-1} of rifampicin (Sigma-Aldrich)(Glandorf et al.,

1992;Berg et al., 2007) and designated *Ds* RIF. This strain was tested in an *in vitro* competition

assay against three sets of competitor strains. Each set comprised two *Dickeya* spp. strains which

either: 1) were killed by the tailocins of *D. solani* A623-S20-A17 (*D. dianthicola* A260-S21-A16 and

D. zeae A661-S21-A17), 2) produced tailocins killing *D. solani* A623-S20-A17 (*D. zeae* A586-S18-

A17 and *D. oryzae* A3-S1-M15) or 3) were killed by the tailocins of *D. solani* A623-S20-A17 but at

the same time produced tailocins targeting this strain (mutual killing)(*D. dadantii* 3937 and *D.*

chrysanthemi A604-S21-A17). All strains were grown overnight in 0.1 TSB medium. Next, the cells

were harvested by centrifugation (3500 RCR, 5 min.), and the pellet was resuspended in a fresh

medium to obtain a turbidity of 0.5 McF. This equaled, on average, 1.4×10^8 CFU ml^{-1} ($\pm 4.5 \times 10^7$),

verified by dilution plating. 150 µl of 0.1 TSB in a single well of a 96-well plate was inoculated (1:1,

25:25 µl) with the suspension of *Ds* RIF and one of the competing strains to ensure the strains had

an even start and avoid tailocins carryover from the initial overnight cultures. Each co-culture was

tested under two conditions: 0.1 TSB alone or 0.1 TSB with 0.003 % (0.88 mM) H$_2$O$_2$ (tailocin-

inductive stress). The cultures were incubated for 22 h at 28°C, with orbital shaking, in the

EPOCH2 reader (BioTek). Culture from each well was dilution-plated on two media types: TSB

agar (for counting the total number of CFUs) and TSB agar with 50 µg mL^{-1} of rifampicin (for

counting the fraction of *Ds* RIF). *Ds* RIF was tested against the wild-type (*wt*) variant of the strain

as a control of the mutant's fitness under the applied experimental conditions. Moreover, the ability

of *Ds* RIF to produce tailocins was verified in a spot assay, as described above. The competition

experiment was performed twice, with two separate co-cultures (replicates) per experiment.

Results

Most *Pectobacterium* spp. and *Dickeya* spp. strains produce phage tail-like particles

TEM imaging of preparations obtained from mitomycin C-induced cultures revealed that 22 out of

25 studied SRP isolates (88%) produce phage tail-like particles (tailocins) (Fig.1). The three

exceptions were *Pectobacterium brasiliense* A143-S20-M16, *Pectobacterium atrosepticum* A597-

S4-A17, and *Dickeya dianthicola* A260-S21-A16. All tailocins had similar morphologies in TEM and

AFM images, regardless of whether they were derived from *Dickeya/Musicola* or *Pectobacterium*

spp. (Fig. 1 and Fig. 2). The average length of tailocins produced by *Pectobacterium* spp. was

163±15 nm. Tailocins produced by *Dickeya/Musicola* spp. were significantly shorter, measuring

153±20 nm (Mann-Whitney test, p<0.05). Similarly, tailocins produced by *Pectobacterium* spp.

were, on average, 2 nm wider than those of *Dickeya/Musicola* species, with the dimensions of

26±2 nm and 24±2 nm, respectively, with the difference being statistically significant (Mann-

Whitney test, p<0.05). Three-dimensional images generated based on AFM scans for seven

representative tailocin particles are shown in Fig. 2.

The susceptible targets of SRP phage tail-like particles vary depending on the producing strain, with several exhibiting inter-genus toxicity

We evaluated the breadth of strains killed by phage tail-like particles produced by the SRP strains.

In total, 702 unique tailocin-mediated interactions (producer-target combinations) were examined

for the 351 unique pairs (individual producer vs. individual target) of SRPs tested (Fig. 3). The

incidence of a strain being susceptible to a particular tailocin was 16.8% (Table 1). This killing

incidence was higher among *Dickeya* strains than among *Pectobacterium* strains (57.1% and

21.6%, respectively) (Table 1). Strains *P. versatile* A73-S18-015 and *P. polaris* A641-S16-A17

were particularly susceptible to tailocins, being killed by those from 8 different strains. In contrast, *P. carotovorum* A280-S2-N16 was resistant to tailocins produced by all strains tested. Twenty-one of the 25 strains produced tailocins that killed at least one strain from the 27 SRPs. Of the 4 strains not exhibiting any killing activity, three (*P. brasiliense* A143-S20-M16, *P. atrosepticum* A597-S4-A17, and *D. dianthicola* A260-S21-A16) lacked visible tailocins in TEM analyses (Fig. 1), while *P. peruviense* A350-S18-N16 produces tailocins (as evidenced by TEM microscopy), these phage tail-like particles did not target any of the 26 SRPs tested (Fig. 3). In contrast, *D. oryzae* A3-S1-M15 and *P. odoriferum* A122-S21-F16 produced tailocins with the broadest target range, each killing 9 out of the 26 strains (33.3%).

We also investigated the ability of tailocins to kill stains in genera different from the producer. *Pectobacterium* strains produced tailocins targeting only other members of the genus. In contrast, 5 out of 8 tested *Dickeya* spp. (62.5%) could kill at least one representative of *Pectobacterium* sp., either a *P. versatile* strain or the closely related *P. odoriferum. M. paradisiaca* NCPPB 2511, the single representative of *Musicola* spp., strongly resembled *Dickeya* spp. both in tailocin susceptibility and the specificity of its tailocins (Fig. 3, Table 1).

Several SRP strains produced tailocins that targeted other tailocin-producing strains, resulting in a phenomenon known as 'mutual killing'

Analysis of tailocin-mediated interactions among the collection of SRPs revealed that 33.9% of the killing events were retaliated (bilateral) – in which strains could kill each other (Table 1). For 20 of the 351 pairs of SRPs tested (5.7%), a given strain pair was mutually inhibitory - a phenomenon we designate 'mutual killing'. Mutual killing was more common within *Dickeya* spp. (65.6% of killing events were mutual) than within *Pectobacterium* spp. (15.2%), and we observed no mutual killing between members of *Dickeya* and *Pectobacterium* (Table 1). The mutual killing was very common (80% of killing events) in mixed pairs of *Dickeya* strains and *M. paradisiaca* NCPPB 2511, reflecting the clustering between strain NCPPB 2511 and *Dickeya* spp.

Tailocins of *Dickeya* spp. and *Pectobacterium* spp. are phylogenetically distinct

Based on the tailocin-associated protein identified by mass spectrometry, we identified gene

clusters responsible for tailocin production in the genomes of the SRP strains. Among the 25

strains examined, tailocin-encoding clusters were absent only in 2 strains - *Pectobacterium*

brasiliense A143-S20-M16 and *Pectobacterium atrosepticum* A597-S4-A17. These two strains,

together with *Dickeya dianthicola* A260-S21-A16, also lacked tailocin production by imaging.

We also performed synteny analyses of the tailocin clusters within the *Pectobacterium* and *Dickeya*

(including *Musicola*) genera (Fig. 4). The cluster in *Pectobacterium* sp. exhibited homology with

that encoding carotovoricin Er (Nguyen et al., 2002) while those in *Dickeya* species and *Musicola*

sp. were genetically similar to those encoding dickeyocin P2D1 (Borowicz et al., 2023).

Furthermore, no SRP strain possessed both carotovorocin Er and P2D1 encoding clusters.

Pectobacterium spp. strains exclusively contained the carotovorocin Er-like cluster, while *Dickeya*

spp. strains solely harbored cluster encoding P2D1-like tailocin. Thus, no strain that contains both

types of tailocins has been identified.

The overall genomic organization of the tailocin clusters was conserved within each genus (Fig. 4).

In *Pectobacterium* species (Fig. 4A), the most prominent difference was the presence of a highly

variable gene annotated as encoding a tail fiber protein/pyocin knob domain-containing protein,

that appeared to contain segments shared among different strains (data not shown). Additional,

albeit less pronounced, variability was observed in the sequence of the gene encoding a tail tape

measure protein. In *P. brasiliense* A143-S20-M16, this cluster was flanked by genes conserved

across other strains; however, instead of a complete tailocin cluster, it contained only the single

gene annotated as a mobile element protein gene. In *P. atrosepticum* A597-S4-A17, this region

included not only conserved flanking genes but also seven genes present in other clusters and one

strain-specific gene. In *Dickeya* species (and *Musicola* sp.) (Fig. 4B), the main differences between

strains included (1) the presence of strain-specific genes likely unrelated to tailocin production, (2)

the orientation of genes not directly involved in tailocin structural components (such as aromatic

acid/H+ transporters and phytase family proteins), (3) sequence variations in intergenic regions

(data not shown), and (4) sequence differences in tail fiber genes. Notably, *D. dianthicola* A260-

S21-A16, which did not produce tailocins, did not exhibit substantial differences in the genomic

region encoding tailocins in other strains. Specifically, the tailocin cluster was present in the

genome and showed high homology to that of other *Dickeya* spp. tailocin clusters.

Carotovoricin and dickeyocin clusters are common in the genomes of *Pectobacterium* spp.

and *Dickeya* spp.

Given that not all SRPs examined produced tailocins nor harbored the tailocin cluster, we

determined the prevalence of tailocin clusters across a broader range of SRP strains to assess the

frequency of this trait better. We examined the complete genomes of 116 *Pectobacterium* spp. for

the presence of the carotovoricin cluster and 74 genomes of *Dickeya* and 1 *Musicola* spp. genome

for the presence of genes encoding dickeyocin. Within the *Pectobacterium* clade, 96 genomes

(83%) harbored homologous tailocin regions, having >70% coverage and identity. Twelve

Pectobacterium genomes had partial alignment (10% of strains), and 9 (8%) showed no evidence

of the tailocin cluster (Dataset S1). Within the many *Dickeya* spp. and the single *Musicola* spp.

genomes examined, 52 (69%) had a cluster that aligned well with the dickeyocin cluster, while 16

(21%) genomes had coverage below 70%, and 7 (9%) genomes had no apparent tailocin cluster

(Dataset S2). None of the 6 complete genomes of the species *P. atrosepticum* present in the NCBI

database had a tailocin cluster.

Tailocin-killing specificity is not aligned with phylogenetic distance in SRP strains

The topology of the dendrograms of the phylogenetic tree of SRP based on MultiLocus Sequence

Analysis (MSLA) was not well aligned with that based on the patterns of tailocin killing position

(Supplementary Fig. S1). This suggests that tailocin activity may be influenced by factors other

than taxonomic relationships.

Constitutive and induced production of tailocins in *Dickeya* and *Pectobacterium* spp.,

Given that our previous study revealed the surprising apparent constitutive basal level of tailocin production in *D. dadantii* 3937 (Borowicz et al., 2023), we explored whether this was also the case in other strains. Tailocin production was thus assessed in 6 strains (3 *Pectobacterium* and 3 *Dickeya* spp.) (*P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16, *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D. zeae* A586-S18-A17). Furthermore, we wished to establish whether reactive oxygen species (ROS), such as H_2O_2, often associated with plant infection (Torres et al., 2006;Reverchon et al., 2016), could induce the production of tailocins in SRPs. Basal levels of tailocin production were observed for all tested strains even in the absence of ROS. However, the basal level of tailocins produced varied substantially between strains, with the highest basal production observed for the model strain *D. dadantii* 3937 (7×10^4 AU mL^{-1}) and the lowest for *P. aquaticum* A411-S4-F17 (5×10^2 AU mL^{-1}) (Fig. 5). The average abundance of tailocin was slightly (43%) higher for *Dickeya* than for *Pectobacterium* spp. (Mann-Whitney test, $p<0.05$) (2.8×10^4 and 1.6×10^4 AU mL^{-1} of culture, respectively).

Mitomycin C (0.1 µg mL^{-1}, control) caused a significant induction (approx. 5 to 20-fold) of tailocin production in all strains except *P. quasiaquaticum* A411-S4-F17 (Fig. 5). The highest induction was observed for *D. zeae* A586-S18-A17 (19-fold) and *D. chrysanthemi* A604-S21-A17 (20-fold). Meanwhile, the induction of tailocins by mitomycin C in *D. dadantii* 3937 was relatively low (5-fold) due to its high production in the absence of mitomycin.

Hydrogen peroxide also induced tailocin production in all strains (approx. 15 to 81-fold) (Fig. 5). For most strains, hydrogen peroxide (0.88 mM, 0.003%) conferred a higher yield of tailocins than that induced by mitomycin C (0.1 µg mL^{-1}). Notably, while no induction of tailocins by mitomycin in *P. quasiaquaticum* A411-S4-F17 was observed, the addition of hydrogen peroxide resulted in the highest increase compared to basal production levels of all strains (81-fold) (Fig. 5).

Detectable levels of phage tail-like particles are produced in rotting potato tuber tissue but not in river water

Tailocin production was explored in SRP in a variety of natural settings in addition to cultural media. Three *Pectobacterium* spp: *P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16, and three *Dickeya* spp.: *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D. zeae* (very recently reclassified as *D. parazeae* (Hugouvieux-Cotte-Pattat and Van Gijsegem, 2021)) (A586-S18-A17) were inoculated into river water (the source of the strains) as well as into potato tubers. Tailocin production was not detected in any strain in river water (Table 2) but was abundant in macerated potato tuber tissue. Tailocins were found not only in the symptomatic tissues caused by those strains that were pathogenic to potato (*P. versatile* A69-S13-O15, *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D. zeae* A586-S18-A17) but also in healthy tuber tissue surrounding the site where non-pathogenic strains (*P. aquaticum* A212-S19-A16 and *P. quasiaquaticum* A411-S4-F17) were inoculated (Table 2) (Supplementary Fig. S2).

Tailocin production is not always sufficient to ensure competitive success against susceptible neighbors

We explored the extent to which tailocin production by *D. solani* A623-S20-A17 (*Ds*) mediated the outcome of two-strain co-cultures with strains susceptible to the strain's tailocins or/and was targeted by tailocins of the competitor (Fig. S3). To monitor changes in the proportion of *Ds* in mixed cultures, we selected a spontaneous rifampicin-resistant mutant of *Ds* (*Ds* RIF), which could be selectively recovered on rifampicin-containing media. Ds RIF retained tailocin production and did not differ from the parental strain in growth rate in 0.1 TSB or ability to compete with the wt *in vitro*. These results indicate that the rifampicin-resistant *Dickeya* spp. strains are not compromised in competitiveness under the applied experimental conditions (Fig. 6). The competitive interactions were explored with three types of competitors: 1) those susceptible to the tailocins of *Ds* RIF, 2) those producing a tailocin inhibitory to *Ds* RiF, or 3) those for which mutual killing with Ds RIF occurs in vitro. The success of a strain was defined as the extent of its end-point dominance in the

co-culture (>90% of total CFU) following its initial inoculation in equal proportions (Fig. S3). Two

competitors that could kill *Ds* RIF with their tailocins prevailed over *Ds* RIF, suggesting that

tailocins contributed to this success. However, Ds RIF outcompeted only a single strain that was

susceptible to its tailocin and only in the presence of H_2O_2. This suggests that tailocins alone do

not guarantee the success of a producer against a susceptible rival.

Discussion

Our study revealed that most SRP strains produce phage tail-like particles. While *P.*

brasiliense A143-S21-A16 and *P. atrosepticum* A597-S4-A17 lacked tailocin genomic clusters, *D.*

dianthicola A260-S21-A16 had a complete cluster but did not produce tailocins. This could be due

to reliance on other competitive strategies (Hibbing et al., 2010) or regulation by specific

environmental factors not present in the experiment (Bellieny-Rabelo et al., 2019). Furthermore,

we assessed the prevalence of tailocin clusters in 190 high-quality, complete genomes of SRP

bacteria and found that 83% of *Pectobacterium* spp. and 69% of *Dickeya* spp. genomes contain

clusters encoding tailocins. This indicates that tailocins are a very common feature among SRP

and may suggest that their production contributes to the ecological success of these bacteria (Ma

et al., 2007). The prevalence of tailocins in SRP aligns with the previous observations documenting

their presence in various bacterial species, including *Pseudomonas* spp. (Stice et al., 2023),

Pantoea spp. (Nagakubo et al., 2021) and *Streptomyces* spp. (Nagakubo et al., 2021).

Most tailocins can kill only closely related strains. This finding has, however, usually been

based on studies of a small set of strains either at the species or genus level (Granato et al.,

2019). Some observations suggest that some phage tail-like particles can kill different species than

the producer (Yao et al., 2017;Principe et al., 2018). Our work interrogated a large group of strains

belonging to three genera for their mutual tailocin-based interactions. While some SRP strains

produce tailocins with a broad range of activity, killing more than 30% of other strains of different

species. Caution, therefore, should be made to accept the current dogma of a narrow killing range

of tailocins.

Interestingly, while some *Dickeya* spp. produced tailocins able to kill *Pectobacterium* spp.

cells, the opposite was not observed in our study. Due to their evolutionary association with

bacteriophage tails (Ghequire and De Mot, 2015), it has been suggested that tailocins might retain

some degree of flexibility in their host range, such as known phage-host interactions (Backman et

al., 2024a). Inter-genus activity may remain an important trait in competition as mixed bacterial

communities involving both genera are frequently observed in the field, and the ability to target a

broader range of competitors could confer a selective advantage (Perombelon, 1988;Charkowski,

2009;2018;Barny et al., 2024).

Likewise, the observation of mutual killing, where two strains reciprocally target each other

with tailocins, has not been extensively documented. The higher incidence of mutual killing within

Dickeya spp. compared to *Pectobacterium* spp. suggests some (yet unknown) ecological factors

may select for this behavior more often in *Dickeya* spp. than in *Pectobacterium* spp. This is

somehow surprising, given that multiple species of SRP bacteria are frequently found together in

the same environments (Toth et al., 2021a;van der Wolf et al., 2021). In such settings, these

various species would be expected to experience similar ecological pressure and benefit from

reciprocal tailocin production (Shyntum et al., 2019).

Phage tail-like particles isolated from both SRP genera exhibited similar morphological

characteristics despite the apparent phylogenetic differences in the origins of tailocin clusters in

Pectobacterium spp. and *Dickeya* spp. suggests that these phage tail-like particles have been

acquired independently in those genera. All of the clusters present in *Dickeya* spp. exhibit

homology to that encoding P2D1 tailocin of *D. dadantii* 3937 (Borowicz et al., 2023), whereas all

clusters present in *Pectobacterium* spp. express homology to the cluster encoding carotovoricin Er

(Yamada et al., 2006). The fact that these phage tail-like particles were observed in all *Dickeya*

spp. and most of the *Pectobacterium* spp. strains tested suggest that the tailocin clusters were

acquired before species diversification in both genera and were subsequently maintained rather

unaltered. As tailocins are large protein complexes and their synthesis should be a burden for a

producer cell, their conservation suggests that tailocin production plays an important role in the

ecology of these bacteria (Glasner et al., 2008;Hugouvieux-Cotte-Pattat, 2016).

The genes encoding tailocin fiber proteins exhibited the lowest homology in the tailocin
clusters both in *Dickeya* spp. and *Pectobacterium* spp. This confirms that fiber proteins play a
pivotal role in determining the specificity of tailocins and their killing range, as reported earlier
(Dams et al., 2019). Fiber proteins are the most important determinants of target recognition in
bacterial cells (Backman et al., 2024a). This specificity is crucial for the success of tailocins as
killing agents, enabling them to discriminate between susceptible and resistant bacterial strains. In
bacteriophages, receptor-binding proteins are known to undergo positive selection to adapt to
variations in surface receptors of target bacteria, leading to high variability even among closely
related strains (Bertozzi Silva et al., 2016). In SRP, the killing spectrum of tailocins, determined by
the sequence of fiber proteins, appears to be more dependent on ecological factors and the
pressure of host-pathogen interactions encountered by the individual strains than the evolutionary
relationships between them.

Although mitomycin C is widely used to trigger tailocin induction under laboratory
conditions, in natural and agricultural environments, the likely inducers of tailocins remain
unknown. It seemed likely that other factors that cause DNA damage and thus activate the SOS
pathway could also serve as inducers of tailocins. In the proof-of-concept experiment, we found
that hydrogen peroxide was a potent inducer of SRP phage tail-like particles. Like mitomycin C,
hydrogen peroxide is known to activate the ROS response (Storz et al., 1990;Erill et al., 2007).
During infection, SRP bacteria often encounter oxidative stress in plant hosts during the immune
response of the plant (Reverchon and Nasser, 2013;Jiang et al., 2016). Therefore, the induction of
tailocins by reactive oxygen species may serve as a mechanism for bacterial survival in their
natural settings, aiding in eliminating competing bacteria under stressful conditions. Indeed, the
highest levels of competition might be expected during the infection process, where the change in
the plant nitch during injection leads to very high bacterial numbers. The ability of plant pathogenic
bacteria, including SRP, to constitutively produce tailocins, as well as under stressful conditions,
such as exposure to ROS, may lead to important ecological consequences. The basal tailocin
production by a fraction of the SRP populations may be seen as an altruistic behavior in which a
small number of bacterial cells sacrifice themselves to confer an advantage for the entire

population over other microorganisms residing in the same location (Griffin et al., 2004).

Alternatively, basal tailocin production may be understood as a division of labor in which

subpopulations of bacteria are 'assigned' to different tasks to grant success to the entire population

in a given niche (Lee et al., 2010;West and Cooper, 2016;Zhang et al., 2016). The ability to

express tailocins in response to such oxidative stress might give SRP a competitive edge, allowing

them to eliminate competitors in the plant environment when they must compete with all other

microorganisms (Backman et al., 2024b).

The variable level of response to mitomycin C and hydrogen peroxide we observed in SRP

strains highlights the still unraveled complexity of tailocin regulation. This suggests that in SRP,

depending on the strain and type of inducer, the production of tailocins is regulated differently in

different strains. Similar observations were made for prophages, where various environmental

factors induced phages with different efficiency (Casjens, 2003;Varani et al., 2013).

The SRP strains addressed in this study were isolated from a single habitat. Despite this,

this collection comprised both aquatic and non-aquatic strains, most of which (except *P. aquaticum*

and *P. quasiaquaticum* strains known to be avirulent on plants) infect plants in agricultural settings

(Ben Moussa et al., 2022). We found that tailocins are produced in infected potato tubers but not in

river water. This aligns with the other studies in which tailocins were found to be produced primarily

in nutrient-rich, competitive environments (Carim et al., 2021). Decaying plant material would be

expected to be a much more nutrient-rich environment than river water. The environmental

conditions in potato tubers might provide several stimuli for tailocin production, including ROS,

plant exudates, cell wall degradation products, and microbial competition. In contrast, water is

neither the common niche nor the environment where competition is likely to be intense. Instead,

water rather serves as a translocation medium from one host to another (Pédron and Van

Gijsegem, 2019;van der Wolf et al., 2021). This observation may be particularly pertinent for SRP

bacteria, which compete with each other and other microorganisms for resources present within

the plant tissues (Reverchon and Nasser, 2013). The potato tuber tissue represents a primary

ecological niche for SRP pathogens (Perombelon and Kelman, 1980;Pérombelon and Salmond,

1995). Consequently, the presence of tailocins in both rotting tissue and the healthy surrounding

tissue suggests that tailocin production may be part of the virulence strategy to minimize the risk of

competitive infections.

Surprisingly, SRP strains belonging to species that are not found on plants (Portier et al.,

2020) and which do not provoke disease symptoms (*vide*: *P. quasiaquaticum* A411-S4-F17 and *P.*

aquaticum A212-S19-A16) did produce tailocins in the plant environment but not in river water. The

low nutrient availability, the cost of production of this complex machinery, and the few direct

competitors of SRP bacteria in river water likely explain the absence of tailocin production in this

environment (Lamichhane and Bartoli, 2015) (Bradford et al., 2013;Toth et al., 2021b). The fact

that these tailocin clusters were maintained in these water-associated species and were induced

by the same stress trigger as plant-associated species suggests that tailocin production is not

solely tied to SRP virulence on plants but could also play a role in other niches where these

bacteria encounter stress. For example, it has been suggested that these water-associated

species could be associated with insect larvae in the water stream (Ben Moussa et al., 2022).

Whether tailocin production is triggered in such an environment remains to be determined.

Multiple factors play roles in the competition between microbes, with success being the sum

of factors/traits of varying importance depending on the environmental context (Hibbing et al.,

2010). In our study of mixed populations of tailocin producers and targets, tailocins were not

always sufficient to ensure the dominance of producers over susceptible strains. The widespread

occurrence of tailocin clusters among SRP led us to conclude that tailocin-mediated success is

often conditional. It is well known that SRP bacteria use diverse weaponry for intra-genus and

intra-genus interactions; therefore, tailocin production may be just one possibility to compete with

kin microbes.

Several factors, such as environmental conditions, strain-specific interactions, and the

relative abundance of producers versus susceptible strains, can influence the outcome of

competitive interactions. This study explored an 'even start' scenario, where tailocin-producing and

susceptible competitors were initially mixed in equal cell numbers. In bacterial populations, even

when one strain has an antagonistic advantage (like producing phage tail-like particles), an equal

start condition can neutralize this advantage because the susceptible strain is not immediately

overwhelmed by that trait (Ross-Gillespie and Kummerli, 2014). Moreover, tailocin production,

unlike the secretion of most bacterial antimicrobials, requires the lysis of the producer cells. Ergo, it

can negatively influence the growth rate of the producer's population. This burden can be assumed

to be negligible when the producer-to-target ratio is high. Therefore, we speculate that tailocin

production may be useful to SRP bacteria to protect already conquered niches from being invaded

by competitors, a process named niche exclusion (Bauer et al., 2018).

Author Contributions

MB: Conceptualization, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing, DMK: Conceptualization, Investigation, Supervision, Visualization, Writing – original draft, Writing – review & editing, MS: Investigation, Methodology, Visualization, MN: Investigation, Methodology, Visualization, Writing – original draft, IM: Investigation, Methodology, Writing – original draft, PC: Methodology, Writing – original draft, JP: Conceptualization, Investigation, Methodology, Writing – review & editing, MAB: Conceptualization, Investigation, Methodology, Writing – review & editing, PYC: Investigation, Methodology, JD: Conceptualization, Investigation, Methodology, RC: Conceptualization, Data curation, Funding acquisition, Resources, Supervision, Writing – original draft, Writing – review & editing

Competing Interest Statement

The authors declare no competing interests.

Data availability statement

All data can be obtained from the corresponding author and will be shared freely upon reasonable request.

Acknowledgments

This research was financially supported by the National Science Center, Poland (Narodowe Centrum Nauki, Polska) via a research grant SONATA BIS 10 (2020/38/E/NZ9/00007) to Robert Czajkowski. The authors would like to express their gratitude to Prof. Steven E. Lindow (University of California-Berkeley, Berkeley, CA, United States) for his comments on the manuscript and his editorial work.

Tables

Table 1. Tailocin-mediated killing events for the tested pool of 27 SRP strains.

Group	Total interactions[A]	Killing events		Mutual killing		
		No.	% total	No.	% total	% killing events
All SRPs	702	118	16.8%	40	5.7%	33.9%
Pectobacterium spp.[B]	306	66	21.6%	10	3.3%	15.2%
Dickeya spp.[C]	56	32	57.1%	21	37.5%	65.6%
Mixed genera[D]	340	19	5.6%	9	2.6%	47.37%
Psp vs *Dsp*	162	0	0%	0	0%	0%
Dsp vs *Psp*	162	6	3.7%	0	0%	0%
Dsp vs *M. par.*	8	5	62.5%	4	50%	80%
M. par vs *Dsp*	8	6	75%	5	62.5%	83.3%
M. par vs *Psp*	18	2	11.1%	0	0%	0%
Psp vs *M. par*	18	0	0%	0	0%	0%

[A] total interactions - the number of combinations where each strain from a subset was tested both for the potential to produce tailocins and for tailocin susceptibility (potential target)
[B] both strains in the tested interaction assigned to *Pectobacterium* spp. (total of 18 strains tested)
[C] both strains in the tested interaction assigned to *Dickeya* spp. (total of 8 strains tested)
[D] each strain of the pair representing a different genus; *Psp* – *Pectobacterium* spp.; *Dsp* – *Dickeya* spp.; *M. par* – *Musicola paradisiaca* NCPPB 2511. The first strain in each combination is the tailocin producer.

Table 2. Production of tailocins in river water and potato tubers. Qualitative tailocin
production in different conditions for various bacterial strains. The strains tested include *P.*
versatile A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16, *D.*
dadantii 3937, *D. chrysanthemi* A604-S21-A17, and *D. zeae* A586-S18-A17. Tailocin
production was assessed in infected potato tubers, in Durance River water, and as a positive
control following Mitomycin C and H_2O_2 induction in TSB (Tryptic Soy Broth). The symbols
represent the following: "+" indicates detectable tailocins in the sample with the applied
method; "-" indicates undetectable tailocins in the sample with the applied method; "*"
indicates that tailocins were detectable if tuber tissue in direct contact with inoculated
bacteria was pooled from three potato tubers.

	Tailocins production			
Strain	**In Durance River water**	**In infected potato tuber**	**Mitomycin C induction in TSB**	**H_2O_2 induction in TSB**
P. versatile A69-S13-O15	-	+	+	+
P. quasiaquaticum A411-S4-F17	-	+*	+	+
P. aquaticum A212-S19-A16	-	+*	+	+
D. dadantii 3937	-	+	+	+
D. chrysanthemi A604-S21-A17	-	+	+	+
D. zeae A586-S18-A17	-	+	+	+

+ tailocins detectable in the sample in the applied method

- tailocins undetectable in the sample in the applied method

** tubers were asymptomatic, but tailocins were detectable in tuber tissue at the
inoculation site; the material was collected and pooled from three potato tubers.*

Figures

657 **Fig. 1 TEM images picturing tailocins isolated from the 27 analyzed SRP strains.**

Fig. 2 Images of the representative tailocins generated based on AFM scans. AFM imaging was performed using the JPK NanoWizard® 4 (NanoScience) in QI™ in the quantitative imaging (QI)™ force spectroscopy mode. Images were processed using JPK Data Processing software. Panel A shows the AFM scan of Dickeyocin P2D1 produced by *D. dadantii* 3937, panels B-H. show the highest quality 3D visualizations of representative tailocin particles, generated based on AFM scans, for the following strains: *D. dadantii* 3937, *D. solani* A623-S20-A17, *D. zeae* A661-S21-A17. *M. paradisiaca* NCPPB 2511, *P. odoriferum* A367-S5-N16, *P. aquaticum* A105-S21-F16, *P. aquaticum* A104-S21-F16.

Fig. 3 Matrix of tailocin-mediated interactions within the group of tested SRP strains.
The tested pool comprised 27 strains: 25 isolates from the Durance River water and two type strains: *D. dadantii* 3937 and *M. paradisiaca* NCPPB 2511 (in bold). Grey-shaded cells indicate the susceptibility of a target strain to the tailocins of a given producer strain (unilateral killing). Unilateral killing where the producer and the target belong to different genera, is additionally marked by a yellow dot. Lillac cells also indicate the susceptibility of the target strain to the investigated tailocin, but in particular situations where the target strain was observed to produce 'retributive' tailocins against the producer (mutual killing). A dedicated column indicates whether tailocins were (Y) or were not (N) detected by microscopic imaging in culture supernatants of the tested strains following treatment with mitomycin C.

Fig. 4 Comparison of tailocin clusters in the analyzed group of SRPs. The upper panel
(A) depicts regions homologous to the carotovoricin-encoding cluster typical for
Pectobacterium spp. (reference: carotovoricin Er accession: AB045036.1, *P. versatile* A69-
S13-O15 RMBQ01000001.1, positions in the genome: 94551-113781). The lower panel (B)
shows regions homologous to dickeyocin P2D1, typical for *Dickeya* spp. and also present in
Musicola paradisiaca NCPPB 2511 (reference: *D. dadantii* 3937 NC_014500.1, positions in
the genome 2730482-2757061).

B

methyl-accepting chemotaxis protein
phage repressor protein CI
hypothetical protein
hypothetical protein
DUF2732 family protein
TraR/DksA fam. transcriptional regulator
replication endonuclease
hypothetical protein
tail protein X
phage tail protein
hypothetical protein
phage baseplate assembly protein V
GPW/gp25 protein
baseplate asssembly protein
phage tail protein I
tail fiber protein
esterase-like act. of phytase fam. protein
arom. acid/H+ symport fam. MFS transp.
DNA endonuclease SmrA
phage tail protein
recombinase family protein
phage tail protein
tail fiber assembly protein
phage tail sheath protein
phage major tail tube protein
GpE family phage tail protein
phage tail assembly protein
hypothetical protein
phage tail protein
VgrG
ogr
guaD

D. dadantii 3937 **(Dickeyocin P2D1)**
D. solani A623-S20-A17
D. dianthicola A260-S21-A16
D. chrysanthemi A604-S21-A17
D. zeae A661-S21-A17
D. zeae A586-S18-A17
D. oryzae A642-S2-A17
D. oryzae A003-S1-M15
M. paradisiaca NCPPB 2511

5 kb

0   Identity (%)   100

baseplate   sheath   tube   spike   fiber/other tail genes   regulatory/replication   other homologous   strain specific genes

Fig. 5 Basal and induced tailocin production levels in six *Pectobacterium* spp. and *Dickeya* spp strains. The graph shows the logarithm of the arbitrary tailocin units per mL of culture (Log AU mL^{-1}) for three strains of *Pectobacterium* spp. (*P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16) and three strains of *Dickeya* spp. (*D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D. zeae* A586-S18-A17). The basal production levels (no induction) are compared to those with mitomycin C or hydrogen peroxide after induction. Averages from 2 experiments, each consisting of 3 separate inductions, are shown, and error bars represent the standard deviation.

Fig. 6 Competition between *D. solani* A623-S20-A17 RIF[R] (*Ds* RIF) and other SRP strains in two-strain co-cultures. The success was defined as the end-point dominance of a given strain in the co-culture (>90% of total CFU), with both strains ensured with an even start (1:1 mix; ca. 10^7 CFU/ml/strain). The theoretical winners, predicted based on tailocin-mediated interactions, were compared with those that were experimentally established. Co-cultures were conducted for 22 h in 0.1 TSB alone or supplemented with 0.88 mM H_2O_2 (tailocin-inductive conditions). SET 1: strains sensitive to tailocins of *Ds* RIF. SET 2: strains producing tailocins against *Ds* RIF. SET 3: strains sensitive to tailocins of *Ds* RIF and producing tailocins targeting this strain (mutual killing). *ctrl* – control experiment; competition between *Ds* RIF and the *wt* variant of this strain.

Supplementary Tables

Table S1. List of bacterial strains used in this study.

Strain	Isolation source, country, year	Reference	GenBank accession no. for genome
Dickeya chrysanthemi A604-S21-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406775.2
Dickeya dianthicola A260-S21-A16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_020406895.2
Dickeya oryzae A003-S1-M15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_020406815.2
Dickeya oryzae A642-S2-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406685.2
Dickeya solani A623-S20-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406975.1
Dickeya parazeae A586-S18-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020520245.1
Dickeya zeae A661-S21-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406575.2
Pectobacterium aquaticum A101-S19-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382625.2
Pectobacterium aquaticum A104-S21-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382595.2
Pectobacterium aquaticum A105-S21-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382585.2
Pectobacterium aquaticum A212-S19-A16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382565.3
Pectobacterium atrosepticum A597-S4-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406655.2
Pectobacterium brasiliense A143-S20-M16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_013449685.1
Pectobacterium carotovorum A077-S18-O15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_020520265.1
Pectobacterium carotovorum A280-S2-N16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_020406635.1
Pectobacterium odoriferum A122-S21-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_013450045.1
Pectobacterium odoriferum A367-S5-N16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_022507245.1
Pectobacterium peruviense A350-S18-N16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003312355.2
Pectobacterium peruviense A97-S13-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003312345.1
Pectobacterium polaris A641-S6-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406835.1
Pectobacterium quasiaquaticum A477-S1-J17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_014946775.2
Pectobacterium quasiaquaticum A411-S4-F17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_014946845.1
Pectobacterium versatile A546-S4-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406615.1
Pectobacterium versatile A69-S13-O15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_004296725.1
Pectobacterium versatile A73-S18-O15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_020865565.1
Dickeya dadantii 3937	*Saintpaulia* sp. France. 1972	(Kotoujansky et al., 1982)	GCA_000147055.1
Musicola paradisiaca NCPPB 2511	*Musa paradisiaca*. Colombia. 1973 *	(Hugouvieux-Cotte-Pattat et al., 2021)	GCA_000400505.1

725 **Table S2. List of sensitive strains used in a spot assay to detect tailocins from the**
726 **given producer strains.**

Tailocin producer	Sensitive strain
P. versatile A69-S13-O15	*P. aquaticum* A105-S21-F16
P. quasiaquaticum A411-S4-F17	*P. atrosepticum* A597-S4-A17
P. aquaticum A212-S19-A16	*P. aquaticum* A105-S21-F16
D. dadantii 3937	*M. paradisiaca* NCPPB 2511
D. chrysanthemi A604-S21-A17	*D. zeae* A586-S18-A17
D. zeae A586-S18-A17 (recently reclassified to *D. parazeae*)	*M. paradisiaca* NCPPB 2511

Supplementary Figures

Supplementary Fig. S1. Comparison of phenotypic and phylogenetic trees. (A)
dendrogram calculated based on the similarity of killing spectra of investigated strain; the
tree is drawn to scale, with branch lengths measured in the distance; (B) multilocus
sequence analysis of core nucleotide coding sequences using the BioNJ algorithm. Branch
lengths and bootstrap values are shown. *Pectobacterium* spp. strains are marked in green
and *Dickeya* spp. strains are marked in red.

Supplementary Fig. S2. Severity of soft rot symptoms caused on potato tubers by six
SRP strains tested for production of tailocins *in planta*. The results are expressed as a
percentage of rotten tissue relative to the total mass of each potato tuber. In the box plot,
whiskers reflect the maximum and minimum values; the box shows the interquartile range
(Q1 to Q3), and the bars inside the boxes represent medians. Points indicate results for
individual samples/tubers (n=6). The asterisks (*) denote statistically significant differences (p
< 0.05) between a sample inoculated with the given strain and the negative control.
Significance was calculated using the one-sample t-test for data sets with normal distribution
and the Wilcoxon test for data not following a normal distribution. Negative control – tubers
inoculated with sterile PBS buffer.

 Supplementary Fig. S3 Schematic representation of the steps included in the
 competition experiment. Created with BioRender.com.

Supplementary Fig. S4 Comparison of growth rates (A) and the survival (B) of the *D. solani* A623-S20-A17 wt and the rifampicin-resistant mutant of this strain in 0.1 TSB alone and supplemented with 0.003 % hydrogen peroxide (H_2O_2). The test was performed in 96-well plates incubated at 28˚C, with shaking, in a multiwell plate reader (Biotek). To assess growth rates (A), 150 µl aliquots of 0.1 TSB medium were inoculated with 3 µl of suspension containing o/n culture diluted to 0.5 McF (low inoculum). Each time point is an average of two replicates. For cell survival (B), the medium was inoculated with a high initial number of cells (ca. 10^7 CFU/ml) applied in the co-culture competition experiments (Fig. 5). The results are the average of two separate experiments, 2 technical replicates each. The obtained results indicate that the wt and RIF^R variants of *D. solani* A623-S20-A17 have similar growth characteristics in the applied conditions and that 0.003% H_2O_2 prevents the growth of the strains when low initial inoculum is used but does not negatively affect the survival of cells in high-density cultures.

Supplementary datasets

Dataset S1 Results of a megablast search for carotovoricin clusters within 116 closed genomes of *Pectobacterium* spp. The carotovoricin sequence from *P. versatile* A73-S18-O15 was used as a query (genome accession: NZ_CP086369.1).

Dataset S2 Results of a megablast search for dickeyocin clusters within 75 closed *Dickeya* and *Musicola* spp. genomes. Dickeyocin P2D1 from *D. dadantii* 3937 was used as a query (genome accession: NC_014500.1).

Dataset S3 Python script for analyses (ChatGPT-4o)

References

Backman, T., Burbano, H.A., and Karasov, T.L. (2024a). Tradeoffs and constraints on the evolution of tailocins. *Trends in Microbiology*.

Backman, T., Latorre, S.M., Symeonidi, E., Muszynski, A., Bleak, E., Eads, L., Martinez-Koury, P.I., Som, S., Hawks, A., Gloss, A.D., Belnap, D.M., Manuel, A.M., Deutschbauer, A.M., Bergelson, J., Azadi, P., Burbano, H.A., and Karasov, T.L. (2024b). A phage tail-like bacteriocin suppresses competitors in metapopulations of pathogenic bacteria. *Science* 384**,** eado0713.

Barny, M.-A., Thieffry, S., Gomes De Faria, C., Thebault, E., and Pédron, J. (2024). Bacterial pathogens dynamic during multi-species infections. *Peer Community Journal* 4.

Bauer, M.A., Kainz, K., Carmona-Gutierrez, D., and Madeo, F. (2018). Microbial wars: Competition in ecological niches and within the microbiome. *Microb Cell* 5**,** 215-219.

Bellieny-Rabelo, D., Tanui, C.K., Miguel, N., Kwenda, S., Shyntum, D.Y., and Moleleki, L.N. (2019). Transcriptome and Comparative Genomics Analyses Reveal New Functional Insights on Key Determinants of Pathogenesis and Interbacterial Competition in Pectobacterium and Dickeya spp. *Appl Environ Microbiol* 85**,** e02050-02018.

Ben Moussa, H., Bertrand, C., Rochelle-Newall, E., Fiorini, S., Pedron, J., and Barny, M.A. (2022). The Diversity and Abundance of Soft Rot Pectobacteriaceae Along the Durance River Stream in the Southeast of France Revealed by Multiple Seasonal Surveys. *Phytopathology* 112**,** 1676-1685.

Berg, G., Grosch, R., and Scherwinski, K. (2007). Risikofolgeabschätzung für den Einsatz mikrobieller Antagonisten: Gibt es Effekte auf Nichtzielorganismen? *Gesunde Pflanzen* 59**,** 107-117.

Bertozzi Silva, J., Storms, Z., and Sauvageau, D. (2016). Host receptors for bacteriophage adsorption. *FEMS Microbiol Lett* 363.

Booth, S.C., Smith, W.P.J., and Foster, K.R. (2023). The evolution of short- and long-range weapons for bacterial competition. *Nat Ecol Evol* 7**,** 2080-2091.

Borowicz, M., Krzyżanowska, D.M., Narajczyk, M., Sobolewska, M., Rajewska, M., Czaplewska, P., Węgrzyn, K., and Czajkowski, R. (2023). Soft rot pathogen Dickeya dadantii 3937 produces tailocins resembling the tails of Peduovirus P2. *Frontiers in Microbiology* 14.

Bradford, S.A., Morales, V.L., Zhang, W., Harvey, R.W., Packman, A.I., Mohanram, A., and Welty, C. (2013). Transport and Fate of Microbial Pathogens in Agricultural Settings. *Critical Reviews in Environmental Science and Technology* 43**,** 775-893.

Brettin, T., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Olsen, G.J., Olson, R., Overbeek, R., Parrello, B., Pusch, G.D., Shukla, M., Thomason, J.A., 3rd, Stevens, R., Vonstein, V., Wattam, A.R., and Xia, F. (2015). RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5**,** 8365.

Carim, S., Azadeh, A.L., Kazakov, A.E., Price, M.N., Walian, P.J., Lui, L.M., Nielsen, T.N., Chakraborty, R., Deutschbauer, A.M., Mutalik, V.K., and Arkin, A.P. (2021). Systematic discovery of pseudomonad genetic factors involved in sensitivity to tailocins. *ISME J* 15**,** 2289-2305.

Casjens, S. (2003). Prophages and bacterial genomics: what have we learned so far? *Mol Microbiol* 49**,** 277-300.

Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17**,** 540-552.

Charkowski, A.O. (2009). Decaying signals: will understanding bacterial-plant communications lead to control of soft rot? *Curr Opin Biotechnol* 20**,** 178-184.

Charkowski, A.O. (2018). The Changing Face of Bacterial Soft-Rot Diseases. *Annu Rev Phytopathol* 56**,** 269-288.

Czajkowski, R., De Boer, W.J., Velvis, H., and Van Der Wolf, J.M. (2010). Systemic colonization of potato plants by a soilborne, green fluorescent protein-tagged strain of *Dickeya* sp. biovar 3. *Phytopathology* 100**,** 134-142.

Dams, D., Brondsted, L., Drulis-Kawa, Z., and Briers, Y. (2019). Engineering of receptor-binding proteins in bacteriophages and phage tail-like bacteriocins. *Biochem Soc Trans* 47**,** 449-460.

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32**,** 1792-1797.

Erill, I., Campoy, S., and Barbe, J. (2007). Aeons of distress: an evolutionary perspective on the bacterial SOS response. *FEMS Microbiol Rev* 31**,** 637-656.

Fiołka, M.J., Czaplewska, P., Macur, K., Buchwald, T., Kutkowska, J., Paduch, R., Kaczyński, Z., Wydrych, J., and Urbanik-Sypniewska, T. (2019). Anti-Candida albicans effect of the protein-carbohydrate fraction obtained from the coelomic fluid of earthworm Dendrobaena veneta. *PLoS One* 14**,** e0212869.

Ghequire, M.G.K., and De Mot, R. (2015). The Tailocin Tale: Peeling off Phage Tails. *Trends Microbiol* 23**,** 587-590.

Gilchrist, C.L.M., and Chooi, Y.H. (2021). clinker & clustermap.js: automatic generation of gene cluster comparison figures. *Bioinformatics* 37**,** 2473-2475.

Glandorf, D.C.M., Brand, I., Bakker, P.a.H.M., and Schippers, B. (1992). Stability of Rifampicin Resistance as a Marker for Root Colonization Studies of Pseudomonas-Putida in the Field. *Plant and Soil* 147**,** 135-142.

Glasner, J.D., Marquez-Villavicencio, M., Kim, H.S., Jahn, C.E., Ma, B., Biehl, B.S., Rissman, A.I., Mole, B., Yi, X., Yang, C.H., Dangl, J.L., Grant, S.R., Perna, N.T., and Charkowski, A.O. (2008). Niche-specificity and the variable fraction of the Pectobacterium pan-genome. *Mol Plant Microbe Interact* 21**,** 1549-1560.

Goldman, A.R., Beer, L.A., Tang, H.Y., Hembach, P., Zayas-Bazan, D., and Speicher, D.W. (2019). Proteome Analysis Using Gel-LC-MS/MS. *Curr Protoc Protein Sci* 96**,** e93.

Golebiowski, A., Pomastowski, P., Rafinska, K., Zuvela, P., Wong, M.W., Pryshchepa, O., Madajski, P., and Buszewski, B. (2022). Functionalization of Alpha-Lactalbumin by Zinc Ions. *ACS Omega* 7**,** 38459-38474.

Gouy, M., Guindon, S., and Gascuel, O. (2010). SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27**,** 221-224.

Granato, E.T., Meiller-Legrand, T.A., and Foster, K.R. (2019). The Evolution and Ecology of Bacterial Warfare. *Curr Biol* 29**,** R521-R537.

Griffin, A.S., West, S.A., and Buckling, A. (2004). Cooperation and competition in pathogenic bacteria. *Nature* 430**,** 1024-1027.

Hibbing, M.E., Fuqua, C., Parsek, M.R., and Peterson, S.B. (2010). Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* 8**,** 15-25.

Hockett, K.L., and Baltrus, D.A. (2017). Use of the Soft-agar Overlay Technique to Screen for Bacterially Produced Inhibitory Compounds. *J Vis Exp*.

Hugouvieux-Cotte-Pattat, N. (2016). "Chapter Four - Metabolism and Virulence Strategies in Dickeya–Host Interactions," in *Progress in Molecular Biology and Translational Science,* eds. M. San Francisco & B. San Francisco. Academic Press), 93-129.

Hugouvieux-Cotte-Pattat, N., Des-Combes, C.J., Briolay, J., and Pritchard, L. (2021). Proposal for the creation of a new genus Musicola gen. nov., reclassification of Dickeya paradisiaca (Samson et al. 2005) as Musicola paradisiaca comb. nov. and description of a new species Musicola keenii sp. nov. *Int J Syst Evol Microbiol* 71.

Hugouvieux-Cotte-Pattat, N., Pédron, J., and Van Gijsegem, F. (2023). Insight into biodiversity of the recently rearranged genus Dickeya. *Frontiers in Plant Science* 14.

Hugouvieux-Cotte-Pattat, N., and Van Gijsegem, F. (2021). Diversity within the Dickeya zeae complex, identification of Dickeya zeae and Dickeya oryzae members, proposal of the novel species Dickeya parazeae sp. nov. *Int J Syst Evol Microbiol* 71.

Jiang, X., Zghidi-Abouzid, O., Oger-Desfeux, C., Hommais, F., Greliche, N., Muskhelishvili, G., Nasser, W., and Reverchon, S. (2016). Global transcriptional response of Dickeya dadantii to environmental stimuli relevant to the plant infection. *Environ Microbiol* 18**,** 3651-3672.

Kotoujansky, A., Lemattre, M., and Boistard, P. (1982). Utilization of a thermosensitive episome bearing transposon TN10 to isolate Hfr donor strains of Erwinia carotovora subsp. chrysanthemi. *J Bacteriol* 150**,** 122-131.

Krzyzanowska, D.M., Maciag, T., Siwinska, J., Krychowiak, M., Jafra, S., and Czajkowski, R. (2019). Compatible Mixture of Bacterial Antagonists Developed to Protect Potato Tubers from Soft Rot Caused by Pectobacterium spp. and Dickeya spp. *Plant Dis* 103**,** 1374-1382.

Lamichhane, J.R., and Bartoli, C. (2015). Plant pathogenic bacteria in open irrigation systems: what risk for crop health? *Plant Pathology* 64**,** 757-766.

Lee, H.H., Molla, M.N., Cantor, C.R., and Collins, J.J. (2010). Bacterial charity work leads to population-wide resistance. *Nature* 467**,** 82-85.

Ma, B., Hibbing, M.E., Kim, H.S., Reedy, R.M., Yedidia, I., Breuer, J., Breuer, J., Glasner, J.D., Perna, N.T., Kelman, A., and Charkowski, A.O. (2007). Host range and molecular phylogenies of the soft rot enterobacterial genera pectobacterium and dickeya. *Phytopathology* 97**,** 1150-1163.

Marquez-Villavicencio, M.D.P., Groves, R.L., and Charkowski, A.O. (2011). Soft Rot Disease Severity Is Affected by Potato Physiology and Pectobacterium taxa. *Plant Dis* 95**,** 232-241.

Miele, V., Penel, S., and Duret, L. (2011). Ultra-fast sequence clustering from similarity networks with SiLiX. *BMC Bioinformatics* 12**,** 116.

Nagakubo, T., Yamamoto, T., Asamizu, S., Toyofuku, M., Nomura, N., and Onaka, H. (2021). Phage tail-like nanostructures affect microbial interactions between Streptomyces and fungi. *Sci Rep* 11**,** 20116.

Nguyen, H.A., Kaneko, J., and Kamio, Y. (2002). Temperature-dependent production of carotovoricin Er and pectin lyase in phytopathogenic Erwinia carotovora subsp. carotovora Er. *Biosci Biotechnol Biochem* 66, 444-447.

Nielsen, F. (2016). "Hierarchical Clustering," in *Introduction to HPC with MPI for Data Science*. (Cham: Springer International Publishing), 195-211.

Pédron, J., and Van Gijsegem, F. (2019). Diversity in the Bacterial Genus Dickeya Grouping Plant Pathogens and Waterways Isolates. *OBM Genetics* 3, 1-1.

Perombelon, M.C.M. (Year). "Ecology of Erwinias causing stem and tuber diseases", in: *Bacterial disease of Potato: report of the planning conference on Bacterial diseases of potato*).

Perombelon, M.C.M., and Kelman, A. (1980). Ecology of the soft rot Erwinias. *Annual Review of Phytopathology* 18, 361-387.

Pérombelon, M.C.M., and Salmond, G.P.C. (1995). Bacterial soft rots. *Pathogenesis and Host Specificity in Plant Diseases* 1, 1-20.

Perombelon, M.C.M., and Van Der Wolf, J.M. (2002). Methods for the detection and quantification of *Erwinia carotovora* subsp. *atroseptica* (*Pectobacterium carotovorum* subsp. *atrosepticum*) on potatoes: a laboratory manual. *Scottish Crop Research Institute Annual Report* 10.

Portier, P., Pédron, J., Taghouti, G., Dutrieux, C., and Barny, M.-A. (2020). Updated Taxonomy of Pectobacterium Genus in the CIRM-CFBP Bacterial Collection: When Newly Described Species Reveal "Old" Endemic Population. *Microorganisms* 8, 1441.

Principe, A., Fernandez, M., Torasso, M., Godino, A., and Fischer, S. (2018). Effectiveness of tailocins produced by Pseudomonas fluorescens SF4c in controlling the bacterial-spot disease in tomatoes caused by Xanthomonas vesicatoria. *Microbiol Res* 212-213, 94-102.

Pritchard, L., Humphris, S., Saddler, G.S., Elphinstone, J.G., Pirhonen, M., and Toth, I.K. (2013). Draft genome sequences of 17 isolates of the plant pathogenic bacterium dickeya. *Genome Announc* 1.

Reverchon, S., Muskhelisvili, G., and Nasser, W. (2016). "Chapter Three - Virulence Program of a Bacterial Plant Pathogen: The Dickeya Model," in *Progress in Molecular Biology and Translational Science,* eds. M. San Francisco & B. San Francisco. Academic Press), 51-92.

Reverchon, S., and Nasser, W. (2013). Dickeya ecology, environment sensing and regulation of virulence programme. *Environ Microbiol Rep* 5, 622-636.

Ross-Gillespie, A., and Kummerli, R. (2014). 'Evolution-proofing' antibacterials. *Evol Med Public Health* 2014, 134-135.

Schmidt, R., and Sinz, A. (2017). Improved single-step enrichment methods of cross-linked products for protein structure analysis and protein interaction mapping. *Anal Bioanal Chem* 409, 2393-2400.

Scholl, D. (2017). Phage Tail-Like Bacteriocins. *Annu Rev Virol* 4, 453-467.

Shyntum, D.Y., Nkomo, N.P., Shingange, N.L., Gricia, A.R., Bellieny-Rabelo, D., and Moleleki, L.N. (2019). The Impact of Type VI Secretion System, Bacteriocins and Antibiotics on Bacterial Competition of Pectobacterium carotovorum subsp. brasiliense and the Regulation of Carbapenem Biosynthesis by Iron and the Ferric-Uptake Regulator. *Front Microbiol* 10, 2379.

Stice, S.P., Jan, H.H., Chen, H.C., Nwosu, L., Shin, G.Y., Weaver, S., Coutinho, T., Kvitko, B.H., and Baltrus, D.A. (2023). Pantailocins: phage-derived bacteriocins from Pantoea ananatis and Pantoea stewartii subsp. indologenes. *Appl Environ Microbiol* 89, e0092923.

Storz, G., Tartaglia, L.A., Farr, S.B., and Ames, B.N. (1990). Bacterial defenses against oxidative stress. *Trends Genet* 6**,** 363-368.

Torres, M.A., Jones, J.D., and Dangl, J.L. (2006). Reactive oxygen species signaling in response to pathogens. *Plant Physiol* 141**,** 373-378.

Toth, I.K., Barny, M.-A., Brurberg, M.B., Condemine, G., Czajkowski, R., Elphinstone, J.G., Helias, V., Johnson, S.B., Moleleki, L.N., and Pirhonen, M. (2021a). Pectobacterium and Dickeya: environment to disease development. *Plant diseases caused by dickeya and pectobacterium species***,** 39-84.

Toth, I.K., Barny, M.-A., Brurberg, M.B., Condemine, G., Czajkowski, R., Elphinstone, J.G., Helias, V., Johnson, S.B., Moleleki, L.N., Pirhonen, M., Rossmann, S., Tsror, L., Van Der Waals, J.E., Van Der Wolf, J.M., Van Gijsegem, F., and Yedidia, I. (2021b). "Pectobacterium and Dickeya: Environment to Disease Development," in *Plant Diseases Caused by Dickeya and Pectobacterium Species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 39-84.

Van Der Wolf, J.M., Acuña, I., De Boer, S.H., Brurberg, M.B., Cahill, G., Charkowski, A.O., Coutinho, T., Davey, T., Dees, M.W., Degefu, Y., Dupuis, B., Elphinstone, J.G., Fan, J., Fazelisanagri, E., Fleming, T., Gerayeli, N., Gorshkov, V., Helias, V., Le Hingrat, Y., Johnson, S.B., Keiser, A., Kellenberger, I., Li, X., Lojkowska, E., Martin, R., Perminow, J.I., Petrova, O., Motyka-Pomagruk, A., Rossmann, S., Schaerer, S., Sledz, W., Toth, I.K., Tsror, L., Van Der Waals, J.E., De Werra, P., and Yedidia, I. (2021). "Diseases Caused by *Pectobacterium* and *Dickeya* apecies around the world," in *Plant diseases caused by Dickeya and Pectobacterium species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 215-261.

Van Gijsegem, F., Toth, I.K., and Van Der Wolf, J.M. (2021). "Soft Rot Pectobacteriaceae: A Brief Overview," in *Plant Diseases Caused by Dickeya and Pectobacterium Species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 1-11.

Varani, A.M., Monteiro-Vitorello, C.B., Nakaya, H.I., and Van Sluys, M.A. (2013). The role of prophage in plant-pathogenic bacteria. *Annu Rev Phytopathol* 51**,** 429-451.

Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., Van Der Walt, S.J., Brett, M., Wilson, J., Millman, K.J., Mayorov, N., Nelson, A.R.J., Jones, E., Kern, R., Larson, E., Carey, C.J., Polat, I., Feng, Y., Moore, E.W., Vanderplas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E.A., Harris, C.R., Archibald, A.M., Ribeiro, A.H., Pedregosa, F., Van Mulbregt, P., and Scipy, C. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* 17**,** 261-272.

Wagner, A. (2022). Competition for nutrients increases invasion resistance during assembly of microbial communities. *Mol Ecol* 31**,** 4188-4203.

West, S.A., and Cooper, G.A. (2016). Division of labour in microorganisms: an evolutionary perspective. *Nat Rev Microbiol* 14**,** 716-723.

Yamada, K., Hirota, M., Niimi, Y., Nguyen, H.A., Takahara, Y., Kamio, Y., and Kaneko, J. (2006). Nucleotide sequences and organization of the genes for carotovoricin (Ctv) from Erwinia carotovora indicate that Ctv evolved from the same ancestor as Salmonella typhi prophage. *Biosci Biotechnol Biochem* 70**,** 2236-2247.

Yao, G.W., Duarte, I., Le, T.T., Carmody, L., Lipuma, J.J., Young, R., and Gonzalez, C.F. (2017). A Broad-Host-Range Tailocin from Burkholderia cenocepacia. *Appl Environ Microbiol* 83.

Zhang, Z., Claessen, D., and Rozen, D.E. (2016). Understanding Microbial Divisions of Labor. *Front Microbiol* 7**,** 2070.