1   RESEARCH ARTICLE

3   Tailocin-mediated interactions in Soft Rot *Pectobacteriaceae*

Marcin Borowicz [a], Dorota M. Krzyżanowska [a], Marta Sobolewska [a], Magdalena Narajczyk [b], Inez

Mruk [c], Paulina Czaplewska [c], Jacques Pédron [d], Marie-Anne Barny [d], Pierre Yves Canto [d], Joanna

Dziadkowiec [e] and Robert Czajkowski [a,*]

[a] Laboratory of Biologically Active Compounds, Intercollegiate Faculty of Biotechnology of UG and

MUG, University of Gdansk, A. Abrahama 58, 80-307 Gdańsk, Poland;

[b] Bioimaging Laboratory, Faculty of Biology, University of Gdansk, Wita Stwosza 59, 80-308

Gdansk, Poland;

[c] Laboratory of Mass Spectrometry-Core Facility Laboratories, Intercollegiate Faculty of

Biotechnology of UG and MUG, University of Gdansk, Antoniego Abrahama 58, 80-307 Gdansk,

Poland;

[d] Institute of Ecology and Environmental Sciences of Paris, Sorbonne Université, INRAE, IRD,

CNRS, UPEC, UMR 7618, Paris, France;

[e] The Njord Centre, Departments of Geosciences and Physics, University of Oslo, Blindern, 0316

Oslo, Norway;

* Correspondence: robert.czajkowski@ug.edu.pl; phone: 0048 58 523 6333

Abstract

Bacteria often carry phage-derived elements within their genomes, some of which can produce phage-like particles (tailocins) used as weapons to kill kin strains in response to environmental conditions. This study investigates the production and activity of such particles by plant pathogenic bacteria belonging to the *Pectobacterium*, *Dickeya,* and *Musicola* genera that often compete for the same niche, providing an attractive model to study the ecological role of tailocins. Transmission electron microscopy (TEM) revealed that most analyzed strains (88%) produced tailocins. Tailocin-mediated killing interactions were assessed across 351 strain pairs, showing that *Dickeya* spp. had a higher incidence of killing (57.1%) than *Pectobacterium* spp. (21.6%). Additionally, *Dickeya* spp. strains exhibited inter-genus activity, targeting both *Pectobacterium* spp. and *Musicola* sp., while *Pectobacterium* spp. tailocins were genus-specific. A so-called "bilateral killing" was observed in 33.9% of interactions, predominantly within *Dickeya* spp. Although tailocins were morphologically indistinguishable between producer strains, genomic analyses identified conserved tailocin gene clusters across both genera, with distinct structural differences between *Pectobacterium* spp. and *Dickeya* spp tailocins. This suggests different origins of these particles. Induction experiments demonstrated that tailocin production was significantly boosted by hydrogen peroxide, supporting the potential role of these particles in bacteria-bacteria competition during infection. Lastly, tailocin production was detectable in infected potato tissue but not in river water, highlighting the ecological relevance of tailocins in plant environments.

Keywords:

bacteriophage, phage tail-like particles, hydrogen peroxide, inter-genus, interactions, *Erwinia chrysanthemi*, *Erwinia carotovora*, prophage

Introduction

In highly competitive and resource-limited environments that bacteria colonize and occupy, their survival and welfare often center on the ability to outcompete neighboring rival cells (Bauer et al., 2018;Wagner, 2022). To gain a competitive edge, bacteria evolve diverse strategies, including producing and utilizing phage tail-like particles (tailocins) - structures that directly resemble the tails of bacteriophages but function independently from viral infection and replication (Booth et al., 2023). These particles are highly specialized killing instruments that, by puncturing the cell envelope, lead to the disruption of the cell membrane following the death of the targeted cell (Scholl, 2017). Until recently, the utilization of phage tail-like particles has been primarily studied in *Pseudomonas* species (Carim et al., 2021). However, there are also reports of tailocins isolated from various other Gram-negative and Gram-positive bacterial species, including human, animal, and plant pathogens, as well as saprophytic bacteria that inhabit a wide range of environments (Backman et al., 2024a).

We previously identified and characterized a new type of phage tail-like particles produced by the plant pathogenic bacterium, *Dickeya dadantii* reference strain 3937 (Borowicz et al., 2023). This tailocin, named dickeyocin P2D1, was resistant to various environmental conditions and expressed broad killing potential against members of different *Dickeya* species but was harmless to *Pectobacterium* species and non-toxic to *Caenorhabditis elegans.* We postulated that the dickeyocin P2D1 targets remain phylogenetically closely related strains. Our study was the first to target the presence and conduct an in-depth characterization of a new phage tail-like particle in SRP bacteria. Previously, only one phage tail-like particle, carotovoricin Er from *Pectobacterium carotovorum*, was described, and its nucleotide sequence was analyzed in detail (Yamada et al., 2006). However, since then, large studies have yet to be done to assess the presence of tailocins in SRP bacteria. Moreover, the carotovoricin Er sequence (Yamada et al., 2006) differs from that of dickeyocin P2D1, indicating a distinct evolutionary origin of these two tailocins.

SRP bacteria (consisting of *Pectobacterium* spp., *Dickeya* spp., and *Musicola* spp., previously collectively described as pectinolytic *Erwinia* spp.) are a particularly interesting model

for studying the production and utilization of phage tail-like particles in the environment.

Pectobacterium spp., *Dickeya* spp., and *Musicola* spp. belong to the same species complex and

are often found together in infected hosts. Consequently, due to their metabolic similarities and

overlapping niches, they are known to compete against each other (Toth et al., 2021a).

Furthermore, these bacteria are known to pass through various ecological niches, including host

and non-host plants, natural and agricultural soils, rainwater, surface water, sewage, and insects

(Van Gijsegem et al., 2021b). In these diverse environments, SRP bacteria often encounter a

diverse array of other species, both closely and distantly related, but their competitive strategies in

these alternative biomes remain unknown (Bellieny-Rabelo et al., 2019).

This study aimed to assess the utilization of phage tail-like particles (tailocins) in SRP

strains obtained from one selected environment – the Durance River in France (Ben Moussa et al.,

2022). Although river water is not recognized as a primary environment where SRP bacteria thrive,

this niche can be considered part of the continuum of distinct ecosystems that support the growth

and survival of SRP. The tested SRP pool included aquatic and non-aquatic strains, which

potentially could interact in water or be transmitted to a common plant host, where they might

interact during infection (Van Gijsegem et al., 2021b).

Our understanding of the competition mechanisms in SRP bacteria mediated by phage tail-

like particles remains poorly understood (Marquez-Villavicencio et al., 2011;Hugouvieux-Cotte-

Pattat, 2016;Hugouvieux-Cotte-Pattat et al., 2023). Specifically, no large studies have been

conducted to assess whether the production of tailocins is common in this group of bacteria and

how it contributes to their overall ecological success (Borowicz et al., 2023). Addressing this gap

remains important for understanding how plant-pathogenic bacteria establish and sustain

dominance in diverse environments, particularly in those niches where microbial competition

between kin strains is expected to be intense. Additionally, we aimed to characterize the isolated

particles and identify gene clusters associated with their synthesis, highlighting the prevalence of

tailocin production among environmental SRP strains and their possible role in the intragenus

competition.

Materials and Methods

Bacterial strains and culture conditions

Bacterial strains included in this work are listed in Supplementary Table S1. The pool comprised 25 SRP isolates from the waters of Durance River (France), collected between 2015 and 2017 (Ben Moussa et al., 2022), as well as two reference strains: *Dickeya dadantii* 3937 (Pritchard et al., 2013) and *Musicola paradisiaca* NCPPB 2511 (Hugouvieux-Cotte-Pattat et al., 2021). The latter strains were a model in our previous study on tailocins (dickeyocins) from *D. dadantii* 3937 (Borowicz et al., 2023). Routine propagation of bacterial cells was performed at 28 ºC in liquid Trypticase Soy Broth (TSB, Oxoid), with agitation (120 rpm), or on solid Trypticase Soy Agar (TSA, Oxoid) under the same growth conditions (Czajkowski et al., 2010).

Induction and purification of tailocins

Tailocin particles were purified as described previously (Borowicz et al., 2023). Briefly, an overnight culture (*ca.* 16 h) of each tested strain was rejuvenated (1:40) in a fresh aliquot of TSB and cultured for 2.5 hours. Next, tailocin production was induced by the addition of 0.1 µg mL^{-1} of mitomycin C (Abcam, Poland). Twenty-four hours post-induction, tailocins were PEG-purified from 10 mL volumes of filtered (0.2 µm, PES membrane, GoogLab™) culture supernatants. Purified particles were stored in phosphate buffer saline (PBS), pH 7.2, at 4 °C.

Microscopic imaging

The morphology of purified tailocin particles was investigated by transmission electron microscopy (TEM) and atomic force microscopy (AFM). TEM imaging was performed as described earlier (Borowicz et al., 2023) using the Tecnai Spirit BioTWIN microscope (FEI). AFM imaging was performed using JPK NanoWizard® 4 (NanoScience), in non-destructive quantitative imaging (QI™) force spectroscopy mode, employing the SCANASYST-AIR probes (f0 7.0 kHz, diameter <12 nm, k: 0.4 N/m) and the SCANASYST-FLUID+ probes (f0 150 kHz, diameter <12 nm, k: 0.7

N/m). Spring constants and sensitivity of the probes were calibrated using the thermal calibration

method. Before imaging, the samples were deposited on freshly cleaved mica surfaces and air-

dried. Images were processed using JPK Data Processing software. Dimensions of tailocins were

calculated based on TEM images. Tailocins from a given strain were considered for measurement

only if more than 4 individual particle images were available for analysis. This included 13 out of 18

Pectobacterium spp. strains, with a median of 10 particles measured per strain, and 7 *Dickeya*

spp., with a median of 17 particles measured per strain.

Killing spectrum and sensitivity assay

Twenty-seven SRP strains, including 25 water isolates (Supplementary Table S1), were tested,

pair by pair, to determine the target range (killing spectrum) of tailocins produced by each group

member. Likewise, the opposite was investigated, namely the sensitivity of strains to tailocins

produced by all other strains in the pool. Both the killing spectrum and sensitivity were tested using

the spot assay as previously described (Hockett and Baltrus, 2017;Yao et al., 2017;Borowicz et al.,

2023). Each combination was tested in duplicate in a single setup, and the entire experiment was

performed twice using independently-obtained batches of tailocins.

SDS-PAGE, ESI LC-MS/MS, and mapping of identified proteins to genomic *loci*

Proteins in PEG-purified tailocin preparations were separated by SDS-PAGE, fragmented, and

analyzed by mass spectrometry, as described in detail previously (Golebiowski et al.,

2022;Borowicz et al., 2023). Briefly, bands were excised from the gel, followed by in-gel trypsin

digestion of the embedded proteins, peptide elution, cleanup, and sample concentration (Schmidt

and Sinz, 2017;Goldman et al., 2019). Next, the peptides were analyzed by ESI LC-MS/MS. This

included separating peptides on the Eksigent microLC column ChromXP C18CL (3 μm, 120 Å, 150

× 0.3 mm). The samples were placed onto the column with a CTC Pal Autosampler (CTC Analytics

AG, Zwinger, Switzerland), and the injection volume was 5 μl. Solvents A and B consisted of 0.1%

(v/v) formic acid (FA) in water and acetonitrile, respectively. LC gradient parameters: 18–50%

solvent B in 30 min (Fiołka et al., 2019). Recording of mass spectra with Triple Tof 5600+ mass

spectrometer with DuoSpray Ion Source (AB SCIEX, Framingham, MA, USA), and identification of

proteins based on the obtained fragmentation spectra using the Peaks Studio 11 (Bioinformatic

Solutions) and the appropriate protein database (Supplementary Table S1) (1% FDR). The list of

annotations for the MS-identified proteins was searched for phage-associated functions, and the

encoding genes were mapped to the genomes of the tested strains. Tailocin-encoding regions

were identified based on the clustering of the identified phage/tailocin-related genes within the

genomes of the analyzed strains and previous studies concerning the structure of the P2D1 tailocin

cluster (Borowicz et al., 2023). As the full sequence of the carotovoricin Er cluster is unavailable,

the originally uploaded sequence (accession: AB017338.2) (Yamada et al., 2006) annotated using

RAST (https://rast.nmpdr.org/rast.cgi) (Brettin et al., 2015) was used for cluster synteny analysis.

Phylogenetic and bioinformatic analyses

Synteny of tailocin clusters

Gene clusters encoding tailocins were extracted from the genomes of *Pectobacterium*, *Dickeya,*

and *Musicola* species, except for *Pectobacterium odoriferum* A122-S21-F16, where the putative

cluster was split across multiple contigs. For *Pectobacterium* spp. analyzed, the genomic region

containing the tailocin cluster was defined as the region between the genes annotated as *tolC*

(locus tag in *P. versatile* A69-S13-O15: EG331_00440) and *ybiB* (locus tag in *P. versatile* A69-

S13-O15: EG331_00565). In contrast, for *Dickeya* species and a single *Musicola* sp. strain, the

genomic region containing the tailocin cluster was identified as the region between the genes

annotated as *methyl-accepting chemotaxis protein* (locus tag in *D. dadantii* 3937:

DDA3937_RS1200) and *guaD* (locus tag in *D. dadantii* 3937: DDA3937_RS12145).

Accession numbers of genomes from which the tailocin regions were extracted are listed in

Supplementary Table 1. The extracted sequences were then subjected to synteny analysis and

visualization using the Clinker tool (Gilchrist and Chooi, 2021). The analysis was performed

separately for *Pectobacterium* and *Dickeya* species, along with a single strain of *Musicola* sp.

Genes within the clusters were color-coded based on their putative roles in tailocin production. In

the visualization, the presence of links between genes indicates sequence identity percentages

greater than 50%. The order of strains in the visualization was determined according to their

phylogenetic relationships. The annotation in the synteny analysis was based on *Pectobacterium*

versatile for *Pectobacterium* spp. and *Dickeya dadantii* 3937 for *Dickeya* spp.

Multilocus sequence analysis

A subset of strains was used for the phylogenetic analysis. The selected strains had to possess the

following features: (1) the tailocin cluster had to be present on a single genomic contig; (2) genes

annotated as fiber and sheath had to be present within the region identified as the tailocin cluster;

and (3) the strain had to demonstrate the ability to target at least one other analyzed strain through

tailocin interaction. Orthologous sequences were clustered into homologous families using the

SiLix software package v1.2.9 (Miele et al., 2011) with an 80% identity threshold and at least 80%

overlap. Orthologous sequences (658 coding sequences) were concatenated, aligned using

MUSCLE (Edgar, 2004) software v5.1, and filtered using the GBLOCK tool (Castresana, 2000).

The alignments were used to build a phylogenetic tree with the BioNJ algorithm with SeaView

software v5.0.5 (Gouy et al., 2010), with 200 bootstrap replications.

Phenotype-based dendrogram

Agglomerative clustering (Nielsen, 2016) was applied to prepare a dendrogram for tested strains

based on the similarity of their killing spectra. The ability to kill a given strain was considered 1, and

the lack of this ability was rated 0. The scikit-learn (Virtanen et al., 2020) and SciPy (Virtanen et al.,

2020) libraries were used for clustering and dendrogram plotting, respectively. A Phyton script for

implementing the above libraries was written with the assistance of ChatGPT-4o (OpenAI)

(Supplementary Dataset 3).

 Prevalence of carotovoricin and dickeyocin clusters in SRP complete genomes

The carotovorocin cluster was blasted against 116 *Pectobacterium* spp. complete, high-quality

genomes available on the NCBI database in June 2024 using the NCBI Megablast algorithm with

default parameters (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The presence of the cluster was

defined according to the 70% sequence coverage and identity thresholds. The available sequence

of the originally described carotovoricin Er cluster was incomplete (accession: AB017338.2).

Therefore, as a reference carotovoricin sequence for Pectobacterium spp. we used the full

nucleotide sequence of the carotovoricin cluster from *P. versatile* A73-S18-O15 (19,528 bp.). For

Dickeya spp. and the single *Musicola* sp. strain, the full nucleotide sequence of the dickeyocin

cluster of *D. dadantii* 3937 strain (26,580 bp) was blasted against a single *Musicola* sp. and 74

Dickeya spp. complete genomes available on the NCBI database in June 2024. Results were

interpreted based on analogous thresholds as described for P2D1 tailocin.

Detection of tailocins in river water and potato tubers

To determine whether SRPs produce detectable levels of tailocins in river water and potato tuber

tissue, we inoculated these environments with bacteria. After incubation, we purified the tailocins

and conducted spot tests as outlined below. Each experiment was performed twice, and detection

was considered positive when all replicates gave positive results. To prepare the inoculation

suspensions, six strains, namely *P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P.*

aquaticum A212-S19-A16, *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, and *D. zeae* A586-

S18-A17, were grown in TSB (Oxoid) for 24 h at 28 °C. The cells were harvested by centrifugation

(4200 RCF, 5 min) and resuspended in PBS buffer. Turbidity of the suspension was adjusted to

0.06 McF (approx. 10^6 CFU mL^{-1}). Five ml of Durance River water filter-sterilized and autoclaved

(0.2 µm, PES membrane) were inoculated with 50 µl of bacterial suspension, three replicates per

strain. Uninoculated water was used as a control. Samples were incubated for 72 h at 15 °C with

shaking (120 rpm) to mimic natural conditions. Following incubation, 2 ml of water from each

sample was centrifuged to remove bacterial cells (8000 RCF, 10 min), and the supernatant was

processed for purification and detection of tailocins. Tubers cv. Gala were surface sterilized by

immersion for 20 min in 5% commercial bleach (ACE, Procter and Gamble), followed by a double

rinse in distilled water and air drying under laminar flow. Each tuber was inoculated by inserting a

pipette tip containing 50 μl of the test suspension into the tuber (up to the level of liquid within the

tip). Tubers inoculated with PBS buffer alone were used as a negative control. Three potato tubers

were tested per treatment. Inoculated tubers were placed in humid boxes (85 to 90% relative

humidity). Samples were incubated at 28°C for 72 h to enable the development of soft rot

symptoms. Next, the tubers were cut at the inoculation site. Tissue macerated by bacteria was

spooned out, weighed, and placed in homogenization bags (Bioreba). PBS buffer supplemented

with 0.02% diethyldithiocarbamic acid (DIECA) (Sigma-Aldrich) was added to the samples (1:2,

w/v) (Perombelon and van Der Wolf, 2002). The samples were homogenized, and 1 mL aliquots

were diluted 1:1 with PBS 0.02% DIECA in new tubes. Bacteria and plant debris were pelleted by

centrifugation (8000 RCF, 10 min, RT), and the supernatant was processed for purification and

detection of tailocins. The supernatants from both experimental setups were filtered (0.2 μm, PES

membrane) and transferred to new tubes containing PEG-8000 (final concentration 10 % m/v).

Samples were incubated overnight at 4°C with shaking. The putative tailocins were collected by

centrifugation (1 h, 16 000 RCF, 4°C) and resuspended in 1/20 volume of the initial sample in PBS.

Tailocins were detected using a spot test (Hockett and Baltrus, 2017;Yao et al., 2017). Five μL

aliquots of purified tailocins were spotted on soft top agar plates, each inoculated with a strain

sensitive to tailocins of the six investigated SRPs (Supplementary Table S2).

Semiquantitative assessment of tailocin production depending on the type of inductive

factor

We investigated whether hydrogen peroxide can induce the production of tailocins in SRPs in a

way similar to mitomycin C. The experiment was conducted on the same group of six strains tested

for tailocin production in river water and potato tuber tissue. Briefly, an overnight culture (ca. 16 h)

of each strain was rejuvenated (1:40) in a fresh aliquot of TSB and cultured for the next 2.5 hours.

After that, tailocin production was induced by the addition of 0.1 μg mL^{-1} mitomycin C (Abcam,

Poland) or 0.003 % (0.88 mM) hydrogen peroxide (Laboratorium Galenowe Olsztyn Sp. z o. o.,

PolandCultures) and cultures with no inductive factor were used as controls. Twenty-four hours

post-induction, tailocins were PEG-purified from 2 mL volumes of filtered (0.2 μm, PES membrane)

culture supernatants and resuspended in 1/10 of the initial volume in PBS (Borowicz et al., 2023).

The concentration of tailocins was estimated by a semiquantitative spot test (Yao et al.,

2017;Borowicz et al., 2023) using tailocin-susceptible strains (Supplementary Table S2). The

reciprocal of the highest dilution causing a visible plaque was defined as the value of the relative

activity in arbitrary units (=1 AU), and the AU per mL of culture was calculated. The experiment

was conducted twice, with tree replicas per combination.

Competition assay *in vitro*

To perform the competition assay, we first selected a spontaneous rifampicin-resistant mutant of *D.*

solani A623-S20-A17 on TSA with 50 μg mL^{-1} of rifampicin (Sigma-Aldrich)(Glandorf et al.,

1992;Berg et al., 2007). Such obtained strain was designated *Ds* RIF and tested in an *in vitro*

competition assay against three sets of competitor strains. Each set comprised two *Dickeya* spp.

strains which either: 1) were killed by the tailocins of *D. solani* A623-S20-A17 (*D. dianthicola* A260-

S21-A16 and *D. zeae* A661-S21-A17), 2) produced tailocins killing *D. solani* A623-S20-A17 (*D.*

zeae A586-S18-A17 and *D. oryzae* A3-S1-M15) or 3) were killed by the tailocins of *D. solani* A623-

S20-A17 but at the same time produced tailocins targeting this strain (bilateral killing)(*D. dadantii*

3937 and *D. chrysanthemi* A604-S21-A17).

All strains were grown overnight in 0.1 TSB medium. Next, the cells were harvested by

centrifugation (3500 RCR, 5 min.), and the pellet was resuspended in a fresh medium to obtain a

turbidity of 0.5 McF. This equaled, on average, 1.4×10^8 CFU ml^{-1} ($\pm 4.5 \times 10^7$), verified by dilution

plating. 150 μl of 0.1 TSB in a single well of a 96-well plate was inoculated (1:1, 25:25 μl) with the

suspension of *Ds* RIF and one of the competing strains to ensure the strains had an even start and

avoid tailocins carryover from the initial overnight cultures. Each co-culture was tested under two

conditions: 0.1 TSB alone or 0.1 TSB with 0.003 % (0.88 mM) H_2O_2 (tailocin-inductive stress). The

cultures were incubated for 22 h at 28°C, with orbital shaking, in the EPOCH2 reader (BioTek).

Culture from each well was dilution-plated on two media types: TSB agar (for counting the total

number of CFUs) and TSB agar with 50 µg mL^{-1} of rifampicin (for counting the fraction of *Ds* RIF).

Ds RIF was tested against the wild-type (*wt*) variant of the strain as a control of the mutant's fitness

under the applied experimental conditions. Moreover, the ability of *Ds* RIF to produce tailocins was

verified in a spot assay, as described above. The competition experiment was performed twice,

with two separate co-cultures (replicates) per experiment.

Results

***Pectobacterium* spp. and *Dickeya* spp. strains from Durance River produce phage tail-like**

particles

TEM imaging of preparations obtained from mitomycin C-induced cultures revealed that 22 out of

25 studied SRP isolates (88%) produce phage tail-like particles (tailocins) (Fig.1). The three

exceptions were *Pectobacterium brasiliense* A143-S20-M16, *Pectobacterium atrosepticum* A597-

S4-A17, and *Dickeya dianthicola* A260-S21-A16, for which no such structures could be detected in

our experimental setup. All detected tailocins showed similar morphology in TEM and AFM images,

regardless of whether they were derived from *Dickeya/Musicola* or *Pectobacterium* spp. (Figures 1

and 2). The average length of tailocins produced by *Pectobacterium* spp. was 163±15 nm.

Tailocins produced by *Dickeya/Musicola* spp. were slightly shorter, measuring 153±20 nm, with the

difference being statistically significant (Mann-Whitney test, $p < 0.05$). Similarly, tailocins produced

by *Pectobacterium* spp. were, on average, 2 nm wider compared to those of *Dickeya/Musicola*

species, with the dimensions amounting to 26±2 nm and 24±2 nm, respectively, with the difference

being statistically significant (Mann-Whitney test, $p < 0.05$). Three-dimensional images generated

based on AFM scans for seven representative tailocin particles are shown in Fig. 2.

The killing spectra of SRP phage tail-like particles vary depending on the strain of origin, with several cases of inter-genus activity

We evaluated in detail the killing spectra of phage tail-like particles produced by studied SRPs strains. In total, we investigated 702 unique tailocin-mediated interactions (producer-target combinations) for 351 unique pairs (individual producer vs. individual target) of SRPs (Fig. 3). The overall incidence of killing events where a strain was found susceptible to a particular tailocin equaled 16.8% (Table 1). The killing incidence was higher in the tested subset of *Dickeya* strains than in the tested *Pectobacterium* strains, at 57.1% and 21.6%, respectively (Table 1). The two most susceptible strains were *P. versatile* A73-S18-015 and *P. polaris* A641-S16-A17, killed by tailocins from 8 different strains. On the opposite side of the spectrum was *P. carotovorum* A280-S2-N16, which was resistant to tailocins produced by all strains in the study. Twenty-one of the 25 environment-derived strains produced tailocins that killed at least one strain from the tested group of 27 SRPs. Out of the 4 strains producing no killing events, three (*P. brasiliense* A143-S20-M16, *P. atrosepticum* A597-S4-A17, and *D. dianthicola* A260-S21-A16) were the ones with no tailocins visible in TEM analyses (Fig. 1), and *P. peruviense* A350-S18-N16 which did not target any of the 27 SRPs tested (Fig. 3). On the contrary, *D. oryzae* A3-S1-M15 and *P. odoriferum* A122-S21-F16 produced tailocins with the broadest killing range, each targeting 9 out of the 27 strains (33.3%).

We also investigated the inter-genus activity of tailocins. *Pectobacterium* strains produced tailocins targeting solely other members of the genus. On the contrary, 5 out of 8 tested *Dickeya* spp. (62.5%) could target at least *one Pectobacterium* sp., in each case, either a *P. versatile* strain or the closely related *P. odoriferum*. *M. paradisiaca* NCPPB 2511, the single representative of *Musicola* spp., strongly clustered with *Dickeya* spp. both regarding tailocin susceptibility and the specificity of the produced tailocins (Fig. 3, Table 1).

Several SRP strains produced tailocins targeting one another – a phenomenon we named 'bilateral killing'

Analysis of tailocin-mediated interactions in the tested pool of SRPs revealed that 33.9% of the killing events were retaliated (Table 1). Ergo, in the case of 20 of the 351 investigated pairs of SRPs (5.7%), the two strains could target and kill one another with their tailocins. We called the phenomenon 'bilateral killing'. Furthermore, the phenomenon of bilateral killing was more common within the tested subgroup of *Dickeya* spp. (65.6% of killing events retaliated) than within the tested pool of *Pectobacterium* spp. (15.2%), and we observed no bilateral killing between the members of *Dickeya* and *Pectobacterium,* where only unilateral targeting was reported (Table 1). The retaliated killing was very common (80% of killing events) for the mixed pairs of *Dickeya* strains and *M. paradisiaca* NCPPB 2511, once again showing clustering between this single NCPPB 2511 strain and *Dickeya* spp.

Tailocins of *Dickeya* spp. and *Pectobacterium* spp. are produced from a different ancestral origin

Based on the tailocin-associated protein identified by mass spectrometry, we identified gene clusters responsible for the tailocin production in the genomes of the analyzed SRP strains. Among the 25 strains examined, tailocin-encoding clusters were absent only in 2 strains from the analyzed SRP pool, namely in *Pectobacterium brasiliense* A143-S20-M16 and *Pectobacterium atrosepticum* A597-S4-A17. These two strains, together with *Dickeya dianthicola* A260-S21-A16, were also found to be negative for tailocin production in microscopic experiments.

We also performed synteny analyses of the tailocin clusters within the *Pectobacterium* and *Dickeya* (including *Musicola*) genera (Fig. 4). These analyses in *Pectobacterium* sp. revealed the homology of the clusters to the one encoding carotovoricin Er (Nguyen et al., 2002) while those in *Dickeya* species and *Musicola* sp. were genetically similar to dickeyocin P2D1 (Borowicz et al., 2023). Furthermore, no SRP strain has been found to possess both two carotovorocin Er and P2D1 clusters tailocin simultaneously. The *Pectobacterium* spp. strains exclusively contained

carotovorocin Er cluster, while *Dickeya* spp. strains solely harbored cluster encoding P2D1 tailocin.

Thus, no strain that contains both types of tailocins has been identified.

The overall genomic organization of the tailocin clusters was conserved within each genus (Fig. 4).

In *Pectobacterium* species (Fig. 4A), the most prominent difference was a highly variable gene

annotated as encoding a tail fiber protein/pyocin knob domain-containing protein, which appeared

to contain segments shuffled among different strains (data not shown). Additional, albeit less

pronounced, variability was observed in the sequence of the gene coding for a tail tape measure

protein. In the case of *P. brasiliense* A143-S20-M16, the cluster region was flanked by genes

conserved across other strains; however, instead of a complete tailocin cluster, it contained only a

single annotated gene: a mobile element protein gene. In *P. atrosepticum* A597-S4-A17, the region

included not only conserved flanking genes but also seven genes present in other clusters and one

strain-specific gene. In *Dickeya* species (and *Musicola* sp.) (Fig. 4B), the main differences between

strains included (1) the presence of strain-specific genes likely unrelated to tailocin production, (2)

the orientation of genes not directly involved in tailocin structural components (such as aromatic

acid/H+ transporters and phytase family proteins), (3) sequence variations in intergenic regions

(data not shown), and (4) partial sequence differences in tail fiber genes. Notably, for *D. dianthicola*

A260-S21-A16, which did not produce tailocins, no significant differences (= the cluster was

present in the genome and expressed a high homology to the other Dickeya spp. tailocin clusters)

were observed in the tailocin region composition in comparison to other studied *Dickeya* spp.

Carotovoricin and dickeyocin clusters are common in the genomes of *Pectobacterium* spp.

and *Dickeya* spp.

In our pool of analyzed SRPs, not all strains produced tailocins, nor did they harbor the tailocin

cluster. This prompted us to determine the prevalence of tailocin clusters across a broader range

of SRP strains. We researched 116 complete genomes of *Pectobacterium* spp. for the presence of

the carotovoricin cluster and 74 genomes of *Dickeya* and 1 *Musicola* spp. genome for the presence

of genes encoding dickeyocin. Within the *Pectobacterium* clade, we identified 96 genomes with

homologous regions, meeting the adopted >70% thresholds for both coverage and identity, making

it 83% of all tested strains. Twelve genomes showed alignment below the threshold criteria (10% of

strains), and 9 (8%) showed no alignment (Dataset S1). Within the joint pool of *Dickeya* spp. and

the single *Musicola* spp. genome, we found 52 genomes with an alignment to dickeyocin cluster

meeting the 70% threshold (69% of strains), 16 genomes with coverage below 70% (21% of

strains), and 7 genomes without alignment (9%) (Dataset S2). Of note, the 6 analyzed complete

genomes of the species *P. atrosepticum* present in the NCBI database do not contain a tailocin

cluster.

Tailocin-killing spectrum activity does not precisely follow the phylogenetic distance of the

analyzed SRP strains

When comparing the topology of two dendrograms, a phylogenetic tree based on MultiLocus

Sequence Analysis (MSLA) and a dendrogram based on the killing spectrum phenotype, our

results indicate that the killing spectrum of tailocins does not correlate strictly with the phylogenetic

relationships among the tested strains. This suggests that tailocin activity may be influenced by

factors other than taxonomic position (Supplementary Fig. S1).

Basal level of tailocin production occurs in both *Dickeya* and *Pectobacterium* spp., and a

significant induction occurs in the presence of hydrogen peroxide

Our previous study reported a basal level of tailocin production in a population of *D. dadantii* 3937

not exposed to tailocin-inducive stress (Borowicz et al., 2023). Here, on a pool of six strains,

including 3 *Pectobacterium* and 3 *Dickeya* spp. (*P. versatile* A69-S13-O15, *P. quasiaquaticum*

A411-S4-F17, *P. aquaticum* A212-S19-A16, *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D.*

zeae A586-S18-A17), we aimed to determine whether the basal level of tailocin production occurs

in other SRPs. Moreover, we wished to establish whether reactive oxygen species (ROS), namely

H_2O_2, induce the production of tailocins in SPRs. As SRPs frequently encounter ROS during

infection, with this setup, we wanted to hypothesize whether tailocins may be produced *in situ*

during infection (Torres et al., 2006;Reverchon et al., 2016). Basal level of tailocin production was

observed for all tested strains. However, the titer of detected tailocins varied between strains, with the highest basal production level observed for our model strain *D. dadantii* 3937 (7×10^4 AU mL^{-1}) and the lowest for *P. aquaticum* A411-S4-F17 (5×10^2 AU mL^{-1}) (Fig. 5). For our group of strains, the average tailocin titer was slightly (43%) higher for *Dickeya* than for *Pectobacterium* spp. (Mann-Whitney test, $p<0.05$), with 2.8×10^4 and 1.6×10^4 AU mL^{-1} of culture, respectively.

Mitomycin C (0.1 µg mL^{-1}, control) caused a significant induction (approx. 5 to 20-fold) of tailocin production in all strains except *P. quasiaquaticum* A411-S4-F17, for which the increase was not statistically significant compared to no induction (Fig. 5). The highest induction was observed for *D. zeae* A586-S18-A17 (19-fold) and *D. chrysanthemi* A604-S21-A17 (20-fold). In the case of *D. dadantii* 3937, the induction was relatively low, only 5-fold. However, this low fold change was not connected with low tailocin titer post-induction but resulted from a high basal production level.

Hydrogen peroxide-induced tailocin production in all tested strains (approx. 15 to 81-fold) (Fig. 5). For most strains, the applied concentration of hydrogen peroxide (0.88 mM, 0.003%) resulted, on average, in a higher yield of tailocins than when induced by mitomycin C (0.1 µg mL^{-1}). The exceptions were *D. chrysanthemi* A604-S21-A17 and *D. zeae* A586-S18-A17, with comparable results for both inducers. Notably, in the case of *P. quasiaquaticum* A411-S4-F17, for which no induction was observed upon treatment with mitomycin C, hydrogen peroxide not only caused induction of tailocin production but also resulted in the highest observed increase compared to basal production levels (81-fold) (Fig. 5).

Detectable levels of phage tail-like particles are produced in rotting potato tuber tissue but not in river water

Six bacterial strains as described earlier, namely three *Pectobacterium* spp: *P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16 and three *Dickeya* spp.: *D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D. zeae* (very recently reclassified to *D. parazeae* (Hugouvieux-Cotte-Pattat and Van Gijsegem, 2021)) (A586-S18-A17) were used to inoculate two types of samples: river water (being the isolation source of the strains) and potato tuber tissue

(with plant tissue being the primary niche for many SRPs). The goal was to determine whether

bacteria in these environments produce noticeable levels of tailocins.

Under the applied conditions, tailocin production was not detected in inoculated water samples

(Table 2.). On the contrary, tailocins were detected in macerated potato tuber tissue, not just in the

case of four strains that caused soft rot symptoms (*P. versatile* A69-S13-O15, *D. dadantii* 3937, *D.*

chrysanthemi A604-S21-A17, *D. zeae* A586-S18-A17) but also in healthy tuber tissue surrounding

the inoculation site in the case of two strains which failed to cause potato tuber rotting (*P.*

aquaticum A212-S19-A16 and *P. quasiaquaticum* A411-S4-F17) (Table 2) (Supplementary Fig.

S2).

Producing tailocins alone does not ensure success against a susceptible strain under

conditions of an 'even start'

We aimed to determine how *D. solani* A623-S20-A17 (*Ds*) performs in two-strain co-cultures with

strains susceptible to the strain's tailocins or/and targeted by tailocins of the competitors (Fig. S3).

To monitor the proportion of *Ds* relative to competing strains in mixed cultures, we selected a

spontaneous rifampicin-resistant mutant of *Ds* (*Ds* RIF). Using *Ds* RIF in the experiments allowed

us to determine the percentage of *Ds* in the co-culture using selective media. Prior to the main

experiment, rifampicin-resistant mutant *Ds* RIF was tested against the wt strain (*D. solani* A623-

S20-A17) to produce tailocins, growth rate, and ability to compete with the wt *in vitro*. The mutant

was not compromised regarding the growth rate in 0.1 TSB (Supplementary Fig. S4), was positive

for tailocin production, and neither strain achieved dominance in a co-culture with its wt variant.

These results indicate that the rifampicin-resistant *Dickeya* spp. strains are not compromised in

competitiveness under the applied experimental conditions (Fig. 6).

Next, the *Ds* RIF strain was set to compete with three types of competitors: susceptible to the

tailocins of *Ds* RIF (SET 1), targeting *Ds* RiF (SET 2), or involved with *Ds* RIF in a bilateral killing-

type interaction (SET 3). The success of a strain was defined as its end-point dominance in the co-

culture (>90% of total CFU) (Fig. S3). Each strain in a co-culture was ensured an even start (1:1

mix). Theoretical prediction of the competition's winner based on tailocin effects was compared to

experimental results. We found that two competitor strains targeting *Ds* RIF with their tailocins had

prevailed over *Ds* RIF (SET 2), according to what would be expected if tailocins had a key role in

the tested setup. However, the Ds RIF outcompeted only a single susceptible strain from SET1

and only in the presence of H_2O_2. This shows that in a competition between SRPs, at least under

the applied *in vitro* conditions and with an even start scenario, tailocins alone do not guarantee the

success of a producer against a susceptible rival.

Discussion

Our study revealed that most SRP strains produce phage tail-like particles, and only three

SRP strains from the pool were tailocin-negative. The *P. brasiliense* A143-S21-A16 and *P.*

atrosepticum A597-S4-A17 lacked tailocin clusters, while *D. dianthicola* A260-S21-A16 had a

complete cluster but did not produce tailocins. This could be due to reliance on other competitive

strategies (Hibbing et al., 2010) or regulation by specific environmental factors not present in the

experiment (Bellieny-Rabelo et al., 2019). Furthermore, we assessed the prevalence of tailocin

clusters in 190 high-quality, complete genomes of SRP bacteria and found that 83% of

Pectobacterium spp. and 69% of *Dickeya* spp. genomes contain clusters encoding tailocins. This

indicates that tailocins are a very common feature among SRP and may suggest that their

production remains important to mediate the ecological success of these bacteria (Ma et al., 2007).

The prevalence of tailocins in SRP aligns with the previous observations documenting their

presence in various bacterial species, including ubiquitously present *Pseudomonas* spp. (Carim et

al., 2021), *Pantoea* spp. (Stice et al., 2023) and *Streptomyces* spp. (Nagakubo et al., 2021). Such

global observations have, however, never been done for Soft Rot *Pectobacteriaceae*.

It has been repeatedly demonstrated that most tailocins exhibit a narrow killing range

typically limited to closely related strains. Still, most of the analysis has been conducted on a small

set of strains either at the species level or the genus level (Granato et al., 2019). Some

observations, however, are that some phage tail-like particles can cross species boundaries (Yao

et al., 2017;Principe et al., 2018). Our work involved a large group of strains encompassing strains

belonging to three different genera tested for mutual tailocin-based interactions. It reveals that

some SRP strains have a broad range of activity, producing tailocins that can kill more than 30% of

the tested strains representing different species. Enlarging the number of tested strains within a

given species complex also expands the spectrum of action of each tested tailocin. One should,

therefore, remain cautious about the accepted narrow killing range of tailocins previously

described.

Interestingly, while some *Dickeya* spp. produced tailocins were able to kill *Pectobacterium*

spp. cells, the opposite was not observed in our study. Due to their evolutionary association with

bacteriophage tails (Ghequire and De Mot, 2015), it is suggested that tailocins might retain some

degree of flexibility in the host range, as known from phage-host interactions (Backman et al.,

2024a). Inter-genus activity may remain an important trait in competition as mixed bacterial

communities involving both genera are frequently observed in the field, and the ability to target a

broader range of competitors could confer a selective advantage (Perombelon, 1988; Charkowski,

2009;2018; Barny et al., 2024).

Likewise, the observation of bilateral killing where two strains reciprocally target each other

with tailocins has not been extensively documented. The higher incidence of bilateral killing within

Dickeya spp. compared to *Pectobacterium* spp. suggests some (yet unknown) ecological factors

may more often promote this behavior in *Dickeya* spp. than in *Pectobacterium* spp. This is

somehow surprising, given that multiple species of SRP bacteria are often found together in the

same environments (Toth et al., 2021a;van der Wolf et al., 2021). In such settings, these various

species would be expected to experience similar ecological pressure and benefit from reciprocal

tailocin production (Shyntum et al., 2019). It may suggest that *Pectobacterium* spp., differently from

Dickeya spp., However, this assumption regarding tailocins still requires experimental validation.

Phage tail-like particles isolated from both SRP genera exhibit similar morphological

characteristics; however, the different origins of tailocin clusters in *Pectobacterium* spp. and

Dickeya spp. indicates that these phage tail-like particles have been acquired independently in

those genera. All analyzed clusters present in *Dickeya* spp. express homology to the one encoding

P2D1 tailocin of *D. dadantii* 3937 (Borowicz et al., 2023), whereas all clusters present in

Pectobacterium spp. express homology to the cluster encoding carotovoricin Er (Yamada et al., 2006). The fact that these phage tail-like particles were observed in all *Dickeya* spp. and most of the *Pectobacterium* spp. strains tested indicated that the tailocin clusters were acquired before species diversification in both genera and were maintained in most species rather untouched. As tailocins are large protein complexes and their synthesis should be a burden for a producer cell, their conservation suggests that tailocin production plays an important role in the ecology of these bacteria (Glasner et al., 2008;Hugouvieux-Cotte-Pattat, 2016).

The genes encoding tailocin fiber proteins expressed the lowest homology in the analyzed tailocin clusters both in *Dickeya* spp. and *Pectobacterium* spp. strains analyzed. This confirms that fiber proteins play a pivotal role in determining the specificity of tailocins and their killing range, as reported earlier (Dams et al., 2019). Fiber proteins are the most important determinants in recognizing target bacterial cells (Backman et al., 2024a). This specificity is crucial for the success of tailocins as killing agents, enabling them to discriminate between susceptible and resistant bacterial strains. In bacteriophages, receptor-binding proteins are known to undergo positive selection to adapt to variations in surface receptors of target bacteria, leading to high variability even among closely related strains (Bertozzi Silva et al., 2016). In SRP, the killing spectrum of tailocins, determined by the sequence of fiber proteins, appears to be more dependent on ecological factors and the pressure of host-pathogen interactions encountered by the individual strains than the evolutionary relationships between them.

Although mitomycin C is widely used to trigger tailocin induction under laboratory conditions, in natural and agricultural environments, the likely inducers of tailocins remain unknown. In the proof-of-concept experiment, we determined that hydrogen peroxide (a reactive oxygen species agent) is a potent inducer of SRP phage tail-like particles. It is well acknowledged that reactive oxygen species are the common signals for the induction of phage tail-like particles in bacteria (Scholl, 2017). In general, hydrogen peroxide is known to activate the ROS response (Storz et al., 1990;Erill et al., 2007), which is sequentially a recognized tailocin induction pathway. During infection, SRP bacteria often encounter oxidative stress in plant hosts when the plant mounts an immune response (Reverchon and Nasser, 2013;Jiang et al., 2016). Therefore, the

induction of tailocins by reactive oxygen species may serve as a mechanism for bacterial survival

in their natural settings, aiding in eliminating competing bacteria under stressful conditions.

Globally, the ability of plant pathogenic bacteria, including SRP, to produce tailocins both at a basal

level and under stress (inducive conditions), such as exposure to ROS, may lead to important

ecological consequences. The ability to regulate tailocin production in response to such oxidative

stress might give SRP a competitive edge, allowing them to eliminate competitors in the plant

environment when they must fight with all other microorganisms to establish a successful infection

(Backman et al., 2024b).

The variable level of response to mitomycin C and hydrogen peroxide we observed in SRP

strains highlights the still unraveled complexity of tailocin regulation. Among the tested strains, *D.*

chrysanthemi A604-S12-A17, *D. zeae (parazeae)* A586-S18-A17, and *P. versatile* A69-S13-O15

showed the highest increase in tailocin production with mitomycin C. In contrast, hydrogen

peroxide led to the highest production in *P. quasiaquaticum* A411-S4-F17. This suggests that in

SRP, depending on the strain and type of inducer, the production of tailocins is slightly differently

regulated. Similar observations were made for prophages, where different environmental factors

induced phages with different efficiency (Casjens, 2003;Varani et al., 2013).

SRP strains tested in this study were isolated from one selected environment. The set

comprised both aquatic and non-aquatic strains, most of which (except *P. aquaticum* and *P.*

quasiaquaticum strains known to be avirulent on plants) cause symptoms in plants when

transmitted to agricultural settings (Ben Moussa et al., 2022). We found that tailocins are produced

in infected potato tubers but not in river water. This aligns with the other studies in which tailocins

were found to be produced in nutrient-rich, competitive environments (Carim et al., 2021), like

decaying plant material, rather than in low-stress environments like river water (Heiman et al.,

2023). The finding that tailocins were detected in (rotting) potato tissue but not in river water

suggests that the environmental conditions in the potato tubers provide a stronger stimulus for

tailocin production, likely due to the presence of ROS, plant exudates, cell wall degradation

products, and microbial competition, as opposed to water which for many strains is neither the

target niche nor the competitive environment but rather serves as a translocation medium from one

host to another (Pédron and Van Gijsegem, 2019;van der Wolf et al., 2021). This observation may

be particularly interesting in the case of SRP bacteria, which compete with each other and other

microorganisms for resources present within the plant tissues (Reverchon and Nasser, 2013). The

potato tuber tissue represents a primary ecological niche for SRP pathogens (Perombelon and

Kelman, 1980;Pérombelon and Salmond, 1995). Consequently, the presence of tailocins in both

rotting tissue and the healthy surrounding tissue suggests that tailocin production may be part of

the virulence strategy to minimize the risk of competitive infections.

Surprisingly, SRP strains belonging to species that are not found on plants (Portier et al.,

2020) and do not provoke evident disease symptoms (*vide*: *P. quasiaquaticum* A411-S4-F17 and

P. aquaticum A212-S19-A16) still produce tailocins in the plant environment but not in river water.

The low nutrient availability, the cost of production of this complex machinery and the few direct

competitors of SRP bacteria in river water likely explain the absence of tailocin production in this

environment (Lamichhane and Bartoli, 2015) (Bradford et al., 2013;Toth et al., 2021b). The fact

that these tailocin clusters were not eliminated from these water-associated species and were

induced by the same trigger as plant-associated species suggests that tailocin production is not

solely tied to SRP virulence on plants but could also serve in another niche where these bacteria

encounter stress. It has been suggested that these water-associated species could be associated

with insect larvae in the water stream (Ben Moussa et al., 2023). Whether tailocin production is

triggered in such an environment remains to be determined.

Multiple factors play roles in the competition between microbes, with success being the sum

of factors/traits of varying importance depending on the environmental context (Hibbing et al.,

2010). In our study, in mixed populations of tailocin producers and targets, tailocins were not

enough to ensure the dominance of producers over susceptible strains under the scenario of 'even

start'. In conjunction with how widespread tailocin clusters are among SRP, this led us to conclude

that tailocin-mediated success is often conditional. It is well known that SRP bacteria use diverse

weaponry for intra-genus and intra-genus interactions; therefore, tailocin production may be just

one possibility to compete with kin microbes.

Several factors can influence the outcome, including environmental conditions, strain-

specific interactions, and the relative abundance of producers versus susceptible strains. This

study explored an 'even start' scenario, where tailocin-producing and susceptible competitors were

mixed in equal cell numbers. It was shown that in bacterial populations, even when one strain has

an antagonistic advantage (like producing phage tail-like particles), an equal start condition can

neutralize this advantage because the susceptible strain is not immediately overwhelmed (Ross-

Gillespie and Kummerli, 2014). Moreover, tailocin production, unlike the secretion of most bacterial

antimicrobials, requires the lysis of the producer cells. Ergo, it can negatively influence the growth

rate of the producer's population. This burden can be assumed negligible when the high producer-

to-target ratio is high. Therefore, we speculate that tailocin production may be useful to SRP

bacteria to protect already conquered niches from invading competitors, a process named niche

exclusion (Bauer et al., 2018).

To conclude, our study considerably advances the understanding of the role of phage tail-

like particles in the ecology of Soft Rot *Pectobacteriaceae* (SRP). Furthermore, our study expands

the knowledge of tailocin function in inter-genus competition among SRPs, their responsiveness to

environmental factors, and their potential role in bacterial survival in various ecological niches.

Author Contributions

MB: Conceptualization, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing, DMK: Conceptualization, Investigation, Supervision, Visualization, Writing – original draft, Writing – review & editing, MS: Investigation, Methodology, Visualization, MN: Investigation, Methodology, Visualization, Writing – original draft, IM: Investigation, Methodology, Writing – original draft, PC: Methodology, Writing – original draft, JP: Conceptualization, Investigation, Methodology, Writing – review & editing, MAB: Conceptualization, Investigation, Methodology, Writing – review & editing, PYC: Investigation, Methodology, JD: Conceptualization, Investigation, Methodology, RC: Conceptualization, Data curation, Funding acquisition, Resources, Supervision, Writing – original draft, Writing – review & editing

Competing Interest Statement

The authors declare no competing interests.

Acknowledgments

This research was financially supported by the National Science Center, Poland (Narodowe Centrum Nauki, Polska) via a research grant SONATA BIS 10 (2020/38/E/NZ9/00007) to Robert Czajkowski.

 Tables

670 **Table 1. Tailocin-mediated killing events for the tested pool of 27 SRP strains.**

Group	Total interactions[A]	Killing events		Bilateral killing		
		No.	% total	No.	% total	% killing events
All SRPs	702	118	16.8%	40	5.7%	33.9%
Pectobacterium spp.[B]	306	66	21.6%	10	3.3%	15.2%
Dickeya spp.[C]	56	32	57.1%	21	37.5%	65.6%
Mixed genera[D]	340	19	5.6%	9	2.6%	47.37%
Psp vs *Dsp*	162	0	0%	0	0%	0%
Dsp vs *Psp*	162	6	3.7%	0	0%	0%
Dsp vs *M. par.*	8	5	62.5%	4	50%	80%
M. par vs *Dsp*	8	6	75%	5	62.5%	83.3%
M. par vs *Psp*	18	2	11.1%	0	0%	0%
Psp vs *M. par*	18	0	0%	0	0%	0%

[A] total interactions - the number of combinations where each strain from a subset was tested both for the
potential to produce tailocins and for tailocin susceptibility (potential target)
[B] both strains in the tested interaction assigned to *Pectobacterium* spp. (total of 18 strains tested)
[C] both strains in the tested interaction assigned to *Dickeya* spp. (total of 8 strains tested)
[D] each strain of the pair representing a different genus; *Psp* – *Pectobacterium* spp.; *Dsp* – *Dickeya* spp.; *M.*
par – *Musicola paradisiaca* NCPPB 2511. The first strain in each combination is the tailocin producer.
678
679
680

Table 2. Production of tailocins in river water and potato tubers. Qualitative tailocin
production in different conditions for various bacterial strains. The strains tested include *P.*
versatile A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16, *D.*
dadantii 3937, *D. chrysanthemi* A604-S21-A17, and *D. zeae* A586-S18-A17. Tailocin
production was assessed in infected potato tubers, in Durance River water, and as a positive
control following Mitomycin C and H_2O_2 induction in TSB (Tryptic Soy Broth). The symbols
represent the following: "+" indicates detectable tailocins in the sample with the applied
method; "-" indicates undetectable tailocins in the sample with the applied method; "*"
indicates that tailocins were detectable if tuber tissue in direct contact with inoculated
bacteria was pooled from three potato tubers.

Strain	Tailocins production			
	In Durance River water	In infected potato tuber	Mitomycin C induction in TSB	H_2O_2 induction in TSB
P. versatile A69-S13-O15	-	+	+	+
P. quasiaquaticum A411-S4-F17	-	+*	+	+
P. aquaticum A212-S19-A16	-	+*	+	+
D. dadantii 3937	-	+	+	+
D. chrysanthemi A604-S21-A17	-	+	+	+
D. zeae A586-S18-A17	-	+	+	+

+ tailocins detectable in the sample in the applied method

- tailocins undetectable in the sample in the applied method

* tubers were asymptomatic, but tailocins were detectable in tuber tissue at the inoculation site; the material was collected and pooled from three potato tubers.

Figures

Fig. 1 TEM images picturing tailocins isolated from the 27 analyzed SRP strains.

Fig. 2 Images of the representative tailocins generated based on AFM scans. AFM imaging was performed using the JPK NanoWizard® 4 (NanoScience) in QI™ in the quantitative imaging (QI)™ force spectroscopy mode. Images were processed using JPK Data Processing software. Panel A shows the AFM scan of Dickeyocin P2D1 produced by *D. dadant*ii 3937, panels B-H. show the highest quality 3D visualizations of representative tailocin particles, generated based on AFM scans, for the following strains: *D. dadantii* 3937, *D. solani* A623-S20-A17, *D. zeae* A661-S21-A17. *M. paradisiaca* NCPPB 2511, *P. odoriferum* A367-S5-N16, *P. aquaticum* A105-S21-F16, *P. aquaticum* A104-S21-F16.

Fig. 3 Matrix of tailocin-mediated interactions within the group of tested SRP strains.
The tested pool comprised 27 strains: 25 isolates from the Durance River water and two type
strains: *D. dadantii* 3937 and *M. paradisiaca* NCPPB 2511 (in bold). Grey-shaded cells
indicate the susceptibility of a target strain to the tailocins of a given producer strain
(unilateral killing). Unilateral killing where the producer and the target belong to different
genera, is additionally marked by a yellow dot. Lillac cells also indicate the susceptibility of
the target strain to the investigated tailocin, but in particular situations where the target strain
was observed to produce 'retributive' tailocins against the producer (bilateral killing). A
dedicated column indicates whether tailocins were (Y) or were not (N) detected by
microscopic imaging in culture supernatants of the tested strains following treatment with
mitomycin C.

Fig. 4 Comparison of tailocin clusters in the analyzed group of SRPs. The upper panel (A) depicts regions homologous to the carotovoricin-encoding cluster typical for *Pectobacterium* spp. (reference: Carotovoricin Er accession: AB045036.1, *P. versatile* A69-S13-O15 RMBQ01000001.1, positions in the genome: 94551-113781). The lower panel (B) shows regions homologous to dickeyocin P2D1, typical for *Dickeya* spp. and also present in *Musicola paradisiaca* NCPPB 2511 (reference: *D. dadantii* 3937 NC_014500.1, positions in the genome 2730482-2757061).

B

Fig. 5 Basal and induced tailocin production levels in six *Pectobacterium* spp. and *Dickeya* spp strains. The graph shows the logarithm of the arbitrary tailocin units per mL of culture (Log AU mL^{-1}) for three strains of *Pectobacterium* spp. (*P. versatile* A69-S13-O15, *P. quasiaquaticum* A411-S4-F17, *P. aquaticum* A212-S19-A16) and three strains of *Dickeya* spp. (*D. dadantii* 3937, *D. chrysanthemi* A604-S21-A17, *D. zeae* A586-S18-A17). The basal production levels (no induction) are compared to those with mitomycin C or hydrogen peroxide after induction. Averages from 2 experiments, each consisting of 3 separate inductions, are shown, and error bars represent the standard deviation.

Fig. 6 Competition between *D. solani* A623-S20-A17 RIF^R (*Ds* RIF) and other SRP strains in two-strain co-cultures. The success was defined as the end-point dominance of a given strain in the co-culture (>90% of total CFU), with both strains ensured with an even start (1:1 mix; ca. 10^7 CFU/ml/strain). The theoretical winners, predicted based on tailocin-mediated interactions, were compared with those that were experimentally established. Co-cultures were conducted for 22 h in 0.1 TSB alone or supplemented with 0.88 mM H_2O_2 (tailocin-inductive conditions). SET 1: strains sensitive to tailocins of *Ds* RIF. SET 2: strains producing tailocins against *Ds* RIF. SET 3: strains sensitive to tailocins of *Ds* RIF and producing tailocins targeting this strain (bilateral killing). *ctrl* – control experiment; competition between *Ds* RIF and the *wt* variant of this strain.

Supplementary Tables

Table S1. List of bacterial strains used in this study.

Strain	Isolation source, country, year	Reference	GenBank accession no. for genome
Dickeya chrysanthemi A604-S21-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406775.2
Dickeya dianthicola A260-S21-A16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_020406895.2
Dickeya oryzae A003-S1-M15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_020406815.2
Dickeya oryzae A642-S2-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406685.2
Dickeya solani A623-S20-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406975.1
Dickeya zeae A586-S18-A17 recently reclassified to *D. parazeae*)	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020520245.1
Dickeya zeae A661-S21-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406575.2
Pectobacterium aquaticum A101-S19-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382625.2
Pectobacterium aquaticum A104-S21-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382595.2
Pectobacterium aquaticum A105-S21-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382585.2
Pectobacterium aquaticum A212-S19-A16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003382565.3
Pectobacterium atrosepticum A597-S4-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406655.2
Pectobacterium brasiliense A143-S20-M16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_013449685.1
Pectobacterium carotovorum A077-S18-O15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_020520265.1
Pectobacterium carotovorum A280-S2-N16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_020406635.1
Pectobacterium odoriferum A122-S21-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_013450045.1
Pectobacterium odoriferum A367-S5-N16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_022507245.1
Pectobacterium peruviense A350-S18-N16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003312355.2
Pectobacterium peruviense A97-S13-F16	Durance river. France. 2016	(Ben Moussa et al., 2022)	GCA_003312345.1
Pectobacterium polaris A641-S6-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406835.1
Pectobacterium quasiaquaticum A477-S1-J17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_014946775.2
Pectobacterium quasiaquaticum A411-S4-F17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_014946845.1
Pectobacterium versatile A546-S4-A17	Durance river. France. 2017	(Ben Moussa et al., 2022)	GCA_020406615.1
Pectobacterium versatile A69-S13-O15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_004296725.1
Pectobacterium versatile A73-S18-O15	Durance river. France. 2015	(Ben Moussa et al., 2022)	GCA_020865565.1
Dickeya dadantii 3937	*Saintpaulia* sp. France. 1972	(Kotoujansky et al., 1982)	GCA_000147055.1
Musicola paradisiaca NCPPB 2511	*Musa paradisiaca*. Colombia. 1973 *	(Hugouvieux-Cotte-Pattat et al., 2021)	GCA_000400505.1

763 **Table S2. List of sensitive strains used in a spot assay to detect tailocins from the**
764 **given producer strains.**

Tailocin producer	Sensitive strain
P. versatile A69-S13-O15	*P. aquaticum* A105-S21-F16
P. quasiaquaticum A411-S4-F17	*P. atrosepticum* A597-S4-A17
P. aquaticum A212-S19-A16	*P. aquaticum* A105-S21-F16
D. dadantii 3937	*M. paradisiaca* NCPPB 2511
D. chrysanthemi A604-S21-A17	*D. zeae* A586-S18-A17
D. zeae A586-S18-A17 (recently reclassified to *D. parazeae*)	*M. paradisiaca* NCPPB 2511

Supplementary Figures

Supplementary Fig. S1. Comparison of phenotypic and phylogenetic trees. (A)
dendrogram calculated based on the similarity of killing spectra of investigated strai; the tree
is drawn to scale, with branch lengths measured in the distance; (B) multilocus sequence
analysis of core nucleotide coding sequences using the BioNJ algorithm. Branch lengths and
bootstrap values are shown. *Pectobacterium* spp. strains are marked in green, and *Dickeya*
spp. strains are marked in red.

Supplementary Fig. S2. Severity of soft rot symptoms caused on potato tubers by six SRP strains tested for production of tailocins *in planta*. The results are expressed as a percentage of rotten tissue relative to the total mass of each potato tuber. In the box plot, whiskers reflect the maximum and minimum values; the box shows the interquartile range (Q1 to Q3), and the bars inside the boxes represent medians. Points indicate results for individual samples/tubers (n=6). The asterisks (*) denote statistically significant differences (p < 0.05) between a sample inoculated with the given strain and the negative control. Significance was calculated using the one-sample t-test for data sets with normal distribution and the Wilcoxon test for data not following a normal distribution. Negative control – tubers inoculated with sterile PBS buffer.

 Supplementary Fig. S3 Schematic representation of the steps included in the
 competition experiment. Created with BioRender.com.

Supplementary Fig. S4 Comparison of growth rates (A) and the survival (B) of the *D. solani* A623-S20-A17 wt and the rifampicin-resistant mutant of this strain in 0.1 TSB alone and supplemented with 0.003 % hydrogen peroxide (H_2O_2). The test was performed in 96-well plates incubated at 28˚C, with shaking, in a multiwell plate reader (Biotek). To assess growth rates (A), 150 µl aliquots of 0.1 TSB medium were inoculated with 3 µl of suspension containing o/n culture diluted to 0.5 McF (low inoculum). Each time point is an average of two replicates. For cell survival (B), the medium was inoculated with a high initial number of cells (ca. 10^7 CFU/ml) applied in the co-culture competition experiments (Fig. 5). The results are the average of two separate experiments, 2 technical replicates each. The obtained results indicate that the wt and RIF^R variants of *D. solani* A623-S20-A17 have similar growth characteristics in the applied conditions and that 0.003% H_2O_2 prevents the growth of the strains when low initial inoculum is used but does not negatively affect the survival of cells in high-density cultures.

Supplementary datasets

Dataset S1 Results of a megablast search for carotovoricin clusters within 116 closed genomes of *Pectobacterium* spp. The carotovoricin sequence from *P. versatile* A73-S18-O15 was used as a query (genome accession: NZ_CP086369.1).

Dataset S2 Results of a megablast search for dickeyocin clusters within 75 closed *Dickeya* and *Musicola* spp. genomes. Dickeyocin P2D1 from *D. dadantii* 3937 was used as a query (genome accession: NC_014500.1).

Dataset S3 Python script for analyses (ChatGPT-4o)

References

Backman, T., Burbano, H.A., and Karasov, T.L. (2024a). Tradeoffs and constraints on the evolution of tailocins. *Trends in Microbiology*.

Backman, T., Latorre, S.M., Symeonidi, E., Muszynski, A., Bleak, E., Eads, L., Martinez-Koury, P.I., Som, S., Hawks, A., Gloss, A.D., Belnap, D.M., Manuel, A.M., Deutschbauer, A.M., Bergelson, J., Azadi, P., Burbano, H.A., and Karasov, T.L. (2024b). A phage tail-like bacteriocin suppresses competitors in metapopulations of pathogenic bacteria. *Science* 384**,** eado0713.

Barny, M.-A., Thieffry, S., Gomes De Faria, C., Thebault, E., and Pédron, J. (2024). Bacterial pathogens dynamic during multi-species infections. *Peer Community Journal* 4.

Bauer, M.A., Kainz, K., Carmona-Gutierrez, D., and Madeo, F. (2018). Microbial wars: Competition in ecological niches and within the microbiome. *Microb Cell* 5**,** 215-219.

Bellieny-Rabelo, D., Tanui, C.K., Miguel, N., Kwenda, S., Shyntum, D.Y., and Moleleki, L.N. (2019). Transcriptome and Comparative Genomics Analyses Reveal New Functional Insights on Key Determinants of Pathogenesis and Interbacterial Competition in Pectobacterium and Dickeya spp. *Appl Environ Microbiol* 85**,** e02050-02018.

Ben Moussa, H., Bertrand, C., Rochelle-Newall, E., Fiorini, S., Pedron, J., and Barny, M.A. (2022). The Diversity and Abundance of Soft Rot Pectobacteriaceae Along the Durance River Stream in the Southeast of France Revealed by Multiple Seasonal Surveys. *Phytopathology* 112**,** 1676-1685.

Berg, G., Grosch, R., and Scherwinski, K. (2007). Risikofolgeabschätzung für den Einsatz mikrobieller Antagonisten: Gibt es Effekte auf Nichtzielorganismen? *Gesunde Pflanzen* 59**,** 107-117.

Bertozzi Silva, J., Storms, Z., and Sauvageau, D. (2016). Host receptors for bacteriophage adsorption. *FEMS Microbiol Lett* 363.

Booth, S.C., Smith, W.P.J., and Foster, K.R. (2023). The evolution of short- and long-range weapons for bacterial competition. *Nat Ecol Evol* 7**,** 2080-2091.

Borowicz, M., Krzyżanowska, D.M., Narajczyk, M., Sobolewska, M., Rajewska, M., Czaplewska, P., Węgrzyn, K., and Czajkowski, R. (2023). Soft rot pathogen Dickeya dadantii 3937 produces tailocins resembling the tails of Peduovirus P2. *Frontiers in Microbiology* 14.

Bradford, S.A., Morales, V.L., Zhang, W., Harvey, R.W., Packman, A.I., Mohanram, A., and Welty, C. (2013). Transport and Fate of Microbial Pathogens in Agricultural Settings. *Critical Reviews in Environmental Science and Technology* 43**,** 775-893.

Brettin, T., Davis, J.J., Disz, T., Edwards, R.A., Gerdes, S., Olsen, G.J., Olson, R., Overbeek, R., Parrello, B., Pusch, G.D., Shukla, M., Thomason, J.A., 3rd, Stevens, R., Vonstein, V., Wattam, A.R., and Xia, F. (2015). RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5**,** 8365.

Carim, S., Azadeh, A.L., Kazakov, A.E., Price, M.N., Walian, P.J., Lui, L.M., Nielsen, T.N., Chakraborty, R., Deutschbauer, A.M., Mutalik, V.K., and Arkin, A.P. (2021). Systematic discovery of pseudomonad genetic factors involved in sensitivity to tailocins. *ISME J* 15**,** 2289-2305.

Casjens, S. (2003). Prophages and bacterial genomics: what have we learned so far? *Mol Microbiol* 49**,** 277-300.

Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17**,** 540-552.

Charkowski, A.O. (2009). Decaying signals: will understanding bacterial-plant communications lead to control of soft rot? *Curr Opin Biotechnol* 20**,** 178-184.

Charkowski, A.O. (2018). The Changing Face of Bacterial Soft-Rot Diseases. *Annu Rev Phytopathol* 56**,** 269-288.

Czajkowski, R., De Boer, W.J., Velvis, H., and Van Der Wolf, J.M. (2010). Systemic colonization of potato plants by a soilborne, green fluorescent protein-tagged strain of *Dickeya* sp. biovar 3. *Phytopathology* 100**,** 134-142.

Dams, D., Brondsted, L., Drulis-Kawa, Z., and Briers, Y. (2019). Engineering of receptor-binding proteins in bacteriophages and phage tail-like bacteriocins. *Biochem Soc Trans* 47**,** 449-460.

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32**,** 1792-1797.

Erill, I., Campoy, S., and Barbe, J. (2007). Aeons of distress: an evolutionary perspective on the bacterial SOS response. *FEMS Microbiol Rev* 31**,** 637-656.

Expert, D., Patrit, O., Shevchik, V.E., Perino, C., Boucher, V., Creze, C., Wenes, E., and Fagard, M. (2018). Dickeya dadantii pectic enzymes necessary for virulence are also responsible for activation of the Arabidopsis thaliana innate immune system. *Mol Plant Pathol* 19**,** 313-327.

Fiołka, M.J., Czaplewska, P., Macur, K., Buchwald, T., Kutkowska, J., Paduch, R., Kaczyński, Z., Wydrych, J., and Urbanik-Sypniewska, T. (2019). Anti-Candida albicans effect of the protein-carbohydrate fraction obtained from the coelomic fluid of earthworm Dendrobaena veneta. *PLoS One* 14**,** e0212869.

Ghequire, M.G.K., and De Mot, R. (2015). The Tailocin Tale: Peeling off Phage Tails. *Trends Microbiol* 23**,** 587-590.

Gilchrist, C.L.M., and Chooi, Y.H. (2021). clinker & clustermap.js: automatic generation of gene cluster comparison figures. *Bioinformatics* 37**,** 2473-2475.

Glandorf, D.C.M., Brand, I., Bakker, P.a.H.M., and Schippers, B. (1992). Stability of Rifampicin Resistance as a Marker for Root Colonization Studies of Pseudomonas-Putida in the Field. *Plant and Soil* 147**,** 135-142.

Glasner, J.D., Marquez-Villavicencio, M., Kim, H.S., Jahn, C.E., Ma, B., Biehl, B.S., Rissman, A.I., Mole, B., Yi, X., Yang, C.H., Dangl, J.L., Grant, S.R., Perna, N.T., and Charkowski, A.O. (2008). Niche-specificity and the variable fraction of the Pectobacterium pan-genome. *Mol Plant Microbe Interact* 21**,** 1549-1560.

Goldman, A.R., Beer, L.A., Tang, H.Y., Hembach, P., Zayas-Bazan, D., and Speicher, D.W. (2019). Proteome Analysis Using Gel-LC-MS/MS. *Curr Protoc Protein Sci* 96**,** e93.

Golebiowski, A., Pomastowski, P., Rafinska, K., Zuvela, P., Wong, M.W., Pryshchepa, O., Madajski, P., and Buszewski, B. (2022). Functionalization of Alpha-Lactalbumin by Zinc Ions. *ACS Omega* 7**,** 38459-38474.

Gouy, M., Guindon, S., and Gascuel, O. (2010). SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* 27**,** 221-224.

Granato, E.T., Meiller-Legrand, T.A., and Foster, K.R. (2019). The Evolution and Ecology of Bacterial Warfare. *Curr Biol* 29**,** R521-R537.

Heiman, C.M., Vacheron, J., and Keel, C. (2023). Evolutionary and ecological role of extracellular contractile injection systems: from threat to weapon. *Front Microbiol* 14**,** 1264877.

Hibbing, M.E., Fuqua, C., Parsek, M.R., and Peterson, S.B. (2010). Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* 8**,** 15-25.

Hockett, K.L., and Baltrus, D.A. (2017). Use of the Soft-agar Overlay Technique to Screen for Bacterially Produced Inhibitory Compounds. *J Vis Exp*.

Hugouvieux-Cotte-Pattat, N. (2016). "Chapter Four - Metabolism and Virulence Strategies in Dickeya–Host Interactions," in *Progress in Molecular Biology and Translational Science,* eds. M. San Francisco & B. San Francisco. Academic Press), 93-129.

Hugouvieux-Cotte-Pattat, N., Des-Combes, C.J., Briolay, J., and Pritchard, L. (2021). Proposal for the creation of a new genus Musicola gen. nov., reclassification of Dickeya paradisiaca (Samson et al. 2005) as Musicola paradisiaca comb. nov. and description of a new species Musicola keenii sp. nov. *Int J Syst Evol Microbiol* 71.

Hugouvieux-Cotte-Pattat, N., Pédron, J., and Van Gijsegem, F. (2023). Insight into biodiversity of the recently rearranged genus Dickeya. *Frontiers in Plant Science* 14.

Jiang, X., Zghidi-Abouzid, O., Oger-Desfeux, C., Hommais, F., Greliche, N., Muskhelishvili, G., Nasser, W., and Reverchon, S. (2016). Global transcriptional response of Dickeya dadantii to environmental stimuli relevant to the plant infection. *Environ Microbiol* 18**,** 3651-3672.

Kotoujansky, A., Lemattre, M., and Boistard, P. (1982). Utilization of a thermosensitive episome bearing transposon TN10 to isolate Hfr donor strains of Erwinia carotovora subsp. chrysanthemi. *J Bacteriol* 150**,** 122-131.

Lamichhane, J.R., and Bartoli, C. (2015). Plant pathogenic bacteria in open irrigation systems: what risk for crop health? *Plant Pathology* 64**,** 757-766.

Ma, B., Hibbing, M.E., Kim, H.S., Reedy, R.M., Yedidia, I., Breuer, J., Breuer, J., Glasner, J.D., Perna, N.T., Kelman, A., and Charkowski, A.O. (2007). Host range and molecular phylogenies of the soft rot enterobacterial genera pectobacterium and dickeya. *Phytopathology* 97**,** 1150-1163.

Marquez-Villavicencio, M.D.P., Groves, R.L., and Charkowski, A.O. (2011). Soft Rot Disease Severity Is Affected by Potato Physiology and Pectobacterium taxa. *Plant Dis* 95**,** 232-241.

Miele, V., Penel, S., and Duret, L. (2011). Ultra-fast sequence clustering from similarity networks with SiLiX. *BMC Bioinformatics* 12**,** 116.

Nagakubo, T., Yamamoto, T., Asamizu, S., Toyofuku, M., Nomura, N., and Onaka, H. (2021). Phage tail-like nanostructures affect microbial interactions between Streptomyces and fungi. *Sci Rep* 11**,** 20116.

Nguyen, H.A., Kaneko, J., and Kamio, Y. (2002). Temperature-dependent production of carotovoricin Er and pectin lyase in phytopathogenic Erwinia carotovora subsp. carotovora Er. *Biosci Biotechnol Biochem* 66**,** 444-447.

Nielsen, F. (2016). "Hierarchical Clustering," in *Introduction to HPC with MPI for Data Science*. (Cham: Springer International Publishing), 195-211.

Pédron, J., and Van Gijsegem, F. (2019). Diversity in the Bacterial Genus Dickeya Grouping Plant Pathogens and Waterways Isolates. *OBM Genetics* 3**,** 1-1.

Perombelon, M.C.M. (Year). "Ecology of Erwinias causing stem and tuber diseases", in: *Bacterial disease of Potato: report of the planning conference on Bacterial diseases of potato*).

Perombelon, M.C.M., and Kelman, A. (1980). Ecology of the soft rot Erwinias. *Annual Review of Phytopathology* 18**,** 361-387.

Pérombelon, M.C.M., and Salmond, G.P.C. (1995). Bacterial soft rots. *Pathogenesis and Host Specificity in Plant Diseases* 1**,** 1-20.

Perombelon, M.C.M., and Van Der Wolf, J.M. (2002). Methods for the detection and quantification of *Erwinia carotovora* subsp. *atroseptica* (*Pectobacterium carotovorum* subsp. *atrosepticum*) on potatoes: a laboratory manual. *Scottish Crop Research Institute Annual Report* 10.

Portier, P., Pédron, J., Taghouti, G., Dutrieux, C., and Barny, M.-A. (2020). Updated Taxonomy of Pectobacterium Genus in the CIRM-CFBP Bacterial Collection: When Newly Described Species Reveal "Old" Endemic Population. *Microorganisms* 8**,** 1441.

Principe, A., Fernandez, M., Torasso, M., Godino, A., and Fischer, S. (2018). Effectiveness of tailocins produced by Pseudomonas fluorescens SF4c in controlling the bacterial-spot disease in tomatoes caused by Xanthomonas vesicatoria. *Microbiol Res* 212-213**,** 94-102.

Pritchard, L., Humphris, S., Saddler, G.S., Elphinstone, J.G., Pirhonen, M., and Toth, I.K. (2013). Draft genome sequences of 17 isolates of the plant pathogenic bacterium dickeya. *Genome Announc* 1.

Reverchon, S., Muskhelisvili, G., and Nasser, W. (2016). "Chapter Three - Virulence Program of a Bacterial Plant Pathogen: The Dickeya Model," in *Progress in Molecular Biology and Translational Science,* eds. M. San Francisco & B. San Francisco. Academic Press), 51-92.

Reverchon, S., and Nasser, W. (2013). Dickeya ecology, environment sensing and regulation of virulence programme. *Environ Microbiol Rep* 5**,** 622-636.

Ross-Gillespie, A., and Kummerli, R. (2014). 'Evolution-proofing' antibacterials. *Evol Med Public Health* 2014**,** 134-135.

Schmidt, R., and Sinz, A. (2017). Improved single-step enrichment methods of cross-linked products for protein structure analysis and protein interaction mapping. *Anal Bioanal Chem* 409**,** 2393-2400.

Scholl, D. (2017). Phage Tail-Like Bacteriocins. *Annu Rev Virol* 4**,** 453-467.

Shyntum, D.Y., Nkomo, N.P., Shingange, N.L., Gricia, A.R., Bellieny-Rabelo, D., and Moleleki, L.N. (2019). The Impact of Type VI Secretion System, Bacteriocins and Antibiotics on Bacterial Competition of Pectobacterium carotovorum subsp. brasiliense and the Regulation of Carbapenem Biosynthesis by Iron and the Ferric-Uptake Regulator. *Front Microbiol* 10**,** 2379.

Stice, S.P., Jan, H.H., Chen, H.C., Nwosu, L., Shin, G.Y., Weaver, S., Coutinho, T., Kvitko, B.H., and Baltrus, D.A. (2023). Pantailocins: phage-derived bacteriocins from Pantoea ananatis and Pantoea stewartii subsp. indologenes. *Appl Environ Microbiol* 89**,** e0092923.

Storz, G., Tartaglia, L.A., Farr, S.B., and Ames, B.N. (1990). Bacterial defenses against oxidative stress. *Trends Genet* 6**,** 363-368.

Torres, M.A., Jones, J.D., and Dangl, J.L. (2006). Reactive oxygen species signaling in response to pathogens. *Plant Physiol* 141**,** 373-378.

Toth, I.K., Barny, M.-A., Brurberg, M.B., Condemine, G., Czajkowski, R., Elphinstone, J.G., Helias, V., Johnson, S.B., Moleleki, L.N., and Pirhonen, M. (2021a). Pectobacterium and Dickeya: environment to disease development. *Plant diseases caused by dickeya and pectobacterium species***,** 39-84.

Toth, I.K., Barny, M.-A., Brurberg, M.B., Condemine, G., Czajkowski, R., Elphinstone, J.G., Helias, V., Johnson, S.B., Moleleki, L.N., Pirhonen, M., Rossmann, S., Tsror, L., Van Der Waals, J.E., Van Der Wolf, J.M., Van Gijsegem, F., and Yedidia, I. (2021b). "Pectobacterium and Dickeya: Environment to Disease Development," in *Plant Diseases Caused by Dickeya and Pectobacterium Species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 39-84.

Van Der Wolf, J.M., Acuña, I., De Boer, S.H., Brurberg, M.B., Cahill, G., Charkowski, A.O., Coutinho, T., Davey, T., Dees, M.W., Degefu, Y., Dupuis, B., Elphinstone, J.G., Fan, J., Fazelisanagri, E., Fleming, T., Gerayeli, N., Gorshkov, V., Helias, V., Le Hingrat, Y., Johnson, S.B., Keiser, A., Kellenberger, I., Li, X., Lojkowska, E., Martin, R., Perminow, J.I., Petrova, O., Motyka-Pomagruk, A., Rossmann, S., Schaerer, S., Sledz, W., Toth, I.K., Tsror, L., Van Der Waals, J.E., De Werra, P., and Yedidia, I. (2021). "Diseases Caused by *Pectobacterium* and *Dickeya* apecies around the world," in *Plant diseases caused by Dickeya and Pectobacterium species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 215-261.

Van Gijsegem, F., Hugouvieux-Cotte-Pattat, N., Kraepiel, Y., Lojkowska, E., Moleleki, L.N., Gorshkov, V., and Yedidia, I. (2021a). "Molecular Interactions of Pectobacterium and Dickeya with Plants," in *Plant Diseases Caused by Dickeya and Pectobacterium Species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 85-147.

Van Gijsegem, F., Toth, I.K., and Van Der Wolf, J.M. (2021b). "Soft Rot Pectobacteriaceae: A Brief Overview," in *Plant Diseases Caused by Dickeya and Pectobacterium Species,* eds. F. Van Gijsegem, J.M. Van Der Wolf & I.K. Toth. (Cham: Springer International Publishing), 1-11.

Varani, A.M., Monteiro-Vitorello, C.B., Nakaya, H.I., and Van Sluys, M.A. (2013). The role of prophage in plant-pathogenic bacteria. *Annu Rev Phytopathol* 51**,** 429-451.

Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., Van Der Walt, S.J., Brett, M., Wilson, J., Millman, K.J., Mayorov, N., Nelson, A.R.J., Jones, E., Kern, R., Larson, E., Carey, C.J., Polat, I., Feng, Y., Moore, E.W., Vanderplas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E.A., Harris, C.R., Archibald, A.M., Ribeiro, A.H., Pedregosa, F., Van Mulbregt, P., and Scipy, C. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* 17**,** 261-272.

Wagner, A. (2022). Competition for nutrients increases invasion resistance during assembly of microbial communities. *Mol Ecol* 31**,** 4188-4203.

Yamada, K., Hirota, M., Niimi, Y., Nguyen, H.A., Takahara, Y., Kamio, Y., and Kaneko, J. (2006). Nucleotide sequences and organization of the genes for carotovoricin (Ctv) from Erwinia carotovora indicate that Ctv evolved from the same ancestor as Salmonella typhi prophage. *Biosci Biotechnol Biochem* 70**,** 2236-2247.

Yao, G.W., Duarte, I., Le, T.T., Carmody, L., Lipuma, J.J., Young, R., and Gonzalez, C.F. (2017). A Broad-Host-Range Tailocin from Burkholderia cenocepacia. *Appl Environ Microbiol* 83.