Tracing Cellular Senescence in Bone: Time-Dependent Changes in Osteocyte Cytoskeleton Mechanics and Morphology

Maryam Tilton [1*], Junhan Liao [1], Chanul Kim [2], Hossein Shaygani [1], Maria Astudillo Potes [3,4],

Domenic Cordova [1], James L. Kirkland [5], Kyle M. Miller [6]

[1] Walker Department of Mechanical Engineering, The University of Texas at Austin, Austin, TX, USA

[2] Department of Biomedical Engineering, The University of Texas at Austin, Austin, TX, USA

[3] Department of Physiology and Biomedical Engineering, Mayo Clinic, Rochester, MN, USA

[4] Department of Orthopedic Surgery, Mayo Clinic, Rochester, MN, USA

[5] Center for Advanced Gerotherapeutics, Division of Endocrinology, Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[6] Department of Molecular Biosciences, The University of Texas at Austin, Austin, TX, USA

*Corresponding author: <u>maryam.tilton@austin.utexas.edu</u>

Abstract

Aging-related bone loss significantly impacts the growing elderly population globally, leading to

debilitating conditions such as osteoporosis. Senescent osteocytes play a crucial role in the aging

process of bone. This longitudinal study examines the impact of continuous local and paracrine

exposure to senescence-associated secretory phenotype (SASP) factors on senescence-associated

biomolecular and biomolecular markers in osteocytes. We found significant cytoskeletal stiffening

in irradiated osteocytes, accompanied by expansion of F-actin areas and a decline in dendritic

integrity. These changes, correlating with alterations in pro-inflammatory cytokine levels and

osteocyte-specific gene expression, support the reliability of biophysical markers for identifying

senescent osteocytes. Notably, local accumulation of SASP factors had a more pronounced

impact on osteocyte properties than paracrine effects, suggesting that the interplay between local

and paracrine exposure could substantially influence cellular aging. This study underscores the

importance of osteocyte mechanical and morphological properties as biophysical markers of

senescence, highlighting their time-dependence and differential effects of local and paracrine

SASP exposure. Collectively, our investigation into biophysical senescence markers offer unique

and reliable functional hallmarks for non-invasive identification of senescent osteocytes,

providing insights that could inform therapeutic strategies to mitigate aging-related bone loss.

1. Introduction

Age-related bone loss impacts over 43 million Americans aged 50 and older, with a projected 47.5% increase in prevalence among those aged 65 and over. [1] This bone loss contributes to osteoporosis and fragility fractures, which remain leading causes of disability, morbidity, and mortality among the elderly. With the global demographic shift towards an aging population, where over 11,000 Americans turn 50 daily and projections indicate that by 2030, all "baby boomers" will be at least 65 years old, [2–4] the urgency to understand the degenerative mechanisms of bone loss becomes a public health priority.

Bone loss is associated with the increased accumulation of senescent cells (SnCs), including senescent osteocytes, and their senescent-associated secretory phenotypes (SASP).[5–8] Indeed, most cells undergoing senescence develop SASPs that include factors with proinflammatory, proapoptotic, and pro-fibrotic effects on neighboring cells.[9] Accumulation of SnCs can have local and systemic detrimental effects due in part to SASP factors.[9,10] To a certain extent, SnCs can be cleared naturally by innate and adaptive immune cells. However, SnCs expressing pro-inflammatory and proapoptotic SASP factors can induce paracrine and endocrine signals, spreading senescence at a rate exceeding immune clearance of existing or newly developed SnCs. This results in accumulation of SnCs at different pathological sites with age. [11] Specifically, SASP factors can induce senescence in normal, non-SnCs both locally and systemically. [12]

SASP factors include pro-inflammatory cytokines, chemokines, growth factors, proteases, and bio-active lipids. [13] In bone, the accumulation of SnCs disrupts the balance between bone formation and resorption, driving the progression of osteoporosis and increasing the risk of fractures. [14,15] The SASP factors secreted by these cells alter the bone microenvironment, enhancing osteoclastic activity and reducing bone density.

The cytoskeleton of cells consists of actin microfilaments, intermediate filaments, and microtubules, providing mechanical support, maintaining cell shape, and playing critical roles in intracellular transport, cell division, and molecular signaling.[16,17] Under mechanical stimulation, the cytoskeleton remodels, including the reorganization of stress fibers, which supports cellular functions by adapting to changes in their mechanical environment.[18–20] Osteocytes, the most abundant skeletal cells, are master regulators of bone homeostasis, being endocrine cells that regulate phosphate metabolism in organs such as the kidney and parathyroid, and most importantly, acting as strain gauges, regulating bone mechanosensation and

mechanotransduction.[21–24] Specifically, the osteocyte cytoskeleton is crucial to their function,

including cell-extracellular matrix (ECM) interactions via focal adhesions [25,26] Previous studies

have demonstrated that SnCs such as senescent mesenchymal stem cells (MSCs) and epithelial

cells, undergo alterations in cytoskeletal architecture, potentially impairing their

mechanotransduction capabilities. These cytoskeletal changes in aging osteocytes offer

opportunities to investigate unique senescence-related biophysical markers, reflecting alterations

in their mechanobiological functions.

Cellular senescence is characterized by essentially irreversible proliferative arrest, altered

chromatin organization, apoptosis resistance, tumor suppressor activation, oncogenic mutation,

and increased protein synthesis.[27–31] Cellular senescence, a state of irreversible cell-cycle arrest,

can be induced by various intrinsic (e.g. DNA damage, telomere shortening or dysfunction,

oncogene activation, loss of tumor suppressor functions) and extrinsic (e.g. ultraviolet radiation

and chemotherapeutic agents) pro-senescence stressors.[32,33] Among these, nuclear DNA damage

is a principal trigger of senescence, as it activates the p53-p21 pathway to halt cell proliferation,

while epigenetic changes drive senescence via the p16-RB pathway. While senescence is

primarily associated with the activation of tumor suppressors, certain oncogenic pathways have

been shown to trigger senescence under specific conditions, further reflecting the complexity of

the process.[34–36] These processes are highly context-dependent, with variability in the stimuli

leading to senescence across different cell types, tissues, and even within heterogeneous cell

populations.[37,38] This variability complicates the identification of a universal marker for SnCs.

For instance, markers such as Cdkn2a/p16^{Ink4a}, which reflects tumor suppressor pathway

activation, and Cdkn1a/p21^{Cip1}, associated with cell cycle arrest, show variable expression levels

in different cellular contexts.[39–42] Senescence associated β-Galactosidase (SA-β-Gal) activity,

while commonly used,[43–45] is neither exclusive to SnCs nor consistent across cell populations

such as MSCs and epithelial cells. [38,46–48] Similarly, chromatin rearrangements and the

formation of senescence-associated heterochromatin foci (SAHF) are observable in some but not

all SnC types.[38,49] Recent reviews emphasize the necessity of combining multiple markers, such

as transcriptional signatures (e.g., SASP-related gene panels) with biophysical and functional

assays, to increase diagnostic accuracy. [37,38,50,51] This multi-marker approach is crucial given the

sexual dimorphism in marker expression reported in human and animal models.

Commonly used pro-senescence stressors to study senescence include ionizing radiation,
oxidative stress induced by reactive oxygen species (ROS), and chemotherapeutic agents such as
doxorubicin.[32,33] These stressors are widely known to induce SASP-related phenotypes,
characterized by the release of cytokines, chemokines, growth factors, and proteases that
influence local and systemic environments.[51–54] Among these stressors, ionizing radiation is
particularly well-suited for in vitro senescence models due to its ability to cause extensive DNA
damage, triggering the DNA damage response (DDR) and activating senescence pathways with
high reproducibility.[7,38,55–57] Previous studies, including our own, have demonstrated that
irradiation effectively models senescence in osteocytes, inducing markers such as *Cdkn2a*
(*p16Ink4a*) and *Cdkn1a* (*p21Cip1*), increasing SA-β-Gal activity, and impairing
mechanobiological properties.[56,57] These characteristics, combined with precise dosing control,
highlight the suitability of irradiation for investigating senescence-associated biophysical
changes in osteocytes.
Osteocyte mechanobiology is fundamentally linked to cytoskeletal mechanics and morphology,
which are essential for maintaining bone health and function. Alterations in cell mechanics and
morphology are not exclusively indicative of senescence but can signal various cellular states or
responses. In this study, we specifically probe these changes in the context of exposure to pro-
senescence stressors (*e.g*., irradiation) in bone cells. By examining these alterations when cells
face such stressors, we aim to establish a novel set of biophysical fingerprints that uniquely
characterize cellular senescence within the bone microenvironment. We specifically explore the
effects of local SASP accumulation over time on these markers, as well as the paracrine
influences of SASP factors through conditioned medium (CM) experiments. By supplying CM to
non-irradiated osteocyte cultures, both immediately and after generating a mature healthy culture,
we aim to mimic the exposure of non-senescent cells to a SASP-rich environment and investigate
their responses. Our findings highlight alterations in cytoskeletal architecture and mechanics
correlated to the local and paracrine effects of pro-senescence stressors, which potentially could
reflect key osteocyte mechanobiological functions. This approach not only enhances our
understanding of the direct and indirect effects of SASP factors but also provides insight into the
aging process of bone through the lens of osteocyte biophysical changes.

2. Methods

2.1 Cell culture and *in vitro* senescence induction

To focus on senescence-associated changes in osteocyte biophysical properties, we employed
immortalized osteocytes, specifically MLO-Y4 cells. The mouse-derived MLO-Y4 cells were
sourced from Prof. Linda Bonewald's lab at Indiana University. This approach mitigated the
variabilities introduced by heterogeneous primary bone cell cultures. Osteocytes were cultured
following established protocols. [58] Briefly, cells were passaged twice before being plated at a
density of 2×10^4 cells/35-mm^2 well. The bottom of the well plates was coated with rat tail
collagen type I (0.5 mg/mL, Sigma-Aldrich) and filter-sterilized 0.02 M acetic acid (diluted from
glacial, ReagentPlus®, ≥99%, Sigma-Aldrich). The growth media consisted of α-Minimum
Essential Medium Eagle (αMEM, Sigma-Aldrich), supplemented with 5% fetal bovine serum
(FBS, Gibco), 5% calf serum (CS, Sigma-Aldrich), [59,60] and 1% penicillin-streptomycin (P/S,
10,000 U/mL, Gibco). Half of the medium volume was replenished every four days. The cells
were incubated at 37°C in a 5% CO_2 humidified chamber at confluency of less than 70% to
minimize disturbance of cell morphology.

Senescence was induced *in vitro* by a single dose of 10 Gy irradiation (CellRad, Precision X-Ray,
Inc.). Irradiation was chosen as a method for inducing senescence due to its ability to rapidly
activate the DDR, leading to permanent cell cycle arrest. This approach aligns with widely
accepted protocols in cellular senescence research.[7,38,55–57] The reproducibility of this method,
combined with its precise dosing control, ensures consistent senescence induction across
experimental groups.[33,38,50–52] Furthermore, irradiation has been widely used to model osteocyte
senescence in vitro, including in MLO-Y4 cells[61] and primary osteocytes[56,57], where it induces
markers of senescence such as *Cdkn2a* (*p16*) and *Cdkn1a* (*p21*), increases SA-β-Gal activity, and
impairs cellular function. These features make irradiation an effective and clinically relevant
model for investigating the impact of senescence on osteocyte mechanobiology. Here, post-
irradiation, cells were assessed for biophysical and biomolecular properties on days 7, 14, and 20
(irradiated study groups: D7, D14, and D20). Non-irradiated cells continuously cultured in fresh
media served as the Control group (Ctrl).

2.2 CM-supplemented groups

To investigate the systemic effects of SASP factors on healthy osteocytes, CM were collected from irradiated osteocyte cultures after 20 days. The CM was centrifuged at 1,500 × g for 5 minutes to remove cellular debris, yielding a consistent and uncontaminated SASP-enriched supernatant. Two groups were established: the immediate CM group (CM-i), where healthy osteocytes were exposed to CM from the start of culture (Day 0), and the delayed CM group (CM-d), where healthy osteocytes were pre-incubated in fresh media for 48 hours before CM supplementation to assess the impact of prior SASP-free growth. Both groups were cultured for 72 hours before biophysical and biomolecular assessments. Cells for imaging and nanoindentation were plated at a density of 1×10^5 cells per 35-mm² well, while those for RT-qPCR were plated at 6×10^5 cells per 35-mm² well, ensuring sufficient material for accurate analysis.

2.3 Single-cell nanoindentation

Single-cell micromechanical testing was performed on 50 random osteocytes across two wells using an optical fiber-based interferometry nanoindenter (Pavone, Optics11 Life). Osteocytes were indented with a probe having a 3 µm radius, which is significantly smaller than their surface areas, and with a stiffness of 0.019 N/m. The nanoindentation was performed at a controlled speed of 30 µm/s, reaching a peak load of 0.01 µN, after which the probe retracted at the same speed. The mechanical properties of osteocyte cytoskeleton, characterized by Young's modulus, were analyzed using Hertzian contact mechanics [62,63]. Indentation curve fitting was conducted using DataViewer V2.5.2 (Optic11 Life) over the linear region on the loading curve, accepting $R^2 > 0.90$ (**Figure S2**).

2.4 Gene expression

Gene expression analysis was conducted according to the manufacturer's protocols. RNA was initially extracted from the cells using TRIzol (Invitrogen) followed by purification with the GeneJET RNA Purification Kit (Thermo Fisher Scientific). The concentration of RNA was measured using a NanoDrop Microvolume Spectrophotometer (Thermo Fisher Scientific). Subsequently, cDNA was synthesized from the purified RNA after genomic DNA (gDNA) digestion, using SuperScript™ IV VILO™ Master Mix with ezDNase (Invitrogen). Real-time

181 quantitative Polymerase Chain Reaction (RT-qPCR) was performed using PowerTrack™ SYBR

182 Green Master Mix (Applied Biosystems) with the specific forward and reverse primers listed in

183 **Supplementary Table S1**. The RT-qPCR reactions were conducted on a Quantstudio 6 Pro

184 system (Applied Biosystems). Gene expression results were normalized to Gapdh as the

185 housekeeping gene using the Comparative Ct ($2^{-\Delta\Delta Ct}$) method.

2.5 SA-β-Gal staining

Following the manufacturer's protocol, cells were fixed and stained with SA-β-Gal (Cell

Signaling Technology). After staining, the cells were incubated at 37°C overnight. Cells

displaying a blue color were considered to be SA-β-Gal positive. Images were captured using a

brightfield microscope equipped with an Olympus XC30 camera, taking five random fields of

view per well. The SA-β-Gal activity was quantified by counting the number of positive cells

and normalizing this count to the total number of cells observed in each micrograph.

2.6 Immunofluorescent (IF) staining

Following the manufacturer's protocol, cells were fixed in 4% paraformaldehyde (PFA) and

stained with Cellmask Orange Actin Tracking Stain (Invitrogen) to visualize filamentous actin

(F-actin). The nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI, Invitrogen).

Fluorescent images were captured at seven fields of view using an inverted fluorescent

microscope (Optics11 Life) equipped with a Nikon 20x objective. Imaging utilized the TRITC

and DAPI channels at 545/570 nm and 350/465 nm (excitation/emission wavelengths),

respectively. ImageJ 2.14.0 was used for postprocessing. The morphologies of nuclei, F-actin,

and dendrites were analyzed and quantified using Cellprofiler 4.2.6 and NeuriteQuant, [64]

respectively. The F-actin area was measured through segmentation and quantification of pixels in

each fluorescent image (**Figure S4**). While NeuriteQuant was originally developed for neuron

analysis, recent studies have demonstrated its application in osteoblasts for the quantification of

dendrite lengths,[65] validating its broader functionality for non-neuronal cells. In this study, the

software was adapted to ensure reliable quantification of osteocyte dendritic processes. Specific

postprocessing parameters were optimized to account for the unique morphology of osteocytes,

including adjustments to thresholding and detection settings, to achieve accurate dendrite

segmentation and length measurements. Lastly, the multinucleation phenomena within the cell

cultures were quantified by counting the number of multinucleated cells and normalizing over
the total number of cells in each micrograph.

2.7 Statistical analysis

All the reported data underwent the normality and statistical analysis using GraphPad Prism 10.
Both one-way and two-way analysis of variance (ANOVA) with Tukey's multiple comparison
tests were employed, each at a 95% confidence interval. One-way ANOVA with Tukey's
multiple comparison tests was utilized for the statistical analysis of SA-β-Gal activity, Young's
moduli, and morphological quantifications. For the RT-qPCR data analyzing gene expression,
either One-way or Two-way ANOVA with Tukey's multiple comparison tests was applied,
depending on the experimental design. Statistical significance was considered to be $p < 0.05$.

3. Results and Discussion

3.1.1 Irradiated osteocytes exhibited time-dependent changes in biomolecular markers

After irradiation, SA-β-Gal positivity in osteocytes increased over time, which was visualized by
increase in cytoplasmic blue staining. The SA-β-Gal-active osteocytes reached close to 100% of
the population on Day 20 post-irradiation ($p < 0.0001$, **Figure 1A**), confirming successful
induction of senescence phenotype by low dose irradiation in these cells.

Gene expression of senescence markers, SASP factors, and osteocyte markers were measured for
all samples. $p16^{Ink4a}$ is a senescence marker that is responsible for regulating the G1/S phase of
the cell cycle, [66] while $p21^{Cip1}$ is another senescence marker involved in inhibiting both the
G1/S and G2/M phase. [67,68] In the irradiated groups, *p16* and *p21* exhibited gradual increases
over time, with *p21* showing an over 2-fold increase compared to the control group by Day 7
(**Figure 1B**). However, these changes were not statistically significant. In contrast, the significant
increase in SA-β-Gal positivity across all time points, along with the upregulation of pro-
inflammatory SASP markers provides robust evidence for senescence induction in irradiated
osteocytes. These findings indicate disruptions of the cell cycle in irradiated osteocytes, which as
typical for senescent cells.

The inflammatory response in osteocytes can be reflected by expression levels of interleukin-6
(*Il-6*) and tumor necrosis factor (*Tnf*). The cytokine *Il-6* is a SASP factor that not only stimulates

osteoclast differentiation and regulates immune responses but also modulates inflammation.
Expression of *Il-6* increased significantly over time in the irradiated groups compared to controls
(p < 0.01, **Figure 1C**), suggesting an escalated inflammatory response in irradiated osteocytes.
This increase in Il-6 has been previously associated with compromised mechanotransduction in
osteocytes [69], linking its accumulation with progressively impaired cellular function [70].
Similarly, expression of Tnf on Day 7 significantly increased, which could potentially signify an
increase in receptor activator of nuclear factor κB ligand (Rankl) production and suppression of
osteoprotegerin (Opg) [71,72], potentially contributing to bone loss [73]. The elevated expression of
both SASP factors in the irradiated groups indicates an increased inflammatory response through
autocrine or paracrine effects. The particularly notable time-dependent accumulation of Il-6
underscores its critical role in the senescence-related progression of bone aging.

The ability for osteocytes to regulate bone remodeling was assessed by measuring the levels of
matrix metalloproteinase (Mmp12) and bone gamma-carboxyglutamate protein (Bglap). Mmp12,
a SASP factor known for its role in ECM degradation, is critical in bone remodeling processes.
Its increased activity is associated with accelerated bone resorption and potentially dysfunctional
bone remodeling [74]. Expression of Mmp12 was elevated in the Irradiated groups over time, with
more than a 20-fold increase by Day 20 compared to the Control (p < 0.0001, **Figure 1C**). This
result suggests intensified ECM degradation and possible remodeling dysfunction in these
groups.

Osteocalcin (*Bglap*) plays a pivotal role in bone formation and remodeling. [75] During bone
formation, *Bglap* enhances matrix mineralization by binding calcium and hydroxyapatite,
contributing to the structural integrity of bone.[76,77] In remodeling, *Bglap* deposited in the bone
matrix indirectly promotes resorption by stimulating osteoclast activity, thereby coordinating the
balance between bone formation and resorption.[76,77] Its dual role in these processes makes it an
ideal marker for investigating the metabolic dynamics of bone. In this study, irradiated
osteocytes showed significantly increased *Bglap* expression (p < 0.0001 for Day 7 and 14, p <
0.001 for Day 20, Figure 1D), suggesting enhanced bone resorption activity and its potential
contribution to overall bone loss during senescence. This finding indicates the potential role of
Bglap as a key metabolic marker in bone remodeling under senescence-associated conditions.

Gap junction protein alpha 1 (*Gja1*) expression can be linked to osteocyte dendrite function in
terms of intercellular communication.[78] Disruption in Gja1 production has been previously
associated with age-related osteoclastogenesis and bone loss.[79] In **Figure 1D**, there were no
statistically significant changes in Gja1 expression among the groups (Ctrl, D7, D14, and D20).
While slight variations were observed, such as a marginal increase in Gja1 expression on Day 20
compared to Day 7, the lack of significant changes in Gja1 expression suggests that intercellular
communication via gap junctions may not be severely impaired under the tested conditions.

The gene expression results evaluated by RT-qPCR suggested that over time, the irradiated group
exhibited an increase in senescence markers and higher levels of pro-inflammatory cytokines.
The increased expression of Bglap further underscored declines in osteocyte mechanobiological
function and potential dysregulation of bone resorption.

Importantly, irradiation-induced senescence primarily represents stress-induced premature
senescence (SIPS) rather than replicative senescence (RS). [37] However, this method offers
distinct advantages, such as rapid senescence induction and precise experimental control.[37]
Unlike RS, which arises gradually through telomere attrition, SIPS bypasses telomere erosion
and directly activates DDR pathways. [37] It should be noted that cellular responses to ionizing
radiation may vary based on cell cycle phase, with proliferating cells being more susceptible to
damage than non-dividing cells.[33,37,80] These factors necessitate careful interpretation of
findings.

Figure 1: Time-dependent changes in biomolecular markers in irradiated osteocytes driven by local

accumulation of SASP factors. A) Representative images of SA-β-Gal staining at various time points for

irradiated cells (left) with quantification of SA-β-Gal active osteocytes (right). Five images per group

were analyzed (n = 5). **B)** RT-qPCR results for senescence, **C)** SASP factors, and **D)** osteocyte markers at

all time points. Results are normalized to *Gapdh* housekeeping gene expression and compared to the

Control group. Sample size for each group is n = 3, presented as mean ± SD. *: $p < 0.05$, **: $p < 0.01$,

***: $p < 0.001$, ****: $p < 0.0001$.

3.1.2 Short-term SASP factor exposure induced biomolecular marker changes in healthy

osteocytes

Gene expression levels from Day 7 served as the control in order to compare the local and

paracrine effects of SASP factors (**Figure 2A**). Expression of *p16* in the CM-i group was similar

to that observed on Day 7, while expression in the CM-d group was lower (0.525-fold).

Conversely, *p21* expression increased in the CM-i group (1.745-fold), with levels in the CM-d

group matching those on Day 7 (see **Figure 2B**). The stable *p16* expression in the CM-i group,

compared to the irradiated groups, suggests a disruption in the S phase of the cell cycle,

indicating that DNA synthesis in osteocytes exposed to CM-i was significantly impacted by

SASP factor exposure. On the other hand, the increase in *p21* levels in the CM-i group suggests

potential dysfunction in mitosis, reflecting disruption in the M phase of the cell cycle. Despite

the changes in p16 and p21 expressions in CM-d group, this group exhibited minimal SA-β-gal
positivity (**Figure S1**).

In the CM-d group (**Figure 2A**), *Il6* and *Mmp12* levels were comparable to those observed on
Day 7, whereas the CM-i group exhibited a 1.509-fold increase in *Il6*, surpassing the levels noted
on Day 14. Conversely, *Tnf* levels in both the CM-i and CM-d groups remained low, aligning
more closely with the values observed on Days 14 and 20 rather than Day 7. *Mmp12* levels in
both CM-supplemented groups showed a notable increase compared to Day 7 (6.198-fold and
4.047-fold, respectively; see **Figure 2C**). These results suggest that *Il6* and *Mmp12* played a
more prominent role in autocrine/paracrine signaling than *Tnf* in the non-irradiated cells within
the CM-supplemented groups. Additionally, the immediate incubation of healthy osteocytes in
SASP-supplemented CM in the CM-i group may have stimulated a more effective uptake and
signaling response to *Il6*, further amplifying its expression. Overall, these findings confirm that
healthy osteocytes in both CM-supplemented groups effectively absorbed SASP factors from
their conditioned media. Gene expression analysis comparing local versus systemic SASP
exposure, normalized to Day 14, further underscores these observations (**Figure S5**). The
senescence marker *p21* was significantly increased in both CM-i and CM-d groups compared to
Day 14, while *p16* expression remained unchanged and non-significant (**Figure S5-A**). SASP-
related cytokines, such as *Il6*, *Mmp12*, and *Tnf*, were comparable to their expression levels on
Day 14 (**Figure S5-B and S5-C**). However, *Il6* and *Mmp12* levels in the CM-d group were
significantly lower compared to Day 20.

Bglap levels in both CM-supplemented groups were significantly lower than those observed at
all local time points, including Day 7, Day 14, and Day 20 (p < 0.05, **Figure 2D**, and **Figure S5-**
D). This finding suggests that the paracrine effects of SASP factors were less effective in
promoting osteocalcin production compared to their local effects. Similarly, Gja1 levels in both
CM-supplemented groups showed an increase from Day 7, with levels in the CM-d group
remaining similar to Day 14, whereas CM-i exhibited a significant increase compared to Day 7
(p < 0.0001, **Figure 2D**, and **Figure S5-D**). This increase might be attributed to the abrupt
exposure of osteocytes in the CM-i group to SASP factors in the CM triggering a more intensive
survival response. Overall, the gene expression results for the CM-supplemented osteocytes
confirm effective uptake of SASP factors and suggest potential stimulation of autocrine/paracrine
SASP signaling pathways. The distinctly different profiles of senescence and osteocyte markers
reflect potential variations in the mechanisms through which paracrine SASP factor exposure
influences non-irradiated osteocytes.

Figure 2: Systemic supplementation with SASP factor-enriched conditioned media induces
senescence phenotypes in non-irradiated osteocytes. **A)** Schematic of the CM-i (immediate

supplementation) and CM-d (delayed supplementation) culture setups. **B)** RT-qPCR results for senescence,

C) SASP factors, and **D)** osteocyte markers across all conditions, normalized to *Gapdh* housekeeping

gene expression and compared to the Day 7 post-irradiation group. Sample size for each group is n = 3,

presented as mean ± SD. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

3.2 Osteocyte morphological changes as biophysical markers for senescence

Various morphological parameters were quantified in our longitudinal study to examine

senescence-associated changes (**Figure 3A**). Nuclear blebbing, observed in the irradiated groups

(**Figure 3B**), indicated nuclear membrane instability. Additionally, nuclear expansion was noted

over time, being particularly significant on Days 14 ($p < 0.0001$) and 20 ($p < 0.001$, **Figure 3C**).

In contrast, osteocytes in both CM-supplemented groups exhibited no significant changes from

the control group, suggesting that short-term exposure to SASP factors may not induce profound

alterations in nuclear morphology. Such stability might be attributed to mechanisms such as

chromosome unwinding within the nuclei, which potentially help mitigate aging-related DNA

damage. The nuclear abnormalities observed in the irradiated groups signify compromised

nuclear function in osteocytes, which could lead to phenotypic variations during protein

translation as previously observed in other cell types [81,82]. These findings should be further

investigated in future studies.

A significantly elevated population of multinucleated osteocytes was observed in the Irradiated

groups, reaching as high as 11% on Day 20 ($p < 0.0001$, **Figure 3A, 3D**). In contrast, both CM-

supplemented groups maintained similarly low levels of multinucleated cells to the Control

group. Since senescent osteocytes often exhibit disrupted cell cycles, the observed

multinucleation may result from incomplete M phases and/or DNA hyper-replication, similar to

findings reported in oncogene-induced senescence studies of primary human fibroblasts [83]. This

hypothesis is supported by the increased expression of p21 over time in the Irradiated groups,

underscoring the progression of cell cycle disruption. The prevalence of multinucleation in

irradiated osteocytes further emphasizes the impact of their local SASP factor accumulation,

which appears to be more consequential than short-term SASP exposure from their

microenvironment.

Figure 3: Alterations in F-actin organization and nuclear morphology are associated with local and systemic exposure to SASP factors. A) Representative IF micrographs showing F-actin and nuclei staining across all experimental conditions, with yellow arrows highlighting multinucleated cells. Scale bar: 50 µm. **B)** Close-up view of nuclear blebbing in irradiated osteocytes, indicated by a yellow arrow. **C)** Average nuclear areas quantified using CellProfiler from IF micrographs across all conditions. **D)** Percentages of multinucleated osteocytes in each experimental group. Quantitative analysis of osteocyte morphology was performed by examining 14 IF micrographs per group across two cell culture wells (n = 14 for all groups, presented as mean ± SD). Significance levels are denoted as *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

F-actin networks in osteocytes from the Irradiated groups exhibited extensive expansion compared to the Control group, characterized by reduced dendricity and non-uniform organization of F-actin structures (**Figures 3A**, **4A**). The actin disassembly (**Figures 3A**, **4A**) and potentially reduced polymerization dynamics are consistent with oxidative stress-induced mechanisms reported in previous studies.[84,85] Specifically, previous studies demonstrated that oxidative stress can alter the actin cytoskeleton through Cys374 oxidation, leading to disruptions in filament stability and polymerization.[84,85] These mechanisms align with our observations and suggest a role for oxidative stress in the observed cytoskeletal changes. However, we

acknowledge that direct oxidative stress measurements were not performed in this study. Future
experiments will incorporate assays, such as superoxide anion fluorescent probe staining, to
directly evaluate reactive oxygen species (ROS) levels and their effects on the osteocyte
cytoskeleton.

A progressive time-dependent increase in the average F-actin area was observed in the Irradiated
groups ($p < 0.0001$), significantly exceeding that of the Control and both CM-supplemented
groups (**Figure 4B**). This increase could be attributed to abnormal filopodia and lamellipodia
formation in senescent cells, which likely contributed to osteocyte flattening and area expansion.
The disorganized and expanded F-actin networks may have resulted in spatially heterogeneous
mechanical properties within the osteocytes.

The temporal accumulation of SASP factors, including *Il6*, *Tnf*, and *Mmp12* (Figure 1C),
suggests that SASP-driven signaling may also play a crucial role in inducing oxidative stress and
contributing to F-actin expansion. Previous studies have demonstrated that pro-inflammatory
cytokines, such as those included in the SASP, can amplify oxidative stress, leading to
cytoskeletal remodeling and impaired cellular function.[17,80,86,87] These findings align with
interpretations in the field but require further investigation to directly link specific SASP factors
to oxidative stress and the biophysical changes observed in osteocytes.

Overall, these findings demonstrate that local SASP factor accumulation significantly altered
osteocyte F-actin morphology, potentially impairing their mechanobiological function. In
contrast, the paracrine effects of SASP factors were less pronounced in influencing F-actin
expansion, likely due to the relatively brief exposure to oxidative stress compared to the longer-
term exposure experienced by osteocytes in the Irradiated groups.

The dendrite lengths of irradiated osteocytes were observed to be generally shorter compared to
the Control group (see **Figure S3**). Specifically, normalized dendrite lengths in the irradiated
groups significantly decreased by Day 14 ($p < 0.0001$) and continued to decline through Day 20
($p < 0.001$, **Figure 4D**). While not statistically significant, dendritic lengths in both CM-
supplemented groups were also reduced compared to the Control cells, aligning with
measurements from Day 7 (**Figure 4D** and **Figure S3**). To mitigate confounding effects due to
variations in F-actin area, dendrite lengths were normalized to soma areas, which was quantified
via NeuriteQuant (**Figure 4C**). This analysis underscores the structural alterations induced by

both local and systemic SASP effects on osteocytes structure and morphology. In particular, the irradiated osteocytes with local SASP accumulation exhibited diminished dendritic network connectivity, which may suggest reduced intercellular communication and mechanosensation capabilities.

Figure 4: Quantitative analyses of F-actin area and dendritic structures in osteocytes following SASP factor exposure compared to non-senescent controls. A) Representative images of F-actin organization in osteocytes from Control and Irradiated groups. Scale bar: Control: 10 µm, Irradiated: 50 µm. **B)** Quantification of average F-actin areas using CellProfiler from IF micrographs across all experimental conditions. **C)** Representative images of semi-automated segmentation and quantification of dendrites and soma performed using NeuriteQuant. Scale bar: 50 µm. **D)** Quantification of average dendrite length normalized to soma area, calculated from IF micrographs of all conditions using NeuriteQuant. All quantitative osteocyte morphology results were obtained by analyzing 14 IF

429   micrographs per group across two cell culture wells (n = 14 for all groups, presented as mean ± SD).

430   Significance levels are indicated as *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

3.3 Osteocyte cytoskeleton stiffening as a biophysical marker for senescence

To characterize the longitudinal changes in cytoskeletal stiffness due to local and paracrine

exposure to SASP factors, we employed live single-cell nanoindentation (**Figures 5A-B**). At all

study time points, osteocytes from the Irradiated groups exhibited significantly higher Young's

modulus (*i.e*. stiffnesses) compared to those in the Control and CM-supplemented groups (see

Figure 5C). The CM-d group displayed no significant change in stiffness compared to the

Control (**Figure 5C**). This increase in cytoskeletal moduli could be linked to the previously

demonstrated expansion in F-actin area (see **Figure 4B**).

The stiffening of the osteocyte cytoskeleton, coupled with the observed senescence-associated

heterogeneity in cytoskeletal organization (**Figure 4A**), suggests a decline in cell motility and

possibly mechanosensory properties. Such degenerative changes potentially lead to deterioration

in interactions of osteocytes with the ECM, a phenomenon similarly reported in studies involving

fibroblasts.[88] The influence of substrate stiffness on cellular behavior, particularly the SASP

secretion in senescent cells, has been previously studied in other cell types.[89] Gresham et al.

demonstrated that compliant substrates (e.g., polyacrylamide hydrogels with a stiffness of 50 kPa)

can mitigate SASP secretion in stress-induced senescent MSCs. [89] This reduction was evident in

key pro-inflammatory markers and DNA damage-associated signals (e.g., γH2AX). Their

findings highlight the importance of substrate mechanical properties in regulating senescence-

associated phenotypes. In our study, all experimental groups were cultured on the same rigid

tissue culture plastic (TCP, ~1 GPa) coated with rat-derived collagen type I, ensuring that

substrate effects were applied uniformly across all groups. This uniformity minimizes substrate

stiffness as a confounding factor in our comparative analyses of cytoskeletal stiffening and

SASP-related alterations between irradiated and control osteocytes. However, it is noteworthy

that the rigid substrate likely exerted greater mechanical resistance compared to compliant

hydrogels or native bone matrix, which may have amplified cytoskeletal stiffening and SASP-

related changes in our study. These findings underscore the importance of exploring compliant

and tunable stiffness substrates in future studies to better replicate the mechanical properties of

the 3D bone microenvironment and refine our understanding of senescence in osteocytes.

In vivo, osteocytes are embedded within the mineralized bone matrix, which exhibits a highly heterogeneous and dynamic mechanical environment.[23,90,91] Cortical bone has a stiffness range of 17–20 GPa, while trabecular bone exhibits stiffness values between 10 and 3,000 MPa, depending on its density and microarchitecture.[90,91] This mechanical environment is critical for osteocyte mechanosensation, as mechanical forces are transmitted through the extracellular matrix and dendritic networks to regulate bone remodeling.[23] The high stiffness of the in vitro TCP substrate differs markedly from the in vivo conditions, potentially amplifying stress-induced phenotypes such as cytoskeletal stiffening and altered mechanotransduction.

These findings indicate that while irradiated osteocytes consistently exhibit cytoskeletal stiffening, systemic exposure to SASP factors results in less significant changes in cytoskeletal stiffness. Future studies employing 3D culture systems with tunable stiffness substrates or more physiologically relevant bone-mimetic scaffolds are needed to validate these findings and assess the role of substrate mechanical properties in osteocyte mechanosensory function.

Figure 5: Irradiated osteocytes exhibit significant increases in cytoskeletal stiffness. A)
Representative image of our live single-cell mechanical testing setup. **B)** Live cells attached to the bottom
of the cell culture plate being indented. Red dots indicate the probe size and locations of indentation. The
main image was obtained via image stitching, and the inset magnifies an individual indentation site for
better visibility. **C)** Young's moduli of osteocytes cytoskeleton measured using Hertzian contact mechanic
model. Data from 50 cells per condition are presented as mean ± SD. Significance levels are denoted as *:
$p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$. **D)** Summary of the study: Irradiation-induced
senescence in osteocytes is characterized by changes in micromechanical properties, morphological
features, and biomolecular markers.

4. Conclusions and Future Directions

This longitudinal study highlights the impact of continuous local SASP factor accumulation, resulting from irradiation, on distinct senescence-associated biophysical markers, supported by corresponding biomolecular changes (**Figure 5D**). Importantly, senescent osteocytes in this study exhibit SASP profiles consistent with those observed in other cell types, including increased secretion of pro-inflammatory cytokines such as *Il-6* and matrix-degrading enzymes such as *Mmp-12* (**Figure 1**).[92] Our findings demonstrate that local SASP accumulation has a pronounced effect on osteocyte cytoskeletal properties, including significant stiffening and heterogeneity in cytoskeletal organization. To build on these findings, future research could incorporate Western blot analysis to validate the observed senescence-associated changes at the protein level. This approach would complement and support the RT-qPCR data by confirming that transcriptional changes in key markers, such as *p16* and *p21*, are reflected at the protein level. Additionally, flow cytometry (FACS) could provide direct evidence of cell cycle disruption by analyzing the distribution of cells across different phases of the cell cycle. Additionally, future studies will expand upon this work by investigating changes in receptors for SASP cues and quantifying the bioactivity of secreted SASP proteins. These analyses will provide deeper insights into the mechanisms through which SASP factors mediate their effects and their functional consequences on cellular and tissue-level remodeling. Such efforts will further inform the development of targeted therapeutic strategies to mitigate the deleterious effects of cellular senescence.

The observed increase in cytoskeletal stiffness in senescent osteocytes also aligns with previous studies in senescent fibroblasts, which exhibit alterations in the cytoskeleton and changes in actin organization leading to altered cell morphology and diminished proliferation and migration.[86] Similarly, studies in MSCs have shown that irradiation-induced senescence leads to altered cytoskeletal and focal adhesion organization and reduced cellular migratory capacity.[93] These parallels highlight the conserved nature of cytoskeletal alterations across different cell types and expand our understanding of osteocyte responses to localized and systemic senescence effects. Notably, in our present study, systemic SASP exposure (CM groups) did not induce substantial changes in cytoskeletal stiffness or morphological properties compared to controls, suggesting a more localized influence of SASP on osteocyte biophysical properties. However, trends observed

in the CM-d group point to the potential role of cellular maturation in modulating SASP uptake,
warranting further investigation.

Osteocyte dysfunction, driven by senescence-associated cytoskeletal changes, plays a pivotal
role in the pathophysiology of osteoporosis.[5,92] Cytoskeletal dynamics are integral to osteocyte
morphology and function, enabling these cells to sense and transduce mechanical signals via
their dendritic processes.[23] The observed cytoskeletal stiffening and altered organization likely
impair mechanosensing, reducing the ability of osteocytes to adapt to mechanical stimuli and
coordinate bone remodeling.[94] Furthermore, SASP, characterized by inflammatory cytokines
and matrix-degrading enzymes, exacerbates bone fragility by promoting osteoclast activity and
inhibiting osteoblast function,[92,95] processes involved in bone remodeling and tightly regulated
by osteocytes. This interplay between cytoskeletal changes and SASP secretion highlights the
potential contribution of senescent osteocytes to the microarchitectural deterioration observed in
osteoporosis, a condition closely linked to age-related imbalance in bone remodeling, reduction
in osteocyte viability/function, and chronic inflammation that contributes to progressive bone
loss and increased fragility over time.[96]

While these findings provide a robust framework for understanding osteocyte dysfunction, the
use of rigid tissue culture plastic (TCP, ~1 GPa) in our study may have amplified stress-induced
phenotypes. In vivo, osteocytes are embedded within a highly heterogeneous bone matrix, with
stiffness values ranging from 17–20 GPa in cortical bone to 10–3,000 MPa in trabecular bone,
depending on density and architecture.[90,91] Furthermore, 2D cultures may not entirely capture
the intercellular interactions and extracellular matrix dynamics found in living tissues, which are
crucial for cellular behavior and function. The pronounced differences between in vitro and in
vivo mechanical environments underscore the importance of future studies using 3D culture
systems or bone-mimetic scaffolds that better replicate physiologically and clinically relevant
conditions. Another limitation of this study was the inability to characterize the same senescent
cells across multiple assays, which would establish direct correlations between senescence
markers and functional outcomes. Single-cell approaches, such as RNA sequencing or multi-
modal profiling, could address this limitation by revealing heterogeneity within senescent cell
populations and uncovering relationships between SASP profiles and biophysical changes.
However, these methods are resource-intensive and currently impractical for large-scale studies.

Future research integrating such advanced assays could provide deeper insights into the dynamic
and heterogeneous nature of senescence, particularly in osteocytes, and their role in bone
remodeling.

To address these limitations, future studies will investigate the relationship between cytoskeletal
stiffening, motility, and mechanosensory properties in osteocytes using live-cell imaging of
dendritic processes and 3D culture models with tunable stiffness. Additionally, combining our
nanoindentation system with real-time PCR or immunocytochemistry will allow us to explore
key mechanosignaling pathways, such as YAP/TAZ and FAK, to elucidate the molecular
mechanisms underlying these changes. The inclusion of proteomic measures in future studies
could enhance the characterization of SASP factors and offer a more comprehensive
understanding of senescence processes. By establishing a causal link between cytoskeletal
stiffening, mechanotransduction impairment, and osteocyte-driven bone remodeling dysfunction,
these efforts aim to validate our findings and uncover potential therapeutic targets for mitigating
prevalent age-related bone diseases like osteoporosis.

Author Contributions

560 MT designed the study, experiments, and concept. JL and MT performed the experiments,
561 researched, analyzed, interpreted the data, and wrote the original manuscript. CK, HS, MAP, and
562 MT analyzed, interpreted the data, and revised manuscript. MT, JLK, and KMM provided
563 resources. All authors reviewed and revised the manuscript. All authors read and approved the
564 final manuscript.

Acknowledgements

566 This research and MT are supported by the Walker Department of Mechanical Engineering at
567 The University of Texas at Austin. KMM is supported by NCI (RO1 CA198279 and CA250905).

Conflict of Interest Statement

569 The authors declare no conflict of interest in regard to this work.

Data Availability

571 The data that support the findings of this study are available on request from the corresponding
572 author.

References

[1]   N. Sarafrazi, *Osteoporosis or Low Bone Mass in Older Adults: United States, 2017-2018*, Atlanta, Georgia, **2021**.

[2]   U. Herbig, M. Ferreira, L. Condel, D. Carey, J. M. Sedivy, *Science (1979)* **2006**, *311*, 1257.

[3]   F. Wang, F. Cai, R. Shi, X.-H. Wang, X.-T. Wu, *Osteoarthritis Cartilage* **2016**, *24*, 398.

[4]   R. J. Pignolo, S. F. Law, A. Chandra, *JBMR Plus* **2021**, *5*, DOI 10.1002/jbm4.10488.

[5]   J. N. Farr, J. Kaur, M. L. Doolittle, S. Khosla, J. Farr, M. Doolittle, *Curr Osteoporos Rep* **2020**, *18*, 559.

[6]   R. J. Pignolo, S. F. Law, A. Chandra, *JBMR Plus* **2021**, *5*, DOI 10.1002/JBM4.10488.

[7]   J. N. Farr, D. G. Fraser, H. Wang, K. Jaehn, M. B. Ogrodnik, M. M. Weivoda, M. T. Drake, T. Tchkonia, N. K. Lebrasseur, J. L. Kirkland, L. F. Bonewald, R. J. Pignolo, D. G. Monroe, S. Khosla, *Journal of Bone and Mineral Research* **2016**, *31*, 1920.

[8]   J. N. Farr, S. Khosla, *Bone* **2019**, *121*, 121.

[9]   S. Chaib, T. Tchkonia, J. L. Kirkland, *Nat Med* **2022**, *28*, 1556.

[10]   J. N. Farr, M. Xu, M. M. Weivoda, D. G. Monroe, D. G. Fraser, J. L. Onken, B. A. Negley, J. G. Sfeir, M. B. Ogrodnik, C. M. Hachfeld, N. K. LeBrasseur, M. T. Drake, R. J. Pignolo, T. Pirtskhalava, T. Tchkonia, M. J. Oursler, J. L. Kirkland, S. Khosla, *Nature Medicine 2017 23:9* **2017**, *23*, 1072.

[11]   S. Chaib, T. Tchkonia, J. L. Kirkland, *Nat Med* **2022**, *28*, 1556.

[12]   U. Tripathi, A. Misra, T. Tchkonia, J. L. Kirkland, *Mech Ageing Dev* **2021**, *198*, 111548.

[13]   S. Lopes-Paciencia, E. Saint-Germain, M.-C. Rowell, A. F. Ruiz, P. Kalegari, G. Ferbeyre, *Cytokine* **2019**, *117*, 15.

[14]   S. Khosla, J. N. Farr, J. L. Kirkland, *J Clin Endocrinol Metab* **2018**, *103*, 1282.

[15]   J. N. Farr, S. Khosla, *Bone* **2019**, *121*, 121.

[16]   F. Chowdhury, B. Huang, N. Wang, *Cytoskeleton* **2021**, *78*, 249.

598   [17]   Y. J. Kim, M. J. Cho, W. D. Yu, M. J. Kim, S. Y. Kim, J. H. Lee, *Cells* **2022**, *11*, DOI
599   10.3390/cells11182896.

600   [18]   A. M. D. Malone, N. N. Batra, G. Shivaram, R. Y. Kwon, L. You, C. H. Kim, J. Rodriguez,
601   K. Jair, C. R. Jacobs, *American Journal of Physiology-Cell Physiology* **2007**, *292*, C1830.

602   [19]   W. M. Jackson, M. J. Jaasma, R. Y. Tang, T. M. Keaveny, *American Journal of*
603   *Physiology-Cell Physiology* **2008**, *295*, C1007.

604   [20]   Q. Wan, E. Cho, H. Yokota, S. Na, *J Bone Miner Metab* **2013**, *31*, 520.

605   [21]   H. Hemmatian, A. D. Bakker, J. Klein-Nulend, G. H. van Lenthe, *Curr Osteoporos Rep*
606   **2017**, *15*, 401.

607   [22]   M. C. M. Li, S. K. H. Chow, R. M. Y. Wong, L. Qin, W. H. Cheung, *J Orthop Translat*
608   **2021**, *29*, 1.

609   [23]   L. Qin, W. Liu, H. Cao, G. Xiao, *Bone Research 2020 8:1* **2020**, *8*, 1.

610   [24]   L. F. Bonewald, *Journal of Bone and Mineral Research* **2011**, *26*, 229.

611   [25]   N. B. M. Rafiq, Y. Nishimura, S. V. Plotnikov, V. Thiagarajan, Z. Zhang, S. Shi, M.
612   Natarajan, V. Viasnoff, P. Kanchanawong, G. E. Jones, A. D. Bershadsky, *Nat Mater* **2019**,
613   *18*, 638.

614   [26]   S. Seetharaman, S. Etienne-Manneville, *J Cell Sci* **2019**, *132*, DOI 10.1242/jcs.232843.

615   [27]   T. Tchkonia, Y. Zhu, J. Van Deursen, J. Campisi, J. L. Kirkland, *J Clin Invest* **2013**, *123*,
616   DOI 10.1172/JCI64098.

617   [28]   N. A. Sims, *Journal of Bone and Mineral Research* **2016**, *31*, 1917.

618   [29]   J. N. Farr, S. Khosla, *Bone* **2019**, *121*, 121.

619   [30]   J. N. Farr, J. Kaur, M. L. Doolittle, S. Khosla, J. Farr, M. Doolittle, *Curr Osteoporos Rep*
620   **2020**, *18*, 559.

[31]   S. Khosla, J. N. Farr, T. Tchkonia, J. L. Kirkland, *Nat Rev Endocrinol* **2020**, *16*, 263.

[32]   A. Hernandez-Segura, J. Nehme, M. Demaria, *Trends Cell Biol* **2018**, *28*, 436.

[33] M. Al-Jumayli, S. L. Brown, I. J. Chetty, M. Extermann, B. Movsas, *Semin Radiat Oncol*
2022, *32*, 172.

[34] P. Ortiz-Montero, A. Londoño-Vallejo, J.-P. Vernot, **n.d.**, DOI 10.1186/s12964-017-0172-3.

[35] C. A. Schmitt, B. Wang, M. Demaria, *Nature Reviews Clinical Oncology 2022 19:10* **2022**,
19, 619.

[36] Z. Yao, B. Murali, Q. Ren, X. Luo, D. V. Faget, T. Cole, B. Ricci, D. Thotala, J. Monahan,
J. M. Van Deursen, D. Baker, R. Faccio, J. K. Schwarz, S. A. Stewart, *Cancer Res* **2020**,
80, 1171.

[37] R. L. Cohn, N. S. Gasek, G. A. Kuchel, M. Xu, *Trends Cell Biol* **2023**, *33*, 9.

[38] M. Ogrodnik, J. Carlos Acosta, P. D. Adams, F. d'Adda di Fagagna, D. J. Baker, C. L.
Bishop, T. Chandra, M. Collado, J. Gil, V. Gorgoulis, F. Gruber, E. Hara, P. Jansen-Dürr,
D. Jurk, S. Khosla, J. L. Kirkland, V. Krizhanovsky, T. Minamino, L. J. Niedernhofer, J. F.
Passos, N. A. R. Ring, H. Redl, P. D. Robbins, F. Rodier, K. Scharffetter-Kochanek, J. M.
Sedivy, E. Sikora, K. Witwer, T. von Zglinicki, M. H. Yun, J. Grillari, M. Demaria, *Cell*
2024, *187*, 4150.

[39] D. A. Alcorta, Y. Xiong, D. Phelps, G. Hannon, D. Beach, J. C. Barrett, *Proceedings of the*
National Academy of Sciences **1996**, *93*, 13742.

[40] M. Fischer, M. Quaas, L. Steiner, K. Engeland, *Nucleic Acids Res* **2016**, *44*, 164.

[41] A. O. Masson, R. Hess, K. O'Brien, K. L. Bertram, P. Tailor, E. Irvine, G. Ren, R. J.
Krawetz, *Mech Ageing Dev* **2015**, *149*, 31.

[42] X. CHEN, W. ZHANG, Y. F. GAO, X. Q. SU, Z. H. ZHAI, *Cell Res* **2002**, *12*, 229.

[43] K. Itahana, J. Campisi, G. P. Dimri, **n.d.**

[44] B. Wang, J. Han, J. H. Elisseeff, M. Demaria, *Nature Reviews Molecular Cell Biology*
2024 **2024**, 1.

[45] B. Y. Lee, J. A. Han, J. S. Im, A. Morrone, K. Johung, E. C. Goodwin, W. J. Kleijer, D.
DiMaio, E. S. Hwang, *Aging Cell* **2006**, *5*, 187.

[46]   M. J. Yousefzadeh, J. Zhao, C. Bukata, E. A. Wade, S. J. McGowan, L. A. Angelini, M. P.
650   Bank, A. U. Gurkar, C. A. McGuckian, M. F. Calubag, J. I. Kato, C. E. Burd, P. D.
651   Robbins, L. J. Niedernhofer, *Aging Cell* **2020**, *19*, DOI 10.1111/acel.13094.

652   [47]   M. Rall-Scharpf, T. W. P. Friedl, S. Biechonski, M. Denkinger, M. Milyavsky, L.
653   Wiesmüller, *Aging* **2021**, *13*, 21066.

654   [48]   B. Wang, L. Wang, N. S. Gasek, C. L. Kuo, J. Nie, T. Kim, P. Yan, J. Zhu, B. L. Torrance,
655   Y. Zhou, L. C. Flores, C. Allen, A. M. Andrade, C. Guo, R. L. Cohn, E. R. Jellison, J. M.
656   Bartley, G. A. Kuchel, S. Li, T. Pirtskhalava, T. Tchkonia, S. Yadav, L. Haynes, J. L.
657   Kirkland, Y. Ikeno, M. Xu, *Cell Metab* **2024**, *36*, 1795.

658   [49]   H. A. Shaban, S. M. Gasser, *Cell Death & Differentiation 2023 32:1* **2023**, *32*, 9.

659   [50]   S. Chaib, T. Tchkonia, J. L. Kirkland, *Nat Med* **2022**, *28*, 1556.

660   [51]   W. Huang, L. T. J. Hickson, A. Eirin, J. L. Kirkland, L. O. Lerman, *Nat Rev Nephrol* **2022**,
661   *18*, 611.

662   [52]   S. Khosla, J. N. Farr, T. Tchkonia, J. L. Kirkland, *Nat Rev Endocrinol* **2020**, *16*, 263.

663   [53]   U. Tripathi, A. Misra, T. Tchkonia, J. L. Kirkland, *Mech Ageing Dev* **2021**, *198*, 111548.

664   [54]   Y. Zhu, T. Tchkonia, T. Pirtskhalava, A. C. Gower, H. Ding, N. Giorgadze, A. K. Palmer, Y.
665   Ikeno, G. B. Hubbard, M. Lenburg, S. P. O'hara, N. F. Larusso, J. D. Miller, C. M. Roos,
666   G. C. Verzosa, N. K. Lebrasseur, J. D. Wren, J. N. Farr, S. Khosla, M. B. Stout, S. J.
667   McGowan, H. Fuhrmann-Stroissnigg, A. U. Gurkar, J. Zhao, D. Colangelo, A. Dorronsoro,
668   Y. Y. Ling, A. S. Barghouthy, D. C. Navarro, T. Sano, P. D. Robbins, L. J. Niedernhofer, J.
669   L. Kirkland, *Aging Cell* **2015**, *14*, 644.

670   [55]   J. N. Farr, M. Xu, M. M. Weivoda, D. G. Monroe, D. G. Fraser, J. L. Onken, B. A. Negley,
671   J. G. Sfeir, M. B. Ogrodnik, C. M. Hachfeld, N. K. LeBrasseur, M. T. Drake, R. J. Pignolo,
672   T. Pirtskhalava, T. Tchkonia, M. J. Oursler, J. L. Kirkland, S. Khosla, *Nature Medicine*
673   *2017 23:9* **2017**, *23*, 1072.

674   [56]   J. N. Farr, D. Saul, M. L. Doolittle, J. Kaur, J. L. Rowsey, S. J. Vos, M. N. Froemming, A.
675   B. Lagnado, Y. Zhu, M. Weivoda, Y. Ikeno, R. J. Pignolo, L. J. Niedernhofer, P. D.

Robbins, D. Jurk, J. F. Passos, N. K. LeBrasseur, T. Tchkonia, J. L. Kirkland, D. G. Monroe, S. Khosla, *J Clin Invest* **2023**, *133*, 366.

[57] M. Tilton, M. Weivoda, M. Astudillo Potes, A. Gingery, A. Y. Liu, T. Tchkonia, L. Lu, J. L. Kirkland, *Aging Cell* **2024**, e14421.

[58] Y. Kato, J. J. Windle, B. A. Koop, G. R. Mundy, L. F. Bonewald, *Journal of Bone and Mineral Research* **1997**, *12*, 2014.

[59] B. Zhang, P. Shang, *STAR Protoc* **2024**, *5*, 103186.

[60] S. L. York, P. Sethu, M. M. Saunders, *Ann Biomed Eng* **2016**, *44*, 1170.

[61] J. D. Gardinier, A. Chougule, C. Zhang, *J Cell Physiol* **2022**, *237*, 2249.

[62] C. A. Grant, D. J. Brockwell, S. E. Radford, N. H. Thomson, *Biophys J* **2009**, *97*, 2985.

[63] A. Demichelis, C. Divieto, L. Mortati, S. Pavarelli, G. Sassi, M. P. Sassi, *J Biomech* **2015**, *48*, 1099.

[64] L. Dehmelt, G. Poplawski, E. Hwang, S. Halpain, *BMC Neurosci* **2011**, *12*, 100.

[65] D. Zauchner, M. Z. Müller, M. Horrer, L. Bissig, F. Zhao, P. Fisch, S. S. Lee, M. Zenobi-Wong, R. Müller, X. H. Qin, *Nature Communications 2024 15:1* **2024**, *15*, 1.

[66] D. A. Alcorta, Y. Xiong, D. Phelps, G. Hannon, D. Beach, J. C. Barrett, *Proceedings of the National Academy of Sciences* **1996**, *93*, 13742.

[67] A. B. Niculescu, X. Chen, M. Smeets, L. Hengst, C. Prives, S. I. Reed, *Mol Cell Biol* **1998**, *18*, 629.

[68] T. Ando, T. Kawabe, H. Ohara, B. Ducommun, M. Itoh, T. Okamoto, *Journal of Biological Chemistry* **2001**, *276*, 42971.

[69] J. D. Gardinier, A. Chougule, C. Zhang, *J Cell Physiol* **2022**, *237*, 2249.

[70] L. Qin, W. Liu, H. Cao, G. Xiao, *Bone Res* **2020**, *8*, 23.

[71] B. F. Boyce, L. Xing, *Arch Biochem Biophys* **2008**, *473*, 139.

[72] M. C. Walsh, Y. Choi, *Front Immunol* **2014**, *5*, DOI 10.3389/fimmu.2014.00511.

[73] G. Schett, S. Hayer, J. Zwerina, K. Redlich, J. S. Smolen, *Nat Clin Pract Rheumatol* **2005**,
1, 47.

[74] S. M. Krane, M. Inada, *Bone* **2008**, *43*, 7.

[75] M. H. Raymond, B. C. Schutte, J. C. Torner, T. L. Burns, M. C. Willing, *Genomics* **1999**,
60, 210.

[76] R. T. Ingram, Y. K. Park, B. L. Clarke, L. A. Fitzpatrick, *J Clin Invest* **1994**, *93*, 989.

[77] J. Li, H. Zhang, C. Yang, Y. Li, Z. Dai, *J Bone Miner Metab* **2016**, *34*, 367.

[78] H. M. Davis, R. Pacheco-Costa, E. G. Atkinson, L. R. Brun, A. R. Gortazar, J. Harris, M.
Hiasa, S. A. Bolarinwa, T. Yoneda, M. Ivan, A. Bruzzaniti, T. Bellido, L. I. Plotkin, *Aging*
Cell **2017**, *16*, 551.

[79] D. Zhao, R. Liu, G. Li, M. Chen, P. Shang, H. Yang, J. X. Jiang, H. Xu, *Front Physiol*
2020, *11*, DOI 10.3389/fphys.2020.00299.

[80] G. La Verde, V. Artiola, V. Panzetta, M. Pugliese, P. A. Netti, S. Fusco, *Biomedicines* **2021**,
9, DOI 10.3390/BIOMEDICINES9091102.

[81] A. Buchwalter, M. W. Hetzer, *Nat Commun* **2017**, *8*, 328.

[82] I. Heckenbach, G. V. Mkrtchyan, M. Ben Ezra, D. Bakula, J. S. Madsen, M. H. Nielsen, D.
Oró, B. Osborne, A. J. Covarrubias, M. L. Idda, M. Gorospe, L. Mortensen, E. Verdin, R.
Westendorp, M. Scheibye-Knudsen, *Nat Aging* **2022**, *2*, 742.

[83] D. Dikovskaya, J. J. Cole, S. M. Mason, C. Nixon, S. A. Karim, L. McGarry, W. Clark, R.
N. Hewitt, M. A. Sammons, J. Zhu, D. Athineos, J. D. G. Leach, F. Marchesi, J. van Tuyn,
S. W. Tait, C. Brock, J. P. Morton, H. Wu, S. L. Berger, K. Blyth, P. D. Adams, *Cell Rep*
2015, *12*, 1483.

[84] A. Milzani, I. DalleDonne, R. Colombo, *Arch Biochem Biophys* **1997**, *339*, 267.

[85] I. Dalle-Donne, R. Rossi, A. Milzani, P. Di Simplicio, R. Colombo, *Free Radic Biol Med*
2001, *31*, 1624.

[86] K. Nishio, A. Inoue, *Histochem Cell Biol* **2005**, *123*, 263.

[87] O. Moujaber, F. Fishbein, N. Omran, Y. Liang, I. Colmegna, J. F. Presley, U. Stochaj, *Cell*
Mol Life Sci **2019**, *76*, 1169.

[88] C. Schulze, F. Wetzel, T. Kueper, A. Malsen, G. Muhr, S. Jaspers, T. Blatt, K.-P. Wittern,
H. Wenck, J. A. Käs, *Biophys J* **2010**, *99*, 2434.

[89] R. C. H. Gresham, A. C. Filler, S. W. Fok, M. Czachor, N. Schmier, C. Pearson, C. Bahney,
J. K. Leach, *J Biomed Mater Res A* **2024**, *112*, 770.

[90] E. Szabo, C. Rimnac, *J Mech Behav Biomed Mater* **2022**, *125*, 104889.

[91] E. F. Morgan, G. U. Unnikrisnan, A. I. Hussein, *Annu Rev Biomed Eng* **2018**, *20*, 119.

[92] M. Wan, E. F. Gray-Gaillard, J. H. Elisseeff, *Bone Research 2021 9:1* **2021**, *9*, 1.

[93] D. Ghosh, C. Mejia-Pena, N. Quach, B. Xuan, A. H. Lee, M. R. Dawson, *J Cell Sci* **2020**,
133, jcs232470.

[94] J. U. A. Choi, A. W. Kijas, J. Lauko, A. E. Rowan, *Front Cell Dev Biol* **2022**, *9*, 1.

[95] M. P. Baar, E. Perdiguero, P. Muñoz-Cánoves, P. L. de Keizer, *Curr Opin Pharmacol* **2018**,
40, 147.

[96] B. Mi, Y. Xiong, S. Knoedler, M. Alfertshofer, A. C. Panayi, H. Wang, S. Lin, G. Li, G.
Liu, *Bone Research 2024 12:1* **2024**, *12*, 1.

Supplementary Materials

Figure S1: Representative SA-β-Gal staining images demonstrating senescence in the CM-d-treated

osteocytes.

Figure S2: Nanoindentation curve fitting over the linear region demonstrates the application of Hertzian
contact mechanics to determine elastic modulus values of osteocytes.

Figure S3: Quantitative analysis of average dendrite lengths in osteocytes under different experimental
conditions. Sample size for each group is n = 3, presented as mean ± SD. *: p < 0.05, **: p < 0.01, ***: p
< 0.001, ****: p < 0.0001.

Figure S4: Example of segmentation (center) and quantification (right) of F-actin micrographs (left)

obtained via IF staining, analyzed using CellProfiler. Scale bar: 50 µm. Units in the quantification image

(right): pixel area.

Figure S5: A) RT-qPCR results of senescence, B-C) SASP factors for all conditions, and D) osteocyte markers, normalized to *Gapdh* housekeeping gene expression and compared to the Day 14 post-irradiation group. n = 3, mean ± SD. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

769 **Table S1:** Primer sequences used for RT-qPCR experiments.

Gene	Forward primer sequences	Reversed primer sequences
Cdkn2a (*p16^{Ink4a}*)	GAACTCTTTCGGTCGTACCC	AGTTCGAATCTGCACCGTAGT
Cdkn1 (*p21^{Cip1}*)	GAACATCTCAGGGCCGAAAA	TGCGCTTGGAGTGATAGAAATC
Mmp 12	GTGCCCGATGTACAGCATCTT	GGTACCGCTTCATCCATCTTG
Il-6	ACCACGGCCTTCCCTACTTC	TTGGGAGTGGTATCCTCTGTGA
Tnf	GTTCTGCAAAGGGAGAGTGG	GCACCTCAGGGAAGAGTCTG
Bglap	CCTGAGTCTGACAAAGCCTTCA	GCCGGAGTCTGTTCACTACCTT
Gja 1	ACAAGTCCTTCCCCATCTCTCA	GGGCACAGACACGAATATGATC
Gapdh	ACTCAAGATTGTCAGCAAT	CCATCCACAGTCTTCTGGGT

770