Phage Steering in the Presence of a Competing Bacterial Pathogen

Authors: Sean Czerwinski [1], James Gurney [1][φ]
Affiliation: 1 School of Biology, Georgia State University, Atlanta, Georgia USA.
φ Corresponding Author jgurney2@gsu.edu

Abstract

The rise of antibiotic-resistant bacteria has necessitated the development of alternative therapeutic strategies such as bacteriophage therapy, where viruses infect bacteria, reducing bacterial burden. However, rapid bacterial resistance to phage treatment remains a critical challenge, potentially leading to failure. Phage Steering, which leverages the evolutionary dynamics between phage and bacteria, offers a novel solution by driving bacteria to evolve away from virulence factors or resistance mechanisms. In this study, we examined whether Phage Steering using bacteriophage Luz19 could function in the presence of a competing pathogen, *Staphylococcus aureus* (USA300), while targeting *Pseudomonas aeruginosa* (PAO1). Through *in vitro* co-evolution experiments with and without the competitor, we observed that Luz19 consistently steered *P. aeruginosa* away from the type IV pilus (T4P), a key virulence factor, without interference from *S. aureus*. Genomic analyses revealed mutations in T4P-associated genes, including *pilR* and *pilZ*, which conferred phage resistance. Our findings suggest that Phage Steering remains effective even in polymicrobial environments, providing a promising avenue for enhancing bacteriophage therapy efficacy in complex infections.

Introduction

The increasing prevalence of antibiotic-resistant bacteria necessitates the exploration of alternative therapeutic approaches. Bacterio(phage) therapy, which utilizes viruses that infect and kill bacteria offers a promising avenue. However, the rapid emergence of phage-resistance *in vitro* raises alarms over phage therapy's long-term efficacy [1, 2]. The antagonistic co-evolutionary dynamics of bacteria-phage interactions generates diversity in both the bacterial and phage populations, with reciprocal evolution of defense vs attack alleles occurring in an almost turn-based competition known as Arms Race Dynamics (ARD) [3-5] wherein bacterial resistance to phage emerges, followed by the evolution of phage infectivity [6-11]. Ultimately, in the lab, ARD culminates in the emergence of a resistant genotype that the phages cannot overcome [8], in most cases coming in the form of *de novo* mutations resulting in the loss or structural and functional attenuation of the specific surface receptors to which the bacteriophage binds [8]. Co-evolutionary dynamics between phage and bacteria can also progress from ARD to exhibit fluctuating selection dynamics in which bacteria resistance is not generalized [12]. The shortcoming of bacteriophages is to be expected, as Lenski and Levin proposed, there is an asymmetry in the evolutionary potential between the bacterium and its phage [8, 9]. This asymmetry is due, in part, to a single spontaneous *de novo* mutation resulting in significant receptor structure alteration and therefore acquisition of resistance for the bacterium, whereas the bacteriophage would need multiple successive *de novo* mutations in order to overcome resistance [13, 14]. Phage Steering seeks to exploit the selection of the host in response to phage pressure. In the event that resistance evolves against the phage, the *de novo* mutations that result should lead to the loss or functional attenuation of the binding receptor [9, 15, 16]. Phage Steering utilizes bacteriophages with known receptors, or factors, that are implicated in antimicrobial resistance or in virulence [15].

The intricate dynamics of bacteria-phage interactions, particularly the rapid emergence of phage resistance, pose significant challenges to the long-term efficacy of bacteriophage therapy. This is

especially critical when considering pathogens like *Pseudomonas aeruginosa* (*PA*), a notorious
Gram-negative bacterium known for its intrinsic multi-drug resistance and its role in chronic,
often life-threatening infections [17]. Furthermore, *PA* infections often occur in a poly-microbial
context, most notably alongside *Staphylococcus aureus* (*SA*), significantly complicating
treatment strategies. These two pathogens are the most prevalent in cystic fibrosis infections,
with studies indicating enhanced virulence when they coexist [18, 19]. The dynamic and often
synergistic interactions between *PA* and *SA* in co-infection models [18, 20-23], emphasize the
complexity of *PA*-associated polymicrobial infections. The community interactions found within
a polymicrobial infection has the potential to greatly impact phage therapies and steering [24,
25]. *PA* is an opportunistic pathogen, and likely did not evolve its vast armament of virulence
factors and antimicrobial resistance mechanisms in order to colonize the CF lung, but rather to
outcompete other micro-organisms in the environment [26-29]. Thus, the question remains
whether Phage Steering will prove efficacious in the presence of competing co-infecting
organisms that exert their own selective pressure via competition or otherwise interaction.
66
In this study, we focus on the interplay between *Pseudomonas aeruginosa* - PAO1 (*PA)*
and *Staphylococcus aureus* - USA300 (*SA*) and the phage Luz19. We employ the phage, Luz19,
to steer PAO1 away from the expression or function of the phage receptor, the type IV pilus
(T4P) [30, 31]. Co-evolving with and without the phage and *SA* present we determined if Luz19
is capable of performing Phage Steering on *PA* during bacterial competition. Our phenotypic
assays of T4P activity and whole genome sequencing of evolved lines show that phage resistance
was mediated by mutations in the Type IV pili system implying Phage Steering still works under
competition with *SA* in LB.
75
76 **Materials and Methods**
77
Bacterial culture and phage preparation.
Pseudomonas aeruginosa (Nottingham *PA*O1) and *SA* (USA300) were independently grown at
37°C with agitation (shaking at 200 rpm) in Lysogeny Broth (LB) media (25g^{-L} LB Broth
(Lennox)). For phage stock quantification, Luz19 phage was diluted in 1X Phosphate Buffer
Saline (PBS) and spot plated onto *PA*O1 soft lawns on Lysogeny Broth Agar (LBA, Agar 15g^{-L})
and incubated at 37°C overnight. Plaque Forming Units (PFUs) were counted, and PFU/mL was
calculated. Phage stock was diluted in 1X PBS. The final concentration of phage stock prior to
initial inoculation was approximately 3.0e9 PFU/mL. Phage inoculation dosage was calculated
for a multiplicity of infection (MOI) of 3, from estimated Colony Forming Unit (CFU) per mL by
multiplying the Optical Density at a wavelength of 600 nm OD_{600} by 8.0e8 (Supplemental data).
88
***In vitro* evolution experiment.**
We investigated bacterial evolutionary response to an anti-virulence bacteriophage, Luz19, over
the course of 10 serial transfers (10 days) in the context of presence or absence of a common
competitor, *SA*. A total of three co-evolution conditions were used, *PA* with *SA*, *PA* with Luz19
phage, and *PA* with both phage and *SA*. A control for serial transfer culturing was included with
only *PA* serially transferred for the total 10 transfers. All co-evolution cultures (5 replicates)
were conducted in 1 ml LB in 24 well plates (Corning)® and incubated at 37°C, shaking, for 24
hours. Bacteria colony forming unit (CFU) estimates for initial and subsequent (for *SA*)
inoculations were made using OD_{600} measurements. For accurate CFU estimation from OD_{600},
*PA*O1 and USA300 suspended cultures were spectrophotometrically measured and then serially
diluted and CFU plate counted. An OD_{600}=1.0 was found to contain approximately 8.0e8 and
1.0e8 CFU/mL for *PA*O1 and USA300 respectively. 50 μL of *PA*O1 (1.6e7 CFU) was inoculated
into wells containing 900 μL of LB media. In conditions with phage, bacteria were exposed to

the 16 μL of the initial ancestral Luz19 phage at an MOI of 3:1 (4.9e7 PFU); from Luz19 stock at
3.0e9 PFU/mL at time point 0 (T0).
104
105   In conditions with competing *SA,* 50 μL of USA300 *(*2.0e6 CFU) was inoculated
106   into the wells containing 900 μL of LB media and 50 μL of newly inoculated PAO1. In +*SA*
107   conditions, 50 μL of *SA*, 5.0e6 CFUs, was re-introduced at each transfer to maintain the
108   selective pressure generated by consistent competition.
109
110   In pilot experiments, we found that *SA* went extinct after no more than 3 transfers when co-
111   cultured with PAO1 at comparable relative concentrations (data not shown). After 24 hours, 100
112   μL was transferred into 900 μL of LB media. 300 μL was stored in 25% final concentration
113   glycerol at − 80˚C.
114
In the case that a replicate culture needed to be restarted, the frozen stock was used to resume
the replicate from the most recently stored time point. Due to human error during transferring
the *PA* (-/-) T5 replicates C, D, E were restarted from T4 for each respective replicates. Likewise,
PA + Luz19 T8 E, restarted from T7 for respective replicate. 1 μl of Frozen culture was inoculated
into 1 mL of LB and incubated at 37°C with shaking for 4 hours. 100 μL was then transferred
into 900 μL of LB and normal culturing conditions resumed. Upon culmination of co-evolution
culturing, all Transfer 10 replicates were frozen in glycerol (25% final) and stored at -80˚C.
122
Preparation for Twitching Motility and Growth Assays.
Frozen replicates were cultured in LB at 37°C shaking overnight. The overnight culture was then
transferred into a 1.5 mL microcentrifuge tube, briefly vortexed to break up aggregates, and then
centrifuged at 17,000 RCF for 2 minutes to pellet the cells, then washed 2 times in 1X PBS and
re-pelleted. Cells were resuspended in 1 mL of LB and further diluted in LB to an OD_{600} of 0.05.
In the case of USA300-positive co-cultures, overnight cultures were streaked onto Pseudomonas
Isolation Agar (PIA) and incubated at 37°C overnight to remove *SA*. Single colonies of isolated
PA were incubated overnight at 37°C in LB.
131
132   **qPCR of Phages.**
133   qPCR of phage stock was performed using the QuantStudio 3 rt PCR system and analyzed with
134   the ThermoFisher applied biosystems app. See SI for list of primers used.
135
Twitching Motility Assays.
We performed a twitching motility assay adapted from Alm & Mattick's 1996 protocol, with the
following modifications [32]. A p20 pipette tip was introduced to the cultures then pushed
through a LB agar plate until the pipette tip contacted plastic at the base of the plate and then
promptly removed. Plates were incubated at 37°C for 20 h. Agar was removed to reveal the
imprint of *PA* growth. Imprints were stained with Crystal Violet for 15 min. Excess stain was
gently washed off using sterile diH_2O. Plates were then air dried. Once dry, the stained imprints
were measured across at the widest point. All phenotypic assays were done in triplicate for each
co-evolution replicate for a total of 15 assay replicates-per-condition.
145
Data transformations (log and Box-Cox using the car package for R 4.2.2 [33]) were attempted,
but normality was not achieved across all treatments according to a Shapiro-Wilk test. Due to
heteroskedasticity we used the Kruskal-Wallis test. Pseudo replication was corrected by first
averaging the measurements taken of triplicates. When significant differences were found, post-
hoc analyses were performed using the Dunn's test with Bonferroni correction for Kruskal-
Wallis. All statistical analysis was performed in R 4.2.2.
152

Growth Assays.
We performed a series of growth assays for *PA* of the four evolution conditions and the ancestor.
All lines of the conditions were prepared following established protocols. They were then
inoculated into 180 μL of LB in individual wells of a 96 well plate (VWR®), at an initial
concentration of approximately 8.9e4 CFU per well. Each of the five lines per condition were
subject to four treatments for the growth assays (All Condition x Line x Treatment combinations
were performed in triplicate): no additional treatment (control), *SA* added at a 1:1 CFU ratio to
PA (8.9e4 CFU), ancestral Luz19 phage added at an MOI of 3 (2.7e5 PFU), or both *SA* and Luz19
added. After inoculation with condition lines and treatments, the 96-well plates were loaded
into an Agilent BioTek BioSpa™ Automated Incubator. The plates were incubated statically at
37°C to promote *PA* growth under the controlled treatment conditions. To monitor bacterial
growth, the plates were automatically transferred every 30 minutes into an Agilent BioTek
Cytation™ for OD_{600} measurements. A total of 49 readings per well were collected over a period
of 24.5 hours. The temporal OD_{600} data were subsequently used to analyze growth dynamics
under each treatment condition.

Growth Analysis.
Analysis of growth assays was performed using the Growthcurver package for R [34].
Growthcurver fits the data points collected for each individual replicate's growth curve to a
standard form of the logistics equation used in ecology and evolution [34]. Growth curves had
the area under the growth curves differentiated for each co-evolution condition subject to four
different growth assay treatments, with standard error bars. AUC is the area under the curve,
calculated by summing the areas beneath the experimental curve obtained from the input
measurements using the Growthcurver package [34].

The growth curves of *PA* from different co-evolution conditions under various treatments was
assessed using the area under the curve (AUC). Statistical analyses were performed to determine
significant differences in growth between conditions and treatments. Data transformations (log
and Box-Cox using 'car' and 'MASS' packages for R 4.4.1 respectively[33]) were attempted but
did not achieve normality across all treatments. We corrected for pseudo replication by
averaging every three technical replicates within each Treatment x Condition to yield
independent pseudo replicate values for AUC. To account for the non-normality observed in our
data, we used the ARTool package for R 4.4.1 to perform a non-parametric two-way analysis of
variance with Treatment and Condition as fixed factors. Subsequent post-hoc analysis was
conducted within the ARTool framework to generate pairwise comparisons (with Bonferroni
correction) for the Treatment x Condition interaction. No differences were observed in the
control evolution line (**SI Fig.1**). Similar analyses were performed on intrinsic mean growth
rate (r) and carrying capacity (k). (**SI Fig. 1 & 2**).

DNA Extraction & Purification.
Single colonies, one from each replicate line, were cultured in LB shaking overnight at 37°C.
DNA extraction, isolation, and purification was performed using the Promega Wizard ®
Genomic DNA Purification Kit following the manufacturer's instructions.

Sequencing.
Whole genome sequencing of single colonies from each replicate was performed by SeqCoast
Genomics (Portsmouth, NH, USA). SeqCoast prepared samples using the Illumina DNA Prep
tagmentation kit and unique dual indexes. Sequencing was performed on the Illumina
NextSeq2000 platform using a 300-cycle flow cell kit to produce a 2x150bp paired reads. 1-2%
PhiX control was spiked into the run to support optimal base calling. Read demultiplexing, read

203 trimming, and run analytics were performed using DRAGEN v4.2.7, an on-board analysis
204 software on the NextSeq2000.
205
Genomic Analysis.
Genomic analyses were completed using breseq 0.38.3 [35] computational pipeline to identify
mutations relative to the reference genome Pseudomonas aeruginosa PAO1 [36]. Sequencing of
our ancestral nPAO1 genome and subsequent breseq analysis was used to identify single
nucleotide polymorphisms (SNPs) that are not present in the reference genome [36]. We
compared the ancestral nPAO1 and the transfer control (*PA* t10) to the evolved *PA* to remove
SNPs that were a result of adaptation to the media.
213
Results
Twitching Motility.
Selection by Luz19 phage attenuates twitching motility function in the presence of competing *SA*
(**Fig. 1**). Twitching motility was attenuated in phage-positive conditions in the presence and
absence of competing *SA* (p < 0.05 for both conditions), relative to the ancestral PAO1. There
was no significant difference between the two positive-phage conditions nor between the
ancestor and the negative/negative culture control. *PA* that was co-cultured with *SA* (*PA+SA*)
did not have a significant difference in twitching motility relative to the ancestral PAO1 (p >
0.05).
223
Growth assays and area under the curve.
Our statistical analysis of growth asked two questions. First – did strains evolved under phage
selection develop resistance. When compared to the ancestor, treatment with the ancestral
phage resulted in significantly increased growth in lines evolved under phage pressure. (**Fig. 2**).
Further, the addition or absence of phage produced no significant difference in growth in lines
evolved with the phage. This is consistent with phage resistance. Our second question, did
adaptation to the media or the presence of *SA* alter phage resistance? Our analysis revealed
highly significant effects of both Treatment (F=132.73, df=3, p<2.22e-16) and Condition
(F=41.83, df=4, p<2.22e-16) on the area under the growth curve, as well as significant
Treatment x Condition interaction (F=37.66, df=12, p<2.22e-16). This indicates that not only do
the individual treatments and co-evolution conditions affect growth outcomes, but the effect of
treatment varies depending on the evolutionary history. (**Fig 2**). Subsequent post-hoc analysis
shows that these results are due to phage resistance but only in lines which evolved under phage
pressure. This increased resistance was observed in both the +Luz19 and +*SA*+Luz19 treatments
compared to all phage-naïve lines, including ancestral PAO1 (p = 6.30e-4 and p = 1.65e-7,
respectively), the PAO1t10 control (p = 1.23e-5 and p = 2.26e-9, respectively), and *PA* evolved
with *SA* (p = 6.47e-5 and p = 1.35e-8, respectively). Importantly, the presence of *S. aureus*
during evolution did not significantly alter phage resistance, our analysis found no significant
difference between the +Luz19 and +*SA*+Luz19 conditions (p > 0.05), indicating that co-
evolution with *S. aureus* did not alter resistance to the phage. Neither *SA* co-culture nor the
media alone increased resistance to the phage relative to the ancestral PAO1 (p > 0.05). These
results consistently demonstrate the enhanced phage resistance in phage positive lines,
regardless of the presence of *SA* during evolution.
247
Genomics.
Sequencing revealed a clear signature of evolution in the presence of the phage (**Fig. 3**). Three
SNP variations between two genes directly involved in type IV pilus biogenesis, *pilR* and *pilZ,*
were defined in the conditions subject to co-evolution with the phage. In the two conditions with
phage, 8 out of 10 replicates (5/5 +Phage, 3/5 +Phage/+*SA*) had SNPs in *pilR*, the response

regulator element of the PilSR two component system, responsible for activation of *pilA*
transcription [37].
255
256 Of the remaining two replicates, one (replicate A) had a 197 bp deletion in *pilZ*. The Δ197 bp *pilZ*
257 replicate showed significantly attenuated twitching motility (p < 0.05) with no significant
258 motility difference between it and the *pilR* mutants (**Fig. 3**).
259
Replicate B of the +*SA*+Luz19 condition lacked detectable SNPs in genes directly related to type
IV pilus biogenesis, structure, or function. However, it did have a unique SNP in *pqsR*, which is
a global virulence regulator involved in quorum sensing [38]. It is currently unclear what
specific impact the *pqsR* mutation has on type IV pili expression or function, though the
replicate did show roughly the same attenuation of twitching motility (**Fig. 1**) and resistant
growth curves (**Fig. 2**)
266
Discussion
Phage Steering is the use of phage to reduce or remove pathogenic behaviors from a bacterial
population. Fundamentally, it works by phage making use of a bacterial product, typically an
outer membrane expressed structure, to complete a full infection cycle. Bacteria are therefore
under a selective pressure to modify the structure – engendering resistance to the phage. This
modification also typically leads to a reduction in effectiveness of the structure in its evolved
function. Should the structure not provide an actual fitness benefit in the environment in which
the phage is applied there is no selective pressure to restore the phenotype. In infections like
those seen in the Cystic Fibrosis lung, many virulence factors of *PA* do not provide a fitness
benefit when considered in isolation, only contributing to nonadaptive virulence, increasing
morbidity of people with CF [39]. This is likely due to *PA* not having evolved in the CF lung, as
the CF lung itself is, in evolutionary terms, incredibly new. Before the 1930s, the majority of
people with CF died before the lung became chronically infected with *PA* [40]. However,
virulence factors such as type IV pili likely evolved to help bacteria deal with inter species
competition [41-43]. Therefore, a central requirement of Phage Steering, that negative fitness
effects are outweighed by the benefit of phage resistance, might not hold true if *PA* is in
competition with other bacteria at the time of phage application.
284
To investigate how Phage Steering works during multi-species competition, we designed a
simple experimental setup that provided the most favorable conditions for it to work. We
conducted a co-evolutionary experiment involving either the phage Luz19 and bacterium *SA*,
both in combination, alone or neither. The major result of our work is that regardless of the
presence of *SA,* the phage still steered *PA* away from the expression of type IV pili (**Fig. 1**). We
measured the phenotypic changes in type IV pili with twitching motility and confirmed phage
resistance emerged as effectively in both treatments that had phage, indicating again that the
presence of *SA* did not alter this outcome. Our growth analysis supports that phage resistance is
similar with or without *SA* present (**Fig. 2**). We did see an increase in variability of phage
resistance in the lines without phage, likely due to weak selection or drift (**Fig. 2C**). Our growth
curves of isolates evolved without phage showed an increased AUC when tested in the presence
of *SA* (**Fig. 2D**). This is due to the growth of the *SA*. Finally, we genomically characterized
individual isolates after the evolution experiment and found the mutations selected by Luz19
were found in pili regulatory genes.
299
Examining these specific mutations in more detail, the gene product of *pilA*, is the major
subunit, pilin (PilA) that comprises the type IV pilus shaft [44, 45]. PilS is located within the
inner membrane of *PA* and is responsible for the detection of surplus pilin/PilA subunits
localized to the inner membrane [37]. In the absence of surplus PilA, PilS autophosphorylates,

and subsequently transfers the phosphate group to the response regulator, PilR [37]. PilR then
binds to a specific sequence proximal to the promoter of *pilA;* and with the assistance of the
alternative sigma factor, σ54, *pilA* is transcribed [37] (**Fig. 4**). Two unique SNPs in *pilR* were
found in the phage-positive replicates, both point substitutions that result in amino acid
changes. Of the eight *pilR* SNPs, seven were C192Y(TGC→TAC) with the sole outlier,
P189Q (CCG→CAG). There was no significant difference between the attenuation of twitching
motility between the two *pilR* mutation variants.
311
The gene product of *pilZ* is a critical regulatory element in the biogenesis of type IV pili [46].
PilZ interacts with PilB, an ATPase localized to the cytoplasmic face of the inner membrane that
provides the energy necessary for transferring PilA subunits from the inner membrane to the
base of the growing pilus polymer for pilus extension (**Fig. 4**) [46]. In the absence of functional
PilZ, PilA subunits accumulate in the membrane but fail to assemble into a pilus shaft, resulting
in loss of surface pili and associated phenotypes such as twitching motility and susceptibility to
type IV pili-dependent phages [46] (**Fig. 4**). While the exact mechanism by which PilZ mediates
PilB activity has yet to be fully elucidated, recent findings have shed significant light on the
process. Hendrix et al. identified PlzR, a novel regulator, which binds to PilZ and modulates its
function. Under elevated c-di-GMP levels, PlzR is transcriptionally induced and binds PilZ,
preventing productive PilZ-PilB interaction, thereby inhibiting pilus assembly [47]. In contrast,
at low c-di-GMP levels, FimX – another c-di-GMP-binding protein – interacts directly with PilB
to promote pilus extension, suggesting that PilZ and FimX act as conditional regulators of the
same extension ATPase [47]. Together, these findings establish PilZ as a central mediator
linking secondary messenger signaling to pilus biogenesis. This layered regulatory model,
proposed by Hendrix et al, is supported by our finding of a 197 bp deletion in PilZ in one of the
Luz19-challenged evolutionary lineages (replicate A of +*SA*+Luz19). This loss-of-function
mutation conferred phage resistance and led to significantly attenuated twitching motility. The
emergence of this deletion under strong phage selection highlights PilZ's essentiality for type IV
pilus biogenesis and thus its disruption serves as a rapid path to type IV pili-dependent phage
resistance.
333
Slightly unexpected is the appearance of a quorum sensing mutation in the *pqsR* regulator,
which is also known as *mvfR* , a regulator in the Pseudomonas Quinolone Signal (PQS) quorum
sensing system. The phage pf, also a type IV pilus binding phage has been shown to generate
mutations within the PQS system [48], therefore there is likely some interaction between the
PQS system and type IV pili. Guo et al. showed that while adding exogenous PQS did not alter
the twitching motility, *pqsR* mutants had attenuated twitching [49]. It is somewhat surprising
that this mutation was seen in the *SA*-positive line. PQS is a pivotal quorum sensing molecule in
PA that regulates biofilm formation along with a plethora of virulence factors [50]. The role that
PQS plays in the killing of *SA* is complex with many PQS regulated factors contributing to *PA*'s
competitive advantage. In brief, PQS has antimicrobial properties against *SA* via the action of 2-
heptyl-4-hydroxyquinoline (HHQ) and 2-heptyl-4-hydroxyquinoline N-oxide (HQNO) [51-56].
In PQS mutants (Δ*pqsA* and Δ*pqsH*), a reduced ability to kill *SA* has been shown relative to WT
PA strains [38, 52]. While PQS plays a critical role in *PA*'s ability to kill *SA*, it is part of a
complex network of QS molecules and virulence factors influenced by the PQS quorum sensing
system. Notably, the antimicrobial activity of PQS and its derivatives is modulated by
environmental factors, particularly iron availability [54]. Consequently, competition studies and
further investigations using media that more accurately mimic *in vivo* nutrient-limited
environments, such as SCFM2, may illuminate potential evolutionary trade-offs or reveal
vulnerabilities in Luz19-resistant *PA* harboring *pqsR* mutations.
353

These phenotypic and genomic data point us towards the conclusion that Phage Steering is effective in the presence of *SA* when targeting the type IV pilus. In future work we will expand this to include other bacterial pathogens and commensals, as well as working in media which better mimics infection sites, such as SCFM2 [39]. We know that the essential genome of bacteria, as measured by TN-seq, is dependent on environment, both the biotic (interspecies competition) and the abiotic [57]. It is possible that infection environments which include competition from pathogens and commensals, a host immune system, and nutritional limitations might impose a fitness cost on the loss of structures which phages can target.

Figures

Fig 1. **Twitching motility compared across evolution conditions.** *PA* twitching motility decreased significantly (p < 0.05) in conditions exposed to Luz19 phage relative to the ancestral PAO1, with no significant difference in twitching motility measured between the positive-phage groups with/without SA (p > 0.05) or between non-phage conditions and the ancestral control p > 0.05). Boxplot ranges are 1.5X interquartile ranges.

380
381
Fig. 2. **Growth assays and area under the curve (AUC). A.) Control treatment AUC boxplots (above) and**
mean growth curves (below). *PA* from the five conditions were grown in LB with no additional challenge – *PA* evolved
alongside *SA* had significantly higher AUC than all other conditions (p < 0.05). **B.) *SA* treatment AUC boxplots**
(above) and mean growth curves (below). *PA* grown in LB with *SA* added. *PA* evolved with *SA* had significantly
greater AUC than the ancestral (p < 0.05) but not significantly different from the transfer control PAO1t10 (p > 0.05).). **C.)**
Luz19 treatment AUC boxplots (above) and mean growth curves (below). Conditions subject to co-evolution
with Luz19 (+Luz19 and +*SA*+Luz19) had significantly greater AUC than those not evolved with the phage (p < 0.05) and
were not significantly different from one another (p > 0.05). There was no significant difference between the AUC of
conditions not previously exposed to the phage (p > 0.05) **D.) *SA* & Luz19 treatment AUC boxplots (above) and**
mean growth curves (below). Conditions subject to co-evolution with Luz19 (+Luz19 and +*SA*+Luz19) had
significantly greater AUC than those not evolved with the phage (p < 0.05) and were not significantly different from one
another (p > 0.05). There was no significant difference between the AUC of conditions not previously exposed to the phage
(p > 0.05). Boxplot ranges are the 1.5X interquartile ranges, error bars on the growth curves are Standard Error.

Fig. 3 **Chromosomal single nucleotide polymorphisms (SNPs) defined for all co-evolution condition replicates and the ancestral PAO1.** SNPs were determined using breseq. **_pilR_**. SNPs of two variations in 5 out of 5 (5/5) +Luz19 co-evo replicates and 3/5 +_SA_+Luz19 co-evo replicates. In all but one of these replicate sequences, the SNP in _pilR_ is C192Y (TGC→TAC). Replicate _B_ of the +Luz19 condition had a different SNP in _pilR,_ P189Q (CCG→CAG). Of the replicates in the two conditions that were subject to co-evolution with Luz19, two lacked a SNP in _pilR,_ both in the +_SA_+Luz19 condition. Replicate _B_ instead had a 197 bp deletion in **_pilZ_** . Replicate _A_ of +_SA_+Luz19 was the only replicate co-evolved alongside Luz19 that sequencing did not reveal a _pil_ gene mutation, instead, a PQS system mutation was identified in **_pqsR._**

Fig 4. **Mutations in key regulatory factors of T4P biogenesis confer phage resistance. A.)** PilR serves as the response regulator of *pilA* transcription. The sensor kinase, PilS, phosphorylates PilR, activating it which in turn activate transcription of *pilA* which encodes for the major pilin subunit, PilA. **B.)** PilZ regulates T4P biogenesis. By binding to the extension ATPase, PilB, activates shaft polymerization. PilZ and FimX are necessary for PilB activation. **C.)** Without PilZ (PilZ*) binding, PilB remains inactive and energy is not generated for shaft assembly. **D.)** Defective PilR does not activate transcription of *pilA*, no PilA subunits are synthesized and thus the shaft is not assembled.

References

1. Betts, A., O. Kaltz, and M.E. Hochberg, *Contrasted coevolutionary dynamics between a bacterial pathogen and its bacteriophages.* Proc Natl Acad Sci U S A, 2014. **111**(30): p. 11109-14.

2. Rohde, C., et al., *Expert Opinion on Three Phage Therapy Related Topics: Bacterial Phage Resistance, Phage Training and Prophages in Bacterial Production Strains.* Viruses, 2018. **10**(4).

3. Medel, R., et al., *Arms Race Coevolution: The Local and Geographical Structure of a Host–Parasite Interaction.* Evolution: Education and Outreach, 2010. **3**(1): p. 26-31.

4. Thompson, J.N., *The geographic mosaic of coevolution.* 2019: University of Chicago Press.

5. Thompson, J.N. and M. Pagel, *Coevolution.* Encyclopedia of Evolution. Oxford University Press, Oxford, 2002.

6. Chao, L., B.R. Levin, and F.M. Stewart, *A Complex Community in a Simple Habitat: An Experimental Study with Bacteria and Phage.* Ecology, 1977. **58**(2): p. 369-378.

7. Horne, M.T., *Coevolution of Escherichia coli and bacteriophages in chemostat culture.* Science, 1970. **168**(3934): p. 992-3.

8. Koskella, B. and M.A. Brockhurst, *Bacteria-phage coevolution as a driver of ecological and evolutionary processes in microbial communities.* FEMS Microbiol Rev, 2014. **38**(5): p. 916-31.

9. Lenski, R.E. and B.R. Levin, *Constraints on the Coevolution of Bacteria and Virulent Phage: A Model, Some Experiments, and Predictions for Natural Communities.* The American Naturalist, 1985. **125**(4): p. 585-602.

10. Schrag, S.J. and J.E. Mittler, *Host-parasite coexistence: the role of spatial refuges in stabilizing bacteria-phage interactions.* The American Naturalist, 1996. **148**(2): p. 348-377.

11. Spanakis, E. and M.T. Horne, *Co-adaptation of Escherichia coli and coliphage lambda vir in continuous culture.* J Gen Microbiol, 1987. **133**(2): p. 353-60.

12. Hall, A.R., et al., *Host-parasite coevolutionary arms races give way to fluctuating selection.* Ecol Lett, 2011. **14**(7): p. 635-42.

13. Hall, Alex R., Pauline D. Scanlan, and A. Buckling, *Bacteria ‐Phage Coevolution and the Emergence of Generalist Pathogens.* The American Naturalist, 2011. **177**(1): p. 44-53.

14. Hall, A.R., et al., *Host-parasite coevolutionary arms races give way to fluctuating selection.* Ecology Letters, 2011. **14**(7): p. 635-642.

15. Gurney, J., et al., *Steering Phages to Combat Bacterial Pathogens.* Trends Microbiol, 2020. **28**(2): p. 85-94.

16. Gurney, J., et al., *Phage steering of antibiotic-resistance evolution in the bacterial pathogen, Pseudomonas aeruginosa.* Evol Med Public Health, 2020. **2020**(1): p. 148-157.

17. Diggle, S.P. and M. Whiteley, *Microbe Profile: Pseudomonas aeruginosa: opportunistic pathogen and lab rat.* Microbiology (Reading), 2020. **166**(1): p. 30-33.

18. DeLeon, S., et al., *Synergistic interactions of Pseudomonas aeruginosa and Staphylococcus aureus in an in vitro wound model*. Infect Immun, 2014. **82**(11): p. 4718-28.

19. Hatziagorou, E., et al., *Changing epidemiology of the respiratory bacteriology of patients with cystic fibrosis-data from the European cystic fibrosis society patient registry*. J Cyst Fibros, 2020. **19**(3): p. 376-383.

20. Dalton, T., et al., *An in vivo polymicrobial biofilm wound infection model to study interspecies interactions*. PLoS One, 2011. **6**(11): p. e27317.

21. Korgaonkar, A., et al., *Community surveillance enhances Pseudomonas aeruginosa virulence during polymicrobial infection*. Proc Natl Acad Sci U S A, 2013. **110**(3): p. 1059-64.

22. Korgaonkar, A.K. and M. Whiteley, *Pseudomonas aeruginosa enhances production of an antimicrobial in response to N-acetylglucosamine and peptidoglycan*. J Bacteriol, 2011. **193**(4): p. 909-17.

23. Sibley, C.D., et al., *Discerning the complexity of community interactions using a Drosophila model of polymicrobial infections*. PLoS Pathog, 2008. **4**(10): p. e1000184.

24. Alseth, E.O., et al., *The impact of phage and phage resistance on microbial community dynamics*. PLOS Biology, 2024. **22**(4): p. e3002346.

25. Blazanin, M. and P.E. Turner, *Community context matters for bacteria-phage ecology and evolution*. ISME J, 2021. **15**(11): p. 3119-3128.

26. Langendonk, R.F., D.R. Neill, and J.L. Fothergill, *The Building Blocks of Antimicrobial Resistance in Pseudomonas aeruginosa: Implications for Current Resistance-Breaking Therapies*. Front Cell Infect Microbiol, 2021. **11**: p. 665759.

27. Fernandez-Billon, M., et al., *Mechanisms of antibiotic resistance in Pseudomonas aeruginosa biofilms*. Biofilm, 2023. **5**: p. 100129.

28. Grosso-Becerra, M.V., et al., *Pseudomonas aeruginosa clinical and environmental isolates constitute a single population with high phenotypic diversity*. BMC Genomics, 2014. **15**: p. 318.

29. Qin, S., et al., *Pseudomonas aeruginosa: pathogenesis, virulence factors, antibiotic resistance, interaction with host, technology advances and emerging therapeutics*. Signal Transduct Target Ther, 2022. **7**(1): p. 199.

30. Chibeu, A., et al., *The adsorption of Pseudomonas aeruginosa bacteriophage phiKMV is dependent on expression regulation of type IV pili genes*. FEMS Microbiol Lett, 2009. **296**(2): p. 210-8.

31. Lammens, E., et al., *Representational Difference Analysis (RDA) of bacteriophage genomes*. J Microbiol Methods, 2009. **77**(2): p. 207-13.

32. Alm, R.A. and J.S. Mattick, *Identification of two genes with prepilin-like leader sequences involved in type 4 fimbrial biogenesis in Pseudomonas aeruginosa*. J Bacteriol, 1996. **178**(13): p. 3809-17.

33. Weisberg, J.F.a.S., *An {R} Companion to Applied Regression*. Third ed. 2019: Sage.

34. Sprouffske, K. and A. Wagner, *Growthcurver: an R package for obtaining interpretable metrics from microbial growth curves*. BMC Bioinformatics, 2016. **17**: p. 172.

554 35. Deatherage, D.E. and J.E. Barrick, *Identification of mutations in laboratory-evolved*
555   *microbes from next-generation sequencing data using breseq.* Methods Mol Biol,
556   2014. **1151**: p. 165-88.

557 36. Stover, C.K., et al., *Complete genome sequence of Pseudomonas aeruginosa PAO1,*
558   *an opportunistic pathogen.* Nature, 2000. **406**(6799): p. 959-64.

559 37. Boyd, J.M. and S. Lory, *Dual function of PilS during transcriptional activation of the*
560   *Pseudomonas aeruginosa pilin subunit gene.* J Bacteriol, 1996. **178**(3): p. 831-9.

561 38. Lin, J., et al., *The Pseudomonas Quinolone Signal (PQS): Not Just for Quorum*
562   *Sensing Anymore.* Front Cell Infect Microbiol, 2018. **8**: p. 230.

563 39. Turner, K.H., et al., *Essential genome of Pseudomonas aeruginosa in cystic fibrosis*
564   *sputum.* Proc Natl Acad Sci U S A, 2015. **112**(13): p. 4110-5.

565 40. Davis, P.B., *Cystic Fibrosis Since 1938.* American Journal of Respiratory and Critical
566   Care Medicine, 2006. **173**(5): p. 475-482.

567 41. Yarrington, K.D., T.N. Shendruk, and D.H. Limoli, *The type IV pilus chemoreceptor*
568   *PilJ controls chemotaxis of one bacterial species towards another.* PLoS Biol, 2024.
569   **22**(2): p. e3002488.

570 42. Limoli, D.H., et al., *Interspecies interactions induce exploratory motility in*
571   *Pseudomonas aeruginosa.* Elife, 2019. **8**.

572 43. Sanchez-Pena, A., et al., *Pseudomonas aeruginosa surface motility and invasion*
573   *into competing communities enhance interspecies antagonism.* mBio, 2024. **15**(9):
574   p. e0095624.

575 44. Sastry, P.A., et al., *Comparative studies of the amino acid and nucleotide sequences*
576   *of pilin derived from Pseudomonas aeruginosa PAK and PAO.* J Bacteriol, 1985.
577   **164**(2): p. 571-7.

578 45. Strom, M.S. and S. Lory, *Mapping of export signals of Pseudomonas aeruginosa pilin*
579   *with alkaline phosphatase fusions.* J Bacteriol, 1987. **169**(7): p. 3181-8.

580 46. Guzzo, C.R., et al., *PILZ Protein Structure and Interactions with PILB and the FIMX*
581   *EAL Domain: Implications for Control of Type IV Pilus Biogenesis.* Journal of
582   Molecular Biology, 2009. **393**(4): p. 848-866.

583 47. Hendrix, H., et al., *PlzR regulates type IV pili assembly in Pseudomonas aeruginosa*
584   *via PilZ binding.* Nat Commun, 2024. **15**(1): p. 8717.

585 48. Schwartzkopf, C.M., et al., *Inhibition of PQS signaling by the Pf bacteriophage*
586   *protein PfsE enhances viral replication in Pseudomonas aeruginosa.* Mol Microbiol,
587   2024. **121**(1): p. 116-128.

588 49. Guo, Q., et al., *PqsR-dependent and PqsR-independent regulation of motility and*
589   *biofilm formation by PQS in Pseudomonas aeruginosa PAO1.* Journal of Basic
590   Microbiology, 2014. **54**(7): p. 633-643.

591 50. Pesci, E.C., et al., *Quinolone signaling in the cell-to-cell communication system of*
592   *Pseudomonas aeruginosa.* Proc Natl Acad Sci U S A, 1999. **96**(20): p. 11229-34.

593 51. Filkins, L.M., et al., *Coculture of Staphylococcus aureus with Pseudomonas*
594   *aeruginosa Drives S. aureus towards Fermentative Metabolism and Reduced*
595   *Viability in a Cystic Fibrosis Model.* J Bacteriol, 2015. **197**(14): p. 2252-64.

52. Haussler, S. and T. Becker, *The pseudomonas quinolone signal (PQS) balances life*
and death in Pseudomonas aeruginosa populations. PLoS Pathog, 2008. **4**(9): p.
e1000166.
53. Nairz, M., et al., *The struggle for iron - a metal at the host-pathogen interface*. Cell
Microbiol, 2010. **12**(12): p. 1691-702.
54. Nguyen, A.T., et al., *Cystic Fibrosis Isolates of Pseudomonas aeruginosa Retain Iron-*
Regulated Antimicrobial Activity against Staphylococcus aureus through the Action
of Multiple Alkylquinolones. Front Microbiol, 2016. **7**: p. 1171.
55. Nguyen, A.T. and A.G. Oglesby-Sherrouse, *Interactions between Pseudomonas*
aeruginosa and Staphylococcus aureus during co-cultivations and polymicrobial
infections. Appl Microbiol Biotechnol, 2016. **100**(14): p. 6141-6148.
56. Otto, B.R., A.M. Verweij-van Vught, and D.M. MacLaren, *Transferrins and heme-*
compounds as iron sources for pathogenic bacteria. Crit Rev Microbiol, 1992. **18**(3):
p. 217-33.
57. Ibberson, C.B., et al., *Co-infecting microorganisms dramatically alter pathogen gene*
essentiality during polymicrobial infection. Nat Microbiol, 2017. **2**: p. 17079.
612
613
614