Spatial learning in feature-impoverished environments in *Drosophila*

Yang Chen[1,2], Robert Alfredson[2], Dorsa Motevalli[2], Ulrich Stern[2], and Chung-Hui Yang[2*]

1. Department of Biology, Duke University, Durham, NC 27708
2. Department of Neurobiology, Duke University Medical School, Durham, NC 27710

*Corresponding Author. Email: rebecca.yang@gmail.com

Abstract

The ability to return to memorized goal locations is essential for animal survival. While it is well documented that animals use visual landmarks to locate goals[1,2], how they navigate spatial learning tasks in environments lacking such landmarks remains poorly understood. Here, using a high-throughput spatial learning task we developed to investigate this question in *Drosophila*, we found that *Drosophila* can simultaneously use self-generated olfactory cues and self-motion cues to learn a spatial goal under visually challenging conditions. Specifically, flies mark a rewarded goal location with self-deposited scents, to which they assign a positive value to, and use these scents and their self-motion cues to guide them back to the goal. This learning process is mediated by the mushroom body(MB) – an olfactory learning center responsible for associating odors with reinforcement[3] –and by PFN neurons, which encode egocentric translational velocity[4,5], a self-motion cue. Intriguingly, when the environment is enriched with prominent external olfactory landmarks, flies shift their strategy, prioritizing these landmarks over self-generated cues. Our findings demonstrate that *Drosophila* can dynamically adapt to environmental complexities when solving spatial learning tasks by creating and integrating internal and external cues, revealing an unexpected level of sophistication in their cognitive capacities.

Introduction

The capacity for returning to memorized goal locations in darkness or without any guidance from prominent visual landmarks is critical for animals to survive in nature, especially for nocturnal animals and animals with no or limited vision. Identifying the behavioral strategies and neural circuits that support navigation in such challenging conditions will provide important insights into the neural basis of simple cognition. In principle, animals can rely purely on self-motion cues to estimate their relative distance and angle to a goal location – i.e., to calculate a home vector – in a process known as path integration, and then use the vector to guide their return to the goal[6,7]. For example, desert ants have been shown to be able to gauge whether they have reached the goal by "counting" the number of steps they have taken[8]. In addition, mammals such as rats[9], mice[10], bats[11], and dogs[12], and insects such as bees[13-16], dung beetles[17-19], and fruit flies[20-23] have all been implicated to use path integration for navigation. The exact neural mechanism by which path integration is implemented in the brain remains poorly understood, but self-motion-sensitive spatially selective neurons, such as heading direction cells[24], likely play a critical role.

In addition to self-motion cues, olfactory cues constitute another major source of information animals could utilize when navigating without access to prominent visual cues. Odors can aid successful navigation to goals in at least two ways. First, animals can return to the goal location by simply following the odor plume – i.e., by beaconing – if the goal happens to emit some innately attractive odors, such as those from a food source[25-29]. Second, animals can also use odors present at fixed locations away from the goal to reset the path integrator[30-32]. For example, a recent study suggests that mice can use odors delivered consistently at specific locations along a linear path to the goal to estimate their relative position to the goal[32]. The circuit mechanism responsible for the first strategy should be relatively straightforward, while that for the second requires integration of olfactory cues and self-motion cues in the path-integration circuit.

In this study, we set out to uncover behavioral and neural strategies animals might adopt when searching for a memorized spatial goal in environments that lack both prominent external landmarks, using *Drosophila melanogaster* as our model. *Drosophila* has been shown to be able to perform visual cues-based spatial learning and path integration, and, importantly, possesses several classes of spatially selective neurons (e.g., head direction cells) in its compact brain[4,5,33-36]. Using a high-throughput spatial learning paradigm we developed where an optogenetically delivered reward was placed in an essentially featureless arena, we found that flies were able to learn the location of the rewarded goal relatively quickly. Interestingly, in addition to making use of self-motion cues, our results suggest that, in parallel, they also exploited scent(s) from substances they deposited themselves, assigning it with a positive valence to help them memorize and locate the goal. Moreover, they adjusted the relative reliance on the two strategies depending on whether salient external olfactory landmarks were available in the environment. Our results highlight this model organism's "creativity" and flexibility in solving a spatial task in landmark-impoverished environments, paving the way for future studies into the underlying molecular and neural mechanisms.

Results

Flies can learn the location of a spatial goal without relying on prominent visual cues

To test whether flies can learn a spatial goal location without using prominent visual cues, we designed a spatial learning task where a fly can collect virtual rewards at a goal location in a rectangularly shaped and otherwise "featureless" arena ("flat chamber") that lacks clear visual landmarks (**Fig.1a**). The task was conducted in a custom-designed high-throughput system (SkinnerSys) – comprising hardware and software (SkinnerTrax[37]) – for behavior-dependent optogenetic stimulation of many flies independently and in parallel. The apparati we used each housed 40 individual flies in separate arenas (**Fig. 1b**). We customized SkinnerTrax to place the goal location – shaped as a small (and invisible) circle – at the center of the arena so that it was not directly associated with specific spatial features (e.g., arena corners) (**Fig. 1a**). Rewards were delivered optogenetically. Specifically, each time the fly crossed over the border of the reward circle from the outside, a 250-ms pulse of light was delivered to optogenetically activate a group of neurons (labeled by *0273-Gal4*) that have been previously shown to be a strong positive reinforcer [38,39] (**Fig. 1c**, **Video S1**). Two features of our design are noteworthy. First, to collect additional rewards, the fly must leave the reward circle and then reenter it, as staying inside will not trigger any rewards (**Fig. 1c**). As such, to continually forage for more rewards, flies likely need to have some memory of the reward location. Second, we paired each experimental fly with a yoked control fly that received a reward at the same time as its experimental fly, but whose positions had no influence over reward delivery (**Fig. 1d**).

The length of the task was 4 hours with three 1-hour training sessions (**Fig. 1d**). Rewards were available only during the training sessions, but not in the pre- or post-training sessions. To assess the spatial learning performance of flies, we first plotted the mean reward index (RI) of the flies over time to quantify their preference for the rewarded circle over a larger control circle (**Fig. 1e**). We found that whereas the RI of yoked controls stayed relatively constant, the RI of the experimental flies increased as the training progressed (**Fig. 1e**), indicating a learned preference for the rewarded location. To visualize flies' learning performance more stringently, we calculated the RI *differences* between each experimental fly and its paired yoked control fly (**Fig. 1f**), aiming to eliminate impacts on RI from non-associative biases induced by repeated reward exposures. As in the case of the RI of the experimental flies, the RI difference between experimental and yoked flies also increased as training progressed (**Fig. 1f**). To simplify the representation of learning performance and its comparisons across genotypes or other manipulations, we introduced the spatial learning index (SLI) – the RI difference between the experimental and yoked flies in the last 10-min sync bucket of training session 3 (see **Methods** for our sync bucket definition). We found that both sighted and blind (*norpA* mutant) flies exhibited a positive SLI in our featureless arenas (**Fig. 1g, h**). Interestingly, learning performance of individual flies did not clearly correlate with their reward preference (Light PI, see **Methods**) and locomotion speed (**Extended Data Fig. 1**). These results suggest flies can learn the goal location in our arenas without relying on any salient visual landmarks.

Fig. 1 Flies can learn a spatial goal without using visual cues. a, Schematic of a featureless flat arena (16mm x 10mm x 3mm) that houses a single fly. During training sessions, the computer would designate a reward circle (d = 2.6mm) at the center of the arena as the goal location. **b,** The high-throughput setup we developed for administering our spatial task. Each setup contains two cameras, one LED board, one infrared back light, and a holder that houses 40 individual arenas. **c,** Our reward rule during the training sessions: only when the fly enters the reward circle from the outside can trigger 1 pulse of light (250ms, 17µW/mm^2, 527nm). **d,** Top: Temporal design of the spatial task. Flies can only receive rewards during the training sessions. Bottom: Positional heatmaps of 20 experimental flies with their 20 yoked control flies at the end of each training session. The fly 'E1' is paired with the fly 'Y1', 'E2' is paired with 'Y2', etc. **e,** The reward index (RI) of experimental flies and their paired yoked control flies (*0273-Gal4>CsChrimson*) at each sync bucket (10 mins, see **Methods)** during the three training sessions. Inset is the schematic of the flat chamber with the reward circle and the control circle (d = 13mm) labeled. Note that we only used males in this study so as to reduce the potential impact of mating experience (virgin vs. mated) and reproductive state (actively egg-laying vs. not) on learning. **f,** The RI difference between experimental and their paired yoked control flies (*0273-Gal4>CsChrimson*) at each syn bucket during training sessions. **g,** Spatial learning index (SLI) of "sighted" (*0273-Gal4>CsChrimson*) flies (n = 83). **h,** SLI of blind (*norpA; 0273-Gal4>CsChrimson*) flies (n = 75). Unless otherwise mentioned, all line plots show means and 95% confidence intervals (shaded), and the numbers above the lines are the sample size at each sync bucket. Error bars in all scatter plots are mean ± S.E.M. One-sample t-test (two-tailed) against 0 was used in **f - h**. Note that throughout the paper *: p<0.05; **: p<0.01; ***: p<0.001; ****: p<0.0001.

Self-motion cues play a role in supporting spatial learning in the absence of prominent visual cues

To begin to identify the potential non-visual cues flies might have used for returning to the unmarked goal location to collect more rewards, we tested the involvement of self-motion cues. Previous studies have suggested that flies can perform path integration[21-23], a strategy where animals rely principally on self-motion cues to calculate the angle and distance (i.e., a vector)

they have traveled from a reference location, and then use the vector to guide them back to the
location. The first indication that flies might have used self-motion cues for locating the goal in our
task came from an examination of animals' trajectories in between rewards (**Fig. 2a**). Specifically,
we calculated the proportion of rewards individual animals collected without touching the arena
walls. Matching our SLI calculation, we then calculated the difference in these proportions
between experimental flies and their paired yoked controls for the last sync bucket of the training
session 3, naming this measure the "no wall contact index." We found that compared to yoked
controls, experimental flies were able to collect a significantly higher proportion of rewards without
touching – and therefore without using – the arena walls, as the mean value was significantly
higher than 0 (**Fig. 2b**).

To further confirm flies' use of self-motion cues for goal locating, we tested the necessity
of proper activities of PFN neurons – a class of spatially selective neurons that comprises two
spatially selective subgroups (PFNd and PFNv) whose populational activities have been shown to
track a fly's egocentric translational velocity[4,5] and to be required for proper local search[5]. We first
silenced the activities of PFNd neurons chronically (by expressing in them the Kir2.1 channel)
and found that this manipulation significantly decreased learning performance, as well as the
proportion of rewards flies collected without wall contacts (**Fig. 2c, d**). Moreover, we also
optogenetically manipulated PFN activity transiently at the goal location via the 250-ms reward
delivery light pulses, as green light is capable of activating both CsChrimson and the light-gated
chloride channel GtACR1[40]. Transiently silencing PFNd neurons, transiently activating them –
which should also disrupt their spatial selectivity – as well as transiently activating PFNv neurons
all reduced learning performance (**Fig. 2e-i**). These manipulations of PFN neurons did not cause
a consistent change in the reward preference or the locomotion speed (**Extended Data Fig. 2**). It
is also worth mentioning that the SLI we use to represent learning should be less sensitive to
unintended consequences of various manipulation, as it indicates the performance *difference*
between an experimental fly and its paired yoked control, both of which underwent the same
manipulation.

We also tested whether EPG neurons may play a role. EPG neurons signal heading
direction and represent a major input into PFN neurons[4,5,35]. Surprisingly, chronically silencing
EPG neurons had no impact on performance (**Fig. 2j, k**). Similarly, transiently silencing or
activating EPG neurons at the goal location did not cause any significant impacts either (**Fig. 2l-**
n). (As in the case of PFNs, we assumed activating the entire EPG population may disrupt its
spatial selectivity.) Thus, the proper heading direction signal represented by the population
activities of EPG neurons appeared unessential for flies to locate the goal in our task, which is
consistent with the finding that EPG neurons are dispensable for local search behavior[41].
Additionally, proper activities of another type of spatially selective neurons that signal allocentric
translational velocity, PFR neurons[4], also appeared to be dispensable (**Extended Data Fig. 3**).
Together, our results suggest that flies use cues that signal egocentric movements but not cues
that signal allocentric orientation or movements to learn our spatial task.

Fig. 2 Flies rely on neurons that signal egocentric translational velocity to learn the task well. a, Left: An in-
between reward trajectory with many wall contacts. Right: An in-between reward trajectory without any wall contact.
b, The no wall contact index of regular flies (*0273-Gal4>CsChrimson*, n = 80). **c,** Expression pattern of the *LexA*
driver line that labels PFNd neurons (*GMR16D01-LexA*). Inset is schematic of PFNd neurons. **d,** Left: SLI of *Ctrl>Kir*
(n = 74) versus *PFNd>Kir* (n = 68) flies. Right: The no wall contact index of *Ctrl>Kir* (n = 74) versus *PFNd>Kir* (n= 66)
flies. Green-filled arena schematic indicates that PFNd neurons were silenced everywhere. Note that throughout this
work, we used *0273-GAL4>CsChrimson* as controls and introduced various transgenes into this background to

manipulate different neurons. Here, *Ctrl>Kir* flies carried the additional transgenes of *empty-lexA* and *lexAop-Kir*
whereas *PFNd>Kir* flies carried the additional transgenes of *16D01-lexA*, and *lexAop-Kir*. **e,** Left: SLI of *Ctrl>GtACR1*
(n = 41) versus *PFNd>GtACR1* (n = 75) flies. Right: The no wall contact index of *Ctrl>GtACR1* (n = 41) versus
PFNd>GtACR1 (n = 74) flies. The green disc in the arena schematic indicates that PFNd neurons were silenced only
when flies received rewards. Note that throughout the paper, we used arenas with either color-filled backgrounds or
center-only circle marks to depict conditions under which neurons of interest were either chronically or transiently
manipulated at the reward circle, respectively. **f,** Confocal image and schematic showing the expression pattern of
the *split-Gal4* driver that labels the PFNd neurons (*SS00078*). **g,** Left: SLI of *Ctrl>CsChrimson* (n = 119) versus
PFNd>CsChrimson (n = 118) flies. Right: The no wall contact index of *Ctrl>CsChrimson* (n = 117) versus
PFNd>CsChrimson (n = 114) flies. **h,** Confocal image and schematic showing the expression pattern of the *split-Gal4*
driver that labels PFNv neurons (*SS52628*). **i,** Left: SLI of *Ctrl>CsChrimson* (n = 98) versus *PFNv>CsChrimson* (n =
119) flies. Right: The no wall contact index of *Ctrl>CsChrimson* (n = 98) versus *PFNv>CsChrimson* (n = 117) flies. **j,**
Confocal image and schematic showing the expression pattern of the *LexA* driver line that labels EPG neurons
(*GMR60D05-LexA*). **k,** SLI of *Ctrl>Kir* (n = 74) versus *EPG>Kir* (n = 72) flies. **l,** SLI of *Ctrl>GtACR1* (n = 81) versus
EPG>GtACR1 (n = 62) flies. **m,** Confocal image and schematic showing the expression pattern of the *split-Gal4*
driver line that labels EPG neurons (*SS00090*). **n,** Left: SLI of *Ctrl>CsChrimson* (n = 119, same as the *Ctrl* for
PFNd>CsChrimson) versus *EPG>CsChrimson* (n = 106) flies. Unless otherwise mentioned, unpaired two-tailed t-test
with Welch's correction was used in all paired comparisons in this paper.
189
Olfaction plays a role in supporting spatial learning in the absence of external odor
landmarks and prominent visual cues

Next, we tested whether flies could use strategies other than ones that depend on self-
motion cues, given that flies with inactivated PFN neurons were still capable of learning, albeit at
a lower level (**Fig. 2c-i**). The flies' antennae are a sensory organ that can detect external non-
visual cues including mechanosensory and olfactory ones. To test if antennae-sensed cues were
used for learning of our task, we first covered both antennae with UV-cured glue. These flies
showed a significant decrease in their learning performance, as did flies with the 3rd segment of
both their antennae surgically removed (**Fig. 3a**), suggesting the importance of one or more of the
antennae-detected cues in enabling learning. To differentiate the relative contributions of
mechanosensory vs. olfactory antennal inputs, we next removed bilaterally just the arista, the
long hair-like structure at the tip of the 3rd antennal segment critical for detecting mechanosensory
but not olfactory cues, and found that these arista-less flies learned normally (**Fig. 3a**). These
results suggest that, surprisingly, olfactory cues appear to be the relevant antennae-detected
cues that promote learning in a task where no external olfactory landmarks were supplied.

To verify that olfactory cues were indeed used for learning, we then tested the
performance of *Orco* mutant flies and found that they had decreased learning performance as
compared to that of controls (**Fig. 3b**). *Orco* mutants are known to be defective in olfaction as the
Orco channel is an obligatory co-receptor for many odorant receptors[42]. The impact of *Orco*
mutation on learning seemed less than that of removing the 3rd segment of the antennae,
possibly because *Orco* is expressed in only ~78% of total olfactory neurons[43]. In a separate
attempt to block the olfactory inputs, we expressed *Kir2.1* in the *Orco*- and the *IR25a*-expressing
neurons simultaneously, which should disrupt the function of more olfactory neurons[43], and also
found a clear decrease in performance (**Fig. 3c**). Again, the olfaction-defective flies we tested did
not show a consistent change in reward preference or locomotion speed (**Extended Data Fig. 4**).

But what might be the source of the odor(s)? Given we did not supply any external odors
and cleaned the arenas thoroughly prior to each experiment, we suspected the odor flies used for
learning might be from deposits they made. To test this idea, we collected samples from the
center of many arenas after the flies had completed the task and sent them for gas
chromatography-mass spectrometry (GC-MS) analysis (**Fig. 3d**). Indeed, several tricosene
isomers, cyclotetracosane, and pentacosene isomers were detected in samples collected from
arenas where flies had undergone training, but not from the clean arenas (**Fig. 3e, Table 1**).

Tricosene and pentacosene are known pheromones for flies[44,45]. These results show that while
we did not provide scents in the arenas, they were not necessarily odorless. To lend further
support to the idea that self-scenting might be important for supporting learning, we attempted to
reduce the transferring of scented substances from flies' bodies onto the arena floor by removing
flies' hind tarsi and covering their genitalia with UV-cured glue. We found that while removal of
hind tarsi did not cause a significant decrease in performance, gluing the genitalia did, and the
decrease was more pronounced when we simultaneously compromised both (**Fig. 3f**). Together
these results suggest that, in addition to using self-motion cues, flies can also use "scent-
marking" as a strategy for returning to the goal location in an environment that lacks prominent
visual and external odor landmarks.

Functional MB is important for learning the spatial goal in the absence of external odor
landmarks and prominent visual cues

Having found that flies deposited potentially scented substances during learning and that
learning depended on a functional olfactory system, we next asked how the scents might be used
for locating the goal. One possibility is that these scents were enriched near the goal such that
flies may find the goal by following scent gradients. Consistent with this idea, we found that the
tendency of flies to make a sharp turn that brings them closer to the goal increased as they were
walking further away from the goal (**Fig. 3g, h**). Importantly, the tendency reduced when we
compromised flies' olfaction or made them less capable of scent marking the arena (**Extended**
Data Fig. 5). Further, we found that naïve flies that had never experienced any trainings did not
show any preference for the goal location when placed in arenas just "scented" by trained flies
(**Extended Data Fig. 6**), suggesting that the deposited scented substances around the goal were
not innately attractive and possibly were afforded positive valence by the presence of reward.

To test the idea that flies could learn to use initially neutral scents from self-deposits for
locating the rewarded goal, we next manipulated the mushroom body (MB), the olfactory
conditioning center where odors can acquire valence by association[3,46]. First, we chemically
ablated the Kenyon cells (KCs), the principal neurons of the MB, by hydroxyurea[47] and found that
flies with a significant portion of their MB ablated showed a substantial decrease in learning
(**Extended Data Fig. 7**). To rule out the possibility that such decrease might be caused by
unintended ablation of brain areas outside of the MB, we also took a genetic approach to
selectively inhibit the KCs. Chronic inhibition of KCs using two different genetic drivers caused a
clear decrease in learning performance in both cases (**Fig. 3i-j**). Interestingly, transiently silencing
KCs specifically at the goal location (again via the 250-ms reward delivery light pulses) also
caused a significant decrease (**Fig. 3k**), in keeping with the notion that flies might be using MB to
associate scents from self-deposits with reward at the goal location. In addition, similar to the
olfaction-defective flies, KCs-silenced flies also showed a decrease in their tendency to make a
sharp turn towards the goal as they moved in a direction away from the goal (**Extended Data Fig.**
8). Lastly, as with the other manipulated flies we have described thus far, flies with non-functional
olfactory learning circuit did not show a consistent change in their preference for reward or
locomotion speed (**Extended Data Fig. 4e-h**). Together, these results suggest that when the goal
is in an environment that lacks prominent external features, flies might mark the rewarded
location with scented deposits, and then use the olfactory learning circuit to stamp the scents with
a positive valence to guide them back to the goal.

Fig. 3 Flies rely on self-deposited chemicals as olfactory landmarks for learning the goal. a, SLI of Ctrl (n = 83, same as flies in Fig1. g), antenna glued (n = 66), 3rd segment of antenna removed (n = 105), and arista removed (n = 56) flies. **b,** SLI of *Orco2/+* (n = 98) versus *Orco1/Orco2* (n = 92) flies. **c,** SLI of control groups *Ir25a & Orco-QF* only (n = 59), *QUAS-Kir* only (n = 88) vs. *Ir25a & Orco-QF>Kir* flies (n = 90). **d,** Steps of collecting deposit samples from the center of the flat chambers. Gray circle area in the center of the flat chamber indicates the transparent film that was used for collecting the odors (see **Methods**). **e,** GC-MS identified several cuticular hydrocarbons that were unique for both experimental and yoked control flies not presented in blank controls (odor samples from empty flat arenas). Triangle mark indicates peak present in the blank control. **f,** SLI of Ctrl (n = 157), hind tarsi removed (n = 146), genitalia glued (n = 103), and hind tarsi removed with genitalia glued simultaneously (n = 104) flies. **g,** Schematic of an analysis that assesses how likely a fly will make a sharp turn (> 90º) within 0.75s arriving at different distances (2 to 6 mm) from the midline of the arena when it moves in a direction away from the goal (see **Methods**). **h,** Comparing the probabilities that such sharp turns were made between the experimental and yoked control flies at different distances during pre-training (Pre, n = 75) and the last sync bucket of training 3 (T3-LSB, n = 76). **i,** Images showing the expression pattern of the *LexA* driver lines we used to label MBKCs. Top: *GMR19B03-LexA*; Bottom: *GMR26E07-LexA*. **j,** Left: SLI of *Ctrl>Kir* (n = 113) versus *MBKC-1>Kir* (n = 102) flies. Right: SLI of *Ctrl>Kir* (n = 74, same as the *Ctrl* for *PFNd>Kir*) versus *MBKC-2>Kir* (n = 66) flies. **k,** Left: SLI of *Ctrl>GtACR1* (n = 82) versus *MBKC-1>GtACR1* (n = 144) flies. Right: SLI of *Ctrl>GtACR1* (n = 82, same as the *Ctrl* for *MBKC-1>GtACR1*) versus *MBKC-2>GtACR1* (n = 153) flies. Welch ANOVA tests with Games-Howell post-hoc test was used for multiple comparisons in panel **a**, **c**, **f.**

Salient external olfactory landmarks can further improve spatial learning performance

Given that flies are likely able to scent mark the goal in an otherwise odorless environment, we next wondered whether flies could also use salient external odor landmarks *away* from the goal to further improve their chances of locating the goal. To answer this question, we enriched the arena by placing strips of 1% agarose into troughs located at the two edges of the arena, away from the goal location (**Fig. 4a, Video S2**) ("agarose chamber"). We found that flies significantly improved their learning performance in the agarose chamber as compared to learning in the flat chamber (**Fig. 4b-f**). Importantly, the enhanced performance was not caused by flies' restricting their search area due to strong innate avoidance of the agarose strips (**Extended Data Fig. 9**). This result suggests flies are able to use salient external landmarks that are consistently "anti-correlated" with the goal to help with locating the goal.

Since agarose can provide both mechanosensory and olfactory cues, we then determined which one of them was responsible for enhancing performance. One observation is consistent with flies using agarose as an olfactory landmark: we found that as the training progressed, the first large turn the flies took towards the goal as they moved away from the goal after receiving a reward, increasingly occurred before physically contacting the agarose (**Extended Data Fig. 10c-d**), suggesting flies might be sensing the agarose from a distance.

To test the idea more directly, we disrupted flies' ability to sense odors by manipulating their antennae, removing their Orco channel, and silencing their olfactory neurons. We found that agarose-induced learning improvement was significantly reduced in all conditions, albeit to different degrees (**Fig. 4g-l, Extended Data Fig. 10a, b**). Inactivating MB severely compromised the agarose-induced performance improvement (**Fig. 4k, l**), too, suggesting MB contributes to learning the goal location by associating agarose odor(s) with value. Further, flies with defective odor-learning systems tended to initiate the first large turn further away from the goal location (when departing from the goal after receipt of a reward) as compared to controls (**Extended Data Fig. 10e-g**), lending support to the notion that flies might have learned to treat the agarose-emitted odors as aversive because they were consistently detected in the absence of reward, thereby limiting the area they needed to search for the goal. Tagging agarose-emitted odors as aversive is in keeping with an earlier finding that odors that are delivered to flies consistently after reward can acquire an aversive quality[48]. Lastly, we found that simultaneously removing flies' hind tarsi and gluing their genitalia no longer significantly impacted learning (**Fig. 4j**), which is quite different from the large decrease in learning performance observed in the flat chamber, indicating that presence of salient external olfactory landmarks can "rescue" flies' learning performance when they were less capable of depositing scents near the goal.

Flies can flexibly adjust their reliance on self-motion cues for learning depending on the availability of external olfactory landmarks in the environment

We have so far shown that flies were able to use both self-motion cues and odor cues from self-deposits for locating the goal in a relatively featureless arena, and to improve their performance when the arena became enriched with external olfactory landmarks. Next, we were curious if flies might adjust their strategies in these two types of arenas where the available spatial cues differ. To begin, we ascertained the parallel nature of the self-motion cues- and olfactory cues-based strategies for locating the goal in the flat chamber by asking whether flies with disabled PFN neurons showed further decreased performance upon their antennae being blocked, and vice versa. Indeed, we found that inhibition of PFN caused a further learning decrease in antennae-glued flies (**Fig. 5a**). Similarly, gluing the antennae caused a further learning decrease in flies with inhibited PFN neurons (**Fig. 5b**). When removing the antennae instead of gluing them, similar results were also found in antennae-removed and PFNd-silenced flies (**Extended Data Fig. 11a, b**). These results support the parallel nature and flies' concurrent adoption of self-motion cue-based and olfactory landmark-based strategies for locating the spatial goal in the "featureless" flat chamber.

Do flies alter their strategies when tasked to learn in environments that contain more useful cues? We found that flies' relative reliance on the olfactory vs. self-motion cues changed in our agarose chamber where the arena was enriched with agarose strips away from the goal. Whereas the olfactory learning circuit remained critical as silencing its components still caused a significant reduction in learning (**Fig. 4g-l**), the contribution of PFN neurons diminished in the agarose chamber, as silencing them caused less of an impact in the agarose chamber compared to the flat chamber (**Fig. 5c, d**). However, PFN neurons can "gain" back their critical role in the agarose chamber when we blocked flies' olfactory input, as inhibiting them caused a more substantial impact in antennae-compromised compared to non-compromised flies (**Fig. 5e, Extended Data Fig. 11c**). These results suggest that the contribution of the self-motion cues-based learning strategy is context-dependent: when the environment is featureless, such strategy plays a more important role and works in parallel with a scent-marking-based strategy mediated by MB; but when the environment is enriched with salient external olfactory landmarks, self-motion cues become more disposable. Thus, flies are able to adopt different strategies for locating goals in visually challenging conditions, depending on the availability of quality non-visual cues.

Fig. 5 Flies can flexibly adjust their strategies for learning the spatial goal depending on the availability of landmarks in the environment. a, SLI of *PFNd>Kir* (n = 68) versus *PFNd>Kir* antenna glued (n = 109) flies in the flat chamber. **b,** SLI of *Ctrl>Kir* antenna glued (n = 106) versus *PFNd>Kir* antenna glued (n = 109) flies in the flat chamber. **c,** Left two columns: SLI of *Ctrl>Kir* (n = 74) versus *PFNd>Kir* (n = 68) flies in the flat chamber. Right two columns: SLI of *Ctrl>Kir* (n = 110) versus *PFNd>Kir* (n = 99) flies in the agarose chamber. Two-way ANOVA was used to calculate the interaction between the effect of silencing PFNd neurons and the chamber type. Šídák's multiple comparisons test was used. **d,** Left two columns: SLI of *Ctrl>GtACR1* (n = 41) versus *PFNd>GtACR1* (n = 75) flies in the flat chamber. Right two columns: SLI of *Ctrl>GtACR1* (n = 56) versus *PFNd>GtACR1* (n = 68) flies in the agarose chamber. Two-way ANOVA was used to calculate the interaction between silencing PFNd neurons and chamber type on SLI. Šídák's multiple comparisons test was used. **e,** Left two columns: SLI of *Ctrl>Kir* (n = 110) versus *PFNd>Kir* (n = 99) flies in the agarose chamber. Right two columns: *Ctrl>Kir* antenna glued (n = 65), and *PFNd>Kir* antenna glued (n = 73) flies in the agarose chamber. Two-way ANOVA was used to calculate the interaction between silencing PFNd neurons and antenna gluing on SLI. Šídák's multiple comparisons test was used. **f,** Our model of how the flies learn a goal location in a featureless environment (left) and in an environment that is enriched with olfactory landmarks (right). The black arrow indicates the integrated path (IP) relative to the goal based on self-motion cues. The green area near the goal indicates the potential scent deposited area, which was assigned a positive value by MB. The red areas indicate the agarose strips, which were assigned a negative value by MB.

Discussion

In this study, we examined the behavioral and neural strategies that flies use to return to a goal location without relying on prominent visual landmarks. Using a high-throughput setup we developed where flies can forage for an optogenetically delivered virtual reward at an unmarked location, our results suggest that when the foraging arena is essentially featureless, flies can learn to use scents from self-deposits along with self-motion information to locate the goal.

When the arena is enriched with external salient olfactory landmarks situated away from the
goal, however, flies switch strategies and principally learn to use odor landmarks that signal the
presence and absence of reward to restrict their search area (**Fig. 5f**). These results highlight
the flexibility and ability of flies to solve a spatial task in a visually challenging environment:
when possible, they can transform the spatial task into primarily an odor association task.

Some key pieces of evidence supporting our claim came from our manipulations of the
odor association center, the mushroom body (MB). Ablating the MB, as well as inhibiting it,
either chronically or transiently, disrupted flies' spatial learning performance. These results,
combined with the finding that olfactory neurons also played a critical role, suggest that the MB
aids learning by assigning value to odor landmarks. Specifically, we propose that in our
featureless flat arena, the MB works by assigning a positive value (appetitiveness) to self-
generated scents near or at the goal. Additionally, in our agarose arena, the MB assigns a
negative value (aversiveness) to odor from agarose located away from the goal, allowing flies to
restrict their search areas to regions with low or no such odor (**Fig. 5f**). Such negative value
assignment to "reward anti-correlated cues" aligns with earlier findings that flies can learn to
treat odors that are consistently experienced following rewards as aversive[48]. Could the MB
work in some capacity other than its conventional association role in supporting learning? It is
conceivable that the MB might facilitate using odor landmarks to reset or correct the path
integrator, especially given that the output neurons of MB (MBON) are known to signal the
central complex (CX)[49,50], where many classes of spatially selective neurons, including PFNs,
are housed. However, our results so far favor some amount of parallel nature of MB-based and
PFN-based strategies, as olfactory input can still promote some learning without functional
PFNs and vice versa.

One of the most unexpected findings of our work is that flies appeared to use scents
from self-deposits to "create" landmarks – and then assigned them meaning – when locating a
rewarded goal in a feature-impoverished environment. This result mirrors a recent finding
showing that dessert ants can create their own visual landmarks when navigating in
environments without obvious ones[51]. While we were not able to directly visualize the scents'
being concentrated at the goal location, our thesis is supported by several lines of evidence.
First, we detected various fly-derived chemicals on the arena floors. Second, preventing flies
from rubbing their bodies, especially their genitalia, reduced learning. Third, disabling olfactory
neurons disrupted learning. Lastly, silencing the olfactory association center, MB, disrupted
learning as well. Our findings raise two interesting questions. First, what is the exact identity of
the substances being used? Some of the chemicals identified by GS-MS are likely candidates,
although it is also possible that the relevant ones were present below the detection threshold.
We cannot provide proof of the use for any of the chemicals for now, partly because the
receptors for many of these chemicals are not known. It is worth noting that *Drosophila* males
have been shown to mark locations where they encounter appetitive food odors with 9-tricosene
to attract egg-laying females[27]. However, 9-tricosene was not the relevant scent used by our
males, or at least not the sole scent, as the mutants for Or7a, the receptor for 9-tricosene, did
not show reduced performance compared to controls (**Extended Data Fig. 12**). Second, how
general might this strategy be? For example, will flies abandon this strategy or use a different
scent depending on the nature of the reward and/or the foraging environment? Curiously,
olfaction was found to be dispensable in a landmark study where the authors demonstrated that
flies can use path integration when navigating towards a goal in darkness[22]. One important
difference between the task described by Kim & Dickinson and ours is that our arena size is
considerably smaller. It is possible that, given the potential low quantity and/or volatility of self-
deposited scents, the strategy we described may be effective only in small and enclosed
environments. As for whether the nature of the reward matters, we found that intact antennae
were still required when we replaced optogenetic activation of *0273-Gal4*-labeled neurons with

activation of another group of neurons (the fructose sensing *Gr43a* neurons), which has also
been shown to promote reward seeking[21] (**Extended Data Fig. 13**). This suggests that the
strategy we observed may generalize to tasks where the goal location is baited with a (virtual)
sugar reward.

Our work suggests that self-motion cues also play a role in aiding flies' returns to the
goal location when the environment lacks prominent visual cues. We showed that flies were
able to collect many rewards without contacting the arena walls. More importantly, the proper
function of neurons known to signal egocentric translational velocity (i.e., PFNs) was critical for
full performance, as well as collecting many rewards without wall contacts. How exactly the flies
use self-motion cues conveyed by PFNs to solve the task remains unclear for now. Intriguingly,
disrupting the function of one target two synapses downstream of PFN neurons, the PFR
neurons, which allow flies to map their egocentric movements onto the allocentric/world frame
(by integrating translational velocity signals from PFNs and heading signals from EPG
neurons)[4], did not produce obvious defects in spatial learning. Similarly, disrupting the function
of the EPG neurons did not significantly impact performance either. The lack of requirement for
EPG and PFR neurons might be because unlike learning task driven by prominent visual cues[52],
our arenas lack any prominent external cues that can engage them reliably. Nonetheless, these
findings raise two questions. First, since EPGs have been shown to provide critical heading
input to PFNs[4,5], it is curious what might be supplying PFNs with directional signals when we
silenced EPGs. One possible source is the noduli[4,5]. Second, since PFRs were dispensable,
too, what might the relevant downstream targets of PFNs be in allowing self-motion to be used
to guide goal-directed movements? We don't have answers for this either, but the recently
identified PFL3 and PFL2 neurons that allow flies to steer their movements according to an
internal goal heading conveyed in FC2 neurons, and hΔC/K neurons that signal goal direction
gated by odor, might be worth testing[34,36,53], as are other anatomical targets delineated in the
connectome.
471
Methods

Fly genotypes

1. To perform the SLI and other behavioral analysis of sighted control flies and flies with
manipulations in antennae (whole antennae glued third segment of antennae removed and
arista removed), hind tarsi removal and gluing genitalia (Fig. 1d-g; 2a, b; 3a, e, f, h; 4b-d, g, j;
Extended Data Fig. 1a, b, d, f; 4a, d; 5a-d, g-i; 9b; 10a, d, e; Table 1): *UAS-CsChrimson/y; +/+;*
0273-Gal4/+.

2. To perform the SLI and other behavioral analysis of blind flies (Fig. 1h; 4e, f; Extended Data
Fig. 1e, 6a, b): *norpA; +; UAS-CsChrimson/0273-Gal4*.

3. To chronically silence target neurons with Kir, the control group flies (Fig. 2d, k; 3j; 4k; 5a, c,
e; Extended Data Fig. 2a; 3b; 4e, f; 8a; 11b, c): *UAS-CsChrimson/y; Empty-LexA/otd-flp; 0273-*
Gal4/LexAop-(FRT)-tdTomato-STOP-(FRT)-eGFP-Kir.

4. To chronically silence PFNd neurons (Fig. 2d; 5a-c, e; Extended Data Fig. 2a; 11 a-c): *UAS-*
CsChrimson/y; 16D01-LexA/otd-flp; 0273-Gal4/LexAop-(FRT)-tdTomato-STOP-(FRT)-eGFP-
Kir.

5. To chronically silence EPG neurons (Fig. 2k): *UAS-CsChrimson/y; 60D05-LexA/otd-flp; 0273-*
Gal4/LexAop-(FRT)-tdTomato-STOP-(FRT)-eGFP-Kir.

6. To chronically silence PFR neurons (Extended Data Fig. 3b): *UAS-CsChrimson/y; 37G12-*
LexA/otd-flp; 0273-Gal4/LexAop-(FRT)-tdTomato-STOP-(FRT)-eGFP-Kir.

7. To chronically silence MBKCs (Fig. 3j; 4k; Extended Data Fig. 4e, f, 8a): *UAS-CsChrimson/y;*
19B03-LexA/otd-flp; 0273-Gal4//LexAop-(FRT)-tdTomato-STOP-(FRT)-eGFP-Kir, and *UAS-*
CsChrimson/y; 26E07-LexA/otd-flp; 0273-Gal4/LexAop-(FRT)-tdTomato-STOP-(FRT)-eGFP-
Kir.

8. To transiently silence target neurons with GtACR1, the control group flies (Fig. 2e, l; 3k; 4l;
5d; Extended Data Fig. 2b; 3c; 4g, h; 8b): *w; ctrl-lexA/lexAop-GtACR1; 0273-Gal4/UAS-*
CsChrimson.

9. To transiently silence PFNd neurons (Fig. 2e; 5d; Extended Data Fig. 2b): *w; 16D01-*
lexA/lexAop-GtACR1; 0273-Gal4/UAS-CsChrimson.

10. To transiently silence EPG neurons (Fig. 2l): *w; 60D05-lexA/lexAop-GtACR1; 0273-*
Gal4/UAS-CsChrimson.

11. To transiently silence PFR neurons (Extended Data Fig. 3c): *w; 37G12-lexA/lexAop-*
GtACR1; 0273-Gal4/UAS-CsChrimson.

12. To transiently silence MBKCs (Fig. 3k; 4l; Extended Data Fig. 4g, h; 8b): *w; 19B03-*
lexA/lexAop-GtACR1; 0273-Gal4/UAS-CsChrimson, and *w; 26E07-lexA/lexAop-GtACR1*; *0273-*
Gal4/UAS-CsChrimson.

13. To transiently activate target neurons, the control group flies (Fig. 2g, i, n; Extended Data
Fig. 2c, d; 3e): *w; UAS-CsC/empty-Gal4-AD; 0273-Gal4/empty-Gal4-DBD*.

14. To transiently activate PFNd neurons (Fig. 2g; Extended Data Fig. 2c): *w; UAS-*
CsChrimson/SS00078; 0273-Gal4/SS00078.

15. To transiently activate PFNv neurons (Fig. 2i; Extended Data Fig. 2d): *w; UAS-*
CsChrimson/SS52628; 0273-Gal4/SS52628.

16. To transiently activate EPG neurons (Fig. 2n): *w; UAS-CsChrimson/SS00090; 0273-*
Gal4/SS00090.

17. To transiently activate PFR neurons (Extended Data Fig. 3e): *w; UAS-*
CsChrimson/SS54549; 0273-Gal4/SS54549.

18. To test the roles of Orco in learning (Fig. 3b; 4h; Extended Data Fig. 4b; 5e; 10f), we used
w; UAS-CsChrimson/+; 0273-Gal4/Orco23130 as the control group, and *w; UAS-CsChrimson/+;*
0273-Gal4, Orco23129/Orco23130 as the experimental group.

19.To chronically silence *Ir25a*-expressing and *Orco*-experssing neurons simultaneously (Fig.
3c; 4i; Extended Data Fig. 4c; 5f; 10b, g), we used *w; UAS-CsChrimson/Ir25a-QF; Orco-QF (BL-*
92400)/0273-Gal4 as control group #1*, w; UAS-CsChrimson/+; 0273-Gal4, QUAS-Kir/+* as
control group #2*, and w; UAS-CsChrimson/Ir25a-QF; 0273-Gal4, QUAS-Kir/Orco-QF* as the
experimental group.

20. To chemically ablate MB (Extended Data Fig. 7): *UAS-CsChrimson/+; 0273-Gal4/+.*

21, To test the roles of Or7a in learning (Extended Data Fig. 12), the control group flies: *UAS-*
CsChrimson/+; 0273-Gal4/+, and flies with *Or7a* mutant: *Or7a/y; UAS-CsChrimson/+; 0273-*
Gal4/+.

22: To test the learning performance of starved flies with optogenetic activation of *Gr43a*
neurons as reward (Extended Data Fig. 13): *UAS-CsChrimson/y; Gr43a-Gal4/+ (II).*

23: To image the expression pattern of driver lines that label PFNd neurons (Fig. 2c, f): *w;*
R16D01-lexA/lexAop-GtACR1; 0273-Gal4, and *w; +/SS00078; UAS-eGFP-Kir/SS00078.*

24: To image the expression pattern of the driver line that labels PFNv neurons (Fig. 2h): *w;*
+/SS52628; UAS-eGFP-Kir/SS52628.

25: To image the expression pattern of driver lines that label EPG neurons (Fig. 2j, m): *w;*
R60D05-lexA/lexAop-GtACR1; 0273-Gal4, and *w; +/SS00090; UAS-eGFP-Kir/SS00090.*

26: To image the expression pattern of driver lines that label MBKCs (Fig. 3i): *w; R19B03-*
lexA/lexAop-GtACR1; 0273-Gal4, and *w; R26E07-lexA/lexAop-GtACR1; 0273-Gal4.*

27: To image the expression pattern of driver lines that label PFR neurons (Extended Data Fig.
3a, d): *w; R37G12-lexA/lexAop-GtACR1; 0273-Gal4*, and *w; +/SS54549; UAS-eGFP-*
Kir/SS54549.

Note that we used the same group of control flies in the following experiments since they were
done at the same time periods:

Ctrl>Kir flies were the same for *PFNd>Kir* and *MBKC-2>Kir* in the flat chamber;

Ctrl>Kir flies were the same for *EPG>Kir* and *PFR>Kir* in the flat chamber;

Ctrl>GtACR1 flies were the same for *MBKC-1>GtACR1* and *MBKC-2>GtACR1* in both the
flat and the agarose chambers;

Ctrl>CsChrimson flies were the same for *PFNd>CsChrimson* and *EPG>CsChrimson.*

549
Fly husbandry

Flies were raised at 25 °C on standard molasses food. We density-controlled the crosses and
supplied them with yeast paste (6 g dry yeast in 10mL 0.5% propionic acid) to ensure the health
of the progeny. Only 10- to 14-day old males were used for our spatial learning tasks. They
were collected on the first day of eclosion into regular food vials for 6-10 days at the density of
12-14 flies/vial, transferred into food vials with 400 µM retinal and kept without light exposure for
2 days, and transferred again into fresh retinal food vials under circadian entrainment (12 hours
light on/off) before being assayed. For *Gr43a-Gal4>CsChrimson* flies, they were starved for 20
hours before the spatial learning task.

559
Non-genetic manipulations on flies

Removing the 3rd segment of antenna, and the arista. The procedures were done using forceps
one day before the assay to give the flies time to recover and adapt.

Gluing the whole antenna. This was done by applying a small piece of UV-cured glue (Bondic)
to cover the entire antenna of the flies under a dissecting scope and exposing them to 5
seconds of UV light (Darkbeam UV Flashlight 365 nm). To allow the flies time to recover and
adapt, these manipulations were performed one day before the assay.

Gluing the genitalia. This was done by applying a small piece of UV-cured glue (Bondic) to
cover the whole genitalia of the flies under a dissecting scope and exposing them to 5 seconds
of UV light (Darkbeam UV Flashlight 365 nm). To limit activation of *0273-GAL4*-labeled neurons
during these manipulations by the illuminator light (Nikon NI-150) of the dissecting scope, we
set its intensity below 30% and restricted the total illumination time to under 3 minutes. Control
flies, which were not glued, were illuminated under the dissecting scope, anesthetized, and
exposed to UV for the same duration as the glued group. As the glue on the genitalia tended to
fall off, we applied it 2 to 3 hours before the task and always inspected for its presence
afterward to ensure it remained intact during the task.

Removing the hind tarsi. This was done using a pair of forceps. Since we used the same control
flies as for the genitalia-glued group, we removed the hind tarsi of the flies 2 to 3 hours before
the task.

MB ablation. MB in the fly brain were chemically ablated following the standard procedure[54] with
the modification that 0- to 1-hour old larvae were fed for 5 hours with yeast paste that contained
50 mg/ml hydroxyurea.
582
Odor sample collection and GC-MS

To collect substances deposited by the flies, we placed a small piece of transparent film (3.5
mm in diameter) at the center of each arena prior to the task and collected it immediately after
training finished. Specifically, 18 pieces from arenas trained with experimental flies, 18 from
arenas trained with yoked controls, and 18 from empty arenas (blank control) were collected,
soaked in three separate vials with 100 µl of hexane each, and sent for gas chromatography-
mass spectrometry (GC-MS) analysis at Alera Labs (https://aleralabs.com/). At Alera Labs,
samples were evaporated and redissolved in 100 µL of hexane containing the internal standard
(methyl nonadecanoate at 5 µg/mL). Then, 3 µL of each sample were analyzed using an Rxi-
5MS column (30 m length, 0.25 mm ID, 0.25 um film thickness) with an inlet temperature of
280 °C and a detector temperature of 300 °C. Helium was used as the carrier gas with a split
10:1 flow rate. The temperature program was set to start at 120 °C, ramping at 6 °C/min to
300 °C, with a 5-minute hold. The mass spectrometer operated in electron ionization (EI) mode
with an m/z scan range of 35-550. All assignments were based on the highest probability
spectral matches.
598
Implementation of the high-throughput spatial learning task

Hardware. *Arenas.* We designed two types of chambers, each containing 40 individual arenas,
for testing flies' ability to locate the spatial goal. To increase the throughput of the task, we have
created 6 setups, enabling us to simultaneously assay 240 flies in parallel. In the flat chambers,
the arena is rectangularly shaped (dimensions: 16 x 10 x 3 mm) and featureless (**Fig. 1a**). In the
agarose chamber, the arena has the same dimensions but includes two troughs at the ends (3 x
10 x 3 mm) that we filled with 1% agarose (**Fig. 4a**). *Optogenetic stimulation.* Light pulses were
delivered by custom LED controller boards with 5-mm 527-nm 15°-viewing angle Cree LEDs
(C503B-GAN-CB0F0791). Board schematic and layout are available at
https://github.com/ulrichstern/SkinnerTrax. We used 25% light intensity for the 250-ms "reward"
pulses for *0273-Gal4>CsChrimson* flies, which corresponds to about 16.4 µW/mm^2 at the arena
floors, while for *Gr43a-Gal4>CsChrimson* flies (Extended Data Fig. 13), we used 125% light
intensity, which corresponding to about 77.3 µW/mm^2 at the arena floors. *Cameras.* Each
apparatus uses two cameras (Microsoft LifeCam Cinema) with their internal IR-block filters
removed and external IR-pass filters (LEE Filters 87) added. *Infrared backlight.* An 850-nm light
pad was placed underneath the arenas and LED controller board to provide background lighting
for tracking. The light pads were custom made by Junlong Display Factory; we asked them to
replace the white LEDs in their Artograph A920 series light pads with 850-nm LEDs.

Software. SkinnerTrax[37] was used to administer the learning task independently for each of the
40 flies per apparatus in parallel. SkinnerTrax's experimental records, which served as input for
follow-on analysis, included the flies' positions, fitted ellipses for each frame, and the times
when each fly was stimulated. The code for SkinnerTrax is available at
https://github.com/ulrichstern/SkinnerTrax.

Handling of flies. Male flies, 10-14 days old and fed standard 400 µM retinal fly food for 4 days,
were maintained under circadian control (12 h light on, 12 h light off) and placed in either flat or

agarose chambers (one fly per arena) within the first 2.5 hours of light onset. The chambers
were placed into the tracking system and shielded from ambient light with a cardboard box.
626
627 Analysis of the high-throughput spatial learning task

All the behavior analysis code used in this paper are available in GitHub at
https://github.com/rcalfredson/nvsl-analysis

Exclusion of flies with poor tracking. To minimize the impact of tracking problems, our analysis
code automatically excluded trajectories for which problems were detected. First, the
percentage of "lost" frames – where the fly could not be detected after the initial onset of
tracking – was calculated. The vast majority of lost frames occurred when flies were walking on
the sidewalls, which appear darker in the frames, making it harder to track flies. Trajectories that
contained more than 10% lost frames were excluded from analysis. Second, tracking may
intermittently switch between the fly (the intended target) and fixed arena locations where
tracking artifacts occur, especially if the fly gets lost. The result is "suspicious jumps" where a fly
incorrectly appears to jump from its true location to the artifact and then back to about the
original location when it is tracked again. If at least 3% of the jumps in a trajectory were
determined to be suspicious, and there were at least three suspicious jumps, the trajectory was
excluded from analysis.

Calculation of the heatmaps. The heatmaps shown in Fig. 1d and 4b were calculated directly by
SkinnerTrax to allow for a quick assessment of the positional preferences of the flies during the
training sessions. Warmer colors indicate more time spent at that position.

Calculation of the reward index (RI). The reward index, which we first introduced in prior work[39],
is defined as follows:

647
$$RI = \frac{\#\ entries\ into\ reward\ circle\ -\ \#\ entries\ into\ control\ circle}{total\ \#\ entries\ into\ the\ two\ circles}$$

The reward and control circles we used are depicted in Fig. 1e.

The time point 0 in the RI over time (and RI difference over time) plots during training
was defined differently from the actual start time of the training session. For each training
session, time 0 marks when the experimental fly first entered the reward circle, receiving a
reward, rather than the start time of the session. This means we excluded the initial part of the
session where the fly had no learning signal yet, and thus the time 0 in the plot can be
considered the session's beginning of learning for each fly. We used the term "sync bucket" to
refer to the synchronization of temporal buckets (intervals) with the first reward of the session.
For example, to calculate the RI for the fifth (last) 10-minute sync bucket, we counted all circle
entries in the 10-minute interval from time point 40 to time point 50 minutes and then applied the
reward index formula.
As we have previously noted[39], the reward index was designed to be insensitive to flies'
speed. This was achieved in two ways. First and foremost, by comparing preference of entering
one circle over another, it is not the absolute number of circle entries that matters (where a fast
animal would have an advantage) but the preference index (where a slow animal that learns
well can be as good as or better than a fast one). Second, to be able to calculate each
preference value reliably, we excluded data points for buckets where the sum of the entries for
the two circles was less than 10. This rule automatically excluded times when flies barely
moved, and a fly that barely moved throughout the experiment would not contribute to any of the
mean reward index scores.

Calculation of the spatial learning index (SLI). To represent the spatial learning ability of a given
group of flies during training, we first calculated the reward index difference between each
experimental fly and its paired yoked control at each sync bucket. See main text for the benefits
of using this difference. The SLI is simply this difference for the last sync bucket of the training
session 3:

$$SLI \ = \ (RI_{experimental \ fly} - RI_{yoked \ control}) \ \text{for the last 10-min sync bucket of T3.}$$

Calculation of light/reward preference. We performed open-loop light preference assays in the
flat chambers to test flies' innate preference for the optogenetically delivered rewards. The
open-loop task comprised a 30-minute pre-training period, a 2-hour training session, and a 30-
minute post-training session. During the training session, only the upper half of the chamber
was illuminated in a repeating cycle of 4 minutes of light on (7% LED intensity, which
corresponds to about 4.63 µW/mm^2 at the arena floors) followed by 4 minutes of light off. We
calculated the light preference as average over all light-on periods as follows:

$$Light \ PI = \frac{Time_{illumated \ side} - Time_{dark \ side}}{Total \ illumination \ time}$$

682
Additional behavioral analyses

Calculation of the no wall contact index. See main text for the calculation of the no wall contact
index. To ensure the index could be calculated reliably, we excluded pairs of flies unless the
experimental fly and its paired yoked control each collected at least 10 rewards during the
period used for the calculation.

Wall contact detection. Wall contact events were determined using a dual-boundary approach,
in which two boundaries were defined for each wall: an event-start boundary 0.7 mm inward
from the wall and an event-end boundary 1.0 mm inward from the wall. A wall contact event
began when the point on the fly's fitted ellipse closest to the wall crossed the event-start
boundary walking toward the wall. The event continued until the point on the ellipse closest to
the wall crossed the event-end boundary walking away from the wall. The boundary offsets were
selected from four options (0.4/0.75 mm, 0.6/0.9 mm, 0.7/1.0 mm and 0.9/1.2 mm) based on
which provided the best agreement between algorithmically detected and manually verified
events.

Calculation of the probability of sharp turns toward the horizontal centerline at different
distances. We aimed to identify sharp turns in the fly's trajectory at virtual boundaries located
symmetrically at distances 2, 3, 4, 5, and 6 mm from the horizontal centerline of the chamber
(Fig. 3g, h, Extended Data Fig. 5, 8). Each distance was analyzed independently, and analysis
was typically performed for the last sync bucket of training 3.

Detection of virtual boundary contact events. Boundary contact events began when the fly's
trajectory (which is based on fitted ellipse centers) crossed a virtual boundary while moving
away from the centerline. The events ended when the trajectory crossed a horizontal line
positioned 0.1 mm inward from the virtual boundary.

Detection of sharp turns. Sharp turns were defined as boundary contact events with a
duration of 0.75 seconds or less and with a cumulative change in velocity angle (the angle of
the velocity vector) in a single direction in excess of 90 degrees. When calculating the
cumulative change in velocity angle, we excluded trajectory edges where the fly's speed
was below 2 mm/s as well as angle changes for the first frame of a wall contact event.

Probability calculation. The probability of sharp turns was calculated as the ratio of the
number of sharp turns to the total number of boundary contact events.

<u>Exclusion criteria and sample size standardization for plotting.</u> To ensure only reliably
calculated probabilities were used, at least 10 boundary contact events were required during
the period used for the calculation. If a fly was excluded at one distance, it was excluded at
all distances, resulting in the same sample size for all distances (e.g., Fig. 3h). Finally,
except for the pre-training analysis in Fig. 3h, we included only flies for which an SLI had
been calculated.

<u>Calculation of the percentage of time spent on agarose (or corresponding) areas.</u> The
percentage of time flies spent on agarose was calculated based on the proportion of frames in
which the ellipse center fell within the rectangles corresponding to the agarose areas (Extended
Data Fig. 9a). For percentage calculations for the last sync bucket of training 3, we included
only flies for which an SLI had been calculated.

<u>Calculation of the distance distribution of the start of the first large turn.</u> We aimed to determine
the distribution of the distance from the chamber center where the flies initiated their first large
turn after their departures from the reward circle (Extended Data Fig. 10c-g).

<u>Detection of large turns.</u> We used the fly's heading angle instead of velocity angle to detect
large turns since the former resulted in fewer false positives, and we skipped frames with
speed below 2 mm/s to exclude trajectory segments for which the flies were resting. We
defined large turns to start with an angle delta of at least 18° and to reach a cumulative
angle delta of at least 90° in the same direction (left or right) as the turn-starting angle delta.
If an angle delta of at least 18° in the opposite direction occurred before reaching 90°, we
started a second angle delta sum to allow for large turns in either direction. Finally, we
excluded turns where 1) wall contact or re-entry into the reward circle happened before
reaching 90°, 2) the total distance traveled during the turn was less than 1 mm, or 3) more
than 70% of the frames of the turn had a speed below 2 mm/s.

<u>Calculation of the distance of the start of the first large turn histograms.</u> Each fly's distance
data was binned into its own histogram. Histograms based on fewer than 20 large turns
were excluded. Extended Data Fig. 10d-g shows mean histograms, averaging the bins
across all flies within a group. Two-tailed Welch's t-test with Bonferroni correction was used
to compare groups for each bin. In addition, we calculated the mean distance for each fly
and compared the means using Two-tailed Welch's t-test.

<u>Calculation of the average speed during learning.</u> To calculate the average speed during
learning (Extended Data Figs. 1a, b, 2, and 4), we used only the last sync bucket of training 3,
excluding sidewall walking to improve the accuracy of the speed calculation. We included only
flies for which an SLI had been calculated.
747

References

1   Lee, S. M., Shin, J. & Lee, I. Significance of visual scene‐based learning in the hippocampal
systems across mammalian species. *Hippocampus* **33**, 505-521 (2023).
2   Ofstad, T. A., Zuker, C. S. & Reiser, M. B. Visual place learning in Drosophila melanogaster.
Nature **474**, 204-207 (2011).
3   Modi, M. N., Shuai, Y. & Turner, G. C. The Drosophila mushroom body: from architecture to
algorithm in a learning circuit. *Annual review of neuroscience* **43**, 465-484 (2020).
4   Lyu, C., Abbott, L. & Maimon, G. Building an allocentric travelling direction signal via vector
computation. *Nature* **601**, 92-97 (2022).
5   Lu, J. *et al.* Transforming representations of movement from body-to world-centric space.
Nature **601**, 98-104 (2022).

6   Müller, M. & Wehner, R. Path integration in desert ants, Cataglyphis fortis. *Proceedings of the*
National Academy of Sciences **85**, 5287-5290 (1988).

7   Heinze, S., Narendra, A. & Cheung, A. Principles of insect path integration. *Current Biology* **28**,
R1043-R1058 (2018).

8   Wittlinger, M., Wehner, R. d. & Wolf, H. The ant odometer: stepping on stilts and stumps.
science **312**, 1965-1967 (2006).

9   Alyan, S. & McNaughton, B. Hippocampectomized rats are capable of homing by path
integration. *Behavioral neuroscience* **113**, 19 (1999).

10   Alyan, S. & Jander, R. Short-range homing in the house mouse, Mus musculus: stages in the
learning of directions. *Animal Behaviour* **48**, 285-298 (1994).

11   Aharon, G., Sadot, M. & Yovel, Y. Bats use path integration rather than acoustic flow to assess
flight distance along flyways. *Current Biology* **27**, 3650-3657. e3653 (2017).

12   Séguinot, V., Cattet, J. & Benhamou, S. Path integration in dogs. *Animal behaviour* **55**, 787-797
(1998).

13   Patel, R. N., Kempenaers, J. & Heinze, S. Vector navigation in walking bumblebees. *Current*
Biology **32**, 2871-2883. e2874 (2022).

14   Srinivasan, M. V., Zhang, S. & Bidwell, N. Visually mediated odometry in honeybees. *Journal of*
Experimental Biology **200**, 2513-2522 (1997).

15   Srinivasan, M. V., Zhang, S., Lehrer, M. & Collett, T. Honeybee navigation en route to the goal:
visual flight control and odometry. *Journal of Experimental Biology* **199**, 237-244 (1996).

16   Frisch, K. v. *The dance language and orientation of bees*. (Harvard University Press, 1993).

17   El Jundi, B. *et al.* A snapshot-based mechanism for celestial orientation. *Current biology* **26**,
1456-1462 (2016).

18   el Jundi, B., Smolka, J., Baird, E., Byrne, M. J. & Dacke, M. Diurnal dung beetles use the intensity
gradient and the polarization pattern of the sky for orientation. *Journal of experimental biology*
217, 2422-2429 (2014).

19   Yilmaz, A., Gagnon, Y., Byrne, M., Baird, E. & Dacke, M. Cold-induced anesthesia impairs path
integration memory in dung beetles. *Current Biology* **32**, 438-444. e433 (2022).

20   Giraldo, Y. M. *et al.* Sun navigation requires compass neurons in Drosophila. *Current Biology* **28**,
2845-2852. e2844 (2018).

21   Corfas, R. A., Sharma, T. & Dickinson, M. H. Diverse food-sensing neurons trigger idiothetic local
search in Drosophila. *Current Biology* **29**, 1660-1668. e1664 (2019).

22   Kim, I. S. & Dickinson, M. H. Idiothetic path integration in the fruit fly Drosophila melanogaster.
Current Biology **27**, 2227-2238. e2223 (2017).

23   Behbahani, A. H., Palmer, E. H., Corfas, R. A. & Dickinson, M. H. Drosophila re-zero their path
integrator at the center of a fictive food patch. *Current Biology* **31**, 4534-4546. e4535 (2021).

24   Hulse, B. K. & Jayaraman, V. Mechanisms underlying the neural computation of head direction.
Annual review of neuroscience **43**, 31-54 (2020).

25   Marin, A. C., Schaefer, A. T. & Ackels, T. Spatial information from the odour environment in
mammalian olfaction. *Cell and Tissue Research* **383**, 473-483 (2021).

26   Jacobs, L. F., Arter, J., Cook, A. & Sulloway, F. J. Olfactory orientation and navigation in humans.
PLoS One **10**, e0129387 (2015).

27   Lin, C.-C., Prokop-Prigge, K. A., Preti, G. & Potter, C. J. Food odors trigger Drosophila males to
deposit a pheromone that guides aggregation and female oviposition decisions. *Elife* **4**, e08688
(2015).

28   Zolin, A. *et al.* Context-dependent representations of movement in Drosophila dopaminergic
reinforcement pathways. *Nature neuroscience* **24**, 1555-1566 (2021).

29   Steele, T. J., Lanz, A. J. & Nagel, K. I. Olfactory navigation in arthropods. *Journal of Comparative*
Physiology A **209**, 467-488 (2023).

30   Hardcastle, K., Ganguli, S. & Giocomo, L. M. Environmental boundaries as an error correction
mechanism for grid cells. *Neuron* **86**, 827-839 (2015).

31   Etienne, A. S., Maurer, R., Boulens, V., Levy, A. & Rowe, T. Resetting the path integrator: a basic
condition for route-based navigation. *Journal of Experimental Biology* **207**, 1491-1508 (2004).

32   Fischler-Ruiz, W. *et al.* Olfactory landmarks and path integration converge to form a cognitive
spatial map. *Neuron* **109**, 4036-4049. e4035 (2021).

33   Green, J. *et al.* A neural circuit architecture for angular integration in Drosophila. *Nature* **546**,
101-106 (2017).

34   Mussells Pires, P., Zhang, L., Parache, V., Abbott, L. & Maimon, G. Converting an allocentric goal
into an egocentric steering signal. *Nature*, 1-11 (2024).

35   Seelig, J. D. & Jayaraman, V. Neural dynamics for landmark orientation and angular path
integration. *Nature* **521**, 186-191 (2015).

36   Westeinde, E. A. *et al.* Transforming a head direction signal into a goal-oriented steering
command. *Nature*, 1-8 (2024).

37   Stern, U. & Yang, C.-H. SkinnerTrax: high-throughput behavior-dependent optogenetic
stimulation of Drosophila. *bioRxiv*, 080614 (2017).

38   Jovanoski, K. D. *et al.* Dopaminergic systems create reward seeking despite adverse
consequences. *Nature* **623**, 356-365 (2023).

39   Stern, U. *et al.* Learning a spatial task by trial and error in Drosophila. *Current Biology* **29**, 2517-
2525. e2515 (2019).

40   Govorunova, E. G., Sineshchekov, O. A., Janz, R., Liu, X. & Spudich, J. L. Natural light-gated anion
channels: A family of microbial rhodopsins for advanced optogenetics. *Science* **349**, 647-650
(2015).

41   Goldschmidt, D. *et al.* Exploration and exploitation are flexibly balanced during local search in
flies. *bioRxiv*, 2024.2006. 2026.600764 (2024).

42   Vosshall, L. B. & Hansson, B. S. A unified nomenclature system for the insect olfactory
coreceptor. *Chemical senses* **36**, 497-498 (2011).

43   Task, D. *et al.* Chemoreceptor co-expression in Drosophila melanogaster olfactory neurons. *Elife*
11, e72599 (2022).

44   Howard, R. W. & Blomquist, G. J. Chemical ecology and biochemistry of insect hydrocarbons.
Annual review of entomology **27**, 149-172 (1982).

45   Howard, R. W. & Blomquist, G. J. Ecological, behavioral, and biochemical aspects of insect
hydrocarbons. *Annu. Rev. Entomol.* **50**, 371-393 (2005).

46   Aso, Y. *et al.* The neuronal architecture of the mushroom body provides a logic for associative
learning. *elife* **3**, e04577 (2014).

47   de Belle, J. S. & Heisenberg, M. Associative odor learning in Drosophila abolished by chemical
ablation of mushroom bodies. *Science* **263**, 692-695 (1994).

48   Handler, A. *et al.* Distinct dopamine receptor pathways underlie the temporal sensitivity of
associative learning. *Cell* **178**, 60-75. e19 (2019).

49   Hulse, B. K. *et al.* A connectome of the Drosophila central complex reveals network motifs
suitable for flexible navigation and context-dependent action selection. *Elife* **10** (2021).

50   Aso, Y. *et al.* Mushroom body output neurons encode valence and guide memory-based action
selection in Drosophila. *elife* **3**, e04580 (2014).

51   Freire, M., Bollig, A. & Knaden, M. Absence of visual cues motivates desert ants to build their
own landmarks. *Current Biology* **33**, 2802-2805. e2802 (2023).

52 Dan, C., Hulse, B. K., Kappagantula, R., Jayaraman, V. & Hermundstad, A. M. A neural circuit
architecture for rapid learning in goal-directed navigation. *Neuron* (2024).
53 Matheson, A. M. *et al.* A neural circuit for wind-guided olfactory navigation. *Nature*
Communications **13**, 4613 (2022).
54 Prokop, A. & Technau, G. M. Normal function of the mushroom body defect gene of Drosophila
is required for the regulation of the number and proliferation of neuroblasts. *Developmental*
biology **161**, 321-337 (1994).

Acknowledgements

862 We would like to thank the Bloomington Stock Center for providing fly strains, the Duke Physics
863 Shop for fabricating the behavior apparati, and Alera Labs (https://aleralabs.com/) for GC-MS
864 analysis. We would also like to thank Drs. Calakos, Nowicki, Volkan, and Dong, and members
865 of the Yang Lab for helpful comments and suggestions. This work was supported by an
866 invertebrate neuroscience research grant awarded to C.-H.Y.

Author contributions

868 Y.C. and C.-H.Y. designed the project. Y.C. performed the experiments and analyzed the data.
869 R.A. and U.S. wrote the code for analyzing the data. U.S. built the behavior-dependent
870 stimulation system. Y.C., R.A., U.S., and C. -H.Y. interpreted the data. D.M. dissected fly brains
871 for imaging. Y.C. wrote the initial draft of the paper, which was then edited by C.-H.Y., U.S., and
872 R.A.

Competing interests

874 The authors declare no competing interests.

Extended Data Fig. 1 SLI of *0273-Gal4>CsChrimson* flies showed weak negative correlation with
average locomotion speed and no correlation with Light PI. a, Average locomotion speed during the
spatial learning task for experimental versus yoked control flies expressing *0273-Gal4>CsChrimson* (n =
83). **b,** *0273-Gal4>CsChrimson* (n = 83) flies showed a very weak negative correlation between average
locomotion speed and SLI. **c,** Top: Temporal structure of the open-loop light (reward) preference assay
(see **Methods**). Bottom: Schematic representation of the flat chamber during the light assay, with the
upper half illuminated during light-on periods. **d,** The positional preference index (Light PI) for illuminated
side in *0273-Gal4>CsChrimson* (n = 36) flies. **e,** Light PI of *0273-GAL4>CsChrimson (*n = 37) flies in
norpA mutant background. **f,** *0273-Gal4>CsChrimson* (n = 70) flies showed no correlation between Light
PI and SLI.

Extended Data Fig. 2 Flies with defective PFNd neurons showed no consistent changes in reward preference (Light PI) or locomotion speed during the spatial learning task. a, Left: Average locomotion speed of *Ctrl>Kir* (n = 74) versus *PFNd>Kir* (n = 68) flies. Right: Light PI of *Ctrl>Kir* (n = 39) versus *PFNd>Kir* (n = 31) flies. **b,** Left: Average locomotion speed of *Ctrl>GtACR1* (n = 41) versus *PFNd>GtACR1* (n = 75) flies. Right: Light PI of *Ctrl>GtACR1* (n = 39) versus *PFNd>GtACR1* (n = 64) flies. **c,** Left: Average locomotion speed of *Ctrl>CsChrimson* (n = 119) versus *PFNd>CsChrimson* (n = 118) flies. Right: Light PI of *Ctrl>CsChrimson* (n = 60) versus *PFNd>CsChrimson* (n = 60) flies. **d,** Left: Average locomotion speed of *Ctrl>CsChrimson* (n = 98) versus *PFNv>CsChrimson* (n = 119) flies. Right: Light PI of *Ctrl>CsChrimson* (n = 40) versus *PFNv>CsChrimson* (n = 39) flies.

Extended Data Fig. 3 Proper activity of allocentric translational velocity-signaling PFR neurons
was dispensable for our spatial task. a, Confocal image and schematic showing the expression pattern
of the *LexA* driver that labels PFR neurons (*GMR37G12-LexA*). **b,** SLI of *Ctrl>Kir* (n = 74, same as the
Ctrl for *EPG>Kir*) versus *PFR>Kir* (n = 57) flies. **c,** SLI of *Ctrl>GtACR1* (n = 79) versus *PFR>GtACR1* flies
(n = 112). **d,** Confocal image and schematic showing the expression pattern of the *split-GAL4* driver that
labels the PFR neurons (*SS54549*). **e,** SLI of *Ctrl>CsChrimson* (n = 140) versus *PFR>CsChrimson* (n =
139) flies.

MBKCs chronically silenced

MBKCs transiently silenced

Extended Data Fig. 4 Flies with disrupted olfactory learning circuit did not show consistent changes in locomotion speed or reward preference. a, Left: Average locomotion speed of Ctrl (n = 83), antenna glued (n = 66), 3rd segment of antenna removed (n = 105), and arista removed (n = 56) flies. Right: Light PI of Ctrl (n = 36), antenna glued (n = 39), 3rd segment of antenna removed (n = 36), and arista removed (n = 38) flies. **b,** Left: Average locomotion speed of *Orco2/+* (n = 98) versus *Orco1/Orco2* (n = 92) flies. Right: Light PI of *Orco2/+* (n = 38) versus *Orco1/Orco2* (n = 37) flies. **c,** Left: Average locomotion speed of control groups *Ir25a & Orco-QF* only (n = 59) and *QUAS-Kir* only (n =88) vs. *Ir25a & Orco-QF>Kir* (n = 90) flies. Right: Light PI of control groups *Ir25a & Orco-QF* only (n = 38) and *QUAS-Kir* only (n =53) vs. *Ir25a & Orco-QF>Kir* (n = 56) flies. **d,** Left: Average locomotion speed of Ctrl (n = 157), hind tarsi removed (n = 146), genitalia glued (n = 103), and flies with both manipulations (n = 104). Right: Light PI of Ctrl (n = 68), hind tarsi removed (n = 75), genitalia glued (n = 76), and flies with both manipulations (n = 57). **e,** Left: Average locomotion speed of *Ctrl>Kir* (n = 113) versus *MBKC-1>Kir* (n = 102) flies. Right: Light PI of *Ctrl>Kir* (n = 38) versus *MBKC-1>Kir* (n = 37) flies. **f,** Left: Average locomotion speed of *Ctrl>Kir* (n = 74, same as the *Ctrl* for *PFNd>Kir*) versus *MBKC-2>Kir* (n = 66) flies. Right: Light PI of *Ctrl>Kir* (n = 39, same as the *Ctrl* for *PFNd>Kir*) versus *MBKC-2>Kir* (n = 34) flies. **g,** Left: Average locomotion speed of *Ctrl>GtACR1* (n = 82) versus *MBKC-1>GtACR1* (n = 144) flies. Right: Light PI of *Ctrl>GtACR1* (n = 39) versus *MBKC-1>GtACR1* (n = 36) flies. **h,** Left: Average locomotion speed of *Ctrl>GtACR1* (n = 82, same as the *Ctrl* for *MBKC-1>GtACR1*) versus *MBKC-2>GtACR1* (n = 153) flies. Right: Light PI of *Ctrl>GtACR1* (n = 39, same as the *Ctrl* for *MBKC-1>GtACR1*) versus *MBKC-2>GtACR1* (n = 40) flies. Welch ANOVA tests with Games-Howell post-hoc test was used for multiple comparisons in panel **a**, **c**, and **d**.

Extended Data Fig. 5 Flies with disrupted olfactory inputs or scent deposition showed reduced
sharp turns toward the goal location. a, Left: Schematic of two boundary crossing events. Middle:

Schematic of a sharp turn event. Right: Example trajectories of sharp turn and non-sharp turn events as
the fly crossed boundaries at a given distance from the goal. **b-j,** Comparison of sharp turn probabilities
toward the goal at different distances from the midline for different groups of flies. **b,** Ctrl (n = 76) versus
antenna glued (n = 64) flies. The same controls were used for the next two panels. **c,** Ctrl (n = 76) versus
3rd segment of antenna removed (n = 102) flies. **d,** Ctrl (n = 76) versus arista removed (n = 55) flies. **e,**
Orco2/+ (n = 98) versus *Orco1/Orco2* (n = 89) flies. **f,** Left: *Ir25a-* & *Orco-QF only* (n = 58) versus *Ir25a-*
QF & *Orco-QF>Kir* (n = 90) flies. Right: *QUAS-Kir only* (n = 88) versus *Ir25a-* & *Orco-QF>Kir* (n = 90)
flies. **g,** Ctrl (n = 129) versus hind tarsi removed (n = 72) flies. The same controls were used for the next
two panels. **h,** Ctrl (n = 129) versus genitalia glued (n = 93) flies. **i,** Ctrl (n = 129) versus both hind tarsi
removed and genitalia glued (n = 59) flies.

Extended Data Fig. 6 Deposited scents were not attractive for naïve flies. a, Proportion of time naïve
flies spent in the 2.5mm radius circle concentric to the reward circle in clean flat chambers (n = 76) versus
the same flies trained after 3 training sessions. **b,** Proportion of time naïve flies spent in the 2.5mm radius
circle concentric to the reward circle in clean flat chambers (n = 76) versus "used" chambers previously
occupied by flies that had just completed 3 training sessions (n = 104). The duration of this preference
assay was 3 minutes.

Extended Data Fig. 7 Flies with HU-ablated mushroom body (MB) showed strongly reduced learning performance. a, Top: Fas-2 staining in the MB lobes of an untreated fly. Bottom: Fas-2 staining in the MB lobes of a hydroxyurea (HU) treated fly. **b,** SLI of Ctrl (n = 62) versus MB-ablated (n = 39) flies. **c,** Light PI of Ctrl (n = 20) versus MB-ablated (n = 20) flies. **d,** Average locomotion speed of Ctrl (n = 62) versus MB-ablated (n = 39) flies.

Extended Data Fig. 8 Flies with silenced MBKCs showed reduced sharp turns toward the goal location. a, Left: Turn probability of *Ctrl>Kir* (n = 110) versus *MBKC-1>Kir* (n = 101) flies. Right: Turn probability of *Ctrl>Kir* (n = 73) flies versus *MBKC-2>Kir* (n = 66) flies. **b,** Left: Turn probability of *Ctrl>GtACR1* (n = 81) flies versus *MBKC-1>GtACR1* (n = 142) flies. Right: Turn probability of *Ctrl>GtACR1* (n = 81, same as the *Ctrl* for *MBKC-1>GtACR1*) flies versus *MBKC-2>GtACR1* (n = 153) flies.

Extended Data Fig. 9 Learning induced avoidance of agarose-occupied areas at the end of the task. a, Schematic of agarose-equivalent areas in the flat chamber (left) and agarose areas in the agarose chamber (right). **b,** Left two columns: Percentage of time flies spent in agarose-equivalent areas in the flat chamber (n = 113) versus agarose areas in the agarose chamber (n = 115) during the pre-training session. The difference was statistically significant but mild. Right two columns: Percentage of time flies spent in agarose-equivalent areas in the flat chamber (n = 83) versus agarose areas in the agarose chamber (n = 105) during the last sync bucket (10 mins) of the training 3. Two-way ANOVA was used to calculate the interaction between chamber types and learning task sessions on the percentage of time flies spent in agarose-equivalent areas. Šídák's multiple comparisons test was used to compare values between the flat and agarose chambers at the given session of the task.

Extended Data Fig. 10 Flies use agarose areas as olfactory landmarks to learn the spatial task. a,
Left two columns: SLI of Ctrl (n = 83) versus 3rd segment of antenna removed (n = 105) flies in the flat
chamber. Right two columns: SLI of Ctrl (n = 105) versus 3rd segment of antenna removed (n = 67) flies in
the agarose chamber. Two-way ANOVA was used to calculate the interaction between 3rd segment
removal and chamber type on SLI. Šídák's multiple comparisons test was used to compare SLI between
Ctrl and antenna removed flies for each given chamber type. **b,** Left three columns: SLI of control groups
Ir25a & Orco-QF only (n = 59) and *QUAS-Kir* only (n = 88) versus *Ir25a* & *Orco-QF>Kir* flies (n = 90) in
the flat chamber. Right three columns: SLI of control groups *Ir25a & Orco-QF* only (n = 67) and *QUAS-Kir*
only (n = 74) versus *Ir25a & Orco-QF>Kir* flies (n = 74) in the agarose chamber. Two-way ANOVA was
used to calculate the interaction between silencing *Ir25a & Orco*-expressing OSNs and chamber type on
SLI. Šídák's multiple comparisons test was used to compare SLI between control groups and *Ir25a* &
Orco-QF>Kir flies for each chamber type. **c,** Schematic of the analysis determining the distribution of
distances where flies initiated their first large turn after departing from the reward circle (see **Methods**). **d,**
Distribution of distances where flies initiated their first large turn after departing from the reward circle in
training session 1 versus session 3 (n = 113) in the agarose chamber. The dotted blue line shows the
shortest distance from the reward circle to positions where the fly could physically contact the agarose
area. Paired two-tailed t-test was used to compare mean distances from the reward circle between the
two groups. **e-g,** Distribution of distances where flies initiated their first large turn departing from the
reward circle goal for different groups of flies in the agarose chamber. **e,** Ctrl (n = 114) versus 3rd segment
of antenna removed (n = 82) flies. **f,** *Orco2/+* (n = 80) versus *Orco1/Orco2* (n = 77) flies. **g,** Left: *Ir25a-QF*
& Orco-QF only (n = 73) versus *Ir25a & Orco-QF>Kir* (n = 83) flies. Right: *QUAS-Kir* only (n = 79) versus
Ir25a & Orco-QF>Kir (n = 83) flies. Stars above each bin show the result of unpaired two-tailed t-tests
with Welch's correction for frequency distribution in each bin. Welch's corrected unpaired two-tailed t-tests
were also used to compare mean distances from the reward circle during the training session for each fly
between groups.

Extended Data Fig. 11 Olfactory cues and self-motion cues independently contribute to spatial learning, with flexible adjustment in cue usage. a, SLI of *PFNd>Kir* (n = 68) versus *PFNd>Kir* with 3rd segment of antenna removed (n = 105) flies in the flat chamber. **b,** *Ctrl>Kir* with 3rd segment of antenna removed (n = 161) versus *PFNd>Kir* with 3rd segment of antenna removed (n = 105) flies in the flat chamber. **c,** Left two columns: SLI of *Ctrl>K*ir (n = 110) versus *PFNd>Kir* (n = 99) flies in the agarose chamber. Right two columns: SLI of *Ctrl>Kir* with 3rd segment of antenna removed (n = 51) versus *PFNd>Kir* with 3rd segment of antenna removed (n = 68) flies in the agarose chamber. Two-way ANOVA was used to calculate the interaction between silencing PFNd neurons and antenna removal on SLI. Šídák's multiple comparisons test was used to compare the SLI between *Ctrl>Kir* and *PFNd>Kir* flies with normal and 3rd segment of antenna removed flies.

1007

1008 **Extended Data Fig. 12 *Or7a* mutant flies did not show decreased spatial learning performance. a,**
1009 *RI difference* between experimental flies and their paired yoked controls of *Ctrl* and *Or7a* mutant flies in
1010 the flat chamber over time.

1011

Extended Data Fig. 13 Optogenetic activation of the *Gr43a-Gal4* labeled sweet-sensing neurons at
the reward circle could also induce learning that depended on intact antennae. a, *RI difference* over
time between unmanipulated experimental and yoked control *Gr43a-Gal4>CsChrimson* flies (solid line)
and between antenna-removed experimental and yoked control *Gr43a-Gal4>CsChrimson* flies (dashed
line). Note that the decline of the RI difference over time might be due to the flies being starved 20 hours
before the task and not receiving any food during the entire task, which might have led to fatigue.

Table 1. List of Putative Observed Compounds identified by GC-MS

1019

Retention Time (min)	Compound ID	Formula	Exp Tracking Mass (ug)	Yoked Tracking Mass (ug)
19.1	Tricosene isomer	$C_{23}H_{46}$	0.28	0.13
19.2	Tricosene isomer	$C_{23}H_{46}$	3.11	1.53
19.4	Tricosene isomer	$C_{23}H_{46}$	0.20	0.10
19.5	Tricosene isomer	$C_{23}H_{46}$	0.78	0.44
20.6	Cyclotetracosane	$C_{24}H_{48}$	0.18	0.09
21.9	Pentacosene isomer	$C_{25}H_{50}$	0.12	0.08
22.0	Pentacosene isomer	$C_{25}H_{50}$	0.65	0.46
22.3	Pentacosene isomer	$C_{25}H_{50}$	0.08	0.05

1020

Supplementary Videos

Video S1. High-throughput spatial learning task with *0273Gal4>CsChrimson* flies in the flat
chamber

Left half: Example video showing 10 experimental flies and their paired yoked control flies undergoing
training in our high-throughput spatial learning task in the flat chamber. Right half: The real-time positional
heatmap of corresponding flies during training. The video is at 2X play back speed.

Video S2. High-throughput spatial learning task with *0273Gal4>CsChrimson* flies in the agarose
chamber

Left half: Example video showing 10 experimental flies and their paired yoked control flies undergoing
training in our high-throughput spatial learning task in the agarose chamber. Right half: The real-time
positional heatmap of corresponding flies during training. The video is at 2X play back speed.