Lgl resets Par complex membrane loading at mitotic exit to polarize neural stem cells

Bryce LaFoya*, Sarah E. Welch*, and Kenneth E. Prehoda[1]

Institute of Molecular Biology
Department of Chemistry and Biochemistry
1229 University of Oregon
Eugene, OR 97403

*contributed equally

[1]Corresponding author: prehoda@uoregon.edu

Summary

The Par complex regulates cell polarity in diverse animal cells, but how its localization is restricted to a

specific membrane domain remains unclear. We investigate how the tumor suppressor Lethal giant

larvae (Lgl) regulates Par complex polarity in *Drosophila* neural stem cells (NSCs). Unlike in epithelial

cells, where Lgl and the Par complex show mutual exclusion, in NSCs the Par complex is apically

polarized while Lgl is cytoplasmic during mitosis. We demonstrate that Lgl's primary function is to

remove the Par complex from the membrane at mitotic exit, resetting polarity for the next cell cycle. In

Lgl-depleted NSCs, the Par complex persists on both apical and basal membranes throughout the cell

cycle. Surprisingly, apical Par complex targeting and polarization occur independently of Lgl, but

residual basal Par complex is sufficient to disrupt asymmetric division. These findings reveal a novel

mechanism of negative polarity regulation by Lgl and highlight the importance of the pre-polarized

state in Par-mediated polarity.

Introduction

Cortical polarity is a fundamental cellular characteristic that underlies diverse biological processes,

including directional cell migration, tissue organization, and asymmetric cell division. The Par complex,

consisting of Par-6 and atypical Protein Kinase C (aPKC), plays a pivotal role in establishing and

maintaining polarity in a wide range of animal cells, from epithelial cells to neurons[1–4]. The proper

localization and function of the Par complex are particularly critical in stem cells, where asymmetric

distribution of cell fate determinants governs the delicate balance between self-renewal and

differentiation. A fundamental question in cell polarity research is how the Par complex itself is

restricted to specific membrane domains. The tumor suppressor Lethal giant larvae (Lgl) plays a key

role in Par polarity by restricting the Par complex to the proper membrane domain[5–8], but how Lgl

controls Par complex localization remains an open question.

Par polarity results from the complex interplay of positive and negative regulators. While positive

regulators like Par-3 (Bazooka in *Drosophila*) and Cdc42 ensure that the Par complex is targeted to the

membrane[9–11], the negative regulation provided by Lgl is thought to arise from mutual exclusion, where

the Par complex and Lgl reciprocally control each other's localization[4]. This model is exemplified by

interphase epithelial cells, where the Par complex is confined to the apical plasma membrane while Lgl

occupies the basolateral domain (Fig. 1A). The importance of this mutual regulation is evident when Lgl

function is lost: in Lgl mutants the Par complex loses its domain specificity and spreads into the

basolateral domain[5,6,8]. Conversely, in Par complex mutants, Lgl is no longer restricted to the

basolateral membrane and becomes uniformly distributed across the entire cell surface, including the

apical domain. The Par complex regulates Lgl localization through phosphorylation by aPKC[12,13].

Phosphorylation reduces Lgl's affinity for the membrane, ensuring that it only binds membrane regions

that lack Par activity. While the Par complex controls Lgl localization by coupling phosphorylation to a

decrease in membrane affinity, little is known about how Lgl regulates Par complex localization.

Like epithelial cells, sensory organ precursor (SOP) cells and neural stem cells (NSCs) require Lgl

function for proper Par complex polarity[14,15]. However, the dynamics of Par complex localization in

these cell types differ significantly from those in epithelial cells (Fig. 1A). In contrast to epithelial cells,

where the Par complex is polarized throughout the cell cycle, the Par complex in SOPs and NSCs is not

polarized during interphase but becomes polarized during mitosis[5,15]. For example, in NSCs the Par

complex localizes to the apical membrane at metaphase where it directs fate determinants like

Prospero to the basal side, ultimately leading to asymmetric sibling cell fates[16,17]. In Lgl mutant SOPs

and NSCs, the Par complex is localized across the entire membrane instead of being restricted to its

proper domain[14], suggesting that Lgl is required for the asymmetric targeting of the Par complex.

Importantly, SOPs and NSCs do not conform to the mutual exclusion model observed in epithelial cells.

While Lgl is removed from the membrane early in mitosis in all these cell types, the critical difference

lies in the timing of Par complex polarization. By metaphase, when the Par complex becomes polarized

in SOPs and NSCs, Lgl is no longer present on the membrane[5,15]. This temporal separation between Lgl

membrane dissociation and Par complex polarization challenges the idea of direct mutual exclusion

between these proteins in SOPs and NSCs, raising questions about how Lgl regulates Par complex

localization in these cell types.

Given that disruptions in cell polarity are frequently associated with tumorigenesis, and that Lgl is a

known tumor suppressor, unraveling its mode of action and localization patterns may offer new

perspectives on the links between polarity, asymmetric division, and cancer development. We

examined Lgl and Par complex dynamics in NSCs to resolve the discrepancy between the mutual

exclusion model and Lgl's function in these cells. Our findings shed light on the molecular basis of Lgl's

function as a negative regulator of Par polarity and provide insights into the broader principles

governing cell polarization in diverse cellular contexts.

Results

Patterned Lgl membrane displacement in early NSC mitosis

Lethal giant larvae (Lgl) is bound to the plasma membrane during interphase and becomes cytoplasmic

during mitosis in neural stem cells (NSCs)[5]. We sought to determine the dynamics of Lgl's transition

between these states including how it is restored to the membrane following mitosis. We examined

Lgl's localization throughout the cell cycle NSCs from *Drosophila* larval brain explants using rapid,

super-resolution spinning disk confocal microscopy (Fig. 1B). We imaged NSCs expressing Lgl-GFP

(expressed by Worniu-GAL4 driven UAS) and RFP-Histone H2A (expressed from its endogenous locus)

to visualize Lgl localization relative to cell cycle progression. Lgl removal from the membrane began in

early prophase, initiating at a small area near the apical pole of the cell (Fig. 1C,D; Video 1). The removal

from the membrane spread rapidly across the cell surface and the apically-initiated pattern gave rise to

a state in which Lgl is polarized to the basal membrane. However, this state is short-lived as Lgl

removal continued without any apparent delay such that it appeared to be completely displaced into

the cytoplasm by late prophase and could be observed both in medial sections and maximum intensity

projections of the full cell volume (Fig. 1E). The full volume data also revealed how Lgl displacement

from the membrane was affected by the membrane structures that are present on the NSC surface[18–20].

These structures were enriched with Lgl, consistent with their higher density of membrane, and Lgl

displacement from the structures proceeded with the same overall dynamics as the surrounding

membrane (Fig. 1E). To confirm that Lgl was completely removed from the basal membrane, we

imaged individual cultured NSCs that had been dissociated from brains. The Lgl dynamics in

dissociated NSCs were consistent with our initial observations of apically-directed membrane clearing

and provided unambiguous evidence that Lgl is indeed completely removed from the membrane by

late prophase (Fig. 1F,G). The patterned Lgl membrane displacement dynamics indicate that removal is

a spatially regulated process, like that observed for SOPs[15], rather than one in which it is simultaneously

removed from across the entire cell surface.

NSCs undergo repeated cycles of asymmetric division such that Lgl must be restored to the membrane

at some point to achieve its interphase localization state. We analyzed Lgl dynamics in both intact brain

and dissociated NSCs to determine when and how Lgl is restored to the membrane following mitosis.

Our results showed that Lgl becomes rapidly restored to the membrane at the completion of mitosis,

beginning near the cytokinetic pore connecting the nascent sibling cells (Fig. 1H-J), like what has been

observed in follicular epithelial cells[21].

Our results provide a detailed characterization of Lgl's dynamic localization throughout the cell cycle in

NSCs, revealing precise spatial and temporal control of its membrane association (Fig. 1K). The

patterned removal of Lgl from the membrane during prophase and its rapid restoration post-mitosis

suggest the existence of tightly regulated mechanisms controlling Lgl localization.

aPKC removes Lgl from the NSC early in mitosis

Lgl can be removed from the membrane by aPKC but also by the mitotic kinase Aurora A[5,6]. We sought

to determine whether one or both kinases are responsible for the patterned removal of Lgl from the

membrane during prophase. To dissect the relative contributions of aPKC and Aurora A to Lgl removal,

we separately inhibited aPKC or Aurora A activity and assessed the effect on Lgl localization dynamics.

We inhibited aPKC by expressing a UAS-controlled aPKC RNAi specifically in larval brain NSCs using

Worniu-GAL4 driven expression. For Aurora A inhibition, we added the specific inhibitor MLN8237[6] to

the larval brain explant culture media.

In NSCs where aPKC was depleted by RNAi, we observed that Lgl removal from the membrane in

mitosis was largely inhibited (Fig. 2A,B; Video 4). Unlike wild type NSCs where we were unable to

detect Lgl at the apical membrane late in prophase, aPKC RNAi NSCs had persistent Lgl membrane

enrichment throughout mitosis. However, we did detect a small decrease in apical Lgl membrane signal

beginning in early prophase indicating that aPKC is not the only kinase that removes Lgl in NSCs,

and/or there is residual aPKC activity in aPKC RNAi NSCs. The retention of Lgl on the membrane in

aPKC-depleted cells indicates that aPKC-mediated phosphorylation is the key mechanism driving Lgl

dissociation from the cell cortex as NSCs enter mitosis. Interestingly, we also observed an increase in

Lgl's membrane enrichment during interphase in aPKC RNAi NSCs (Fig. 2A,B), suggesting that Lgl

exchanges between the membrane and cytoplasm due to phosphorylation and dephosphorylation in

wild type NSCs.

Consistent with our results indicating that aPKC is the dominant kinase responsible for displacement of

Lgl from the membrane in NSCs, we found that inhibition of Aurora A activity using MLN8237, did not

cause a significant difference in Lgl removal from the membrane compared to wild-type NSCs (Fig.

2C,D; Video 5). The patterned removal of Lgl from the membrane proceeded with similar dynamics and

extent as observed in untreated cells. However, we did observe cytokinesis defects in drug treated

NSCs (Fig. 2C; Video 5) indicating that Aurora A was successfully inhibited. We conclude that, despite

its known ability to phosphorylate Lgl, Aurora A does not participate in removal of Lgl from the NSC

membrane. This conclusion is consistent with a previous study that found the Aurora A phosphorylation

sites on Lgl were dispensable for its removal from the membrane in NSCs[5].

aPKC patterns Lgl membrane clearance

Given our findings implicating aPKC as the primary kinase responsible for Lgl removal from the

membrane during prophase, we next sought to determine whether the spatial and temporal dynamics

of aPKC localization are consistent with the observed pattern of Lgl removal. The recruitment of aPKC

to the membrane in early prophase is a complex, multistep process where aPKC is initially localized to

small domains on the apical surface before coalescing into an apical cap by metaphase[22,23]. To

determine the interplay between Lgl's disappearance and aPKC's appearance on the membrane, we

simultaneously imaged aPKC-GFP (expressed from its endogenous promoter) and Lgl-mCherry

(expressed by Worniu-GAL4 driven UAS) in NSCs. This approach allowed us to directly correlate the

spatial and temporal dynamics of aPKC recruitment with the pattern of Lgl removal from the cell

membrane. We observed that Lgl removal at the apical pole of the NSC was tightly correlated with the

appearance of aPKC at that site (Fig. 3A,B; Video 6). The correlation between aPKC's appearance and

Lgl's disappearance from the membrane continued until aPKC was fully targeted to the apical

membrane. However, Lgl's removal from the membrane continued after aPKC targeting was

completed. We conclude that aPKC removes Lgl from the membrane both directly, by removing it from

membrane regions that it occupies, and indirectly. Indirect removal of basal Lgl by apical aPKC is likely

influenced by factors such as the rate of Lgl diffusion along the plasma membrane.

Lgl excludes basal aPKC by regulating aPKC interphase membrane association

The mutual exclusion model, which posits that Lgl and the Par complex reciprocally control each other's

localization by occupying an opposing membrane domain, does not explain the behavior of these

proteins in NSCs or SOPs. The primary inconsistency is the metaphase polarized state when the Par

complex polarized while Lgl is cytoplasmic. However, it is also not known whether Lgl regulates aPKC's

cytoplasmic localization in interphase. To gain insight into how Lgl regulates Par complex localization in

these contexts, we examined aPKC localization in NSCs expressing Lgl RNAi (Worniu-GAL4 driven

UAS). In wild-type NSCs, aPKC is typically excluded from the cell membrane during interphase.

However, in Lgl-depleted NSCs, we observed a key difference from wild type NSCs where aPKC rapidly

returns to the cytoplasm at the end of mitosis. Rather than rapidly dissipating from the membrane,

aPKC remained on the membrane throughout interphase in the absence of Lgl (Figure 4A,B; Video 7).

This observation indicates that Lgl plays a crucial role in regulating the interphase membrane

localization of aPKC, preventing its premature association with the cell cortex.

As Lgl RNAi NSCs progressed into mitosis, we observed aPKC at the basal membrane and this signal

persisted through metaphase (Fig. 4A,C; Video 7). Importantly, our results suggest that the presence of

aPKC on the basal membrane during metaphase in Lgl-depleted NSCs is not due to new recruitment

during mitosis. Instead, it appears to result from aPKC that remained associated with the membrane

throughout interphase. Despite the aberrant basal localization of aPKC in Lgl-depleted NSCs, we

observed that the positive signals responsible for recruiting aPKC to the apical membrane in wild-type

NSCs remain in Lgl RNAi cells. We observed an increase in apical aPKC from early prophase to

metaphase indicating that Lgl is not required for apical recruitment of the Par complex in NSCs. In fact,

even though aPKC is present on the basal membrane, the recruitment to the apical membrane leads to

metaphase polarity (Fig. 4A,D; Video 7). Thus, the defects caused by loss of Lgl function aren't strictly

due to the loss of Par polarity, but instead are due to the presence of basal Par complex. These results

provide important insights into how Lgl regulates Par complex localization in NSCs, revealing a more

nuanced picture than that suggested by the mutual exclusion model. Lgl appears to play a critical role in

preventing premature membrane association during interphase by ensuring proper removal from the

membrane during mitotic exit (Fig. 4E).

Discussion

How Lethal giant larvae (Lgl) controls Par complex polarity is a key outstanding question in animal cell

polarity. Evidence from localization studies and mutant phenotypes in epithelial cells support a mutual

exclusion model in which Lgl and the Par complex control one another's localization by occupying a

reciprocal membrane domain[4,24]. This model suggests that depolarization of the Par complex in the

absence of Lgl function could result from defects in asymmetric targeting – rather than being targeted

to a polarized membrane domain, the Par complex is targeted uniformly to the membrane. However,

this model fails to explain Par polarity in sensory organ precursor cells (SOPs) and neural stem cells

(NSCs), where Lgl is cytoplasmic during mitosis while the Par complex remains polarized[5,15]. We aimed

to resolve this discrepancy and elucidate the mechanism by which Lgl polarizes the Par complex in

NSCs when it isn't present on the membrane during mitosis. Our results support a model distinct from

mutual exclusion, where Lgl plays a key role in establishing the starting point for NSC and SOP polarity

in which the Par complex is not present on the membrane.

Our results indicate that Lgl is not required for polarized targeting of the Par complex. We observed

enrichment at the apical membrane in NSCs lacking Lgl function. This led to a state in which the Par

complex is present on the basal membrane, but more concentrated at the apical membrane. Thus, Lgl

isn't strictly required for Par polarity, since the Par complex is asymmetric in Lgl RNAi NSCs. Instead,

Lgl is critical for ensuring that the Par complex isn't present on the basal membrane when apical

targeting of the complex begins in early prophase. Basal Par complex, even at the reduced levels we

observed compared to apically-enriched complex, is sufficient to disrupt the polarity of downstream

components such as Miranda, which become cytoplasmic in *lgl* mutants[14].

These findings suggest a revised model for Lgl's role in controlling Par complex polarity in NSCs. Rather

than acting through mutual exclusion during mitosis, Lgl primarily functions to ensure the complete

removal of the Par complex from the membrane at the end of mitosis. This removal is crucial for

resetting the cell's polarity state, allowing for proper polarization in the subsequent cell cycle. Our

results highlight the importance of considering the entire cell cycle when studying polarity regulation.

The effect of Lgl on Par complex localization during interphase has profound consequences for mitotic

polarization, even though Lgl itself is cytoplasmic during this phase. This study also underscores the

context-dependent nature of polarity regulation. While the mutual exclusion model may accurately

describe Lgl-Par complex interactions in epithelial cells, it does not fully capture the dynamics in NSCs.

The mechanism by which Lgl removes aPKC from the membrane at the end of mitosis remains an

intriguing question. There are several possibilities based on our current understanding of Lgl and aPKC.

One potential mechanism is that Lgl, upon its rapid return to the membrane post-mitosis, could act as a

competitive inhibitor for aPKC membrane binding sites. Lgl's strong affinity for the membrane[12,13]

might allow it to outcompete aPKC and displace it from the membrane. Alternatively, Lgl could recruit

or activate a phosphatase that dephosphorylates aPKC or its binding partners, thereby reducing aPKC's

affinity for the membrane. Further investigation into these potential mechanisms will be crucial for fully

understanding Lgl's role in polarity regulation.

Resource Availability

Lead Contact

Contact the Lead Contact, Kenneth Prehoda (prehoda@uoregon.edu), for further information or
to request resources and reagents.

Materials Availability

No new reagents were generated in this study.

Data and Code Availability

Raw data available from the corresponding author on request.

Experimental Model and Subject Details

Fly Strains

Tissue specific expression of UAS controlled transgenes in NSCs was achived using a Worniu-GAL4 driver line. Lgl dynamics were imaged using UAS-Lgl-GFP and the membrane marker UAS-PLCδ-PH-mCherry, which expresses the pleckstrin homology domain of human PLCδ tagged with mCherry and binds to the plasma membrane lipid phosphoinositide $PI(4,5)P_2$ as a fiduciary marker. The specific mitotic stages were monitored using the chromosome marker His2A-RFP. aPKC and Lgl co-imaging was achieved using UAS-Lgl-mCherry to monitor Lgl localization. and the BAC-encoded-GFP-aPKC [25] was used to monitor aPKC localization. Worniu-Gal4 was used to express UAS-RNAi against aPKC and Lgl in NSCs. Knockdown was enhanced through the coexpression of UAS-Dicer2.

Method Details

Live Imaging
Third instar larvae were incubated at 30 ℃ for at least 48 hours prior to imaging and dissection. The brain lobes and ventral nerve cord were dissected in Schneider's Insect Media (SIM) from each larvae, and mounted on a sterile poly-D-lysine coated 35mm glass bottom dish (ibidi Cat#81156) containing modified minimal hemolymph-like solution (HL3.1). Each brain was imaged using a Nikon Eclipse Ti-2 Yokogawa CSU-W1 SoRa spinning disk microscope with dual Photometrics Prime BSI sCMOS cameras, utilizing a 60x water immersion objective. GFP-tagged proteins were visualized using 488 nm illumination, while RFP and mCherry tags were excited with 561 nm light. Super-resolution images were captured via SoRa (super-resolution through optical photon reassignment) optics, as described by Azuma and Kei (2015). Neuroblasts were distinguished based on size, central nervous system location, and expression of the Worniu-GAL4 neuroblast-specific driver line.

To acquire "dissociated NSCs", *Drosophila* central nervous systems were collected from third instar larvae and incubated for 30 min at 30°C in Schneider's Insect Medium containing 1mg/mL Papain and 1mg/mL Collagenase. After incubation, mechanical dissociation was achieved by flushing medium/brains through a pipette tip. The resulting cell suspension was then pelleted using gentle centrifugation and washed 3 times with culture medium consisting of Schneider's Insect Medium containing 10% synthetic fetal bovine serum, 20mM Glutamine, 0.05mg/mL

252 Glutathione, and 0.01mg/mL Insulin. After the final wash step, cells were resuspended and
253 plated in the previously described culture medium and imaged.
254
255 **Pharmacological Inhibition**
256
257 The selective Aurora A inhibitor MLN8237 (Alisertib) was used at a working concentration of
258 15µm solubilized in DMSO in all relevant experiments.
259
260 **Image Processing and Analysis**
261 Imaging data was processed using ImageJ (FIJI package). For some movies, the bleach
262 correction tool was used to correct for photobleaching.
263
264 Quantification for Figure 1D To quantify patterned Lgl removal, NSCs were analyzed by
265 identifying the frame where the apical membrane-to-cytoplasm ratio neared 1, indicating
266 complete Lgl removal. A 5-pixel line perpendicular to the apical membrane was used to capture
267 the membrane signal, and a cytoplasmic value was obtained from a square region excluding the
268 nuclear envelope. This ratio was measured at both the apical and lateral regions.
269
270 Quantification for Figure 1H: To quantify patterned Lgl removal in dissociated NSCs (n=3), the
271 same method of measuring the membrane to cytoplasmic ratio as was used in figure 1D was
272 used, in which the frame where the apical membrane to cytoplasm ratio neared 1 was identified,
273 and the apical membrane to cytoplasmic ratio and the equatorial membrane to cytoplasmic ratio
274 was measured to show depletion of Lgl.
275
Quantification for Figure 1G: For basal Lgl removal in dissociated NSCs, the same
quantification method as used for the previous figure was applied. The basal membrane-to-
cytoplasm ratio was plotted relative to metaphase, which was defined as the frame before
anaphase onset. Anaphase onset was defined as the frame where chromosome separation from
the metaphase plate first occurred, using His2A-RFP-tagged chromosomes or UAS-Lgl-GFP
after nuclear envelope breakdown as a chromosomal marker.
282
Quantification for Figure 1J: To quantify Lgl reloading, UAS-Lgl-GFP dissociated neural stem
cells (NSCs) were used, as these provided a clearer view of the basal side compared to in-tissue
cells. In these cells, Lgl appears to be recruited first to the basal cortex. The analysis identified
the first frame during anaphase in which Lgl was fully cytoplasmic. Anaphase was defined as
any time after the chromosomes had visibly separated. This frame was designated as the "Pre-
reloading" time point. The next key frame was where Lgl reloading was observed at both the
midbody and lateral cortex, which was labeled as the "Reloading" time point. The final frame of
each movie, where Lgl was confirmed to be uniformly cortical, was taken as the "Cortical" time
point. At each of these time points, the membrane-to-cytoplasm Lgl ratio was measured using the
previously described method at three locations: the midbody, the lateral (equatorial) cortex, and
the apical cortex. The lateral cortex was defined as the equator of the cell, and the apical side was
positioned opposite the midbody. The membrane-to-cytoplasm ratios at the midbody, lateral, and
apical regions were quantified at the pre-reloading, reloading, and cortical time points for each
UAS-Lgl-GFP NSC.

Quantification for Figure 2B: To quantify ectopic membrane-bound Lgl in the aPKC RNAi
condition, UAS-Lgl-GFP cells were first analyzed to confirm the typical pattern of Lgl
localization. Lgl was fully membrane-bound during interphase (15 minutes before nuclear
envelope breakdown, NEB), partially removed in early mitosis (7 minutes before NEB), and
completely removed by metaphase (defined as one frame before anaphase onset). The
membrane-to-cytoplasmic Lgl ratio was measured at these time points for both the control
(UAS-Lgl-GFP n =5) and aPKC RNAi (n=5), using the quantification method described in
Figure 1.

Quantification for Figure 2D: To quantify the cortical removal of Lgl during aurA inhibition,
each movie was analyzed to identify two key time points: (1) the last frame before nuclear
envelope breakdown (NEB) where Lgl was fully cortical, and (2) the frame where Lgl was
completely cleared in relation to anaphase onset in DMSO control cells. The earliest usable time
point was 6 minutes and 40 seconds before NEB. Lgl was fully removed by metaphase, defined
as the frame immediately preceding anaphase onset. These time points were designated as "Early
Mitosis" and "Metaphase," respectively. The membrane-to-cytoplasmic ratio at the apical region
was measured using previously described methods for both time points. Comparisons were made
between aurA-inhibited cells (n=3) and DMSO control cells (n=3) for the early mitosis and
metaphase time points.

Quantifications for Figure 3B: To quantify the simultaneous release of Lgl and recruitment of
aPKC at the apical pole, the frame marking the initial clearance of Lgl from the apical membrane
was selected for analysis. In this frame, the apical membrane-to-cytoplasmic ratio was calculated
for cells co-expressing BAC aPKC-GFP and UAS-Lgl-mCherry using previously established
methods (N=3).

Quantification for Figure 4B: To quantify aPKC-GFP in the apical region in the Lgl-RNAi
condition, the same timepoints as the aPKC-RNAi experiment in figure 2 were used. It was first
confirmed that at interphase (15 minutes pre NEB) that wildtype aPKC was completely
cytoplasmic. It was then confirmed at early mitosis (7 minutes pre NEB) that aPKC was just
beginning to localize to the apical membrane. By metaphase, which again was defined as the
frame before anaphase onset, wildtype aPKC is strongly localized at the apical crescent. The
apical membrane to cytoplasmic ratio was measured for both Lgl-RNAi cells (n=3) and WT-
aPKC cells (n=3).
328
329   Quantifications for Figure 4C: To quantify ectopic basal aPKC in the Lgl-RNAi condition, it was
330   determined in WT aPKC cells that aPKC was fully polarized on the apical membrane, and
331   completely absent from the basal membrane by metaphase, which was determined to be the
332   frame before anaphase onset.
333
334   **Statistical Analysis**

335   Gardner-Altman estimation plots and 95% confidence intervals of datasets were prepared using
336   the DABEST package [26]. Statistical details can be found in the relevant methods section and
337   figure legends.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Model: Fly Strains		
BacaPKC-GFP	François Schweisguth Lab; (Besson et al., 2015)	PMID: 25843034
UAS-Lgl-GFP	Juergen A. Knoblich Lab; (Writz-Peitz et al., 2008)	PMID: 25484300
Endogenous Lgl-GFP	Bloomington *Drosophila* Stock Center (BDSC)	BDSC Cat#63183 RRID: BDSC 63183
His2a-RFP (chromosome marker)	Bloomington *Drosophila* Stock Center (BDSC)	BDSC Cat# 23650 RRID: BDSC 23650
Worniu-GAL4 (NSC driver line)	Chris Q. Doe Lab	
aPKC-RNAi	Bloomington *Drosophila* Stock Center (BDSC)	BDSC Cat#67204 RRID: BDSC 67204
Lgl-RNAi	Bloomington *Drosophila* Stock Center (BDSC)	BDSC Cat#38989 RRID: BDSC 38989
UAS-Lgl-mCherry	Bloomington *Drosophila* Stock Center (BDSC)	BDSC Cat#31776 RRID: BDSC 31776
UAS-PLCδ-PH-mCherry (membrane marker)	Bloomington *Drosophila* Stock Center (BDSC)	BDSC Cat#51658 RRID: BDSC 51658
Pharmacological Inhibitors		
MLN8237 (Alisertib)	Selleckchem	Cat#S1133

Acknowledgments

We thank Adam Fries for maintaining the microscope used in this study. This work was supported by NIH grants R35GM127092 and K99GM147601.

Author Contributions

357 B.L. and K.E.P. designed the experiments. B.L. performed the experiments. R.R.P. contributed
358 to the String inhibition and String staining experiments. B.L. and K.E.P analyzed the data,
359 prepared the figures, and wrote the manuscript.

360 **Declaration of Interests**

361 The authors have no competing interests to declare.
362
Figure Legends
Figure 1 Patterned Lgl displacement and reloading during neural stem cell asymmetric division
(A) Mutual exclusion model for Par polarity. In this model, the Par complex (Par-6 and atypical
Protein Kinase C or aPKC) forms a mutually exclusive membrane domain with Lethal giant
larvae (Lgl) where the presence of one on the membrane excludes the other. This model fails to
account for Drosophila neural stem cells (NSCs) and sensory organ precursors where the Par
complex is polarized during mitosis and Lgl is cytoplasmic.
(B) Imaging Drosophila larval brain neural stem cells (NSCs). The larval central nervous system
(brain lobes and ventral nerve cord) are isolated from L3 stage larvae and the NSCs, identified
by their relatively larger size, are either imaged in the intact CNS or the tissue is dissociated and
isolated NSCs are imaged.
(C) Removal of Lgl from the plasma membrane in early mitosis in an NSC from the intact central
nervous system. Frames from Video 1 are shown with Lgl-GFP (expressed from Worniu-GAL4
driven UAS) and Histone H2A (His2a-RFP) with the NSC marked with asterisk. Time is shown
relative to nuclear envelope breakdown.
(D) Patterned removal of Lgl from the plasma membrane in early mitosis. Lgl's membrane
enrichment relative to the cytoplasm is shown for 10 different NSCs at the apical and lateral
region in early mitosis when Lgl was first cleared from the apical membrane. Gardner-Altman
estimation plot shows the paired measurements for each NSC. The 95% confidence interval is
shown in the mean difference comparison along with the bootstrap distribution from which it was
derived.
(E) Lgl dynamics across the cell cycle and in the full cell volume. Frames from Video 2 are
shown with views of both a medial optical section and the full volume of the NSC reconstructed
using a maximum intensity projection (MIP) of optical sections across the cell.
(F) Removal of Lgl from the plasma membrane in early mitosis in an NSC dissociated from the
central nervous system. Frames from Video 3 are shown as in (C).
(G) Removal of Lgl from the basal membrane. The time when basal Lgl membrane signal was
cleared (membrane to cytoplasmic signal was approximately one) relative to metaphase is
shown for three dissociated NSCs.
(H) Lgl reloading to the plasma membrane in early interphase in an NSC from the intact central
nervous system. Frames from Video 1 are shown as in (C).
(I) Lgl reloading to the plasma membrane in early interphase in an NSC dissociated from the
central nervous system. Frames from Video 3 are shown as in (C).
(J) Lgl patterned membrane reloading in early interphase. Lgl membrane enrichment relative to
the cytoplasm is shown for telophase, early interphase, and interphase measured at the
midbody, along with the lateral and apical membranes.
(K) Schematic of Lgl membrane dynamics in NSCs. Lgl is bound across the NSC membrane in
interphase and is removed starting at the apical pole in early mitosis until it is completely
removed in prophase. Membrane reloading begins in early interphase starting near the
midbody.
403
Figure 2 aPKC but not Aurora A is required for Lgl removal from the NSC membrane in early
mitosis

(A) Lgl dynamics in an NSC from the intact central nervous system expressing aPKC RNAi.
Frames from Video 4 are shown with Lgl-GFP and aPKC RNAi (both expressed from Worniu-
GAL4 driven UAS). Time is shown relative to nuclear envelope breakdown. The same
timepoints are shown from an NSC not expressing aPKC RNAi ("WT") for comparison.
(B) aPKC is required for Lgl removal from the membrane in early mitosis. The apical Lgl
membrane enrichment relative to the cytoplasm is shown for aPKC RNAi or WT NSCs in
interphase, prophase, and metaphase.
(C) Lgl dynamics in an NSC from the intact central nervous system incubated with the specific
Aurora A inhibitor MLN8237. Frames from Video 5 are shown with Lgl-GFP and aPKC RNAi
(both expressed from Worniu-GAL4 driven UAS). Arrowheads highlight furrow retraction in late
cytokinesis. Time is shown relative to anaphase onset. Similar timepoints are shown from an
NSC incubated with the solvent used for the inhibitor ("DMSO").
(D) No detectable effect of Aurora A inhibition on Lgl membrane displacement in early mitosis.
The apical Lgl membrane enrichment is shown for drug- or solvent-only-treated NSCs prophase
and metaphase.
421
Figure 3 Patterned Lgl removal from the apical and basal NSC membrane by apical aPKC
(A) Lgl and aPKC dynamics in an NSC from the intact central nervous system. Frames from
Video 6 are shown with aPKC-GFP (expressed from its native promoter) and Lgl-mCherry
(expressed from Worniu-GAL4 driven UAS). Time is shown relative to nuclear envelope
breakdown.
(B) aPKC is enriched at the apical membrane when Lgl is first cleared. The apical Lgl and aPKC
membrane enrichments relative to the cytoplasm are shown for three different NSC divisions.
429
Figure 4 Lgl regulates aPKC membrane removal at mitotic exit
(A) aPKC dynamics in an NSC from the intact central nervous system expressing Lgl RNAi.
Frames from Video 7 are shown with aPKC-GFP (expressed from its native promoter) from an
NSC expressing Lgl RNAi (from Worniu-GAL4 driven UAS). Arrowheads highlight basal aPKC
near mitotic entry. Time is shown relative to nuclear envelope breakdown. An NSC not
expressing Lgl RNAi ("WT") is shown at similar time points for comparison.
(B) Lgl is required to remove aPKC from the membrane in interphase. Comparison of apical
aPKC membrane enrichment between NSCs expressing and not expressing Lgl RNAi at
interphase, prophase, and metaphase.
(C) Lgl is required to remove aPKC from the basal membrane at metaphase. Comparison of
basal aPKC membrane enrichment between NSCs expressing and not expressing Lgl RNAi at
metaphase. Gardner-Altman estimation plot is shown for three different NSC divisions. The
error bars represent one standard deviation (gap is mean); the bar in the mean difference
comparison represents bootstrap 95% confidence interval.
(D) Lgl is not required for metaphase aPKC polarity. The ratio of apical to basal membrane
aPKC at metaphase is shown for three different NSC divisions. The error bars represent one
standard deviation (gap is mean).
(E) Model for Lgl function in NSC asymmetric division. Lgl plays a key role in early interphase by
removing the Par complex from the membrane. When mitosis begins Lgl's clearing of the Par
complex ensures that only apically targeted complex is present on the membrane allowing fate
determinants to polarize to the basal membrane. Without Lgl, the Par complex persists on the
membrane through interphase. During mitosis the Par complex is targeted to the apical
membrane as in wild type NSCs, but the presence of Lgl on the membrane before recruitment
leads to a polarized state where residual Par complex remains on the basal membrane prevents
fate determinants from binding the basal membrane.
455
456   **Video Legends**

Video 1: Lgl dynamics in NSCs

Super-resolution videos of in-tissue NSCs expressing UAS-Lgl-GFP (Lgl) and His2a-RFP
(His2a). Time relative to anaphase onset is indicated. 3 medical section videos of dividing in-
tissue NSCs are shown.

Video 2: Lgl and membrane dynamics in NSCs

Super-resolution videos of in-tissue NSCs expressing UAS-Lgl-GFP (Lgl) and the membrane
marker PH- PLCδ-mCherry (membrane). Time relative to anaphase onset is indicated. 3D max
intensity projections and medial sections are shown for 3 dividing NSCs.

Video 3: Lgl dynamics in dissociated NSCs

Super-resolution time-lapse videos of dissociated NSCs expressing UAS-Lgl-GFP (Lgl) and
His2a-RFP (His2a). Time relative to anaphase onset is indicated. 3 medial section videos of
dividing dissociated NSCs are shown.

Video 4 The effects of aPKC knockdown on Lgl dynamics

Super-resolution videos of in-tissue NSCs expressing UAS-Lgl-GFP (Lgl-GFP) and aPKC
RNAi. Time relative to anaphase onset is indicated. Medial sections are shown for 6 dividing
NSCs.

Video 5: The effects of Aurora A inhibition on Lgl dynamics

Part 1: Lgl dynamics in NSCs treated with Aurora A inhibitor. Super-resolution videos of in
tissue NSCs expressing UAS-Lgl-GFP (Lgl) and His2a-RFP (His2a). Time relative to anaphase
onset is indicated. Medial sections are shown for 3 dividing NSCs.

Part 2: Lgl dynamics in NSCs treated with DMSO. Super-resolution videos of in-tissue NSCs
expressing UAS-Lgl-GFP (Lgl) and His2a-RFP (His2a). Time relative to anaphase onset is
indicated. Medial sections are shown for 3 dividing NSCs.

Video 6: Dual imaging of Lgl and aPKC dynamics in NSCs NSCs expression aPKC-GFP and
Lgl-mCherry

Super-resolution videos of in-tissue NSCs expressing aPKC-GFP (aPKC) and Lgl-mCherry
(Lgl). Time relative to anaphase onset is indicated. Medial sections are shown for 3 dividing
NSCs.

Video 7: The effects of Lgl knockdown on aPKC dynamics

502 Part 1: aPKC dynamics in NSCs expressing Lgl-RNAi. Super-resolution videos of in-tissue
503 NSCs expressing BacaPKC-GFP (aPKC) and Lgl-RNAi. Time relative to anaphase onset is
504 indicated. Medial sections are shown for 3 dividing NSCs.
505
506 Part 2: aPKC dynamics in dividing NSCs (Wild-type control). Super-resolution videos of in-
507 tissue NSCs expressing BacaPKC-GFP (aPKC). Time relative to anaphase onset is indicated.
508 Medial sections are shown for 3 dividing NSCs.
509

References
1. Lang, C. F. & Munro, E. The PAR proteins: from molecular circuits to dynamic self-stabilizing cell polarity. *Dev.*

Camb. Engl. **144**, 3405–3416 (2017).

2. Venkei, Z. G. & Yamashita, Y. M. Emerging mechanisms of asymmetric stem cell division. *J. Cell Biol.* (2018)

doi:10.1083/jcb.201807037.

3. Sunchu, B. & Cabernard, C. Principles and mechanisms of asymmetric cell division. *Dev. Camb. Engl.* **147**,

dev167650 (2020).

4. Buckley, C. E. & St Johnston, D. Apical-basal polarity and the control of epithelial form and function. *Nat. Rev.*

Mol. Cell Biol. **23**, 559–577 (2022).

5. Bell, G. P., Fletcher, G. C., Brain, R. & Thompson, B. J. Aurora kinases phosphorylate Lgl to induce mitotic

spindle orientation in Drosophila epithelia. *Curr. Biol. CB* **25**, 61–68 (2015).

6. Carvalho, C. A., Moreira, S., Ventura, G., Sunkel, C. E. & Morais-de-Sá, E. Aurora A triggers Lgl cortical release

during symmetric division to control planar spindle orientation. *Curr. Biol. CB* **25**, 53–60 (2015).

7. Grifoni, D. *et al.* aPKCzeta cortical loading is associated with Lgl cytoplasmic release and tumor growth in

Drosophila and human epithelia. *Oncogene* **26**, 5960–5965 (2007).

8. Yamanaka, T. *et al.* Lgl mediates apical domain disassembly by suppressing the PAR-3-aPKC-PAR-6 complex

to orient apical membrane polarity. *J. Cell Sci.* **119**, 2107–2118 (2006).

9. Holly, R. W., Jones, K. & Prehoda, K. E. A Conserved PDZ-Binding Motif in aPKC Interacts with Par-3 and

Mediates Cortical Polarity. *Curr. Biol. CB* **30**, 893-898.e5 (2020).

10. Atwood, S. X., Chabu, C., Penkert, R. R., Doe, C. Q. & Prehoda, K. E. Cdc42 acts downstream of Bazooka to

regulate neuroblast polarity through Par-6 aPKC. *J. Cell Sci.* **120**, 3200–3206 (2007).

11. Hutterer, A., Betschinger, J., Petronczki, M. & Knoblich, J. A. Sequential Roles of Cdc42, Par-6, aPKC, and Lgl

in the Establishment of Epithelial Polarity during Drosophila Embryogenesis. **6**, 845–854 (2004).

12. Bailey, M. J. & Prehoda, K. E. Establishment of Par-Polarized Cortical Domains via Phosphoregulated

Membrane Motifs. *Dev. Cell* **35**, 199–210 (2015).

13. Dong, W. *et al.* A conserved polybasic domain mediates plasma membrane targeting of Lgl and its regulation

by hypoxia. *J. Cell Biol.* **211**, 273–286 (2015).

14. Peng, C. Y., Manning, L., Albertson, R. & Doe, C. Q. The tumour-suppressor genes lgl and dlg regulate basal

protein targeting in Drosophila neuroblasts. *Nature* **408**, 596–600 (2000).

15. Wirtz-Peitz, F., Nishimura, T. & Knoblich, J. A. Linking cell cycle to asymmetric division: Aurora-A

phosphorylates the Par complex to regulate Numb localization. *Cell* **135**, 161–173 (2008).

16. Prehoda, K. E. Polarization of Drosophila neuroblasts during asymmetric division. *Cold Spring Harb. Perspect.*

Biol. **1**, a001388 (2009).

17. Knoblich, J. A. Asymmetric cell division: recent developments and their implications for tumour biology. *Nat.*

Rev. Mol. Cell Biol. **11**, 849–860 (2010).

18. LaFoya, B. & Prehoda, K. E. Actin-dependent membrane polarization reveals the mechanical nature of the

neuroblast polarity cycle. *Cell Rep.* **35**, 109146 (2021).

19. LaFoya, B. & Prehoda, K. E. Consumption of a polarized membrane reservoir drives asymmetric membrane

expansion during the unequal divisions of neural stem cells. *Dev. Cell* S1534-5807(23)00159–4 (2023)

doi:10.1016/j.devcel.2023.04.006.

20. Penkert, R. R. *et al.* The Drosophila neuroblast polarity cycle at a glance. *J. Cell Sci.* **137**, jcs261789 (2024).

21. Moreira, S. *et al.* PP1-Mediated Dephosphorylation of Lgl Controls Apical-basal Polarity. *Cell Rep.* **26**, 293-

301.e7 (2019).

22. Oon, C. H. & Prehoda, K. E. Asymmetric recruitment and actin-dependent cortical flows drive the neuroblast

polarity cycle. *eLife* **8**, (2019).

23. Oon, C. H. & Prehoda, K. E. Phases of cortical actomyosin dynamics coupled to the neuroblast polarity cycle.

eLife **10**, e66574 (2021).

24. Lee, C.-Y., Robinson, K. J. & Doe, C. Q. Lgl, Pins and aPKC regulate neuroblast self-renewal versus

differentiation. *Nature* **439**, 594–598 (2006).

25. Besson, C. *et al.* Planar Cell Polarity Breaks the Symmetry of PAR Protein Distribution prior to Mitosis in

Drosophila Sensory Organ Precursor Cells. *Curr. Biol. CB* **25**, 1104–1110 (2015).

26. Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond P values: data analysis with

estimation graphics. *Nat. Methods* **16**, 565–566 (2019).

A Mutual exclusion model for Par polarity

interphase / mitosis

epithelial cell
Par complex
Lgl

stem cell

B
Drosophila larva → larval central nervous system → image CNS neural stem cells (NSCs) → dissociate tissue → image cultured neural stem cells (NSCs)

D
Lgl Membrane:cytoplasm at initial apical clearing
apical N = 10 / lateral N = 10
Paired mean difference
lat minus apical

G
Basal clearing time relative to metaphase (sec)
= 1

neural stem cells imaged in intact central nervous system

C Lgl
prophase −19:00 / −6:00 | metaphase 5:00 | anaphase 8:00
5µm *
Lgl His2a

H Lgl
telophase 0:00 | interphase 6:00 / 10:00 / 29:00
5µm
Lgl His2a

E Lgl medial
interphase −19:00 | prophase −7:00 / 0:00 | metaphase 6:00 | anaphase 15:00
Lgl MIP

neural stem cells imaged after dissociation

F Lgl
prophase −17:20 / −4:00 | metaphase 7:00 | anaphase 10:20
5µm
Lgl His2a

I Lgl
telophase 12:00 / 15:00 | interphase 18:00 / 26:20
5µm
Lgl His2a

J
Lgl Membrane:Cytoplasmic
Midbody / Lateral / Apical
telophase | Early interphase | interphase

K NSC Lgl membrane removal (early mitosis)

NSC Lgl membrane reloading (early interphase)

A

| interphase | prophase | metaphase | anaphase | interphase |

Lgl RNAi — aPKC

-15:00   -7:00   0:00   5:00   8:00   10:00

5μm

WT — aPKC

-15:00   -7:00   0:00   5:00   9:00   10:00

5μm

B

Apical Lgl
membrane:cytoplasmic

- Lgl-RNAi
- WT

Interphase   Prophase   Metaphase

C

Basal aPKC at metaphase
membrane:cytoplasmic

Lgl-RNAi   WT   WT minus Lgl-RNAi

mean difference

D

aPKC polarity at metaphase
apical:basal

Lgl-RNAi   WT

E

interphase   mitosis

WT NSC

Lgl — Par complex — fate determinants

Par complex removed early interphase

Par complex recruited apically

fate determinants polarized

Lgl⁻ NSC

Par complex **NOT** removed early interphase

Par complex recruited apically

fate determinants cytoplasmic