Thrifty phenotypes in ants: Extending a human developmental hypothesis to a superorganism

Érik Plante[1,2], Ehab Abouheif[2,3]* & Jean-Philippe Lessard[1]*

1. Department of Biology, Concordia University, Montreal, Quebec, Canada.
2. Department of Biology, McGill University, Montreal, Quebec, Canada.
3. Centre of Evolutionary & Organismal Biology, Women's Hospital, School of Medicine, Zhejiang University, Hangzhou, China.

* abouheif@zju.edu.cn (E.A.); jp.lessard@concordia.ca (J.-P.L.)

Competing interests:
We declare we have no competing interests.

Funding:
This research was supported by Natural Sciences and Engineering Research Council Discovery Grant to J.-P.L. (RGPIN-2022-04716) and E.A (RGPIN-2020-05175). É.P. was supported by the Concordia University Faculty of Arts and Science Graduate Doctoral Award and the Quebec Centre for Biodiversity Science Excellence Awards. The Centre for Northern Studies also provided support through access to its research infrastructure. Additional support was provided by the Wladimir A. Smirnoff Fellowship, a joint initiative of the Entomological Society of Quebec, the Quebec Society for the Protection of Plants, and Natural Resources Canada – Canadian Forest Service.

ABSTRACT

Adaptation to extreme environments is commonly assumed to occur through microevolutionary

change. However, studies of high-altitude human populations show that organisms can also

respond to nutritional stress through developmental plasticity, producing thrifty phenotypes that

prioritize essential function over costly structures. Whether this adaptive strategy extends beyond

humans, and how it operates in natural systems, remains largely unexplored. Here, we test the

thrifty phenotype hypothesis (TPH) in a widely distributed superorganism, the carpenter ant

Camponotus herculeanus, which inhabits some of the most environmentally challenging regions

of the Northern Hemisphere. Colonies comprise inexpensive minor workers and energetically

costly major workers, providing a powerful system for examining plastic investment under

resource limitation. We quantified relationships between caste structure and climate and

conducted a common-garden experiment to test the TPH. We show that the proportion of major

workers declines with increasing latitude, independently of body size and colony size, and is best

predicted by the number of days, annually, during which workers can nurse brood. Experimental

results further demonstrate that colonies rapidly and plastically adjust caste structure in response

to environmental conditions. Our findings reveal that thrifty phenotypes can emerge in

superorganisms and could represent a conserved developmental response to environmental stress.

By extending a central hypothesis from human biology to social insects, this work provides a

unifying framework for understanding how developmental plasticity shapes adaptation across

levels of biological organization.

INTRODUCTION

Adaptation can be challenging in stressful environments where both energy and key

nutrients are chronically limited. The evolutionary and developmental processes enabling

organisms to adapt to stressful environments remain poorly understood [1–4]. Organisms

may respond through evolutionary changes [5,6], phenotypic plasticity [7–9], or a

combination of both. Populations of humans and other organisms inhabiting high altitude

mountains or polar latitudes have been the focus of research on adaptation to extremely

stressful environments[10–14]. Studies on high-altitude human populations, such as those

of the Andean and Himalayan mountains [15,16], have revealed that nutritional stress

during development can induce plastic adjustments consistent with thrifty phenotypes

[17–20]. The *thrifty phenotype hypothesis* (TPH) [17] posits that nutrient shortages

during development lead to selective investment in organs, where the growth of critical

organs (e.g. heart, brain) is preserved at the expense of peripheral growth, resulting in

shortened limbs (e.g., reduced ulna and limb length) relative to overall body size

[15,16,21]. However, it remains unclear whether non-human organisms evolve thrifty

phenotypes to adapt to stressful environments, and which external factors may drive their

expression [22,23].

The TPH was originally proposed in medical literature to explain type 2 diabetes

through the lens of developmental plasticity [17]. Malnutrition during fetal development

can alter growth patterns and metabolism [17,24–26,9,27,28,18,29–31]. When such

individuals experience abundant nutrition later in life, the thrifty phenotype may become

maladaptive, increasing the risk of type 2 diabetes and other health problems, including

obesity and cardiovascular disease [17,24,25,9,27–31]. However, when environmental

conditions are persistently stressful throughout an individual's entire lifetime, such as at

high altitudes or latitudes, then thrifty phenotypes are predicted to provide an adaptive

advantage. In such stressful environments, individuals must store sufficient energy during

the growing season to survive long winters [22]. However, identifying environmental

drivers of thrifty phenotypes in humans is complicated by long developmental periods

and cultural buffering that obscure environmental effects [32].

The TPH highlights the existence of a sensitive period or window during

development, during which environmental conditions are sensed, and organisms

plastically adjust organ and tissue growth to match the perceived environment [22].

Adaptive developmental plasticity is a core principle of the field of ecological

evolutionary developmental biology (eco-evo-devo) [33,34]. Although the TPH has been

tested only in humans, eco-evo-devo predicts that such developmentally sensitive periods

may be more conserved than might be expected, and may therefore apply more broadly

across animals [35]. This prediction is based on the existence of a highly conserved

developmental toolkit that is shared across metazoan animals, in which a single genome

produces differentiated cell types, tissues, and organs in response to internal and external

cues [36,37]. Extending the TPH beyond humans therefore predicts the evolution of a

sensitive developmental window across animals that detects nutrient limitation and

reallocates resources toward essential organs at the expense of others [38,39]. Thus, in

eco-evo-devo terms, the TPH can be reframed as the expression of conserved

developmental windows through which external cues canalize growth trajectories,

generating thrifty phenotypes across a wider range of organisms [40].

Eusocial colonies provide a natural system to test the TPH, because a single

genome produces morphologically distinct individuals or castes in response to

environmental cues [41–43], making colonies superorganisms with traits analogous to

organ-level traits in multicellular organisms [44,45]. Moreover, colony-level phenotypes,

including caste structure, are directly observable [38,39]. Colony size (number of

individuals) and colony structure (caste, age, or size distribution) are two phenotypic

traits with adaptive value [46–48] that vary geographically [49–51], and respond to

resource availability [41]. Although most species have a monomorphic worker caste,

meaning that workers show little variation in size or head-to-body scaling, several species

have evolved worker polymorphism, in which workers show dramatic variation in size

and head-to-body scaling [51]. Worker polymorphism is therefore particularly

informative because it reflects developmental allocation decisions in response to

environmental conditions [50,51].

Broadly distributed species may encounter extremely stressful conditions at the

edge of their range, imposing selective pressures on those edge populations that are

distinct from those of core populations [52–55]. We therefore test the *thrifty phenotype*

hypothesis (TPH) by investigating phenotypic variation among populations of a widely

distributed superorganism, the carpenter ant *Camponotus herculeanus* (Linnaeus, 1758),

whose northern range limit occurs in extremely stressful environmental conditions across

the northern hemisphere. Eusocial colonies of *C. herculeanus* have a worker caste

consisting of thousands of small-headed minor workers specializing in nursing and

foraging, and larger-headed, more energetically costly, major workers specializing in

defense and food storage [56]. Subcaste identity is determined by a developmental switch

mediated by Juvenile Hormone (JH) during the last larval instar [57,58]. Variation in the

proportion of majors and minors results from this developmental reprogramming,

reflecting larval allocation decisions and can be influenced by environmental conditions.

The TPH, when applied to eusocial organisms, predicts that colonies experiencing

resource limitation should maintain overall colony size by plastically reducing investment

in costly individuals, such as majors.

The TPH has been tested on human populations and via lab experiments on

rodents [59,60]. Here, we performed the first test of TPH in natural systems outside

humans by using whole-colony sampling across the temperate and boreal forest biomes

of eastern Canada—by felling nesting trees and collecting all individuals—and relating

colony worker caste structure to variation in micro- and macro-climate. Specifically, we

collected 26 whole colonies of *C. herculeanus* distributed along an 8° latitudinal gradient

from temperate forest to the northern edge of the boreal forest in eastern Canada (figure

2). This range encompasses both the core and northern edge of the species' distribution.

We quantified colony size, worker body size and caste structure, and estimated local

environmental variables at each colony. To determine whether caste structure is

evolutionarily fixed or plastic, five colonies from the core and four colonies from the

northern edge were transplanted into a common rearing environment and assessed after

one larval generation. Together, these approaches provide the first comprehensive test of

the TPH in a non-vertebrate, offering insights into how developmental plasticity shapes

colony-level adaptation under extreme environments.

RESULTS

The TPH predicts a reduced investment in costly major workers (i.e., reduced subcaste

ratio) in stressful environments. Our data support the TPH, such that colonies in colder,

drier environments invested less in major workers (electronic supplementary material,

figure S1, table S4), whereas mean major worker body (electronic supplementary

material, figure S1, table S4) and colony size (R^2_{adj} = -0.04, df = 24, $F < 0.01$, $p = 0.983$)

were not related to regional climate, indicating that colonies maintain consistent body and

colony size regardless of environmental stress.

To test whether the thrifty colony phenotype arises from developmental plasticity

or from microevolutionary adaptation, we reared colonies from the southern and northern

ends of the gradient for one generation in a common controlled environment and

compared caste structure before and after rearing. Microevolutionary adaptation would

predict that colony traits are genetically fixed and remain unchanged under uniform

conditions.

After one generation, southern colonies showed no significant change in caste

structure ($t = -0.31$, df = 4.89, $p = 0.77$; figure 5a), whereas northern colonies showed a

significant increase in the proportion of major workers ($t = -2.82$, df = 4.14, $p = 0.045$;

figure 5a). These results indicate that the thrifty colony phenotype in northern

environments is driven by developmental plasticity. Moreover, after rearing, northern and

southern colonies converged in caste structure ($t = 0.95$, df = 4.06, $p = 0.40$), indicating

plastic adjustment under uniform environmental conditions. In contrast, mean major

worker body size did not change in either southern ($t = -0.99$, df = 7.57, $p = 0.35$; figure

5b) or northern colonies ($t = 0.06$, df $= 4.10$, $p = 0.95$; figure *5b*), consistent with the

earlier results that body size is not environmentally plastic.

Having established that the thrifty phenotype arises from plasticity, we next

examined which environmental factors best predict caste structure. Proportion of major

workers was most strongly associated with regional climate, consistent with TPH

predictions. Colonies invested less in costly major workers in colder, drier environments

(figure 3; electronic supplementary material, table S4). While colony size did not vary

with climate, it contributed to predicting other colony-level traits. Mean major worker

body size was positively related to colony size (electronic supplementary material, figure

S1, table S4). Worker body size variance increases with colony size, whereas head size

variance was negatively associated with climate and positively with colony size. Head-to-

body ratio variance showed a weak relationship with regional climate only.

We further tested which temperature factor best predicted the major worker

proportion using separate GAMs. The best fit model included the temperature threshold

for colony activity, measured as the number of days above $0°$, and colony size (electronic

supplementary material, table S5). Both predictors showed significant relationships with

the proportion of major workers (figure 4a). The second-best model included the

temperature threshold for larval development, measured as the number of days above $9°C$

and colony size (figure 4b; electronic supplementary material, table S5).

Finally, we explored whether colony size is influenced by environmental factors

other than climate. Across 26 colonies, colony size ranged from 151 to 6,557 workers

(mean $\pm$ SE $= 1,809 \pm 299$). Most colonies nested in black spruce (*Picea mariana* (Mill.)

BSP; ~78%), with others found in balsam fir (*Abies balsamea* (L.) Mill.; ~7%), eastern

white cedar (*Thuja occidentalis* L.; ~7%), eastern white pine (*Pinus strobus* L.; ~4%),

and tamarack (*Larix laricina* (Du Roi) K. Koch; ~4%). Colony size was positively related

to nesting tree diameter (R^2_{adj} = 0.29, df = 23, F = 10.86, p = 0.003) and negatively related

to the presence of strong competitors, *Formica* spp. (t = 2.77, df = 19.66, p = 0.01;

electronic supplementary material, figure S3). No significant effects were observed for

the presence of *Myrmica* (t = -1.16, df = 18.26, p = 0.26; electronic supplementary

material, figure S3) or *Leptothorax* spp. (t = 1.01, df = 23.52, p = 0.32; electronic

supplementary material, figure S3). For additional results on competition estimation, see

electronic supplementary material, S5.

DISCUSSION

Our results support the TPH, which provides a general mechanism by which

(super)organisms adapt to extremely stressful environments. We show that eusocial

colonies of a geographically widespread ant species consistently reduce their investment

in costly major workers when confronted with short growing seasons and most likely,

nutritional shortage. Moreover, colonies converged in caste structure after one generation

in laboratory settings, under common environmental conditions, which shows that caste

structure is a plastic colony-level trait. Overall, our study lends support to the hypothesis

that organisms experiencing nutritional shortage in early life stages can plastically

reprogram development to rapidly adapt their phenotype to ambient condition.

The thrifty phenotype as an adaptive response to energetic constraints

The reduction in major worker proportion we observed in northern latitudes, where the

growing season is short and food resource are scarce, aligns with the idea that those

stressful conditions experienced early during larval development favor energy-saving

strategies. Moreover, the absence of relationship between proportion of majors and

colony size indicates a reallocation of resources within colonies rather than a general

reduction in whole-colony productivity. Given that large colonies have lower per-capita

metabolic demand than small colonies [61,62], and that major workers consume more

energy per unit of biomass than minor workers [63], energetic shortage in northern

colonies could lead to a strategy where colonies maintain their size but reduce the

production of major workers. Producing fewer major workers may therefore help

colonies maintain overall size while conserving energy, much like individual organisms

reduce allocation to costly organs under nutritional stress, which is consistent with

patterns observed in vertebrate models of the thrifty phenotype [38,39].

Our study provides insight into the environmental factors that could cause a

restructuring of the worker caste. We expected that micro-climate could limit the

production of majors by either limiting the amount or quality of resources available to

larvae [41,64,65] or limiting the time for development [66,67]. The number of days with

temperatures above 0°C was the best predictor of geographic variation in caste structure,

indicating that constraints on worker activity, rather than duration of larval development

per se, most likely limit the production of energetically costly majors. Thermal

constraints on worker activity could limit the amount of time that workers devote to

foraging and nursing larvae yearly, possibly reducing the amount or quality of food

received by larvae [68–73]. In sum, these results further support the role of nutrient

shortage in shaping whole-colony phenotype, possibly via developmental plasticity.

The TPH posits that reduced investment in costly structures is plastic and occurs

during development. Consistent with this idea, colonies collected from northern latitudes

where foraging and nursing activity of ant workers is limited by the short growing season,

produced more majors when reared under warmer and stable controlled conditions with

abundant resources, whereas southern colonies produced fewer majors under the same

conditions. After one generation under controlled conditions, colonies from different

climatic origins converged in caste structure, demonstrating that the reduced investment

in major workers observed in the field is environmentally induced rather than genetically

fixed. In contrast, body size remained unchanged after rearing, supporting the idea that it

is developmentally canalized or under stronger genetic control, and thus decoupled from

variation in major worker investment [74]. These results contrast with previous work

where population-level differences in worker caste structure across climatic gradients

were largely genetically determined [50], possibly due to the much less stressful

conditions experienced by ants in this study conducted at more southern latitudes. In sum,

our common garden experiment supports the view that organisms can rapidly change

their phenotype to match ambient conditions via plastic developmental changes rather

than microevolution. Taken together, studies on colony-level adaptation suggests that the

balance between plastic and genetic mechanisms regulating worker polymorphism may

be evolutionary conserved at the genus level [75–79], with *Camponotus* generally

favoring plastic responses to environmental variation [58,64,80–82]. More broadly, our

results highlight that adaptive demography in ants can rely heavily on plasticity when

environmental predictability is low, and that the relative contribution of plastic versus

genetically fixed mechanisms may differ among taxa and environmental contexts.

The thrifty phenotype as a conserved evolutionary response to environmental stress

Environmentally induced worker caste adjustment could reflect a general and

evolutionary conserved mechanism by which organisms rapidly adapt to environmental

stress via developmental plasticity [83]. It has been suggested that extremely stressful

environments could shape phenotype owing to environmentally sensitive developmental

windows during which fine-tuning of developmental pathways to cope with current or

anticipated conditions can occur [84]. This view is consistent with evolutionary

conserved developmental windows documented in vertebrates and other taxa, where

environmental cues, like temperature or nutrient availability, modify developmental

trajectories [85,86]. In social insects, and ants in particular, caste differentiation is known

to be influenced by temperature, nutrition and hormonal switches [41,57], suggesting that

similar conserved developmental mechanisms could underlie the plastic response we

observed.

While most studies of the TPH have focused on humans, parallels exist in other

model organisms, including rodents. For example, experimental work in mice shows that

early-life nutritional restriction results in altered metabolic and growth profiles consistent

with a thrifty strategy [60]. In rats, maternal undernutrition induces a clearly defined

"thrifty liver programme" characterized by reduced relative liver growth, inflammatory

and transcriptomic changes interpreted explicitly within the thrifty phenotype framework

[59]. These examples, alongside our findings in natural ant colonies, suggest that the

thrifty phenotype may reflect a broad and evolutionarily conserved response to

environmental unpredictability across both solitary and superorganisms.

The thrifty phenotype as an adaptation to unpredictable environments

The thrifty colony phenotype could therefore not only be an energy-saving plastic

adaptation, but also as a finely tuned developmental strategy to cope with unpredictable

resource availability [4]. The thrifty colony phenotype in *C. herculeanus* illustrates how

colonies can adjust caste allocation in a time-sensitive, predictive manner (Figure 6).

Bipotent larvae enter diapause during fall, then overwinter, and resume development in

spring, when early-spring nutritional conditions possibly provide the critical cue for caste

reprogramming. This cue possibly determines whether larvae develop into costly major

workers or less expensive minors, effectively predicting the resource environment for the

coming active season [58,82]. In populations inhabiting northern latitudes, persistent

nutrient scarcity at this early life stage could reprogram larval development to produce

fewer majors, generating a thrifty colony phenotype well-suited to short, resource-limited

summers. However, if spring cues indicate scarcity while nutritional resources turn out to

be abundant during summer, colonies may be maladapted. That is, their low proportion of

majors could limit their ability to exploit abundant resources and could even increase

mortality due to competition with other nearby colonies. Conversely, when spring cues

signal abundant resources, larvae develop more majors, which is advantageous if

conditions remain favorable but potentially costly to maintain if resources decline later in

the season. The thrifty colony phenotype could therefore be a predictive developmental

adjustment, with fitness payoffs dependant on the reliability of environmental cues across

life stages.

Environmental cue–response systems are predicted by models of predictive

adaptive plasticity and seasonal polyphenism, in which developmental trajectories are

adjusted based on early-life environmental signals to optimize performance in anticipated

conditions [87]. When environmental predictability is low, these same systems can

produce mismatch costs, a form of developmental bet-hedging that may help explain

variation in colony success at the northern range edge of the species, where

environmental conditions are expected to be extremely stressful [88].

Building on previous theory about colony-level adaptation

Our findings further shed new light on the validity of one of the key hypotheses proposed

to explain geographic variation in body and colony size in eusocial organisms. The

starvation resistance hypothesis predicts that colonies experiencing long winters should

rely on larger workers and/or larger colonies to store fat and nutrients, thereby buffering

winter mortality [89,90]. In our study, however, neither body nor colony size varied with

climate. Only caste structure, namely the proportion of major workers produced by

colonies, co-varied with climatic conditions. This contrasts with Yang et al. (2004), who

found that in colonies of *Pheidole morrisii* Forel, 1886, the proportion of majors declined

in more seasonal climates, but that the average size of majors increased [50]; in those

populations, majors in seasonal climates often functioned as repletes for fat storage [91].

Together, these studies suggest that both the *starvation resistance hypothesis*- and the

TPH-related patterns can occur depending on the species and ecological context, whereas

our results are more consistent with the TPH.

Conclusions

Our study presents the results of the first empirical test of the TPH performed in natural

systems outside humans. The findings of developmentally plastic thrifty phenotypes in

superorganisms inhabiting northern latitudes could reflect both energetic constraints and

adaptive, predictive responses to environmental stress, similar to what is observed in

humans, supports the view that the thrifty phenotype strategy is evolutionary conserved

between insects and humans [23]. The potentially conserved developmental plasticity

observed here in a superorganism may offer a powerful model to study how stress-

responsive phenotypes evolve in complex, long-lived organisms, including humans [28].

METHODS

Study sites

The study took place in the province of Quebec, Canada (figure 2). Study sites spanned

from temperate deciduous forest of southern to the northern limit of the boreal forest

(aka, tree line). The southern sites were in the Mauricie (46.6°N, -72.5°W) and National

Capital (46.8°N, -71.5°W) regions whereas the northern sites were above 52°N in

Northern Quebec region. Along this environmental gradient, mean annual temperature

ranges from 3.93°C to -2.79°C, minimum annual temperature of coldest month ranges

from -18.10°C to -28.70°C, and annual precipitation ranges from 1383 mm to 572 mm

[92]. The growing season are approximately 140 to 150 days in the south and 100 to 110

days in the north [93].

Study organism

We used *C. herculeanus* (figure 1) as a study organism because it is the most broadly

distributed, polymorphic eusocial species in eastern Canada for which it is possible to

collect entire colonies. *C. herculeanus* is widely distributed throughout the Holarctic

realm [94]. In the Nearctic region, entire colonies of this species can often be found

inside a single tree, stump or log and frequently occur in peat bogs [95]. It builds its nest

by digging chambers inside dead or live wood. In addition, since peat bogs are frequently

flooded, the colonies rarely ramify into very complex tunnel systems or satellite nests in

these habitats, thus allowing the complete collecting of the colony. *C. herculeanus* is a

monogynous (single egg-laying queen), slow-growing species and the most cold-tolerant

ant species in the world [95,96]. Based on Wilson's classification [56], *C. herculeanus*

exhibits continuous worker polymorphism, specifically triphasic allometry, producing

clear minor and major worker subcastes (in line with recent evidence in *Camponotus*

[58]). Individuals falling between these endpoints (intermediate workers or medias) are

best interpreted as vestigial products of developmental threshold slack rather than a

distinct subcaste. For details on identification of the major worker subcaste, see electronic

supplementary material, S1.

Colony collecting for observational study

For the observational study aiming to document several colony-level traits, we collected

26 colonies in peat bogs along the environmental gradient. We only collected colonies

including at least 150 workers. Colonies with less that 150 workers were not collected

because they are considered incipient and rarely contain major workers [97]. When

excavating the colony, we collected all workers, broods and alates we observed. Colonies

were collected at least 1km apart. We collected the northern colonies (~52°N to ~54°N)

from mid-July to early August, and southern colonies (~46°N to ~52°N) in both mid-July

and early August of summer 2021. These dates were selected as to not bias our estimation

of colony traits because of winter mortality. They correspond with periods when the

yearly generation was hatched in the colonies [98,99]. For more details on colony

collecting, see electronic supplementary material, S2.

Colony collecting for common garden experiment

To test whether the variation in the caste structure along the gradient is due to

microevolution or developmental plasticity, we collected nine entire colonies along the

same gradient and using the same methodology as in the observational study. In 2022,

we collected five colonies in the southern sites (September 16th to 25th), and four in the

northern sites (August 4th to 9th). Colony size was estimated visually by counting the

number of occupied plaster nests as to not stress or hurt workers. The average colony size

was similar (1,667 ±333 (*SE*) workers) to the average size of all colonies we ever

collected along this gradient (1,772 ±299 (*SE*) workers) (includes unpublished

observations).

Estimating regional climate

To estimate variation in regional climate, we extracted several georeferenced bioclimatic

variables that could possibly influence worker polymorphism. Bioclimatic variables were

extracted for each site where a colony was collected using WorldClim data at 1 km^2

spatial resolution, which included mean annual temperature, mean diurnal range,

isothermality, temperature seasonality, maximum temperature of warmest month,

minimum temperature of coldest month, temperature annual range, and annual

precipitation [92,100]. We performed a principal component analysis (PCA) on the eight

bioclimatic variables listed above to reduce the dimensionality of the data and collinearity

[101]. We extracted the first axis, which captured 76% of the variance (electronic

supplementary material, table S1).

Estimating local conditions

The size of nesting spaces can impose limits on colony size in carpenter ants and possibly

the degree of worker polymorphism [102,103]. We therefore measured nesting tree

diameter at the base of the tree trunk. We also estimated canopy openness using a

spherical densiometer (Forest Densiometers, Model-C, Bartlesville, Oklahoma, USA) as

it relates to the amount of light reaching the nest, thereby altering its temperature and the

amount of energy and food resources available. In turn, temperature can influence larval

growth rate [104] and whether some larvae might pupate or enter diapause [105], which

could influence worker polymorphism and colony size in *C. herculeanus* colonies.

Estimating local temperature

To estimate the temperature regime experienced by colonies, we used Thermocron

iButton DS1921G-F5# integrated thermometer and iButton data-logger (Maxim

Integrated, San Jose, California, USA). They were deployed at 18 of the 26 colonies

collected in 2021 due to resource and logistical limitations. Temperature was measured

every 255 minutes (maximum length for the data-logger model) for 12 months. We then

estimated the number of days during which brood development is possible (above 9°C)

and the number of days during which worker activity is possible (above 0°C) (for more

details, see electronic supplementary material, S3).

Estimating the intensity of competition

The presence of competitive ant species near *C. herculeanus* colonies could influence

major worker production, as major workers can have a role for defense of the colony

[98]. We assessed the presence of potential interspecific ant competitors by active visual

observation and opening wood debris within a radius of ten meters around each colony

for 15 minutes between 10:00AM and 5:00PM on days without precipitations. We

sampled workers of all species found with a mouth aspirator or manually. Species

identifications were confirmed under a microscope by a trained ant taxonomist.

Morphometric measurements

To estimate colony-level traits, we took morphometric measurements on 200 randomly

sampled workers in each colony, except for one colony comprising a total of 151

workers, for which we measured all the workers. We froze collected colonies within five

days of collecting to avoid influencing the proportion of major workers or any other

colony-level traits. To avoid selection bias, we created all the subsets on the same day

using the same method by mixing all the workers of each colony together in a plastic

container with ethanol for two minutes and randomly collecting 200 workers using

forceps.

We took photos of morphological traits using a digital microscope (Brand:

Koolertron, Model: AD106S). We digitalized the dorsal and lateral views of each

specimen and obtained morphometric measurements using an image processing software

(ImageJ version 1.53e). For each worker, we measured head width (maximum width of

head including eyes) and body size (Weber's length, i.e. maximum diagonal length of

mesosoma) (electronic supplementary material, figure S4).

Measuring colony-level traits

We first estimated colony size by counting each collected worker using forceps and a

tally counter. We then took morphometric measurements to quantify five different

colony-level traits describing worker polymorphism, including the proportion of major

workers, mean major worker body size, worker body size variance, worker head size

variance and worker head-to-body ratio variance. We quantified five different traits

related to worker polymorphism because each of them related to different functions for

colony adaptation (table 1). We calculated the mean size of major workers as the

average of body size measurements (i.e., Weber's length) of major workers in the entire

colony. We calculated the variance of worker body size (i.e. Weber's length), head size,

and the ratio between head width and worker body size for each of the sampled ant

colonies. We used variance because it is the parameter that best captured the variation

of each trait between the colonies (electronic supplementary material, table S3) and

represents how large the variation of worker morphology is in comparison to the other

workers of a same colony.

Common garden experiment

We ran the common garden experiment from August 10^{th}, 2022, to June 21^{st}, 2023. For

detailed information on the experimental conditions, see electronic supplementary

material, S4. We focused this experiment on the proportion of major workers as a

response variable because it was the only colony-level trait that varied independently

from colony size and was predicted by an environmental variable. To obtain a

representative estimation of the proportions of major workers without harming the

colonies, we randomly selected 200 workers in each colony upon collection in nature.

To do so, for each colony, we placed all adult workers in a same plastic box and

randomly sampled them using forceps while not looking directly at the sample and

shaking the box every 30 seconds to avoid worker clustering. This procedure ensured

that our subsample was not biased toward a particular subcaste that tend to cluster or

that might be inadvertently chosen when sampling visually [106]. Then, to compare the

colony phenotypes, we waited until brood develops into adulthood after diapause in a

common rearing environment and randomly resampled 200 workers using the same

method after one larval generation.

Statistical analyses

Environmental predictors of colony-level traits

To select the variables to be used in a multivariate normal additive model relating worker

polymorphism to environmental factors, we ran separate generalized additive models

(GAMs) for each of the five colony-level traits [107]. We included the predictor variables

of regional climate, local climate, nesting tree diameter, colony size and competition (i.e.,

presence of competitor). After fitting a separate GAM for each trait, only the significant

predictor variables were kept for the multivariate normal GAM.

The multivariate normal GAM fits a separate regression for each response

variable, as a covariance matrix estimating the covariance between response residuals.

The output includes a separate linear predictor for the mean for each response and the

covariance matrix. The conditional mean in a GAM is estimated as a linear combination

of nonlinear basis functions, that form a smooth curve. GAMs are adequate for selecting

variables because p-values can be used for testing each variable for equality to zero for

deciding on which candidate variables to be removed from the model.

Multivariate normal GAMs can be used to assess the relationship between many

variables and are robust to autocorrelation. Using only the selected variables, we fitted a

single multivariate GAM [108]. To fit this multivariate model, we used the five colony-

level traits as response variables representative of ant worker polymorphism (table 1). We

used "colony size" and "regional climate" as fixed effect predictor variables for each

response because they were the significant predictors in the previous approach using

separate GAMs for each colony-level trait. We used linear predictors instead of

smoothing terms based on the model linear relationships between response and predictors

residuals. Finally, we assessed the relationship between each colony trait and the

predictors, and we plotted the partial effects as a visual aid for the direction of the

relationships (electronic supplementary material, figure S6). We fitted the model using

the *gam* function in the *mgcv* package [109].

To test which temperature factor best predict the proportion of major workers, we

ran four separate GAMs with the proportion of major workers and colony size as

response variables and the number of days above 0 °C, the number of days above 9°C,

the regional mean annual temperature (°C) and the regional mean diurnal range (°C) as

predictors. Then, we determined which was the best-fitting model based on the lowest

corrected Akaike's information criteria (AICc) value, adjusted for small sample size. We

calculated AICc values using the *AICc* function in the *AICcmodavg* package [110].

Environmental predictors of colony size

To assess the relationship between colony size and the environmental variables

(predictors) we used a linear model using the *lm* function. We performed Welch two-

sample t tests to test whether colony size differed in the presence of certain ant genera

using the t.test function.

Plasticity of caste structure

During the common garden experiment, to test whether the proportion of major workers

or mean major worker body size significantly changed from the moment of collection to

after one larval generation in a common rearing environment, we performed Welch two-

sample t tests for each group (North and South) using the *t.test* function. We also tested

whether the proportion of major workers in both groups were significantly different from

each other using the same method. All statistical analyzes were performed in R version

4.2.0 [111].

ACKNOWLEDGEMENTS

We are especially grateful to D. Ouellette (McGill University) for his assistance during

fieldwork and for mounting specimens in figure 1. We also thank E.J. Pedersen, E.

Despland, and G. Muñoz (Concordia University) for their advice on analytical

approaches, as well as P.-M. Brousseau (Concordia University) and members of the

Abouheif Lab (McGill University) for their helpful comments on the manuscript. We also

thank K. Marshall (University of British Columbia) and R. Khelifa (Concordia

University) for valuable discussions. We acknowledge the Centre for Northern Studies for

providing access to research infrastructure and the McGill University Phytotron facility

for support of the common garden experiment.

REFERENCES

1.   Hoffmann AA, Parsons PA. Evolutionary genetics and environmental stress.

Oxford University Press; 1991.

2.   Chevin L-M, Hoffmann AA. Evolution of phenotypic plasticity in extreme

environments. Philosophical Transactions of the Royal Society B: Biological

Sciences. 2017;372: 20160138. doi:10.1098/rstb.2016.0138

3.   Fitt RN, Palmer S, Hand C, Travis JM, Lancaster LT. Towards an interactive,

process-based approach to understanding range shifts: developmental and

environmental dependencies matter. Ecography. 2019;42: 201–210.

4.   Usui T, Lerner D, Eckert I, Angert AL, Garroway CJ, Hargreaves A, et al. The

evolution of plasticity at geographic range edges. Trends in Ecology & Evolution.

2023;38: 831–842.

5.   Bontrager M, Usui T, Lee-Yaw JA, Anstett DN, Branch HA, Hargreaves AL, et al.

Adaptation across geographic ranges is consistent with strong selection in marginal

climates and legacies of range expansion. Evolution. 2021;75: 1316–1333.

doi:10.1111/evo.14231

6.   Angert AL, Bontrager MG, Ågren J. What do we really know about adaptation at

range edges? Annual Review of Ecology, Evolution, and Systematics. 2020;51:

341–361. doi:10.1146/annurev-ecolsys-012120-091002

7.   Snell-Rood EC, Van Dyken JD, Cruickshank T, Wade MJ, Moczek AP. Toward a

population genetic framework of developmental evolution: the costs, limits, and

consequences of phenotypic plasticity. Bioessays. 2010;32: 71–81.

8.   West-Eberhard MJ. Developmental plasticity and evolution. Oxford University

Press; 2003.

9.   Gluckman PD, Hanson MA. Living with the past: evolution, development, and

patterns of disease. Science. 2004;305: 1733–1736.

10. Leppäluoto J, Hassi J. Human physiological adaptations to the arctic climate.

Arctic. 1991; 139–145.

11. Katzmarzyk PT, Leonard WR. Climatic influences on human body size and

proportions: ecological adaptations and secular trends. American Journal of

Physical Anthropology: The Official Publication of the American Association of

Physical Anthropologists. 1998;106: 483–503.

12. Snodgrass JJ, Leonard WR, Tarskaia LA, Alekseev VP, Krivoshapkin VG. Basal

metabolic rate in the Yakut (Sakha) of Siberia. American Journal of Human

Biology: The Official Journal of the Human Biology Association. 2005;17: 155–

172.

13. Daanen HA, Van Marken Lichtenbelt WD. Human whole body cold adaptation.

Temperature. 2016;3: 104–118.

14.   Frost P. The problem of vitamin D scarcity: Cultural and genetic solutions by

indigenous arctic and tropical peoples. Nutrients. 2022;14: 4071.

15.   Pomeroy E, Stock JT, Stanojevic S, Miranda JJ, Cole TJ, Wells JCK. Trade-offs in

relative limb length among Peruvian children: extending the thrifty phenotype

hypothesis to limb proportions. PLoS One. 2012;7: e51795.

doi:10.1371/journal.pone.0051795

16.   Payne S, Kumar Bc R, Pomeroy E, Macintosh A, Stock J. Thrifty phenotype versus

cold adaptation: trade-offs in upper limb proportions of Himalayan populations of

Nepal. Royal Society Open Science. 2018;5: 172174. doi:10.1098/rsos.172174

17.   Hales CN, Barker DJP. Type 2 (non-insulin-dependent) diabetes mellitus: the

thrifty phenotype hypothesis. Diabetologia. 1992;35: 595–601.

18.   Kuzawa CW. Fetal origins of developmental plasticity: are fetal cues reliable

predictors of future nutritional environments? American Journal of Human

Biology. 2005;17: 5–21.

19.   Li L, Dangour AD, Power C. Early life influences on adult leg and trunk length in

the 1958 British birth cohort. American Journal of Human Biology: The Official

Journal of the Human Biology Association. 2007;19: 836–843.

20.   Ríos L, Terán JM, Varea C, Bogin B. Plasticity in the growth of body segments in

relation to height-for-age and maternal education in Guatemala. American Journal

of Human Biology. 2020;32: e23376.

21.   Holliday TW. Body proportions in Late Pleistocene Europe and modern human

592   origins. Journal of human evolution. 1997;32: 423–448.

593   22.   Gluckman PD, Hanson MA, Spencer HG. Predictive adaptive responses and

594   human evolution. Trends in ecology & evolution. 2005;20: 527–533.

595   23.   Monaghan P. Early growth conditions, phenotypic development and environmental

596   change. Philosophical Transactions of the Royal Society B: Biological Sciences.

597   2007;363: 1635–1645. doi:10.1098/rstb.2007.0011

598   24.   Hales CN, Barker DJP. The thrifty phenotype hypothesis: Type 2 diabetes. British

599   Medical Bulletin. 2001;60: 5–20. doi:10.1093/bmb/60.1.5

600   25.   Barker DJ. The fetal and infant origins of adult disease. BMJ. 1990;301: 1111.

601   doi:10.1136/bmj.301.6761.1111

602   26.   Bateson P, Barker D, Clutton-Brock T, Deb D, D'Udine B, Foley RA, et al.

603   Developmental plasticity and human health. Nature. 2004;430: 419–421.

604   27.   Gluckman PD, Hanson MA, Beedle AS. Early life events and their consequences

605   for later disease: a life history and evolutionary perspective. American journal of

606   human biology. 2007;19: 1–19.

607   28.   Gluckman PD, Hanson MA, Bateson P, Beedle AS, Law CM, Bhutta ZA, et al.

608   Towards a new developmental synthesis: adaptive developmental plasticity and

609   human disease. The Lancet. 2009;373: 1654–1657.

610   29.   Prentice AM, Rayco-Solon P, Moore SE. Insights from the developing world:

611   thrifty genotypes and thrifty phenotypes. Proceedings of the Nutrition Society.

612   2005;64: 153–161.

613   30.   Wells JCK. The thrifty phenotype as an adaptive maternal effect. Biological

614   Reviews. 2007;82: 143–172.

615   31.   Wells JC. The thrifty phenotype: an adaptation in growth or metabolism? American

616   Journal of Human Biology. 2011;23: 65–75.

617   32.   Beall CM, Jablonski NG, Steegmann AT. Human adaptation to climate:

618   temperature, ultraviolet radiation, and altitude. Human biology: an evolutionary

619   and biocultural perspective. 2012; 175–250. doi:10.1002/9781118108062.ch6

620   33.   Abouheif E, Favé M-J, Ibarrarán-Viniegra AS, Lesoway MP, Rafiqi AM,

621   Rajakumar R. Eco-evo-devo: the time has come. Advances in experimental

622   medicine and biology. 2014;781: 107–125. doi:10.1007/978-94-007-7347-9_6

623   34.   Beldade P, Mateus ARA, Keller RA. Evolution and molecular mechanisms of

624   adaptive developmental plasticity. Molecular Ecology. 2011;20: 1347–1363.

625   doi:10.1111/j.1365-294X.2011.05016.x

626   35.   Buchanan KL, Meillère A, Jessop TS. Early life nutrition and the programming of

627   the phenotype. Development strategies and biodiversity: Darwinian fitness and

628   evolution in the Anthropocene. Springer; 2022. pp. 161–214.

629   36.   Carroll SB. Evo-devo and an expanding evolutionary synthesis: a genetic theory of

630   morphological evolution. Cell. 2008;134: 25–36.

631   37.   Davidson EH, Erwin DH. Gene regulatory networks and the evolution of animal

632   body plans. Science. 2006;311: 796–800.

633   38.   Wells JCK. The thrifty phenotype as an adaptive maternal effect. Biological

634   Reviews. 2007;82: 143–172. doi:10.1111/j.1469-185X.2006.00007.x

635   39.   Petry CJ, Desai M, Ozanne SE, Hales CN. Early and late nutritional windows for

636   diabetes susceptibility. Proceedings of the Nutrition Society. 1997;56: 233–242.

637   40.   Low FM, Gluckman PD, Hanson MA. Developmental plasticity, epigenetics and

638   human health. Evolutionary Biology. 2012;39: 650–665.

639   41.   Lillico-Ouachour A, Abouheif E. Regulation, development, and evolution of caste

640   ratios in the hyperdiverse ant genus *Pheidole*. Current Opinion in Insect Science.

641   2017;19: 43–51. doi:10.1016/j.cois.2016.11.003

642   42.   McGlynn TP, Owen JP. Food supplementation alters caste allocation in a natural

643   population of *Pheidole flavens*, a dimorphic leaf-litter dwelling ant. Insectes

644   Sociaux. 2002;49: 8–14. doi:10.1007/s00040-002-8270-6

645   43.   Passera L, Roncin E, Kaufmann B, Keller L. Increased soldier production in ant

646   colonies exposed to intraspecific competition. Nature. 1996;379: 630–631.

647   doi:10.1038/379630a0

648   44.   Hölldobler B, Wilson EO. The superorganism: the beauty, elegance, and
649   strangeness of insect societies. WW Norton & Company; 2009.

650   45.   Yang AS. Thinking outside the Embryo: The Superorganism as a Model for
651   EvoDevo Studies. Biological Theory. 2007;2: 398–408.
652   doi:10.1162/biot.2007.2.4.398

653   46.   Gardner A, Grafen A. Capturing the superorganism: a formal theory of group
654   adaptation. Journal of evolutionary biology. 2009;22: 659–671. doi:10.1111/j.1420-
655   9101.2008.01681.x

656   47.   Michod RE. The theory of kin selection. Annual Review of Ecology and
657   Systematics. 1982;13: 23–55.

658   48.   Bell-Roberts L, Turner JFR, Werner GDA, Downing PA, Ross L, West SA. Larger
659   colony sizes favoured the evolution of more worker castes in ants. Nature ecology
660   & evolution. 2024;8: 1959–1971.

661   49.   Kaspari M, Vargo EL. Colony size as a buffer against seasonality: Bergmann's rule
662   in social insects. The American Naturalist. 1995;145: 610–632.

663   50.   Yang AS, Martin CH, Nijhout HF. Geographic variation of caste structure among
664   ant populations. Current Biology. 2004;14: 514–519.
665   doi:10.1016/j.cub.2004.03.005

666   51.   La Richelière F, Muñoz G, Guénard B, Dunn RR, Economo EP, Powell S, et al.
667   Warm and arid regions of the world are hotspots of superorganism complexity.

Proceedings of the Royal Society B. 2022;289: 20211899.

doi:10.1098/rspb.2021.1899

52.   Nguyen AD, Brown M, Zitnay J, Cahan SH, Gotelli NJ, Arnett A, et al. Trade-offs

in cold resistance at the northern range edge of the common woodland ant

Aphaenogaster picea (Formicidae). The American Naturalist. 2019;194: E151–

E163. doi:10.1086/705939

53.   Hardie DC, Hutchings JA. Evolutionary ecology at the extremes of species' ranges.

Environmental Reviews. 2010;18: 1–20.

54.   Sexton JP, McIntyre PJ, Angert AL, Rice KJ. Evolution and ecology of species

range limits. Annu Rev Ecol Evol Syst. 2009;40: 415–436.

55.   Bridle JR, Vines TH. Limits to evolution at range margins: when and why does

adaptation fail? Trends in ecology & evolution. 2007;22: 140–147.

56.   Wilson EO. The origin and evolution of polymorphism in ants. The Quarterly

Review of Biology. 1953;28: 136–156.

57.   Wheeler D. The developmental basis of worker caste polymorphism in ants. The

American Naturalist. 1991;138: 1218–1238. doi:10.1086/285279

58.   MacMillan O, Singer J, Perrakis S, Craig A, Ntanga D, Qiu D, et al. Juvenile

Hormone Signalling Underlies the Switchpoint and Differentiation of Soldiers in

Camponotus floridanus. bioRxiv. 2025; 2025.08.25.672237.

doi:10.1101/2025.08.25.672237

688 59. Ellis PJ, Morris TJ, Skinner BM, Sargent CA, Vickers MH, Gluckman PD, et al.

689   Thrifty metabolic programming in rats is induced by both maternal undernutrition

690   and postnatal leptin treatment, but masked in the presence of both: implications for

691   models of developmental programming. BMC genomics. 2014;15: 49.

692 60. Sauce B, Goes CP, Forti I, O. do Monte BG, Watanabe IM, Cunha J, et al. A link

693   between thrifty phenotype and maternal care across two generations of intercrossed

694   mice. PLoS One. 2017;12: e0177954.

695 61. Ferral N, Holloway K, Li M, Yin Z, Hou C. Heterogeneous activity causes a

696   nonlinear increase in the group energy use of ant workers isolated from queen and

697   brood. Insect Science. 2018;25: 487–498.

698 62. Guo X, Lynch C, Fewell JH, Harrison JF. Ontogeny of energy use in harvester ant

699   colonies, and the metabolic expense of colony growth. Proceedings B. 2025;292:

700   20242534.

701 63. Cannon CA. Demography, cold hardiness, and nutrient reserves of overwintering

702   nests of the carpenter ant *Camponotus pennsylvanicus* (De Geer) (Hymenoptera:

703   Formicidae). Virginia Polytechnic Institute and State University. 1990.

704 64. Ma R, Zhang L, He H. Eco-Morphological Responses of Camponotus japonicus

705   (Hymenoptera: Formicidae) to Varied Climates and Habitats. Insects. 2024;15:

706   719.

707   65.   Smith C, Anderson K, Tillberg C, Gadau J, Suarez A. Caste determination in a

708   polymorphic social insect: nutritional, social, and genetic factors. The American

709   Naturalist. 2008;172: 497–507.

710   66.   Blanckenhorn WU, Bauerfeind SS, Berger D, Davidowitz G, Fox CW, Guillaume

711   F, et al. Life history traits, but not body size, vary systematically along latitudinal

712   gradients on three continents in the widespread yellow dung fly. Ecography.

713   2018;41: 2080–2091. doi:https://doi.org/10.1111/ecog.03752

714   67.   Molet M, Péronnet R, Couette S, Canovas C, Doums C. Effect of temperature and

715   social environment on worker size in the ant Temnothorax nylanderi. Journal of

716   Thermal Biology. 2017;67: 22–29.

717   68.   Jayatilaka P, Narendra A, Reid SF, Cooper P, Zeil J. Different effects of

718   temperature on foraging activity schedules in sympatric Myrmecia ants. Journal of

719   Experimental Biology. 2011;214: 2730–2738. doi:10.1242/jeb.053710

720   69.   Spicer ME, Stark AY, Adams BJ, Kneale R, Kaspari M, Yanoviak SP. Thermal

721   constraints on foraging of tropical canopy ants. Oecologia. 2017;183: 1007–1017.

722   70.   Sepúlveda Y, Nicholls E, Schuett W, Goulson D. Heatwave-like events affect drone

723   production and brood-care behaviour in bumblebees. PeerJ. 2024;12: e17135.

724   71.   Nürnberger F, Härtel S, Steffan-Dewenter I. The influence of temperature and

725   photoperiod on the timing of brood onset in hibernating honey bee colonies. PeerJ.

726   2018;6: e4801.

72.   Wögler L, Kurze C. Experimental short-term heatwaves negatively impact body

weight gain and survival during larval development in Bombus terrestris

L.(Hymenoptera: Apidae). Biology Open. 2025;14: bio061781.

73.   Mitesser O, Weissel N, Strohm E, Poethke H-J. The evolution of activity breaks in

the nest cycle of annual eusocial bees: a model of delayed exponential growth.

BMC Evolutionary Biology. 2006;6: 45. doi:10.1186/1471-2148-6-45

74.   Abouheif E. Ant caste evo-devo: it's not all about size. Trends in Ecology &

Evolution. 2021;36: 668–670. doi:10.1016/j.tree.2021.04.002

75.   Abouheif E, Wray GA. Evolution of the gene network underlying wing

polyphenism in ants. Science. 2002;297: 249–252.

76.   Favé MJ, Johnson RA, Cover S, Handschuh S, Metscher BD, Müller GB, et al.

Past climate change on Sky Islands drives novelty in a core developmental gene

network and its phenotype. BMC Evolutionary Biology. 2015;15: 1–21.

doi:10.1186/s12862-015-0448-4

77.   Qiu B, Dai X, Li P, Larsen RS, Li R, Price AL, et al. Canalized gene expression

during development mediates caste differentiation in ants. Nature ecology &

evolution. 2022;6: 1753–1765.

78.   Vasquez-Correa A, Arnet J, Chen T, Abouheif E. Latent gene network expression

underlies the reversion of a polyphenic trait in the worker caste of ants. bioRxiv.

2025; 2025–11.

79.   Vizueta J, Xiong Z, Ding G, Larsen RS, Ran H, Gao Q, et al. Adaptive radiation
748   and social evolution of the ants. Cell. 2025.

749   80.   Ma R, Zhang L, Yang L, Tang L, Zhang X, Wei C, et al. Environmental Variation
750   Contributes to Head Phenotypes in Workers of Camponotus japonicus
751   (Hymenoptera: Formicidae). Ecology and Evolution. 2025;15: e71940.

752   81.   Azevedo-Silva M, Sendoya SF, Côrtes MC, Longo PA, Nogueira A, Mori GM, et
753   al. Sugar-Rich Resources Mediate Geographic Morphological Variation in a
754   Dominant, Neotropical Savanna Ant. Journal of Biogeography. 2025;52: e15111.

755   82.   Chen T, Rafiqi AM, Rajakumar A, Milat NS, Lamouret T, Plante É, et al. A
756   Developmental Table for the Florida Carpenter Ant *Camponotus floridanus*:
757   Establishing Foundations for Mechanistic Studies of Development and Evolution
758   in Ants. 2025. doi:10.32388/OJQC0N

759   83.   Mirth CK, Saunders TE, Amourda C. Growing up in a changing world:
760   environmental regulation of development in insects. Annual Review of
761   Entomology. 2021;66: 81–99.

762   84.   Snell-Rood EC, Ehlman SM. Ecology and evolution of plasticity. Phenotypic
763   plasticity & evolution. 2021; 139–160.

764   85.   Hou Z, Fuiman LA. Nutritional programming in fishes: insights from mammalian
765   studies. Reviews in Fish Biology and Fisheries. 2020;30: 67–92.

86. Jonsson B, Jonsson N, Hansen MM. Knock-on effects of environmental influences

during embryonic development of ectothermic vertebrates. The Quarterly Review

of Biology. 2022;97: 95–139.

87. Bateson P, Gluckman P, Hanson M. The biology of developmental plasticity and

the Predictive Adaptive Response hypothesis. The Journal of physiology.

2014;592: 2357–2368.

88. Scheiner SM. Bet-hedging as a complex interaction among developmental

instability, environmental heterogeneity, dispersal, and life-history strategy.

Ecology and Evolution. 2014;4: 505–515.

89. Wilson EO. The soldier of the ant *Camponotus* (*Colobopsis*) *fraxinicola* as a

trophic caste. Psyche. 1974;81: 182–188. doi:10.1155/1974/17428

90. Dussutour A, Poissonnier L-A, Buhl C, Simpson SJ. Resistance to nutritional stress

in ants: when being fat is advantageous. Journal of Experimental Biology.

2016;219: 824–833. doi:10.1242/jeb.136234

91. Yang AS. Seasonality, division of labor, and dynamics of colony-level nutrient

storage in the ant *Pheidole morrisi*. Insectes Sociaux. 2006;53: 456–462.

doi:10.1007/s00040-005-0896-3

92. Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces

for global land areas. International Journal of Climatology. 2017;37: 4302–4315.

doi:https://doi.org/10.1002/joc.5086

93.   Agriculture and Agri-Food Canada. Length of growing season - Quebec. Ottawa,
787   Ontario: Prepared by Agriculture and Agri-Food Canada's Agri-Environmental
788   Services Branch (AESB), in partnership with Research Branch and Environment
789   Canada's National Service Office-Agriculture; 2010.

790   94.   Janicki J, Narula N, Ziegler M, Guénard B, Economo EP. Visualizing and
791   interacting with large-volume biodiversity data using client–server web-mapping
792   applications: The design and implementation of antmaps. org. Ecological
793   informatics. 2016;32: 185–193. doi:10.1016/j.ecoinf.2016.02.006

794   95.   Ellison AM, Gotelli NJ, Farnsworth EJ, Alpert GD. A field guide to the ants of
795   New England. Yale University Press; 2012.

796   96.   Akre RD, Hansen LD, Myhre EA. Colony size and polygyny in carpenter ants
797   (Hymenoptera: Formicidae). Journal of the Kansas Entomological Society.
798   1994;67: 1–9.

799   97.   Gibson RL. Soldier production in *Camponotus novaeboracensis* during colony
800   growth. Insectes Sociaux. 1989;36: 28–41. doi:10.1007/BF02225878

801   98.   Passera L. Production des soldats dans les sociétés sortant d'hibernation chez la
802   fourmi *Pheidole pallidula* (Nyl.) (Formicidae, Myrmicinae). Insectes sociaux.
803   1977;24: 131–146.

804   99.   Walker J, Stamps J. A test of optimal caste ratio theory using the ant *Camponotus*
805   (*Colobopsis*) *impressus*. Ecology. 1986;67: 1052–1062. doi:10.2307/1939828

806   100.   Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution

807   interpolated climate surfaces for global land areas. International Journal of

808   Climatology: A Journal of the Royal Meteorological Society. 2005;25: 1965–1978.

809   doi:10.1002/joc.1276

810   101.   Bair E, Hastie T, Paul D, Tibshirani R. Prediction by supervised principal

811   components. Journal of the American Statistical Association. 2006;101: 119–137.

812   doi:10.1198/016214505000000628

813   102.   Vesnic A, Škrijelj R, Trožić-Borovac S, Tomanović Ž. Diversity and nesting

814   preferences of *Camponotus lateralis* group species on Western Balkan Peninsula

815   (Hymenoptera: Formicidae). Journal of the Entomological Research Society.

816   2017;19: 73–82.

817   103.   Chen Y, Hansen LD, Brown JJ. Nesting sites of the carpenter ant, *Camponotus*

818   *vicinus* (Mayr) (Hymenoptera: Formicidae) in northern Idaho. Environmental

819   Entomology. 2002;31: 1037–1042. doi:10.1603/0046-225X-31.6.1037

820   104.   Porter SD. Impact of temperature on colony growth and developmental rates of the

821   ant, *Solenopsis invicta*. Journal of Insect Physiology. 1988;34: 1127–1133.

822   doi:10.1016/0022-1910(88)90215-6

823   105.   Lopatina EB, Kipyatkov VE. The influence of temperature on brood development

824   in the incipient colonies of the ants *Camponotus herculeanus* (L.) and *Camponotus*

825   *xerxes* Forel (Hymenoptera, Formicidae). Proceedings of the Colloquia on Social

826   Insects. 1993. pp. 61–74.

106. Jo H, Song W, Jablonski PG, Lee S. Exploration and Aggregation of Differently

Sized Camponotus japonicus Worker Ants. 2025.

107. Wood SN. Generalized additive models: An introduction with R. Generalized

Additive Models. 2017;10: 9781315370279.

108. Wood SN, Pya N, Säfken B. Smoothing parameter and model selection for general

smooth models. Journal of the American Statistical Association. 2016;111: 1548–

1563. doi:10.1080/01621459.2016.1180986

109. Wood S, Wood MS. Package 'mgcv.' R package version. 2015;1: 729.

110. Mazerolle MJ, Mazerolle MMJ. Package 'AICcmodavg.' See https://cran uib

no/web/packages/AICcmodavg/AICcmodavg pdf. 2020.

111. R Core Team. R: A Language and Environment for Statistical Computing. Vienna,

Austria: R Foundation for Statistical Computing; 2024.

112. Wilson EO. The organization of colony defense in the ant *Pheidole dentata* Mayr

(Hymenoptera: Formicidae). Behavioral Ecology and Sociobiology. 1976;1: 63–

81. doi:doi.org/10.1007/BF00299953

113. Molet M, Maicher V, Peeters C. Bigger helpers in the ant *Cataglyphis bombycina*:

Increased worker polymorphism or novel soldier caste? PLoS One. 2014;9:

e84929. doi:10.1371/journal.pone.0084929

114.   Hasegawa E. Nest defense and early production of the major workers in the

846   dimorphic ant *Colobopsis nipponicus* (Wheeler) (Hymenoptera : Formicidae).

847   Behavioral Ecology and Sociobiology. 1993;33: 73–77. doi:10.1007/BF00171658

848   115.   Metzl C, Wheeler D, Abouheif E. Wilhelm Goetsch (1887 - 1960): Pioneering

849   studies on the development and evolution of the soldier caste in social insects.

850   Myrmecological News. 2018;26: 81–96. doi:10.25849/myrmecol.news_026:081

851   116.   Lamon B, Topoff H. Avoiding predation by army ants: defensive behaviours of

852   three ant species of the genus *Camponotus*. Animal Behaviour. 1981;29: 1070–

853   1081. doi:10.1016/S0003-3472(81)80060-7

854   117.   Wilson EO. Pheidole in the New World: a dominant, hyperdiverse ant genus.

855   Harvard University Press; 2003.

856   118.   Cerdá X, Retana J, Cerda X. Links between worker polymorphism and thermal

857   biology in a thermophilic ant species. Oikos. 1997;78: 467. doi:10.2307/3545608

858   119.   Bernstein RA. The adaptive value of polymorphism in an alpine Ant, *Formica

859   neorufibarbis* Gelid A. Wheeler. Psyche (New York). 1976;83: 180–184.

860   doi:10.1155/1976/52542

861   120.   Wendt CF, Verble-Pearson R. Critical thermal maxima and body size positively

862   correlate in red imported fire ants, *Solenopsis invicta*. The Southwestern Naturalist.

863   2016;61: 79–83. doi:10.1894/0038-4909-61.1.79

864   121. Porter SD, Tschinkel WR. Foraging in *Solenopsis invicta* (Hymenoptera:

865   Formicidae): effects of weather and season. Environmental Entomology. 1987;16:

866   802–808. doi:10.1093/ee/16.3.802

867   122. Cerdá X, Retana J. Alternative strategies by thermophilic ants to cope with extreme

868   heat: Individual versus colony level traits. Oikos. 2000;89: 155–163.

869   doi:10.1034/j.1600-0706.2000.890117.x

870   123. O'Donnell S, Bulova S, Caponera V, Oxman K, Giladi I. Species differ in worker

871   body size effects on critical thermal limits in seed-harvesting desert ants (*Messor*

872   *ebeninus* and *M. arenarius*). Insectes Sociaux. 2020;67: 473–479.

873   doi:10.1007/s00040-020-00782-5

874   124. Hood WG, Tshchinkel WR. Desiccation resistance in arboreal and terrestrial ants.

875   Physiological Entomology. 1990;15: 23–35. doi:10.1111/j.1365-

876   3032.1990.tb00489.x

877   125. Kaspari M. Body size and microclimate use in Neotropical granivorous ants.

878   Oecologia. 1993;96: 500–507. doi:10.1007/BF00320507

879   126. Powell S, Franks NR. Ecology and the evolution of worker morphological

880   diversity: a comparative analysis with *Eciton* army ants. Functional Ecology.

881   2006;20: 1105–1114.

882   127. Retana J, Cerdá X. Worker size polymorphism conditioning size matching in two

883   sympatric seed-harvesting ants. Oikos. 1994;71: 261–266. doi:10.2307/3546274

128. Kaspari M. Worker size and seed size selection by harvester ants in a Neotropical forest. Oecologia. 1996;105: 397–404. doi:10.1007/BF00328743

129. Willott SJ, Compton SG, Incoll LD. Foraging, food selection and worker size in the seed harvesting ant *Messor bouvieri*. Oecologia. 2000;125: 35–44. doi:10.1007/PL00008889

130. Wilson EO. The relation between caste ratios and division of labor in the ant genus *Pheidole* (Hymenoptera: Formicidae). Behavioral Ecology and Sociobiology. 1984;16: 89–98. doi:10.1007/BF00293108

131. Sarty M, Abbott K, Lester P. Habitat complexity facilitates coexistence in a tropical ant community. Oecologia. 2006;149: 465–473. doi:10.1007/s00442-006-0453-9

132. Kaspari M, Weiser MD. The size–grain hypothesis and interspecific scaling in ants. Functional Ecology. 1999;13: 530–538. doi:10.1046/j.1365-2435.1999.00343.x

133. Farji-Brener A, Barrantes G, Ruggiero A. Environmental rugosity, body size and access to food: A test of the size-grain hypothesis in tropical litter ants. Oikos. 2004;104: 165–171. doi:10.1111/j.0030-1299.2004.12740.x

134. Parr ZJE, Parr CL, Chown SL. The size-grain hypothesis: a phylogenetic and field test. Ecological Entomology. 2003;28: 475–481. doi:10.1046/j.1365-2311.2003.00529.x

135. Centorame M, Angelino D, Bonanni R, Fanfani A. Static and evolutionary allometry in the Italian endemic ant species *Cataglyphis italica* (Emery 1906).

Ethology Ecology & Evolution. 2020;32: 16–28.

doi:10.1080/03949370.2019.1639080

136. Hölldobler B, Wilson EO. The ants. Harvard University Press; 1990.

137. Araujo MB, Tschinkel WR. Worker allometry in relation to colony size and social

form in the fire ant *Solenopsis invicta*. Journal of Insect Science. 2010;10: 1–10.

doi:10.1673/031.010.9401

138. Schöning C, Kinuthia W, Franks NR, Lindström J. Evolution of Allometries in the

Worker Caste of *Dorylus* Army Ants. Oikos. 2005;110: 231–240.

doi:10.1111/j.0030-1299.2005.13672.x

139. Dinerstein E, Olson D, Joshi A, Vynne C, Burgess ND, Wikramanayake E, et al.

An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm.

BioScience. 2017;67: 534–545. doi:10.1093/biosci/bix014

140. Shik JZ. The metabolic costs of building ant colonies from variably sized subunits.

Behavioral Ecology and Sociobiology. 2010;64: 1981–1990.

141. Coto ZN, Traniello JF. Social brain energetics: Ergonomic efficiency,

neurometabolic scaling, and metabolic polyphenism in ants. Integrative and

comparative biology. 2022;62: 1471–1478.

142. Porter SD, Tschinkel WR. Fire ant polymorphism: the ergonomics of brood

production. Behavioral Ecology and Sociobiology. 1985;16: 323–336.

143. Kaspari M, Byrne MM. Caste allocation in litter *Pheidole*: lessons from plant

defense theory. Behavioral Ecology and Sociobiology. 1995;37: 255–263.

144. Kaspari M, Weiser MD. Energy, taxonomic aggregation, and the geography of ant

abundance. Ecography. 2012;35: 65–72.

145. McGlynn TP, Diamond SE, Dunn RR. Tradeoffs in the evolution of caste and body

size in the hyperdiverse ant genus Pheidole. PLoS One. 2012;7: e48202.

TABLES

Table 1. Rationale for using different ant colony-level traits of the worker polymorphism.

Colony trait	Functions for the colony
Proportion of major workers	**Defense of the colony** [50,89,112–117] **Food storage** [89,91]
Major worker mean body size	**Thermal tolerance** [118–123] **Desiccation** [124,125]
Worker head and body size variance	**Dietary composition** [126–129] **Division of labor** [56,57,130]
Worker head-to-body ratio variance	**Habitat complexity** [131–134] **Division of labor** [135–138]

935 **FIGURES**

2 mm

938 **Figure 1.** *Camponotus herculeanus* worker polymorphism (left-to-right) from minors to
939 medias and majors. All workers originate from the same colony and were photographed
940 at the same scale. These specimens represent the possible worker morphological range for
941 a given large colony of the species in the study sites in the province of Quebec, Canada.

943
944 **Figure 2.** Location (*n* = 26) of the *Camponotus herculeanus* colonies used for the
945 observational study in Eastern Canada (province of Quebec) across biomes [139].

947
948   **Figure 3.** The (A) proportion of major workers and the (B) number of workers in
949   *Camponotus herculeanus* colonies (*n* = 26) in relation to latitude.

Figure 4. The proportion of major workers in relation to micro- and macro-climate for (*n* = 18) *Camponotus herculeanus* nest site locations. Micro-climatic variables (A & B) were obtained from Thermochron iButton Devices positioned adjacent to each nest. (A) colony activity is estimated as the number of days above 0°C and (B) larval development is estimated as the number of days above 9°C. These temperature measures were obtained between August 21st, 2021, and June 30th, 2022. Macro-climatic variables (C & D) were (C) regional mean annual temperature and (D) regional mean diurnal range for the years 1970 to 2000 [92].

Figure 5. Results from the common garden experiment. Initial (green) and final (orange)
(A) proportion of major workers and their average body size (B) in (n = 9) *Camponotus*
herculeanus colonies collected from the southern (A-E) and northern (F-I) parts of the
environmental gradient.

Figure 6. The thrifty phenotype as an adaptation to unpredictable environments in
Camponotus herculeanus. The figure illustrates colony-level developmental plasticity
leading to thrifty phenotype in ant colonies and its nutritional context-dependent
consequences on colony survival. Young (early-instar) larvae develop from eggs during
fall, enter diapause and receive environmental cues in spring that influence caste
determination, potentially mediated by colony-level nutritional conditions. In northern
colonies, which experience nutrient-poor environments during the spring, colonies favor
the production of cost-effective minor workers over costly major workers (i.e., thrifty
phenotype), which may buffer colonies against resource scarcity during nutrient-poor
summers, while maintaining colony size. In southern colonies, larvae benefit from
nutrient-rich springs, leading to developmental reprogramming and production of more
major workers, which should enhance colony fitness during nutrient-rich summers while
maintaining colony size. Solid arrows in the figure indicate putative causal pathways
from environmental cues to subcaste outcomes.