Typhi Mykrobe: fast and accurate lineage identification and antimicrobial resistance genotyping directly from sequence reads for the typhoid fever agent *Salmonella* Typhi

Authors:

Danielle J. Ingle[1], Jane Hawkey[2,3], Martin Hunt[4,5,6,7], Zamin Iqbal[4,8], Jacqueline A. Keane[9] Ayorinde O. Afolayan[10,11], Niyaz Ahmed[12], Saadia Andleeb[13], Philip M. Ashton[14,15], Isaac I. Bogoch[16], Megan E. Carey[17,18], Marie Anne Chattaway[19], John A. Crump[20], Paula Diaz Guevara[21], Benjamin P. Howden[1,22,23], Hidemasa Izumiya[24], Jobin John Jacob[25], Louise M. Judd[1,23], Arti Kapil[26], Karen H. Keddy[27], Justin Y. Kim[28], Myron M. Levine[29], Masatomo Morita[24], Satheesh Nair[19], Sophie Octavia[30], Iruka N. Okeke[10], Precious E. Osadebamwen[10], Sadia Isfat Ara Rahman[31], Assaf Rokney[32], David A. Rasko[29,33,34], Varun Shamanna[35], Michael J. Sikorski[29,33,], Anthony M. Smith[36,37], Gabriel T. Sunmonu[10], Kaitlin A. Tagg[28], Ryan R. Wick[1], Zoe A. Dyson[17,38]*, Kathryn E. Holt[2,17]* and Global Typhoid Genomics Consortium

* Indicates equal contribution

Affiliations:

The Department of Microbiology and Immunology, The University of Melbourne at The Peter Doherty Institute for Infection and Immunity, Melbourne, Australia

Department of Infectious Diseases, School of Translational Medicine, Monash University, Melbourne, Victoria 3004, Australia

Centre to Impact AMR, Monash University, Clayton, Victoria, Australia

European Molecular Biology Laboratory - European Bioinformatics Institute, Hinxton, Cambridge, United Kingdom

Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

National Institute of Health Research Oxford Biomedical Research Centre, John Radcliffe Hospital, Headley Way, Oxford, United Kingdom

Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance, University of Oxford, Oxford, United Kingdom.

Milner Centre for Evolution, University of Bath, United Kingdom

Department of Medicine, University of Cambridge, Cambridge Institute of Therapeutic Immunology and Infectious Disease, Jeffrey Cheah Biomedical Centre, Cambridge Biomedical Campus, United Kingdom

10 Faculty of Pharmacy and College of Medicine, University of Ibadan, Ibadan, Nigeria

11 Institute for Infection Prevention and Control, Medical Centre - University of Freiburg,

Freiburg, Germany

12 Pathogen Biology Laboratory, Department of Biotechnology and Bioinformatics, University of

Hyderabad, Hyderabad, India

13 Atta-ur-Rahman School of Applied Biosciences, National University of Sciences and

Technology, Pakistan

14 Malawi Liverpool Wellcome Programme, Blantyre, Malawi

15 Institute of Infection, Ecology and Veterinary Sciences, University of Liverpool, Liverpool, UK

16 Department of Medicine, University of Toronto, Canada

17 Department Infection Biology, London School of Hygiene and Tropical Medicine, London, UK

18 IAVI, Chelsea & Westminster Hospital, London, United Kingdom

19 Gastrointestinal Bacteria Reference Unit, United Kingdom Health Security Agency, London,

United Kingdom

20 Centre for International Health, University of Otago, Dunedin, New Zealand

21 Grupo de Microbiologia, Instituto Nacional de Salud, Bogota, Colombia

22 Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology &

Immunology, The University of Melbourne at The Peter Doherty Institute for Infection and

Immunity, Melbourne, Australia

23 Innovation Hub, Centre for Pathogen Genomics, University of Melbourne (Doherty Institute)

24 Department of Bacteriology I, National Institute of Infectious Diseases, Tokyo, Japan

25 Department of Clinical Microbiology, Christian Medical College, Vellore, India

26 All India Institute of Medical Sciences, New Delhi, India

27 Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of

Pretoria, South Africa

28 Centers for Disease Control and Prevention, Atlanta, United States of America

29 Center for Vaccine Development and Global Health, University of Maryland School of

Medicine, Baltimore, Maryland, United States of America

30 School of Biotechnology and Biomolecular Sciences, University of New South Wales,

Sydney, Australia

31 International Centre for Diarrhoeal Disease Research Bangladesh (icddr,b), Bangladesh

32 Ministry of Health, Jerusalem, Israel

33 Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore,

Maryland, United States of America

34 University of Maryland School of Dentistry, Baltimore, Maryland, United States of America

35 Central Research Laboratory, KIMS, Bengaluru, India

36 Centre for Enteric Diseases, National Institute for Communicable Diseases, Johannesburg,

South Africa

37 Department of Medical Microbiology, Faculty of Health Sciences, University of Pretoria,

Pretoria, South Africa

38 Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, United Kingdom

Correspondence:

Danielle J. Ingle: danielle.ingle@unimelb.edu.au

Zoe A. Dyson: zoe.dyson@lshtm.ac.uk

Kathryn E. Holt: kat.holt@lshtm.ac.uk

Key words: Typhoid fever, *Salmonella* Typhi, Antimicrobial Resistance (AMR), Whole Genome

Sequencing (WGS), pathogen sequencing, genotyping, lineage

Abstract

Background

Typhoid fever results from systemic infection with *Salmonella enterica* serovar Typhi (Typhi) and
causes 10 million illnesses annually. Disease control relies on prevention (water, sanitation, and
hygiene interventions or vaccination) and effective antimicrobial treatment. Antimicrobial
resistant (AMR) Typhi lineages have emerged and become established in many parts of the
world. Knowledge of local pathogen populations informed by genomic surveillance, including of
lineages (defined by the GenoTyphi scheme) and AMR determinants, is increasingly used to
inform local treatment guidelines and to inform vaccination strategy. Current tools for genotyping
Typhi require multiple read alignment or assembly steps and have not been validated for
analysis of data generated with Oxford Nanopore Technologies (ONT) long-read sequencing
devices. Here, we introduce Typhi Mykrobe, a command line software tool for rapid genotyping
of Typhi lineages, AMR determinants, and plasmid replicons direct from sequencing reads.

Results
We validated Typhi Mykrobe lineage genotyping by comparison with the current standard read
mapping-based approach and demonstrated 99.8% concordance across nearly 13,000
genomes sequenced with Illumina platforms. For the few isolates with discordant calls, we show
that Typhi Mykrobe results are better supported by the evidence from raw sequence read data
than the results generated using the mapping-based approach. We also demonstrate 99.9%
concordance for detection of AMR determinants compared with the current standard assembly-
based approach, with similar results for plasmid marker detection. Typhi Mykrobe predicts
clinical resistance categorisation (S/I/R) for eight drug classes, and we show strong agreement
with phenotypic categorisations generated from reference laboratory minimum inhibitory
concentration (MIC) data for n=1,572 Illumina-sequenced isolates (>99% agreement within one
doubling dilution). We show strong concordance (>96% for genotype and >98% for AMR and
plasmid) between calls made from ONT reads and those made from Illumina reads for isolates
sequenced on both platforms (n =93 genomes). Typhi Mykrobe takes less than a minute per
sample and is available at https://github.com/typhoidgenomics/genotyphi.

Conclusions
Typhi Mykrobe provides rapid and sensitive genotyping of Typhi genomes direct from Illumina
and ONT reads, although lower accuracy was observed for R9 ONT data. It demonstrated
accurate assignment of GenoTyphi lineage, detection of AMR determinants and prediction of
corresponding AMR phenotypes, and identification of plasmid replicons.

Background

Typhoid fever is caused by systemic infection with the human-restricted bacterium *Salmonella enterica* subspecies *enterica* serovar Typhi (Typhi) [1]. More than 10 million typhoid fever illnesses occur annually (mainly in South Asia), associated with 100,000 deaths, and the risk of complications and death increases without effective antimicrobial therapy [2]. The World Health Organization AWaRe (Access, Watch, Reserve) Antibiotic guidance recommends typhoid fever is treated with ciprofloxacin, unless resistance is prevalent locally, in which case azithromycin is recommended to treat uncomplicated disease and intravenous ceftriaxone is recommended for treatment of severe disease [3,4]. A high proportion of typhoid fever is caused by antimicrobial resistant (AMR) Typhi, and nearly all Typhi isolated in South Asia have been ciprofloxacin non-susceptible for over two decades [5,6]. Extensively drug resistant (XDR, resistant to amoxicillin/ampicillin, chloramphenicol, trimethoprim-sulfamethoxazole, ciprofloxacin, and ceftriaxone) Typhi, for which the only treatment options are oral azithromycin or intravenous carbapenems, have been circulating in Pakistan for nearly a decade with travel-associated infections reported globally [7–9]. Concerningly, XDR Typhi with no travel links has been reported in USA and China [7–10], and XDR isolates with additional resistance to azithromycin and carbapenems were recently reported in Pakistan [11].

Tracking the emergence and spread of AMR Typhi through whole genome sequencing (WGS) is important to inform public health control measures such as deployment of typhoid conjugate vaccines (TCVs) [12], investigating and controlling outbreaks [10], and informing the development of antimicrobial treatment guidance [3], including management of travel-associated infections in non-endemic countries [4]. Genetic determinants of AMR are well-understood in Typhi [13,14], and include both acquired genes that are plasmid-borne or chromosomally integrated [15], and point mutations in core genes. These include *acrB* for azithromycin resistance [16], and quinolone resistance-determining regions (QRDR) of *gyrA*, *gyrB,* and *parC* for ciprofloxacin non-susceptibility and resistance [13,17]. Recently, carbapenem-resistant Typhi was described, due to acquisition of a plasmid-borne carbapenemase gene [11]. Typhi is highly clonal, which has hampered subtyping in the past; however, its population structure has been well defined using WGS [15], and the GenoTyphi framework for genotyping and variant nomenclature first proposed in 2016 [18] has been widely adopted by the global community to facilitate identification of emerging variants and communication across laboratories and settings [13,14,19–21].

The GenoTyphi scheme was originally defined based on nearly 2,000 WGS isolates from over

60 countries [18]. It uses a hierarchical framework based on the global phylogeny, stratifying the

Typhi population structure into four primary lineages, 16 clades, and 49 subclades (henceforth

referred to as genotypes), based on lineage-specific marker single nucleotide variants (SNVs).

A schematic mapping the hierarchical genotype labels to the Typhi phylogeny is shown in

Supplementary Figure 1. Under the GenoTyphi scheme, the lineage formerly known as 'H58'

was designated genotype 4.3.1, and is further divided into multiple higher-resolution genotypes

with the prefix 4.3.1 including 4.3.1.1, 4.3.1.2 (see **Supplementary Figure 1**). The H58 lineage,

or genotype 4.3.1 and associated sub-genotypes, has been associated with the spread of

multidrug resistance (MDR, resistant to amoxicillin/ampicillin, chloramphenicol, trimethoprim-

sulfamethoxazole) Typhi throughout Asia and into Eastern Africa [15]. The scheme was later

extended to provide further discriminatory power for identifying additional subpopulations of

epidemiological importance [21–27], such as the XDR Typhi lineage associated with outbreaks

in Pakistan (genotype 4.3.1.1.P1) [26,27]. The scheme currently includes 87 genotypes [28],

and its future maintenance and development will be managed by a working group within the

Global Typhoid Genomics Consortium (https://www.typhoidgenomics.org/).

The GenoTyphi scheme specification consists of a set of marker SNVs defining the hierarchical

genotypes, which in principle can be used by any bioinformatics tool to assign genotypes to new

genomes. The original GenoTyphi paper [18] was accompanied by a Python-based pipeline

(available at: https://github.com/typhoidgenomics/genotyphi) designed for typing Illumina WGS

data against the scheme. This implementation requires users to first map short reads or

assemblies to the reference sequence of Typhi strain CT18 (accession AL513382.1) [29], then

provide as input the resulting binary alignment map (BAM) or variant call file (VCF). Marker

SNVs were identified from these input files and used to assign genomes to genotypes.

Functionality was extended to include detection of mutations associated with reduced

susceptibility to ciprofloxacin or azithromycin [22]. This mapping-based approach had various

limitations, largely due to the need for users to pre-map their data to a specific reference, which

was cumbersome (e.g. there is wide variation in the upstream mapping tools used and some

users had issues with mapping tools creating incompatible output files, the BAM files are large

and inefficient to process), and AMR analysis was restricted to mutation detection. The

GenoTyphi scheme has also been implemented in other genotyping tools, including Typhi

PathogenWatch [13], which works by analysing assembled genomes, and the split-*k*-mer-based

BioHansel tool, which does not include AMR detection [30].

186 Here, we present a novel approach to rapid genotyping of both AMR and GenoTyphi lineages

187 from reads, implemented in the Mykrobe open-source software framework [31], dubbed Typhi

188 Mykrobe. Input into this new tool accepts raw WGS reads (FASTQ files), and has been tested

189 with reads generated by ONT and Illumina platforms. The tool performs rapid (i) lineage

190 genotyping using the GenoTyphi framework; (ii) detection of a broad range of clinically relevant

191 molecular determinants of AMR, including both acquired genes and mutations; and (iii)

192 detection of plasmid replicon markers common to Typhi. We validated GenoTyphi lineage

193 assignment from reads by comparison with the original mapping-based implementation, using

194 12,839 Typhi genomes [14]. We assessed AMR genotype calls by comparison with assembly-

195 based genotype calls from PathogenWatch, and with antimicrobial susceptibility phenotypes for

196 4,018 isolates. Finally, we assessed performance on long read data generated using Oxford

197 Nanopore Technologies (ONT) MinION devices, using isolates with matched ONT and Illumina

198 data.

199 **Implementation**

200 As outlined in the Introduction, GenoTyphi is a typing scheme specific to Typhi, which specifies

201 marker SNVs suitable to identify specific lineages. In principle GenoTyphi can be used with any

202 genotyping tool that is capable of searching for the marker SNVs in sequence data. GenoTyphi

203 provides the framework directing genotyping tools what SNVs to look for and how to interpret

204 them in terms of lineages to report. Mykrobe is a genotyping tool, which uses a *k*-mer-based

205 approach to identify marker SNVs in sequence reads. It was developed for typing

206 *Mycobacterium tuberculosis* and *Staphylococcus aureus* [32–34], but can in principle be used

207 with any genotyping scheme. Here we present a ready-to-run code base that allows users to

208 type genomes from raw reads, using Mykrobe, against the GenoTyphi framework to identify and

209 report known lineages – which we label "Typhi Mykrobe". This tool additionally includes AMR

210 and plasmid typing frameworks, and generates predictions of resistance to the drugs

211 recommended for clinical treatment of typhoid. Details of this Implementation are described

212 below.

214 **Genotyping targets**

215 The current scope of the GenoTyphi scheme as discussed here (n=87 genotypes) is illustrated

216 in **Supplementary Figure 1**. This includes five new genotypes (3.5.4.1, 3.5.4.2, 3.5.4.3,

217 4.3.1.2.1, 4.3.1.2.1.1) recently described in a technical report [28]. The full scheme specification

is available in the GenoTyphi repository (https://github.com/typhoidgenomics/genotyphi) in the

file Genotype_specification.csv.

The AMR and plasmid targets included in Typhi Mykrobe are summarised in **Figure 1**. These

include sequence probes targeted to detect presence/absence of epidemiologically relevant

mobile AMR determinants [13,14], conferring resistance to the currently recommended drugs

ciprofloxacin, ceftriaxone, and azithromycin, as well as older drugs ampicillin, chloramphenicol,

trimethoprim-sulfamethoxazole, and tetracycline, as well as carbapenems. Fourteen SNVs

associated with resistance to ciprofloxacin and azithromycin are also included. In addition, Typhi

Mykrobe includes probes to detect the presence or absence of 13 plasmid replicons previously

reported in Typhi [14], including 11 associated with AMR plasmids, the cryptic plasmid pHCM2,

and linear plasmid pBSSB1, which carries the alternative flagellin z66 [35]. A marker SNV to

identify the plasmid sequence type 6 (PST6) lineage of the dominant IncHI1 plasmid is also

included [36]. The full list of AMR targets is available in the GenoTyphi repository

(https://github.com/typhoidgenomics/genotyphi) in the file AMR_genes_mutations_plasmids.csv.

Implementation in Mykrobe

The Mykrobe software, first developed in 2015 for *Mycobacterium tuberculosis* and

Staphylococcus aureus [32], uses a *k*-mer-based approach to identify marker SNVs and the

presence or absence of target genes using 'probe' sequences to screen against [31]. These

data are then used to assign hierarchical genotypes and report AMR determinants. Probe sets

have been previously developed and implemented for other bacterial pathogens including

Mycobacterium tuberculosis [32–34], *Staphylococcus aureus* [31], *Shigella sonnei* [37], and

Salmonella Paratyphi B [38]. Mykrobe is available at https://github.com/Mykrobe-tools/mykrobe.

Mykrobe probe sets were created for genotype and AMR marker SNVs using the *mykrobe*

variants make-probes command using a *k*-mer size of 21 and the Typhi CT18 reference

genome (accession: AL513382.1). The nested hierarchical relationships of the marker SNVs

were specified and Mykrobe used this information to identify the best supported genotype call

(e.g., a call of '4.3.1.1' is typically supported by detection of nested markers that demarcate

different levels of the hierarchy: 4, 4.3.1, 4.3.1.1). Briefly, probe sets for gene presence or

absence were created using a Python script that extracted the relevant AMR sequences. Details

of the specific commands and Python scripts are provided in

https://github.com/typhoidgenomics/genotyphi. Probe sets were also created to confirm input

read sets as *S. enterica* using the presence of marker gene *invA*, involved in invasion of

intestinal epithelium cells by *S. enterica*, and serovar Typhi based on the 26 known multi-locus

sequence types (STs) (see **Figure 1**). Further details of the preliminary testing and

implementation of the Typhi probe sets are available in the technical report

https://zenodo.org/doi/10.5281/zenodo.7407984. The Typhi Mykrobe probe sets and typing

panel are available at https://doi.org/10.6084/m9.figshare.27102334.

Summarising Mykrobe outputs

To support interpretation of the Mykrobe output, we developed a Python-based parser script

parse_typhi_mykrobe.py, which takes as input a set of JavaScript Object Notation (JSON) files

output by Mykrobe, and summarises them in a tab-delimited table with one row per genome,

and columns to indicate the genotype calls (see **Figure 1**). Also included are several quality-

control fields reported by Mykrobe, including read-level support for each marker SNV

contributing to the final genotype call ('node support'), and for any additional markers detected.

This information is helpful for identifying contaminated samples (see examples in Results). The

parser script also assigns a summary 'confidence' level in the final GenoTyphi lineage call,

based on the Mykrobe quality scores for each nested component marker that makes up the final

genotype call. In brief, a weak' confidence call is assigned when one or more marker SNVs are

of low quality, defined as Mykrobe quality score of 0 or Mykrobe quality score of 0.5 with

minority support (i.e., <50% of reads) for the derived allele. A 'moderate' confidence call is

assigned if one (and only one) marker SNV has a Mykrobe quality score of 0.5 but with majority

support (i.e. ≥50% of reads) for the derived allele. Otherwise, the genotype call is assigned

'strong' confidence.

The parser script summarises the AMR markers detected by Mykrobe in the form of a pseudo-

antibiogram with one column per drug. Each drug column contains either 'S' (indicating no

markers detected) or 'R' (detected markers). For ciprofloxacin, it is useful to distinguish between

low-level resistance (MIC 0.06-0.5 mg/L), which is associated with a single mutation or acquired

gene and high-level resistance (MIC >0.5 mg/L), which is associated with combinations of two

or more determinants [14,17], because while both groups are associated with increased fever

clearance times and probability of treatment failure compared with susceptible strains, high-level

resistance is associated with worse outcomes than low-level resistance [39]. Current guidelines

and terminology for these groups vary, but for simplicity, the Mykrobe parser reports these

groups as 'I' (for intermediate) and 'R', respectively. This matches the current CLSI guidelines

(https://clsi.org/standards/products/microbiology/documents/m100/). However, it should be

noted that (i) EUCAST defines the clinical breakpoint for 'R' in *Salmonella* spp. as MIC >0.06

mg/L (https://www.eucast.org/clinical_breakpoints), which includes both low-level and high-level

resistance; and (ii) previous literature has referred to the low-level range as 'decreased

ciprofloxacin susceptibility' rather than 'I' [40]. In addition to this antibiogram-style summary,

individual AMR and plasmid markers are reported in their own columns, coded as 1=present

and 0=absent (as seen in **Supplementary Table 1,** available at

https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe).

All code and instructions for genotyping isolates with GenoTyphi using Typhi Mykrobe are freely

available in the GitHub repository at https://github.com/typhoidgenomics/genotyphi (DOI:

10.5281/zenodo.13859721).

Validation data

Validation on Illumina short-read data was conducted using the Global Typhoid Genomics

Consortium dataset described in Carey et al [14] . Reads were mapped to the CT18 reference

genome (accession: AL513382) [29] using bwa-mem v0.7.17 via the Centre for Genomic

Pathogen Surveillance (CGPS) mapping pipeline v1.2.2

(https://gitlab.com/cgps/ghru/pipelines/snp_phylogeny), to generate alignments in BAM format.

Assemblies were generated using the CGPS assembly pipeline v2.1.0

(https://gitlab.com/cgps/ghru/pipelines/dsl2/pipelines/assembly/) which utilises the SPAdes

assembler (v3.12.0) [41,42]. A total of 141 isolates were excluded due to excessive unclustered

heterozygous SNVs indicative of mixed cultures (n=92), assembly issues (n=46) or both (n=3),

resulting in 12,839 isolates for Mykrobe validation (listed in **Supplementary Table 2,** available

at https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe). The genomes of

isolates called as 'not Typhi' were characterised for presence of the marker genes of *invA* and

the seven MLST genes using Blast in Bandage for *invA* [43] and MLST with the 'senterica'

database (https://github.com/tseemann/mlst). The dataset covered 85 of the 87 defined

genotypes (exceptions being genotypes 1 and 2.3, which represent internal nodes of the

hierarchy for which no genomes have been observed). Individual calls are given in

Supplementary Table 1 and **Supplementary Table 2.**

Validation of genotype calls

Typhi Mykrobe v0.12.1 was run on all (n=12,839) Illumina FASTQ files using the new Typhi panel (v20221208) and the results were summarised using *parse_typhi_mykrobe.py*. To validate Typhi Mykrobe's GenoTyphi genotype calls, we compared them with those called by the original mapping-based approach. To do this, we used BAM file read alignments generated from the same n=12,839 FASTQs as described above as input to the Python script genotyphi.py (v2.0; available at https://github.com/typhoidgenomics/genotyphi) [18,22,28]. To validate the AMR and plasmid calls, we compared them to those called in corresponding genome assemblies by Pathogenwatch (available for n=11,992 genomes, assemblies for the other genomes were excluded as they did not meet Pathogenwatch quality-control criteria [13]). We downloaded the GenoTyphi, AMR, and plasmid replicon typing reports generated by Basic Local Alignment Search Tool (BLAST) searching assemblies for these genomes from Pathogenwatch on 7 March 2023 (GenoTyphi v20221208 specification for lineage genotypes). Categorical agreement (percentage of genomes where both tools agreed on the same call of presence or absence) was calculated separately for each marker, and collectively by drug (percentage of genomes where both tools agreed that at least one marker was present or that no markers were present) (**Supplementary Table 3,** available at https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe). Probes for plasmid-borne carbapenemase genes circulating in Enterobacterales (*bla*NDM-5, *bla*KPC-2, *bla*VIM-1, *bla*IMP-27, *bla*OXA-48) were added to GenoTyphi v20240407 (DOI: 10.5281/zenodo.13859721) following the report of *bla*NDM-5 in Typhi from Pakistan [11], and validated using the genomes of the two distinct morphologic variants from that study (accessions: SRR22801766 and SRR22801806). Comparisons and statistical analyses were conducted in R v4.1.0 using tidyverse v1.3.1 [44,45] .

Benchmarking of run time for Mykrobe

We further investigated the average run time for Typhi Mykrobe to demonstrate the rapid time to a result. Here we used the short-read data from 100 Typhi genomes. These were randomly selected but included at least one genome from each genotype and a diversity of AMR and plasmid profiles. The `mykrobe predict` command was run for the 100 Typhi genomes on three different computers where both the run time and RAM were recorded. The number of threads was also varied from 1 to 64 threads for the two HPC and 1 to 16 for the Mac laptop.

Accuracy of AMR phenotype prediction

Antimicrobial susceptibility testing (AST) data were available for n=3,970 of the Illumina-

sequenced isolates, sourced from three separate datasets (**Supplementary Table 4,** available

at https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe). Minimum

inhibitory concentration (MIC) data were available from the UK Health Security Agency

(UKHSA) for n=852 isolates following EUCAST standards and interpretive criteria (v10, 2020)

[46] and the US Centers for Disease Control and Prevention (US CDC) for n=720 isolates

following CLSI standards and interpretive criteria (M100, 2023) [47]. Disk diffusion data

following EUCAST standards and interpretive criteria (v8.0, 2018) were available for n=2,446

isolates collected and analysed from three countries as part of the Surveillance for Enteric Fever

in Asia Project (SEAP), with multiple source labs contributing AST data and isolates for

sequencing [6] . Typhi Mykrobe AMR calls following the S/I/R categorisation described above

were compared with the phenotypic antimicrobial susceptibility S/I/R categorisation using the

relevant CLSI or EUCAST standards for each data set, for the currently recommended drugs

azithromycin and ciprofloxacin and ceftriaxone, and the older drugs ampicillin, chloramphenicol,

trimethoprim-sulfamethoxazole, and tetracycline (note UKHSA tested amoxicillin rather than

ampicillin; and SEAP did not test tetracycline). To assess concordance between genotype and

phenotype, we followed the principles used for assessment of new AST devices by the US Food

and Drug Administration (FDA), which considers (i) categorical agreement, (ii) major error rate

(defined as the proportion of S isolates that test R with the new method [48,49]), and (iii) very

major error rate (defined as the proportion of R isolates that test S with the new method)

(**Supplementary Table 5,** available at

https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe). The FDA standard

requires >90% overall categorical agreement (within one doubling dilution), <3% major errors for

each drug, and <1.5% very major errors for each drug; hence, we adopted these thresholds to

assess the accuracy of the genotype-based prediction of phenotype. For drugs with high error

rates, we investigated the relationship between individual resistance markers and phenotype,

which we visualised using upset plots generated in R v4.1.0 using the ComplexUpset package

v1.3.3 (10.5281/zenodo.3700590). For five isolates with unexplained trimethoprim-

sulfamethoxazole resistance, we used nucleotide blast search of the genome assemblies to

confirm the presence of wild-type chromosomal *folA* and *folP* (100% nucleotide match to

reference sequences from Typhi strain Ty2) and screened the assemblies for other AMR

determinants using CARD RGI v6.0.3 [50].

Matched ONT and Illumina validation data

For 98 isolates, long-read ONT data were available in addition to the Illumina data; these included 68 isolates from Carey et al 2023 [14] and 30 novel isolates from this study. Sample-level details including accessions and technical specifications such as library, flow cell, device, basecaller, and ONT processing (including demultiplexing, trimming and filtering), are available in **Supplementary Table 6,** available at https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe. These varied between isolates including different flow cells (e.g., R9.4.1 and R10.4.1) and base calling methods. Read depth for each ONT dataset was assessed by using seqtk v1.2-r94 (https://github.com/lh3/seqtk) to determine total number of bases per read set, which was divided by the length of the CT18 reference genome (4,809,037 bases). For five genomes with read depth <30×, Kraken2 was used for the initial read-level classification of ONT data [51]. These five genomes were also aligned to the CT18 reference genome using minimap2 [52,53] to determine the read coverage of the *invA* and the seven core genes in the *Salmonella enterica* MLST scheme [54]. ONT reads were assembled for these five genomes and in addition to four genomes to resolve discrepant plasmid calls. Assembly was done using Flye v2.9.1-b1780 [55] with default parameters and yielding genome sizes of the expected 4.8 Mbp. Due to the high read depth (784×), the ONT data for SRR17299234 was subsampled prior to assembly using Filtlong v0.2.1 (https://github.com/rrwick/Filtlong) to --target_bases 480000000 (resulting in 99.81× depth). The genomes of isolates called as 'not Typhi' were characterised for presence of the marker genes of *invA* and the seven MLST genes using Blast in Bandage for *invA* [43] and mlst with the 'senterica' database (https://github.com/tseemann/mlst).

Results

Validation of GenoTyphi genotype calling

To validate Typhi Mykrobe for assigning hierarchical GenoTyphi genotypes, we ran it on n=12,839 Illumina readsets (covering all existing types) and compared the resulting genotypes to those called using the original mapping-based implementation. Five genomes were typed as 'not Typhi' by Mykrobe **(Supplementary Table 1**)**.** Four of these genomes lacked the *invA* gene but MLST profiles were detected that were consistent with those on Pathogenwatch. In the remaining isolate, DRR071001, the *invA* gene was detected but no allele was reported for *hisD* in the MLST scheme, consistent with the MLST profiles from Pathogenwatch. Hence, these five isolates were reported as 'not Typhi' and not analysed further. Of the remaining n=12,834

genomes, Mykrobe called the same single genotype as the mapping-based approach for

n=12,807 (99.79%). For a further n=17 genomes, Mykrobe and the mapping-based pipeline

gave concordant reports of two genotypes being present, consistent with mixed samples (in the

Typhi Mykrobe output this is reported as one primary genotype plus additional markers, while in

the mapping-based pipeline the result is reported as a list of supported genotypes). Eight

genomes belonged to genotype 3.5.3, which was nested in 3.5.4, breaking the usual

hierarchical structure of the scheme. Typhi Mykrobe correctly reported these genomes as

genotype 3.5.3 with support for both markers 3.5.4 and 3.5.3, as it has this nested relationship

explicitly encoded in its hierarchy; whereas the mapping-based pipeline does not have this logic

and incorrectly reported these genomes as genotype 3.5.4.

There were two instances of genuine disagreement in the genotype calls generated by Typhi

Mykrobe as compared to the mapping-based approach. One isolate (ERR5243665) was called

as genotype 2 by Mykrobe with strong confidence (no reads support for any alternative alleles

or markers). It was genotyped as 0.1.3, albeit with low support, by the mapping-based

approach, but the genotype 2 marker was also detected. This isolate clustered with other

genotype 2 isolates in a distance-based phylogeny of genomes from the same study [6] (see

tree at https://pathogen.watch/collection/0z5knw9jic9b-guevara-et-al-2021), suggesting that the

Mykrobe call is correct. The other isolate (ERR2663783) was called as genotype 4.3.1.1 by

Typhi Mykrobe with strong confidence. It was reported as "4.3.1.1,4.3.1.3.Bdq" (with low

support) by the mapping approach; however, the support for the 4.3.1.3.Bdq marker was low.

This isolate clustered closely with other 4.3.1.1 isolates from the same study [25](see tree at

https://pathogen.watch/collection/b2k6dym3bdyq-tanmoy2018), which is consistent with the

Mykrobe call.

We demonstrate that the run time for Typhi Mykrobe's `mykrobe predict` command on a modern

computer was <1minute to complete (**Supplementary Figure 2**). The `mykrobe predict`

command can be run with multiple threads using the `--threads`/`-t` option. Up to ~4 threads will

increase performance. Using multiple threads on a very fast CPU, the run time for each genome

was reduced to seconds (**Supplementary Figure 2**). Further, `mykrobe predict` is very

memory-efficient and will typically use less than 100 MB of RAM per genome.

Validation of AMR genotyping

To assess the accuracy of Mykrobe for detecting the presence of AMR genes and mutations, we compared genotype calls from Mykrobe (called from *k*-mer analysis of reads) with those from Pathogenwatch (called from BLAST analysis of genome assemblies of the same read sets). We found very strong categorical agreement between the two methods at the individual marker level, with 99.88% of the 299,004 marker-genome combinations yielding the same call (summarised in **Table 2,** full details in **Supplementary Table 2**). High concordance was evident for all genetic markers of AMR (98.56–100%, **Table 2**), as well as plasmid replicon markers (99.74–100%, **Table 3**). The test dataset included all AMR and plasmid markers in GenoTyphi v2.0 implemented Typhi Mykrobe v0.12.1. The *ereA*, *ermB* and carbapenemases are very rarely reported in Typhi but were included in the Mykrobe panel in case they emerge.

Both Mykrobe and Pathogenwatch summarise AMR data in terms of a binary phenotype categorization (S/R), assuming the presence of a marker associated with resistance to a drug implies resistance 'R', with the exception of ciprofloxacin, where presence of any marker is categorised as 'I' and specific combinations of markers are categorised as 'R', resulting in three categories S/I/R (see Methods). Categorical agreement was very high between the drug-level categorisations reported by both genotyping tools, with >99.7% agreement (n=2 to 33 discordant genomes) for each drug (**Table 2, Supplementary Table 3**), and for subsequent categorisation as MDR (99.86%, n=17 discordant) or XDR (99.97%, n=4 discordant).

Validation of AMR phenotype prediction

To assess the accuracy of Mykrobe's genotype-based clinical resistance categorisations, we compared them with AST data, which was available for a subset of n=4,018 isolates (see Methods, **Supplementary Table 4**). These isolates were from three data sets, using different methods (MIC with EUCAST standards, MIC with CLSI standards, disk diffusion with EUCAST standards), hence we analyzed the results from each dataset separately. Categorical agreement was high (>95%) for all drugs across all three datasets, and very high (>99%) for the two sets of reference-laboratory MIC measures (**Figure 2, Supplementary Figure 5**).The only exception was US CDC categorisation of low-level ciprofloxacin resistance (using the CLSI 'I' threshold, ≥0.125 to 0.5 mg/L), which showed 97.36% agreement with Mykrobe calls of 'I' or 'R'. This was mainly due to the presence of major errors (susceptible isolates with AMR determinants detected, thus reported as R by Mykrobe, see Methods), whereby n=17 isolates tested susceptible to ciprofloxacin (n=14 with MIC=0.06 mg/L, n=3 with MIC=0.015 mg/L) but

Mykrobe detected a known QRDR mutation (**Figure 3b**); categorical agreement within one
doubling dilution was 99.4%.

Major errors were infrequent (<3% prevalence) for all drugs in all datasets (**Figure 2**), except for
ciprofloxacin (**Supplementary Table 5**). Most of the ciprofloxacin major errors were in isolates
whose MIC or disk zone measurement were close to the breakpoint (**Figures 3, 4c**); all carried
well-characterized molecular resistance determinants (QRDR mutations and/or *qnr* genes, see
Figures 3, 4c), and in all but one case the same determinants were also detected in the
corresponding genome assemblies by Pathogenwatch (details below).

Very major errors (resistant isolates with no AMR determinants detected, thus reported as S by
Mykrobe) were also infrequent (<1.5% prevalence) for the reference-laboratory MIC datasets,
but errors were higher in the disk diffusion dataset (**Figure 2**). The only drugs showing major or
very major errors compared with the reference-laboratory MIC data were ciprofloxacin and
trimethoprim-sulfamethoxazole (**Figure 2**). For ciprofloxacin, the US CDC categorisation of
high-level resistance (using the CLSI 'R' threshold, ≥1 mg/L) yielded four very major errors (R
isolate called as 'I' by Mykrobe). Two had a single mutation detected (GyrA-S83F) and two had
double QRDR mutations (GyrA-S83F plus GyrA-D87N or ParC-S80I) (see **Figure 3**);
Pathogenwatch calls agreed in all instances. Importantly, all four isolates had MIC=1 mg/L, i.e.
they agreed with the genotype-based call within one doubling dilution. UKHSA uses the
EUCAST standards, which have a single (low-level) breakpoint for ciprofloxacin (R >0.06 mg/L),
and this yielded just two major errors (2.9%) and two very major errors (0.26%) (see **Figure 3**).

For trimethoprim-sulfamethoxazole, Mykrobe analysis of the US CDC dataset yielded a single
error (R isolate called as S) but the MIC was 4 mg/L, i.e. within one doubling dilution, and the
isolate had no *sul* or *dfr* genes that were identified by Mykrobe or Pathogenwatch. Mykrobe
analysis of the UKHSA dataset yielded five errors for trimethoprim-sulfamethoxazole; all were R
isolates called as S by Mykrobe, which detected *sul* genes but no *dfrA* genes. Interestingly, all
five isolates tested susceptible to trimethoprim alone (MIC ≤0.5, well below the >4 mg/L
breakpoint), supporting the absence of *dfrA* genes; these were also not detected by
Pathogenwatch analysis of the corresponding genome assemblies, or by UKHSA's internal
pipeline [44]. Further investigation of these genome assemblies confirmed the presence of wild-
type *folA* and *folP*, and no additional determinants were reported by CARD RGI (v6.0.3).

Mykrobe analysis of the SEAP dataset showed lower overall categorical agreement with AST

phenotypes (95.3–98.2%). As the testing was done using disk diffusion, it was not possible to

assess the categorical agreement within one doubling dilution of a MIC. However, most of the

errors were within 2–4 mm of the breakpoint (see **Figure 4**), and in all but one instance, the

Mykrobe genotype result matched that called by Pathogenwatch from the corresponding

genome assemblies. The only exception was a ciprofloxacin resistant isolate (ERR4325960),

which Mykrobe reported as carrying GyrA-S83F with no other determinants, but Pathogenwatch

detected both GyrA-S83F and *qnrS* in the assembly.

Validation on nanopore reads

We explored the accuracy of Typhi Mykrobe for genotyping ONT long-read data by comparing

results from 98 pairs of ONT to Illumina reads generated from the same isolates. Data were

contributed by members of the Global Typhoid Genomics Consortium and generated using

different library preparation methods and sequencing devices (see **Supplementary Table 6**).

Mykrobe called one Illumina read set and five ONT read sets as non-Typhi (mean ONT read

depths estimated as 0.04×, 4.06×, 8.01×, 26.40×, 27.85×, all sequenced on R9 flowcells), these

were excluded from analysis of genotype accuracy as no genotype is called on isolates

identified as non-Typhi.

Among the n=92 Typhi genomes called from both ONT and Illumina reads, the Typhi Mykrobe

genotype lineage calls matched for n=89 (96.7%). For all three discrepant isolates, (all

sequenced on R.9 or R.9.4 flowcells), the set of SNV markers detected from ONT reads closely

matched those detected from Illumina reads, but with one or two markers missing from the

ONT-based calls and no unexpected markers reported resulting in final genotype calls of 4.3.1.1

vs 4.3.1, 4.3.1.2 vs 4.3.1, 2.1.7.2 vs 2.1.7. For the missing markers, Mykrobe reported low and

variable ONT read depths, with ranges 3–9, 4–31 and 1–14, suggesting that the errors are likely

due to lack of informative ONT reads at the marker loci. Notably, the overall ONT read depths

for these were reasonable (mean 35×, 170× and 90× respectively), suggesting the lack of

informative reads for genotyping may be caused by contamination from non-Typhi reads, or

high error rates resulting in low rates of *k*-mer matching.

In this study, the five ONT genomes that had 'not Typhi' calls were further explored. These were

all classified as *Salmonella enterica*. However, Typhi was not the top serovar from read-level

classification, instead *Salmonella* Infantis was the top serovar for BZD62J, while *Salmonella*

Typhimurium was the top serovar for the remaining four. Further, the presence of the marker
genes of *invA* and the seven MLST genes were detected in the genomes at read depths
consistent with the genome mean read depth reported in **Supplementary Table 6**, with the
gene *invA* was detected in three of the five genomes from Blast analysis of the Flye assemblies.
The gene *invA* was not detected in the 611427 ONT assembly and only had a read depth of
0.43×, and BZD62J failed to assemble due to insufficient reads and was found to have no reads
mapping to *invA*. MLST analysis of the three complete genomes with *invA* detected reported,
were found to have partial matches to known alleles for all seven MLST genes and no reported
MLST profile. Of note, these five genomes had low read depth (0.04–28×), sequencing on older
R9 flowcells, and basecalling with older versions of guppy (**Supplementary Table 6**). The
failure of Typhi Mykrobe to identify these as Typhi is likely due to a combination of potential
contamination, low read depth and high error rates resulted in low rates of *k*-mer matching to
the probes. This highlights the importance of QC of the read data, as Typhi Mykrobe provides
some metrics for confidence of the calls, but does not itself perform detailed QC.
556
AMR marker detection from ONT reads was highly concordant with Illumina-based calls, with
98.77% overall agreement across the n=736 isolate-drug combinations (92 isolates across 8
drugs; see **Supplementary Table 7**). There were nine genomes with disagreement of AMR
calls; five of these had a marker detected in Illumina reads but not ONT reads (n=1 *catA1*, n=1
qnr, n=3 QRDR mutations), three had GyrB-464F detected in ONT reads but not Illumina, and
one had GyrB-S464F detected in Illumina reads and GyrA-S83F in ONT. In all nine cases, the
Typhi Mykrobe call from Illumina reads matched that from Pathogenwatch analysis of the
corresponding assemblies of Illumina reads. There was no overlap between ONT read sets with
AMR discrepancies and those with lineage genotype discrepancies. Overall, ONT-based
resistance prediction showed complete agreement with Illumina for six of the eight drugs, 98.9%
agreement for chloramphenicol resistance, and 91% for ciprofloxacin resistance.
568
Plasmid marker detection from ONT data was also highly concordant with Illumina-based calls,
with 99.67% overall agreement across the n=1,196 isolate-marker combinations (92 isolates
across 13 markers; see **Supplementary Table 8**). Four genomes had discrepant plasmid
marker calls, all with a marker detected from Illumina reads but not from ONT reads. One isolate
(DRR070993) had a typical IncHI1 MDR plasmid profile reported from Illumina reads (IncHI1A,
IncHI1BR27, IncHI1_ST6 indicating presence of IncHI1-ST6 plasmid, with *catA1*, *dfrA7*, *bla*TEM-1,
sul1, *sul2*), but the ONT read profile was missing the *IncHI1A* and *catA1* markers (and missing

catA1 in the corresponding Illumina assembly and a mismatch for *IncHI1A*). The other three

isolates had pHCM2 detected in the Illumina reads and in the corresponding assemblies [14] but

not in ONT reads; for these isolates the ONT and Illumina sequencing was done on different

DNA extracts so this may reflect plasmid loss in the culture used for ONT sequencing.

Discussion

Typhi Mykrobe provides detection and genotyping of key features of clinical and public health

importance, including assigning lineages, AMR determinants, and plasmid replicons (**Figure 1**).

Compared with other tools for Typhi genotyping, Mykrobe has the advantage of not requiring

any pre-processing, aligning, or assembling of reads, but instead takes as inputs raw FASTQ

files. This simplifies bioinformatics workflows and reduces time-to-result as Typhi Mykrobe can

return a full genotype profile in typically <1 minute. Importantly, although Typhi Mykrobe is not a

QC tool *per se*, it provides useful data on the read-level support for individual genotype markers,

which can be used to help identify potentially mixed samples. As a command-line tool that is

easy to install locally using Bioconda, Typhi Mykrobe is suitable for research and public health

laboratory settings, facilitates standardized reporting of Typhi genotypes without the need to

share sequence data, and enables broader access to bioinformatics functionality across a

variety of settings.

Typhi Mykrobe showed very high genotyping accuracy, for GenoTyphi lineage assignment

(**Table 1**), and detection of AMR and plasmid markers (**Tables 2-3**), compared with existing

genotyping tools. Of note, the included IncY, pO111 and IncFIB for the cryptic pHCM2 replicon

targets have recently been shown to be sometimes carried on phage-plasmids [56,57], in

addition to conjugative plasmids such as the IncY plasmids associated with XDR Typhi [26].

Future work is required to identify additional targets to delineate the phage-plasmids, and the

Typhi Mykrobe software can be readily updated to include new markers.

In addition, we show that the panel of AMR markers currently included in Typhi Mykrobe can be

used to predict AMR phenotypes with high accuracy, demonstrating very high concordance with

S/I/R calls made from reference-laboratory MICs (**Figure 2**). Concordance was slightly lower

when compared with research laboratory disk diffusion data. However, as the phenotypic

resistance prevalence rates were similar across the three datasets, it is reasonable to assume

that the higher error rates in the disk diffusion dataset compared with the two reference-

laboratory MIC datasets most likely reflect comparatively lower accuracy of the disk diffusion

phenotype measurements, which are known to yield lower accuracy for some drugs such as

azithromycin [58,59] and in this case were completed across multiple laboratories. There are

currently no formal standards for assessing the accuracy of AMR phenotype predictions from

genomic data[60]; however, applying the standards for FDA licensing of new AST devices,

Typhi Mykrobe easily passed the target threshold for categorical agreement (>99% agreement

within one doubling dilution for reference-laboratory MIC data, >95% agreement for disk-

diffusion data; vs >90% target) (**Figure 2**).

Typhi Mykrobe also showed very low rates of major errors and very major errors compared with

the reference-laboratory MIC data, and for nearly all drugs these fell below the acceptable

thresholds for a new AST device (<3% and <1.5%, respectively) (**Figure 2**). The exception was

trimethoprim-sulfamethoxazole, for which five UKHSA isolates tested resistant but Typhi

Mykrobe predicted as susceptible due to a lack of *dfrA* genes, which are thought to be required

for resistance to trimethoprim-sulfamethoxazole along with *sul* genes, which were detected in

these genomes. Other genotyping tools (Typhi Pathogenwatch, CARD RGI) also could not

identify any known mechanisms of resistance to trimethoprim or trimethoprim-sulfamethoxazole

in these genomes, and the isolates tested sensitive to trimethoprim. Therefore, we hypothesise

these isolates may harbour efflux mutations or a novel mechanism of resistance to

trimethoprim-sulfamethoxazole that does not result in resistance to trimethoprim alone. Further

analyses to determine the exact resistance mechanism was beyond the scope of this study.

Ciprofloxacin showed very low error rates in the UKHSA MIC data, interpreted using the single

low-level resistance threshold (MIC >0.06 mg/L) recommended by EUCAST, but slightly higher

error rates for the US CDC MIC data, which was interpreted using low-level and high-level

resistance thresholds (although still below the acceptable thresholds within one doubling

dilution). Carbapenemase genes were added to the Typhi Mykrobe panel, but this resistance is

newly emerging and there was only a single carbapenem-resistant isolate sequenced at the

time of testing (in which *bla*NDM-5 was correctly identified by Typhi Mykrobe), therefore we

could not yet assess accuracy of phenotype prediction for carbapenems.

Finally, we demonstrated that Typhi Mykrobe can be used to genotype ONT reads, showing

high concordance with results obtained from the analysis of Illumina reads generated from the

same isolates (96.74% agreement on GenoTyphi lineage calls, 98.77% on AMR markers and

99.67% on plasmid markers). We did not have suitable data to directly assess ONT-based AMR

predictions with AST data. ONT-based and Illumina-based drug-level predictions of AST

phenotypes showed high agreement; however, future studies assessing ONT data compared to
reference-laboratory MIC data would be useful to confirm the accuracy of ONT data for
phenotype prediction. As one might expect, many of the errors occurred in isolates with lower
read depth. However, as the matched ONT-Illumina data available for testing were contributed
by different laboratories that used different ONT protocols and devices for library preparation,
sequencing, and base calling, we were unable to perform a detailed assessment of the impact
of read depth and quality on genotyping performance. The accuracy of ONT reads has greatly
increased in recent years and is approaching Illumina read accuracy of ~99%; hence modern
ONT data would be expected to perform similarly to Illumina data [61]
(doi.org/10.5281/zenodo.10397817). Future studies assessing the impact of read depth and
base calling methods on the performance of Typhi Mykrobe on ONT data would be desirable to
help guide laboratories on the most appropriate methods and target yield for ONT experiments
aiming to produce reliable genomic surveillance data for Typhi.

Conclusions

Typhi Mykrobe provides rapid and accurate genotyping of Typhi genomes directly from Illumina
reads, including (i) assignment of GenoTyphi lineage; (ii) detection of AMR determinants and
prediction of corresponding AMR phenotypes demonstrating >99% categorical agreement with
reference-laboratory MIC data; and (iii) detection of plasmid replicons. The software also
performs well on direct genotyping from ONT reads information, although further investigation is
needed to assess the accuracy of AMR phenotype prediction.

Other sections

List of abbreviations

AMR: antimicrobial resistance
AST: antimicrobial susceptibility testing
CARD: Comprehensive Antibiotic Resistance Database
CLSI: Clinical and Laboratory Standards Institute
EUCAST: The European Committee on Antimicrobial Susceptibility Testing
FDA: United States Food and Drug Administration
MDR: multidrug resistance
MIC: minimum inhibitory concentration

ONT: Oxford Nanopore Technologies

QRDR: quinolone resistance-determining regions

SEAP: Surveillance for Enteric Fever in Asia Project

SNV: single nucleotide variant

UKHSA: United Kingdom Health Security Agency

US CDC: United States of America Centers for Disease Control and Prevention

WGS: whole genome sequencing

XDR: extensively drug resistant

Availability and requirements

Project name: GenoTyphi

Project home page: https://github.com/typhoidgenomics/genotyphi

Operating system(s): Platform independent

Programming language: Python3 or higher

Other requirements: Mykrobe

License: GNU General Public License

Any restrictions to use by non-academics: None

Project name: Mykrobe

Project home page: https://github.com/Mykrobe-tools/mykrobe

Operating system(s): Platform independent

Programming language: Python3 or higher

Other requirements: anytree 2.8.0 or higher, Biopython 1.79 or higher, PyVCF3 1.0.3 or higher,

requests 2.27.1 or higher, mongoengine 0.24.1 or higher, cython 0.29.28 or higher, numpy

1.22.0 or higher, pytest

License: MIT License

Any restrictions to use by non-academics: None.

Declarations

Ethics approval and consent to participate

Each contributing study or surveillance program obtained local ethical and governance

approvals, as reported in the primary publication for each dataset. Inclusion of data that were

not yet public domain by August 2021 was approved by the Observational / Interventions

Research Ethics Committee of the London School of Hygiene and Tropical Medicine (ref

#26408), on the basis of details provided on the local ethical approvals for sample and data

collection (listed in Carey et al, 2023, Supplementary file 1).

Consent for publication

Not applicable

Availability of data and materials

All code and instructions for genotyping isolates with GenoTyphi using Mykrobe is available in

the GitHub repository at https://github.com/typhoidgenomics/genotyphi. The version reported in

this paper is v2.1 (DOI: 10.5281/zenodo.13859721)

The datasets and analysis code used in this article are available in a GitHub repository at

https://github.com/typhoidgenomics/TyphoidGenomicsConsortiumMykrobe.

Competing interests

MML was a co-developer of a Trivalent Salmonella (Enteritidis/Typhimurium/Typhi Vi) conjugate

vaccine with Bharat Biotech International and the Wellcome Trust. MML has also received

payments from Pfizer for consultancy work. MML holds US patents for "Compositions and

Methods for Producing Bacterial Conjugate Vaccines". MML was a member of a NIH DSMB that

oversaw US government-funded efficacy trials of COVID-19 vaccines. DSMB was disbanded

after several vaccines were given Emergency Use Authorization. MML was a member of the

Vaccines and Related Biological Products Advisory Committee of the FDA. IIB was a consultant

for the Weapons Threat Reduction Program, Global Affairs Canada. AJP has been involved an

Oxford University partnership with AZ for development of COVID-19 vaccines. AJP has received

payments for consultancy work from Shionogi. AJP is chair of DHSC's Joint Committee on

Vaccination and Immunisation, is a chair of WHOs Salmonella TAG, and was a member of

WHOs SAGE. AJP received support from MRNA – Moderna. KLC has received payments from

Pfizer for presentations and travel support from BD. INO has received payments from the

Wellcome Trust for consultancy work and receives royalties for books or book chapters

published via Springer, Cornell University Press, and Oxford University Press. INO has received

travel support from BMGF, ESCMID, and the American ASM and has held leadership or

advisory roles for Wellcome SEDRIC, the BMGF surveillance advisory group, the Thomas

Bassir Biomedical Foundation, and International Centre for Antimicrobial Resistance Solutions

(ICARS) Technical Advisory Forum. ZI has received travel support from ETH Zurich.

Funding

DJI was supported by an Emerging Leadership Fellowship from the National Health and Medical

Research Council (NHMRC) of Australia (GNT1195210). ZAD received funding from the

European Union's Horizon 2020 research and innovation programme under the Marie

Sklodowska-Curie (grant agreement No 845681). KEH received funding from the Wellcome

Trust (grant 226432/Z/22/Z). INO was supported by Official Development Assistance (ODA)

funding from the NIHR (grant number #16/136/111) and a Calestous Juma Science Leadership

Fellowship from the BMGF (INV-036234). AMS is supported by the SEQAFRICA project, funded

by the Department of Health and Social Care's Fleming Fund using UK aid. MJS was supported

by BMGF (INV-000049, OPP1161058) and National Institute of Allergy and Infectious Diseases

(NIAID) of the National Institutes of Health (F30AI156973). JAC was supported by the Bill &

Melinda Gates Foundation Typhoid Vaccine Acceleration Consortium (TyVAC, INV-030857).

BPH was supported by a NHMRC Leadership Fellowship (GNT1196103). KAT and this work

are supported by the Centers for Disease Control and Prevention. MML was supported by

BMGF (OPP1194582, INV-000049, and INV-029806). MH was supported by the National

Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated

Infections and Antimicrobial Resistance at Oxford University in partnership with the UK Health

Security Agency (UKHSA) (NIHR200915), and supported by the NIHR Biomedical Research

Centre, Oxford. DAR is supported by the NIH (U19 AI110802-01, NIH T32AI162579). JJJ was

supported by grants from Bill and Melinda Gates Foundation, USA (Investment ID INV-009497

OPP1159351) for the Project, National Surveillance System for Enteric Fever in India. RRW is

supported by an ARC Discovery Early Career Researcher Award (DE250100677). JH was

supported by an Emerging Leadership Fellowship from the NHMRC of Australia (GNT2034741).

This research was funded in part by the Wellcome Trust. For the purpose of open access, the

author has applied a CC BY public copyright licence to any Author Accepted Manuscript version

arising from this submission

Authors' Contributions

DJI, JH, ZAD and KEH – Conceptualization.

DJI, JAK, ZAD and KEH - Data curation.

JH, MH, ZI, ZAD and KEH – Software

DJI, JH, RRW, ZAD and KEH - Methodology

DJI, JK, AOA, NA, SA, PMA, IIB, MEC, MAC, JAC, PDG, HI, JJJ, AK, KHK, MM, SN, SO, INO,

SIAR, AR, DR, VS, MJS, AMS, GTS, KAT, RRW, ZAD and KEH - Formal analysis

DJI, JH, AOA, NA, SA, PMA, IIB, MC, MAC, JAC, PDG, BPH, HI, JJJ, LMJ, AK, KHK, JK, MML,

MM, SN, SO, INO, PEO, SIAR, AR, DAR, VS, MJS, AMS, GTS, KAT, RRW, ZAD and KEH -

Investigation,

DJI and KEH - Visualization,

DJI, ZAD and KEH - Writing – original draft,

DJI, JH, ZI, MH, JAK, AOA, NA, SA, PMA, IIB, MEC, MAC, JC, PDG, BPH, HI, JJJ, LMJ, AK,

KHK, JK, ML, MM, SN, SO, INO, PEO, SIAR, AR, DR, VS, MJS, AMS, GTS, KT, RRW, ZAD

and KEH - Writing – review and editing

All authors read and approved the final manuscript.

Acknowledgements

MAC and SN are affiliated to the National Institute for Health Research Health Protection

Research Unit (NIHR HPRU) in Genomics and Enabling Data at University of Warwick in

partnership with the UK Health Security Agency (UKHSA). MAC and SN are based at UKHSA.

MH is affiliated to the NIHR Health Protection Research Unit in Healthcare Associated Infections

and Antimicrobial Resistance at Oxford University. The views expressed are those of the

author(s) and not necessarily those of the NIHR, the Department of Health and Social Care or

the UK Health Security Agency.

SIAR from the icddr,b is grateful to the Governments of Bangladesh, and Canada for providing

unrestricted support.

NA is grateful for support from ICMR India (IIRP-2023-2500).

KAT thanks the epidemiology and laboratory partners in state and local health departments. The

findings and conclusions in this report are those of the authors and do not necessarily represent

the official position of the Centers for Disease Control and Prevention. Names of specific

vendors, manufacturers, or products are included for public health and informational purposes;

inclusion does not imply endorsement of the vendors, manufacturers, or products by the

Centers for Disease Control and Prevention or the US Department of Health and Human

Services.

AMS thanks all participants of the GERMS-SA Laboratory Surveillance Network, South Africa,

for submission of clinical isolates of Salmonella species to the National Institute for

Communicable Diseases; the surveillance network includes laboratories belonging to the

Department of Health (NHLS laboratories) and laboratories that form part of the private sector.

We thank all the lab members involved in SEFI reference lab activities at CMC Vellore

implicated in culture maintenance and ONT data generation.

This work was supported by Monash eResearch capabilities, including the high performance

computer M3 and Research Data Storage.

Global Typhoid Genomics Consortium Group Authorship

Ali H. Abbas[1], Safina Abdul Razzak[2], Mabel K. Aworh[3], Stephen Baker[4], Buddha Basnyat[5], Ka

Lip Chew[6], Thomas C. Darton[7], Andrew R. Greenhill[8], Madhu Gupta[9], Mochammad Hatta[10],

Aamer Ikram[11], Dasaratha Ramaiah Jinka[12], A. C. Lauer[13], Octavie Lunguya[14], Jaspreet

Mahindroo[15], Tapfumanei Mashe[16], Elli Mylona[17], Maria Pardos de la Gandara[18], Andrew J.

Pollard[19], Saikt Rahman[20], Elrashdy M Redwan[21,22,23], Jean Pierre Rutanga[24], Jivan Shakya[25],

Arif Tanmoy[26], Mathew S. Thomas[27], Sandra Van Puyvelde[28], Jacqueline M Wright[29]

1 Department of Microbiology, Faculty of Veterinary Medicine, University of Kufa, Kufa- Najaf

Street, Najaf, Iraq

2 Department of Paediatrics and Child Health, Aga Khan University, P.O. Box 3500, National

Stadium Rd, Karachi, Sindh, Pakistan.

3 North Carolina State University, 1060 William Moore Drive, Raleigh, NC 27607, USA

4 University of Cambridge, JCBC, Puddicome way, Cambridge, CB2 0AW, United Kingdom

5 OUCRU-Nepal, Jhamshikhel, Lalitpur, Nepal

6 National University Hospital, 5 Lower Kent Ridge Road, Singapore 119074, Republic of

Singapore

7 Clinical Infection Research Group, School of Medicine and Population Health, University of

Sheffield, Western Bank, Sheffield, S10 2TN, United Kingdom

8 Federation University Australia, Gippsland Campus, Churchill, Victoria, Australia

9 Postgraduate Institute of Medical Education and Research, sector 12, Chandigarh. CMC

Vellore, India

10 Department of Molecular Biology and Immunology, Faculty of Medicine, Hasanuddin

University, Makassar, 90245, Indonesia

11 Fauji Foundation, Rawal Road, Rawalpindi, Pakistan

12 RDT Hospital, Bathalapalli, Srisathya Sai District, 515661, India

13 Centers for Disease Control and Prevention, Atlanta, United States of America

14 Department of Microbiology, Institut National de Recherche Biomédicale, Kinshasa,

Democratic Republic of the Congo

15 MRC Centre for Global Infectious Disease Analysis, School of Public Health, Imperial

College London, London, W12 0BZ, United Kingdom

16 World Health Organisation, Zimbabwe

17 Cambridge Institute of Therapeutic Immunology and Infectious Disease (CITIID), University

of Cambridge, Cambridge CB1 0AW, United Kingdom

18 Institut Pasteur, France

19 Oxford Vaccine Group, Department of Paediatrics, University of Oxford and the NIHR Oxford

Biomedical Research Centre, Oxford, UK

20 RDT Hospital, Bathalapalli, Srisathya Sai District, 515661, India

21 Department of Biological Science, Faculty of Science, King Abdulaziz University, Jeddah,

Saudi Arabia

22 Centre of Excellence in Bionanoscience Research, King Abdulaziz University, Jeddah

21589, Saudi Arabia

23 Therapeutic and Protective Proteins Laboratory, Protein Research Department, Genetic

Engineering and Biotechnology Research Institute, City of Scientific Research and

Technological Applications, New Borg EL-Arab 21934, Alexandria, Egypt.

24 University of Rwanda, KK 737 street, Kigali, B.P 3900, Rwanda

25 Institute for Research in Science and Technology, Sachal-2, Lalitpur, Nepal

26 Child Health Research Foundation, Dhaka, Bangladesh

27 CMAI, ICMDA

28 University of Antwerp, Antwerp, Belgium

29 Institute of Environmental Sciences and Research 27 Creyke Road, Christchurch, 8041, New
Zealand

References

1. Meiring JE, Khanam F, Basnyat B, Charles RC, Crump JA, Debellut F, et al. Typhoid fever.
Nat Rev Dis Prim. 2023;9: 71. doi:10.1038/s41572-023-00480-z

2. Collaborators G 2017 T and P, Stanaway JD, Reiner RC, Blacker BF, Goldberg EM, Khalil IA,
et al. The global burden of typhoid and paratyphoid fevers: a systematic analysis for the Global
Burden of Disease Study 2017. Lancet Infect Dis. 2019;19: 369–381. doi:10.1016/s1473-
3099(18)30685-6

3. The WHO AWaRe (Access, Watch, Reserve) antibiotic book. Geneva: World Health
Organization; 2022. Licence: CC BY-NC-SA 3.0 IGO.

4. Nabarro LE, McCann N, Herdman MT, Dugan C, Ladhani S, Patel D, et al. British infection
association guidelines for the diagnosis and management of enteric fever in England. J Infect.
2022;84: 469–489. doi:10.1016/j.jinf.2022.01.014

5. Collaborators G 2021 F, Vollset SE, Ababneh HS, Abate YH, Abbafati C, Abbasgholizadeh R,
et al. Burden of disease scenarios for 204 countries and territories, 2022–2050: a forecasting
analysis for the Global Burden of Disease Study 2021. Lancet. 2024;403: 2204–2256.
doi:10.1016/s0140-6736(24)00685-8

6. da Silva KE, Tanmoy AM, Pragasam AK, Iqbal J, Sajib MSI, Mutreja A, et al. The
international and intercontinental spread and expansion of antimicrobial-resistant *Salmonella*
Typhi: a genomic epidemiology study. Lancet Microbe. 2022;3: e567–e577. doi:10.1016/s2666-
5247(22)00093-3

7. Hughes MJ, Birhane MG, Dorough L, Reynolds JL, Caidi H, Tagg KA, et al. Extensively Drug-
Resistant Typhoid Fever in the United States. Open Forum Infect Dis. 2021;8: ofab572.
doi:10.1093/ofid/ofab572

8. Walker J, Chaguza C, Grubaugh ND, Carey M, Baker S, Khan K, et al. Assessing the global
risk of typhoid outbreaks caused by extensively drug resistant *Salmonella* Typhi. Nat Commun.
2023;14: 6502. doi:10.1038/s41467-023-42353-9

9. Posen HJ, Wong W, Farrar DS, Campigotto A, Chan T, Barker KR, et al. Travel-associated
extensively drug-resistant typhoid fever: a case series to inform management in non-endemic
regions. J Travel Med. 2022;30: taac086. doi:10.1093/jtm/taac086

10. Wang Y, Lu D, Jin Y, Wang H, Lyu B, Zhang X, et al. Extensively Drug-Resistant (XDR)
Salmonella Typhi Outbreak by Waterborne Infection — Beijing Municipality, China, January–
February 2022. China CDC Wkly. 2022;4: 254–258. doi:10.46234/ccdcw2022.062

11. Nizamuddin S, Khan EA, Chattaway MA, Godbole G. Early Release - Case of Carbapenem-
Resistant *Salmonella* Typhi Infection, Pakistan, 2022 - Volume 29, Number 11—November
2023 - Emerging Infectious Diseases journal - CDC. Emerg Infect Dis. 2023;29: 2395–2397.
doi:10.3201/eid2911.230499

12. Dyson ZA, Klemm EJ, Palmer S, Dougan G. Antibiotic Resistance and Typhoid. Clin Infect
Dis. 2019;68: S165–S170. doi:10.1093/cid/ciy1111

13. Argimón S, Yeats CA, Goater RJ, Abudahab K, Taylor B, Underwood A, et al. A global
resource for genomic predictions of antimicrobial resistance and surveillance of *Salmonella*
Typhi at pathogenwatch. Nat Commun. 2021;12: 2879. doi:10.1038/s41467-021-23091-2

14. Carey ME, Dyson ZA, Ingle DJ, Amir A, Aworh MK, Chattaway MA, et al. Global diversity
and antimicrobial resistance of typhoid fever pathogens: Insights from a meta-analysis of 13,000
Salmonella Typhi genomes. eLife. 2023;12: e85867. doi:10.7554/elife.85867

15. Wong VK, Baker S, Pickard DJ, Parkhill J, Page AJ, Feasey NA, et al. Phylogeographical
analysis of the dominant multidrug-resistant H58 clade of *Salmonella* Typhi identifies inter- and
intracontinental transmission events. Nat Genet. 2015;47: 632–639. doi:10.1038/ng.3281

16. Hooda Y, Sajib MSI, Rahman H, Luby SP, Bondy-Denomy J, Santosham M, et al. Molecular
mechanism of azithromycin resistance among typhoidal *Salmonella* strains in Bangladesh
identified through passive pediatric surveillance. Plos Neglect Trop D. 2019;13: e0007868.
doi:10.1371/journal.pntd.0007868

17. Day MR, Doumith M, Nascimento VD, Nair S, Ashton PM, Jenkins C, et al. Comparison of
phenotypic and WGS-derived antimicrobial resistance profiles of *Salmonella enterica* serovars
Typhi and Paratyphi. J Antimicrob Chemoth. 2017;73: 365–372. doi:10.1093/jac/dkx379

18. Wong VK, Baker S, Connor TR, Pickard D, Page AJ, Dave J, et al. An extended genotyping
framework for *Salmonella enterica* serovar Typhi, the cause of human typhoid. Nat Commun.
2016;7: 12827. doi:10.1038/ncomms12827

19. Ingle DJ, Andersson P, Valcanis M, Wilmot M, Easton M, Lane C, et al. Genomic
Epidemiology and Antimicrobial Resistance Mechanisms of Imported Typhoid in Australia.
Antimicrob Agents Ch. 2021;65: e01200-21. doi:10.1128/aac.01200-21

20. Ikhimiukor OO, Oaikhena AO, Afolayan AO, Fadeyi A, Kehinde A, Ogunleye VO, et al.
Genomic characterization of invasive typhoidal and non-typhoidal *Salmonella* in southwestern
Nigeria. PLoS Neglected Trop Dis. 2022;16: e0010716. doi:10.1371/journal.pntd.0010716

21. Sikorski MJ, Hazen TH, Desai SN, Nimarota-Brown S, Tupua S, Sialeipata M, et al.
Persistence of Rare *Salmonella* Typhi Genotypes Susceptible to First-Line Antibiotics in the
Remote Islands of Samoa. Mbio. 2022;13: e01920-22. doi:10.1128/mbio.01920-22

22. Dyson ZA, Holt KE. Five years of GenoTyphi: updates to the global *Salmonella* Typhi
genotyping framework. J Infect Dis. 2021;224: jiab414. doi:10.1093/infdis/jiab414

23. Britto CD, Dyson ZA, Duchene S, Carter MJ, Gurung M, Kelly DF, et al. Laboratory and
molecular surveillance of paediatric typhoidal *Salmonella* in Nepal: Antimicrobial resistance and
implications for vaccine policy. Plos Neglect Trop D. 2018;12: e0006408.
doi:10.1371/journal.pntd.0006408

24. Rahman SIA, Dyson ZA, Klemm EJ, Khanam F, Holt KE, Chowdhury EK, et al. Population
structure and antimicrobial resistance patterns of Salmonella Typhi isolates in urban Dhaka,
Bangladesh from 2004 to 2016. Plos Neglect Trop D. 2020;14: e0008036.
doi:10.1371/journal.pntd.0008036

25. Tanmoy AM, Westeel E, Bruyne KD, Goris J, Rajoharison A, Sajib MSI, et al. *Salmonella*
enterica Serovar Typhi in Bangladesh: Exploration of Genomic Diversity and Antimicrobial
Resistance. Mbio. 2018;9: e02112-18. doi:10.1128/mbio.02112-18

26. Klemm EJ, Shakoor S, Page AJ, Qamar FN, Judge K, Saeed DK, et al. Emergence of an
Extensively Drug-Resistant *Salmonella enterica* Serovar Typhi Clone Harboring a Promiscuous
Plasmid Encoding Resistance to Fluoroquinolones and Third-Generation Cephalosporins. Mbio.
2018;9: e00105-18. doi:10.1128/mbio.00105-18

27. Kariuki S, Dyson ZA, Mbae C, Ngetich R, Kavai SM, Wairimu C, et al. Multiple introductions
of multidrug-resistant typhoid associated with acute infection and asymptomatic carriage,
Kenya. eLife. 2021;10: e67852. doi:10.7554/elife.67852

28. Ingle DJ, Hawkey J, Hunt M, Dyson ZA, Holt KE. Technical Report: GenoTyphi
Implementation in Mykrobe. Zenodo 2022. Available from: https://zenodo.org/records/7407985

29. Parkhill J, Dougan G, James KD, Thomson NR, Pickard D, Wain J, et al. Complete genome
sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. Nature.
2001;413: 848–852. doi:10.1038/35101607

30. Labbé G, Kruczkiewicz P, Robertson J, Mabon P, Schonfeld J, Kein D, et al. Rapid and
accurate SNP genotyping of clonal bacterial pathogens with BioHansel. Microb Genom. 2021;7:
000651. doi:10.1099/mgen.0.000651

31. Hunt M, Bradley P, Lapierre SG, Heys S, Thomsit M, Hall MB, et al. Antibiotic resistance
prediction for *Mycobacterium tuberculosis* from genome sequence data with Mykrobe.
Wellcome Open Res. 2019;4: 191. doi:10.12688/wellcomeopenres.15603.1

32. Bradley P, Gordon NC, Walker TM, Dunn L, Heys S, Huang B, et al. Rapid antibiotic-
resistance predictions from genome sequence data for *Staphylococcus aureus* and
Mycobacterium tuberculosis. Nat Commun. 2015;6: 10063. doi:10.1038/ncomms10063

33. Walker TM, Miotto P, Köser CU, Fowler PW, Knaggs J, Iqbal Z, et al. The 2021 WHO
catalogue of *Mycobacterium tuberculosis* complex mutations associated with drug resistance: a
genotypic analysis. Lancet Microbe. 2022;3: e265–e273. doi:10.1016/s2666-5247(21)00301-3

34. Hall MB, Coin LJM. Assessment of the 2021 WHO *Mycobacterium tuberculosis* drug
resistance mutation catalogue on an independent dataset. Lancet Microbe. 2022;3: e645.
doi:10.1016/s2666-5247(22)00151-3

35. Baker S, Holt K, Whitehead S, Goodhead I, Perkins T, Stocker B, et al. A linear plasmid
truncation induces unidirectional flagellar phase change in H:z66 positive *Salmonella* Typhi. Mol
Microbiol. 2007;66: 1207–1218. doi:10.1111/j.1365-2958.2007.05995.x

36. Holt KE, Phan MD, Baker S, Duy PT, Nga TVT, Nair S, et al. Emergence of a Globally
Dominant IncHI1 Plasmid Type Associated with Multiple Drug Resistant Typhoid. Plos Neglect
Trop D. 2011;5: e1245. doi:10.1371/journal.pntd.0001245

37. Hawkey J, Paranagama K, Baker KS, Bengtsson RJ, Weill F-X, Thomson NR, et al. Global
population structure and genotyping framework for genomic surveillance of the major dysentery
pathogen, *Shigella sonnei*. Nat Commun. 2021;12: 2684. doi:10.1038/s41467-021-22700-4

38. Hawkey J, Frézal L, Dien AT, Zhukova A, Brown D, Chattaway MA, et al. Genomic
perspective on the bacillus causing paratyphoid B fever. Nat Commun. 2024;15: 10143.
doi:10.1038/s41467-024-54418-4

39. Thanh DP, Karkey A, Dongol S, Thi NH, Thompson CN, Rabaa MA, et al. A novel
ciprofloxacin-resistant subclade of H58 *Salmonella* Typhi is associated with fluoroquinolone
treatment failure. Elife. 2016;5: e14003. doi:10.7554/elife.14003

40. Crump JA, Kretsinger K, Gay K, Hoekstra RM, Vugia DJ, Hurd S, et al. Clinical Response
and Outcome of Infection with *Salmonella enterica* Serotype Typhi with Decreased
Susceptibility to Fluoroquinolones: a United States FoodNet Multicenter Retrospective Cohort
Study. Antimicrob Agents Chemother. 2008;52: 1278–1284. doi:10.1128/aac.01509-07

41. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New
Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J Comput Biol.
2012;19: 455–477. doi:10.1089/cmb.2012.0021

42. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo
Assembler. Curr Protoc Bioinform. 2020;70: e102. doi:10.1002/cpbi.102

43. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo
genome assemblies. Bioinformatics. 2015;31: 3350–3352. doi:10.1093/bioinformatics/btv383

44. Wickham H. ggplot2. Wiley Interdiscip Rev: Comput Stat. 2011;3: 180–185.
doi:10.1002/wics.147

45. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the
Tidyverse. J Open Source Softw. 2019;4: 1686. doi:10.21105/joss.01686

46. Chattaway MA, Gentle A, Nair S, Tingley L, Day M, Mohamed I, et al. Phylogenomics and
antimicrobial resistance of Salmonella Typhi and Paratyphi A, B and C in England, 2016–2019.
Microb Genom. 2021;7. doi:10.1099/mgen.0.000633

47. CDC. National Antimicrobial Resistance Monitoring System (NARMS) Now: Human Data.
Atlanta, Georgia: U.S. Department of Health and Human Services, CDC [Internet]. Available
from: https://www.cdc.gov/narmsnow

48. Reller LB, Weinstein M, Jorgensen JH, Ferraro MJ. Antimicrobial Susceptibility Testing: A
Review of General Principles and Contemporary Practices. Clin Infect Dis. 2009;49: 1749–1755.
doi:10.1086/647952

49. U.S Food and Drug Administration. Antimicrobial Susceptibility Test (AST) Systems - Class
II Special Controls Guidance for Industry and FDA. [Internet]. Available
https://www.fda.gov/media/88069/download.

50. Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA, et al. CARD 2023:
expanded curation, support for machine learning, and resistome prediction at the
Comprehensive Antibiotic Resistance Database. Nucleic Acids Res. 2022;51: D690–D699.
doi:10.1093/nar/gkac920

51. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol.
2019;20: 257. doi:10.1186/s13059-019-1891-0

52. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34:
3094–3100. doi:10.1093/bioinformatics/bty191

53. Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics. 2021;37:
4572–4574. doi:10.1093/bioinformatics/btab705

54. Achtman M, Wain J, Weill F-X, Nair S, Zhou Z, Sangal V, et al. Multilocus Sequence Typing
as a Replacement for Serotyping in *Salmonella enterica*. Plos Pathog. 2012;8: e1002776.
doi:10.1371/journal.ppat.1002776

55. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using
repeat graphs. Nat Biotechnol. 2019;37: 540–546. doi:10.1038/s41587-019-0072-8

56. Nair S, Barker CR, Bird M, Greig DR, Collins C, Painset A, et al. Presence of phage-
plasmids in multiple serovars of *Salmonella enterica*. Microb Genom. 2024;10: 001247.
doi:10.1099/mgen.0.001247

57. Greig DR, Bird MT, Chattaway MA, Langridge GC, Waters EV, Ribeca P, et al.
Characterization of a P1-bacteriophage-like plasmid (phage-plasmid) harbouring bla CTX-M-15
in *Salmonella enterica* serovar Typhi. Microb Genom. 2022;8: mgen000913.
doi:10.1099/mgen.0.000913

58. Goldblatt J, Ward A, Yusuf M, Day M, Godbole G, Morris-Jones S. Azithromycin
susceptibility testing for *Salmonella enterica* isolates: discordances in results using MIC gradient
strips. J Antimicrob Chemother. 2020;75: 1820–1823. doi:10.1093/jac/dkaa097

59. Jabeen K, Saleem S, Nizamuddin S, Arshad F, Jahan S, Hasnain F, et al. Reporting of
Azithromycin Activity against Clinical Isolates of Extensively Drug-Resistant Salmonella enterica
Serovar Typhi. Am J Trop Med Hyg. 2023;108: 942–947. doi:10.4269/ajtmh.22-0557

60. Ellington MJ, Ekelund O, Aarestrup FM, Canton R, Doumith M, Giske C, et al. The role of
whole genome sequencing in antimicrobial susceptibility testing of bacteria: report from the
EUCAST Subcommittee. Clin Microbiol Infect. 2017;23: 2–22. doi:10.1016/j.cmi.2016.11.012

61. Wick R. Yet another ONT accuracy test: Dorado v0.5.0. Available from:
doi.org/10.5281/zenodo.10397817

Tables and Figures
1045

1046 **Table 1. Validation of genotype calls**
1047 Summary of Typhi Mykrobe genotype calls (GenoTyphi scheme) for 12,839 Illumina readsets,
1048 compared with mapping-based genotype calls.

Typhi Mykrobe vs mapping-based genotype calls	Genomes (N)
invA-negative, thus designated non-Typhi and not genotyped by Mykrobe	5
Single genotype, exact match	12,807
Two genotypes, exact match (reported differently)	17
Genotype 3.5.3, mismatch (reported incorrectly by mapping pipeline as 3.5.4 due to breaking hierarchy assumption)	8
Discordant result (low support reported by mapping-based pipeline, phylogenetic tree supports Mykrobe call)	2

1049

Table 2. Validation of AMR genotyping (vs assembly based)

Comparison of AMR genotyping by Typhi Mykrobe (*k*-mer-based analysis of reads) vs Pathogenwatch (based on nucleotide blast search of assemblies), for n=11,992 isolates. Complete details of the call comparisons are given in **Supplementary Table 3**. Categorical agreement at drug level is calculated based on agreement of the binary S/R categorisation of the 11,992 isolates by each method (i.e. 'Concordant' means Typhi Mykrobe and Typhi Pathogenwatch made the same call of S or R based on absence or presence, respectively, of known genetic determinants; 'Discordant' means one method identified determinants and called R, while the other did not and called S). *For ciprofloxacin, both tools categorise as S/I/R as defined in Methods, and categorical agreement was calculated across these three categories. **Results for *dfrA* genes are summarised together, and as some genomes have multiple *dfrA* alleles the total observations is greater than the number of isolates (see details in **Supplementary Table 3**).

Drug	Agreement (drug level)	Genetic marker	Concordant	Discordant	Agreement (marker level)
Ampicillin	99.72%	*bla*TEM-1	11,958	34	99.72%
		*bla*OXA-7	11,991	1	99.99%
Azithromycin	99.98%	AcrB-717	11,991	1	99.99%
		mphA	11,992	0	100.00%
Ceftriaxone	99.97%	*bla*CTX-M	11,990	3	99.97%
		*bla*SHV-12	11,991	1	99.99%
		*bla*OXA-134	11,992	0	100.00%
		ampC	11,992	0	100.00%
Chloramphenicol	99.89%	*catA1*	11,979	13	99.89%
		cmlA1	11,992	0	100.00%
Ciprofloxacin*	99.72%	*qnrB*	11,979	13	99.89%
		qnrD	11,992	0	100.00%
		qnrS	11,987	5	99.96%
		GyrA-83	11,986	6	99.95%
		GyrA-87	11,986	10	99.92%
		GyrB-464	11,493	8	99.93%
		ParC-80	11,976	17	99.86%
		ParC-84	11,992	0	100.00%
Sulfonamides	99.81%	*sul1*	11,971	21	99.82%
		sul2	11,819	173	98.56%
Trimethoprim	99.83%	*dfrA* genes**	12,049	22	99.82%
Tetracycline	99.85%	*tetA(A)*	11,986	6	99.95%
		tetA(B)	11,980	12	99.90%
		tetA(C)	11,970	0	100.00%
		tetA(D)	11,970	0	100.00%
-	-	**Total**	**299,004**	**346**	**99.88%**

1064 **Table 3 - Validation of plasmid replicon detection (vs assembly based).**
1065 Comparison of plasmid replicon markers identified in the genomes by Typhi Mykrobe (*k*-mer-
1066 based analysis of reads) vs Pathogenwatch (PW, based on nucleotide blast search of
1067 assemblies), for n=11,992 isolates.

Plasmid replicon marker	Both	Mykrobe only	PW only	Agreement
IncFIAHI1	251	0	0	100%
IncFIB(K)	175	0	1	99.99%
IncFIB(pHCM2)	1406	26	5	99.74%
IncHI1A	872	9	3	99.90%
IncHI1B	874	7	1	99.93%
IncHI2A	3	0	0	100.00%
IncI1	5	0	0	100.00%
IncL/M	1	0	0	100.00%
IncN	100	2	0	99.98%
IncX3	21	0	0	100.00%
IncY	666	0	0	100.00%

1068
1069

A. Typhi Mykrobe pipeline

B. AMR determinants included in Typhi Mykrobe

Drug	Acquired genes	Point mutations
ampicillin	*bla*TEM-1, *bla*OXA-7	
azithromycin	*ereA, ermB, mphA*	AcrB-717
carbapenems	*bla*KPC-2, *bla*NDM-5, *bla*VIM-1, *bla*IMP-27, *bla*OXA-48	
ceftriaxone	*ampC1, bla*CTX-M-15, *bla*OXA-134, *bla*SHV-12	
chloramphenicol	*catA1, cmlA1*	
ciprofloxacin	*qnrB1, qnrD1, qnrS1*	GyrA-83, GyrA-87, GyrB-464F, ParC-80, ParC-84
sulfonamides	*sul1, sul2*	
trimethoprim	*dfrA1, dfrA5, dfrA7, dfrA14, dfrA15, dfrA17, dfrA18*	
tetracycline	*tetA(A), tetA(B), tetA(C), tetA(D)*	

C. Plasmids included in Typhi Mykrobe

Group	Plasmid probe target
AMR-linked replicons	IncFIBK, IncHI2A, IncI1, IncL/M, IncN, IncX1, IncX3, IncY
IncHI1 targets	IncHI1A, IncHI1BR27, IncFIAHI1 (rep markers); C19241A (marker of PST6 lineage)
Non-AMR plasmids	IncFIB_pHCM2 (cryptic), pBSSB1 (z66 flagellin), pO111

1070
1071
1072 **Figure 1: Typhi Mykrobe functionality**

1073 A) A schematic overview of the Typhi Mykrobe pipeline. B) Table of epidemiologically important

1074 AMR determinants targeted for detection/typing by Typhi Mykrobe, grouped by drug. C) Table of

1075 epidemiologically important plasmid replicons targeted for detection by Typhi Mykrobe.

1076
1077

Figure 2: Validation of Typhi Mykrobe AMR predictions vs AMR phenotypes

Summary of Typhi Mykrobe's genotype-based clinical resistance categorisations compared with susceptibility phenotypes, for 4,018 isolates with publicly available matched genome and phenotype data. Comparisons are summarised as categorical agreement, and calculated separately for the three different source datasets, which used different phenotyping methods and interpretive standards (US CDC: MIC, CLSI; UKHSA: MIC, EUCAST; SEAP: disk diffusion, EUCAST; see **Methods**). Major errors were defined as susceptible isolates with AMR determinants detected (reported as R by Mykrobe). Very major errors were defined as resistant isolates with no AMR determinants detected (reported as S by Mykrobe). Error bars are shown that represent 95% confidence interval of proportion. Note UKHSA tested amoxicillin rather than ampicillin, the result is reported in the ampicillin row.

1091

Figure 3: Typhi Mykrobe genotypes vs minimum inhibitory concentration from two

reference laboratories, for ciprofloxacin.

a) UKHSA dataset, using EUCAST. **b)** US CDC dataset, using CLSI. Each column represents

the set of isolates in which Typhi Mykrobe identified a unique combination of genetic

determinants, indicated in the panel at the bottom. For each column, the violin plots show the

distribution of ciprofloxacin MIC values, and stacked barplots show the proportion of genomes

called as S, I, or R (coloured as per inset legend, and labelled with counts). The solid horizontal

lines on the violin plots mark the R breakpoint, the dashed line marks the CLSI I breakpoint

used for the US CDC dataset.

1101

1102

Figure 4: Typhi Mykrobe genotypes vs disk diffusion data from the SEAP study, for three
drug classes

Data for three drugs are shown in panels **a)** Azithromycin, **b)** Ceftriaxone, and **c)** Ciprofloxacin.
Each column represents the set of isolates in which Typhi Mykrobe identified a unique
combination of genetic determinants, indicated in the panel at the bottom. For each column, the
violin plots show the distribution of disk diffusion values for azithromycin, ceftriaxone or
ciprofloxacin. The stacked barplots show the proportion of genomes called as S, I, or R
(coloured as per inset legend, and labelled with counts). The solid horizontal lines on the violin
plots mark the R EUCAST breakpoint used for the SEAP disk diffusion data. The dashed line
marks the EUCAST breakpoint for I.

1113