Proximity proteomics reveals a co-evolved LRRK2-regulatory network linked

to centrosomes

Marita Eckert[1,4,5], Pasquale Miglionico[2,5], Francesca Izzi[1], Natalia De Oliveira Rosa[2], Marius Ueffing[3],

Francesco Raimondi[2*] and Christian Johannes Gloeckner[1,3*]

[1]German Center for Neurodegenerative Diseases, 72076 Tübingen, Germany;

[2]Laboratorio di Biologia Bio@SNS Scuola Normale Superiore, 56126 Pisa, Italy;

[3]Center for Ophthalmology, Institute for Ophthalmic Research, University of Tübingen, 72076

Tübingen, Germany

[4]Graduate School of Cellular and Molecular Neuroscience, University of Tübingen, 72074 Tübingen,

Germany.

[5]contributed equally to this work.

Corresponding authors:

Christian Johannes Gloeckner (johannes.gloeckner@dzne.de)

Francesco Raimondi (francesco.raimondi@sns.it)

Abstract

The Leucine-rich repeat kinase 2 (LRRK2) not only plays a vital role in familial forms of Parkinson's

disease (PD) but is also considered as a risk factor for idiopathic PD. Its multi-domain architecture

enables a fine-tuned regulation of its biological function by orchestrating intra- and inter-molecular

interactions. Here, we present BioID proximity proteomes of LRRK2 revealing new interactors, which

we further characterized by a novel evolutionary and structural bioinformatics pipeline. Co-

evolutionary analysis of the protein-protein interaction network identified a structural and functional

module enriched in cytoskeletal components linked to the centrosome and microtubules.

Furthermore, structural modelling of binary interactions via AlphaFold-multimer revealed distinct

groups of interactors engaging with LRRK2 dependent on specific conformations and epitopes.

Furthermore, we identified distinct changes in the LRRK2 proximity proteome induced by the type I

kinase inhibitor MLi-2 or by co-expression of the LRRK2 upstream effector RAB29. Dependent on its

state of activity and conformation, these protein-protein interactions link LRRK2 to defined cellular

sub-compartments, including centriolar satellites as well as vesicular sub-compartments.

Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disease after Alzheimer's, it occurs mostly sporadically and is distributed with high frequency in the older generations. PD is characterized by a loss of dopaminergic neurons in the *substantia nigra* leading to the motor symptoms associated with the disease (Fahn, 2003). Several therapies targeting the symptoms, such as dopamine substitution, are practiced but so far, no causative therapy exists. About 5-10% of PD cases are inheritable (Gasser et al., 2011). *LRRK2*, localized within the PARK8 locus and coding for the Leucine-rich repeat kinase 2, is one of the genes associated with familial forms of the disease but also represents a risk factor in idiopathic PD (Paisan-Ruiz et al., 2004; Simon-Sanchez et al., 2009; Zimprich et al., 2004). The multi-domain protein LRRK2 is an active protein kinase and, at least, mutations with confirmed disease segregation, including the most common G2019S variant within its kinase domain, augment its kinase activity (Kalogeropulou et al., 2022). This makes LRRK2 an important drug target (Kluss et al., 2022b). ATP-competitive LRRK2-specific kinase inhibitors for PD-treatment are under development and in clinical trials (Jennings et al., 2022). Nevertheless, structural changes of the lung seen in non-human primates indicate a considerable small therapeutic window and additionally challenge a long-term treatment with these compounds necessary to allow a life-long reduction of augmented LRRK2 pathway activation (Miller et al., 2023). For this reason, the identification of alternative drug targets within the LRRK2-associated pathways would be a potential solution for a better control of pathophysiological alterations associated with pathogenic forms of this protein, minimizing the risk of drug-induced lung pathologies. In addition, a larger group of patients beyond LRRK2 mutation carriers might benefit from targeting more common PD-associated pathways.

Various functional studies have demonstrated a function of LRRK2 in vesicle trafficking, including presynaptic vesicles as well as autophagy and lysosomal pathways ((Piccoli et al., 2011); reviewed in: (Follett and Farrer, 2021)). In agreement with these cellular functions, LRRK2 phosphorylates a subset of Rab proteins, small G-proteins involved in vesicular transport, such as Rab8a and Rab10, which is elevated by PD-associated LRRK2 variants (Steger et al., 2017).

In common with other Roco proteins, LRRK2 features a remarkable complexity (Wauters et al., 2019). In fact, LRRK2 contains multiple scaffolding domains, not only being involved in keeping inactive LRRK2 in an auto-inhibited state, as recently demonstrated by high-resolution structures of LRRK2 (Deniston et al., 2020; Myasnikov et al., 2021), but are also involved in protein-protein interactions making LRRK2 a signaling hub possibly integrating multiple up- and down-stream signals (reviewed in: (Manzoni et al., 2015)). These studies suggest that the Roco protein LRRK2 is a tightly controlled protein kinase serving as hub and signal integrator and is most likely involved in context-specific, spatially controlled cellular responses that appear evolutionarily conserved. Quite some effort has therefore been spent on the investigation of the protein-protein interaction networks of LRRK2, mainly by using affinity-

based methods. Data of various studies were curated in public databases such as IntAct or STRING and were also subject to meta-analysis to dissect functional LRRK2 signaling networks (Gloeckner and Porras, 2020; Manzoni et al., 2015; Zhao et al., 2024; Zhao et al., 2023). However, affinity-based methods face limits, given their bias for rather stable complexes which miss dynamic/ transient interactions. We therefore used the BioID proximity labelling method to detect new interaction partners of LRRK2. By systematically applying the BioID approach with three different BioID tags, we describe the proximity proteome of LRRK2 under basal conditions. In addition, by a newly designed bioinformatics workflow that estimates co-evolution of interacting partners based on their co-presence in sequence genomes, we could demonstrate that the resulting BioID datasets are characterized by higher co-evolution values similarly to confirmed direct interactions. Through this approach, we have identified a cluster of interactors featuring a high degree of co-evolution to LRRK2 which is enriched in cytoskeletal proteins involved in centrosomal and ciliary dynamics. We predicted the 3D complexes mediated by LRRK2 via AF-multimer, which also revealed distinct conformations and interfaces associated to specific processes. Finally, to further stratify the interactome of LRRK2, we looked at experimentally-defined conformational states of LRRK2, e.g. by investigating the context-specific proximity proteome upon co-expression of the LRRK2 upstream effector RAB29 and upon stabilization of an active-like kinase conformation by inhibition of LRRK2 with a type I ATP-competitive inhibitor (Taylor et al., 2020; Wauters et al., 2019). Inhibition of LRRK2 with the type I inhibitor MLi-2 but not with the type II inhibitor GZD-824 lead to an enrichment of a highly specific sub interactome associated with centriolar satellites.

Methods

Cloning and generation of the stable miniTurbo-LRRK2 cell line

All used proximity tags have been ordered as synthetic genes (Eurofins) and subcloned via *Eco*RI and *Xho*I into pcDNA3.0 (Invitrogen) allowing the expression of fusion proteins in mammalian cells under the strong viral CMV promoter. Gateway Destination clones were generated by cloning a Gateway cassette (Invitrogen) into an *Eco*RV site at the 3' end and in frame of the different BioID-tags [BioID (Roux et al., 2012) (referred as BioID1), BioID2 (Kim et al., 2016), miniTurbo (Branon et al., 2018)]. For better and comparable detection of the fusion proteins, all tags contained an N-terminal HA-tag. The LRRK2 expression clones were subsequently generated by LR cloning (Invitrogen) based on ENTRY clones described earlier (Guaitoli et al., 2016). The CDS of human of RAB29 has been ordered as synthetic gene strand (Eurofins) with flanking attB sites to allow its direct cloning into pDONR201 by BP cloning and subsequently cloned into an in-house (N)FLAG-HA Gateway destination clone by the LR reaction. The HEK293 cell line (ACC-305) stably expressing miniTurbo-LRRK2 has been generated as previously described (Gloeckner et al., 2007). Briefly, single colonies have been selected and tested

99 after transient transfection and selection via G418 (Sigma). The Flag-HA-CYLD vector was kindly

100 provided by Wade Harper (plasmid #22544, Addgene) (Sowa et al., 2009), the pCMV poly HA-ubiquitin

101 construct was kindly provided by D. Bohmann (Treier et al., 1994) and the pRK5-HA-Ubiquitin-K63

102 vector was kindly provided by T. Dawson (plasmind #17606, Addgene) (Lim et al., 2005).

Cell culture and proximity labelling

HEK239T (ACC-635) or HEK293 stably expressing miniTurbo-LRRK2 were grown in 14 cm dishes in full

Dulbecco's modified medium (DMEM) supplemented with penicillin streptomycin and if not

mentioned otherwise with 10% FBS at 37°C in 5% CO2. The cells were grown for 8 hrs to 1 day reaching

a density of 50-80% until transfection with a self-made PEI reagent (Guaitoli et al., 2016). For each

condition, 4-6 plates have been used. If not stated otherwise, for biotinylation, cells have been

incubated with 50μM biotin (Sigma) in serum-free medium for up to 16 hours. In case of the mini turbo

tag, cells have been incubated with 50μM biotin after o/n starvation for 2h. For the determination of

the LRRK2-inhibitor dependent proximity proteomes, a HEK293 line stably expressing mT-LRRK2 was

either incubated with 1μM MLi-2, 1μM GZD-824 or DMSO (vehicle) 1h prior and during 2hrs biotin

incubation. For the analysis of RAB29-dependent proximity proteomes, mT-LRRK2 was either co-

expressed with HA-RAB29 or an empty pcDNA3.0 vector.

Affinity purification of biotinylated proteins

For cell lysis, cell culture dishes were transferred to ice and culture media was removed. The cells were

rinsed two times with 5ml ice-cold PBS. The cells were covered with 1ml lysis buffer [0.5% Nonidet P-

40 Substitute (Merck), protease inhibitor cocktail (Roche), and 0.1% deoxycholic acid (Sigma-Aldrich)

in TBS (30mM Tris-HCl, pH7.4 and 150mM NaCl)] per 14cm dish and detached with a cell scraper. The

cell suspension was collected and incubated for 20-40min on an end-over-end shaker at 4°C. Lysate

were cleared by centrifugation (15,000xg for 10min at 4°C). The protein concentration of cleared

lysates was determined photometrically using a nanodrop (Thermo-Fisher Scientific) or with the BCA

assay (Thermo-Fisher Scientific) following the manufacturer's instructions. Biotinylated proteins were

purified by affinity enrichment using Strep-Tactin Superflow resin (IBA). To this end, the combined

lysates for each condition were incubated with an adjusted resin volume of 25μl packed beads per 14

cm cell culture dish. The resin was equilibrated by washing 3x with 500μl lysis buffer. The lysates were

incubated with the resin overnight at 4°C under constant agitation. The resin was sedimented by

centrifugation at 5,000xg for 3 min at 4°C, transferred to micoSpin columns (Cytiva) and subsequently

washed 3x in 500μl lysis buffer followed by two washes in 500μl 1x TBS to remove traces of SDS.

Biotinylated proteins were eluted by incubation with 400μl desthiobiotin elution buffer (2.5M, IBA).

The desthiobiotin eluates were precipitated by chloroform/ methanol and directly subjected to tryptic

proteolysis following standard protocols (Gloeckner et al., 2009) and stored at -80°C.

CYLD co-expression experiment

For the analysis of CYLD co-expression, cells were co-transfected with LRRK2 Y1699C, CYLD, and

ubiquitin (either in WT form or K63) plasmids. As negative control, CYLD plasmid was replaced by

pCDNA3 empty vector, to maintain the plasmid concentration constant among the different

conditions. For 6-wells, a total of 500ng of plasmid DNA was used for each well; for 14 cm plates, a

total of 8µg of plasmid DNA was used for each plate. The plasmids ratios were LRRK2:CYLD:Ubiquitin

1:3:1. Transfection and cell lysis were performed as described above.

SDS PAGE and Western Blots

For functional evaluation of the bait proteins and to determine the biotinylation efficiency, samples

were subjected to SDS PAGE and analyzed by Western-blotting on PVDF membranes. To assess blotting

efficiency and equal loading, the membranes were stained with Ponceau S. Prior to blocking of the

membranes in 5% nonfat dry milk powder (BioRad) or 5% BSA (Carl Roth) in TBS with 0.1% Tween 20

(TBST) for 1-3h at RT, membranes were de-stained in TBST. Membranes were afterwards incubated

with primary antibody solutions overnight at 4°C under mild agitation. The primary antibody solution

was prepared for rat α-LRRK2 mAb (24D8 clone, hybridoma supernatant, generated in house (Carrion

et al., 2017)) 1:1000in 5% milk in TBST), rabbit α-pS935-LRRK2 mAb (Abcam, ab133450; 1:100,000 in

5% BSA in TBST), rabbit α-RAB10 mAb (Abcam, ab181367; 1:5,000 in 5% milk in TBST), rabbit α-pT73-

RAB10 mAb (Abcam, ab230261; 1:1,000 in 5% BSA in TBST), and mouse α-HA-tag (Roche, 3F10 clone,

Rat 1gG1; 1:1,000 in 5% milk in TBST). Membranes were washed 5x 5min in TBST prior to incubation

with the HRP-conjugated secondary antibodies α-rat (Jackson Immuno-Research, 112-036-062;

1:7,500) and α-rabbit (Jackson Immuno-Research, 111-036-045; 1:7,500), respectively. For the

detection of biotinylated proteins, membranes were incubated with Pierce™ High Sensitivity

Streptavidin-HRP (Thermo Scientific, 1:20,000 in 1% BSA in TBST) overnight at 4°C and washed the next

day 5x 5 min in TBST. Blots were developed with Pierce ECL Plus Western Blotting Substrate following

the manufacturer's protocol (Thermo Scientific) and exposed to Amersham Hyperfilm™ ECL (GE

Healthcare).

Mass spectrometry

160 In solution proteolysis was performed as previously described (Gloeckner et al., 2009). Briefly,

161 chloroform/ methanol precipitates were dissolved in 50mM ammonium bicarbonate pH8.0. Cysteines

162 were reduced and alkylated by treatment with 100mM DTT and 300mM Iodoacetamide, respectively

163 following the incubation with trypsin (Sigma-Aldrich) overnight at 37°C. Samples were desalted by

micropipette C-18 solid-phase extraction tips (StageTip, Thermo Scientific) prior to LC-MSMS analysis. Peptides were subsequently analyzed by LC-MSMS using a nanoflow HPLC system (Ultimate 3000 RSLC; Thermo-Fisher Scientific) coupled to an Orbitrap Q-Exactive plus (Thermo-Fisher Scientific) tandem mass spectrometer. Peptides were separated by reversed C-18 chromatography and 120-minute gradients. MS1 spectra were acquired in the Orbitrap at 70K resolution. After selection of the 10 most intense precursor ions from the MS1 scans for HCD fragmentation (Top 10 method), MS2 spectra were acquired at 17.5K. RAW MS files were analyzed with Maxquant (version 2.0.3.1), enabling label-free-quantification (LFQ with no normalization) and "match between runs" (Cox et al., 2014). For the embedded Andromeda search engine, trypsin has been selected as enzyme and carbamylation was set as fixed modification N-terminal acetylation was set as variable modification. The search was performed against the human subset of the Uniprot/Swissprot database (release 2021_04, 20375 entries). The MaxQuant output was further processed by Perseus and SaintExpress.

Perseus post-processing

For post-analysis, the protein groups were further processed by Perseus (v. 1.6.7). LFQ intensities of the individual were assigned to the conditions and potential contaminants, reverse hits as well as IDs only identified by site were filtered out. For further analysis, only IDs with a maximum number of missing values of n-1 in at least one condition were considered. Remaining missing values were replaced by group-wise imputation based on a normal distribution (log2 values) around the detection limit (width=0.3; down shift=1.8). Intensities of the individual experiments were normalized by their medians and significantly enriched proteins (condition vs control) were determined by a T-test combined with an FDR filter (p=0.05, S0=0.1).

SaintExpress post-processing

We followed a previously described protocol (http://proteomics.fi (Liu et al., 2020)) to select the most confident interactors using the statistical framework of SAINTExpress (Teo et al., 2014), as well as a reference knowledgebase for contaminants (i.e. CRAPome (Mellacheruvu et al., 2013)), to filter out most likely false negative interactors. In particular, we generated the "Bait", "Prey" and "Interaction" files to be subjected to the analysis pipeline from the MaxQuant "*proteinGroups.txt*" file by employing the following columns: "Bait" file (IP name=EV_* or LRRK2_*, bait name=LRRK2, indicator for test and negative control= C=EV_* ;T=LRRK2_*); "Interaction" file (IP name = EV_*/LRRK_*, bait name=LRRK2, prey name=interaction partner gene symbol, spectral count or intensity values=LFQ intensity); "Prey" file (prey (protein)name= interaction partner Uniprot accession, prey protein length=Sequence length, prey gene name= gene symbol). The proteins that were not classified through the CRAPome filter, were retained together with those passing the filter.

197 We unified the interactor list from the three BioID experiments. We finally retained a list of 208

198 interactors which passed the two filters. Only 45 proteins in the filtered interactor list are already

199 present in IntAct.

Graphical representation and network annotation

PPI networks were analyzed by Cytoscape version 3.8.2 (Shannon et al., 2003) and were mapped to

available interaction from IntAct using the Cytoscape IntAct app (Ragueneau et al., 2021). Networks

were annotated with STRINGDB terms via the 'Stringify' function of the StringApp (Doncheva et al.,

2019) and functional enrichment retrieved via the same app. We annotated the networks using the

Auto Annotate plug-in (Kucera et al., 2016) using the Glay clustering method and calculated the labels

using the 'Adjacent Words' of the WordCloud algorithm considering significantly enriched GO

Biological processes and Cellular Compartments in which each gene of the network participated. The

Venn diagram shown in figure 1A has been generated with JVenn provided by the ProteoRE server

(Galaxy Version 2021.05.12) (Galaxy, 2024).

Co-evolution based PPI stratification

211 Co-evolution of two proteins A and B was measured as the Jaccard similarity coefficient of the set of

212 genomes containing an orthologous sequence of A and the set of genomes containing an orthologous

213 sequence of B. Orthologous sequences were collected from the Orthologous MAtrix (OMA) database

214 (Altenhoff et al., 2021) using a Python script written in-house using the PyOMADB client (Kaleb et al.,

215 2019) (Kaleb et al., 2019) (December 13, 2023, database version July 2023).

AF-multimer prediction of binary interactions

We used Alphafold-Multimer v2.3.2 (Evans et al., 2021) to generate 3D models of full-length LRRK2

binary interaction complexes for all the interactors shorter than 2000 residues. The databases required

to run AlphaFold-Multimer were downloaded on 12 January 2023 and the predictions were run with

the flag --max_template_date=01-01-1900, to avoid the usage of any available experimental

templates. Among the 5 models generated for each LRRK2 binary complex, only the best one was

considered for further analysis. The score used to evaluate the models was the default one used by

AlphaFold-Multimer (0.2*pTM + 0.8*ipTM).

Interaction fingerprint

AlphaFold produces probabilities for the C_β-C_β distances between residues *i* and *j* to fall into a series of

distance bins. For each pair of residues, we calculated the expected value of such distance distribution,

yielding a matrix of dimension l_{LRRK2} x l_n, corresponding to the sequence lengths of LRRK2 and the nth

interacting partner, respectively Subsequently, we pooled the distogram on the interaction partner's

side and we assigned to each residue of LRRK2 the minimum expected distance to any residue of the

interactor. We refer to this vector of minimum expected distances as the interaction fingerprint of a

given interacting protein.

RMSD calculation

We compared LRRK2 conformations by performing Root Mean Square Deviation (RMSD)-based

clustering. To calculate the RMSD we fitted the structures using the C_α atoms of the LRRK2 chains from

position 983 to the end of the sequence. The N-terminal portion of the protein was excluded because

it's very flexible. Calculations were performed using the Superimposer function of the PDB Biopython

module (Cock et al., 2009) (version 1.83) through customized scripts and pipeline (Matic et al., 2023).

Clustering

The clustering was performed using the Ward method with the *clustermap* function from *seaborn*

python library (version 0.11.2). Results were displayed through matplotlib (https://matplotlib.org/)

and seaborn (https://seaborn.pydata.org/) libraries using customized python scripts. The number of

clusters chosen was one that maximized the mean Silhouette Coefficient of all samples (Rousseeuw,

1987) , as computed by the *silhouette_score* function in *scikit-learn* (Pedregosa et al., 2011)*.*

Results

Definition of cellular LRRK2 proximity proteomes with BioID1, BioID2 and miniTurbo.

The elucidation of comprehensive protein-protein networks represents a powerful tool not only to

identify signaling pathways associated with a protein of interest but also to connect different

pathogenic mutations in a disease network or identify novel disease-associated proteins through

proximity to known proteins (Boldt et al., 2016; Kiel et al., 2011). Finally, new drug targets as well as

disease biomarkers might arise from this approach. For this reason, various studies addressed the

interatomic networks of LRRK2 in a different cellular context or model organism (Islam et al., 2016;

Meixner et al., 2011; Piccoli et al., 2011; Piccoli et al., 2014). However, most of the used technologies

rely on immuno-precipitation or pull-downs, which have a bias for considerable stable PPIs while

transient interactions remain underrepresented. In the last years, proximity-based methods, e.g.

BioID, have been developed allowing the identification of PPIs under physiological conditions and

within cells also covering transient interactions. The BioID approach is based on a variant of the

bacterial biotin ligase BirA, which has been converted to a promiscuous labeling tool by a single point

mutation within its active center allowing the release of the reactive intermediate biotinyl-AMP. To determine the LRRK2 proximity proteome, we have generated different N-terminal fusion proteins. Besides the original BioID (BioID1) tag (Roux et al., 2012), we used the BioID2 (Kim et al., 2016) as well as miniTurbo, a shorter version of the TurboID tag, lacking the BirA DNA binding domain (Branon et al., 2018). For reasons of comparability, all tags have been cloned in the same way, containing an N-terminal HA tag for detection and a 3x(GGGS) linker between the tag and the protein of interest inserted C-terminally of the tag by Gateway-cloning. All constructs were functionally assessed to demonstrate biotinylation activity and to exclude that the tag interferes with the activity of LRRK2. Western blot analysis shown in Supplemental Figure 1 demonstrates that all constructs allowed sufficient biotinylation in cells. As expected, BioID2 and miniTurbo showed a higher labeling efficiency compared to BioID1. We also confirmed that the fusion proteins are still able to phosphorylate the physiological LRRK2 substrates by probing for Rab10 phosphorylated on Threonine-73. As far as their transient expression is concerned, all constructs were able to phosphorylate endogenous Rab10 in HEK293T cells at comparable LRRK2 expression levels (see: Supplemental Figure 1).

After initial functional tests, all three fusion constructs were used to determine their interactomes after transient expression in HEK293T cells. To increase the stringency of the approach, a tag-only control was used as reference. For the relative comparison of the LRRK2 vs. the tag-only control, the expression levels were adjusted via pre-experiments and always confirmed by Western blot analysis of the cell lysates. In contrast to comparing plus/minus biotin conditions, the tag-only control allows to efficiently exclude natively biotinylated proteins as their degree of biotinylation is also increased by the biotin treatment of the cells already by the experimental design. The data of all three proximity proteomes were analyzed by SaintExpress and, alternatively, by labelfree quantification (LFQ) combined with an FDR-controlled T-test as implemented in Perseus (Tyanova et al., 2016). To further filter and analyze the data set, we implemented a bioinformatics pipeline based on co-evolution and structural modeling by Alphafold-Multimer (Figure 1A). With BioID1, BioID2 and miniTurbo, 168, 312 and 241 proteins were identified, respectively, as being significantly enriched in the LRRK2-containing condition over the tag-only control by the LFQ approach (Figure 1B; Supplemental Table 1). SaintExpress (Teo et al., 2014) and subsequent filtering against the Crapome data of commonly found false-positive hits (Mellacheruvu et al., 2013), however, appeared to be the more stringent analysis compared to MS1-based label-free quantification. Based on this analysis, we created a non-redundant set of 208 unique interactors, hereinafter referred to as "BioID set" which was submitted to the bioinformatics post-processing pipeline.

Functional-enrichment analysis of the LRRK2 proximity proteome

We compared the LRRK2 BioID interactome with a reference LRRK2 interactome available from IntAct

(Orchard et al., 2014). We found that only 54 interactors in the filtered BioID set previously reported

in association with LRRK2, while the vast majority (154) were not yet reported for their interaction

with LRRK2 (Figure 2A,B). By integrating the LRRK2 BioID interactome with the network of known

interactions from Intact it is possible to define several modules of proteins interacting with each other

in addition to LRRK2 (Figure 2A). Among the enriched intracellular and cytosolic compartment, we

found terms related to microtubules (e.g. Microtubule Cytoskeleton, FDR=5.83E-15) and Centrosome

(FDR= 1.43E-13), emphasizing the importance of LRRK2 in microtubule organization (Figure 2C) and

representing the largest module of interacting proteins in the network (Figure 2A). In addition to the

enrichment of microtubule or cytoskeleton associated processes (e.g. FDR =6.04E-10), we also found

processes related to cell cycle and mitosis (e.g. "Regulation of mitotic cell cycle", FDR=9.71E-08) (Figure

2D).

Multiple protein domains instances are significantly enriched within the list of identified LRRK2

interaction partners (Figure 2E). In particular, we found that the Armadillo-type fold instance is the

most enriched one (FDR= 9.58E-06), which is found in 21 distinct interactors, followed by other

domains with lower, but still significant, degree of enrichment, such as "Forkhead-associated (FHA)

domain" (Almawi et al., 2017), found in six distinct proteins, the "DIX domain" (Ehebauer and Arias,

2009; Kafka et al., 2014) found in four proteins, the "K Homology domain, type 1" (Grishin, 2001) found

in six proteins and "Ubiquitin-like domain superfamily" (FDR<0.05), the latter being found in 11 distinct

interacting proteins. Interestingly, LRRK2 and its Drosophila orthologue dLRRK have been

demonstrated to functionally interact with the Forkhead box transcription factor FoxO1/dFoxO. In

particular, LRRK2-mediated phosphorylation of FoxO1 enhances its transcriptional activity leading to

neuronal toxicity (Kanao et al., 2010).

Analysis of co-evolution reveals a cluster of proteins involved in centrosome/ microtubular dynamics

We analyzed the extent of co-evolution of LRRK2 with its interactors to prioritize the most conserved

relationships associated to LRRK2's functions. We derived a metric for the co-evolution between two

interacting proteins based on the degree of co-presence, assessed through a Jaccard score, of

interacting gene pairs in sequenced genomes available from a reference orthology database (e.g. OMA,

see Methods). We found that such simple co-evolutionary metric is informative about non-enzymatic,

direct interactions as well as stable complexes at a human interactome-wide level (Miglionico et al.,

2024, in prep.; Meldal et al., 2022; Orchard et al., 2014) (Supplementary Table 2). We found similar

trends when focusing on LRRK2 interactions reported in IntAct, with "non-enzymatic direct" PPIs being

characterized by a significantly higher Jaccard score compared to those categorized as "association" or

"physical interaction" (Figure 3A). Intriguingly, we found that the distribution of co-evolution metrics

calculated for BioID interactors is significantly higher than most IntAct categories (i.e. "association",

"physical", "enzymatic direct") and it is as high as the non-enzymatic direct interactions (Figure 3A).

This result suggests that interactors captured within the BioID dataset are more likely to represent

stable, direct interaction partners of LRRK2 with whom they co-evolved. Likewise, simple co-evolution

scores are predictive for LRRK2 direct interactions, particularly the non-enzymatic direct ones (Figure

3B; ROC AUC=0.715). In the case of the LRRK2 interactome, we found that this co-evolutionary metric

is more predictive of non-enzymatic direct interaction compared to another co-evolution estimate, i.e.

Mutual Information (ROC AUC=0.66; Supplemental Figure 2), which is instead more predictive at a

human proteome scale (Miglionico et al., 2024, in prep.).

We used the co-evolution metric to cluster LRRK2 and its interactors, which revealed groups of proteins

characterized by reciprocal high co-evolution (Figure 3C). The different groups of interactors are also

characterized by specific biological processes, which are overall mutual exclusive to each other,

suggesting that clusters of co-evolved proteins underpin specific functional interactions and biological

process (Supplemental Figure 3). Notably, the cluster containing LRRK2 (i.e. Cluster 2) is characterized

by on average higher co-evolution and several highly co-evolved proteins (Figure 3C). The top 10

proteins of Cluster 2 showing highest co-evolution are summarized in Table 1. By mapping members

of this cluster onto the STRING functional interaction network, using default parameters (Figure 3D,

see Methods), we found functional interactions for most of the members, including LRRK2. Overall,

STRING yielded a highly significant PPI enrichment (P-value=1.0E-16), suggesting that these proteins

are more likely to interact with themselves than with randomly selected proteins. Moreover, analysis

of the network topology revealed two distinct submodules: one entailing LRRK2 and proteins mainly

involved in Wnt signaling (FDR= 0.0013; Figure 3E); the second one with proteins involved in various

processes related to cytoskeleton ("Protein localization to microtubule cytoskeleton" FDR=2.18E-05),

centrosome ("Loss of Nlp from mitotic centrosomes" FDR=0.00086) and Centriole (FDR= 1.11E-05).

Overall, cluster 2 showed particular enrichment of compartments and cellular component terms, such

as "Microtubule Cytoskeleton" (FDR=3.52E-16) and "Centrosome" (FDR=7.84E-13) (Supplemental

Table 3). Interestingly, the single most co-evolved protein to LRRK2 is CYLD, a deubiquitinase that acts

as a negative regulator of dopamine neuron survival in Parkinson's disease (Pirooznia et al., 2022), but

that has not yet been reported in direct interaction with LRRK2. Interestingly, CYLD appears as a critical

bridging element between the two main submodules within Cluster 2 (Figure 3D).

On the other hand, cluster 1 is characterized by distinct processes, such as FAR/SIN/STRIPAK complex

(P-value= 0.0012) in cluster 1. Taken together, these results suggest that our co-evolution-based

approach was able to identify a group of LRRK2 interacting proteins, mostly associated to microtubule

and centrosome functions via the formation of stable, supramolecular complexes with LRRK2.

Likewise, co-evolution-based clustering of LRRK2 interactors from IntAct shows interactors such as
RAB29, RAB12, LRRK1 or SNCA among the most co-evolved partners which clustered with LRRK2
(Supplemental Figure 4), confirming the tight evolutionary constraint on LRRK2's functional regulation.

We validated the functional role of the interaction between LRRK2 and CYLD by assessing whether the
latter can increase the stability of the former. Interestingly, we could demonstrate that the co-
expression of CYLD increased the stability of Y1999C, a pathogenic variant of LRRK2 showing high levels
of Rab phosphorylation but impaired protein stability (Figure 3F). Given that CYLD is a K-63 selective
ubiquitin deubiquitinase, we co-expressed HA-tagged ubiquitin constructs, either containing a tandem
repeat of WT ubiquitin or a ubiquitin construct only allowing lysine 63 linkage as the other common
linkage lysines were preplaced by arginine (Kaleb et al., 2019). In agreement with previous
observations demonstrating that CYLD is a K-63 selective DUB, CYLD-mediated stabilization was also
greater in the presence of the K63-ubiquitin construct. The condition without CYLD co-expression
showed overall higher LRRK2 levels compared to the HA-WT ubiquitin condition.

In conclusion, our functional data indicate that the co-expression of CYLD can increase the protein
stability of LRRK2.

AF-based modelling reveals a discrete repertoire of conformations and interfaces shaping the LRRK2
interactome

We predicted 3D complexes for binary interactions of LRRK2 and 196 partners in the BioID dataset via
AlphaFold-multimer (AF) and characterized their interfaces and conformations (Figure 4A).
We structurally aligned LRRK2 conformations in the different complexes by superimposing the
monomeric LRRK2 chains and then calculating the pairwise distance of all the possible pairs via Root
Mean Squared Deviation (RMSD) (Figure 4A; Methods). We performed hierarchical clustering of LRRK2
structures based on RMSD, through which we identified two main clusters (Figure 4B). A smaller one
is characterized by conformations with lower deviation from the inactive LRRK2 structure (PDB ID:
7LHW), which we termed "locked" (Figure 4B-D). The main structural hallmarks characterizing the
"locked" conformation and inherited by the experimental inactive LRRK2 structure (PDB:7LHW)
(Myasnikov et al., 2021), are the LRR wrapping around the catalytic domains, particularly the kinase,
and a contact formed by the extended loop within the LRR repeats, which includes the hinge helix, and
the C-terminal WD40 (Myasnikov et al., 2021). A bigger cluster comprised conformations that are
characterized by a larger deviation from the inactive structure, due to a major rearrangement of the
LRR, ROC and COR domains, which we termed "unlocked" (Figure 4B-D). Such conformations are
reminiscent of the ones that we observed in the predicted LRRK2-RAB10 complex, which we validated
as the AF-predicted conformation best accommodating experimental XL-MS data (Guaitoli et al.,

2023). Intriguingly, we found that proteins interacting with LRRK2 in these two alternative
conformations are associated with distinct evolutionary and functional properties. Indeed, interactors
engaging with LRRK2 in the "locked" conformations are characterized by a significantly lower RMSD
(P-value=3.81E-20; Figure 4D) and higher co-evolution (P-value=1.33E-3; Figure 4E). The two groups
are also characterized by largely exclusively enriched processes (Figure 4F). In particular, the
interactors engaging with LRRK2 in the locked conformation are significantly associated to processes
linked to centriolar satellites (P-value=4.76E-10) and cilium assembly (P-value= 0.0001) (Figure 4G;
Supplemental Table 4). Those interacting with the unlocked conformations are associated with a
higher number of enriched processes, including cluster-specific ones such as cytoskeleton (P-value=
4.53E-09) or the microtubule organizing center MTOC (P-value= 6.40E-07). Hence, the predicted LRRK2
conformational switch might be relevant in orchestrating the interaction with partners associated with
distinct functions.

We also explored patterns of engagement of interactors with the different regions of the LRRK2 multi-
domain architecture. To this end, we considered the distribution of distance probabilities (i.e.
distograms) generated by AF for each binary interaction and mapped the most likely interacting
residues of the partners with respect to LRRK2 positions. We used such partner-specific "interaction
fingerprint" to cluster the interactome, through which we identified four clusters of proteins named
after the specific LRRK2 3D interfaces involved in the interaction (Figure 5A; see Methods): "Core"
comprises partners interacting with the LRR-ROC-COR-KIN domains (Figure 5A,B), "WD40" entails
complexes with the omonymous domain (Figure 5A,C), "ARM" complexes with a new site within the
Armadillo domain (Figure 5A,D) and "Other" complexes binding in alternative modalities or with no
specific binding interface (Figure 5A). The complexes predicted to link to the "Core" group have a
significantly higher AF confidence scores with respect to clusters of complexes engaging with either
WD40 domain (P-value=1.14E-8) and in "Other" modality (P-value=4.86E-14; Figure 5E). Complexes of
the Core group are also characterized by significantly higher RMSD from interactors of the WD40 one
(P-value=4.05E-3; Figure 5F), which bind LRRK2 in both locked and unlocked conformations (see
below). In the Core cluster we found established LRRK2 interactors, such as DVL1-2, confirming the
known interaction mechanism through the ROC-COR domains (Sancho et al., 2009). In the same
cluster, we also found novel interactors discovered within the BioID set, such as e.g. CYLD interacting
through the Core interface in a similar way. A second cluster displayed higher likelihood of interaction
with the WD40 domain (Figure 5A,C), with predicted complexes characterized by a similar interface
involving the binding of a short, disordered peptide stretch to a conserved crevice formed by β-
propeller loops of the WD40 domain (Figure 5C). WD40 domain binders might bind to LRRK2 in either
locked or unlocked conformation (Figure 5A, F), as also suggested by a bi-modal distribution of the
LRRK2 chain RMSD (Figure 5F) These interactors also displayed significantly higher co-evolution with

respect to "Other" LRRK2 interactors (P-value=5.57E-4; Figure 5G). The ARM cluster involves proteins

showing binding preferences for a new docking site on the Armadillo domain, centered around

residues 500-600 and mediating the interaction with proteins such as AXIN1 (Figure 5A,D). The larger

cluster (Other) contains proteins interacting with weaker likelihood and variable docking sites (Figure

5A). Among these, we also found proteins engaging with two more ARM docking sites, including one

(around aa 240) which has been recently shown to interact with RAB12 (Dhekne et al., 2023; Li et al.,

2024) and mediate a RAB29-independent LRRK2 Kinase activity activation (Purlyte et al., 2018; Zhu et

al., 2023), which is predicted to interact with proteins such as KIF16B. A third ARM binding region,

encompassing aa 354-444, corresponds to the RAB29 interaction interface found in the recently

determined RAB29-bound, tetrameric LRRK2 structure (Zhu et al., 2023), and is the predicted interface

for proteins such as NBR1 and MON2.

Overall, these results further support the hypothesis that interactors *de-novo* identified by proximity

labelling may physically interact with LRRK2 domains through specific conformations and contact

interfaces. Interactors lacking a strong interaction probability in the distogram might represent

proteins interacting with LRRK2 either transiently or indirectly.

Type I but not type II inhibitors lead to conformational changes directing LRRK2 to centriolar satellites

As mentioned above, LRRK2 exists in multiple conformations. Effector-mediated oligomerization to an

asymmetric tetramer has been demonstrated to switch LRRK2 to an active conformation (Zhu et al.,

2023). In addition, ATP-competitive kinase inhibitors can stabilize different conformations. While type

I inhibitors stabilize the kinase domain in an active-like conformation, type II inhibitors stabilize the

inactive conformation of the kinase domain (Deniston et al., 2020; Raig et al., 2024; Zhu et al., 2024).

Interestingly, type I but not type II inhibitors lead to a rapid dephosphorylation of LRRK2 phospho-sites

in an interdomain space downstream of the hinge-helix which locks LRRK2 in a compact inactive fold

by binding the C-terminal WD40 domain (Kalogeropulou et al., 2022; Myasnikov et al., 2021). We

therefore reasoned that specific types of inhibitors could be used to discriminate the proximity

proteome of the active versus inactive conformation of LRRK2. For this reason, we compared a DMSO

control (Figure 6B) with the highly potent LRRK2-specific type I kinase inhibitor, MLi-2 with the type II

inhibitor GZD-824 (Figure 6A), which was previously shown to inhibit LRRK2 in cells (Kalogeropulou et

al., 2022; Tasegian et al., 2021). By Western-blot, we confirmed, that LRRK2 is sufficiently inhibited at

a final concentration of 1µM in a HEK293 line stably expressing miniTurbo-tagged LRRK2 (Supplemental

Figure 5). A quantitative comparison of the different conditions based on an LFQ analysis combined

with an FDR-controlled T-test revealed that only upon treatment of the cells with MLi-2 but not with

GZD-824, we identified a set of proteins in the LRRK2 proximity proteome almost exclusively enriched

in proteins of the cilia assembly/ centriolar satellites (Figure 6A-E). Intriguingly, interactors determined

in presence of the MLi-2 inhibitor are characterized by a significantly higher co-evolution (Figure 6F)

compared to other BioID interactors. The complete list of significantly enriched proteins (MLi-2 vs.

DMSO) is shown in Table 2. Additional information is provided in Supplemental Table 1.

RAB29 co-expression directs LRRK2 to well established lysosomal protein networks

It has been previously shown that LRRK2 interacts with the RAB29/RAB32/RAB38 family of small G-

proteins and subsequently locates to post-Golgi endomembranes (Beilina et al., 2014; McGrath et al.,

2021; Vides et al., 2022; Waschbusch et al., 2014). Furthermore, RAB29 not only recruits LRRK2 to

lysosomal membranes but also induces its tetramerization (Zhu et al., 2023). For this reason, we

performed the BioID experiment in dependence of RAB29 co-expression to determine PPI networks

associated with the activated oligomeric state of LRRK2. In addition to mT-LRRK2, either RAB29 or an

empty vector were co-expressed. The resulting differential proximity proteome showed differentially

enriched proteins only for the condition where RAB29 was co-expressed, demonstrating that a small

portion of LRRK2 was recruited by RAB29 with lesser impact on the total proximity proteome (Figure

6G). This differential proteome was enriched in proteins involved in vesicular transport, including the

post-Golgi/ lysosomal compartment as revealed by an LFQ analysis combined with an FDR-controlled

T-test (Figure 6H,I). In particular, known LRRK2 effectors, namely RAB8A (Steger et al., 2017) and

SPAG9 (Boecker and Holzbaur, 2021; Bonet-Ponce et al., 2020; Kluss et al., 2022a) were among the

enriched proteins (Figure 6G,H). Furthermore, VAMP7, which is part of a SNARE complex mediating

autophagosome-lysosome fusion (Jian et al., 2024), was among the significantly enriched proteins

(Figure 6G,H). VAMP7-LRRK2 physical interaction has recently been described (Filippini et al., 2023).

The complete list of significantly enriched proteins is shown in Table 3. Additional information is

provided in Supplemental Table 1.

MLi-2 and RAB29 differently modulate the shape the LRRK2 interactome

We also modelled the 3D structures of the binary complexes between LRRK2 and partners from both

the MLi-2 and RAB29 interactomes and analyzed complex interfaces as well as LRRK2's conformations

(Vides et al., 2022).

Interface-based clustering suggests that MLi-2 interactors engage with LRRK2 through specific domain

regions, particularly the ones involving terminal domains (Supplemental Figure 6A), while they are

predicted to engage little with the catalytic domains, particularly the kinase, which is consistent with

the MLi-2 inhibitor function, while in the RAB29 interactome we see a more variable pattern of

engaged interfaces. In particular, predictions for this interactome revealed a cluster of proteins

interacting with LRRK2 through the ROC-COR-KIN domains, including the RAB8A substrate

(Supplemental Figure 6B), suggesting that in the context of RAB29 overexpression several partners

might bind in a substrate-like manner. Notably, this interaction modality is not observed for predicted binary complexes of the MLi-2 interactome, confirming that the inhibitor prevents LRRK2 to bind substrate-like partners. Moreover, we further validated the capability of the distogram-based interface-fingerprint to discriminate functionally relevant LRRK2 interactors. We predicted the 3D complexes of LRRK2 with known interacting Rabs, and we found out that Rab substrates clustered separately from interacting RABs with distinct functions (e.g. RAB29 and RAB32; Supplemental Figure 7B).

From a LRRK2 conformational standpoint, we pooled the RAB29 and MLi-2 interactomes and clustered LRRK2 conformations based on RMSD, which similarly revealed a locked and an unlocked cluster. The interactors of RAB29 interactome are predicted to bind LRRK2 in both conformations, being particularly enriched in the unlocked (Figure 7A). On the other hand, we found a significant enrichment of MLi-2 interactors in the locked LRRK2 conformation (Fisher Exact Test P-value=0.023), further suggesting that the inhibitor is predicted to favor the engagement with interaction partners which, unlike substrates, do not require the interaction with the kinase domain (Figure 7A). Also, for these interactors, the locked conformation has a significantly lower RMSD (P-value=3.85E-6; Figure 7D) and a tendency towards higher coevolution (Figure 7E). Notably, when performing the enrichment of interaction partners grouped based on conformational clustering, we could find even more cluster-specific enrichment, with only one process in common between the partners in the locked and unlocked conformations (Figure 7F, G; Supplementary Table 6).

Discussion

The systematic analysis of the interactomes of disease-associated proteins is a powerful tool to dissect molecular networks and to identify pathways relevant for the molecular pathophysiology, for example demonstrated by the comprehensive interactomic network underlying the assembly and homeostasis of primary cilia (Boldt et al., 2016). Given its complex domain structure and tight regulation, LRRK2 is most likely representing a scaffold protein orchestrating distinct signaling complexes, which makes the analysis of its PPI networks an attractive approach to dissect context-specific signaling pathways (Gloeckner and Porras, 2020). While a wealth of data is available for affinity-based identification of protein-protein interactions, which have a bias for rather stable complexes, there is an unmet need for the dissection of less stable and transient PPIs. The rapid development of biotin-based proximity labelling technologies allowed to get information about this dark space the interactome (Qin et al., 2021) and, in consequence, BioID approaches emerged to widely used methods to dissect PPI networks. Most recently, proximity and affinity-based methods have even been shown to provide complementary information of the interactomes (Moreira et al., 2023). We therefore systematically

studied the proximity interactome of LRRK2 applying three different BioID tags, BioID1, BioID2 and

miniTurbo. To identify direct LRRK2 interactors with high confidence form the proximity proteomes,

we applied a newly developed bioinformatic pipeline relaying on co-evolution, *co-expression and*

protein domain architecture to predict interaction type (Miglionico et al., 2024, in prep.). This pipeline

allowed the identification of a group of proteins within the proximity proteome showing a high degree

of co-evolution with LRRK2 which is highly enriched in proteins associated with the centrosome. This

is particularly interesting as pathogenic LRRK2 has previously been shown to lead to a centrosome

cohesion defect as well as to a shortening of primary cilia (Dhekne et al., 2018; Khan et al., 2024; Lara

Ordonez et al., 2022). Furthermore, defects in primary cilia emerges as a new pathophysiological

scheme connecting different genes involved in mendelian forms of Parkinson's disease (Mohd Rafiq et

al., 2024; Schmidt et al., 2022). Strikingly, in the cluster of proteins with higher co-evolution to LRRK2

(Cluster 2), the single protein showing the highest co-evolutionary association was the deubiquitinase

CYLD. The K-63 selective deubiquitinase CYLD is a key effector in the biology of primary cilia.

Furthermore, preliminary functional data indicate that CYLD, when co expressed, is able to increase

expression levels of the pathogenic variant LRRK2 Y1699C which shows a decreased protein stability.

In fact, this DUB has been shown to control apical docking of basal bodies in ciliated epithelial cells

(Eguether et al., 2014). It also regulates centrosomal satellites proteostasis by counteracting the E3

ubiquitin ligase MIB1 (Douanne et al., 2019). Furthermore, CYLD has been shown to deubiquitinate

Cep70 and to inactivate HDAC6 thereby mediating ciliogenesis (Yang et al., 2014). Interestingly, a

missense mutation has recently been shown to be a causative gene for FTD-ALS (Dobson-Stone et al.,

2020). Most importantly, as the deubiquitinase CYLD is part of the PINK1/parkin pathway to regulate

cellular levels of PARIS (ZNF746), it is a critical regulator of mitochondrial biogenesis thereby acting as

a negative regulator of dopamine neuron survival in Parkinson's disease. In fact, CYLD inhibition has

been shown to ameliorate mitochondrial pathologies in cellular PD models which makes CYLD a

promising target for therapeutic intervention for PD and other neurodegenerative diseases (Pirooznia

et al., 2022). The E3 ligase parkin preserves mitochondrial homeostasis via the PARIS/PGC-1a axis

(Stevens et al., 2015; Zheng et al., 2017). Noteworthy, the PD risk variant LRRK2 G2385R has recently

been shown to alter mitochondrial biogenesis via the PGC-1a pathway, leading to reduced PGC-1a

levels similarly to a PARKIN knock-out in DA neurons (Kumar et al., 2020; Xue et al., 2023). PARKIN-

deficient DA neurons show increased PARIS and decreased PGC-1a levels (Kumar et al., 2020). In

conclusion, this indicates that LRRK2/CYLD signaling at one hand and Parkin signaling at the other hand

might converge at the level of PARIS.

The second most co-evolved interactor found in Cluster 2 was the cilia-associated protein CEP85, which

plays a role in centrosome disjunction (Chen et al., 2019). This is of particular interest as pathogenic

LRRK2 variants lead to a centrosomal cohesion phenotype in patient-derived cells (Lara Ordonez et al.,

2019). Notably, the cohesion phenotype is also seen in EBV-transformed lymphoblastoid cell lines from

iPD patients which can be reverted by LRRK2 kinase inhibition (Naaldijk et al., 2024). Importantly,

Cluster 2 also contains the autophagy cargo protein p62/sequestosome-1. This protein has previously

been identified as LRRK2 interactor and substrate (Kalogeropulou et al., 2018; Park et al., 2016).

SQSTM1 is also linked to PD pathophysiology via the Pink1/Parkin pathway (Shin et al., 2020). In

addition, Cluster 2 points towards other pathways which have been previously linked to PD pathology.

In particular, SNAP47 is part of the STX17-SNAP47-VAMP7/8 pathway, which has been shown to be

involved in selective autophagy – a process not only essential for the homeostasis of primary cilia but

also for mitophagy (Jian et al., 2024). Phospho-ubiquitin chains generated by PINK1/Parkin activity can

recruit autophagy adaptors including the ubiquitin adaptor SQSTM1/p62 linking the PINK1/Parkin-

dependent pathway to mitophagy (Geisler et al., 2010). All three proteins participate in the

recruitment of the autophagy machinery to aged or dysfunctional mitochondria, which are eventually

enclosed inside an autophagosome and subjected to the lysosome for degradation (Wilhelm et al.,

2022).

Interestingly, also the kinesin motor protein and PX-domain containing protein KIF16B was among the

top hits which is involved in vesicle trafficking and receptor recycling. It binds to PdIns(3)P by its unique

PX domain and promotes the circulating cargo movement from early endosomes to the plasma

membrane (Li et al., 2020). Interestingly, it has been found to be associated with lipid droplets

(Bersuker et al., 2018). In turn, increased RAB8a phosphorylation by pathogenic LRRK2 variants, has

been shown to promote lipid storage (Yu et al., 2018), indicating that the functional interaction of

LRRK2 and KIF16B might contribute to alterations in the neuronal lipid homeostasis associated with PD

pathology (Hallett et al., 2019).

Furthermore, induced by conformational changes upon MLi-2 inhibition, associated with a rapid and

almost complete dephosphorylation of LRRK2 in a disordered domain downstream of the hinge helix,

we identified a molecular network around PCM1, OFD1 and CEP131 strongly suggesting a link between

LRRK2 and centriolar satellites. Strikingly, the proteins interacting with LRRK2 in presence of MLi-2

displayed the highest degree of co-evolution, suggesting that the connection between LRRK2 and

centriolar satellites is highly conserved. The conformational changes induced by binding of type I

inhibitors might result in a phase transition of LRRK2 into centriolar satellites preventing 14-3-3 binding

and thereby shifting the equilibrium towards the dephosphorylated protein. A similar mechanism has

been shown for CEP131, which is extracted from centriolar satellites by phosphorylation and

subsequent binding of 14-3-3 (Tollenaere et al., 2015). Centriolar satellites not only play an essential

role in the biogenesis of primary cilia (Hall et al., 2023) but also play a role in the degradation of

proteins by driving the assembly of aggresomes (Prosser et al., 2022). In fact, also LRRK2 has been

shown to interfere with aggresome formation preceding autophagic clearance (Bang et al., 2016).
Nevertheless, one caveat might be seen in the fact that LRRK2 inhibition with type I inhibitors induces
its destabilization and degradation.

Structural prediction of LRRK2 binary complexes with AF revealed distinct interfaces within each
determined interactome. While several interactors preferentially engage with the catalytic core of the
protein, including the ROC and KIN domains, others are more likely to interact on terminal domains
(e.g. ARM or WD40). We predicted the interaction on the kinase domain for several interactors of the
BioID interactome as well as the one obtained in the presence of RAB29 over-expression, including the
known substrate RAB8A. The engagement in a substrate-like fashion with the catalytic core entails the
opening of the LRRK2 structure, with the detachment of the LRR and rearrangement of the entire N-
term domains. We previously shown that this conformation, that we similarly predicted for the LRRK2:
RAB10 complex, is compatible with distance restraint obtained by mass spectrometric mapping of
chemical crosslinking sites via XL-MS (Guaitoli et al., 2023). Moreover, if superimposed on the active-
like conformation of the most recent LRRK2 tetrameric structure, such predicted complexes fit very
well on the rest of the oligomer, with no clashes between the modelled LRRK2 chain and the rest of
the tetramer, further supporting the plausibility of such complexes (Guaitoli et al., 2023). We speculate
that partners engaging in a such a way might represent putative substrate or might regulate the kinase
domain function. Strikingly, we rarely observed such interaction interface for interactors enriched in
presence of the MLi-2 inhibitor, which bind more frequently to the locked conformation of LRRK2.

In conclusion, combining proximity proteomics with co-evolutionary and structural information helped
identifying distinct molecular networks linking LRRK2 to centrosomal biology and primary cilia. The
data may foster future functional analysis to father dissect these pathways to get a detailed picture of
the molecular pathophysiology underlying the cilia phenotypes caused by genetic variants in different
genes underlying familial forms of Parkinson's disease.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium
via the PRIDE (Perez-Riverol et al., 2022) partner repository with the dataset identifier PXD048832 and
10.6019/PXD048832 (BioID1, BioID2, miniTurbo), PXD048806 and 10.6019/PXD048806 (LRRK2
proximity-proteomes MLi-2 vs. GZD-824) as well as PXD048808 and 10.6019/PXD048808 (RAB29 co-
expression). The data will be released with the final publication of the peer-reviewed manuscript. The
BioID interaction network and associated co-evolutionary and structural analysis can accessed at the

following URL: (https://dip.bioinfolab.sns.it/explore/). All data presented in the preprint version of the manuscript, including MS RAW files, can be obtained upon request.

Author contributions

M.E. and F.I. performed BioID experiments. M.E., F.I., P.M., F.R. and C.J.G. analyzed the primary BioID data. P.M. and F.R. performed the post-analysis of primary BioID data. P.M., N.D.R. and F.R. wrote the code for the bioinformatic workflow. C.J.G., F.R. and M.U. designed the study. M.E., F.I., P.M., F.R. and C.J.G. wrote the manuscript.

Acknowledgements

We are grateful to the staff of the Core Facility for Medical Proteomics at the medical faculty of the University of Tübingen for technical assistance. Furthermore, we gratefully acknowledge computational resources of the Center for High Performance Computing (CHPC) at SNS. CJG received funds from The Michael J. Fox Foundation for Parkinson's Research (grant ID: MJFF-8068.02, MJFF-8068.04 and MJFF-023920) and iMed – the Helmholtz Initiative on Personalized Medicine.

Literature

Almawi, A.W., Matthews, L.A., and Guarne, A. (2017). FHA domains: Phosphopeptide binding and beyond. *Prog Biophys Mol Biol 127*, 105-110 doi: 10.1016/j.pbiomolbio.2016.12.003.

Altenhoff, A.M., Train, C.M., Gilbert, K.J., Mediratta, I., Mendes de Farias, T., Moi, D., Nevers, Y., Radoykova, H.S., Rossier, V., Warwick Vesztrocy, A., Glover, N.M., and Dessimoz, C. (2021). OMA orthology in 2021: website overhaul, conserved isoforms, ancestral gene order and more. *Nucleic acids research 49*, D373-D379 doi: 10.1093/nar/gkaa1007.

Bang, Y., Kim, K.S., Seol, W., and Choi, H.J. (2016). LRRK2 interferes with aggresome formation for autophagic clearance. *Mol Cell Neurosci 75*, 71-80 doi: 10.1016/j.mcn.2016.06.007.

Beilina, A., Rudenko, I.N., Kaganovich, A., Civiero, L., Chau, H., Kalia, S.K., Kalia, L.V., Lobbestael, E., Chia, R., Ndukwe, K., Ding, J., Nalls, M.A., International Parkinson's Disease Genomics, C., North American Brain Expression, C., Olszewski, M., Hauser, D.N., Kumaran, R., Lozano, A.M., Baekelandt, V., Greene, L.E., Taymans, J.M., Greggio, E., and Cookson, M.R. (2014). Unbiased screen for interactors of leucine-rich repeat kinase 2 supports a common pathway for sporadic and familial Parkinson disease. *Proceedings of the National Academy of Sciences of the United States of America 111*, 2626-2631 doi: 10.1073/pnas.1318306111.

Bersuker, K., Peterson, C.W.H., To, M., Sahl, S.J., Savikhin, V., Grossman, E.A., Nomura, D.K., and Olzmann, J.A. (2018). A Proximity Labeling Strategy Provides Insights into the Composition and Dynamics of Lipid Droplet Proteomes. *Dev Cell 44*, 97-112 e117 doi: 10.1016/j.devcel.2017.11.020.

Boecker, C.A., and Holzbaur, E.L.F. (2021). Hyperactive LRRK2 kinase impairs the trafficking of axonal
autophagosomes. *Autophagy 17*, 2043-2045 doi: 10.1080/15548627.2021.1936933.

Boldt, K., van Reeuwijk, J., Lu, Q., Koutroumpas, K., Nguyen, T.M., Texier, Y., van Beersum, S.E., Horn,
N., Willer, J.R., Mans, D.A., Dougherty, G., Lamers, I.J., Coene, K.L., Arts, H.H., Betts, M.J., Beyer, T.,
Bolat, E., Gloeckner, C.J., Haidari, K., Hetterschijt, L., Iaconis, D., Jenkins, D., Klose, F., Knapp, B*., et
al.* (2016). An organelle-specific protein landscape identifies novel diseases and molecular
mechanisms. *Nature communications 7*, 11491 doi: 10.1038/ncomms11491.

Bonet-Ponce, L., Beilina, A., Williamson, C.D., Lindberg, E., Kluss, J.H., Saez-Atienzar, S., Landeck, N.,
Kumaran, R., Mamais, A., Bleck, C.K.E., Li, Y., and Cookson, M.R. (2020). LRRK2 mediates tubulation
and vesicle sorting from lysosomes. *Sci Adv 6* doi: 10.1126/sciadv.abb2454.

Branon, T.C., Bosch, J.A., Sanchez, A.D., Udeshi, N.D., Svinkina, T., Carr, S.A., Feldman, J.L., Perrimon,
N., and Ting, A.Y. (2018). Efficient proximity labeling in living cells and organisms with TurboID. *Nat
Biotechnol 36*, 880-887 doi: 10.1038/nbt.4201.

Carrion, M.D.P., Marsicano, S., Daniele, F., Marte, A., Pischedda, F., Di Cairano, E., Piovesana, E., von
Zweydorf, F., Kremmer, E., Gloeckner, C.J., Onofri, F., Perego, C., and Piccoli, G. (2017). The LRRK2
G2385R variant is a partial loss-of-function mutation that affects synaptic vesicle trafficking through
altered protein interactions. *Sci Rep 7*, 5377 doi: 10.1038/s41598-017-05760-9.

Chen, C., Xu, Z., Zhang, T., Lin, L., Lu, M., Xie, C., and Yu, X. (2019). Cep85 Relays Plk1 Activity to
Phosphorylated Nek2A for Its Timely Activation in Centrosome Disjunction. *iScience 11*, 114-133
doi: 10.1016/j.isci.2018.12.013.

Cock, P.J., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I., Hamelryck, T., Kauff,
F., Wilczynski, B., and de Hoon, M.J. (2009). Biopython: freely available Python tools for
computational molecular biology and bioinformatics. *Bioinformatics 25*, 1422-1423 doi:
10.1093/bioinformatics/btp163.

Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate proteome-wide
label-free quantification by delayed normalization and maximal peptide ratio extraction, termed
MaxLFQ. *Molecular & cellular proteomics : MCP 13*, 2513-2526 doi: 10.1074/mcp.M113.031591.

Deniston, C.K., Salogiannis, J., Mathea, S., Snead, D.M., Lahiri, I., Matyszewski, M., Donosa, O.,
Watanabe, R., Bohning, J., Shiau, A.K., Knapp, S., Villa, E., Reck-Peterson, S.L., and Leschziner, A.E.
(2020). Structure of LRRK2 in Parkinson's disease and model for microtubule interaction. *Nature
588*, 344-349 doi: 10.1038/s41586-020-2673-2.

Dhekne, H.S., Tonelli, F., Yeshaw, W.M., Chiang, C.Y., Limouse, C., Jaimon, E., Purlyte, E., Alessi, D.R.,
and Pfeffer, S.R. (2023). Genome-wide screen reveals Rab12 GTPase as a critical activator of
Parkinson's disease-linked LRRK2 kinase. *Elife 12* doi: 10.7554/eLife.87098.

Dhekne, H.S., Yanatori, I., Gomez, R.C., Tonelli, F., Diez, F., Schule, B., Steger, M., Alessi, D.R., and Pfeffer, S.R. (2018). A pathway for Parkinson's Disease LRRK2 kinase to block primary cilia and Sonic hedgehog signaling in the brain. *Elife 7* doi: 10.7554/eLife.40202.

Dobson-Stone, C., Hallupp, M., Shahheydari, H., Ragagnin, A.M.G., Chatterton, Z., Carew-Jones, F., Shepherd, C.E., Stefen, H., Paric, E., Fath, T., Thompson, E.M., Blumbergs, P., Short, C.L., Field, C.D., Panegyres, P.K., Hecker, J., Nicholson, G., Shaw, A.D., Fullerton, J.M., Luty, A.A., Schofield, P.R., Brooks, W.S., Rajan, N., Bennett, M.F.*, et al.* (2020). CYLD is a causative gene for frontotemporal dementia - amyotrophic lateral sclerosis. *Brain 143*, 783-799 doi: 10.1093/brain/awaa039.

Doncheva, N.T., Morris, J.H., Gorodkin, J., and Jensen, L.J. (2019). Cytoscape StringApp: Network Analysis and Visualization of Proteomics Data. *Journal of proteome research 18*, 623-632 doi: 10.1021/acs.jproteome.8b00702.

Douanne, T., Andre-Gregoire, G., Thys, A., Trillet, K., Gavard, J., and Bidere, N. (2019). CYLD Regulates Centriolar Satellites Proteostasis by Counteracting the E3 Ligase MIB1. *Cell Rep 27*, 1657-1665 e1654 doi: 10.1016/j.celrep.2019.04.036.

Eguether, T., Ermolaeva, M.A., Zhao, Y., Bonnet, M.C., Jain, A., Pasparakis, M., Courtois, G., and Tassin, A.M. (2014). The deubiquitinating enzyme CYLD controls apical docking of basal bodies in ciliated epithelial cells. *Nature communications 5*, 4585 doi: 10.1038/ncomms5585.

Ehebauer, M.T., and Arias, A.M. (2009). The structural and functional determinants of the Axin and Dishevelled DIX domains. *BMC Struct Biol 9*, 70 doi: 10.1186/1472-6807-9-70.

Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A., Bates, R., Blackwell, S., Yim, J., Ronneberger, O., Bodenstein, S., Zielinski, M., Bridgland, A., Potapenko, A., Cowie, A., Tunyasuvunakool, K., Jain, R., Clancy, E., Kohli, P., Jumper, J., and Hassabis, D. (2021). Protein complex prediction with AlphaFold-Multimer. Preprint available at: *bioRxiv* doi: 10.1101/2021.10.04.463034.

Fahn, S. (2003). Description of Parkinson's disease as a clinical syndrome. *Annals of the New York Academy of Sciences 991*, 1-14 doi: 10.1111/j.1749-6632.2003.tb07458.x.

Filippini, F., Nola, S., Zahraoui, A., Roger, K., Esmaili, M., Sun, J., Wojnacki, J., Vlieghe, A., Bun, P., Blanchon, S., Rain, J.C., Taymans, J.M., Chartier-Harlin, M.C., Guerrera, C., and Galli, T. (2023). Secretion of VGF relies on the interplay between LRRK2 and post-Golgi v-SNAREs. *Cell Rep 42*, 112221 doi: 10.1016/j.celrep.2023.112221.

Follett, J., and Farrer, M.J. (2021). LRRK2; a dynamic regulator of cellular trafficking. *Brain Res 1761*, 147394 doi: 10.1016/j.brainres.2021.147394.

Galaxy, C. (2024). The Galaxy platform for accessible, reproducible, and collaborative data analyses: 2024 update. *Nucleic acids research 52*, W83-W94 doi: 10.1093/nar/gkae410.

Gasser, T., Hardy, J., and Mizuno, Y. (2011). Milestones in PD genetics. *Mov Disord 26*, 1042-1048 doi:
10.1002/mds.23637.

Geisler, S., Holmstrom, K.M., Skujat, D., Fiesel, F.C., Rothfuss, O.C., Kahle, P.J., and Springer, W. (2010).
PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat Cell Biol 12*, 119-
131 doi: 10.1038/ncb2012.

Gloeckner, C.J., Boldt, K., Schumacher, A., Roepman, R., and Ueffing, M. (2007). A novel tandem affinity
purification strategy for the efficient isolation and characterisation of native protein complexes.
Proteomics 7, 4228-4234 doi: 10.1002/pmic.200700038.

Gloeckner, C.J., Boldt, K., and Ueffing, M. (2009). Strep/FLAG tandem affinity purification (SF-TAP) to
study protein interactions. *Current protocols in protein science / editorial board, John E Coligan [et
al] Chapter 19*, Unit19 20 doi: 10.1002/0471140864.ps1920s57.

Gloeckner, C.J., and Porras, P. (2020). Guilt-by-Association - Functional Insights Gained From Studying
the LRRK2 Interactome. *Front Neurosci 14*, 485 doi: 10.3389/fnins.2020.00485.

Grishin, N.V. (2001). KH domain: one motif, two folds. *Nucleic acids research 29*, 638-643 doi:
10.1093/nar/29.3.638.

Guaitoli, G., Raimondi, F., Gilsbach, B.K., Gomez-Llorente, Y., Deyaert, E., Renzi, F., Li, X., Schaffner, A.,
Jagtap, P.K., Boldt, K., von Zweydorf, F., Gotthardt, K., Lorimer, D.D., Yue, Z., Burgin, A., Janjic, N.,
Sattler, M., Versees, W., Ueffing, M., Ubarretxena-Belandia, I., Kortholt, A., and Gloeckner, C.J.
(2016). Structural model of the dimeric Parkinson's protein LRRK2 reveals a compact architecture
involving distant interdomain contacts. *Proceedings of the National Academy of Sciences of the
United States of America 113*, E4357-4366 doi: 10.1073/pnas.1523708113.

Guaitoli, G., Zhang, X., Saitta, F., Miglionico, P., Silbermann, L.M., Ho, F.Y., von Zweydorf, F., Signorelli,
M., Tych, K., Fessas, D., Raimondi, F., Kortholt, A., and Gloeckner, C.J. (2023). Biophysical analysis
reveals autophosphorylation as an important regulator of LRRK2 dimerization. Preprint available at:
bioRxiv doi: 10.1101/2023.08.11.549911.

Hall, E.A., Kumar, D., Prosser, S.L., Yeyati, P.L., Herranz-Perez, V., Garcia-Verdugo, J.M., Rose, L., McKie,
L., Dodd, D.O., Tennant, P.A., Megaw, R., Murphy, L.C., Ferreira, M.F., Grimes, G., Williams, L.,
Quidwai, T., Pelletier, L., Reiter, J.F., and Mill, P. (2023). Centriolar satellites expedite mother
centriole remodeling to promote ciliogenesis. *Elife 12* doi: 10.7554/eLife.79299.

Hallett, P.J., Engelender, S., and Isacson, O. (2019). Lipid and immune abnormalities causing age-
dependent neurodegeneration and Parkinson's disease. *J Neuroinflammation 16*, 153 doi:
10.1186/s12974-019-1532-2.

Hori, A., and Toda, T. (2017). Regulation of centriolar satellite integrity and its physiology. *Cellular and
molecular life sciences : CMLS 74*, 213-229 doi: 10.1007/s00018-016-2315-x.

Islam, M.S., Nolte, H., Jacob, W., Ziegler, A.B., Putz, S., Grosjean, Y., Szczepanowska, K., Trifunovic, A.,

Braun, T., Heumann, H., Heumann, R., Hovemann, B., Moore, D.J., and Kruger, M. (2016). Human

R1441C LRRK2 regulates the synaptic vesicle proteome and phosphoproteome in a Drosophila

model of Parkinson's disease. *Human molecular genetics 25*, 5365-5382 doi: 10.1093/hmg/ddw352.

Jennings, D., Huntwork-Rodriguez, S., Henry, A.G., Sasaki, J.C., Meisner, R., Diaz, D., Solanoy, H., Wang,

X., Negrou, E., Bondar, V.V., Ghosh, R., Maloney, M.T., Propson, N.E., Zhu, Y., Maciuca, R.D., Harris,

L., Kay, A., LeWitt, P., King, T.A., Kern, D., Ellenbogen, A., Goodman, I., Siderowf, A., Aldred, J.*, et al.*

(2022). Preclinical and clinical evaluation of the LRRK2 inhibitor DNL201 for Parkinson's disease.

Science translational medicine 14, eabj2658 doi: 10.1126/scitranslmed.abj2658.

Jian, F., Wang, S., Tian, R., Wang, Y., Li, C., Li, Y., Wang, S., Fang, C., Ma, C., and Rong, Y. (2024). The

STX17-SNAP47-VAMP7/VAMP8 complex is the default SNARE complex mediating autophagosome-

lysosome fusion. *Cell Res* doi: 10.1038/s41422-023-00916-x.

Kafka, A., Basic-Kinda, S., and Pecina-Slaus, N. (2014). The cellular story of dishevelleds. *Croat Med J*

55, 459-467 doi: 10.3325/cmj.2014.55.459.

Kaleb, K., Vesztrocy, A.W., Altenhoff, A., and Dessimoz, C. (2019). Expanding the Orthologous Matrix

(OMA) programmatic interfaces: REST API and the OmaDB packages for R and Python. *F1000Res 8*,

42 doi: 10.12688/f1000research.17548.2.

Kalogeropulou, A.F., Purlyte, E., Tonelli, F., Lange, S.M., Wightman, M., Prescott, A.R., Padmanabhan,

S., Sammler, E., and Alessi, D.R. (2022). Impact of 100 LRRK2 variants linked to Parkinson's Disease

on kinase activity and microtubule binding. *The Biochemical journal* doi: 10.1042/BCJ20220161.

Kalogeropulou, A.F., Zhao, J., Bolliger, M.F., Memou, A., Narasimha, S., Molitor, T.P., Wilson, W.H.,

Rideout, H.J., and Nichols, R.J. (2018). P62/SQSTM1 is a novel leucine-rich repeat kinase 2 (LRRK2)

substrate that enhances neuronal toxicity. *The Biochemical journal 475*, 1271-1293 doi:

10.1042/BCJ20170699.

Kanao, T., Venderova, K., Park, D.S., Unterman, T., Lu, B., and Imai, Y. (2010). Activation of FoxO by

LRRK2 induces expression of proapoptotic proteins and alters survival of postmitotic dopaminergic

neuron in Drosophila. *Human molecular genetics 19*, 3747-3758 doi: 10.1093/hmg/ddq289.

Khan, S.S., Jaimon, E., Lin, Y.E., Nikoloff, J., Tonelli, F., Alessi, D.R., and Pfeffer, S.R. (2024). Loss of

primary cilia and dopaminergic neuroprotection in pathogenic LRRK2-driven and idiopathic

Parkinson's disease. Preprint available at: *bioRxiv* doi: 10.1101/2024.01.15.575737.

Kiel, C., Vogt, A., Campagna, A., Chatr-aryamontri, A., Swiatek-de Lange, M., Beer, M., Bolz, S., Mack,

A.F., Kinkl, N., Cesareni, G., Serrano, L., and Ueffing, M. (2011). Structural and functional protein

network analyses predict novel signaling functions for rhodopsin. *Molecular systems biology 7*, 551

doi: 10.1038/msb.2011.83.

Kim, D.I., Jensen, S.C., Noble, K.A., Kc, B., Roux, K.H., Motamedchaboki, K., and Roux, K.J. (2016). An

improved smaller biotin ligase for BioID proximity labeling. *Mol Biol Cell 27*, 1188-1196 doi:

10.1091/mbc.E15-12-0844.

Kluss, J.H., Bonet-Ponce, L., Lewis, P.A., and Cookson, M.R. (2022a). Directing LRRK2 to membranes of

the endolysosomal pathway triggers RAB phosphorylation and JIP4 recruitment. *Neurobiol Dis 170*,

105769 doi: 10.1016/j.nbd.2022.105769.

Kluss, J.H., Lewis, P.A., and Greggio, E. (2022b). Leucine-rich repeat kinase 2 (LRRK2): an update on the

potential therapeutic target for Parkinson's disease. *Expert Opin Ther Targets 26*, 537-546 doi:

10.1080/14728222.2022.2082937.

Kucera, M., Isserlin, R., Arkhangorodsky, A., and Bader, G.D. (2016). AutoAnnotate: A Cytoscape app

for summarizing networks with semantic annotations. *F1000Res 5*, 1717 doi:

10.12688/f1000research.9090.1.

Kumar, M., Acevedo-Cintron, J., Jhaldiyal, A., Wang, H., Andrabi, S.A., Eacker, S., Karuppagounder, S.S.,

Brahmachari, S., Chen, R., Kim, H., Ko, H.S., Dawson, V.L., and Dawson, T.M. (2020). Defects in

Mitochondrial Biogenesis Drive Mitochondrial Alterations in PARKIN-Deficient Human Dopamine

Neurons. *Stem Cell Reports 15*, 629-645 doi: 10.1016/j.stemcr.2020.07.013.

Lara Ordonez, A.J., Fasiczka, R., Fernandez, B., Naaldijk, Y., Fdez, E., Blanca Ramirez, M., Phan, S.,

Boassa, D., and Hilfiker, S. (2022). The LRRK2 signaling network converges on a centriolar phospho-

Rab10/RILPL1 complex to cause deficits in centrosome cohesion and cell polarization. *Biol Open 11*

doi: 10.1242/bio.059468.

Lara Ordonez, A.J., Fernandez, B., Fdez, E., Romo-Lozano, M., Madero-Perez, J., Lobbestael, E.,

Baekelandt, V., Aiastui, A., Lopez de Munain, A., Melrose, H.L., Civiero, L., and Hilfiker, S. (2019).

RAB8, RAB10 and RILPL1 contribute to both LRRK2 kinase-mediated centrosomal cohesion and

ciliogenesis deficits. *Human molecular genetics 28*, 3552-3568 doi: 10.1093/hmg/ddz201.

Li, B.J., Chen, H., Jiang, S.S., Wang, C.Y., Tuo, Q.H., Long, S.Y., Zhang, C.P., and Liao, D.F. (2020). PX

Domain-Containing Kinesin KIF16B and Microtubule-Dependent Intracellular Movements. *J Membr

Biol 253*, 101-108 doi: 10.1007/s00232-020-00110-9.

Li, X., Zhu, H., Huang, B.T., Li, X., Kim, H., Tan, H., Zhang, Y., Choi, I., Peng, J., Xu, P., Sun, J., and Yue, Z.

(2024). RAB12-LRRK2 Complex Suppresses Primary Ciliogenesis and Regulates Centrosome

Homeostasis in Astrocytes. Preprint available at: *bioRxiv*, 2024.2007.2017.603999 doi:

10.1101/2024.07.17.603999.

Lim, K.L., Chew, K.C., Tan, J.M., Wang, C., Chung, K.K., Zhang, Y., Tanaka, Y., Smith, W., Engelender, S.,

Ross, C.A., Dawson, V.L., and Dawson, T.M. (2005). Parkin mediates nonclassical, proteasomal-

independent ubiquitination of synphilin-1: implications for Lewy body formation. *J Neurosci 25*,

2002-2009 doi: 10.1523/JNEUROSCI.4474-04.2005.

Liu, X., Salokas, K., Weldatsadik, R.G., Gawriyski, L., and Varjosalo, M. (2020). Combined proximity
labeling and affinity purification-mass spectrometry workflow for mapping and visualizing protein
interaction networks. *Nature protocols 15*, 3182-3211 doi: 10.1038/s41596-020-0365-x.

Manzoni, C., Denny, P., Lovering, R.C., and Lewis, P.A. (2015). Computational analysis of the LRRK2
interactome. *PeerJ 3*, e778 doi: 10.7717/peerj.778.

Matic, M., Miglionico, P., Tatsumi, M., Inoue, A., and Raimondi, F. (2023). GPCRome-wide analysis of
G-protein-coupling diversity using a computational biology approach. *Nature communications 14*,
4361 doi: 10.1038/s41467-023-40045-y.

McGrath, E., Waschbusch, D., Baker, B.M., and Khan, A.R. (2021). LRRK2 binds to the Rab32 subfamily
in a GTP-dependent manner via its armadillo domain. *Small GTPases 12*, 133-146 doi:
10.1080/21541248.2019.1666623.

Meixner, A., Boldt, K., Van Troys, M., Askenazi, M., Gloeckner, C.J., Bauer, M., Marto, J.A., Ampe, C.,
Kinkl, N., and Ueffing, M. (2011). A QUICK screen for Lrrk2 interaction partners--leucine-rich repeat
kinase 2 is involved in actin cytoskeleton dynamics. *Molecular & cellular proteomics : MCP 10*, M110
001172 doi: 10.1074/mcp.M110.001172.

Meldal, B.H.M., Perfetto, L., Combe, C., Lubiana, T., Ferreira Cavalcante, J.V., Bye, A.J.H., Waagmeester,
A., Del-Toro, N., Shrivastava, A., Barrera, E., Wong, E., Mlecnik, B., Bindea, G., Panneerselvam, K.,
Willighagen, E., Rappsilber, J., Porras, P., Hermjakob, H., and Orchard, S. (2022). Complex Portal
2022: new curation frontiers. *Nucleic acids research 50*, D578-D586 doi: 10.1093/nar/gkab991.

Mellacheruvu, D., Wright, Z., Couzens, A.L., Lambert, J.P., St-Denis, N.A., Li, T., Miteva, Y.V., Hauri, S.,
Sardiu, M.E., Low, T.Y., Halim, V.A., Bagshaw, R.D., Hubner, N.C., Al-Hakim, A., Bouchard, A.,
Faubert, D., Fermin, D., Dunham, W.H., Goudreault, M., Lin, Z.Y., Badillo, B.G., Pawson, T., Durocher,
D., Coulombe, B.*, et al.* (2013). The CRAPome: a contaminant repository for affinity purification-
mass spectrometry data. *Nature methods 10*, 730-736 doi: 10.1038/nmeth.2557.

Miglionico, P., Nemati Fard, L.A., Gloeckner, C.J., and Raimondi, F. (2024, in prep.). Predicition and
discovery of protein-protein direct interactions and stable complexes based on gene co-expression
and co-evolution. *bioRxiv*, preprint in preparation.

Miller, G.K., Kuruvilla, S., Jacob, B., LaFranco-Scheuch, L., Bakthavatchalu, V., Flor, J., Flor, K., Ziegler,
J., Reichard, C., Manfre, P., Firner, S., McNutt, T., Quay, D., Bellum, S., Doto, G., Ciaccio, P.J.,
Pearson, K., Valentine, J., Fuller, P., Fell, M., Tsuchiya, T., Williamson, T., and Wollenberg, G. (2023).
Effects of LRRK2 Inhibitors in Nonhuman Primates. *Toxicol Pathol*, 1926233231205895 doi:
10.1177/01926233231205895.

Mohd Rafiq, N., Fujise, K., Rosenfeld, M.S., Xu, P., and De Camilli, P. (2024). Parkinsonism Sac domain
mutation in Synaptojanin-1 affects ciliary properties in iPSC-derived dopaminergic neurons.

Proceedings of the National Academy of Sciences of the United States of America 121, e2318943121 doi: 10.1073/pnas.2318943121.

Moreira, C., Kelemen, C.D., Obado, S.O., Zahedifard, F., Zhang, N., Holetz, F.B., Gauglitz, L., Dallagiovanna, B., Field, M.C., Kramer, S., and Zoltner, M. (2023). Impact of inherent biases built into proteomic techniques: Proximity labeling and affinity capture compared. *The Journal of biological chemistry 299*, 102726 doi: 10.1016/j.jbc.2022.102726.

Myasnikov, A., Zhu, H., Hixson, P., Xie, B., Yu, K., Pitre, A., Peng, J., and Sun, J. (2021). Structural analysis of the full-length human LRRK2. *Cell 184*, 3519-3527 e3510 doi: 10.1016/j.cell.2021.05.004.

Naaldijk, Y., Fernandez, B., Fasiczka, R., Fdez, E., Leghay, C., Croitoru, I., Kwok, J.B., Boulesnane, Y., Vizeneux, A., Mutez, E., Calvez, C., Destee, A., Taymans, J.M., Aragon, A.V., Yarza, A.B., Padmanabhan, S., Delgado, M., Alcalay, R.N., Chatterton, Z., Dzamko, N., Halliday, G., Ruiz-Martinez, J., Chartier-Harlin, M.C., and Hilfiker, S. (2024). A potential patient stratification biomarker for Parkinson s disease based on LRRK2 kinase-mediated centrosomal alterations in peripheral blood-derived cells. *NPJ Parkinsons Dis 10*, 12 doi: 10.1038/s41531-023-00624-8.

Orchard, S., Ammari, M., Aranda, B., Breuza, L., Briganti, L., Broackes-Carter, F., Campbell, N.H., Chavali, G., Chen, C., del-Toro, N., Duesbury, M., Dumousseau, M., Galeota, E., Hinz, U., Iannuccelli, M., Jagannathan, S., Jimenez, R., Khadake, J., Lagreid, A., Licata, L., Lovering, R.C., Meldal, B., Melidoni, A.N., Milagros, M.*, et al.* (2014). The MIntAct project--IntAct as a common curation platform for 11 molecular interaction databases. *Nucleic acids research 42*, D358-363 doi: 10.1093/nar/gkt1115.

Paisan-Ruiz, C., Jain, S., Evans, E.W., Gilks, W.P., Simon, J., van der Brug, M., Lopez de Munain, A., Aparicio, S., Gil, A.M., Khan, N., Johnson, J., Martinez, J.R., Nicholl, D., Marti Carrera, I., Pena, A.S., de Silva, R., Lees, A., Marti-Masso, J.F., Perez-Tur, J., Wood, N.W., and Singleton, A.B. (2004). Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron 44*, 595-600 doi: 10.1016/j.neuron.2004.10.023.

Park, S., Han, S., Choi, I., Kim, B., Park, S.P., Joe, E.H., and Suh, Y.H. (2016). Interplay between Leucine-Rich Repeat Kinase 2 (LRRK2) and p62/SQSTM-1 in Selective Autophagy. *PloS one 11*, e0163029 doi: 10.1371/journal.pone.0163029.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., B., T., Griesel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., D., C., M., B., Perrot, M., and E., D. (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research 12*, 2825-2830 doi:

Perez-Riverol, Y., Bai, J., Bandla, C., Garcia-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S., Kundu, D.J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., Walzer, M., Wang, S., Brazma, A., and Vizcaino, J.A. (2022). The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic acids research 50*, D543-D552 doi: 10.1093/nar/gkab1038.

Piccoli, G., Condliffe, S.B., Bauer, M., Giesert, F., Boldt, K., De Astis, S., Meixner, A., Sarioglu, H., Vogt-
Weisenhorn, D.M., Wurst, W., Gloeckner, C.J., Matteoli, M., Sala, C., and Ueffing, M. (2011). LRRK2
controls synaptic vesicle storage and mobilization within the recycling pool. *J Neurosci 31*, 2225-
2237 doi: 10.1523/JNEUROSCI.3730-10.2011.

Piccoli, G., Onofri, F., Cirnaru, M.D., Kaiser, C.J., Jagtap, P., Kastenmuller, A., Pischedda, F., Marte, A.,
von Zweydorf, F., Vogt, A., Giesert, F., Pan, L., Antonucci, F., Kiel, C., Zhang, M., Weinkauf, S., Sattler,
M., Sala, C., Matteoli, M., Ueffing, M., and Gloeckner, C.J. (2014). Leucine-rich repeat kinase 2 binds
to neuronal vesicles through protein interactions mediated by its C-terminal WD40 domain. *Mol*
Cell Biol 34, 2147-2161 doi: 10.1128/MCB.00914-13.

Pirooznia, S.K., Wang, H., Panicker, N., Kumar, M., Neifert, S., Dar, M.A., Lau, E., Kang, B.G., Redding-
Ochoa, J., Troncoso, J.C., Dawson, V.L., and Dawson, T.M. (2022). Deubiquitinase CYLD acts as a
negative regulator of dopamine neuron survival in Parkinson's disease. *Sci Adv 8*, eabh1824 doi:
10.1126/sciadv.abh1824.

Prosser, S.L., Tkach, J., Gheiratmand, L., Kim, J., Raught, B., Morrison, C.G., and Pelletier, L. (2022).
Aggresome assembly at the centrosome is driven by CP110-CEP97-CEP290 and centriolar satellites.
Nat Cell Biol 24, 483-496 doi: 10.1038/s41556-022-00869-0.

Purlyte, E., Dhekne, H.S., Sarhan, A.R., Gomez, R., Lis, P., Wightman, M., Martinez, T.N., Tonelli, F.,
Pfeffer, S.R., and Alessi, D.R. (2018). Rab29 activation of the Parkinson's disease-associated LRRK2
kinase. *The EMBO journal 37*, 1-18 doi: 10.15252/embj.201798099.

Qin, W., Cho, K.F., Cavanagh, P.E., and Ting, A.Y. (2021). Deciphering molecular interactions by
proximity labeling. *Nature methods 18*, 133-143 doi: 10.1038/s41592-020-01010-5.

Ragueneau, E., Shrivastava, A., Morris, J.H., Del-Toro, N., Hermjakob, H., and Porras, P. (2021). IntAct
App: a Cytoscape application for molecular interaction network visualization and analysis.
Bioinformatics 37, 3684-3685 doi: 10.1093/bioinformatics/btab319.

Raig, N.D., Surridge, K.J., Sanz-Murillo, M., Dederer, V., Kramer, A., Schwalm, M.P., Elson, L., Chatterjee,
D., Mathea, S., Hanke, T., Leschziner, A.E., Reck-Peterson, S.L., and Knapp, S. (2024). Type-II kinase
inhibitors that target Parkinson's Disease-associated LRRK2. Preprint available at: *bioRxiv*,
2024.2009.2017.613365 doi: 10.1101/2024.09.17.613365.

Rousseeuw, P.J. (1987). Silhouettes: A graphical aid to the interpretation and validation of cluster
analysis. *Journal of Computational and Applied Mathematics 20*, 53-65 doi: 10.1016/0377-
0427(87)90125-7.

Roux, K.J., Kim, D.I., Raida, M., and Burke, B. (2012). A promiscuous biotin ligase fusion protein
identifies proximal and interacting proteins in mammalian cells. *J Cell Biol 196*, 801-810 doi:
10.1083/jcb.201112098.

Sancho, R.M., Law, B.M., and Harvey, K. (2009). Mutations in the LRRK2 Roc-COR tandem domain link
Parkinson's disease to Wnt signalling pathways. *Human molecular genetics 18*, 3955-3968 doi:
10.1093/hmg/ddp337.

Schmidt, S., Luecken, M.D., Trumbach, D., Hembach, S., Niedermeier, K.M., Wenck, N., Pflugler, K.,
Stautner, C., Bottcher, A., Lickert, H., Ramirez-Suastegui, C., Ahmad, R., Ziller, M.J., Fitzgerald, J.C.,
Ruf, V., van de Berg, W.D.J., Jonker, A.J., Gasser, T., Winner, B., Winkler, J., Vogt Weisenhorn, D.M.,
Giesert, F., Theis, F.J., and Wurst, W. (2022). Primary cilia and SHH signaling impairments in human
and mouse models of Parkinson's disease. *Nature communications 13*, 4819 doi: 10.1038/s41467-
022-32229-9.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and
Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res 13*, 2498-2504 doi: 10.1101/gr.1239303.

Shin, W.H., Park, J.H., and Chung, K.C. (2020). The central regulator p62 between ubiquitin proteasome
system and autophagy and its role in the mitophagy and Parkinson's disease. *BMB Rep 53*, 56-63
doi: 10.5483/BMBRep.2020.53.1.283.

Simon-Sanchez, J., Schulte, C., Bras, J.M., Sharma, M., Gibbs, J.R., Berg, D., Paisan-Ruiz, C., Lichtner, P.,
Scholz, S.W., Hernandez, D.G., Kruger, R., Federoff, M., Klein, C., Goate, A., Perlmutter, J., Bonin,
M., Nalls, M.A., Illig, T., Gieger, C., Houlden, H., Steffens, M., Okun, M.S., Racette, B.A., Cookson,
M.R.*, et al.* (2009). Genome-wide association study reveals genetic risk underlying Parkinson's
disease. *Nat Genet 41*, 1308-1312 doi: 10.1038/ng.487.

Sowa, M.E., Bennett, E.J., Gygi, S.P., and Harper, J.W. (2009). Defining the human deubiquitinating
enzyme interaction landscape. *Cell 138*, 389-403 doi: 10.1016/j.cell.2009.04.042.

Steger, M., Diez, F., Dhekne, H.S., Lis, P., Nirujogi, R.S., Karayel, O., Tonelli, F., Martinez, T.N., Lorentzen,
E., Pfeffer, S.R., Alessi, D.R., and Mann, M. (2017). Systematic proteomic analysis of LRRK2-
mediated Rab GTPase phosphorylation establishes a connection to ciliogenesis. *Elife 6* doi:
10.7554/eLife.31012.

Stevens, D.A., Lee, Y., Kang, H.C., Lee, B.D., Lee, Y.I., Bower, A., Jiang, H., Kang, S.U., Andrabi, S.A.,
Dawson, V.L., Shin, J.H., and Dawson, T.M. (2015). Parkin loss leads to PARIS-dependent declines in
mitochondrial mass and respiration. *Proceedings of the National Academy of Sciences of the United
States of America 112*, 11696-11701 doi: 10.1073/pnas.1500624112.

Tasegian, A., Singh, F., Ganley, I.G., Reith, A.D., and Alessi, D.R. (2021). Impact of Type II LRRK2
inhibitors on signaling and mitophagy. *The Biochemical journal 478*, 3555-3573 doi:
10.1042/BCJ20210375.

Taylor, S.S., Kaila-Sharma, P., Weng, J.H., Aoto, P., Schmidt, S.H., Knapp, S., Mathea, S., and Herberg,

F.W. (2020). Kinase Domain Is a Dynamic Hub for Driving LRRK2 Allostery. *Frontiers in molecular*

neuroscience 13, 538219 doi: 10.3389/fnmol.2020.538219.

Teo, G., Liu, G., Zhang, J., Nesvizhskii, A.I., Gingras, A.C., and Choi, H. (2014). SAINTexpress:

improvements and additional features in Significance Analysis of INTeractome software. *J*

Proteomics 100, 37-43 doi: 10.1016/j.jprot.2013.10.023.

Tollenaere, M.A.X., Villumsen, B.H., Blasius, M., Nielsen, J.C., Wagner, S.A., Bartek, J., Beli, P., Mailand,

N., and Bekker-Jensen, S. (2015). p38- and MK2-dependent signalling promotes stress-induced

centriolar satellite remodelling via 14-3-3-dependent sequestration of CEP131/AZI1. *Nature*

communications 6, 10075 doi: 10.1038/ncomms10075.

Treier, M., Staszewski, L.M., and Bohmann, D. (1994). Ubiquitin-dependent c-Jun degradation in vivo

is mediated by the delta domain. *Cell 78*, 787-798 doi: 10.1016/s0092-8674(94)90502-9.

Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and Cox, J. (2016). The

Perseus computational platform for comprehensive analysis of (prote)omics data. *Nature methods*

13, 731-740 doi: 10.1038/nmeth.3901.

Vides, E.G., Adhikari, A., Chiang, C.Y., Lis, P., Purlyte, E., Limouse, C., Shumate, J.L., Spinola-Lasso, E.,

Dhekne, H.S., Alessi, D.R., and Pfeffer, S.R. (2022). A feed-forward pathway drives LRRK2 kinase

membrane recruitment and activation. *Elife 11* doi: 10.7554/eLife.79771.

Waschbusch, D., Michels, H., Strassheim, S., Ossendorf, E., Kessler, D., Gloeckner, C.J., and Barnekow,

A. (2014). LRRK2 transport is regulated by its novel interacting partner Rab32. *PloS one 9*, e111632

doi: 10.1371/journal.pone.0111632.

Wauters, L., Versees, W., and Kortholt, A. (2019). Roco Proteins: GTPases with a Baroque Structure and

Mechanism. *Int J Mol Sci 20* doi: 10.3390/ijms20010147.

Wilhelm, L.P., Zapata-Munoz, J., Villarejo-Zori, B., Pellegrin, S., Freire, C.M., Toye, A.M., Boya, P., and

Ganley, I.G. (2022). BNIP3L/NIX regulates both mitophagy and pexophagy. *The EMBO journal 41*,

e111115 doi: 10.15252/embj.2022111115.

Xue, J., Zhang, J., Zhang, J., Liu, J., Wang, F., Li, K., and Liu, C. (2023). The Parkinson's disease-associated

mutation LRRK2 G2385R alters mitochondrial biogenesis via the PGC-1alpha-TFAM pathway.

Mitochondrion 73, 10-18 doi: 10.1016/j.mito.2023.09.002.

Yang, Y., Ran, J., Liu, M., Li, D., Li, Y., Shi, X., Meng, D., Pan, J., Ou, G., Aneja, R., Sun, S.C., and Zhou, J.

(2014). CYLD mediates ciliogenesis in multiple organs by deubiquitinating Cep70 and inactivating

HDAC6. *Cell Res 24*, 1342-1353 doi: 10.1038/cr.2014.136.

Yu, M., Arshad, M., Wang, W., Zhao, D., Xu, L., and Zhou, L. (2018). LRRK2 mediated Rab8a

phosphorylation promotes lipid storage. *Lipids Health Dis 17*, 34 doi: 10.1186/s12944-018-0684-x.

Zhao, Y., Bracher-Smith, M., Li, Y., Harvey, K., Escott-Price, V., Lewis, P.A., and Manzoni, C. (2024).
Transcriptomics and weighted protein network analyses of the LRRK2 protein interactome reveal
distinct molecular signatures for sporadic and LRRK2 Parkinson's Disease. *NPJ Parkinsons Dis 10*,
144 doi: 10.1038/s41531-024-00761-8.

Zhao, Y., Vavouraki, N., Lovering, R.C., Escott-Price, V., Harvey, K., Lewis, P.A., and Manzoni, C. (2023).
Tissue specific LRRK2 interactomes reveal a distinct striatal functional unit. *PLoS Comput Biol 19*,
e1010847 doi: 10.1371/journal.pcbi.1010847.

Zheng, L., Bernard-Marissal, N., Moullan, N., D'Amico, D., Auwerx, J., Moore, D.J., Knott, G., Aebischer,
P., and Schneider, B.L. (2017). Parkin functionally interacts with PGC-1α to preserve mitochondria
and protect dopaminergic neurons. *Human molecular genetics 26*, 582-598 doi:
10.1093/hmg/ddw418.

Zhu, H., Hixson, P., Ma, W., and Sun, J. (2024). Pharmacology of LRRK2 with type I and II kinase
inhibitors revealed by cryo-EM. *Cell Discov 10*, 10 doi: 10.1038/s41421-023-00639-8.

Zhu, H., Tonelli, F., Turk, M., Prescott, A., Alessi, D.R., and Sun, J. (2023). Rab29-dependent
asymmetrical activation of leucine-rich repeat kinase 2. *Science 382*, 1404-1411 doi:
10.1126/science.adi9926.

Zimprich, A., Biskup, S., Leitner, P., Lichtner, P., Farrer, M., Lincoln, S., Kachergus, J., Hulihan, M., Uitti,
R.J., Calne, D.B., Stoessl, A.J., Pfeiffer, R.F., Patenge, N., Carbajal, I.C., Vieregge, P., Asmus, F., Muller-
Myhsok, B., Dickson, D.W., Meitinger, T., Strom, T.M., Wszolek, Z.K., and Gasser, T. (2004).
Mutations in LRRK2 cause autosomal-dominant parkinsonism with pleomorphic pathology. *Neuron*
44, 601-607 doi: 10.1016/j.neuron.2004.11.005.

Tables:

Accession	Gene name	Protein name	Jaccard score
Q5S007	LRRK2	Leucine-rich repeat serine/threonine-protein kinase 2 (bait)	1.000000
Q9NQC7	**CYLD**	**Ubiquitin carboxyl-terminal hydrolase CYLD**	**0.848485**
Q6P2H3	CEP85	Centrosomal protein of 85 kDa	0.846626
Q96JH7	VCPIP1	Deubiquitinating protein VCPIP1	0.834286
Q96L93	KIF16B	Kinesin-like protein KIF16B	0.832335
P78312	FAM193A	Protein FAM193A	0.817647
Q8NEY1	NAV1	Neuron navigator 1	0.817073
Q7Z2T5	TRMT1L	TRMT1-like protein	0.815951
Q5SZL2	CEP85L	Centrosomal protein of 85 kDa-like	0.810127
O95684	CEP43	Centrosomal protein 43	0.809816
Q96RS0	TGS1	Trimethylguanosine synthase	0.798742

Table 1: 10 IDs with the highest Jaccard score underlying identified "Cluster 2". The DUB CYLD (bold) shows highest co-evolution to LRRK2.

1030

Accession	Protein name	Gene name	p-value	Log2 T-test difference
P81605	Dermcidin	DCD	2.91844	3.3
Q9UPN4	Centrosomal protein of 131 kDa	CEP131	4.15494	2.2
Q49A88	Coiled-coil domain-containing protein 14	CCDC14	5.39373	2.1
Q9Y2D8	Afadin- and alpha-actinin-binding protein	SSX2IP	2.98866	2.1
O75665	Oral-facial-digital syndrome 1 protein	OFD1	7.92350	2.0
Q15154	Pericentriolar material 1 protein	PCM1	6.62987	1.8
Q9UHB6	LIM domain and actin-binding protein 1	LIMA1	2.96451	1.7
Q8WXW3	Progesterone-induced-blocking factor 1	PIBF1	4.62801	1.5
Q6WCQ1	Myosin phosphatase Rho-interacting protein	MPRIP	7.40967	1.5
Q14596	Next to BRCA1 gene 1 protein	NBR1	3.34347	1.3
Q5TB80	Centrosomal protein of 162 kDa	CEP162	3.46643	1.3
Q8WWI1	LIM domain only protein 7	LMO7	5.84664	1.3

1031

1032 **Table 2:** Enriched proteins in the LRRK2 proximity proteome upon MLi-2 treatment (FDR-controlled T-
1033 test).

1034

Accession	Protein name	Gene name	p-value (-log10)	Log2 T-test difference
O14966	Ras-related protein Rab-7L1	RAB29	9,72545	10,5
Q8IZA0	Dyslexia-associated protein KIAA0319-like protein	KIAA0319L	5,00209	4,9
Q6WKZ4	Rab11 family-interacting protein 1	RAB11FIP1	4,56815	4,6
O15126	Secretory carrier-associated membrane protein 1	SCAMP1	4,09385	3,5
Q9UIQ6	Leucyl-cystinyl aminopeptidase;Leucyl-cystinyl aminopeptidase, pregnancy serum form	LNPEP	4,33552	3,4
Q9H2J7	Sodium-dependent neutral amino acid transporter B(0)AT2	SLC6A15	5,06972	2,9
Q96QD8	Sodium-coupled neutral amino acid transporter 2	SLC38A2	2,89271	2,9
Q13586	Stromal interaction molecule 1	STIM1	3,12115	2,8
P98194	Calcium-transporting ATPase type 2C member 1	ATP2C1	5,00347	2,6
Q9H2H9	Sodium-coupled neutral amino acid transporter 1	SLC38A1	2,37723	2,6
O75976	Carboxypeptidase D	CPD	3,20952	2,6
Q14126	Desmoglein-2	DSG2	4,31421	2,5
P51809	Vesicle-associated membrane protein 7	VAMP7	4,02209	2,5
Q8NDI1	EH domain-binding protein 1	EHBP1	2,37346	2,5
Q8TBA6	Golgin subfamily A member 5	GOLGA5	3,0259	2,4
Q9BXF6	Rab11 family-interacting protein 5	RAB11FIP5	2,9101	2,4
Q7L804	Rab11 family-interacting protein 2	RAB11FIP2	3,33154	2,2
Q6IAA8	Ragulator complex protein LAMTOR1	LAMTOR1	3,23604	2,1
P61006	Ras-related protein Rab-8A	RAB8A	3,56486	2,1
P11717	Cation-independent mannose-6-phosphate receptor	IGF2R	5,71543	2,0
Q13190	Syntaxin-5	STX5	3,88884	2,0
Q15836;P63027	Vesicle-associated membrane protein 3;Vesicle-associated membrane protein 2	VAMP3;VAMP2	3,67019	2,0
Q92692	Nectin-2	PVRL2	3,06334	2,0
Q6NXT4	Zinc transporter 6	SLC30A6	3,47208	1,8
Q96A65	Exocyst complex component 4	EXOC4	2,51947	1,7
Q14141	Septin-6	Sept6	3,87782	1,7
Q04656	Copper-transporting ATPase 1	ATP7A	2,77415	1,7
Q7Z2W4	Zinc finger CCCH-type antiviral protein 1	ZC3HAV1	2,62567	1,6
P78310	Coxsackievirus and adenovirus receptor	CXADR	2,87956	1,4
P08195	4F2 cell-surface antigen heavy chain	SLC3A2	2,43497	1,4
P02786	Transferrin receptor protein 1;Transferrin receptor protein 1, serum form	TFRC	3,21618	1,3
Q07960	Rho GTPase-activating protein 1	ARHGAP1	2,65929	1,3
O60271	C-Jun-amino-terminal kinase-interacting protein 4	SPAG9	3,80826	1,3
P42167	Lamina-associated polypeptide 2, isoforms beta/gamma;Thymopoietin;Thymopentin	TMPO	3,54449	1,2
Q9BSJ8	Extended synaptotagmin-1	ESYT1	3,59064	1,0
Q14739	Lamin-B receptor	LBR	3,12029	0,8

1035

1036 **Table 3:** Enriched proteins in the RAB29-depended LRRK2 proximity proteome (FDR-controlled T-test).

1037

1038 **Figures**

1039

1040 **Figure 1:** (A) Workflow of the BioID analysis and post processing of the data. (B) Overlap between the

1041 different BioID approaches. Enriched proteins were determined by relative label-free quantification

1042 (combined with an FDR-controlled T-test, p=0.05) of BioID LRRK2 vs the corresponding tag-only

1043 control: BioID1-LRRK2/BioID1: N=8, BioID2-LRRK2/BioID2: N=9, miniTurbo-LRRK2/miniTurbo: N=8.

1044

Figure 2: LRRK2 interactome functional enrichment. A) the LRRK2 BioID interaction network was mapped on known PPIs from IntAct via Cytoscape. The edges' color and thickness for mapped interactions are proportional to IntAct MI score, else they are colored in gray with constant thickness. Modules within the network have been identified through the Glay method of the Autoannotate app and nodes colored accordingly. Labels have been calculated using the 'Adjacent Words' of the WordCloud algorithm, considering the significantly enriched (FDR < 0.01) GO Biological processes and cellular compartments in which each given gene of the network participated; B) Venn diagram showing the overlap between the BioID interaction sets and known LRRK2 interactions in IntAct; barplots

1053   showing the top 15 **most** significant terms obtained through String's functional enrichment of C)

1054   Compartments, D) GO Biological Process and D) Interpro domains.

1055

Figure 3: Co-evolution analysis of LRRK2 interactome. (A) Distribution of co-evolution metric (i.e. Jaccard score) among the different LRRK2 interaction sets, including IntAct interaction types and BIOID. P-values have been computed with the Wilcoxon rank-sum, two-sided test with Bonferroni correction (* $P < 0.05$, ** $P < 0.001$, *** $P < 0.001$, **** $P < 0.0001$). Violin plots show the median as the white central dot and first and third quartiles as bounds of the thicker black line (box); the whiskers extend to the last data point within 1.5 times the interquartile range (IQR) from the box's boundaries ; (B) predictiveness of LRRK2 **interaction type** via co-evolution (C) Heatmap showing the hierarchical clustering based on the degree of co-evolution (Jaccard score) of the LRRK2 proximity proteome; (D) String network of the proteins most co-evolved with LRRK2 participating to cluster 2. Nodes are colored in gray or light-gray according to modules membership determined with the MCL approach available from the Cytoscape clusterMaker app. Node rims are colored according to enriched processes; (E) Dot plot showing top 15 significantly enriched (FDR < 0.01) categories from the string network of cluster 2. Values on the x axis display the ratio of genes of the term participating to the network over the total number of genes in that category. The ratio of the dot is proportional to the number of genes of that category in the network, while the node color is the

1071   darker the more significant is the category enrichment. Only categories with at least 5 genes have been

1072   displayed. Representative, enriched categories have been colored with the same color scheme as in

1073   panel (D); (F) Effect of CYLD co-expression on LRRK2 stability in HEK293T. Cells were transfected with

1074   LRRK2 Y1699C and either with WT-ubiquitin (WT-Ub) or Ubiquitin-K63 (Ub-63) which can only be

1075   Ubiquitin-conjugated at lysine 63. In addition, CYLD or a vector control (pcDNA3.0) were co-expressed.

Figure 4: 3D complexes prediction and analysis of LRRK2 interactome (A) workflow of the structural

bioinformatics analysis of LRRK2 interactome; (B) on the left the heatmap showing the hierarchical

clustering based on pairwise RMSD calculation of LRRK2's chain in the different complexes. On the

right two representative conformations of LRRK2 in the "locked" and "unlocked" conformation to mark

the two main clusters; (C) cartoon representation of "locked" (tomato) and "unlocked" (light green)

superimposed LRRK2 conformations; (D) violin plot of the distribution of RMSD calculated with respect

to the reference LRRK2 experimental structure (PDB 7LHW) for the locked and unlocked clusters. Color

scheme as in (C); (E) distribution of the Jaccard score for the locked and unlocked clusters. Color

scheme as in (C). P-values for violin plots in D and E have been computed with the Wicoxon

rank-sum, two-sided test (* $P < 0.05$, ** $P < 0.001$, *** $P < 0.001$, **** $P < 0.0001$). Violin plots

show the median as the white central dot and first and third quartiles as bounds of the thicker

black line (box); the whiskers extend to the last data point within 1.5 times the interquartile

range (IQR) from the box's boundaries; (F) Venn diagram of the enriched terms in the locked and

unlocked clusters. Color scheme as in (C); (E) Top 10 exclusively enriched (FDR<0.01) terms in the

locked and unlocked clusters.

Figure 5: Interface analysis of LRRK2's 3D interactome (A) interface fingerprint of the LRRK2 complexes via distogram calculation. Cells contain the minimum expected distance obtained from the distograms of any residue of each interactor (row) to every residue in LRRK2 (column). Rows are color annotated based on confidence score, Jaccard and RMSD from reference LRRK2 structure (PDB: 7LHW); (B) representative structure of the Core cluster (i.e. LRRK2-ANAPC5 complex); (C) representative structures of the WD40 cluster; (D) representative structures of the ARM binding mode cluster (i.e.

LRRK2- DCUN1D4 complex); distribution among different interface cluster of the (E) confidence score,

(F) RMSD and (G) Jaccard score descriptors. P-values have been computed with the Wicoxon rank-

sum, two-sided test with Bonferroni correction (* $P < 0.05$, ** $P < 0.001$, *** $P < 0.001$, ****

$P < 0.0001$). Violin plots show the median as the white central dot and first and third quartiles

as bounds of the thicker black line (box); the whiskers extend to the last data point within 1.5

times the interquartile range (IQR) from the box's boundaries.

1105

Figure 6: Different effects of type I and type II inhibitors and RAB29 on the LRRK2 proximity proteome.
(A) MLi-2 vs DMSO. (B) MLi-2 vs GZD-824; (C) Stringified network of the MLi-2 interactome from (A) mapped to IntAct known interactions. Edges' color and thickness for mapped interactions are proportional to IntAct MI score, else they are colored in gray with constant thickness. Node rims are colored according to enriched processes; (D) Dot plot showing top 15 significantly enriched (FDR < 0.01) categories from the LRRK2-MLi-2 interactome. Values on the x axis display the ratio of genes of the term participating to the network over the total number of genes in that category. The ratio of the dot is proportional to the number of genes of that category in the network, while the node color is the darker the more significant is the category enrichment. Only categories with at least 5 genes have been displayed. Representative, enriched categories have been colored with the same color scheme as in panel C. (E) Scheme highlighting the function of centriolar satellites and the core protein PCM1 in ciliation ((Hori and Toda, 2017), modified) (F) Violin plot reporting the Jaccard Score distribution in different data sets. P-values have been computed with the Wicoxon rank-sum, two-sided test with Bonferroni correction (* *P* < 0.05). Violin plots show the median as the white central dot and first and third quartiles as bounds of the thicker black line (box); the whiskers extend to

the last data point within 1.5 times the interquartile range (IQR) from the box's boundaries;

(G) mt-LRRK2 proximity proteome in dependence of RAB29 co-expression: RAB29 redirects LRRK2 to

lysosomal PPI networks; (H) Stringified network of the RAB29 over-expression interactome from (G)

mapped to IntAct known interactions. Edges' color and thickness for mapped interactions are

proportional to IntAct MI score, else they are colored in gray with constant thickness. Node rims are

colored according to enriched processes; (I) Dot plot showing top 15 significantly enriched (FDR < 0.01)

categories from the LRRK2-RAB29 over-expression interactome. Values on the x axis display the ratio

of genes of the term participating to the network over the total number of genes in that category. The

ratio of the dot is proportional to the number of genes of that category in the network, while the node

color is the darker the more significant is the category enrichment. Only categories with at least 5

genes have been displayed. Representative, enriched categories have been colored with the same

color scheme as in panel H.

1133

Figure 7: AF-structural modeling of MLi-2 and Rab29 interactomes. (A) Hierarchical clustering of pairwise RMSD of LRRK2 chain in the different complexes; B) LRRK2-RAB8A predicted complex; (C) LRRK2-RAB29 predicted complex; (D) violin plot of the distribution of RMSD calculated with respect to the reference LRRK2 experimental structure (PDB 7LHW) for the locked and unlocked clusters. Color scheme as in 4C; (E) distribution of the Jaccard score for the locked and unlocked clusters. Color scheme as in 4C. P-values in violin plots in (D) and (E) have been computed with the Wicoxon rank-sum, two-sided test (* $P < 0.05$, ** $P < 0.001$, *** $P < 0.001$, **** $P < 0.0001$). Violin plots show the median as the white central dot and first and third quartiles as bounds of the thicker black line (box); the whiskers extend to the last data point within 1.5 times the interquartile range (IQR) from the box's boundaries; (F) Venn diagram of the enriched terms in the locked and

unlocked clusters. Color scheme as in 4C; (G) Top exclusively enriched (FDR<0.01) terms in the locked

and unlocked clusters. Color scheme as in 4C.

Supplemental Information

Supplemental tables (2 separate files):

Supplemental Table 1: LRRK2 proximity proteomes obtained with BioID1, BioID2 and miniTurbo as

well as the Rab29-dependent LRRK2-mT and the inhibitor-dependent LRRK2-mT proximity proteomes.

Supplemental Table 2: co-evolutionary and structural features of BioID interactors

Supplemental Table 3: over-representation analysis of protein clusters defined via co-evolution

Supplemental Table 4: over-representation analysis of protein clusters defined via RMSD clustering of

AF-multimer predicted complexes

Supplemental Table 5: over-representation analysis of protein clusters defined via Interaction

fingerprint clustering of AF-multimer predicted complexes

Supplemental Table 6: over-representation analysis of protein clusters defined via RMSD clustering of

AF-multimer predicted complexes of the RAB29-overexpression and MLi-2 interactomes

1158

Supplemental Figure 1: Western blot, pRab10, pS935 and biotinylation – comparison BioID1, BioID2, miniTurbo. To ensure functional BioID-fusion proteins, LRRK2 phosphorylation at S935 and Rab10 phosphorylation was assessed. (A) The different BioID LRRK2 constructs show sufficient biotinylation of proteins as well as MLi-2-responsive Rab10 phosphorylation. (B) All constructs are phosphorylated at S935 (top panel). Replication of LRRK2-mediated Rab10 phosphorylation (last panel) demonstrating functional BioID fusion proteins.

1165

Supplemental Figure 2: ROC Curve of prediction with Mutual Information (MI) of LRRK2's IntAct

interaction types.

1168

1169 **Supplemental Figure 3:** heatmap with cluster specific, significantly enriched processes (FDR < 0.01;

1170 StringDB) obtained from coevolution (Jaccard)-based clustering of the LRRK2 BioID interactome.

1171

Supplemental Figure 4: heatmap displaying the coevolution (Jaccard)-based hierarchical clustering of the LRRK2's IntAct interactome. Colored bars provide IntAct's interaction type annotation.

1174

1175

1176 **Supplemental Figure 5:** Western blot demonstrating sufficient LRRK2 inhibition for MLi-2 and GZD-824

1177 in a HEK293 line stably expressing miniTurbo LRRK2. Both inhibitors show a sufficient inhibition of

1178 Rab10 phosphorylation at 1µM. While the type I inhibitor shows the characteristic loss of LRRK2 pS935

1179 phosphorylation, the type II inhibitor GZD-824 does not alter pS935 phosphorylation (Tasegian et al.,

1180 2021).

1181

Supplemental Figure 6: (A) interaction fingerprint clustering of LRRK2's modulators (MLi-2 and RAB29)

interactomes. Cells contain the minimum expected distance obtained from the distograms of any

residue of each interactor (row) to every residue in LRRK2 (column). Rows are color annotated based

on confidence score, Jaccard, RMSD from reference LRRK2 structure (PDB: 7LHW) and experiment type

(either MLi-2 or RAB29 over-expression); (B) interaction fingerprint of LRRK2 and interacting RABs

(from IntAct).