# Genomic stability in the Galápagos *Scalesia* adaptive radiation: Consistent transposable element accumulation despite hybridization and ecological niche shifts

José Cerca[1,2], Patricia Jaramillo Díaz[3,4,5], Clément Goubert[6], Heidi Yang[7], Vanessa C. Bieker[1], Mario Fernández-Mazuecos[8,9], Pablo Vargas[10], Rowan Schley[11], Siyu Li[12], Juan Ernesto Guevara-Andino[13], Bent Petersen[14,15], Gitte Petersen[16], Neelima R. Sinha[12], Lene R. Nielsen[17], James H. Leebens-Mack[18], Gonzalo Rivas-Torres[19], Loren H. Rieseberg[20], Michael D. Martin[1]

1 - Department of Natural History, University Museum, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

2 - Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo, Oslo, Norway

3 . Estación Científica Charles Darwin, Fundación Charles Darwin, Santa Cruz, Galápagos, Ecuador.

4. Department of Botany and Plant Physiology, University of Málaga, Málaga, Spain.

5. IUCN SSC Galapagos Plant Specialist Group, Puerto Ayora, Galapagos 200102, Ecuador.

6. McGill Genome Centre, McGill University, Montreal H3A 0G1, QC, Canada

7. Department of Ecology & Evolutionary Biology, University of California, Los Angeles, Los Angeles, California, United States of America

8. Departamento de Biología (Botánica), Facultad de Ciencias, Universidad Autónoma de Madrid, Calle Darwin 2, 28049 Madrid, Spain

9. Centro de Investigación en Biodiversidad y Cambio Global, Universidad Autónoma de Madrid (CIBC-UAM), Calle Darwin 2, 28049 Madrid, Spain

10. Biodiversity and Conservation, Real Jardín Botánico, 28014 Madrid, Spain

11. Department of Geography, University of Exeter, Laver Building, North Park Road, Exeter, Devon, UK

12. Department of Plant Biology, University of California, Davis, Davis, CA, 95616, USA

13. Grupo de Investigación en Ecología y Evolución en los Trópicos-EETrop- Universidad de las Américas, Quito-Ecuador

14. Center for Evolutionary Hologenomics, Globe Institute, University of Copenhagen, DK-1353, Copenhagen, Denmark

15. Centre of Excellence for Omics-Driven Computational Biodiscovery (COMBio), Faculty of Applied Sciences, AIMST University, Kedah, Malaysia

16. Department of Ecology, Environment and Plant Sciences, Stockholm University, 106 91 Stockholm,

Sweden

17. Department of Geosciences and Natural Resource Management, University of Copenhagen,

Rolighedsvej 23, 1958 Frederiksberg C, Denmark

18. Department of Plant Biology, University of Georgia, Athens, GA, 30602, USA

19. Colegio de Ciencias Biológicas y Ambientales, Galapagos Science Center, Universidad San Francisco de

Quito USFQ, Quito 170901, Ecuador

20. Department of Botany and Biodiversity Research Centre, University of British Columbia, Vancouver,

BC, V6T 1Z4, Canada

**Abstract**

Transposable elements (TEs) have been hypothesized to play a pivotal role in driving diversification by

facilitating the emergence of novel phenotypes and the accumulation of divergence between species. The

rapid speciation and ecological diversification characteristic of adaptive radiations offer a unique

opportunity to examine the link between TE accumulation and speciation, diversification and adaptation.

Here, focusing on all 15 species of the genus *Scalesia* (Asteraceae), a radiation endemic to the Galápagos

Islands, we test whether hybridization or shifts in ecological niche are associated with changes in TE

accumulation in genomes. Our analyses reveal little to no variation in TE accumulation among *Scalesia*

species nor its hybrid populations. Shifts in ecological niches, linked to climatic variation, did not result in

discernible changes in TE accumulation, a surprising finding given the anticipated selective pressure

imposed by aridity, a factor often linked to genome size reduction. We found no distinct patterns in the

temporal accumulation of TEs, with the exception of a taxon (var. *atractyloides*) of *Scalesia atractyloides*.

Our findings challenge the assertion that TEs have directly driven diversification, speciation and local

adaptation. Instead, TEs may simply be 'along for the ride,' rather than actively contributing to plant

diversification.

## Introduction

Transposable elements are DNA sequences capable of relocating within the genome (McClintock 1984). Through their mobility and ability to replicate, TEs are able to generate a diverse range of mutations encompassing changes in promoter gene sequences, frameshifts, intronic modifications and gene duplications within their hosts (Choi and Lee 2020; Kidwell and Lisch 1997; Bennetzen and Wang 2014; Bourque et al. 2018; Pimpinelli and Piacentini 2020; Oliver, McComb, and Greene 2013; Almeida et al. 2022). Some of these mutations have been linked to the evolution of novel phenotypes. For instance, the evolution of the melanic phenotype of the peppered moth (*Biston betularia*) showcases the potential functional consequences of TE insertions as the insertion of the 'carb-TE' in an intronic region of the *cortex* gene alters its expression, likely contributing to the melanic phenotype (Van't Hof et al. 2016).

Advances in genomic sequencing technology and processing algorithms have greatly facilitated the detection and characterization of TEs, particularly in non-model organisms with limited genomic resources (Clement Goubert et al. 2022). The evident adaptive potential of TEs in shaping phenotypes (Kratochwil et al. 2022; Conart et al. 2022; Bennetzen and Wang 2014), along with the evidence of substantial variation in TE diversity and accumulation across different reference genomes (e.g. (Cerca et al. 2021; Schley et al. 2022), has intensified interest in a potential relationship between TEs and diversification. One particularly intriguing hypothesis is that lineages with high speciation rates also exhibit elevated rates of TE accumulation on their genomes (Ricci et al. 2018; Pimpinelli and Piacentini 2020; Oliver and Greene 2009, 2012; Belyayev 2014; Ray et al. 2008; Falcon, Tanaka, and Rodriguez-Terrones 2023; Pace and Feschotte 2007; de Boer et al. 2007). However, a key limitation of these studies has been their reliance on a small number of genomes, often with limited taxonomic coverage while encompassing large clades, and primarily based on correlations. Consequently, there may be an overestimation of the role of TEs as major drivers of speciation and ecological diversification at macroevolutionary scales.

Adaptive radiations represent evolutionary experiments where extensive ecological variation aligns with rapid speciation, and they have been a cornerstone of our understanding of evolutionary change (Cerca et al. 2023; Gillespie et al. 2020; Schluter 2000). Adaptive radiations can also yield valuable insights into the interplay between TEs and evolutionary success; however, the evidence available is conflicting and largely originated from cichlid radiations. For example, a study examining four species within an African cichlid adaptive radiation found higher TE counts in the genomes and transcriptomes of the radiating species compared to non-radiating lineages (Brawand et al. 2014). More recent evidence, however, suggested a lack of correlation between species richness at the tribe level in the Lake Tanganyika adaptive radiation and TE content across 245 genomes (Ronco et al. 2021). Similarly, cichlid lineages in the Americas had no

discernible differences in TE content when contrasting a radiating lineage (*Amphilophus citrinellus*) with a

non-radiating lineage (*Archocentrus centrarchus*; (Xiong et al. 2021). Notably, a golden phenotype in the

cichlid genus *Amphilophus* is associated with a transposon-derived inverted repeat in the *goldentouch* gene

(Kratochwil et al. 2022). This conflicting body of evidence underscores the necessity for further exploration

of TEs as facilitators of diversifications, particularly extending beyond the scope of cichlid radiations.

The *Scalesia* radiation (Asteraceae, Asterales, Magnoliidae), endemic to the Galápagos archipelago,

presents an excellent opportunity to investigate the potential relationship between TEs and diversification.

This genus comprises ~15 species that evolved from a single common ancestor which diversified about 1

million years ago (Fernández-Mazuecos et al. 2020). The recent origin of this radiation may facilitate the

reconstruction and tracking of TE accumulation over time. *Scalesia* exhibits remarkable ecological diversity,

particularly in its occupation of various climate zones (*hereafter* referred to as climate niches) (Blaschke and

Sanders 2009). The genus inhabits a wide range of ecological niches and soil types (Figure 1), from coastal or

littoral zones characterized by saltwater habitats, aridity, low precipitation, elevated temperatures, and

intense solar radiation, to more humid zones typically found at higher elevations, which experience

increased rainfall throughout the year, lower temperatures, and frequent cloud cover. This ecological

variation provides an opportunity to explore whether climatic niches exert selective pressures that influence

TE dynamics, as observed in plants under arid conditions (Schley et al. 2022). Specifically, the accumulation

of TEs can result in an increase in genome size, which is associated with larger stomatal cells with impaired

gas exchange efficiency and poor water economy (Franks and Beerling 2009). Thus, natural selection may

play a role in regulating genome size in species adapted to arid conditions, such as many *Scalesia* species.

Furthermore, hybrid populations exist in the radiation (Fernández-Mazuecos et al. 2020), allowing testing

whether hybridization triggers genome deregulation by inducing TE activity (L. Wang et al. 2023; Petrov et

al. 1995; Serrato-Capuchina et al. 2020; Rubin, Kidwell, and Bingham 1982; Jiang et al. 2003; Parisod et al.

2010). Finally, the reference genome assembly for the group, *Scalesia atractyloides*, is composed of over 80%

of repetitive content and TEs – making this lineage ideal to understand TE accumulation dynamics (Cerca

et al. 2022).

Building on the premise that TEs play a role in diversification dynamics, here we test three

hypotheses: *(H1)* Lineages undergoing rapid speciation and broadening of ecological variation exhibit an

increased rate of lineage- or taxon-specific TE accumulation/expansion on their genomes; *(H2)* Shifts in

climatic niches result in a change of selective regimes, with lineages in arid areas experiencing stronger

selection for TE removal as a result of selection for smaller cells; and *(H3)*. Hybrids have an increased

accumulation of TEs as a result of genome deregulation. To test these hypotheses, we employed whole-

genome resequencing of all 15 *Scalesia* species, four *Scalesia* hybrid populations and outgroup species, and

contrasted genomic results with ecological data (Blaschke and Sanders 2009; Perez et al. 2023).

# Materials and Methods

## Study species and data collection

*Scalesia* (Asteraceae) is an endemic genus to the Galápagos Islands consisting of approximately 15

species derived from a single common ancestor. The divergence of *Scalesia* from members of a continental

sister lineage, *Pappobolus*, occurred around 3 million years ago, while most diversification of the group

occurred within the last million years (Fernández-Mazuecos et al. 2020). *Scalesia* occupies a wide array of

ecological niches and soil types (Figure 1) (Blaschke and Sanders 2009), displaying clear environment-

phenotype associations (Perez et al. 2023; Fernández-Mazuecos et al. 2020), as expected in an adaptive

radiation (Schluter 2000). In this study, we sequenced samples representing the entire *Scalesia* genus,

including: *S. affinis, S. aspera, S. atractyloides* (var. *atractyloides* and *darwinii*), *S. baurii* (subsp. *baurii* and

*hopkinsii*), *S. cordata, S. crockeri, S. divisa, S. gordilloi, S. helleri, S. incisa, S. microcephala, S. pedunculata, S.*

*retroflexa, S. stewartii,* and *S. villosa* (Supplementary Table 01). Additionally, we carried out sequencing of

samples from populations that show evidence of extensive hybridization (Bieker, *in prep*). The hybridizing

pairs include only pairs of sister species: *S. retroflexa × helleri, S. crockerii × aspera, S. incisa × divisa,* and *S.*

*stewartii × atractyloides* (Supplementary Table 01). We incorporated individuals from the genus *Pappobolus*

from the South American continent as the outgroup (sister group) in some analyses (Blaschke and Sanders

2009; Fernández-Mazuecos et al. 2020), including *P. ecuadoriensis, P. hypargyreus, P. juncosae, P. lehmanii,*

and *P. nigrescens*. Detailed information and accession numbers regarding the sampling of outgroup species

can be found in Fernández-Mazuecos et al. (2020).

## Data generation (library preparation and sequencing)

We extracted genomic DNA using the DNeasy 96 Plant Kit (Qiagen) following the manufacturer's

protocol with minor modifications. These modifications consisted of grinding dried leaf tissues in a Qiagen

collection microtube with a 3-mm tungsten carbide bead and repeating the membrane washing step with an

additional 800 µl of buffer AW2 to remove residual greenish color from the samples. We eluted the DNA

from the membranes using 50 µl of buffer AE to obtain a high-concentration DNA yield. The DNA

extracts were then sent to the commercial provider Novogene UK for shearing to a mean insert size of 350

bp, double-stranded DNA genomic library preparation, indexing and amplification via PCR, and 150-bp

paired-end sequencing on the Illumina NovaSeq 6000 platform. The average sequencing depth was ~6x

(low-coverage). Three samples used in this study were obtained from herbarium specimens and were

prepared for genomic sequencing independently. Leaf tissue was disrupted using a Tissue Lyser II

(Quiagen), followed by DNA extraction using the DNeasy Plant Mini Kit following the manufacturer's

protocol except for an additional   overnight incubation step with 20 µL of proteinase K after cell lysis as

described in (Martin et al. 2014). DNA was eluted using 62 µL buffer AE. Extracts were then sent to the

commercial provider Novogene for double-stranded DNA genomic library preparation, indexing and

amplification via PCR, and 150-bp paired-end sequencing on the Illumina NovaSeq 6000 platform.  The

raw sequencing data generated for this study can be found at the European Nucleotide Archive under

accession code PRJEB70770.

## Low-coverage phylogenomics

We started by conducting a distance-based phylogenetic reconstruction, accounting for low-

coverage sequencing, to investigate the evolutionary history of the *Scalesia* genus and to establish a

backbone for subsequent analyses and interpretations. We performed an additional phylogenetic

reconstruction, including hybrid populations to confirm the phylogenetic placement of these hybrids. We

started by quality-filtering the raw Illumina data by identifying and removing adapters and low-quality

basepairs using *AdapterRemoval v2.3.2* (Schubert, Lindgreen, and Orlando 2016). Subsequently, we used

*Skmer v3.2.1*, a pipeline designed to estimate genetic distances between genomes through a *k*-mer-based

approach. *Skmer* offers a robust approach for phylogenetic reconstruction without requiring an assembly

and accounts for low-coverage of the samples (Marçais and Kingsford 2011; Ondov et al. 2016; Sarmashghi

et al. 2019; Rachtman et al. 2022). Running *Skmer* involved running the module 'skmer reference' to

process the adapter-trimmed sequencing data, followed by 'skmer subsample' to create 100 subsamples of

the libraries, and 'skmer correct' to correct the distance matrices based on these subsample replicates. To

construct a distance-based phylogenetic tree we employed *fastme v2.1.5* (Lefort, Desper, and Gascuel 2015).

## TE library construction

We downloaded the reference genome of the sunflower (*Helianthus annuus*) from its official

website (https://www.sunflowergenome.org/; version Ha412HOv2.0-20181130), and used it to generate a

sunflower-based TE and repeat library using *Repeat Modeler v2.0.2* (Flynn et al. 2020). Sunflower was

selected as the reference species because it has a high-quality reference genome, and it is equally evolutionary

divergent from *Scalesia* and *Pappobolus*. Building a TE and repeat library based on *Scalesia* alone could

introduce biases when conducting comparisons with the outgroup taxa (but see below). We concatenated

the sunflower-based library with Repbase (2017), and used the concatenated library as the base library for

subsequent TE detections using *dnaPipeTE*.

We used *dnaPipeTE* (Clément Goubert et al. 2015) to extract and characterize TEs and repeat

sequences. *dnaPipeTE* leverages the RNA-seq assembly pipeline Trinity to extract, quantify and assemble

sequences, including TEs and other repeats. While the primary objective of *dnaPipeTE* is to construct

complete TE and repeat sequences, it also detects and quantifies partially assembled TE and repeats (which

suffice for estimating the relative abundances on which we rely in this study). Upon the completion of the

assembly and collection process, *dnaPipeTE* categorises TEs and repeats into different classes based on their

homology with the TE and repeat library. In the established 'concatenated library', TE classes included

DNA, Helitron, Long interspersed nuclear element (LINE), Long terminal reads (LTR), rRNA, and Short

interspersed nuclear elements (SINE). Leveraging *RepeatMasker*, *dnaPipeTE* identifies low-complexity

sequences, simple repeats, and satellites, guided by a specific coverage threshold. *dnaPipeTE* has been

successfully applied to a diverse array of species, including arthropods, mollusks, and vertebrates (reviewed

in (Clément Goubert 2023)), even revealing substantial variation in TE and repeat content at the

population-level (e.g. (Lerat et al. 2019)). We ran *dnaPipeTE* on single-end forward reads for each

individual separately, downscaling the data to 0.1 coverage, specifying a genome size of 3.9 Gbp (Cerca et al.

2022), two Trinity assembly iterations, and a minimal query percentage of 0.2 (Clément Goubert et al.

2015).

## Exploration of patterns

We extracted three outputs from *dnaPipeTE*: 1) counts of TEs and repeats, which allowed a direct

comparison of the number of TEs and repeats accumulated across species; 2) repeat landscape plots from all

resequenced samples, which offered insights into the accumulation of TEs through time (to complement

this data, we also ran a repeat accumulation analysis in the chromosome-resolved *Scalesia atractyloides*

genome assembly (Cerca et al. 2022) using RepeatMasker; and 3) fasta files containing TE and repeat

sequences. Using the sequences in a fasta format, we used *OrthoFinder* to group TEs and repeats based on

sequence similarity (Emms and Kelly 2015, 2019), and we examined the number of TE/repeat orthogroups

shared between the outgroup species and *Scalesia* using *UpSet plots* and *Venn diagrams*. To test for

differences between groups, we performed a Mann–Whitney U test comparing hybrids *vs* non-hybrids, and

humid (*S. cordata, S. microcephala, S. pedunculata*) *vs* arid (*S. villosa, S. stewartii, S. atractyloides, S. crockeri,*

*S. aspera, S. retroflexa, S. helleri, S. gordilloi, S. incisa, S. divisa*).

We considered the possibility that the limited variation in TE and repeat content observed in some

of our results was due to the use of a sunflower-based library, which might be too divergent from *Scalesia*

and *Pappobolus*. To address this concern, we repeated all the aforementioned analyses using a *Scalesia*-

specific library instead of the sunflower library. To obtain the *Scalesia*-specific library, we ran

*RepeatModeler* on the *Scalesia atractyloides* genome (Cerca et al. 2022) and repeated all steps above (i.e.

concatenating the output from *RepeatModeler* with Repbase, running *DNApipeTE* analyses using the same

parameters, and running analyses involving the exploration of TE and repeats). Analyses of the *Scalesia*-

specific library found similar patterns to the main results (Supplementary Figure 01).

# Results

**Figure 01** - Phylogenetic relationships and transposable element (TE) variation in the *Scalesia* radiation

and the outgroup (genus *Pappobolus*). a) Neighbor-joining phylogeny of *Scalesia* and the outgroup. Hybrid

samples are shown below and were not included in the phylogenetic reconstruction. b) Stacked bar plot

illustrating TE and repeat groups (for the analysis displayed here we used the sunflower library; the analysis

using a *Scalesia*-based library is found in Supplementary Figure 01). Color-coding of the different repeat

and TE groups is found in the rightmost part of the figure, where 'na' denotes unidentified TEs. c)

Labelling of the different islands, environments, and soils occupied by different *Scalesia* species (for hybrids

we include only island labelling). Environment and soil labelling follow (Blaschke and Sanders 2009), and

black dots indicate uncertainty in environmental characterization.

The phylogenetic tree is characterized by mostly short internal branches in *Scalesia*, as expected in a

rapid radiation. *Scalesia villosa* is positioned as the sister species to all the remaining *Scalesia* species (Figure

01A; Supplementary Figure 02). *S. stewartii* and *S. atractyloides* form a clade which is sister to the remaining

12 species. *S. affinis*, *S. microcephala*, and *S. cordata* are grouped together as a clade being sister to a clade

including nine *Scalesia* species, namely *S. crockeri*, *S. aspera*, *S. retroflexa*, *S. helleri*, *S. baurii*, *S.*

*pedunculata*, *S. gordilloi*, *S. incisa,* and *S. divisa* (Figure 01A; Supplementary Figure 02). The phylogenetic

analysis including hybrid populations placed all hybrids as sister to one of the parental species

(Supplementary Figure 03).

We observed little variation in TE counts among different species, both in relative and absolute

terms (Figure 01B; Supplementary Table 02). The outgroup lineages displayed a higher count of TEs and

repeats, with an average of 323,234,351 bp (max 328,160,920 bp, min 318,480,284 bp, standard deviation

3,752,276 bp). *Scalesia* species (excluding hybrids) exhibited a lower TE and repeat content, with an average

of 266,935,585 bp (max 271,943,828 bp, min 259,698,199 bp, standard deviation 3,610,348 bp). Hybrid

*Scalesia* populations displayed a TE and repeat content averaging 269,790,871 bp (max 274,454,284 bp,

min 265,429,355 bp, standard deviation 4,796,678 bp; Figure 1B). Both Mann-Whitney U tests were non-

significant (humid *vs* arid $P$ = 0.1264; hybrid *vs* non-hybrid $P$ = 0.8972)

Long terminal repeat (LTR) retrotransposons were the largest class of TEs found in the data.

*Scalesia* species had an average of 129,559,033 bp of their genomes composed of LTRs (max 130,687,970

bp, min 106,894,927 bp, standard deviation 5,858,406 bp). The outgroup species had a mean of

172,863,512 bp of their genomes composed of LTRs (max 178,110,682 bp, min 166,411,171 bp, standard

deviation 4,342,796 bp), while hybrid *Scalesia* populations had 129,559,033 bp of LTRs (max 135,389,443

bp, min 122,925,507 bp, standard deviation 6,440,151 bp). In the second largest class of TEs, DNA

elements, *Scalesia* had an average of 13,256,247 bp composed of DNA elements (max 14,561,033 bp, min

11,743,596 bp, standard deviation 783,002 bp), while hybrid populations had 13,385,339 bp (max

14,377,105 bp, min 12,542,486 bp, standard deviation 853,847 bp). In contrast, the outgroup exhibited

fewer DNA elements, with an average of 7,703,936 bp (max 8,258,881 bp, min 7,308,486 bp, standard

deviation 348,650 bp).

**Repeat landscape plots**

**Figure 02** - Repeat/transposable element landscape plot. The *x-axis* shows the sequence divergence

(Kimura substitution level), an indirect measure of TE divergence/age, the *y-axis* shows the portion of the

genome covered by TEs and repeats. Assuming a constant mutation rate, higher Kimura substitution levels

indicate older TE ages. The outgroup (genus *Pappobolus*) is plotted in green, and all *Scalesia* species in

orange with the exception of *S. atractyloides* var. *atractyloides* in blue since it has a distinct repeat

accumulation pattern.

The repeat landscape plot shows that all *Scalesia* species exhibit a similar pattern of TE and repeat

accumulation, with the exception of *S. atractyloides* var. *atractyloides* (Figure 02; Supplementary Figure 04).

All outgroup species have a similar accumulation of TEs, which is distinct from that of *Scalesia*. Specifically,

there is a younger accumulation of TEs in the outgroup relative to *Scalesia* (Figure 02; Supplementary

Figure 04). In all the landscape plots, the TE/repeat accumulation peak of *Scalesia* is at a Kimura

substitution level of approximately 7, while the genome assembly of *S. atractyloides* is around 12

(Supplementary Figure 05). The repeat landscape plots decomposed in different TE and repeat classes show

that the accumulation of TEs and repeats does not differ when comparing the outgroup (Supplementary

Figure 06), *Scalesia* (Supplementary Figure 07-09), and *Scalesia* hybrids (Supplementary Figure 10).

**Organization of genome-specific elements**

**Figure 03** - UpSet plot of long terminal repeat elements (LTR) identified within the *Scalesia* radiation

(other TE and repeat groups are found in Supplementary Figures 11-15). The *UpSet* plot is an alternative

visualisation to a Venn diagram where the y-axis on the top graphic shows a frequency of LTR orthogroups

(groups of closely related LTRs) and the x-axis shows different combinations of the data (the first column

includes orthogroups present in all *Scalesia*, the second column are orthogroups exclusive to S. *retroflexa x*

*helleri* hybrids, the third column are orthogroups exclusive to *S. atractyloides* var. *atractyloides*). Dark grey

circles highlight species combinations.

Using *OrthoFinder*, we grouped TEs and repeats into orthogroups, sets of orthologous sequences

derived from the last common ancestor of all species under consideration (Figure 02; Supplementary Figures

11-15). In almost all the comparisons, we found that the majority of TE orthogroups are present in all

*Scalesia*. The sole exception was in the satellite elements, for which we detected very few orthologs. In the

largest class, LTR, we found 4,367 orthogroups present in all *Scalesia* genomes, while only 142-195

orthogroups were found to be unique to a single genome (Figure 03).

**Figure 04** - TE superfamily counts for different *Scalesia* species. The count scale is different for

different TE super families, namely *LTR/Copia* and *LTR/Gypsy*, due to the large numbers of these elements.

Superfamily-level analyses show that TE figures were remarkably even in *Scalesia* (Figure 04) and

the outgroup (Supplementary Table 16). With the exception of LTR/Gypsy and LTR/Copia, all TE

counts were below 30,000 elements and were remarkably even across *Scalesia* species. LTR/Gypsy and

LTR/Copia included 250,000-500,000 elements (Figure 04).

**Figure 05** - Representation of orthogroups present in *Scalesia* (yellow) or the outgroup (blue). Shared

orthogroups are given as overlap for different TEs and repeats.

We used *OrthoFinder* to identify orthogroups of TEs and repeats in five randomly chosen *Scalesia*

species and five outgroup *Pappobolus* species, and parsed the results to compare counts of *Scalesia*-specific

orthogroups, outgroup-specific orthogroups, and orthogroups shared between the two genera (Figure 05).

We chose to downscale the data to five random species because we observed imbalances in estimations that

arose from analysing the 15 *Scalesia* species in comparison to only five outgroup species (two specimens of

*P. hypargyreus* were included; Figure 05). Since we downscaled the data to only five *Scalesia* species, we

repeated the analyses three times, finding that the results were consistent across the three iterations

(Supplementary Table 03). The largest three classes of TEs (LTR, DNA and Helitron) had a majority of its

orthogroups common to *Scalesia* and the outgroup. *Scalesia* had more private SINEs than the outgroup

(Figure 05).

## Discussion

Transposable elements (TEs) play a crucial role in shaping genomes and phenotype diversity. Here,

we tested whether rapid diversification, shifts in climatic niches or hybridization are associated with an

increase of TE accumulation in the genomes of the genus *Scalesia*, a radiation endemic to the Galápagos

(Figure 01). Our results show that TE or repeat accumulation is consistent across the *Scalesia* phylogeny,

with no major changes across species (Figure 01) or over time (with one exception; Figure 02). We also

found no evidence of lineage-specific TE or repeat expansions, as most TE groups were found across all

species (Figure 03), in similar numbers across the explored genomes (Figure 04), and shared with the

outgroup (Figure 05). Overall, our findings indicate that neither rapid diversification, hybridization, nor

ecological niche shifts are significantly associated with TE and repeat content variation within the *Scalesia*

adaptive radiation.

**Limited variation in transposable element content in *Scalesia* despite rapid diversification**

We found no major differences in TE or repeat content in different *Scalesia* species in any of the

analyses performed and focusing on phylogenetic comparisons, accumulation over evolutionary time, and

class-specific expansions (Figures 01-5). The lack of variation may be attributable to the methods used, but

this seems unlikely since *dnaPipeTE* retrieved different results for *Pappobolus* and *Scalesia*, suggesting that

the two lineages have accumulated different TEs and repeats after their split, about 3 million years ago

(Fernández-Mazuecos et al. 2020). *dnaPipeTE* has been shown to detect population-level differences

(Kapun et al. 2020; Lerat et al. 2019; Blommaert et al. 2019; Vargas-Chavez et al. 2022), and between closely

related species across the Tree of Life (Talla et al. 2017; Jaron et al. 2021; Wong et al. 2019; Hill and

Betancourt 2018; Parisot et al. 2021; Lata et al. 2022; Castro et al. 2020). One tempting explanation for the

lack of variation is that *dnaPipeTE* may be biased to find conserved old elements, thereby explaining the

observed patterns. However, this is likely not the case as *dnaPipeTE* is very sensitive in detecting recent TE

families, being less sensitive to older TE families due to increased divergence and short reads that limits

alignments. For instance, when applied to the human genome, *dnaPipeTE* is biased towards more recent

TEs (see supplementary data included in (Clément Goubert et al. 2015). As a result, we exclude method-

orient bias, and we suggest that TEs are under phylogenetic inertia, as shown for various *Drosophila* species

(Sessegolo, Burlet, and Haudry 2016). Our findings suggest that the fundamental transposable element

composition remains conserved across *Scalesia* lineages.

The lack of variation TE content in *Scalesia* is noteworthy. In our dataset, we observe a variation of

7.23 % in LTRs, and 0.28 % of variation in DNA elements. This variation is very little when compared to

other works focusing on congeneric species. For instance, in the plant genus *Passifora*, Tekay LTR-elements

varied between 8.52 - 35.71 % and Angela LTR-elements between 1.96 - 29.05% (absolute estimates

considering the whole genome; (Sader et al. 2021). In oaks (*Quercus*) LTRs varied between 34.07 - 55.05 %

(relative TE content; (Mascagni et al. 2019). In *Rhodnius* bugs, with different ecologies, Tc1/Mariner

elements varied between 14.3 - 25 % (Castro et al. 2020), while in *Diabrotica* beetles MITE elements varied

between 150 - 610 MegaBasepairs (Lata et al. 2022). In this regard, the variation in *Scalesia* is diminished

and noteworthy.

One potential explanation for the observed lack of species-specific TEs is the high repeat content of

the *Scalesia* genome. Our previous analyses revealed that approximately 80% of the *Scalesia atractyloides*

genome is composed of repetitive sequences (Cerca et al. 2022). This high repeat content could be due to

the presence of numerous repeat families that actively replicate with a strong phylogenetic signal, rather

than a dynamic turnover of species-specific repeat families with a low copy number. In this regard, Novák

and colleagues (2020) found that TE content reaches a plateau at around 80% across angiosperms with

genome sizes between 5 and 10 Gbp (Novák et al. 2020). It is possible that a plateau of TE content could be

maintained, with few families represented at high numbers, and that a dynamic turnover occurs within

families, rather than between TE families.

Our findings challenge the notion that lineages with high speciation rates also display elevated rates

of TE content accumulation. One caveat to this interpretation is that the sister lineage to *Scalesia*, including

*Pappobolus*, may have undergone similar diversification on the mainland (Fernández-Mazuecos et al. 2020).

Accordingly, we hypothesize that the entire *Pappobolus-Scalesia* clade possesses ancestrally high TE content

. However, recent TE activity is observed as in both lineages most TEs are relatively young (Figure 02). In

any case, the role of TEs in adaptive radiation remains debated, with some studies demonstrating their

importance as drivers of radiation (Feiner 2016; Carleton et al. 2020), while others proposing a less

significant role (Xiong et al. 2021; Ronco et al. 2021). This discrepancy could stem from the scale and data

employed. Specifically, the correlation between TE content and diversification based on limited data may

have led to an overestimation of TEs' role. For instance, an initial analysis of four transcriptomes and

genomes of East African cichlids, compared to tilapia and other teleosts, suggested a higher accumulation of

TE counts during the radiation (Brawand et al. 2014). In contrast, recent evidence involving analyses of

hundreds of genomes found no relationship between species diversity and TEs within different tribes of the

Tanganyika radiation (Ronco et al. 2021). However, this does not negate the compelling experimental

evidence supporting the role of TEs in driving the evolution of specific phenotypic change, such as the

elegant findings of Kratochwil and colleagues demonstrating that a TE insertion triggered the emergence of

a novel phenotype, the cichlid gold morph (Kratochwil et al. 2022). Two conclusions can be drawn from

our results and the literature. First, it is necessary to conduct additional studies that functionally validate the

evidence and the mechanisms underlying TE insertions and their impact on phenotypic evolution. Second,

caution should be taken with potential overestimation of the role of TEs as major driving forces in

speciation and ecological diversification at the macroevolutionary scale in all cases (e.g. (Ricci et al. 2018;

Pace and Feschotte 2007; Ray et al. 2008; Pritham and Feschotte 2007; de Boer et al. 2007; Falcon, Tanaka,

and Rodriguez-Terrones 2023; Carducci et al. 2019; Belyayev 2014; Oliver and Greene 2009, 2012; Alioto

et al. 2020)). This is particularly evident given the limitations of existing studies, which often rely on limited

genomic data, correlational analyses, and a narrow taxonomic scope. Some of these correlations

encompassing macroevolutionary scales might have overstated their impact on adaptive evolution,

speciation, and their direct connection to evolutionary success. Indeed, it can be argued that TEs are

unlikely to enhance diversification unless genetic variation is limiting, such as in a bottleneck – as expected

in lineages such as *Scalesia*. While TEs offer a convenient explanation for rapid evolution, it is likely that

they are merely "along for the ride" and may only be purged when they reach problematic levels of

replication.

**TEs and ecological variation**

The minimal variation in TE accumulation observed in *Scalesia* suggests that the climatic niche

does not significantly influence TE content within the radiation. In adaptive radiations, closely related

species typically occupy distinct ecological niches (Gillespie et al. 2020; Schluter 2000; Arnegard et al. 2014).

In *Scalesia*, species exhibit distinct climatic niches (Figure 01) (Blaschke and Sanders 2009; Fernández-

Mazuecos et al. 2020; Perez et al. 2023), with temperature and the presence of arid environments being key

factors shaping the evolution of the radiation (Fernández-Mazuecos et al. 2020; Cerca et al. 2022; Perez et al.

2023). Various studies have demonstrated that aridity exerts a significant selective pressure on plant genome

size (e.g. (Doyle and Coate 2019; Schley et al. 2022). In arid regions, many plant species face strong selective

pressures to maintain smaller genome sizes (Doyle and Coate 2019), which correlate with smaller cell size

(Beaulieu et al. 2008). Smaller cells improve water economy through smaller stomata that respond more

efficiently to water stress, enhance gas exchange regulation, and optimize photosynthetic performance

(Franks and Beerling 2009; Lawson and Blatt 2014; Roddy et al. 2020). Given that arid environments favor

smaller cells, aridity is hypothesized to be an important selective pressure regulating genome size (Schley et

al. 2022). Therefore, it would be expected that TEs are being purged in species occupying arid coastal areas

(e.g. *S. affinis*), where temperatures frequently reach 40°C throughout the year and water stress is pervasive,

compared to species found in the humid highlands (e.g., *S. pedunculata*), where conditions are consistently

moist and cooler. This leads to an alternative hypothesis: perhaps selection is unable to effectively remove

transposable elements. The removal of a single TE, only a few kilobases in size, might be 'invisible' in a

genome of 4 gigabases (X. Wang et al. 2021).

Interestingly, the inclusion of an infraspecific taxon in the study helped observe a distinct pattern at

the microevolutionary scale, as *S. atractyloides* var. *atractyloides* exhibits a lower transposable element

content (Figure 02). This observation aligns with an emerging evolutionary change occurring in the arid

volcanic regions of Santiago Island, where a shift in leaf pubescence has also been observed (Wiggins and

Porter 1971)

**TEs and hybridization**

The four hybrids examined in this study exhibited similar TE content to non-hybrid populations,

excluding the possibility of genome deregulation through TE-release as a result of hybridization.

Traditionally, hybridization has been perceived as a detrimental force to the integrity of species (Mayr

1940), and it has been proposed that hybridization may lead to genome destabilisation by promoting TE

activity (Zhang et al. 2018; McClintock 1984). The presence of more TEs in hybrid populations would be

expected in a scenario involving genome shock triggered by multi-generational hybridization. In such cases,

increased TE transcription should lead to more TEs insertions in the hybrid genomes, and a consequent

increase of genome size. However, the case in *Scalesia*, as evidenced by the absence of supporting patterns in

the repeat accumulation curves (Figure 02), the lack of superfamily expansion or super-family level (Figure

03), and absolute accumulation (Figure 01). These findings are in agreement with those of (Kawakami et al.

2011), who demonstrated no evidence of LTR retrotransposon proliferation when comparing multiple

hybrid populations of *Helianthus annuus* (sunflower) × *H. petiolaris*. This result suggests that post-

transcriptional repression mechanisms may prevent the proliferation of LTRs (Kawakami et al. 2011). A

similar scenario may be at play in *Scalesia*, but additional data would be required to confirm this hypothesis.

## Acknowledgements

We were supported by the Research Council of Norway, project 287327. We used the supercomputer Saga, managed by Sigma2 (National e-infrastructure services of Norway; projects nn10082k, and nn9449k). Samples fromf *Scalesia* populations were originally obtained in accordance with Galápagos National Park research permit number PC-001/98 PNG, and later subjected to normalisation procedures authorized by the Ecuadorian Ministry of the Environment genetic permit number MAAE-DBI-CM-2021-0213. This publication is contribution number 2613 of the Charles Darwin Foundation for the Galápagos Islands. We are grateful to Alexander Suh, William B. Reinar and Irene Julca for support in TE analyses and for their helpful comments. We are also thankful to Ross McCauley for providing the *Pappobolus* samples and to Justine Villalba for assisting with the sampling of the *Scalesia* herbarium specimens.

## References

Alioto, Tyler, Konstantinos G. Alexiou, Amélie Bardil, Fabio Barteri, Raúl Castanera, Fernando Cruz, Amit Dhingra, et al. 2020. "Transposons Played a Major Role in the Diversification between the Closely Related Almond and Peach Genomes: Results from the Almond Genome Sequence." *The Plant Journal: For Cell and Molecular Biology* 101 (2): 455–72.

Almeida, Miguel Vasconcelos, Grégoire Vernaz, Audrey L. K. Putman, and Eric A. Miska. 2022. "Taming Transposable Elements in Vertebrates: From Epigenetic Silencing to Domestication." *Trends in Genetics: TIG* 38 (6): 529–53.

Arnegard, Matthew E., Matthew D. McGee, Blake Matthews, Kerry B. Marchinko, Gina L. Conte, Sahriar Kabir, Nicole Bedford, et al. 2014. "Genetics of Ecological Divergence during Speciation." *Nature* 511 (7509): 307–11.

Beaulieu, Jeremy M., Ilia J. Leitch, Sunil Patel, Arjun Pendharkar, and Charles A. Knight. 2008. "Genome Size Is a Strong Predictor of Cell Size and Stomatal Density in Angiosperms." *The New Phytologist* 179 (4): 975–86.

Belyayev, A. 2014. "Bursts of Transposable Elements as an Evolutionary Driving Force." *Journal of Evolutionary Biology* 27 (12): 2573–84.

Bennetzen, Jeffrey L., and Hao Wang. 2014. "The Contributions of Transposable Elements to the Structure, Function, and Evolution of Plant Genomes." *Annual Review of Plant Biology* 65 (February):505–30.

Blaschke, Jeremy D., and Roger W. Sanders. 2009. "PRELIMINARY INSIGHTS INTO THE PHYLOGENY AND SPECIATION OF SCALESIA (ASTERACEAE), GALÁPAGOS ISLANDS." *Journal of the Botanical Research Institute of Texas* 3 (1): 177–91.

Blommaert, J., S. Riss, B. Hecox-Lea, D. B. Mark Welch, and C. P. Stelzer. 2019. "Small, but Surprisingly Repetitive Genomes: Transposon Expansion and Not Polyploidy Has Driven a Doubling in Genome Size in a Metazoan Species Complex." *BMC Genomics* 20 (1): 466.

Boer, Johan G. de, Ryosuke Yazawa, William S. Davidson, and Ben F. Koop. 2007. "Bursts and Horizontal Evolution of DNA Transposons in the Speciation of Pseudotetraploid Salmonids." *BMC Genomics* 8

(November):422.

Bourque, Guillaume, Kathleen H. Burns, Mary Gehring, Vera Gorbunova, Andrei Seluanov, Molly
Hammell, Michaël Imbeault, et al. 2018. "Ten Things You Should Know about Transposable
Elements." *Genome Biology* 19 (1): 199.

Brawand, D., C. E. Wagner, Y. I. Li, M. Malinsky, and I. Keller. 2014. "The Genomic Substrate for
Adaptive Radiation in African Cichlid Fish." *Nature*. https://www.nature.com/articles/nature13726.

Carducci, F., M. A. Biscotti, M. Barucca, and A. Canapa. 2019. "Transposable Elements in Vertebrates:
Species Evolution and Environmental Adaptation." *The European Zoological Journal* 86 (1): 497–503.

Carleton, Karen L., Matthew A. Conte, Milan Malinsky, Sri Pratima Nandamuri, Benjamin A. Sandkam,
Joana I. Meier, Salome Mwaiko, Ole Seehausen, and Thomas D. Kocher. 2020. "Movement of
Transposable Elements Contributes to Cichlid Diversity." *Molecular Ecology* 29 (24): 4956–69.

Castro, Marcelo R. J., Clément Goubert, Fernando A. Monteiro, Cristina Vieira, and Claudia M. A.
Carareto. 2020. "Homology-Free Detection of Transposable Elements Unveils Their Dynamics in
Three Ecologically Distinct Rhodnius Species." *Genes* 11 (2).
https://doi.org/10.3390/genes11020170.

Cerca, José, Ellie E. Armstrong, Joel Vizueta, Rosa Fernández, Dimitar Dimitrov, Bent Petersen, Stefan
Prost, Julio Rozas, Dmitri Petrov, and Rosemary G. Gillespie. 2021. "The Tetragnatha Kauaiensis
Genome Sheds Light on the Origins of Genomic Novelty in Spiders." *Genome Biology and Evolution*
13 (12). https://doi.org/10.1093/gbe/evab262.

Cerca, José, Darko D. Cotoras, Vanessa C. Bieker, Rishi De-Kayne, Pablo Vargas, Mario Fernández-
Mazuecos, Julia López-Delgado, et al. 2023. "Evolutionary Genomics of Oceanic Island Radiations."
*Trends in Ecology & Evolution*, March. https://doi.org/10.1016/j.tree.2023.02.003.

Cerca, José, Bent Petersen, José Miguel Lazaro-Guevara, Angel Rivera-Colón, Siri Birkeland, Joel Vizueta,
Siyu Li, et al. 2022. "The Genomic Basis of the Plant Island Syndrome in Darwin's Giant Daisies."
*Nature Communications* 13 (1): 3729.

Choi, Jae Young, and Yuh Chwen G. Lee. 2020. "Double-Edged Sword: The Evolutionary Consequences
of the Epigenetic Silencing of Transposable Elements." *PLoS Genetics* 16 (7): e1008872.

Conart, Corentin, Nathanaelle Saclier, Fabrice Foucher, Clément Goubert, Aurélie Rius-Bony, Saretta N.
Paramita, Sandrine Moja, et al. 2022. "Duplication and Specialization of NUDX1 in Rosaceae Led to
Geraniol Production in Rose Petals." *Molecular Biology and Evolution* 39 (2).
https://doi.org/10.1093/molbev/msac002.

Doyle, Jeff J., and Jeremy E. Coate. 2019. "Polyploidy, the Nucleotype, and Novelty: The Impact of
Genome Doubling on the Biology of the Cell." *International Journal of Plant Sciences* 180 (1): 1–52.

Emms, David M., and Steven Kelly. 2015. "OrthoFinder: Solving Fundamental Biases in Whole Genome
Comparisons Dramatically Improves Orthogroup Inference Accuracy." *Genome Biology* 16
(August):157.

———. 2019. "OrthoFinder: Phylogenetic Orthology Inference for Comparative Genomics." *Genome
Biology* 20 (1): 238.

Falcon, Francisco, Elly M. Tanaka, and Diego Rodriguez-Terrones. 2023. "Transposon Waves at the Water-
to-Land Transition." *Current Opinion in Genetics & Development* 81 (August):102059.

Feiner, Nathalie. 2016. "Accumulation of Transposable Elements in Hox Gene Clusters during Adaptive
Radiation of Anolis Lizards." *Proceedings. Biological Sciences / The Royal Society* 283 (1840).
https://doi.org/10.1098/rspb.2016.1555.

Fernández-Mazuecos, Mario, Pablo Vargas, Ross A. McCauley, David Monjas, Ana Otero, Jaime A.
Chaves, Juan Ernesto Guevara Andino, and Gonzalo Rivas-Torres. 2020. "The Radiation of Darwin's
Giant Daisies in the Galápagos Islands." *Current Biology: CB* 30 (24): 4989–98.e7.

Flynn, Jullien M., Robert Hubley, Clément Goubert, Jeb Rosen, Andrew G. Clark, Cédric Feschotte, and
Arian F. Smit. 2020. "RepeatModeler2 for Automated Genomic Discovery of Transposable Element

Families." *Proceedings of the National Academy of Sciences of the United States of America* 117 (17):
9451–57.

Franks, Peter J., and David J. Beerling. 2009. "Maximum Leaf Conductance Driven by CO2 Effects on
Stomatal Size and Density over Geologic Time." *Proceedings of the National Academy of Sciences of the*
*United States of America* 106 (25): 10343–47.

Gillespie, Rosemary G., Gordon M. Bennett, Luc De Meester, Jeffrey L. Feder, Robert C. Fleischer, Luke J.
Harmon, Andrew P. Hendry, et al. 2020. "Comparing Adaptive Radiations Across Space, Time, and
Taxa." *The Journal of Heredity* 111 (1): 1–20.

Goubert, Clément. 2023. "Assembly-Free Detection and Quantification of Transposable Elements with
dnaPipeTE." *Methods in Molecular Biology* 2607:25–43.

Goubert, Clement, Rory J. Craig, Agustin F. Bilat, Valentina Peona, Aaron A. Vogan, and Anna V.
Protasio. 2022. "A Beginner's Guide to Manual Curation of Transposable Elements." *Mobile DNA* 13
(1): 7.

Goubert, Clément, Laurent Modolo, Cristina Vieira, Claire ValienteMoro, Patrick Mavingui, and Matthieu
Boulesteix. 2015. "De Novo Assembly and Annotation of the Asian Tiger Mosquito (Aedes
Albopictus) Repeatome with dnaPipeTE from Raw Genomic Reads and Comparative Analysis with
the Yellow Fever Mosquito (Aedes Aegypti)." *Genome Biology and Evolution* 7 (4): 1192–1205.

Hill, Tom, and Andrea J. Betancourt. 2018. "Extensive Exchange of Transposable Elements in the
Drosophila Pseudoobscura Group." *Mobile DNA* 9 (June):20.

Jaron, Kamil S., Jens Bast, Reuben W. Nowell, T. Rhyker Ranallo-Benavidez, Marc Robinson-Rechavi, and
Tanja Schwander. 2021. "Genomic Features of Parthenogenetic Animals." *The Journal of Heredity*
112 (1): 19–33.

Jiang, Ning, Zhirong Bao, Xiaoyu Zhang, Hirohiko Hirochika, Sean R. Eddy, Susan R. McCouch, and
Susan R. Wessler. 2003. "An Active DNA Transposon Family in Rice." *Nature* 421 (6919): 163–67.

Kapun, Martin, Maite G. Barrón, Fabian Staubach, Darren J. Obbard, R. Axel W. Wiberg, Jorge Vieira,
Clément Goubert, et al. 2020. "Genomic Analysis of European Drosophila Melanogaster Populations
Reveals Longitudinal Structure, Continent-Wide Selection, and Previously Unknown DNA Viruses."
*Molecular Biology and Evolution* 37 (9): 2661–78.

Kawakami, Takeshi, Preeti Dhakal, Angela N. Katterhenry, Chelsea A. Heatherington, and Mark C.
Ungerer. 2011. "Transposable Element Proliferation and Genome Expansion Are Rare in
Contemporary Sunflower Hybrid Populations despite Widespread Transcriptional Activity of LTR
Retrotransposons." *Genome Biology and Evolution* 3 (January):156–67.

Kidwell, M. G., and D. Lisch. 1997. "Transposable Elements as Sources of Variation in Animals and
Plants." *Proceedings of the National Academy of Sciences of the United States of America* 94 (15): 7704–
11.

Kratochwil, Claudius F., Andreas F. Kautt, Alexander Nater, Andreas Härer, Yipeng Liang, Frederico
Henning, and Axel Meyer. 2022. "An Intronic Transposon Insertion Associates with a Trans-Species
Color Polymorphism in Midas Cichlid Fishes." *Nature Communications* 13 (1): 1–8.

Lata, Dimpal, Brad S. Coates, Kimberly K. O. Walden, Hugh M. Robertson, and Nicholas J. Miller. 2022.
"Genome Size Evolution in the Beetle Genus Diabrotica." *G3* 12 (4).
https://doi.org/10.1093/g3journal/jkac052.

Lawson, Tracy, and Michael R. Blatt. 2014. "Stomatal Size, Speed, and Responsiveness Impact on
Photosynthesis and Water Use Efficiency." *Plant Physiology* 164 (4): 1556–70.

Lefort, Vincent, Richard Desper, and Olivier Gascuel. 2015. "FastME 2.0: A Comprehensive, Accurate,
and Fast Distance-Based Phylogeny Inference Program." *Molecular Biology and Evolution* 32 (10):
2798–2800.

Lerat, Emmanuelle, Clément Goubert, Sara Guirao-Rico, Miriam Merenciano, Anne-Béatrice Dufour,
Cristina Vieira, and Josefa González. 2019. "Population-Specific Dynamics and Selection Patterns of

Transposable Element Insertions in European Natural Populations." *Molecular Ecology* 28 (6): 1506–22.

Marçais, Guillaume, and Carl Kingsford. 2011. "A Fast, Lock-Free Approach for Efficient Parallel Counting of Occurrences of K-Mers." *Bioinformatics* 27 (6): 764–70.

Martin, Michael D., Elizabeth A. Zimmer, Morten T. Olsen, Andrew D. Foote, M. Thomas P. Gilbert, and Grace S. Brush. 2014. "Herbarium Specimens Reveal a Historical Shift in Phylogeographic Structure of Common Ragweed during Native Range Disturbance." *Molecular Ecology* 23 (7): 1701–16.

Mascagni, Flavia, Alberto Vangelisti, Tommaso Giordani, Andrea Cavallini, and Lucia Natali. 2019. "A Computational Comparative Study of the Repetitive DNA in the Genus Quercus L." *Tree Genetics & Genomes* 16 (1): 11.

Mayr, Ernst. 1940. "Systematics and the Origin of Species from the Point of View of a Zoologist (New York, 1942; New Edn, Cambridge, Mass., 1999), 173--85; Mayr,"." *Speciation Phenomena in Birds", American Naturalist* 74:249–78.

McClintock, B. 1984. "The Significance of Responses of the Genome to Challenge." *Science* 226 (4676): 792–801.

Novák, Petr, Maïté S. Guignard, Pavel Neumann, Laura J. Kelly, Jelena Mlinarec, Andrea Koblížková, Steven Dodsworth, et al. 2020. "Repeat-Sequence Turnover Shifts Fundamentally in Species with Large Genomes." *Nature Plants* 6 (11): 1325–29.

Oliver, Keith R., and Wayne K. Greene. 2009. "Transposable Elements: Powerful Facilitators of Evolution." *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology* 31 (7): 703–14.

———. 2012. "Transposable Elements and Viruses as Factors in Adaptation and Evolution: An Expansion and Strengthening of the TE-Thrust Hypothesis." *Ecology and Evolution*. https://doi.org/10.1002/ece3.400.

Oliver, Keith R., Jen A. McComb, and Wayne K. Greene. 2013. "Transposable Elements: Powerful Contributors to Angiosperm Evolution and Diversity." *Genome Biology and Evolution* 5 (10): 1886–1901.

Ondov, Brian D., Todd J. Treangen, Páll Melsted, Adam B. Mallonee, Nicholas H. Bergman, Sergey Koren, and Adam M. Phillippy. 2016. "Mash: Fast Genome and Metagenome Distance Estimation Using MinHash." *Genome Biology* 17 (1): 132.

Pace, John K., 2nd, and Cédric Feschotte. 2007. "The Evolutionary History of Human DNA Transposons: Evidence for Intense Activity in the Primate Lineage." *Genome Research* 17 (4): 422–32.

Parisod, Christian, Karine Alix, Jérémy Just, Maud Petit, Véronique Sarilar, Corinne Mhiri, Malika Ainouche, Boulos Chalhoub, and Marie-Angèle Grandbastien. 2010. "Impact of Transposable Elements on the Organization and Function of Allopolyploid Genomes." *The New Phytologist* 186 (1): 37–45.

Parisot, Nicolas, Carlos Vargas-Chávez, Clément Goubert, Patrice Baa-Puyoulet, Séverine Balmand, Louis Beranger, Caroline Blanc, et al. 2021. "The Transposable Element-Rich Genome of the Cereal Pest Sitophilus Oryzae." *BMC Biology* 19 (1): 241.

Perez, Timothy M., Juan Ernesto Guevara Andino, Gonzalo Rivas-Torres, and Kenneth J. Feeley. 2023. "Climate Constrains Photosynthetic Strategies in Darwin's Daisies: A Test of the Climatic Variability and Jack-of-All-Trades Hypotheses." *The American Naturalist* 201 (1): 78–90.

Petrov, D. A., J. L. Schutzman, D. L. Hartl, and E. R. Lozovskaya. 1995. "Diverse Transposable Elements Are Mobilized in Hybrid Dysgenesis in Drosophila Virilis." *Proceedings of the National Academy of Sciences of the United States of America* 92 (17): 8050–54.

Pimpinelli, Sergio, and Lucia Piacentini. 2020. "Environmental Change and the Evolution of Genomes: Transposable Elements as Translators of Phenotypic Plasticity into Genotypic Variability." *Functional Ecology* 34 (2): 428–41.

Pritham, Ellen J., and Cédric Feschotte. 2007. "Massive Amplification of Rolling-Circle Transposons in the Lineage of the Bat Myotis Lucifugus." *Proceedings of the National Academy of Sciences of the United States of America* 104 (6): 1895–1900.

Rachtman, Eleonora, Shahab Sarmashghi, Vineet Bafna, and Siavash Mirarab. 2022. "Quantifying the Uncertainty of Assembly-Free Genome-Wide Distance Estimates and Phylogenetic Relationships Using Subsampling." *Cell Systems* 13 (10): 817–29.e3.

Ray, David A., Cedric Feschotte, Heidi J. T. Pagan, Jeremy D. Smith, Ellen J. Pritham, Peter Arensburger, Peter W. Atkinson, and Nancy L. Craig. 2008. "Multiple Waves of Recent DNA Transposon Activity in the Bat, Myotis Lucifugus." *Genome Research* 18 (5): 717–28.

Ricci, Marco, Valentina Peona, Etienne Guichard, Cristian Taccioli, and Alessio Boattini. 2018. "Transposable Elements Activity Is Positively Related to Rate of Speciation in Mammals." *Journal of Molecular Evolution* 86 (5): 303–10.

Roddy, Adam B., Guillaume Théroux-Rancourt, Tito Abbo, Joseph W. Benedetti, Craig R. Brodersen, Mariana Castro, Silvia Castro, et al. 2020. "The Scaling of Genome Size and Cell Size Limits Maximum Rates of Photosynthesis with Implications for Ecological Strategies." *International Journal of Plant Sciences* 181 (1): 75–87.

Ronco, Fabrizia, Michael Matschiner, Astrid Böhne, Anna Boila, Heinz H. Büscher, Athimed El Taher, Adrian Indermaur, et al. 2021. "Drivers and Dynamics of a Massive Adaptive Radiation in Cichlid Fishes." *Nature* 589 (7840): 76–81.

Rubin, G. M., M. G. Kidwell, and P. M. Bingham. 1982. "The Molecular Basis of P-M Hybrid Dysgenesis: The Nature of Induced Mutations." *Cell* 29 (3): 987–94.

Sader, Mariela, Magdalena Vaio, Luiz Augusto Cauz-Santos, Marcelo Carnier Dornelas, Maria Lucia Carneiro Vieira, Natoniel Melo, and Andrea Pedrosa-Harand. 2021. "Large vs Small Genomes in Passiflora: The Influence of the Mobilome and the Satellitome." *Planta* 253 (4): 86.

Sarmashghi, Shahab, Kristine Bohmann, M. Thomas P Gilbert, Vineet Bafna, and Siavash Mirarab. 2019. "Skmer: Assembly-Free and Alignment-Free Sample Identification Using Genome Skims." *Genome Biology* 20 (1): 34.

Schley, Rowan J., Jaume Pellicer, Xue-Jun Ge, Craig Barrett, Sidonie Bellot, Maïté S. Guignard, Petr Novák, et al. 2022. "The Ecology of Palm Genomes: Repeat-Associated Genome Size Expansion Is Constrained by Aridity." *The New Phytologist* 236 (2): 433–46.

Schluter, D. 2000. "The Ecology of Adaptive Radiation." https://books.google.com/books?hl=en&lr=&id=Q1wxNmLAL10C&oi=fnd&pg=PR7&dq=schluter+2000+ecology+of+adaptive&ots=4sJ7RGaMvq&sig=TMKvoFMaMWtAYiKX8J8wy0Qal70.

Schubert, Mikkel, Stinus Lindgreen, and Ludovic Orlando. 2016. "AdapterRemoval v2: Rapid Adapter Trimming, Identification, and Read Merging." *BMC Research Notes* 9 (February):88.

Serrato-Capuchina, Antonio, Jeremy Wang, Eric Earley, David Peede, Kristin Isbell, and Daniel R. Matute. 2020. "Paternally Inherited P-Element Copy Number Affects the Magnitude of Hybrid Dysgenesis in Drosophila Simulans and D. Melanogaster." *Genome Biology and Evolution* 12 (6): 808–26.

Sessegolo, Camille, Nelly Burlet, and Annabelle Haudry. 2016. "Strong Phylogenetic Inertia on Genome Size and Transposable Element Content among 26 Species of Flies." *Biology Letters* 12 (8). https://doi.org/10.1098/rsbl.2016.0407.

Talla, Venkat, Alexander Suh, Faheema Kalsoom, Vlad Dinca, Roger Vila, Magne Friberg, Christer Wiklund, and Niclas Backström. 2017. "Rapid Increase in Genome Size as a Consequence of Transposable Element Hyperactivity in Wood-White (Leptidea) Butterflies." *Genome Biology and Evolution* 9 (10): 2491–2505.

Van't Hof, Arjen E., Pascal Campagne, Daniel J. Rigden, Carl J. Yung, Jessica Lingley, Michael A. Quail, Neil Hall, Alistair C. Darby, and Ilik J. Saccheri. 2016. "The Industrial Melanism Mutation in British Peppered Moths Is a Transposable Element." *Nature* 534 (7605): 102–5.

Vargas-Chavez, Carlos, Neil Michel Longo Pendy, Sandrine E. Nsango, Laura Aguilera, Diego Ayala, and
Josefa González. 2022. "Transposable Element Variants and Their Potential Adaptive Impact in
Urban Populations of the Malaria Vector Anopheles Coluzzii." *Genome Research* 32 (1): 189–202.
Wang, Luyang, Shuo Zhang, Savana Hadjipanteli, Lorissa Saiz, Lisa Nguyen, Efren Silva, and Erin Kelleher.
2023. "P-Element Invasion Fuels Molecular Adaptation in Laboratory Populations of Drosophila
Melanogaster." *Evolution; International Journal of Organic Evolution* 77 (4): 980–94.
Wang, Xiaotong, Joseph A. Morton, Jaume Pellicer, Ilia J. Leitch, and Andrew R. Leitch. 2021. "Genome
Downsizing after Polyploidy: Mechanisms, Rates and Selection Pressures." *The Plant Journal: For Cell*
*and Molecular Biology* 107 (4): 1003–15.
Wiggins, Ira L., and Duncan M. Porter. 1971. *Flora of the Galápagos Islands.* Stanford University Press.
Wong, Wai Yee, Oleg Simakov, Diane M. Bridge, Paulyn Cartwright, Anthony J. Bellantuono, Anne Kuhn,
Thomas W. Holstein, Charles N. David, Robert E. Steele, and Daniel E. Martínez. 2019. "Expansion
of a Single Transposable Element Family Is Associated with Genome-Size Increase and Radiation in
the Genus *Hydra.*" *Proceedings of the National Academy of Sciences of the United States of America* 116
(46): 22915–17.
Xiong, Peiwen, C. Darrin Hulsey, Carmelo Fruciano, Wai Y. Wong, Alexander Nater, Andreas F. Kautt,
Oleg Simakov, et al. 2021. "The Comparative Genomic Landscape of Adaptive Radiation in Crater
Lake Cichlid Fishes." *Molecular Ecology* 30 (4): 955–72.
Zhang, Min, Xin-Kai Liu, Wen Fan, Dan-Feng Yan, Nai-Sheng Zhong, Ji-Yin Gao, and Wen-Ju Zhang.
2018. "Transcriptome Analysis Reveals Hybridization-Induced Genome Shock in an Interspecific F1
Hybrid from Camellia." *Genome / National Research Council Canada = Genome / Conseil National de*
*Recherches Canada* 61 (7): 477–85.

696