1 **Single Nuclei Sequencing Reveals Intratumoral Cellular Heterogeneity and Replication**
2 **Stress in Adrenocortical Carcinoma**
3

4 Liudmila V. Popova, PhD[1];* Elizabeth A. R. Garfinkle, PhD[2];* Daniel M. Chopyk, MD, PhD[1];*
5 Jaye B. Navarro[2], Adithe Rivaldi[2], Yaoling Shu[1], MD, MS; Elena Lomonosova, PhD[4], John E.
6 Phay, MD[1]; Barbra S. Miller, MD[1]; Swati Sattuwar, MD[3]; Mary Mullen, MD[4]; Elaine R. Mardis,
7 PhD[2, 5]; Katherine E. Miller, PhD[2, 5]; Priya H. Dedhia, MD, PhD[1, 6, 7†]
8
9
10 [1]Division of Surgical Oncology, The Ohio State University and Arthur G. James
11 Comprehensive Cancer Center, Columbus, Ohio, USA.
12
13 [2]The Steve and Cindy Rasmussen Institute for Genomic Medicine at Nationwide Children's
14 Hospital, Columbus, Ohio, USA.
15
16 [3]Department of Pathology, The Ohio State University, Columbus, Ohio, USA
17
18 [4]Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, Alvin J.
19 Siteman Cancer Center, Washington University School of Medicine, St Louis, MO, USA.
20
21 [5]Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH,
22 USA
23
24 [6]Translational Therapeutics Program, The Ohio State University and Arthur G. James
25 Comprehensive Cancer Center, Columbus, Ohio, USA.
26
27 [7]Center for Cancer Engineering, The Ohio State University, Columbus, Ohio, USA.
28
29
30 *Authors contributed equally to this manuscript.
31
32 †Correspondence
33
Priya Dedhia, MD, PhD
Division of Surgical Oncology
The Ohio State University and Arthur G. James Comprehensive Cancer Center
Department of Surgery Division of Surgical Oncology
N924 Doan Hall, 410 W. 10th Avenue
Columbus, OH 43210
priya.dedhia@osumc.edu
41

ABSTRACT

Adrenocortical carcinoma (ACC) is a rare endocrine malignancy with a poor prognosis and limited treatment options. Bulk genomic characterization of ACC has not yielded obvious therapeutic or immunotherapeutic targets, yet novel therapies are needed. We hypothesized that elucidating the intratumoral cellular heterogeneity by single nuclei RNA sequencing analyses would yield insights into potential therapeutic vulnerabilities of this disease. In addition to characterizing the immune cell and fibroblast landscape, our analyses of single nuclei gene expression profiles identified an adrenal cortex cell cluster exhibiting a program of replication stress and DNA damage response in primary and metastatic ACC. *In vitro* assessment of replication stress and DNA damage response using an ACC cell line and a series of newly-derived hormonally active patient-derived tumor organoids revealed ATR sensitivity. These findings provide novel mechanistic insight into ACC biology and suggest that an underlying dependency on ATR may be leveraged therapeutically in advanced ACC.

149 / 150 words

INTRODUCTION

Adrenocortical carcinoma (ACC) is a rare, aggressive malignancy that originates in the adrenal cortex and has a 5-year survival of ~6% for patients with stage IV disease.[1] While margin negative surgical resection is the most desirable frontline treatment, many patients present with metastatic disease necessitating treatment with systemic agents. In those able to undergo resection, most experience tumor recurrence.[2–5] Mitotane, an adrenolytic agent that also blocks cortisol synthesis, remains the only FDA-approved therapeutic agent specifically for treatment of advanced ACC. However, attaining therapeutic levels of mitotane is challenging due to significant toxicity and harmful side effects.[2,4] Administration of chemotherapeutic drugs - etoposide, doxorubicin, and cisplatin - in conjunction with mitotane improve outcomes, but combination therapy increases median progression-free survival by only 5 months and adds significant toxicity.[2,6] Immunotherapy has shown promise in certain cancers, yet its efficacy in ACC remains limited.[7] Thus, there is a critical need to understand molecular mechanisms driving ACC tumor progression to improve therapy.

Bulk sequencing strategies have allowed for comprehensive genomic characterization of ACC. Previous studies using whole exome sequencing, bulk mRNA sequencing, micro RNA sequencing, DNA copy number analysis, and DNA methylation analysis have enabled molecular classifications of ACC that correlate with distinct clinical outcomes.[8,9] Despite significant advances in understanding the genomic landscape of ACC, effective treatment options are still lacking. Single-cell RNA sequencing offers a powerful, complementary technique to bulk sequencing strategies by profiling individual cells, and thus enabling analysis of intratumoral cellular heterogeneity, a key factor that contributes to poor outcomes in many other cancers.[10–13] The rationale for this approach in ACC includes evidence of heterogeneity in nuclear localization of β-catenin in ACC[14] and identification of ACC tumor cell subpopulations with unique metabolomic signatures (e.g. increased pentose phosphate pathway activity) using spatial metabolomics.[15] In addition, recent work demonstrated intratumoral heterogeneity in ACC at single-nuclei resolution.[16] These observations suggest important cellular differences in ACC and identify a prospective cell of origin; however, they have not identified new therapeutic targets. Furthermore, prior single nuclei RNA sequencing (snRNAseq) of ACC consisted primarily of early-stage ACC, which is often amenable to surgical resection. Thus, we hypothesized that elucidating the underlying cellular biology at the single cell level in advanced ACC has the potential to identify novel therapeutic

vulnerabilities in this disease. In addition, prior snRNAseq of ACC demonstrated enrichment of

fibroblasts and immune cells compared to normal adrenal tissue but did not fully characterize

these populations to identify potential therapeutic targets.[16] Our snRNAseq analyses of

advanced ACC patient specimens further define the immune landscape and discover a

population of fibroblasts associated with ACC. We also identify distinct population of adrenal

cortex cells in ACC primary and metastatic samples, characterized by a unique transcriptomic

program of proliferation, replication stress, and DNA damage response. Incorporating an ACC

cell line and our novel ACC patient-derived tumor organoids (PTOs) in functional studies, we

uncover ataxia telangiectasia and Rad3-related protein (ATR) dependency, a promising

therapeutic strategy for this deadly disease.

RESULTS

snRNAseq reveals heterogeneous subpopulations in ACC

To profile cellular subpopulations in normal adrenal, adrenal adenoma, and ACC tissues, we

performed snRNAseq on 11 specimens collected from 9 patients. The samples included

normal adrenal tissue (n = 1), adrenal adenoma (AA, n = 2), primary ACC (PACC, n = 4), and

metastatic ACC (MACC, n = 4, Figures 1A, 1B, and Table 1). After standard preprocessing

steps, 4,735 nuclei from normal adrenal, 14,481 nuclei from AA, 23,837 nuclei from PACC

samples, and 19,151 nuclei from MACC samples were used for analysis. Samples were

integrated using the RPCA algorithm in Seurat and unbiased clustering resulted in 10 distinct

cell populations (Figure 1C). To define adrenal cortex cells, we calculated average module

scores (AC module scores) based on expression of adrenal cortex and ACC biomarkers,

NR5A1 encoding the protein SF1, *INHA*, and *MLANA*.[17–19] This analysis identified 6 adrenal

cortex clusters with high AC module scores and 4 clusters with low AC module scores. The 4

clusters with low AC module scores were identified as lymphocytes (*PTPRC*)[20], myeloid cells

(*ITGAM*)[21], fibroblasts (*SERPINE1* and *IGFBP7*)[22,23], and endothelial cells (*VWF* and

FLT1).[24,25] In addition to increased AC module scores, adrenal cortex clusters lacked

expression of *PTPRC*, *ITGAM*, *SERPINE1*, *IGFBP7*, *VWF*, or *FLT1* (Figure 1D and

Supplementary Figure S1). We also calculated cell type proportions and determined that

normal adrenal samples were characterized by the presence of 7 different cell clusters, AA

samples were characterized by the presence of 9 different cell clusters, and both PACC and

MACC samples were defined by the presence of 10 different cell clusters. (Figure 1E).

Overall, these data suggest that the cellular biology of ACC is defined by unique clusters

when compared to normal adrenal or adrenal adenoma.

ACC exhibits a unique tumor immune microenvironment

To characterize the immune landscape in adrenal tissues, we subset the lymphocyte and

myeloid cell clusters (Figure 1), reprocessed, integrated using Harmony[26], and performed

unbiased reclustering. Eight immune cell types were identified including macrophages,

granulocytes, regulatory T cells, CD4+ and CD8+ T cells, NK cells, B cells, and plasma cells

(Figure 2A). Across all specimens, macrophages were the most common immune cell (Figure

2B, Supplementary Tables S1A-2A). Key immunocyte-specific genes were plotted in a dot plot

and used to guide immune cell typing (Figure 2C).

We first focused on the macrophage cluster, as these were the dominant immune population.

1,661 total macrophages were subset, reprocessed, and reclustered. From this subset, an

additional 51 lymphocytes were identified and removed leaving 1,610 total macrophages for

further analysis (Figure 2D). We assessed macrophage polarity by expression of

proinflammatory or M1 markers, *CD80*, *CD86*, and *FCGR1A* (also known as *CD64*),[27,28] and

immunosuppressive or M2 markers, *MRC1*, *CD163*, and *MARCO*[29–31]. Although we noted no

significant differences in the proportion of M1 macrophages across all tissues, we found a

significant decrease in M2 macrophages in PACC and MACC specimens compared to normal

and AA specimens (false discovery rate (FDR) <0.05, Figure 2E). Prior work correlated

increased M2 macrophage infiltration with worse prognosis in other solid tumors; however,

more recent evidence indicates plasticity in M1 and M2 polarization.[32] Furthermore,

macrophage expression of *SPP1* was strongly associated with clinical prognosis independent

of classic M1 versus M2 macrophage markers.[33,34] Additionally, SPP1+ macrophages have

been associated with an immunosuppressive phenotype.[32,35,36] Therefore, we assessed *SPP1*

expression and found a significant increase in SPP1+ macrophages (FDR = 0.03) in PACC

and MACC specimens compared to normal and AA (Figure 2E, Supplementary Tables S1B-

2B). Our findings suggest that SPP1+ macrophages are associated with ACC tumor

progression.

We next analyzed the expression of immune checkpoint molecules expressed by
macrophages to identify potential strategies for immunotherapy. We plotted the average
expression of pro-inflammatory (*CD40* and *CD80*) and immunosuppressive (*CD274* or *PDL1*,
LGALS9, *PVR*, *SIRPA,* and *CD80*) markers (Figure 2G).[37–44] CD80 has a dual role as either a
costimulatory or coinhibitory ligand depending on whether it binds CD28 or CTLA-4,
respectively, on neighboring T cells.[44] We found that PACC-associated macrophages had
higher average expression of *CD274,* and *LGALS9*, and MACC-associated macrophages had
higher average expression of *PVR*, and *SIRPA*. PACC-associated macrophages also
exhibited elevated expression of *CD40* and *CD80* relative to the benign tissues, suggesting
these markers may serve as potential targets for immunotherapy. Taken together, our data
suggest a changing macrophage landscape from normal adrenal to ACC that increasingly
presents an immunosuppressive environment.
172
Analysis of lymphoid populations demonstrated that both T and NK cells were reduced or
absent in most ACC specimens. Interestingly, PACC 4, a tumor from a patient without clinical
or biochemical evidence of excess cortisol secretion, had the highest numbers of T and NK
cells compared to other samples (Supplementary Table S2A). This finding is consistent with
the association between hypercortisolism and reduced T cell infiltration in ACC (Table 1 and
Figure 1B).[8,45] Therefore, our analysis of CD8+, CD4+, and NK cells were primarily
represented by PACC 4. First, we subset CD8+ T cells and plotted average expression of key
immunoregulatory molecules (Supplementary Figure S2). Our analysis revealed that CD8+ T
cells in the ACC samples compared to those in benign tissue had reduced expression of
markers associated with immune activating functions (*CD28* and *CD40L*G) and increased
expression of markers of T cell exhaustion (*CTLA-4*, *PDCD1*, *TIGIT*, *LAG3*, and *HAVCR2*).[46–
48] We next assessed expression of these markers within the CD4+ T cell subset and found
that expression of *CD28*, *CTLA-4*, *TIGIT*, and *LAG3* was increased whereas expression of
CD40LG, *PDCD1*, and *HAVCR2* was decreased in ACC relative to normal adrenal and AA
(Supplementary Figure S2). Taken together, our analyses demonstrate signs of T cell
exhaustion in ACC which was more pronounced in CD8+ compared to CD4+ T cells. Finally,
analysis of the NK cells revealed increased expression of *TIGIT*[49] and the NK inhibitory
receptor *KLRC1*[50] in ACC relative to normal adrenal and AA specimens (Supplementary
Figure S2). Collectively, these findings indicate that hypercortisolism in ACC reduces T cell
infiltration, and thus ACC is characterized by an immunosuppressive microenvironment, which

may require precision therapeutic strategies that enhance immune infiltration in combination

with immunotherapy to achieve efficacy.

ACC is associated with a unique fibroblast subpopulation

We next subset fibroblasts to determine if malignant specimens are associated with changes

in fibroblast subpopulations. Initially we identified 10 fibroblast clusters. Four clusters including

adipocytes were excluded from further analysis, and the remaining fibroblast data were

reprocessed and reintegrated to identify 5 fibroblast clusters (Figure 3A-B). These clusters

were annotated based on marker gene expression (Figure 3C) and fibroblast markers (Figure

3D).[51,52] Fibroblast cluster 3.1 was present in all specimens and was the predominant cluster

in PACC and MACC (Figure 3B). Characterized by high expression of *ACTA2*, *MMP11*, and

HAS2, cluster 3.1 likely represents myofibroblastic cancer-associated fibroblasts

(myCAFs).[53,54] Further analysis of fibroblast cluster 3.1 in PACC specimens for known cancer-

associated fibroblasts markers[55] demonstrated increased levels of *PDGFRB, FAP, VIM,*

POSTN, HAS2, CXCL8, CXCL2, LMNA, and HAS 1. MACC specimens expressed increased

CAV1, CFD, MMP11, COL1A1, which are additional markers of cancer-associated fibroblasts

(Figure 3E). These findings suggest that ACC is characterized by the expansion of a distinct

fibroblast subpopulation at both primary and metastatic sites.

Adrenal cortex clusters demonstrate unique transcriptional programs

To further characterize the 6 adrenal cortex clusters, we subset, reintegrated, and performed

unbiased clustering of these data, which resulted in identification of 7 adrenal cortex clusters

and one adrenal medulla cluster (defined by *CHGA* expression, Supplementary Figure S3).[56]

Adrenal medulla cells were predominantly found in normal adrenal tissue (150 cells). After

excluding adrenal medulla cells, we reintegrated and reclustered adrenal cortex cells to reveal

7 final adrenal cortex clusters (AC 3.1-3.7) based on marker gene expression (Figures 4A and

B). The majority of the cells in cluster 3.7 originated from PACC 3, the only specimen with

excess DHEAS, androstenedione, and testosterone secretion. Consistent with these clinical

findings, AC 3.7 expressed high levels of *CYP11A1* and *CYP17A1* which are involved in

testosterone, DHEAS, and androstenedione production. (Figure 4C, Table 1, and

Supplementary Figure S4).[57,58] We then assessed subpopulation frequency across different

sample types and found that clusters AC 3.5 and AC 3.6 were significantly enriched in PACC

and MACC specimens (FDR < 0.0001, Figure 4D). In order to identify gene expression

changes associated with tumor progression, we calculated average module scores for each of

the clusters for Hallmark pathways downloaded from MSigDB[59,60] (Figure 4E and

Supplementary Figure S5). In these clusters, in which cell cycle regression was performed, we

found that cluster AC 3.6 was characterized by the highest module scores for the following

hallmark pathways: E2F targets, G2M checkpoint, mitotic spindle, PI3K AKT mTOR signaling,

and DNA repair. Hence, AC 3.6 bears a gene expression signature of dysfunctional DNA

replication and repair, indicating potential mechanistic underpinnings of malignancy and tumor

progression. Cluster AC 3.5 was defined by TGF-β signaling, unfolded protein response, and

glycolysis. The unfolded protein response and glycolysis signatures in cluster 3.5 were

enriched only in MACC, whereas the DNA damage repair signature in cluster AC 3.6 was

present in all PACC and MACC samples (Supplementary Figure S5). We also found that

cluster AC 3.3 was defined by the Wnt/β catenin signaling pathway, which is commonly

dysregulated in ACC.[2] Cluster AC 3.2 was defined by cholesterol homeostasis, indicating a

potential role in steroid metabolism.

Replication stress and DNA damage repair signature is enriched in cluster AC 3.6

To better understand the role of replication stress and DNA damage repair pathways in ACC,

we used REACTOME pathways downloaded from the MSigDB database to generate module

scores for each of the identified clusters.[59,61] We observed that cluster AC 3.6 in PACC and

MACC was characterized by the highest module scores associated with activation of ATR in

response to replication stress, homology-directed repair, non-homologous end joining,

mismatch repair, and base excision repair (Figure 5A). Additionally, we observed that

replication stress and DNA repair pathways were enriched in PACC and MACC specimens

(Supplementary Figure S6). We next used QIAGEN Ingenuity Pathway Analysis (IPA)[62] to

compare cluster AC 3.6 to other clusters found in PACC and MACC specimens and again

identified that cell cycle checkpoints and homology-directed repair through homologous

recombination or single-strand annealing were among the top 10 upregulated pathways

(Figure 5B). Replication stress and chromosome instability have been previously associated

with tumorigenesis and tumor progression.[63] To elucidate genomic instability in AC clusters,

we used inferCNV[64] to identify copy-number variation (CNV) in tumor specimens on a per-

cluster basis. We found that cluster AC 3.6 had the highest total number of CNVs, including

both copy number gains and losses compared to other clusters (Figure 5C-E). This

observation suggests that cluster AC 3.6, which demonstrates dysregulated replication stress

and DNA damage repair response as well as genomic instability, may reflect the underlying

biology driving ACC tumor development and progression.

ATR-dependent replication stress in ACC

To assess for the presence of replication stress and DNA damage in ACC, we performed

immunofluorescence on patient specimens (MACC 2 and MACC 3) for the DNA damage

marker, ☐H2AX and the replication stress marker phosphorylated RPA. Both proteins localize

to form foci, or clusters, in regions of DNA damage or replication stress.[65] Both ACC

specimens exhibited ☐H2AX and RPA foci (Figure 5F, Supplementary Figure S7,

Supplementary Table S3). We next sought to evaluate DNA damage in the human NCI-

H295R ACC cell line to establish an *in vitro* model for further mechanistic studies and found

☐H2AX foci indicating a cellular response to DNA damage in the absence of cytotoxic

treatments (Figure 6A, Supplementary Figure S8).

To assess for potential therapeutic vulnerabilities within the DNA damage and replication

stress response in ACC, we assessed sensitivity of NCI-H295R to small molecule inhibitors of

ATR kinase, elimusertib and VX-803, and of ATM kinase, AZ32 and KU60019. NCI-H295R

cells were sensitive to both ATR inhibitors with IC_{50} values ranging from 23 to 42 nM (Figure

6B). In contrast, ATM inhibition with AZ32 and KU60019 had minimal cytotoxic effects, even at

concentrations of 10,000 nM (Figure 6C). Prior studies have identified IC50 values ranging

from 100 to 300 nM and 500 to 20,000 nM for ATR monotherapy and ATM monotherapy

respectively in cell lines for other solid tumors.[66–69] ATR is recruited to sites of replication

stress and enacts downstream signaling via phosphorylation of specific targets to arrest the

cell cycle and ensure repair of damaged replication forks.[70] Therefore, we next assessed the

response of NCI-H295R cells to inhibition of enzymes in the DNA repair response

downstream of ATR, including checkpoint kinase 1 (CHK1, a direct target of ATR)[70] and

Rad51, a key regulator of homologous recombination that is not a direct ATR target.[71]

Inhibition of CHK1 with rabusertib or PF47736 resulted in a modest effect on viability with IC_{50}

values between 1000 to 10,000 nM (Figure 6D), which is higher than previously reported IC_{50}

values in cell lines considered sensitive to CHK1 monotherapy.[69,72,73] Inhibitors of Rad51

including RI-1 (specifically inhibits Rad51) and SAHA (HDAC inhibitor which inhibits Rad51),

had minimal effect on viability except at highest concentrations (50000 nM for RI-1 and 10000

nM for SAHA, Figure 6E). Finally, inhibition of poly-ADP-ribose polymerase (PARP), a DNA

repair protein critical for single strand break repair and base excision repair pathways, with

olaparib had no effect on NCI-H295R viability (Figure 6F). Prior studies have identified IC_{50}

values for olaparib ranging from 100 to 20000 nM in sensitive cell lines.[74,75] Together, these

data indicate the presence of replication stress and specific sensitivity to ATR inhibition in an

ACC cell line.

PTOs are 3-D cultures comprised of multiple cell types that can recapitulate the heterogenetic

nature of the intratumoral environment and can model many aspects of tumor complexity *in*

vitro.[76,77] To further assess the role of replication stress in ACC tumorigenesis, we generated

hormonally active ACC PTOs from patient 5 (**Table 1**, **OSU1**) and a second ACC patient with

biochemical evidence of hypercortisolism (**OSU2**). Neither patient received neoadjuvant

cytotoxic therapies prior to surgery. Both ACC PTO lines proliferated long-term in culture,

expressed high levels of the adrenal cortex marker SF1, and produced cortisol (Figures 7A-

C). Similar to the NCI-H295R cell line, both untreated ACC PTO lines demonstrated □H2AX

foci (Figure 7B, Supplementary figure S9). To determine if ACC PTOs exhibit sensitivity to

ATR inhibition, we treated the ACC PTOs with elimusertib and VX-803, which reduced viability

at IC_{50} values of 139 to 326 nM and 2000 to 3000 nM, respectively (Figure 7D). We also

treated the ACC PTOs with the ATM inhibitors, AZ32 and KU60019, neither of which caused

any appreciable reduction in viability (Figure 7D). These findings reinforce a critical role of

ATR in ACC and further indicate a potential role for ATR inhibition in treatment of ACC.

DISCUSSION

In this study, we use snRNA sequencing to demonstrate changes in immune, fibroblast and

adrenal cortex landscape in PACC and MACC specimens relative to non-malignant adrenal

cortex tissues. As such, we have identified several potential therapeutic approaches for further

investigation in ACC clinical management, as outlined below. Immunotherapy options for ACC

have been characterized by poor responses.[7] Our data indicate that macrophages are the

dominant immune cell and that SPP1+ macrophages are increased in ACC. Recent studies

demonstrate that SPP1+ macrophages are associated with tumor progression and worse

clinical outcomes.[32,33,36,78,79] In addition, ACC macrophages expressed markers of

immunosuppression. Collectively, these data suggest targeting macrophages with antibodies

or small molecules, such as blockade of the CD47-SIRPα axis or anti-PDL1 antibodies[38], to

induce activation and tumor cell phagocytosis may be a viable immunotherapy strategy in
ACC. In contrast, we found lymphoid infiltration only in the absence of excess cortisol
production, suggesting that immunotherapy alone may be more effective in non-functional
ACCs and that concurrent cortisol blockade may improve effectiveness of immunotherapy in
the presence of hypercortisolism.

Interestingly, our analysis of fibroblasts showed a loss of fibroblast subpopulations in ACC. In
addition, the same fibroblast population was present in primary and metastatic ACC. These
findings suggest that this fibroblast subpopulation may migrate from the primary tumor to the
site of metastasis and serve as a premetastatic niche to support tumor cells as in other
cancers.[80] Fibroblasts in ACC expressed increased *FAP* and *PDGFRB*. Targeting FAP and
PDGFRB resulted in tumor reduction in preclinical studies of other cancers.[81–83] Further
studies are necessary to elucidate the role of fibroblasts in ACC tumor progression and
identify potential therapeutic targets.

We found that ACC was characterized by a unique adrenal cortex subpopulation defined by a
signature for replication stress and DNA damage response pathways. This subpopulation may
correspond to the ACC M subpopulation identified by Tourigny et al.[16] Although Tourigny et al.
identified a signature for mitosis in the ACC M subpopulation, they did not identify a signature
for replication stress or DNA repair. Discrepancies in study findings may be attributed to
differences in sequencing methodologies and patient population. While Tourigny et al.[16]
predominantly included early-stage tumors, which can provide insight into the developmental
origin of ACC, our study included stage 4 tumors, which may provide insights for developing
targeted therapies. Other studies have established that genes implicated in DNA damage and
cell cycle regulation are overexpressed in ACC and are associated with worse overall survival
in patients.[8,84] In addition, Gara et al. have shown that *APOBEC3B,* which is known to induce
DNA damage, is overexpressed in ACC tumors with *TP53* mutations.[85] Prior work has also
shown that metastatic ACC exhibits a higher mutation burden compared to primary tumors,[86]
indicative of genomic instability. In addition to identifying replication stress signature using
snRNAseq data, our work mechanistically evaluated these pathways in ACC.

Cancer cells experience chronic replication stress due to oncogene activation and sustained
proliferative signaling. Furthermore, many chemotherapeutic agents induce replication stress
by depleting nucleoside pools required for DNA replication or causing direct DNA damage in

tumor cells.[88] ATR is a primary upstream orchestrator of the replications stress response
pathway. By inducing G2-S phase cell arrest and DNA repair as well as replication fork
stabilization, ATR allows cells to adapt to and prevent replication-stress induced cell
death.[70,89,90]

Our findings suggest that ATR can be therapeutically targeted in ACC irrespective of cortisol
secretion. ATR inhibitors drive cells with increased DNA damage to prematurely enter mitosis,
resulting in cell death. In addition to preclinical studies, some ATR inhibitors have also been
investigated in clinical trials as single agents or in conjunction with other therapeutic
modalities, such as radiotherapy or chemotherapy. In advanced solid tumors, ATR inhibitors
as monotherapy are well-tolerated and have resulted in partial or complete response.[93,94]
Other preclinical studies and clinical trials show ATR inhibitors and chemotherapeutic agents
can work synergistically to show efficacy in triple negative breast cancer, melanoma, platinum-
resistant high-grade serous ovarian cancer, and other advanced solid tumors.[91,94–99] Thus, by
demonstrating ATR-dependent replication stress in ACC, we identify a promising novel
treatment strategy for this deadly disease.

Identifying biomarkers to predict ATR sensitivity in ACC can improve potential efficacy. In non-
small cell lung cancer (NSCLC), loss of BRG1, a chromatin remodeling enzyme, was
identified as a biomarker for efficacy of ATR inhibition.[100] Loss of ARID1A, genomic instability,
ATM/G1 pathway abnormalities, and high tumor inflammation have also been identified as
potential biomarkers of sensitivity to ATR inhibition.[92] Although some preclinical models,
including chronic lymphocytic leukemia, have demonstrated ATR sensitivity in conjunction with
TP53 mutation, this relationship was not seen in the aforementioned clinical trial of ATR
inhibition in solid tumors.[102–104] Importantly, *TP53* inactivating mutations are present in 21-33%
of ACC,[8,105] so future work elucidating the role of p53 inactivation or other biomarkers in the
context of ATR sensitivity are necessary.

CHK1 is a target of ATR that plays a prominent role in the ATR signaling pathway by
activating checkpoints, delaying cell cycle progression, and enabling DNA repair to
proceed.[89,106] Our data show that the NCI-H295R cell line and ACC PTOs exhibit notable
sensitivity to ATR inhibition and modest sensitivity to CHK1 inhibition. These data suggest that
ATR-dependence in ACC may in part occur via alternative downstream targets, such as
SMARCAL1, which functions to promote replication fork stability and fork restart.[107] Additional

non-CHK1 targets of ATR, which are implicated in the replication stress response, include
WRN and FANCI.[108,109] Further studies are needed to determine if these targets play a role in
ACC tumor progression.
Taken together, our results demonstrate diversity of transcriptional signatures in ACC and
implicate several possible strategies for targeted treatment of ACC. Moreover, our results
exemplify how elucidation of cellular heterogeneity combined with mechanistic evaluation in
PTOs might direct therapeutic decisions and improve patient outcomes in ACC and other rare
cancers.
405
3899/4000 words

METHODS

Sample collection

Patients were consented for collection of specimens and clinical information using a study
protocol approved by The Ohio State University Institutional Review Board. Normal adrenal
and adrenal adenoma (AA) specimens were collected from patients diagnosed with adrenal
hypercortisolism and included one normal adrenal-adrenal adenoma pair. Primary (PACC)
and metastatic ACC (MACC) samples were collected from patients diagnosed with ACC and
included one tumor-metastasis pair. After collection, samples were frozen for further
processing. We performed snRNAseq as our samples included previously banked frozen
samples.[110] Somatic and germline mutations are reported for patients who underwent genetic
testing using Tempus (Chicago, IL), Foundation Medicine (Boston, MA), or CustomNext-
Cancer testing from Ambry Genetics (Aliso Viejo, CA). Sample details are summarized in
Table 1.

Single nuclei RNA sequencing

Nuclei isolation and library preparation
Nuclei were isolated from frozen tissue as described in Lacar et al. and sorted using the
Bigfoot Spectral Cell Sorter (Thermo Fisher Scientific, Waltham, MA). For each sample,
16,000 nuclei were sorted directly into the 10X Genomics master mix to load approximately
10,000 nuclei into a 10x Genomics Chromium device for microfluidic-based capture of single
nuclei.[111] Reverse transcription, cDNA amplification and library preparation were performed
according to the manufacturer's protocol for Chromium Next GEM Single Cell 3' Gene
Expression kit. Resulting libraries were sequenced on an Illumina NovaSeq 6000 instrument
to generate paired-end sequencing data.

Data preprocessing
Fastq files were generated using the mkfastq command from 10x Genomic CellRanger.
Alignment to the GRCh38 human reference, filtering, barcode counting, and UMI counting
steps were performed using the 10x Genomics CellRanger software suite v7.0 following the
default parameters.

Quality control, filtering, and doublet removal

Downstream analyses were performed using Seurat v4 for R 4,1,1 and 4.2.2.[19] For quality

control purposes, individual objects were filtered to remove nuclei that contained more than

5% of mitochondrial transcripts. In addition, data were filtered to only include nuclei that

contained a minimum of 200 detected genes for all the samples and a maximum of 8000

genes (for Normal and AA 1 samples), a maximum of 7000 genes (for AA 2 sample), a

maximum of 13000 genes (for PACC 1 sample), a maximum of 12000 genes (for PACC 2

sample), a maximum of 9000 genes (for PACC 3, PACC 4, and MACC 4 samples), a

maximum of 15000 genes (for MACC 1 sample), a maximum of 11500 genes (for MACC 2

sample), and a maximum of 14000 (for MACC 3 sample). Upper filtering cutoffs were selected

based on differences in data distribution between samples. As a result of filtering, the

following numbers of nuclei were excluded from further analysis: 17 nuclei (Normal sample),

49 nuclei (AA 1 and MACC 1 samples), 29 nuclei (AA 2 sample), 70 nuclei (PACC 1 sample),

381 nuclei (PACC 2 sample), 328 nuclei (PACC 3) sample, 171 nuclei (PACC 4 sample), 31

nuclei (MACC 2 sample), 553 nuclei (MACC 3 sample), and 5 nuclei (MACC 4 sample).

Doublets were detected and removed from individual objects using the DoubletFinder

package v2.0 for R.[112]

Normalization and cell cycle regression

To control for cell-cycle effects on transcriptional programs, we performed cell cycle

regression. After DoubletFinder, cell cycle scores were calculated for the individual objects

using the CellCycleScoring() function based on the list of cell cycle marker genes published in

Tirosh et al.[113] Normalization and variance stabilization were performed on the individual

objects using the SCTransform() function from the sctransform package v0.3 for R with

"vst.flavor" argument set to v2 and "vars.to.regress" argument set to regress the percentage of

mitochondrial content, G2/M and S cell cycle scores.[114]

Data integration

To minimize patient-to-patient variability, samples were integrated using the RPCA algorithm

in Seurat.[115]

Dimensionality reduction and clustering

Principal component analysis (PCA) was performed on the integrated Seurat object. The first

30 principal components were used for dimensionality reduction via the Uniform Manifold

Approximation and Projection (UMAP) method. Clusters were defined using the FindClusters()

function for a series of resolutions between 0.1 and 1. Optimal clustering resolution of 0.1 was

selected based on cluster stability, as in Long et al., using the clustree package v0.5 in

R.[116,117]

Cell type annotation

FindAllMarkers() function in Seurat was used to define marker genes for further cluster

annotation. Dot plots were generated using the DotPlot() function in Seurat with the argument

"scale" set to the default value. To complement the FindAllMarkers() approach and annotate

adrenal cortex clusters more precisely, we used the AddModuleScore() function in Seurat to

calculate module scores based on the expression of *NR5A1* (which encodes the protein

steroidogenic factor 1, SF1*), INHA*, and *MLANA*, which are all adrenal cortex biomarkers.[17,18]

Adrenal cortex clusters were characterized by higher adrenal cortex biomarker module scores

when compared to lymphocyte, myeloid cell, endothelial, and fibroblast clusters.

Subclustering of immune cells

Myeloid- and lymphocyte-typed clusters were subset from the initial integrated Seurat object

and re-processed using SCTransform as described above. Samples were then merged and

re-integrated using Harmony[26] at a resolution of 0.2. Dimensionality reduction and clustering

steps were followed as described above. This process was repeated after non-immune cells

were subset out. ggplot2[118] was used to generate a histogram of immunocyte proportions and

the FindAllMarkers() function in Seurat was used to define marker genes for cluster

annotations displayed using the DotPlot() function. The macrophage-typed clusters were then

further subset, re-processed, and re-integrated using Harmony at a resolution of 0.2. This

process was repeated after excluding a small number of CD3-expressing lymphocytes.

Dimensionality reduction, clustering steps, and cell type annotation were performed as

described above. ggplot2[118] was used to generate a histogram of macrophage subtypes for

each sample type (normal adrenal, AA, PACC, MACC). The function propeller() from the

speckle[119] package for R was used to compare subtype proportions across sample types.

False discovery rate (FDR) is reported. Average gene expression was calculated across each

sample type and scaled average expression was displayed in a heatmap using the

DoHeatmap() function. There were not enough cells for the CD8+ T cell, CD4+ T cell, and

natural killer (NK) cell clusters to subset and reprocess, re-integrate, and recluster. Therefore,

these lymphocytes were subset from the macrophage and leukocyte re-integrated object and

average gene expression was calculated across each sample type and displayed in a
heatmap as described above.
511

Subclustering of fibroblasts
The fibroblast cluster was subset from the initial integrated Seurat object, reprocessed, and
re-integrated using Harmony to increase clustering resolution. After dimensionality reduction
and clustering, optimal resolution of 0.2 was selected as described above. As a result, 10
clusters were identified. Four clusters including adipocytes were excluded, after which the
fibroblasts were reprocessed and re-integrating using Harmony. PACC 2 was excluded from
analysis due to having only 5 fibroblasts. After dimensionality reduction and clustering, optimal
resolution of 0.1 was selected as described above. As a result, 5 clusters of fibroblasts were
identified. UMAP projection was displayed using the DimPlot() function in Seurat. Marker
genes were identified using the FindAllMarkers() function and visualized using the DotPlot()
function in Seurat. Average gene expression was calculated across sample types and
displayed in a heatmap as described above.
524

Subclustering of adrenal cortex cells
To increase clustering resolution, adrenal cortex cells were subset from the integrated Seurat
object and reprocessed using normalization, cell-cycle regression, integration using RPCA,
dimensionality reduction, and clustering steps as in processing the initial Seurat object.
Optimal clustering resolution of 0.1 was selected as described above. FindAllMarkers()
function was used to identify marker genes for each of the clusters. Marker genes were
visualized using the DotPlot() function as described above. In addition, adrenal cortex module
scores were calculated for each of the clusters as described above. As a result, 6 adrenal
cortex clusters and 1 cluster of adrenal medulla were identified. To further increase clustering
resolution, the adrenal medulla cluster was excluded, and adrenal cortex cells were again
subset from the integrated Seurat object and reprocessed using normalization, cell-cycle
regression, integration, dimensionality reduction, and clustering steps as described above.
Optimal resolution of 0.1 was selected as described above. As a result, 7 clusters of adrenal
cortex cells were identified; these cluster assignments were used for all downstream analyses.
539

540 *Cell type proportions*
541 To compare cell type proportions of adrenal cortex cells across the samples, we used the
542 propeller method from the speckle package v0.0.3 in R.[119]

Pathway analysis

To characterize adrenal cortex clusters based on gene expression patterns, we used the AddModuleScore() function from the Seurat package to calculate module scores using a selection of Hallmark pathways downloaded from MSigDB using the msigdbr package v7.5 in R.[59,60] We further calculated average module scores for each of the clusters to generate the heatmap for Figure 4C. We also calculated average module scores for each of the clusters on a per-tumor type basis to generate the heatmap for Supplementary Figure S4. We performed a similar analysis using a selection of replication stress and DNA damage response REACTOME pathways downloaded from MsigDB[59–61] We calculated module scores as in Wu et al.[35] for each of the selected REACTOME pathways, and then calculated average module scores for each cluster on a per-tumor type basis (for Figure 5A) and on a per-sample basis (for supplementary Figure S5), which were displayed on a heatmap. To generate heatmaps, we used the pheatmap() function from the pheatmap package v.1.0 in R with the argument "scale" set to "row".[120] For all heatmaps, clusters with fewer than 50 cells were excluded from analysis. Additionally, we used QIAGEN Ingenuity Pathways Analysis (IPA) (QIAGEN Inc., https://digitalinsights.qiagen.com/IPD) to carry out pathway enrichment analysis for Figure 5B.[62]

InferCNV

We used inferCNV[64] v1.8 to infer copy number alterations in the tumor samples. We used the HMM_CNV_predictions.HMMi6.hmm_mode-samples.Pnorm_0.5.pred_cnv_regions.datfile to quantify the number of chromosomal aberrations per sample. To compare the number of chromosomal aberrations between clusters, we used the chisq.test() function in R.

Data and code availability

snRNAseq data generated for this study has been deposited to dbGAP, accession phs003764. Code that was used to analyze the snRNAseq data is available upon request.

Immunofluorescence of ACC

Hematoxylin and eosin–stained slides of formalin fixed paraffin embedded (FFPE) samples were screened for diagnosis, cellularity, and necrosis by a board-certified pathologist. FFPE blocks were cut into 4-µm sections and deparaffinized in organic solvents. Slides were

dehydrated, submerged in DAKO Antigen Retrieval Buffer (pH 9.0) (Agilent Technologies,
Santa Clara, CA), and incubated at 110°C in a steam rice cooker for 30 minutes. The slides
were then cooled on ice for 30 minutes, followed by 5 minutes in distilled water. As previously
described[121], samples were permeabilized with DAKO Wash Buffer (Agilent Technologies),
then blocked with blocking buffer (DAKO Wash buffer; 1% BSA). Sections were stained with
mouse anti-□H2AX (Millipore-Sigma, St.Louis, MO, 05-636) at 1:500 and rabbit anti-RPA
(Bethyl Laboratories, Montgomery, TX, A300-245A-M) at 1:200. Secondary incubation was
performed with goat anti-mouse Alexa fluor 488 and goat anti-rabbit Alexa fluor 647. DAPI
(Millipore-Sigma) was added, and the slides were dehydrated with increasing concentrations
of ethanol. Samples were then mounted with ProLong Gold anti-fade reagent (Thermo Fisher
Scientific) and stored at −20°C. Stained samples were imaged on a Leica Thunder Imager
Microscope.
589
590 **Cell culture**
591
NCI-H295R
The NCI-H295R ACC cell line was originally purchased from American Type Culture
Collection (ATCC, Manassas, VA). NCI-H295R cells were grown in DMEM/F12 with L-
glutamine and 15 mM HEPES without phenol red (Thermo Fisher Scientific) supplemented
with 2.5% Nu-Serum™ (Corning, Corning, NY), 1% penicillin-streptomycin (Sigma-Millipore),
and 1x ITS Premix Universal Cell Culture Supplement (Corning). Cells were split
approximately once per week and all experiments utilized cells below 20 passages.
599
ACC patient-derived tumor organoid (PTO) generation
PTOs were generated from a tumor harvested from a patient diagnosed with ACC (patient 5 in
Table 1). Briefly, tumor tissue was minced then digested with 0.5 mg/mL collagenase II
(Worthington Biochemical, Freehold, NJ) in DMEM/F12 (Cytiva, Marlborough, MA). Tumor
cells were then seeded at $5\text{-}10\text{x}10^4$ cells per well in 24-well plates in 80% Matrigel® (Corning),
with DMEM/F12 supplemented with 10 nM HEPES (Cytiva), 1x Glutamax, 1x B27 Supplement
(both Thermo Fisher Scientific), 1.25 mM n-acetylcysteine, 10 mM nicotinamide, 10 nM
[Leu15]-Gastrin, 0.5 μM A-83-01, 1 μM PGE2 (all Sigma-Millipore), 0.5 μg/mL R-Spondin-1,
50 ng/mL EGF (both R&D Systems, Minneapolis, MN), 0.1 μg/mL FGF-10, 30 ng/mL
recombinant murine Wnt3a, and 0.1 μg/mL recombinant human Noggin (all Thermo Fisher

Scientific). PTOs were routinely passaged upon confluency. Brightfield images of organoid
colonies were captured using an EVOS™ M5000 microscope (Thermo Fisher Scientific).

Immunofluorescence of cultured cells and ACC PTOs

NCI-H295R cells were seeded at 1.5 x 10^5 cells per cm^2 on top of 12 mm diameter poly-L-
lysine coated glass coverslips (Electron Microscopy Sciences, Hatfield, PA) placed in wells of
a 24-well cell culture plate. After 24 hours, cells were washed with PBS, fixed with 4%
paraformaldehyde (Thermo Fisher Scientific), and permeabilized with 0.5% Triton X-100. After
blocking with 10% normal goat serum (Abcam, Cambridge, UK) in 0.1% Triton X-100, cells
were stained with mouse anti-γH2AX (Cell Signaling Technologies, Danvers, MA,
CST#80312) at 1:250 followed by secondary incubation with goat anti-mouse Alexa Fluor$^®$
488 antibody at 1:500 (Abcam #ab150117) and phalloidin-Alexa Fluor$^®$ 594 at 1:1000 (Abcam
#ab176757). After mounting with Fluoroshield mounting medium with DAPI (Abcam), images
were obtained on a Nikon AXR confocal microscope (Nikon, Tokyo, Japan). Cells incubated
without primary antibodies were used as negative controls for NCI-H295R
immunofluorescence experiments.

ACC PTOs were removed from Matrigel domes and stained in a similar manner, and then
mounted on the 0.2-0.4 μm double cavity well microscope slides (Electron Microscopy ciences
#71883-79) and imaged similarly to NCI-H295R cells. Primary antibodies were incubated at
the following concentrations overnight at 4C: rabbit anti-SF-1 1:250 (Abcam #ab51074),
mouse anti-γH2AX 1:250 (Cell Signaling Technologies CST#80312), or equivalent isotype
controls (Abcam #ab172730, Cell Signaling Technologies CST#5415). The following
secondary antibodies were used: goat anti-rabbit Alexa Fluor$^®$ 488 1:1000 (Abcam
#ab150081), goat anti-mouse Alexa Fluor$^®$ 546 1:500 (Thermo Fisher Scientific #A-11003)
phalloidin-Alexa Fluor$^®$ 647 1:400 (Thermo Fisher Scientific #A2287). Comparable isotype
antibodies (Cell Signaling Technologies #5145 and Abcam #ab172730) were used as
negative controls for ACC PTO immunofluorescence experiments.

Cytotoxicity assays

NCI-H295R Cytotoxicity Assays

NCI-H295R cells were seeded at 1×10^4 cells per well in a 96-well plate and allowed to adhere

for 24 hours. Media was then replaced with fresh media containing specified drugs or

equivalent DMSO vehicle control (Corning). Cell viability was assessed by water-soluble

tertozolium-8 (WST-8) based Cell Counting Kit-8 assay (Dojindo, Mashiki, Kumamoto, Japan)

per the manufacturer's instructions at 4 days. Absorbance readings were conducted using a

SpectraMax® iD5 plate reader and SoftMax® Pro Software (Molecular Devices, San Jose, CA).

All assays were performed in biologic triplicates with 3-5 replicates per condition. The

following drugs were used at concentrations ranging from 1nM to 50uM: KU60019 (Selleck

Chemicals, Houston, TX), RI-1 (Thermo Fisher Scientific), suberoylanilide hydroxamic acid

(SAHA, aka vorinostat, Thermo Fisher Scientific), rabusertib (Selleck Chemicals), PF47736

(Selleck Chemicals), olaparib (Cell Signaling Technologies). Elimusertib (Selleck Chemicals),

VX-803 (Selleck Chemicals), and AZ32 (Thermo Fisher Scientific). Mean viability of each drug

treatment group was measured as a percentage of baseline DMSO vehicle treated cell

absorbance. The half-maximal inhibitory concentration (IC_{50}) for each drug was calculated by

performing a four-parameter nonlinear best fit analysis of log concentration by percent viability

on GraphPad Prism® 7.0a software (GraphPad Software, La Jolla, CA). Differences in viability

across each concentration of a particular drug were compared using one-factor ANOVA with

post-hoc multiple comparison analysis by Dunnett's test in GraphPad Prism® 7.0a software

(GraphPad Software). All data are reported as means ± SEM with statistical significance set

as $p < 0.05$.

ACC PTO cytotoxicity assays

Single-cell suspensions were generated by incubating well-established ACC organoid domes

with 1 mg/mL dispase II (Thermo Fisher Scientific). 1.5×10^4 cells per well were seeded in 96-

well plates in 80% Matrigel® as described above. Three days after seeding, media was

replaced with media containing specified drugs or DMSO vehicle control. Organoid viability

was assessed at 4 days by WST-8 assay as described above using the cell free Matrigel®

domes for background absorbance readings. Differences in viability across each

concentration of a drug were compared using one-factor ANOVA with post-hoc multiple

comparison analysis by Dunnett's test in GraphPad Prism® 7.0a software (GraphPad

Software, La Jolla, CA). Each drug was tested in at least 3 independent experiments with 5

replicates per experiment.

Cortisol ELISA

Single cell suspensions of NCI-H295R cells and ACC organoid-derived cells were generated

as described above. 3×10^{5} cells were plated and medium was collected at 72-hours. Cortisol

was measured by ELISA per manufacturer's instructions (Abcam #ab108665). Each

experiment was conducted in triplicate and read on BioTek Synergy H1 microplate reader

using Gen5 v3.10 software (Agilent Technologies). Cortisol concentrations were compared

between groups using two-tailed independent student's t-test. All data are reported as

means ± SEM with statistical significance set was $p < 0.05$.

AUTHOR CONTRIBUTION STATEMENT

Conceptualization: P.H.D.

Methodology: P.H.D. and K.E.M.

Investigation: L.V.P., E.A.R.G., D.M.C., J.B.N., A.R., Y.S., and E.L.

Specimen Acquisition: P.H.D., J.E.P., and B.S.M.

Specimen Pathology Review: S.S.

Formal analysis: L.V.P., E.A.R.G., D.M.C., Y.S., E.L.

Data curation: L.V.P., E.A.R.G., and K.E.M.

Funding acquisition: P.H.D.

Project administration: P.H.D.

Supervision: E.R.M., K.E.M. and P.H.D.

Validation: K.E.M. and P.H.D.

Visualization: L.V.P., P.H.D.

Writing—original draft: L.V.P., E.A.R.G., and D.M.C.

Writing—review and editing: L.V.P., E.A.R.G., D.M.C., B.S.M., M.M., K.E.M., and P.H.D.

ACKNOWLEDGMENTS

This project was supported by Ohio Cancer Research, The Victory Over Cancer Foundation,

DOD W81XWH-22-PRCRP-CDA-SO, NIH R01CA279997, and NIH R21CA277083 (PHD).

LVP is supported by the Pelotonia Scholars Program. DMC is supported by training grant PF-

23-1036284-01-CDP via the American Cancer Society. Any opinions, findings, and

conclusions expressed in this material are those of the author(s) and do not necessarily reflect

those of the NIH, Pelotonia Scholars Program, OSU, or the American Cancer Society. We

acknowledge resources from the Campus Microscopy and Imaging Facility (CMIF),

Biospecimen Services Shared Resource and the Microscopy Shared Resource. These shared

resources are supported in part by the Cancer Center Support Grant P30 CA016058, National

Cancer Institute, Bethesda, MD. We also acknowledge Matthew Ringel, MD, Gary Hammer,

MD, PhD, Timothy Frankel, MD, PhD, Wayne Miles, PhD, and Abby Green, MD, PhD for

insightful discussions.

REFERENCES

1. Else T, Williams AR, Sabolch A, Jolly S, Miller BS, Hammer GD. Adjuvant therapies and patient and tumor characteristics associated with survival of adult patients with adrenocortical carcinoma. *J Clin Endocrinol Metab*. 2014;99(2):455-461. doi:10.1210/jc.2013-2856

2. Else T, Kim AC, Sabolch A, et al. Adrenocortical carcinoma. *Endocr Rev*. 2014;35(2):282-326. doi:10.1210/er.2013-1029

3. Bilimoria KY, Shen WT, Elaraj D, et al. Adrenocortical carcinoma in the United States: treatment utilization and prognostic factors. *Cancer*. 2008;113(11):3130-3136. doi:10.1002/cncr.23886

4. Vaidya A, Nehs M, Kilbridge K. Treatment of Adrenocortical Carcinoma. *Surg Pathol Clin*. 2019;12(4):997-1006. doi:10.1016/j.path.2019.08.010

5. Kenney L, Hughes M. Adrenocortical Carcinoma: Role of Adjuvant and Neoadjuvant Therapy. *Surg Oncol Clin N Am*. 2023;32(2):279-287. doi:10.1016/j.soc.2022.10.005

6. Fassnacht M, Terzolo M, Allolio B, et al. Combination chemotherapy in advanced adrenocortical carcinoma. *N Engl J Med*. 2012;366(23):2189-2197. doi:10.1056/NEJMoa1200966

7. Araujo-Castro M, Pascual-Corrales E, Molina-Cerrillo J, Alonso-Gordoa T. Immunotherapy in Adrenocortical Carcinoma: Predictors of Response, Efficacy, Safety, and Mechanisms of Resistance. *Biomedicines*. 2021;9(3):304. doi:10.3390/biomedicines9030304

8. Zheng S, Cherniack AD, Dewal N, et al. Comprehensive Pan-Genomic Characterization of Adrenocortical Carcinoma. *Cancer Cell*. 2016;29(5):723-736. doi:10.1016/j.ccell.2016.04.002

9. Assié G, Letouzé E, Fassnacht M, et al. Integrated genomic characterization of adrenocortical carcinoma. *Nat Genet*. 2014;46(6):607-612. doi:10.1038/ng.2953

10. McGranahan N, Swanton C. Clonal Heterogeneity and Tumor Evolution: Past, Present, and the Future. *Cell*. 2017;168(4):613-628. doi:10.1016/j.cell.2017.01.018

11. Lei Y, Tang R, Xu J, et al. Applications of single-cell sequencing in cancer research: progress and perspectives. *J Hematol OncolJ Hematol Oncol*. 2021;14(1):91. doi:10.1186/s13045-021-01105-2

12. Bischoff P, Trinks A, Wiederspahn J, et al. The single-cell transcriptional landscape of lung carcinoid tumors. *Int J Cancer*. 2022;150(12):2058-2071. doi:10.1002/ijc.33995

13. Pu W, Shi X, Yu P, et al. Single-cell transcriptomic analysis of the tumor ecosystems
underlying initiation and progression of papillary thyroid carcinoma. *Nat Commun*.
2021;12(1):6058. doi:10.1038/s41467-021-26343-3

14. Penny MK, Lerario AM, Basham KJ, et al. Targeting Oncogenic Wnt/β-Catenin Signaling
in Adrenocortical Carcinoma Disrupts ECM Expression and Impairs Tumor Growth.
Cancers. 2023;15(14):3559. doi:10.3390/cancers15143559

15. Wang Q, Sun N, Meixner R, et al. Metabolic heterogeneity in adrenocortical carcinoma
impacts patient outcomes. *JCI Insight*. 2023;8(16):e167007. doi:10.1172/jci.insight.167007

16. Tourigny DS, Altieri B, Secener KA, et al. Cellular landscape of adrenocortical carcinoma
at single-nuclei resolution. *Mol Cell Endocrinol*. 2024;590:112272.
doi:10.1016/j.mce.2024.112272

17. Sbiera S, Schmull S, Assie G, et al. High diagnostic and prognostic value of steroidogenic
factor-1 expression in adrenal tumors. *J Clin Endocrinol Metab*. 2010;95(10):E161-171.
doi:10.1210/jc.2010-0653

18. Mizdrak M, Tičinović Kurir T, Božić J. The Role of Biomarkers in Adrenocortical
Carcinoma: A Review of Current Evidence and Future Perspectives. *Biomedicines*.
2021;9(2):174. doi:10.3390/biomedicines9020174

19. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data.
Cell. 2021;184(13):3573-3587.e29. doi:10.1016/j.cell.2021.04.048

20. Al Barashdi MA, Ali A, McMullin MF, Mills K. Protein tyrosine phosphatase receptor
type C (PTPRC or CD45). *J Clin Pathol*. 2021;74(9):548-552. doi:10.1136/jclinpath-2020-
206927

21. Broz ML, Krummel MF. The emerging understanding of myeloid cells as partners and
targets in tumor rejection. *Cancer Immunol Res*. 2015;3(4):313-319. doi:10.1158/2326-
6066.CIR-15-0041

22. Li D, Xia L, Huang P, et al. Cancer-associated fibroblast-secreted IGFBP7 promotes
gastric cancer by enhancing tumor associated macrophage infiltration via
FGF2/FGFR1/PI3K/AKT axis. *Cell Death Discov*. 2023;9(1):17. doi:10.1038/s41420-023-
01336-x

23. Sakamoto H, Koma YI, Higashino N, et al. PAI-1 derived from cancer-associated
fibroblasts in esophageal squamous cell carcinoma promotes the invasion of cancer cells
and the migration of macrophages. *Lab Investig J Tech Methods Pathol*. 2021;101(3):353-
368. doi:10.1038/s41374-020-00512-2

24. Sawano A, Iwai S, Sakurai Y, et al. Flt-1, vascular endothelial growth factor receptor 1, is
a novel cell surface marker for the lineage of monocyte-macrophages in humans. *Blood*.
2001;97(3):785-791. doi:10.1182/blood.v97.3.785

25. Lopes da Silva M, Cutler DF. von Willebrand factor multimerization and the polarity of
secretory pathways in endothelial cells. *Blood*. 2016;128(2):277-285. doi:10.1182/blood-
2015-10-677054

26. Korsunsky I, Millard N, Fan J, et al. Fast, sensitive and accurate integration of single-cell
data with Harmony. *Nat Methods*. 2019;16(12):1289-1296. doi:10.1038/s41592-019-0619-
0

27. Prenen H, Mazzone M. Tumor-associated macrophages: a short compendium. *Cell Mol
Life Sci CMLS*. 2019;76(8):1447-1458. doi:10.1007/s00018-018-2997-3

28. Akinrinmade OA, Chetty S, Daramola AK, Islam MU, Thepen T, Barth S. CD64: An
Attractive Immunotherapeutic Target for M1-type Macrophage Mediated Chronic
Inflammatory Diseases. *Biomedicines*. 2017;5(3):56. doi:10.3390/biomedicines5030056

29. Martinez FO, Helming L, Milde R, et al. Genetic programs expressed in resting and IL-4
alternatively activated mouse and human macrophages: similarities and differences. *Blood*.
2013;121(9):e57-69. doi:10.1182/blood-2012-06-436212

30. Lu J, Ma L. The role of tumor-associated macrophages in the development, metastasis and
treatment of breast cancer. *Pathol Res Pract*. 2020;216(9):153085.
doi:10.1016/j.prp.2020.153085

31. Georgoudaki AM, Prokopec KE, Boura VF, et al. Reprogramming Tumor-Associated
Macrophages by Antibody Targeting Inhibits Cancer Progression and Metastasis. *Cell Rep*.
2016;15(9):2000-2011. doi:10.1016/j.celrep.2016.04.084

32. Nasir I, McGuinness C, Poh AR, Ernst M, Darcy PK, Britt KL. Tumor macrophage
functional heterogeneity can inform the development of novel cancer therapies. *Trends
Immunol*. 2023;44(12):971-985. doi:10.1016/j.it.2023.10.007

33. Bill R, Wirapati P, Messemaker M, et al. CXCL9:SPP1 macrophage polarity identifies a
network of cellular programs that control human cancers. *Science*. 2023;381(6657):515-
524. doi:10.1126/science.ade2292

34. Cheng S, Li Z, Gao R, et al. A pan-cancer single-cell transcriptional atlas of tumor
infiltrating myeloid cells. *Cell*. 2021;184(3):792-809.e23. doi:10.1016/j.cell.2021.01.010

35. Wu SZ, Al-Eryani G, Roden DL, et al. A single-cell and spatially resolved atlas of human
breast cancers. *Nat Genet*. 2021;53(9):1334-1347. doi:10.1038/s41588-021-00911-1

36. Ozato Y, Kojima Y, Kobayashi Y, et al. Spatial and single-cell transcriptomics decipher
the cellular environment containing HLA-G+ cancer cells and SPP1+ macrophages in
colorectal cancer. *Cell Rep*. 2023;42(1):111929. doi:10.1016/j.celrep.2022.111929

37. Shinchi Y, Ishizuka S, Komohara Y, et al. The expression of PD-1 ligand 1 on
macrophages and its clinical impacts and mechanisms in lung adenocarcinoma. *Cancer
Immunol Immunother CII*. 2022;71(11):2645-2661. doi:10.1007/s00262-022-03187-4

38. Mantovani A, Allavena P, Marchesi F, Garlanda C. Macrophages as tools and targets in
cancer therapy. *Nat Rev Drug Discov*. 2022;21(11):799-820. doi:10.1038/s41573-022-
00520-5

39. Steele NG, Carpenter ES, Kemp SB, et al. Multimodal Mapping of the Tumor and
Peripheral Blood Immune Landscape in Human Pancreatic Cancer. *Nat Cancer*.
2020;1(11):1097-1112. doi:10.1038/s43018-020-00121-4

40. Zhang H, Liu L, Liu J, et al. Roles of tumor-associated macrophages in anti-PD-1/PD-L1
immunotherapy for solid cancers. *Mol Cancer*. 2023;22(1):58. doi:10.1186/s12943-023-
01725-x

41. Ji K, Zhang Y, Jiang S, et al. SIRPα blockade improves the antitumor immunity of
radiotherapy in colorectal cancer. *Cell Death Discov*. 2023;9(1):180. doi:10.1038/s41420-
023-01472-4

42. Zhang CX, Huang DJ, Baloche V, et al. Galectin-9 promotes a suppressive
microenvironment in human cancer by enhancing STING degradation. *Oncogenesis*.
2020;9(7):65. doi:10.1038/s41389-020-00248-0

43. Dong Y, Sun Q, Zhang X. PD-1 and its ligands are important immune checkpoints in
cancer. *Oncotarget*. 2017;8(2):2171-2186. doi:10.18632/oncotarget.13895

44. Thiel M, Wolfs MJ, Bauer S, et al. Efficiency of T-cell costimulation by CD80 and CD86
cross-linking correlates with calcium entry. *Immunology*. 2010;129(1):28-40.
doi:10.1111/j.1365-2567.2009.03155.x

45. Landwehr LS, Altieri B, Schreiner J, et al. Interplay between glucocorticoids and tumor-
infiltrating lymphocytes on the prognosis of adrenocortical carcinoma. *J Immunother*
Cancer. 2020;8(1):e000469. doi:10.1136/jitc-2019-000469

46. Friedlaender A, Addeo A, Banna G. New emerging targets in cancer immunotherapy: the
role of TIM3. *ESMO Open*. 2019;4(Suppl 3):e000497. doi:10.1136/esmoopen-2019-
000497

47. Thommen DS, Schumacher TN. T Cell Dysfunction in Cancer. *Cancer Cell*.
2018;33(4):547-562. doi:10.1016/j.ccell.2018.03.012

48. Liu PS, Chen YT, Li X, et al. CD40 signal rewires fatty acid and glutamine metabolism for
stimulating macrophage anti-tumorigenic functions. *Nat Immunol*. 2023;24(3):452-462.
doi:10.1038/s41590-023-01430-3

49. Zhang Q, Bi J, Zheng X, et al. Blockade of the checkpoint receptor TIGIT prevents NK
cell exhaustion and elicits potent anti-tumor immunity. *Nat Immunol*. 2018;19(7):723-732.
doi:10.1038/s41590-018-0132-0

50. Crinier A, Dumas PY, Escalière B, et al. Single-cell profiling reveals the trajectories of
natural killer cell differentiation in bone marrow and a stress signature induced by acute

myeloid leukemia. *Cell Mol Immunol*. 2021;18(5):1290-1304. doi:10.1038/s41423-020-
00574-8

51. Lendahl U, Muhl L, Betsholtz C. Identification, discrimination and heterogeneity of
fibroblasts. *Nat Commun*. 2022;13(1):3409. doi:10.1038/s41467-022-30633-9

52. Li S, Lu R, Shu L, et al. An integrated map of fibroblastic populations in human colon
mucosa and cancer tissues. *Commun Biol*. 2022;5(1):1326. doi:10.1038/s42003-022-
04298-5

53. Yang D, Liu J, Qian H, Zhuang Q. Cancer-associated fibroblasts: from basic science to
anticancer therapy. *Exp Mol Med*. 2023;55(7):1322-1332. doi:10.1038/s12276-023-01013-
0

54. Menezes S, Okail MH, Jalil SMA, Kocher HM, Cameron AJM. Cancer-associated
fibroblasts in pancreatic cancer: new subtypes, new markers, new targets. *J Pathol*.
2022;257(4):526-544. doi:10.1002/path.5926

55. Han C, Liu T, Yin R. Biomarkers for cancer-associated fibroblasts. *Biomark Res*.
2020;8(1):64. doi:10.1186/s40364-020-00245-w

56. Bedoya-Reina OC, Li W, Arceo M, et al. Single-nuclei transcriptomes from human adrenal
gland reveal distinct cellular identities of low and high-risk neuroblastoma tumors. *Nat
Commun*. 2021;12(1):5309. doi:10.1038/s41467-021-24870-7

57. Ye L, Su ZJ, Ge RS. Inhibitors of testosterone biosynthetic and metabolic activation
enzymes. *Mol Basel Switz*. 2011;16(12):9983-10001. doi:10.3390/molecules16129983

58. Neunzig J, Bernhardt R. Dehydroepiandrosterone sulfate (DHEAS) stimulates the first step
in the biosynthesis of steroid hormones. *PloS One*. 2014;9(2):e89727.
doi:10.1371/journal.pone.0089727

59. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. *Bioinforma Oxf Engl*. 2011;27(12):1739-
1740. doi:10.1093/bioinformatics/btr260

60. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The
Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst*.
2015;1(6):417-425. doi:10.1016/j.cels.2015.12.004

61. Gillespie M, Jassal B, Stephan R, et al. The reactome pathway knowledgebase 2022.
Nucleic Acids Res. 2022;50(D1):D687-D692. doi:10.1093/nar/gkab1028

62. Krämer A, Green J, Pollard J, Tugendreich S. Causal analysis approaches in Ingenuity
Pathway Analysis. *Bioinforma Oxf Engl*. 2014;30(4):523-530.
doi:10.1093/bioinformatics/btt703

63. Gaillard H, García-Muse T, Aguilera A. Replication stress and cancer. *Nat Rev Cancer*.
2015;15(5):276-289. doi:10.1038/nrc3916

64. Tickle T, Tirosh I, Georgescu C, Brown M, Haas B. inferCNV of the Trinity CTAT
Project. Published online 2019. https://github.com/broadinstitute/inferCNV

65. Yang SF, Nelson CB, Wells JK, et al. ZNF827 is a single-stranded DNA binding protein
that regulates the ATR-CHK1 DNA damage response pathway. *Nat Commun*.
2024;15(1):2210. doi:10.1038/s41467-024-46578-0

66. Wengner AM, Siemeister G, Lücking U, et al. The Novel ATR Inhibitor BAY 1895344 Is
Efficacious as Monotherapy and Combined with DNA Damage-Inducing or Repair-
Compromising Therapies in Preclinical Cancer Models. *Mol Cancer Ther*. 2020;19(1):26-
38. doi:10.1158/1535-7163.MCT-19-0019

67. Shu J, Wang X, Yang X, Zhao G. ATM inhibitor KU60019 synergistically sensitizes lung
cancer cells to topoisomerase II poisons by multiple mechanisms. *Sci Rep*. 2023;13(1):882.
doi:10.1038/s41598-023-28185-z

68. Scotto L, Serrano XJ, Zullo K, et al. ATM inhibition overcomes resistance to histone
deacetylase inhibitor due to p21 induction and cell cycle arrest. *Oncotarget*.
2020;11(37):3432-3442. doi:10.18632/oncotarget.27723

69. Wang C, Tang M, Chen Z, et al. Genetic vulnerabilities upon inhibition of DNA damage
response. *Nucleic Acids Res*. 2021;49(14):8214-8231. doi:10.1093/nar/gkab643

70. da Costa AABA, Chowdhury D, Shapiro GI, D'Andrea AD, Konstantinopoulos PA.
Targeting replication stress in cancer therapy. *Nat Rev Drug Discov*. 2023;22(1):38-58.
doi:10.1038/s41573-022-00558-5

71. Ito M, Fujita Y, Shinohara A. Positive and negative regulators of RAD51/DMC1 in
homologous recombination and DNA replication. *DNA Repair*. 2024;134:103613.
doi:10.1016/j.dnarep.2023.103613

72. Yang X, Pan Y, Qiu Z, et al. RNF126 as a Biomarker of a Poor Prognosis in Invasive
Breast Cancer and CHEK1 Inhibitor Efficacy in Breast Cancer Cells. *Clin Cancer Res Off*
J Am Assoc Cancer Res. 2018;24(7):1629-1643. doi:10.1158/1078-0432.CCR-17-2242

73. Corrales-Guerrero S, Cui T, Castro-Aceituno V, et al. Inhibition of RRM2 radiosensitizes
glioblastoma and uncovers synthetic lethality in combination with targeting CHK1. *Cancer*
Lett. 2023;570:216308. doi:10.1016/j.canlet.2023.216308

74. Fleury H, Malaquin N, Tu V, et al. Exploiting interconnected synthetic lethal interactions
between PARP inhibition and cancer cell reversible senescence. *Nat Commun*.
2019;10(1):2556. doi:10.1038/s41467-019-10460-1

75. Cahuzac M, Langlois P, Péant B, Fleury H, Mes-Masson AM, Saad F. Pre-activation of
autophagy impacts response to olaparib in prostate cancer cells. *Commun Biol*.
2022;5(1):251. doi:10.1038/s42003-022-03210-5

76. Tuveson D, Clevers H. Cancer modeling meets human organoid technology. *Science*.
2019;364(6444):952-955. doi:10.1126/science.aaw6985

77. Dedhia PH, Bertaux-Skeirik N, Zavros Y, Spence JR. Organoid Models of Human
Gastrointestinal Development and Disease. *Gastroenterology*. 2016;150(5):1098-1112.
doi:10.1053/j.gastro.2015.12.042

78. Pittet MJ, Michielin O, Migliorini D. Clinical relevance of tumour-associated
macrophages. *Nat Rev Clin Oncol*. 2022;19(6):402-421. doi:10.1038/s41571-022-00620-6

79. Wu J, Shen Y, Zeng G, Liang Y, Liao G. SPP1+ TAM subpopulations in tumor
microenvironment promote intravasation and metastasis of head and neck squamous cell
carcinoma. *Cancer Gene Ther*. 2024;31(2):311-321. doi:10.1038/s41417-023-00704-0

80. Wang Z, Liu J, Huang H, et al. Metastasis-associated fibroblasts: an emerging target for
metastatic cancer. *Biomark Res*. 2021;9(1):47. doi:10.1186/s40364-021-00305-9

81. Lee HO, Mullins SR, Franco-Barraza J, Valianou M, Cukierman E, Cheng JD. FAP-
overexpressing fibroblasts produce an extracellular matrix that enhances invasive velocity
and directionality of pancreatic cancer cells. *BMC Cancer*. 2011;11:245.
doi:10.1186/1471-2407-11-245

82. Lo A, Wang LCS, Scholler J, et al. Tumor-Promoting Desmoplasia Is Disrupted by
Depleting FAP-Expressing Stromal Cells. *Cancer Res*. 2015;75(14):2800-2810.
doi:10.1158/0008-5472.CAN-14-3041

83. Voltà-Durán E, Alba-Castellón L, Serna N, et al. High-precision targeting and destruction
of cancer-associated PDGFR-β+ stromal fibroblasts through self-assembling, protein-only
nanoparticles. *Acta Biomater*. 2023;170:543-555. doi:10.1016/j.actbio.2023.09.001

84. Subramanian C, Cohen MS. Over expression of DNA damage and cell cycle dependent
proteins are associated with poor survival in patients with adrenocortical carcinoma.
Surgery. 2019;165(1):202-210. doi:10.1016/j.surg.2018.04.080

85. Gara SK, Tyagi MV, Patel DT, et al. GATA3 and APOBEC3B are prognostic markers in
adrenocortical carcinoma and APOBEC3B is directly transcriptionally regulated by
GATA3. *Oncotarget*. 2020;11(36):3354-3370. doi:10.18632/oncotarget.27703

86. Gara SK, Lack J, Zhang L, Harris E, Cam M, Kebebew E. Metastatic adrenocortical
carcinoma displays higher mutation rate and tumor heterogeneity than primary tumors. *Nat
Commun*. 2018;9(1):4172. doi:10.1038/s41467-018-06366-z

87. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*.
2011;144(5):646-674. doi:10.1016/j.cell.2011.02.013

88. Ubhi T, Brown GW. Exploiting DNA Replication Stress for Cancer Treatment. *Cancer*
Res. 2019;79(8):1730-1739. doi:10.1158/0008-5472.CAN-18-3631

89. Saldivar JC, Cortez D, Cimprich KA. The essential kinase ATR: ensuring faithful
duplication of a challenging genome. *Nat Rev Mol Cell Biol*. 2017;18(10):622-636.
doi:10.1038/nrm.2017.67

90. Lecona E, Fernandez-Capetillo O. Targeting ATR in cancer. *Nat Rev Cancer*.
2018;18(9):586-595. doi:10.1038/s41568-018-0034-3

91. Yano K, Shiotani B. Emerging strategies for cancer therapy by ATR inhibitors. *Cancer Sci*.
2023;114(7):2709-2721. doi:10.1111/cas.15845

92. Dillon MT, Guevara J, Mohammed K, et al. Durable responses to ATR inhibition with
ceralasertib in tumors with genomic defects and high inflammation. *J Clin Invest*.
2024;134(2):e175369. doi:10.1172/JCI175369

93. Yap TA, Tan DSP, Terbuch A, et al. First-in-Human Trial of the Oral Ataxia
Telangiectasia and RAD3-Related (ATR) Inhibitor BAY 1895344 in Patients with
Advanced Solid Tumors. *Cancer Discov*. 2021;11(1):80-91. doi:10.1158/2159-8290.CD-
20-0868

94. Yap TA, O'Carrigan B, Penney MS, et al. Phase I Trial of First-in-Class ATR Inhibitor
M6620 (VX-970) as Monotherapy or in Combination With Carboplatin in Patients With
Advanced Solid Tumors. *J Clin Oncol Off J Am Soc Clin Oncol*. 2020;38(27):3195-3204.
doi:10.1200/JCO.19.02404

95. Middleton MR, Dean E, Evans TRJ, et al. Phase 1 study of the ATR inhibitor berzosertib
(formerly M6620, VX-970) combined with gemcitabine ± cisplatin in patients with
advanced solid tumours. *Br J Cancer*. 2021;125(4):510-519. doi:10.1038/s41416-021-
01405-x

96. Shapiro GI, Wesolowski R, Devoe C, et al. Phase 1 study of the ATR inhibitor berzosertib
in combination with cisplatin in patients with advanced solid tumours. *Br J Cancer*.
2021;125(4):520-527. doi:10.1038/s41416-021-01406-w

97. Telli ML, Tolaney SM, Shapiro GI, et al. Phase 1b study of berzosertib and cisplatin in
patients with advanced triple-negative breast cancer. *NPJ Breast Cancer*. 2022;8(1):45.
doi:10.1038/s41523-022-00406-0

98. Kim ST, Smith SA, Mortimer P, et al. Phase I Study of Ceralasertib (AZD6738), a Novel
DNA Damage Repair Agent, in Combination with Weekly Paclitaxel in Refractory Cancer.
Clin Cancer Res Off J Am Assoc Cancer Res. 2021;27(17):4700-4709. doi:10.1158/1078-
0432.CCR-21-0251

99. Konstantinopoulos PA, Cheng SC, Wahner Hendrickson AE, et al. Berzosertib plus
gemcitabine versus gemcitabine alone in platinum-resistant high-grade serous ovarian

cancer: a multicentre, open-label, randomised, phase 2 trial. *Lancet Oncol*. 2020;21(7):957-968. doi:10.1016/S1470-2045(20)30180-7

100. Gupta M, Concepcion CP, Fahey CG, et al. BRG1 Loss Predisposes Lung Cancers to Replicative Stress and ATR Dependency. *Cancer Res*. 2020;80(18):3841-3854. doi:10.1158/0008-5472.CAN-20-1744

101. Borenäs M, Umapathy G, Lind DE, et al. ALK signaling primes the DNA damage response sensitizing ALK-driven neuroblastoma to therapeutic ATR inhibition. *Proc Natl Acad Sci U S A*. 2024;121(1):e2315242121. doi:10.1073/pnas.2315242121

102. Kwok M, Davies N, Agathanggelou A, et al. ATR inhibition induces synthetic lethality and overcomes chemoresistance in TP53- or ATM-defective chronic lymphocytic leukemia cells. *Blood*. 2016;127(5):582-595. doi:10.1182/blood-2015-05-644872

103. Toledo LI, Murga M, Zur R, et al. A cell-based screen identifies ATR inhibitors with synthetic lethal properties for cancer-associated mutations. *Nat Struct Mol Biol*. 2011;18(6):721-727. doi:10.1038/nsmb.2076

104. Reaper PM, Griffiths MR, Long JM, et al. Selective killing of ATM- or p53-deficient cancer cells through inhibition of ATR. *Nat Chem Biol*. 2011;7(7):428-430. doi:10.1038/nchembio.573

105. Libè R, Groussin L, Tissier F, et al. Somatic TP53 mutations are relatively rare among adrenocortical cancers with the frequent 17p13 loss of heterozygosity. *Clin Cancer Res Off J Am Assoc Cancer Res*. 2007;13(3):844-850. doi:10.1158/1078-0432.CCR-06-2085

106. Neizer-Ashun F, Bhattacharya R. Reality CHEK: Understanding the biology and clinical potential of CHK1. *Cancer Lett*. 2021;497:202-211. doi:10.1016/j.canlet.2020.09.016

107. Couch FB, Bansbach CE, Driscoll R, et al. ATR phosphorylates SMARCAL1 to prevent replication fork collapse. *Genes Dev*. 2013;27(14):1610-1623. doi:10.1101/gad.214080.113

108. Chen YH, Jones MJK, Yin Y, et al. ATR-mediated phosphorylation of FANCI regulates dormant origin firing in response to replication stress. *Mol Cell*. 2015;58(2):323-338. doi:10.1016/j.molcel.2015.02.031

109. Ammazzalorso F, Pirzio LM, Bignami M, Franchitto A, Pichierri P. ATR and ATM differently regulate WRN to prevent DSBs at stalled replication forks and promote replication fork recovery. *EMBO J*. 2010;29(18):3156-3169. doi:10.1038/emboj.2010.205

110. Slyper M, Porter CBM, Ashenberg O, et al. A single-cell and single-nucleus RNA-Seq toolbox for fresh and frozen human tumors. *Nat Med*. 2020;26(5):792-802. doi:10.1038/s41591-020-0844-1

1035   111.   Lacar B, Linker SB, Jaeger BN, et al. Nuclear RNA-seq of single neurons reveals
1036   molecular signatures of activation. *Nat Commun*. 2016;7:11022.
1037   doi:10.1038/ncomms11022

1038   112.   McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-
1039   Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst*. 2019;8(4):329-
1040   337.e4. doi:10.1016/j.cels.2019.03.003

1041   113.   Tirosh I, Izar B, Prakadan SM, et al. Dissecting the multicellular ecosystem of
1042   metastatic melanoma by single-cell RNA-seq. *Science*. 2016;352(6282):189-196.
1043   doi:10.1126/science.aad0501

1044   114.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-
1045   seq data using regularized negative binomial regression. *Genome Biol*. 2019;20(1):296.
1046   doi:10.1186/s13059-019-1874-1

1047   115.   Stuart T, Butler A, Hoffman P, et al. Comprehensive Integration of Single-Cell Data.
1048   *Cell*. 2019;177(7):1888-1902.e21. doi:10.1016/j.cell.2019.05.031

1049   116.   Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at
1050   multiple resolutions. *GigaScience*. 2018;7(7):giy083. doi:10.1093/gigascience/giy083

1051   117.   Long Z, Sun C, Tang M, et al. Single-cell multiomics analysis reveals regulatory
1052   programs in clear cell renal cell carcinoma. *Cell Discov*. 2022;8(1):68.
1053   doi:10.1038/s41421-022-00415-0

1054   118.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Published online 2016.
1055   https://ggplot2.tidyverse.org.

1056   119.   Phipson B, Sim CB, Porrello ER, Hewitt AW, Powell J, Oshlack A. propeller: testing
1057   for differences in cell type proportions in single cell data. *Bioinforma Oxf Engl*.
1058   2022;38(20):4720-4726. doi:10.1093/bioinformatics/btac582

1059   120.   Kolde R. pheatmap: Pretty Heatmaps. Published online 2019. https://CRAN.R-
1060   project.org/package=pheatmap

1061   121.   van Biljon L, Fashemi B, Rodriguez J, et al. Visualizing DNA Damage Repair Proteins
1062   in Patient-Derived Ovarian Cancer Organoids via Immunofluorescence Assays. *J Vis Exp*
1063   *JoVE*. 2023;(192). doi:10.3791/64881

1064

1065 **FIGURE LEGENDS**

1066

Sample ID	Sample Name	Sample Description	Paired	Organ Site	Age at Surgery	Sex	Excess hormone secretion	Neoadjuvant Treatment	ENSAT Stage	KI67 Index	Somatic Mutations	Germline Mutations
Patient 1	NORMAL	Normal Adrenal	Yes	Adrenal Gland	35	F	None	N/A	N/A	N/A	N/A	N/A
	AA 1	Adrenal Adenoma	Yes	Adrenal Gland			Cortisol	N/A	N/A	N/A	N/A	N/A
Patient 2	AA 2	Adrenal Adenoma	No	Adrenal Gland	34	M	Cortisol	N/A	N/A	N/A	N/A	N/A
Patient 3	PACC 1	Primary ACC	Yes	Adrenal Gland	61	M	Cortisol	Etoposide, cisplatin	4	No data available	No data available	No data available
	MACC 1	Metastatic ACC	Yes	Liver						50%	*CHEK*2 p.Ile157Thr [1] *CTNNB1* p.Ser33Cys *TP53* p.Arg267Trp *DUSP16-ASXL1* fusion (D3;A11*)	No data available
Patient 4	PACC 2	Primary ACC	No	Adrenal Gland	54	F	Cortisol, DHEAS	Mitotane started 12 days prior to surgery	4	15-20%	*DAXX* p.Glu583fs*19 [1] *TP53* p.Tyr220fs*27 *BCORL1* p.Trp14*	No data available
Patient 5	PACC 3	Primary ACC	No	Adrenal Gland	25	F	Cortisol, androstenedione, testosterone, DHEAS	None	4	20%	*TERT* promoter c.124C>T [2] *CDKN2A* Copy Number Loss *CDKN2B* Copy Number Loss	No data available
Patient 6	PACC 4	Primary ACC	No	Adrenal Gland	54	F	None	None	4	No data available	*CTNNB1* p.Ser545Pro [2] *TP53* p.Asn200fs LOF	No data available
Patient 7	MACC 2	Metastatic ACC	No	Liver	57	M	Cortisol	Mitotane	4	20% (for primary tumor)	No data available	No clinically significant variants detected
Patient 8	MACC 3	Metastatic ACC	No	Lung	45	F	Cortisol	Mitotane	4	35% (for primary tumor)	No data available	No clinically significant variants detected
Patient 9	MACC 4	Metastatic ACC	No	Cervical Lymph Node	38	M	Cortisol, Androstenedione	2 months of mitotane treatment 2 years prior to surgery	4	1-10% (for primary tumor)	No data available	Variants of unknown significance detected *DICER* p.Trp32Cys *EGLN1* p.Pro378Ser

1067

1068

1069 **Table 1. Summary of Patient Specimens.** AA – adrenal adenoma, PACC – primary ACC,

1070 MACC – metastatic ACC, DHEAS - dehydroepiandrosterone sulfate. [1] - Foundation Medicine

1071 Testing, [2] - Tempus Testing.

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

Figure 1. Single-Nuclei RNA Sequencing Reveals Intratumoral Heterogeneity in Normal Adrenal, Adrenal Adenoma, Primary ACC, and Metastatic ACC. A. Overview of the experimental workflow. B. Summary of the samples and their hormone secretion profiles. C. UMAP projection of the integrated Seurat object showing that 10 clusters of 5 distinct cell types were identified. D. Dot plot demonstrating marker genes that were used to annotate the identified clusters. For the dot plot, scaled average expression is displayed. E. Bar plot showing cell type proportions of the identified cell populations across different sample types. DHEAS - dehydroepiandrosterone sulfate, AC – adrenal cortex.

Figure 2. Single-Nuclei RNA-Sequencing Uncovers Infiltrating Immunocyte Populations.

A. UMAP projection of subset lymphocyte and macrophage clusters from Figure 1C. B. Bar

plot display of percent immunocyte composition per sample type. C. Dot plot displaying key

differentially expressed cell typing markers. Scaled average expression is displayed. D. UMAP

projection of subset macrophage clusters from Figure 2A. E. Bar plot display of percent

macrophage subtypes per sample type. F. Dot plot displaying key differentially expressed cell

typing markers. Scaled average expression is displayed. G. Heatmap displaying average

expression of key macrophage immune checkpoint ligands for each sample type. Scaled

average expression is displayed.

1113

1114

1115

1116

Figure 3. Single-Nuclei Sequencing Identifies Cancer-Associated Fibroblasts in ACC. A.

UMAP projection demonstrating that further subclustering of fibroblasts identified in Figure 1C

results in identification of 5 fibroblast clusters, denoted 3.1 through 3.5. B. Bar plot

demonstrating cell type proportions of each of the identified fibroblast clusters (* - FDR <=

0.05, ***- FDR <= 0.001), C. Dot plot demonstrating top marker genes which were used to

annotate the identified clusters. Top 5 markers per cluster are displayed. D. Dot plot

demonstrating expression of known of fibroblast marker genes. For both dot plots, scaled

average expression is displayed. E. Heatmap showing expression of selected fibroblast

markers in fibroblast cluster 3.1. For the heatmap, scaled average expression is displayed.

FDR – false discovery rate.

1129

1130

1131

1132

1133

1134
1135

Figure 4. Single-Nuclei RNA Sequencing Demonstrates Intratumoral Heterogeneity of
Adrenal Cortex Cells. A. UMAP projection showing that further subclustering of adrenal
cortex cells results in identification of 7 adrenal cortex clusters, denoted AC 3.1 through AC
3.7. B. Dot plot demonstrating top 5 marker genes which were used to annotate each of the
identified clusters. For the dot plot, scaled average expression is displayed. C. Bar plot
showing cell type proportions of each of the adrenal cortex clusters (*** - FDR <= 0.001). D.
Bar plot demonstrating cell type proportions of each of the identified clusters on a per-sample
basis. E. Heatmap demonstrating average module scores for a selection of Hallmark
pathways downloaded from MSigDB for each of the identified clusters. In the heatmap,
average module scores are scaled on a per-row basis.

1146

1147

Figure 5. DNA Repair and Replication Pathways Are Enriched in Cluster AC 3.6. A.

Heatmap demonstrating average module scores for REACTOME replication stress and DNA

repair pathways downloaded from MSigDB. Clusters with fewer than 50 cells were excluded

from the analysis. In the heatmap, average module scores are scaled per row. B. Bar plot

demonstrating Ingenuity Pathway Analysis pathways upregulated and downregulated in

cluster AC 3.6. Pathways are ordered by the z-score, and top 10 upregulated and top 3

downregulated pathways are displayed. Only 3 pathways were found to be downregulated in

cluster AC 3.6. C. Bar plot showing total numbers of copy number variations (CNVs) inferred

by InferCNV on a per-cluster basis. Chi-square test was used to calculate statistical

significance (**** - FDR <= 0.0001). D and E. Bar plots showing the total number of gains (D)

and losses (E) inferred by inferCNV on a per-cluster basis. For both plots, chi-square test was

used to calculate statistical significance (**** - FDR <= 0.0001). F. Representative

immunofluorescence images of formalin fixed paraffin embedded (FFPE) tissues from two

patients stained for γH2AX (green), pRPA (red), and DAPI nuclear stain (blue). Scale bars, 10

μm. Zoom-in panels show details of nuclei, gH2AX foci and pRPA foci.

1163

1164

Figure 6. NCI-H295R Adrenocortical Cell Line Exhibits DNA Replication Stress Response. A. Immunofluorescence demonstrating that DNA damage or replication stress is present in NCI-H295R cells in the absence of cytotoxic agents as determined by the presence of intranuclear H2AX foci. In addition to the anti- H2AX antibody, the cells were stained with DAPI as nuclear stain and phalloidin. Images were obtained using a 20X objective on a Nikon® AXR confocal microscope. Insets represent a cropped portion of the same fields at a 5X higher magnification. B through F. Water-soluble tetrazolium dye cell viability assays demonstrating sensitivity of the NCI-H29R5 cell line to various drugs targeting replication stress and DNA damage repair pathways: elimusertib and VX-803 (ATR inhibitors, 6B), AZ32 and KU60019 (ATM inhibitors, 6C), rabusertib and PF47736 (CHK1 inhibitors, 6D), RI-1 and SAHA (RAD51 inhibitors, 6E) and olaparib (PARP inhibitor, 6F). For each of the drug treatments, top panels display raw data and bottom panels display 4-parameter non-linear lines of best fit. G. Schematic summarizing replication stress and DNA damage repair pathways and drug treatments targeting those pathways. nd – non-determined. SAHA -

 suberoylanilide hydroxamic acid, ns – not significant, * - p<0.05, ** - p < 0.01, *** - p<0.001,

 **** - p<0.0001.

Figure 7. ACC PTOs exhibit DNA Replication Stress Response. A. ACC PTOs proliferate in culture long-term. Brightfield images of two ACC PTO lines (**OSU1** and **OSU2**) were taken using a 20X objective >6 months after original tumor cell isolation. B. Immunofluorescence demonstrating that ACC PTOs express steroidogenic factor 1 (SF-1) and also showing that DNA damage or replication stress is present in ACC PTOs in the absence of cytotoxic agents as determined by the presence of intranuclear ☐H2AX foci. In addition to the anti- H2AX antibody and anti-SF-1 antibody, the cells were stained with DAPI as nuclear stain and phalloidin. Images were taken using a 20X objective on a Nikon® AXR confocal microscope. Insets represent a cropped portion of the same fields at a 5X higher magnification. C. ELISA demonstrates that ACC PTOs secrete cortisol. D. Water-soluble tetrazolium dye cell viability assays demonstrate that ACC PTOs exhibit sensitivity to ATR but not ATM inhibition. For each of the drug treatments, left panel displays raw data and right panel displays 4-parameter non-linear lines of best fit. nd - not determined, ns – not significant, * - p<0.05, ** - p < 0.01, *** - p<0.001, **** - p<0.0001.

1197