Improving the visualization of viruses in soil

Amar D. Parvate[1], Trinidad Alfaro[1], Regan McDearis[1], Amy Zimmerman[1], Kirsten Hofmockel[1], William C. Nelson[1], James E. Evans[1,3]*

[1]Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland, WA 99354, USA
[2]School of Biological Sciences, Washington State University, Pullman, WA 99354, USA
*Corresponding Author: James E. Evans (james.evans@pnnl.gov)

KEYWORDS: Soil sample, cryo-EM, virus diversity

ABSTRACT

Viruses are numerically the most abundant forms on Earth, and most are present in soil. Even though viruses are highly abundant in soil and critical to rhizosphere function, visualizing the diverse morphotypes within soil has been challenging. The difficulty is primarily due to the heterogenous nature of isolated suspensions that typically contain nanometer to micron scale debris which renders protein crystallography for structural studies unfeasible and hinders cryo-electron microscopy due to ice thickness and contrast issues. Here we employed and compared a simple spin filtration method to cleanup solutions of extracted viruses for direct observation with cryo-electron microscopy. Although relatively simple, the method employs common physical biochemical separation steps to remove large and small debris which dramatically improves image quality and preservation of structural features to permit visualizing morphotypes not typically seen with conventional negative stain approaches. In addition to tailed and non-tailed polyhedral phages, several under reported or novel morphotypes of soil viruses are directly visualized as a particle library with both 2D and 3D information.

INTRODUCTION

Viruses are numerically the most abundant forms on Earth with estimates of up to 10^{31} total virus particles worldwide [1, 2] and are being explored as alternatives to pesticides, antibiotics or as source of genes encoding molecules of biotechnological importance for commercialization purposes [3-5]. Interestingly, the majority of viruses are believed to be present in soil where they are estimated to range between 10^5 to 10^9/ g and outnumber bacteria by 5-1,000 fold [6]. Viruses are known to integrate themselves into complex soil ecosystems and interact with microbial communities [7]. In the case of agricultural land, multiple factors impact the soil virome including the use of fertilizers, the type of fertilizer, as well as type of crops grown [8].

Furthermore, the soil type, moisture content, season of sampling, and depth of sampling can all

impact the number and types of viruses observed from sample to sample [9]. This has largely

limited scientific study to viruses or phages which are known to either positively or negatively

impact human activities, livestock, or crops, and to a limited number of viruses which have been

successfully isolated and propagated through a host [10, 11]. As of 2023, soil viromes constituted

<1% of the total viromes available in public databases, making them some of the least studied

forms of life on the planet [12].

"Soil" is a deceptively simple term applied to an extremely complex layer of matter covering

the entire planet which harbors its own ecosystems consisting of everything from viruses to

unicellular microbes, fungi as well as the rhizosphere. Soil contains solid, liquid and gaseous

phases with stoichiometries varying over time and space. For soil virome studies, typically soil in

the top few inches is utilized [2]. Viruses regulate microbial communities in soil environments in

cycles, seasonally and spatiotemporally and may influence soil nutrient cycles in lands under

cultivation [2, 13]. Negative stain TEM is the dominant approach used in already documented

studies highlighting the morphological diversity of soil viruses with nanoscale imaging. Major

benefits for negative stain TEM are that it is cheap and easy to perform, the resulting heavy metal

stained grids are stable at room temperature for years and can be reimaged multiple times, and

in general this method requires ~10-100x less sample than other approaches like cryo-electron

microscopy. One of the most comprehensive examples of high-quality negative stain TEM

analysis of the soil virome was reported recently and focused on the discovery and

characterization of giant virus-like particles in forest soil[14]. While they presented stunning

images, the authors also acknowledged that their method of negative staining with heavy metal

and high salt content in conjunction with a chemical fixation step and dehydration could perturb

the viral ultrastructure resulting in shrinking and other artifacts. They also mention that "it is not

possible to prove the viral nature of any particles analyzed only by negative staining TEM,

regardless of how similar to published virus structures they may look," which is partly due to the

inability of negative staining to capture the full structural details and whole 3D structure of large

particles. This is a widely recognized limitation of negative staining which generally fails to

preserve high resolution features beyond ~16 Angstrom for non-crystalline samples, fails to stain

sugar or glycan surfaces, and fails to capture the full height and context of the virus components

depending on stain depth and local area reproducibility. Several researchers have also reported

that the method of extraction influences the virus morphotypes observed using negative stain

electron microscopy [6]. The difference in extraction methods includes but is not limited to factors

like buffer composition, type of soil, and whether a de-clumping step has been utilized, as these
can interfere with the uniformity of the negative stain.

Due to these historic challenges for direct morphological analysis of soil viromes in native-
like and hydrated (not chemically fixed or dehydrated) conditions, many studies are currently
exploring the soil virome with only genomic methods[15, 16]. However, both genomic and
morphological characterization remain critical aspects for taxonomic classification of viral
Families, and both are needed to fully understand the diversity of viruses present in soil [17].
Cryo-electron microscopy (cryo-EM) is a well-established technique for imaging biological
samples in near-native suspension conditions (i.e. monodisperse suspensions without any use of
fixatives or heavy metal salt stains) and allows the preservation of atomic details while permitting
direct visualization of material up to 0.8-1 micrometer in thickness which is an ideal range for most
virus particles. Here we present a sample preparation approach that quickly and simply converts
native soil isolates into a suspension compatible with high-resolution cryo-EM workflows for
visualizing the soil virome with 2D and 3D analyses. Using this workflow, we observed several
different morphotypes of tail-less and tailed phages with symmetrical heads and tail components,
as well as a diverse array of other morphotypes including round, elongated and polymorphic
viruses, and some morphotypes not previously reported using direct visualization.

METHODS
Field Site Description
Soil was collected on 7 March 2023 from the Tall Wheatgrass Irrigation Field Trial at the
Washington State University Irrigated Agriculture Research and Extension Center (WSU IAREC)
in Prosser, WA. As described previously [18], the site has aridisol soils with low organic matter
content and alkaline pH and was planted with Tall Wheatgrass in 2018. Irrigation treatments from
25% to 100% Field Water Capacity (FWC) have been ongoing since spring 2019. Soil was
collected from a random location within plot 23, which is planted with the Alkar cultivar of Tall
Wheatgrass and irrigated at 25% FWC, using a 5 cm diameter hammer corer. The soil from 5-15
cm depth was used in this study. Soil was transported to the laboratory on blue ice, where it was
homogenized and sieved (<2 mm) prior to further sample processing.

Enrichment of virus-like particles from soil
Virus-sized particles were enriched from freshly collected soils (maintained at 4ºC after
homogenization) following methods adapted from Santos-Medellín et al. [13]. For each sample,
20 g of soil was weighed into 50 mL tubes. Each tube received 18 mL of protein supplemented

phosphate-buffered saline solution (PPBS: 2% bovine serum albumin, 10% phosphate-buffered
saline, 1% potassium citrate, and 150 mM MgSO4) and was mixed at 300 rpm for 10 minutes at
4°C. After mixing, tubes were centrifuged at 4,000 x g for 10 minutes at 4°C. The supernatant
was collected, and the soil pellet was resuspended in 18 mL PPBS, mixed as above, and
centrifuged. These steps were repeated once more. The pooled supernatant was then centrifuged
twice at 10,000 x g for 8 minutes at 4°C for removal of remaining soil particles. The supernatant
was then filtered through a 0.22 µm filter for removal of bacterial cells. To concentrate the virus-
like particles (VLP), the filtrate was ultracentrifuged in a 45 Ti rotor at 32,000 x g for 3 hours at
4°C, and the supernatant discarded. To resuspend the VLP pellet, 300 µL of cell culture grade
water was added directly to the surface of the pellet, incubated at 4°C for 1 hour at 150 rpm, and
gently resuspended via pipetting.

Clean-up step before TEM imaging
The resulting VLP suspensions were subsampled (40 µL) for direct visualization by cryo-EM. Each
suspension was diluted ~9-10X in TBD buffer (350-400 µL) and loaded on a Pleuriselect 1 µm
membrane filter (Cat # SKU 46-10001-15) to separate large micron size debris. Samples were
centrifuged and the filtrate was loaded on a 300 kDa MWCO Amicon spin concentrator (Millipore)
to further concentrate the retentate to ~30 µL by centrifugation. The filtrate was then loaded on a
100 kDa MWCO spin concentrator and centrifuged. The retentate from this step was concentrated
to 30 µL. Alternatively, the filtrate from the Pleuriselect filter was loaded directly onto a 100 kDa
MWCO filter and the retentate was concentrated to 30 µL. All centrifugation steps were performed
at 5000-10,000 x g at 4°C for 15 min.

Grid preparation for TEM imaging
Aliquots of soil samples before the Pleuriselect step, and retentates from the 300 and 100 kDa
MWCO filters were used for preparing cryo-EM grids. Briefly, 3 µL of sample was loaded onto
glow discharged holey carbon grids (Quantifoil, 300 mesh, R1/2). Samples were blotted for 3s
and plunge frozen in liquid ethane using a Leica EM GP2 to obtain vitrified ice for imaging. For
negative staining, 2 µL suspension from the 300 kDa MWCO retentate was loaded on holey
carbon grids and excess sample was blotted away. Staining was performed using Nano-W™ (Ted
Pella Inc.), following conventional procedures of negative stained sample preparation [18]

TEM, cryo-EM and cryo-ET

Samples were loaded on a 300 keV (Thermo Fisher Scientific) Titan Krios and all screening and
data collection was performed using standard EPU software. Images were collected either at
nominal magnifications ranging from 135x, 580x, 11500x for low dose screening and 42000x for
data collection using a K3 direct electron detector (Gatan Inc.) fitted with a Bioquantum energy
filter and a slit width of 20 eV. Dose was not optimized for low dose screening images, but
exposure was kept to 1 s. For data collection, movies were collected at a pixel size of 1.1 Å in
counting mode, using a defocus of -1 to -3 μm, at an exposure of 2-2.5 s exposure across 32
frames resulting in a total dose of 25-28 e$^-$/Å^2/s. Motion correction was performed using standard
parameters in CryoSparc v4.2 and micrographs were Fourier cropped to 0.25 x [19]. Motion
corrected micrographs were inspected and sorted manually in Fiji [20] to visualize different virus
morphotypes. For cryo-ET, tilt series were collected from +/- 60° in 3° increments with a total dose
of 100 e$^-$/Å^2 using a dose symmetric scheme. Each tilt was collected as a movie with 3 subframes,
and an exposure of 0.36s per tilt. Tilt series were collected at a defocus of – 4 μm. Motion
correction and restacking of the tilt series was performed using MotionCorr2 using scripts written
in house [21]. Individual tilt images were then [22]denoised using TOPAZ and tilt-series were
imported in IMOD 4.12.11, binned 4-8x and tomograms were reconstructed using standard
reconstruction workflows using the ETomo suite [23]. Missing wedge correction was performed
using Isonet [24]. and final visualization of the reconstructed volumes was done in 3dmod and
ChimeraX [25].
156
RESULTS
Conventional negative stain TEM introduces artifacts and occludes some native details
Using a soil sample from a managed agricultural field site in Prosser, WA which was
enriched for viruses using an approached optimized for downstream genomics analysis, we
performed Negative Stain TEM to get an initial image of the sample. While the typical fields of
view for Negative Stain TEM can allow researchers to pick and choose the best local area and
particle of interest, we chose random locations to provide more reproducible examples of best
imaging conditions. As seen in Figure S1, two random locations within the same ~50 micron x 50
micron grid square show vastly different backgrounds. While similar elongated symmetrical
particles with intact tails can be identified in both images with the arrowheads, the two particles
show a putatively intact virus with genomic content still inside the capsid (red arrowhead) and a
particle that clearly has stain inside the capsid indicating the DNA/RNA content was already
released or the particle itself is damaged and not intact (white arrowhead). Other particle
morphologies can be seen but the images also show aggregation, stain artifacts and particle

spatial overlap. For example, while a bottle-shaped phage is seen in **Figure S1A** (white asterisk)
the stain prevented observing any high-resolution features like surface glycoproteins which are
expected to be present on that morphotype[26, 27].

Filtration improves cryo-EM imaging of viruses
Suspensions of native viruses isolated from soil generally have low abundance compared
to the amount of material needed for protein crystallography and they contain nanometer to
micron scale debris which complicate cryo-EM. Debris that is usually not a problem in
conventional negative stain electron microscopy or multi-omics becomes a hindrance in cryo-EM
for observing any viruses or virus like particles (VLPs) as the largest structures tend to set the
local ice thickness gradients (**Figure 1, grid *1**).

Figure 1 Workflow cartoon of methods to prepare soil samples for cryo-EM. Virus extraction
- Soil samples were processed per Santos-Medellin et al to extract viruses. Clean-up for Cryo-
EM – Suspensions of viruses natively isolated from soil contain viruses as well as nanometer to
micrometer size debris. Soil suspensions frozen on grids as-is and imaged with cryo-EM
contained lots of debris and thick ice which hindered visualization of virus particles (*1). However,
soil suspensions passed through a 1 µm pore size Pleuriselect filter got rid of larger debris in the
retentate and allowed the filtrate containing the viruses and smaller debris to be subsequently
further concentrated and fractionated using 2 separate approaches. In the first approach,
sequential filtering steps help separate different size viruses. Here the filtrate from the Pleuriselect
filter can be concentrated through a 300 kDa MWCO filter to concentrate viruses and get rid of
nanometer sized debris. This sample resulted in thin ice and facilitated visualization of viruses.

Cryo-EM images collected on the retentate (*2) capture most of the virus diversity. Filtrate from
the 300 kDa MWCO filter can be further processed through a 100 kDa MWCO filter. Imaging the
retentate (*3) showed fewer total viruses that were mostly smaller virus like particles, while
imaging the filtrate showed mostly junk protein. If desired the sequential approach allows for
visualizing the differently sized viruses. Alternatively filtrate from the Pleuriselect membrane can
be directly loaded on a 100 kDa MWCO filter. The retentate (*4) from this approach captures the
entire mixture of small and large viruses while and most small protein debris is removed to provide
quality thin ice to directly visualize the viral diversity. (Created using BioRender.com).
202

Using direct freezing of the as-delivered initial suspension, we observed thick ice and the
presence of large aggregates and mineral debris despite the original preparation for viral
enrichment including filtration through a 0.22 µm pore size membrane filter (**Figure 2** and **S2**).
We therefore employed a cleanup step to get rid of debris and aggregates that might occlude or
hinder visualization of viruses and VLPs in cryo-EM imaging. In principle, cryo-EM imaging with
a phase plate and energy filter can image specimens up to 1 micron in thickness. Our cleanup
step therefore consisted of centrifuging the suspension through a 1 µm pore size membrane
Pleuriselect filter to separate out the larger micron-sized debris. We then subject the cleaned
filtrate to simultaneous concentration and size selection filtration using 300 or 100 kDa MWCO
spin columns to further improve the sampling efficiency and image quality.
Following this simple cleanup, we observed that at lower magnifications (135x-580x) the
atlases and montages appeared roughly similar between the as-delivered and final cleaned up
samples **(Figure S2 A-C, S3A-C)**. However, images and montages collected at 11,500 x **(Figure**
S2D, S3D) showed clear differences where the as-delivered sample depicted high amounts of
debris which was absent from the cleaned samples **(Figure S2E, S3E).**

Figure 2 – Ice quality comparison of grids frozen before and after the cleanup step. (A) A
cryo-EM image at 42000 x nominal magnification showing overall devitrified crystalline ice. The

221 inset is a zoomed in view of a phage (blue boundary) which shows the crystalline ice hindering
222 visualization of the phage morphology. (B) A cryo-EM image from the sample post cleanup which
223 shows vitreous ice that allows visualization of multiple phage and virus particles (blue triangles).
224 The inset is zoomed in view of a phage in blue boundary showing glycoprotein spikes on an
225 unidentified elongated virus. Scale bar = 100 nm
226

Any protein macromolecules like viruses frozen for cryo-EM should be ideally present in
a layer of uniform vitreous ice only slightly thicker than the target macromolecule to maximize
contrast and resolution. To compare ice quality on the grids with virus suspensions frozen as-
delivered and after filtration, we collected images at 42,000x where the presence of ice
crystallization artifacts from uneven vitrification would be obvious. Crystalline ice has unit cell with
dimensions around 3.4 Angstroms which results in prominent spots visible in the Fast Fourier
Transform of the image or in severe cases the lattice is visible in the direct image. Due to this,
images which have crystalline ice cannot be used for high resolution structural studies. As
expected, images of the as-delivered virus suspension showed aggregates, debris as well as
devitrified ice **(Figure 2A)** with multiple microdomains showing tell-tale signs of crystalline ice.
While a bacilliform phage like particle was visible in the field of view, no high-resolution features
could be clearly seen even in magnified views **(Figure 2A inset)**. However, after filtration and
removal of the larger debris, the cryo-EM image (**Figure 2B**) of the virus suspension showed
uniform vitrified ice (confirmed by absence of any detectable lattice in the direct image nor any
diffraction spots in the FFT of the image, data not shown)**.** The vitreous ice, with minimal debris
from particle aggregation, allowed visualization of several phages and one elongated virus in the
field of view. A zoomed in view of the virus particle **(Figure 2B, inset)** showed similar morphology
as that in Figure 2A but here the vitreous ice allowed better visualization of the capsid and other
features.
To estimate how many viruses may have escaped observation in the 300 kDa MWCO
retentate, we further concentrated the 300 kDa MWCO filtrate over a 100 kDa MWCO spin
concentrator and imaged the resulting retentate (Figure 1, *3). Unsurprisingly, we mostly
observed vlps with diameters between 20-50 nm and an occasional virus. By quantifying the
number of observed phages seen in roughly equivalent datasets of around 1100 captured
micrographs from each of the 300 kDa MWCO and 100 kDa MWCO retentates, we concluded
that the 300 kDa spin concentration step is efficient to capture most (>90%) of the viral diversity
from our current sample and the 100 kDa spin concentration post 300 kDa spin concentration
may not be absolutely necessary beyond getting rid of small debris that can obscure high-
resolution imaging of smaller viruses. Thus, if the main goal is to evaluate the diversity of the

viromes, a combination of Pleuriselect 1 micrometer filtration followed directly by 100 kDa MWCO

filtration (Figure 1, *4) provides the best option for capturing all viruses and vlps present in solution

while removing both large and small debris that hinder high-quality imaging.

Direct visualization of viral diversity in soil extracts

While previous negative stain TEM studies [28-32] of soil extracts and high-resolution

cryo-EM structural characterization of purified viruses have provided detailed classification of

symmetric viruses, we decided to classify any detected viruses based on known and unknown

morphotypes to fully capture the diversity. The most readily identifiable morphology consisted of

tail-less polyhedral phages and phages with tails of varying length **(Figure 3A-B)**. Figure 3B.1

shows a bacilliform phage while 3B.2 shows a myovirus-like phage with relatively thicker tail, and

Figure 3B.3 a podovirus-like phage with a very short tail. Figure 3B.5 shows a siphovirus-like

phage where the head, tail and tail fibers are clearly visible. Our workflow also clearly enables

visualization of features like contractile and non-contractile tails **(Figure 3B.2, 3B.1,5)** as well as

glycoproteins on several symmetrical phages **(Figure 3B.4).** Diameters for the icosahedral phage

heads ranged from 50-80 nm whereas the bacilliform phages had 100-120 nm long heads with

40 nm diameter. Several images showed the presence of long thin tube-like structures with 10-

20 nm diameter and of varying lengths (200 nm – 1 μm) which could not be visually assigned to

any group of viruses **(Figure 3C.1,2, Figure S4 C)** and could be remnants of [31]broken tails or

contractile sheaths or something else currently undetermined. Another major morphotype

observed had an overall round shape, diameter <50 nm and no visible glycoproteins on the

surfaces **(Figure 3C.3,4,5, Figure S5 C)**.

In addition to the tailed and polyhedral viruses which are readily seen with negative stain

TEM and above with the cryo-EM workflow described here, we also identified several virus

morphotypes which have only be occasionally reported [28] and some novel ones which have

never been directly visualized before. These novel morphotypes could belong to any of the

multiple genera reported from soil viromes which currently have no morphology associated with

them. These virus morphotypes were broadly divided into groups based on either shape, size or

other visual similarities **(Figure 4)**. The presence of an intact membrane and visible glycoprotein

spikes of any shape and size were considered as pre-requisites for a visual "feature" observed in

an image to be counted as a virus. Round viruses with diameters between 50-200 nm and visually

different surface glycoprotein spikes made up the numerically second largest group (**Figure 4.1).**

A diverse group of morphotypes showed the presence of a very thick coat surrounding the viral

capsid/core **(Figure 4.2-6)**. The sizes of these viruses ranged from 20-157 nm with coat thickness

from 15-35 nm. We observed an inovirus like morphotype (caterpillar like) that had a cylindrical

central core dotted with putative glycoprotein spikes orthogonal to the axial virus core **(Figure**

4.7,8). Bottle shaped viruses with diverse morphologies **(Figure 4.9, 4.10**) and filamentous

viruses **(Figure 4.11)** were also occasionally observed. Interestingly, the images of bottle shaped

viruses by cryo-EM clearly show a glycoprotein-like coat that is not visualized in a similar negative

stain image (compare Figure 4.10 with Figure S1A white asterisk). Another group of viruses with

diameters ranging from 30 to 85 nm showed the presence of fiber like glycoprotein like projections

between 31-47 nm in length (here called LG viruses) **(Figure 4.12)**. The LG viruses are visually

similar to morphotypes described by Fischer *et al* as viruses with capsid fibers[14]. Of note, the

size/dimension of virus morphotypes did not have a linear correlation with either the measured

thickness of the external coat or length of the glycoprotein projections. Some polymorphic viruses

with irregular shapes were also observed which have not been reported before in literature

(Figure 4.13,14). While Figures 3 and 4 document the library of 2D morphotypes identified by this

imaging approach, we also summarize the results in Figure 5 to illustrate the diversity more

graphically.

**Figure 3 Cryo-EM images of symmetric viruses in filtrate of the sample 100 kDa MWCO
filter – A)** Cryo-EM image showing multiple phages with symmetric shapes. Tailed and tail-less
phages along with other icosahedral viruses. Phages display both icosahedral and bullet shaped
heads. B) Zoomed in view of 3 select phages from (A) shown. 1. A bacilliform phage showing a

long non contractile tail and siphovirus like morphology. 2. Phage showing a thick contractile tail having myovirus like morphology. 3. Podovirus like phage showing an extremely short tail (green triangle) (Phages in 1,2,4 are zoomed in views of phages from 3A). 4) Tail-less phage with glycoproteins visible on the virus surface. 5. Phage showing intact head (red triangle), contractile tail (green triangle) as well as tail fibers (blue triangle). 6. Phage showing a slightly rounded head relative to the phage shown in 3A.6. C) 1 and 2 are tube like VLPs of differing diameters abundantly observed in images. 3, 4, and 5 are VLPs which are <50 nm in diameter and have no visible glycoproteins on the surface. Scale bar = 50 nm for B and C.

Figure 4- Cryo-EM images of polymorphic viruses in filtrate of the sample 100 kDa MWCO filter. 1) Example of a round phage. 2-6) Diverse uncharacterized phages displaying a wide range of diameters and a thick outer coating/layer external to the virus core (red triangle). 7-8) Caterpillar like appearance of a virus with long thin central core and glycoproteins on the surface putatively belonging to the Inoviridae Family. 9-10) Tube shaped viruses with diverse morphologies 11) Elongated or filamentous phage. 12) Phage showing unique long projections which could be

glycoproteins or glycans (LG, red arrows) with the white parenthesis indicating the length of the
glycoproteins. 12) A polymorphic vlp showing a head and body morphology on either side of a
narrow neck (white triangle) 13) Polymorphic phage having plasmavirus like irregular morphology.
Scale bar 100 nm.

One of the main benefits of cryo-EM, besides imaging samples in near-native conditions

without crystallization, is that it can be used to both produce a 2D library of images as above and

obtain 3D volumetric information using tomography. Since the polymorphic viruses have not been

widely studied and their overall shape can be hard to recognize in 3D from a single 2D projection,

we performed cryo-electron tomography on a small-subset of the viruses **(Figures S4 and S5)**.

This approach allows tracing the contiguous and intact membranes for the polymorphic phages

as well as detecting glycoprotein extensions and in some cases density inside the viruses that

could correspond to packed DNA/RNA.

Figure 5 - Cartoon representation of virus morphotypes in current study. *The brown/red morphotypes have been reported in previous studies and were also observed in the current*

cryo-EM analyses. The blue morphotypes are mostly uncharacterized, diverse phages some of which have not yet been reported in literature. (Created using Biorender.com)

Discussion

Negative staining and conventional TEM has been widely used to visualize viruses and VLPs from soil environments, but a number of technical and operational problems have plagued this technique and hindered viral research. While negative stained TEM is a quick and easy way to visualize viruses, a recent report delved into the gradual decline in quality of electron microscopy images seen over the years by evaluating hundreds of publications where the loss was largely ascribed to a loss of expertise and training [33]. Additionally, previous methods for extracting viruses from soil have sometimes been too harsh or biased towards tailed phages [34]. Most studies of TEM imaging of tailed viruses isolated from soil have reported broken tails or loss of tails, where the frequency of loss of tail is inversely proportional to the tail length [6, 35]. Here we have demonstrated an alternate technique for visualizing viruses from challenging samples that employs highly selective but gentle filtration and cryo-EM to mitigate or eliminate many of the problems that plague negative stain TEM images.

So far, cryo-EM of soil viruses has been restricted to studies where a single virus isolated from soil sample can be propagated and purified in quantities amenable for cryo-EM single particle studies [36]. Here we overcome this challenge/bottleneck and demonstrate the ability to visualize the native viral community isolated directly from soil using cryo-EM by using common and simple physical separation techniques. We used microcentrifuge spin columns to simultaneously remove unwanted debris and to concentrate the sample to increase likelihood of detection on the grid which overcame two of the main bottlenecks. The resulting grids were then imaged using semi-automated cryo-EM screening and data collection where we able to detect at least one virus or VLP in ~70% of the images collected on the 300 kDa MWCO retentate. Since most of the virus suspension loaded on a grid gets blotted away prior to vitrification, the virus library we have obtained **(Figure 5)** cannot yet be truly representative and it is possible that several morphotypes could not be imaged. However, this drawback also extends to libraries obtained using negative stained TEM and our work sets the foundation for being able to visualize the viral diversity with cryo-EM while future work can focus on optimizing the loading and sampling efficiency to capture more of the diversity. For example, multiblot methods are known to increase the local concentration of particles trapped in cryo-EM grid holes even when using wicking, while lossless sample deposition and vitrification can be accomplished using the Vitrojet [37] and other similar preparative instruments.

Another way to validate the diversity is linking the observed cryo-EM 2D and 3D data with
omics analysis on the same samples. While beyond the focus of the current study, orthogonal
multi-omics would be needed to provide more information about the morphotypes imaged in this
study to confirm that a novel virus morphotype is truly an infectious virus. For example, gene
transfer agents (GTAs) appear and function like phages but carry fragments of cellular genome
resulting in horizontal gene transfer. It would be very challenging to distinguish between a
bacteriophage and a GTA purely based on 2D images [38]. Additionally, some of the round virus
morphotypes we have observed may be a result of cellular blebbing. Without further orthogonal
studies, it is not possible to unambiguously assign all phage morphotypes to the Class
Caudoviricetes [17].

Given estimates of the diversity of viruses in soil, there is ample opportunity for this work
to be replicated on soil samples from numerous diverse locations, during different times of the
year to obtain a broader representation of viruses resident in soil. Further, for higher resolution
studies, soil virus extractions could be processed using the virus reduction approaches used in
aquatic virology to selectively propagate a few viruses in high copy number [39, 40]. Complete
purification and isolation of these viruses would not be needed; their selective amplification would
provide suitable samples for high-resolution cryo-EM/ET studies using sub-tomogram averaging,
while also allowing verification of host-virus infectivity. Hypothetically, a dataset with just a few
hundred symmetric virus particles could be used to obtain 2D classes and sub-nanometer
resolution 3D maps. The selectively amplified viruses could be used to infect target host cells and
samples can be frozen at various stages of infection to perform cryo-ET, which will provide
structural details of interaction of host-virus systems from soil in unprecedented resolution.

Acknowledgement and Funding:
The research was performed using EMSL (grid.436923.9), a DOE Office of Science User Facility
sponsored by the Biological and Environmental Research program located at PNNL. A portion of
this research was performed on project award (10.46936/fics.proj.2022.60449/60008585) under
the FICUS program and used resources at the DOE Joint Genome Institute and the
Environmental Molecular Sciences Laboratory, which are DOE Office of Science User Facilities.
Both facilities are sponsored by the Biological and Environmental Research program and
operated under Contract Nos. DE-AC02-05CH11231 (JGI) and DE-AC05-76RL01830 (EMSL). A
portion of this research was supported by the DOE Office of Biological and Environmental
Research FWP 74915 and D7G33.

Conflict of Interest

The authors declare no conflict of interest.

Author Contributions

JEE, ADP conceived this project. ADP, RWM, TA, AEZ performed all the experiments and ADP performed all the data analyses. JEE, WCN and KSH provided supervision and research support (funding). All the authors contributed to drafting and editing the manuscript.

References

1. Bar-On, Y.M., R. Phillips, and R. Milo, *The biomass distribution on Earth.* Proc Natl Acad Sci U S A, 2018. **115**(25): p. 6506-6511.
2. Williamson, K.E., et al., *Viruses in Soil Ecosystems: An Unknown Quantity Within an Unexplored Territory.* Annu Rev Virol, 2017. **4**(1): p. 201-219.
3. Duan, Y., et al., *Bacteriophage targeting of gut bacterium attenuates alcoholic liver disease.* Nature, 2019. **575**(7783): p. 505-511.
4. Adriaenssens, E.M., et al., *T4-related bacteriophage LIMEstone isolates for the control of soft rot on potato caused by 'Dickeya solani'.* PLoS One, 2012. **7**(3): p. e33227.
5. Jansson, J.K. and R. Wu, *Soil viral diversity, ecology and climate change.* Nat Rev Microbiol, 2023. **21**(5): p. 296-311.
6. Williamson, K.E., et al., *Estimates of viral abundance in soils are strongly influenced by extraction and enumeration methods.* Biology and Fertility of Soils, 2013. **49**(7): p. 857-869.
7. Santos-Medellín, C., et al., *Viral but not bacterial community succession is characterized by extreme turnover shortly after rewetting dry soils.* bioRxiv, 2023.
8. Chen, L., et al., *Effect of different long-term fertilization regimes on the viral community in an agricultural soil of Southern China.* European Journal of Soil Biology, 2014. **62**: p. 121-126.
9. Narr, A., et al., *Soil Viral Communities Vary Temporally and along a Land Use Transect as Revealed by Virus-Like Particle Counting and a Modified Community Fingerprinting Approach (fRAPD).* Front Microbiol, 2017. **8**: p. 1975.
10. Noteborn, W.E.M., et al., *Using cryogenic electron microscopy methods to gain insight into structure and initial host attachment of the flagellotropic bacteriophage 7-7-1.* bioRxiv, 2025.
11. Almeida, G.M.d.F., et al., *Genomic and structural insights into Jyvaskylavirus, the first giant virus isolated from Finland.* eLife, 2025. **13**.
12. Graham, E.B., et al., *A global atlas of soil viruses reveals unexplored biodiversity and potential biogeochemical impacts.* Nat Microbiol, 2024. **9**(7): p. 1873-1883.
13. Santos-Medellin, C., et al., *Spatial turnover of soil viral populations and genotypes overlain by cohesive responses to moisture in grasslands.* Proc Natl Acad Sci U S A, 2022. **119**(45): p. e2209132119.
14. Fischer, M.G., U. Mersdorf, and J.L. Blanchard, *Amazing structural diversity of giant virus-like particles in forest soil.* bioRxiv, 2023.
15. Hazard, C., et al., *Beneath the surface: Unsolved questions in soil virus ecology.* Soil Biology & Biochemistry, 2025. **205**.

16. Roux, S. and J.B. Emerson, *Diversity in the soil virosphere: to infinity and beyond?* Trends
Microbiol, 2022. **30**(11): p. 1025-1035.
17. Turner, D., et al., *Abolishment of morphology-based taxa and change to binomial species*
names: 2022 taxonomy update of the ICTV bacterial viruses subcommittee. Arch Virol,
2023. **168**(2): p. 74.
18. Naylor, D., et al., *Interactive effects of depth and differential irrigation on soil microbiome*
composition and functioning. Frontiers in Microbiomes, 2023. **2**.
19. Punjani, A., et al., *cryoSPARC- algorithms for rapid unsupervised cryo-EM structure*
determination. Nature Methods, 2017. **14**: p. 290-297.
20. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat
Methods, 2012. **9**(7): p. 676-82.
21. Zheng, S.Q., et al., *MotionCor2: anisotropic correction of beam-induced motion for*
improved cryo-electron microscopy. Nat Methods, 2017. **14**(4): p. 331-332.
22. Bepler, T., et al., *Topaz-Denoise: general deep denoising models for cryoEM and cryoET.*
Nat Commun, 2020. **11**(1): p. 5208.
23. Kremer, J.R., D.N. Mastronarde, and J.R. McIntosh, *Computer visualization of three-*
dimensional image data using IMOD. J Struct Biol, 1996. **116**(1): p. 71-6.
24. Liu, Y.T., et al., *Isotropic reconstruction for electron tomography with deep learning.* Nat
Commun, 2022. **13**(1): p. 6482.
25. Pettersen, E.F., et al., *UCSF ChimeraX: Structure visualization for researchers, educators,*
and developers. Protein Sci, 2021. **30**(1): p. 70-82.
26. Prangishvili, D., P. Forterre, and R.A. Garrett, *Viruses of the Archaea: a unifying view.* Nat
Rev Microbiol, 2006. **4**(11): p. 837-48.
27. Wang, F., et al., *Spindle-shaped archaeal viruses evolved from rod-shaped ancestors to*
package a larger genome. Cell, 2022. **185**(8): p. 1297-1307 e11.
28. Ackermann, H.W., *5500 Phages examined in the electron microscope.* Arch Virol, 2007.
152(2): p. 227-43.
29. Swanson, M.M., et al., *Viruses in soils: morphological diversity and abundance in the*
rhizosphere. Annals of Applied Biology, 2009. **155**(1): p. 51-60.
30. Reavy, B., et al., *Distinct circular single-stranded DNA viruses exist in different soil types.*
Appl Environ Microbiol, 2015. **81**(12): p. 3934-45.
31. Zheng, J., et al., *A Capsid Structure of Ralstonia solanacearum podoviridae GP4*
with a Triangulation Number T = 9. Viruses, 2022. **14**(11).
32. Effantin, G., et al., *Cryo-electron microscopy three-dimensional structure of the jumbo*
phage PhiRSL1 infecting the phytopathogen Ralstonia solanacearum. Structure, 2013.
21(2): p. 298-305.
33. Ackermann, H.W., *Sad State of Phage Electron Microscopy. Please Shoot the Messenger.*
Microorganisms, 2013. **2**(1): p. 1-10.
34. Williamson, K.E., K.E. Wommack, and M. Radosevich, *Sampling natural viral communities*
from soil for culture-independent analyses. Appl Environ Microbiol, 2003. **69**(11): p. 6628-
33.
35. Williamson, K.E., R.R. Helton, and K.E. Wommack, *Bias in bacteriophage morphological*
classification by transmission electron microscopy due to breakage or loss of tail
structures. Microsc Res Tech, 2012. **75**(4): p. 452-7.
36. Okamoto, K., et al., *Structural variability and complexity of the giant Pithovirus sibericum*
particle revealed by high-voltage electron cryo-tomography and energy-filtered electron
cryo-microscopy. Sci Rep, 2017. **7**(1): p. 13291.
37. Ravelli, R.B.G., et al., *Cryo-EM structures from sub-nl volumes using pin-printing and jet*
vitrification. Nat Commun, 2020. **11**(1): p. 2563.
38. Bardy, P., et al., *Structure and mechanism of DNA delivery of a gene transfer agent.* Nat
Commun, 2020. **11**(1): p. 3034.

39.   Wei, W., et al., *Reduced bacterial mortality and enhanced viral productivity during sinking*
in the ocean. ISME J, 2022. **16**(6): p. 1668-1675.
40.   Wilhelm, S.W., S.M. Brigden, and C.A. Suttle, *A dilution technique for the direct*
measurement of viral production: a comparison in stratified and tidally mixed coastal
waters. Microb Ecol, 2002. **43**(1): p. 168-73.

Supplementary Figures

Figure S1 Negative stained image of viruses natively isolated after the Pleuriselect step –
Conventional negatively stained images of viruses collected at 42000 x nominal magnification. A)
Example of a good negative staining where only the background has been stained and phage
shape is clearly visible (white triangle) B) Example of an overstained image where stain seems to
have penetrated inside the phage head (read triangle). The images show a lot of background junk
and loss of high-resolution features. Area in blue boundary shows aggregates in the field of view
which prevents identifying any VLPs due to spatial overlap.

Figure S2 – Comparison of scales of imaging for virus samples before purification. A) A
stitched atlas images collected at very low magnification showing the entire cryo-EM grid showing.
Squares on the grid are colored from relatively thick (purple) to thin (blue) with the blue colored
squares having ice thickness ideal for cryo-EM imaging to B) A zoomed in area of the atlas level
magnification shows no visible difference in the grid with samples before or after purification. C)
A higher mag view of the grid with an individual square (blue border and inset) that shows no
visible difference in ice quality. D) A 2x2 montage of individual images collected at 11500x
magnification for the sample before Pleuriselect cleanup. At this magnification, the ice begins to
show irregularities. E) Zoomed in image of the individual image in the montage shows presence
of debris (red triangle) as well as crystalline ice (red arrow). Scale bar for D and I – 500 nm. Each
circular hole is 1 μm in diameter.

Figure S3 - Comparison of scales of imaging for virus samples after purification. A) stitched atlas images collected at very low magnification showing the entire cryo-EM grid showing. Squares on the grid are colored from relatively thick (purple) to thin (blue) with the blue colored squares having ice thickness ideal for cryo-EM imaging to B) A zoomed in area of the atlas level magnification shows no visible difference in the grid with samples before or after purification. C) A higher mag view of the grid with an individual square (blue border and inset) that shows no visible difference in ice quality. D) A 2x2 montage of individual images collected at 11500x magnification for the sample after Pleuriselect cleanup. E) Zoomed in image of the individual image in the montage shows no detectable. No visible large debris or crystalline ice seen in the montage or the zoomed in image. Scale bar for D and I – 500 nm. Each circular hole is 1 µm in diameter.

Figure S4 - Cryo-tomographic imaging of viruses natively isolated from soil. A)Zero-degree
tilt image of the tomogram with diverse morphotypes. B) Slice through the reconstructed

tomogram with 3 morphotypes in in the red, blue and yellow boundary. C) Successive Z slices
through the tomogram for each of the individual morphotypes tracing appearance and
disappearance of the viruses/vlps. False color rendering of the virus/vlps are shown as visualized
using ChimeraX. Red – a comma or bean shaped vlp showing intact membranes and glycoprotein
spikes visible on its surface. Blue – An elongated virus morphotype with blebs internal to its
membrane and glycoprotein spikes on the surface. Pink – A tube like morphotype observed with
uncharacterized material internal to the tube. Beige – Morphotype with a thick coat surrounding
the central core. Slices in pink and beige boundaries have overlapping regions. All scale bars =
25 nm

Figure S5 Cryo-tomographic imaging of vlps natively isolated from soil. . A)Zero-degree tilt
image of a tomogram with diverse morphotypes/vlps B) Slice through the reconstructed tomogram
with 3 morphotypes in in the green, violet and gold. C) Successive Z slices through the tomogram

565   for each of the individual morphotypes tracing appearance and disappearance of the viruses/vlps.
566   False color rendering of the virus/vlps are shown as visualized using ChimeraX. Green – A round
567   morphotype with a small core and a thick coat. Violet– A curved morphotype. Gold – Two round
568   vlps very close to each other. No glycoproteins visible on their surface. All scale bars = 50 nm

569