CXXC-finger protein 1 associates with FOXP3 to stabilize homeostasis and suppressive functions of regulatory T cells

Xiaoyu Meng, [1,2,3†] Yezhang Zhu, [4,5†] Kuai Liu, [1,2,3†] Yuxi Wang,[6] Xiaoqian Liu, [1,2,3] Chenxin Liu,[2] Yan Zeng, [1,2,3] Shuai Wang, [1,2,3] Xianzhi Gao, [1,2,3] Xin Shen, [7] Jing Chen, [8] Sijue Tao, [6] Qianying Xu, [2] Linjia Dong, [9] Li Shen, [10,11,*] and Lie Wang[1,2,3,6,12,**]

[1]Institute of Immunology and Bone Marrow Transplantation Center, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.

[2]Zhejiang University School of Medicine, Hangzhou, China.

[3]Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, China.

[4]Department of Hematology, Tongji Hospital, School of Medicine, Tongji University, Shanghai, China

[5]Shanghai Key Laboratory of Signaling and Disease Research, Frontier Science Center of Stem Cell Research, National Stem Cell Translational Resource Center, School of Life Sciences and Technology, Tongji University, Shanghai, China

[6]Laboratory Animal Center, Zhejiang University, Hangzhou, China.

[7]Co-Facility Center, Zhejiang University School of Medicine, Hangzhou, China

[8]Department of Gastrointestinal Surgery, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.

[9]School of Basic Medical Sciences and Forensic Medicine, Hangzhou Medical College,

Hangzhou, China

[10]MOE Key Laboratory of Biosystems Homeostasis & Protection and Zhejiang Provincial Key

Laboratory for Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang University,

Hangzhou, China.

[11]Department of Orthopedics Surgery, the Second Affiliated Hospital, School of Medicine,

Zhejiang University, Hangzhou, China

[12]Future Health Laboratory, Innovation Center of Yangtze River Delta, Zhejiang University,

Jiaxing, China.

[†] These authors contributed equally to this work.

* Correspondence: wanglie@zju.edu.cn; li_shen@zju.edu.cn.

Abstract

FOXP3-expressing regulatory T (T_{reg}) cells play a pivotal role in maintaining immune

homeostasis and tolerance, with their activation being crucial for preventing various

inflammatory responses. However, the mechanisms governing the epigenetic program in T_{reg}

cells during their dynamic activation remain unclear. In this study, we demonstrate that CXXC

finger protein 1 (CXXC1) interacts with the transcription factor FOXP3 and facilitates the

regulation of target genes by modulating H3K4me3 deposition. *Cxxc1* deletion in T_{reg} cells leads

to   severe   inflammatory   disease   and   spontaneous   T-cell   activation,   with   impaired

immunosuppressive function. As a transcriptional regulator, CXXC1 promotes the expression of

key T_{reg} functional markers under steady-state conditions, which are essential for the

maintenance of T_{reg} cell homeostasis and their suppressive functions. Epigenetically, CXXC1

binds to the genomic regulatory regions of T_{reg} program genes in mouse T_{reg} cells, overlapping

with FOXP3 binding sites. Given its critical role in T_{reg} cell homeostasis, CXXC1 presents itself

as a promising therapeutic target for autoimmune diseases.

Introduction

Regulatory T (T_{reg}) cells are a distinct subset of $CD4^+$ T cells that play a critical role in

maintaining immune homeostasis and self-tolerance by suppressing excessive or aberrant

immune responses to foreign or self-antigens [1-3]. These cells can be further categorized into

thymus-derived regulatory T cells (tT_{reg} cells), periphery-derived T_{reg} cells (pT_{reg} cells), and

induced T_{reg} cells (iT_{reg} cells) [4]. They uniquely express the transcription factor FOXP3, a member

of the forkhead winged-helix family, which is essential for T_{reg} cell lineage commitment and

suppressive function [5,6]. Deletion or mutation of the *Foxp3* gene leads to a range of

immunological disorders, including allergies, immunopathology, and autoimmune diseases in

both mice and humans [7,8].

It is well established that FOXP3 recruits various cofactors to form complexes that either

promote or repress the expression of downstream genes, with histone and DNA modifications

playing pivotal roles in this process. FOXP3 can activate or repress the transcription of key

regulators of T_{reg} cell activation and function by recruiting the histone acetyltransferases (HATs)

or histone deacetylases (HDACs)[9,10]. Notably, FOXP3-bound sites exhibit significant enrichment

of H3K27me3, a modification essential for FOXP3-mediated repressive chromatin remodeling

under inflammatory conditions[11]. However, the direct role of FOXP3 as a transcriptional

activator through interactions with epigenetic regulators, particularly via modulation of H3K4

trimethylation, remains poorly documented.

In mammals, six proteins have been identified that catalyze H3K4 methylation. These proteins

contain the SET domain and include MLL1 (KMT2A), MLL2 (KMT2B), MLL3 (KMT2C),

MLL4 (KMT2D), SETD1A, and SETD1B [12,13]. For example, MLL1 plays a critical role as an

epigenetic regulator in T_{reg} cell activation and functional specialization[14]. Additionally, Placek *et*

al. demonstrated that MLL4 is essential for T_{reg} cell development by catalyzing H3K4me1 at

distant unbound enhancers through chromatin looping [15]. H3K4me3, which is enriched at the

transcription start site (TSS) and the CpG island (CGI), converts chromatin into active

euchromatin by recruiting activating factors [16]. CXXC finger protein 1(CXXC1, also known as

CFP1), which contains a SET1 interaction domain (SID), is required for binding to the histone

H3K4 methyltransferases SETD1A and SETD1B [17]. Previous studies have shown that CXXC1

regulates promoter patterns during T cell maturation, GM-CSF-derived macrophage

phagocytosis, TH17 cell differentiation, and the function of ILC3 cells with aging by modulating

H3K4me3 modification[18-21]. Despite the well-documented role of CXXC1 in various immune

effector cells, its role in T_{reg} cells remains unclear.

Here, we demonstrate that CXXC1 interacts with FOXP3 and enhances the expression of FOXP3

target genes. T_{reg} cell-specific deletion of *Cxxc1* triggers systemic autoimmunity, accompanied

by multiorgan inflammation, ultimately resulting in early-onset fatal inflammatory disease in

mice. *Cxxc1*-deficient T_{reg} cells exhibit a significant disadvantage in proliferation and

homeostasis, even in non-inflammatory mice where coexisting wild-type (WT) T_{reg} cells were

present. Moreover, *Cxxc1*-deficient T_{reg} cells display intrinsic defects in the expression of key

suppression molecules, including CTLA-4, CD25, ICOS, and GITR. Consistent with these

findings, mechanistic studies strongly suggest that CXXC1 functions as an essential cofactor of

FOXP3, playing a crucial role in sustaining H3K4me3 modifications at key T_{reg} cell genes.

Materials and Methods

Mice

All mice used in this study were bred for a minimum of seven generations on a C57BL/6

background. Mouse experiments, or cells from mice of the same genotype, compared littermates

or age-matched control animals. The *Cxxc1*$^{fl/fl}$ mouse strain has been previously described [22]. The

Foxp3^{cre} mice (JAX,016959) were generously provided by Bin Li (Shanghai Jiao Tong

University School of Medicine, Shanghai, China). CD45.1 (NM-KI-210226) mice were

purchased from the Nanjing Biomedical Research Institute of Nanjing University. *Rag1*$^{-/-}$ mice

(stock#T004753) were purchased from GemPharmatech. 2D2 (MOG35-55-specific TCR

transgenic) mice were graciously supplied by Prof. Linrong Lu (Zhejiang University School of

Medicine, Hangzhou, Zhejiang, China). *Foxp3*^{cre} (WT) and *Foxp3*^{cre} *Cxxc1*$^{fl/fl}$ (cKO) were

utilized at 3 weeks old unless otherwise stated. All mice were housed in the Zhejiang University

Laboratory Animal Center under specific pathogen-free conditions, and all animal experimental

procedures were approved by the Zhejiang University Animal Care and Use Committee

(approval no.ZJU20230246).

Cell culture

102   HEK 293T cells were obtained from ATCC, and Plat E cells were kindly provided by Prof.

103   Xiaolong Liu (Shanghai Institutes for Biological Sciences). Both cell lines were cultured in

104   Dulbecco-modified Eagle medium (DMEM) containing 10% (v/v) fetal bovine serum (FBS),

105   supplemented with 1% penicillin/streptomycin.

Immunofluorescence (IF) Staining

As previously described [23], coverslips were treated with a 0.01% poly-L-lysine solution (P4707;

Sigma) for 10 minutes, air-dried, and then coated with $CD4^+$ YFP^+ T_{reg} cells. The cells were then

fixed in 4% formaldehyde for 15 minutes at room temperature, permeabilized with 0.2% Triton

X-100, and blocked with 1% BSA. Antibodies against CXXC1 (ab198977; Abcam) and

FOXP3(17-5773-82; Invitrogen) were diluted in Image iT FX signal enhancer (I3693;

Invitrogen) and incubated with cells overnight at 4 °C. After washing with phosphate-buffered

saline (PBS), the cells were incubated with a goat anti-rabbit antibody Alexa Fluor 594

(1:250;10015289; Invitrogen) secondary antibody and stained with DAPI (200 ng/mL; D523;

Dojindo). Slides were washed with PBS and sealed with an antifade solution(P36934; Invitrogen)

before imaging with an Olympus FV3000 fluorescence microscope. The images were visualized

using the FV31-SW software.

Co-immunoprecipitation and western blot

Harvest the appropriate transfected cell lines and primary cells from culture and wash them with

ice-cold PBS. Lyse the cells in NETN300 buffer (300 mM NaCl, 0.5 mM EDTA, 0.5% (v/v)

NP-40, 20 mM Tris-HCl pH 8.0) supplemented with a protease inhibitor (1:100, P8340,

Sigma-Aldrich) and PMSF (1 mM) on ice for 10 minutes. Take a small portion of the whole-cell

lysate as input, and incubate the remaining lysate with either Anti-FLAG M2 Beads (M8823;

Sigma) or Anti-HA Beads (HY-K0201; MCE) on a rotator at 4°C overnight. For the endogenous

Co-IP assay targeting CXXC1 and FOXP3, incubate the cell lysate with protein G magnetic

beads along with anti-CXXC1 (ab198977; Abcam) or anti-FOXP3 (14-4774-82; Invitrogen)

antibodies on a rotator at 4°C overnight. Wash the beads three times with IP buffer (100 mM

NaCl, 0.5 mM EDTA, 0.5% (v/v) NP-40, 20 mM Tris-HCl pH 8.0) to remove non-specific

binding. Boil the washed beads with 1× Laemmli sample buffer (1610747; Bio-Rad) to elute the

bound proteins. Separate the denatured proteins by SDS-PAGE. Transfer the separated proteins

onto PVDF membranes (Millipore) for immunoblotting.Immunoblot the PVDF membranes

(IPVH00010) with the following antibodies: anti-CXXC1 (1:1000; ab198977; Abcam),

anti-FOXP3 (1:500; 14-7979-80; Invitrogen), anti-FlAG (1:1000; 14793; Cell Signaling

Technology), anti-HA (1:1000; 3724S; Cell Signaling Technology).Detect the immunoblotted

proteins using a secondary HRP-conjugated goat anti-rabbit antibody (1:1000; HA1001-100;

Huabio) and visualize the bands using an appropriate detection method.

Flow cytometry

Spleen and lymph nodes were harvested from 3-week-old mice. Tissues were placed in a dish

containing PBS with 2% fetal bovine serum (FBS). The tissues were minced into small pieces

and then filtered through a 70 µM nylon filter to obtain a single-cell suspension. For surface

marker analysis, cells were incubated for 15 minutes with purified anti-mouse CD16/32 antibody

(101320; BioLegend) to block Fc receptors. After blocking, cells were stained with the indicated

antibodies for surface markers. To determine cytokine expression, cells were stimulated for 4

hours at 37□ with phorbol12-myristate13-acetate (50ng/mL; S1819; Beyotime), ionomycin

(1mg/ml; S1672; Beyotime), and brefeldin A (BFA;00-4506-51; Invitrogen). After stimulation,

cells were labeled with a fixable viability dye and surface markers. Cells were then fixed and

permeabilized according to the manufacturer's instructions (00-8222-49; Invitrogen). For

transcription factor staining, samples were fixed using the FOXP3/Transcription Factor Staining

Buffer Set (00-5523-00; Invitrogen). Flow cytometry was performed using a BD Fortessa (BD

Biosciences). Flow cytometry data were acquired and analyzed using FlowJo software.

The following antibodies were purchased from Invitrogen or BioLegend: Zombie Violet f-

ixable viability(423113), Zombie NIR fixable(423105), CD25(PC61), CD8α(53-6.7), CD62

L(MEL-14), PD-1(J43), CD44(IM7), IL-17A(TC1118H10), KL-RG1(2F1), CD4 (GK1.5), T

CRβ(H57-597), IFN-γ(XMG1.2), FOXP3(FJK-16s), CTLA-4(UC10-4B9), ICOS(15F9), GIT

R(DTA-1), LAG3(C9B7W), TCRVβ11(RR3-15), CD45RB(C363-16A) CD69 (H1.2F3), MH

CII(M5/114.15.2), CD80(16-10A1), T-bet(eBio4B10), IKZF4(ESB7C2), IL-4(11B11), CCR7

(4B12), CD73(TY/11.8).

Real-time PCR

Total RNA was extracted from T_{reg} cells using the RNAAiso Plus(9109; Takara) reagent

according to the manufacturer's instructions, and cDNA synthesis was performed using the Prime

Script RT Reagent Kit (Takara). TB Green Premix Ex Taq (RR420A; Takara) was used for

quantitative real-time PCR (qPCR). The expression levels of target mRNA were normalized to

the level of *β-actin* expression. The primers for qPCR are as follows:

Cxxc1 qPCR Forward: ATCCGGGAATGGTACTGTCG

Cxxc1 qPCR Reverse: CTGTGGAGAAGATTTGTGGG

β-actin qPCR Forward: CTGTCCCTGTATGCCTCTG

β-actin qPCR Reverse: ATGTCACGCACGATTTCC

CD4$^+$T and YFP$^+$ T_{reg} cells adoptive transfer in Experimental autoimmune

encephalomyelitis (EAE)

CD4$^+$YFP$^+$ T_{reg} cells from *Foxp3*[cre] and *Foxp3*[cre] *Cxxc1*[fl/fl] mice were enriched using the Mouse

CD4 T Cell Isolation Kit (480005; Biolegend) and then sorted using the BD Aria II flow

cytometer. Naïve CD4$^+$T cells from 2D2 (MOG35-55-specificTCR transgenic) mice were

isolated using the Mouse CD4 Naïve T cell Isolation Kit480039 (480039; Biolegend). As

previously described[24], 2D2 naïve CD4$^+$ T cells alone (5 x 10^5 per mouse), or 2D2 naïve CD4$^+$ T

cells (5x 10^5 per mouse) together with WT or $Foxp3^{cre}Cxxc1^{fl/fl}$ T_{reg} cells (2x10^5 per mouse), were

transferred into $Rag1^{-/-}$ mice via the tail vein. One day after cell transfer, the recipient mice were

inoculated   subcutaneously   (s.c.)   with   200μg   MOG35-55   peptide

(MEVGWYRSPFSRVVHLYRNGK; GenemeSynthesis) emulsified in complete Freund's

adjuvant (CFA) (F5506; Sigma). Intravenous administration of 200 ng of Pertussis toxin (181;

List Biological Laboratories) was performed on days 0 and 2 after peptide inoculation. The

severity of EAE was monitored and blindly graded using a clinical score from 0 to 5: 0, no

clinical signs; 1, limp tail; 2, paraparesis (weakness, incomplete paralysis of one or two hind

limbs); 3, paraplegia (complete paralysis of two hind limbs); 4, paraplegia with forelimb

weakness or paralysis; 5, dying or death.

Isolation lymphocytes from the central nervous system（CNS）

On day 14 after EAE induction, mice were perfused with transcardially administered PBS to

eliminate contaminating blood cells in the central nervous system (CNS). The forebrain and

cerebellum were dissected to expose the spinal cord, which was then carefully removed from the

spinal canal. The fresh spinal cord was harvested and cut into 2 mm pieces. The CNS tissue

pieces were homogenized using a syringe and passed through a 70 µM cell strainer to obtain a

single-cell   suspension.   The   single-cell   suspension   was   digested   with   collagenase   D

(2μg/ml;11088858001; Roche) and deoxyribonuclease I (DNase I; 1 μg/ml; DN25;

Sigma-Aldrich) at 37°C for 20 minutes under rotation. After digestion, the cell suspension was

centrifuged to pellet the cells. The cell pellets were resuspended in 40% Percoll and layered onto

a discontinuous Percoll gradient. Centrifugation at 80% Percoll allowed for the separation of

cells at the 40–80% Percoll interface, which were collected as CNS mononuclear cells. The

collected CNS mononuclear cells were washed with PBS to remove any remaining Percoll. CNS

mononuclear cells were stimulated for 4 hours with PMA and ionomycin in the presence of

Brefeldin A to induce cytokine production. After stimulation, cells were fixed, rendered

permeable, and stained with appropriate antibodies for intracellular cytokine detection.

Histological analyses

The lungs, skin, liver, and colon were excised from three-week-old mice. Prior to histological

analysis, the samples were fixed in formalin, embedded in paraffin, and stained with hematoxylin

and eosin. For CNS histology, spinal cords were fixed in 4% paraformaldehyde,

paraffin-embedded, sectioned, and stained with Luxol Fast Blue and hematoxylin and eosin

(H&E). To examine colon histology, colons from $Rag1^{-/-}$ hosts were similarly processed and

stained with H&E.

Adoptive transfer colitis model

Colitis was induced following the protocol described[25]. In brief, CD4$^+$ YFP$^+$ T$_{reg}$ cells were

isolated from 3-week-old CD45.2$^+$$Foxp3^{cre}Cxxc1^{fl/fl}$ and CD45.2$^+$ $Foxp3^{cre}$ mice. A total of 2x10^5

T$_{reg}$ cells from each group were mixed with 4x10^5 T$_{eff}$ cells (CD45.1$^+$CD4$^+$CD45RB^{hi}) sorted

from CD45.1$^+$ mice and transferred into the $Rag1^{-/-}$ mice via intraperitoneal injection. T$_{eff}$ cells

alone were transferred as a control group. Mouse body weight was measured weekly

post-adoptive transfer. The percentage change in body weight was calculated by comparing the

current weight with the initial weight on day 0. Mice were euthanized when any had reached 80%

of their initial body weight. The large intestines were sectioned into 4μm thick slices and stained

with hematoxylin.

In vitro T$_{reg}$ suppression assay

Naïve CD4$^+$ T cells isolated from WT mice were labeled with CFSE (C34554; Invitrogen).

CD4$^+$YFP$^+$ T$_{reg}$ cells from *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice were cultured with naïve CD4$^+$ T

cells (1□10^5 cells) at various ratios in the presence of 2μg/mL anti-CD3(16-0031-85; Invitrogen)

and 3μg/mL anti-CD28(16-0281-85; Invitrogen). On day 3, cells were analyzed by flow

cytometry.

CUT&Tag

CUT&Tag assays of CD4$^+$YFP$^+$ T$_{reg}$ cells were conducted as previously described[26]. Briefly,

approximately 1x10^5 single cells were carefully pipetted into wash buffer twice. The pelleted

cells were resuspended in wash buffer, activated concanavalin (BP531; Bangs Laboratories), and

incubated for 15 minutes at room temperature. Cells bound to the beads were resuspended in

Dig-Wash Buffer and incubated with a 1:50 dilution of primary antibodies (rabbit anti-H3K4me3,

Active Motif,39016; rabbit anti-CXXC1, abcam, ab198977; normal IgG, Cell Signaling, 2729) at

4 □ overnight. The beads were incubated with a secondary antibody (goat anti-rabbit IgG;

SAB3700883; Sigma-Aldrish) diluted 1:100 in Dig-Wash buffer for 60 minutes at room

temperature. Cells were treated with Hyperactive pG-Tn5 Transposase (S602; Vazyme) diluted in

Dig-300 Buffer for 1 hour at room temperature. The cells were subsequently resuspended in

Tagmentation buffer (10mM MgCl2 in Dig-300 Buffer) and incubated at 37□ for 1 hour. To halt

tagmentation, 10μl was spiked with 0.5M EDTA, 3μl with 10%SDS, and 3μl with 20mg/ml

Proteinase K and incubated at 55□ for 1h. DNA library amplification was performed according

to the manufacturer's instructions and purified using VAHTS DNA Clean Beads (N411; Vazyme).

Libraries were sequenced on the Illumina NovaSeq platform (Annoroad Gene Technology).

CUT&Tag and ChIP-seq data analysis

FOXP3 ChIP-seq data was obtained from GSE121279. H3K27me3 ChIP-seq data was obtained

from GSE14254. CUT&Tag and ChIP-seq reads were trimmed to 50 bp and aligned against the

mouse genome build mm9 using Bowtie2(v2.3.4.1) with default parameters. All PCR duplicates

and unmapped reads were removed. Peak calling was performed using MACS2(v2.1.1.20160309)

and signal tracks for each sample were generated using the 'wigToBigWig' utility of UCSC. We

classified the H3k4me3 peaks around TSSs into three groups: broad (>5 kb), medium (1-5kb),

and narrow (<1 kb). The top 5% of the widest peaks were considered as broad peaks. The

average intensity profiles were generated using deepTools (v2.5.4). Motif analysis was

performed using the 'findmotifsGenome.pl' command inHomer2 package. Epigenetic factors

were identified using the Epigenetic Factor Database(https://epifactors.autosome.org/) and then

screened for those that exclusively regulate the expression of their target genes by modulating

the deposition of H3K4me3.

Genomic distribution was analyzed using the "genomation" R package. GO pathway analysis

was performed using the "clusterProfiler" R package. The sequencing information of CUT&Tag

data used in this study is summarized in Supplementary Table S1.

Clustering analysis

257 Promoters were defined as ±2 kb regions flanking the annotated TSS. Reads in promoters were

258 counted using the "coverage" command in bedtools (v2.26.0) and further normalized to RPKM.

259 The k-means clustering of H3K4me3 and H3K27me3 enrichment at promoters was conducted

260 using the "kmeans" function in R.

RNA-seq and data analysis

Total RNA was extracted from sorted CD4$\square$YFP$\square$ T$_{reg}$ cells using the RNeasy Plus Mini Kit

(Qiagen, #74134), following the manufacturer's protocol. RNA-Seq libraries were constructed

and sequenced by Haplox (Nanchang, China), using an Illumina platform with paired-end reads

of 150 bp. RNA-seq data was obtained from GSE82076. Raw reads were trimmed to 50 bp and

mapped to the mouse genome (mm9) using TopHat (v2.1.1) with default parameters. Only

uniquely mapped reads were kept for downstream analysis. The RNA abundance of each gene

was quantified using Cufflinks (v2.2.1).

Single-Cell RNA-Sequencing

A total of 300,000 sort-purified CD4$^+$YFP$^+$ T$_{reg}$ cells from *Foxp3*[cre] and *Foxp3*[cre] *Cxxc1*[fl/fl] were

resuspended in BD Pharmingen Stain Buffer (FBS) (554656; BD). Single cells were isolated

using a chromium controller (BD platform, BD Bioscience) according to the manufacturer's

instructions, as previously described[27]. The single cells were labeled with sample tags using the

BD Mouse Immune Single-Cell Multiplexing Kit (633793; BD). Following standard protocols,

cDNA amplification and library construction were performed to generate scRNA-seq libraries.

Targeted scRNA-seq data processing

The raw FASTQ files were processed by BD Rhapsody using the Targeted analysis pipeline.

After alignment and filtering, the distribution-based error correction(DBEC)-adjusted molecules

were loaded into R studio(version 4.3.2). All subsequent analyses were performed using the

package Seurat (version 4.4.0) with default parameters. Specifically, the scRNA-seq data counts

were log-normalized. All targeted genes were scaled and then were used for Principal

components analysis (PCA). The batch effects were removed by the HarmonyMatrix function in

the Harmony package (version 1.1.0). The first 20 principal components were used to calculate

nonlinear dimensionality reduction using RunUMAP. Differential gene expression (DEGs)

between clusters was assessed using the FindAllMarkers function. The clusters were then

annotated based on DEGs. Barplots were generated using ggplot2 (version 3.4.4). Heatmaps

were generated using pheatmap (version 1.0.12).

Analysis of the single-cell TCR-seq repertoire

Raw V(D)J fastq reads were processed using BD Rhapsody Pipeline and then were analyzed

using scRepertoire (version 1.12.0). The TCR clonotype was called using the nucleotide

sequence of the CDR3 region for both TCR alpha and beta chains. For cells with multiple chains,

the top two clonotypes with the highest expression were selected for downstream analysis.

Clonal overlap between different cell types was calculated using the clonalOverlap function of

scRepertoire. A clonotype was defined as expansion if it could be detected in at least two cells.

scRNA-seq trajectory analysis

UMAP embeddings obtained from the Seurat package were projected into the Slingshot(version

2.10.0) package to construct pseudotime Trajectories for T_{reg} cells. Naïve subsets were set as the

root state.

Whole genome bisulfite sequencing (WGBS) and data analysis

Sorted $CD4^+YFP^+$ T_{reg} cells($3\square10^6$) were lysed in cell lysis buffer to release DNA. The

bisulfite-treated DNA was used to prepare the sequencing library. DNA libraries were transferred

to the Illumina Platform for sequencing using 150 bp paired-end reads. Raw reads were trimmed

using TrimGalore (v0.4.4) with default parameters. Subsequently, the reads were mapped against

the mm9 reference genome using Bismark v0.19.0 with parameters '--bowtie2'. PCR duplicates

were removed and the methylation levels were calculated using 'bismark_methylation_extractor'.

We calculated the mean CpG methylation levels of various genome elements: promoter, 5′-UTR,

exon, intron, 3′-UTR, genebody, intergenic, CGIs, and repeats using in-house scripts. The

sequencing information of WGBS data used in this study is summarized in Supplementary Table

S2.

Statistical analysis

311   The statistical significance analysis was performed using Prism 8.0 (GraphPad). Error bars are

312   presented as mean $\pm$ SD. *P* values of < 0.05 were deemed statistically significant (*P < 0.05, **P

313   < 0.01, ***P < 0.001 and ****P < 0.000). Statistical analyses were performed with unpaired

314   Student's *t*-test or two-way ANOVA and Holm–Sidak post hoc test.

Results

FOXP3 binds regulatory loci primed for activation and repression in T_{reg} cells

FOXP3-mediated gene expression is well-recognized, with several studies highlighting its dual

role as both a transcriptional activator and repressor [28-32]. However, the regulation of T_{reg}-specific

gene transcription by FOXP3 through intermediary mechanisms, such as epigenetic factors,

remains incompletely understood. To address this, we employed CUT&Tag to generate global

H3K4me3 maps in T_{reg} cells and compared these with an H3K27me3 ChIP-Seq dataset from Wei

et al.[33] at previously identified FOXP3-bound loci(Konopacki *et al.* 2019)[34]. This comparison

aimed to identify FOXP3-dependent genes associated with H3K4me3 and H3K27me3 deposition.

As expected, H3K4me3 was enriched at gene promoters (fig. S1, A and B). A Venn diagram

revealed a significant overlap between FOXP3 binding sites and H3K4me3 peaks, with minimal

overlap with H3K27me3 peaks (Fig. 1, A and B). The overlapping regions between FOXP3

binding sites and H3K4me3 or H3K27me3 peaks were predominantly located at promoters (Fig.

1, A and B). We then clustered the promoters into four groups based on the enrichment of

H3K4me3 and H3K27me3. Clusters 1 and 3 showed strong enrichment of H3K4me3; cluster 2

was enriched with H3K27me3; and cluster 4 showed weak enrichment of both modifications.

FOXP3 enrichment analysis revealed that FOXP3 preferentially binds to the promoters of

clusters 1 and 3, which display high levels of H3K4me3 (Fig. 1C). Correspondingly, genes in

these clusters exhibited high transcription levels, as demonstrated by our reanalysis of previously

published RNA-seq data (Oh *et al.* 2017) shown in the right panels of Figure 1D[35]. In contrast,

genes with H3K27me3 enrichment at their promoters were transcribed at low levels. Gene

Ontology (GO) analysis of these four clusters revealed that cluster 1 was significantly enriched

in genes involved in mRNA processing, covalent chromatin modification, and histone

modification, while cluster 3 was enriched in genes related to DNA repair and mitochondrion

organization (Fig. 1E). Cluster 2, on the other hand, was enriched in the pattern specification

process, whereas cluster 4 showed no significant correlation with T_{reg} cells. As expected,

signature T_{reg} cell genes such as *Tnfrsf18*, *Nrp1*, *Stat5a*, *Lag3*, *Icos*, and *Pdcd1* were enriched in

clusters 1 and 3, which exhibited strong H3K4me3 marks (fig. S1C). Conversely, FOXP3-bound

sites in cluster 2 were specifically enriched   for H3K27me3, including genes like *Hic1*, *Trp73,*

and *Rnf157*, which are associated with inflammatory responses(fig. S1C). These observations

support the hypothesis that FOXP3 primarily functions as an activator by promoting H3K4me3

deposition in T_{reg} cells.

CXXC1 interacts with FOXP3 and binds H3K4me3-enriched sites in T$_{reg}$ cells

We conducted an enrichment analysis of known motifs at the overlapping peaks of FOXP3 ChIP-seq and H3K4me3 CUT&Tag in T$_{reg}$ cells to identify epigenetic factors that directly interact with FOXP3 to mediate chromatin remodeling and transcriptional reprogramming. Motif analysis of the overlapping peaks between FOXP3 binding sites and regions enriched in H3K4me3 revealed that, in addition to transcription factors, the most abundant motif associated with H3K4me3 was the epigenetic factor CXXC1(fig. S2A). To investigate this further, we performed CUT&Tag for endogenous CXXC1 in T$_{reg}$ cells to examine the genome-wide co-occupancy of CXXC1 and FOXP3. Over half of these CXXC1 binding sites were located at promoter regions (fig. S2B). Additionally, CXXC1 exhibited strong binding at transcription start site (TSS) and CpG islands (CGIs)(Fig. 2A and fig. S2C). As illustrated by the Venn diagram (Fig. 2B), more than half of the FOXP3-bound genes and H3K4me3-enriched genes were also bound by CXXC1. Similarly, more than half of CXXC1 peaks were overlapped with FOXP3 peaks (fig. S2D). Furthermore, the CXXC1-specific and FOXP3-specific binding sites also demonstrated modest binding of FOXP3 and CXXC1, respectively (Fig. 2C). These findings indicate that FOXP3 and CXXC1 share a substantial number of target genes in T$_{reg}$ cells. To confirm this interaction, we further validated the reciprocal immunoprecipitation of both endogenous and exogenous CXXC1 and FOXP3 (Fig. 2D and fig. S2E). An immunofluorescence assay revealed predominant colocalization of CXXC1 with FOXP3 in the nucleus (Fig. 2E). Overall, these results suggest that CXXC1 primarily functions as a coactivator

367   of FOXP3-driven transcription in T_{reg} cells.

Complete ablation of *Cxxc1* in T_{reg} cells leads to a fatal autoimmune disease

To investigate the role of CXXC1 in T_{reg}-cell homeostasis and function, we generated

Foxp3^Cre*Cxxc1*^fl/fl mice (conditional knockout [cKO] mice) by crossing *Cxxc1*^fl/fl with

Foxp3^YFP-Cre [36] mice, thereby specifically deleting *Cxxc1* in T_{reg} cells. The effective depletion of

Cxxc1 in T_{reg} cells was confirmed through quantitative PCR (qPCR) and Western blotting (fig.

S3A). Notably, cKO mice appeared normal at birth but later exhibited spontaneous mortality

starting around three weeks of age (Fig. 3A). Deletion of *Cxxc1* in T_{reg} cells led to the

development of severe inflammatory disease, characterized by reduced body size, stooped

posture, crusting of the eyelids, ears, and tail, and skin ulceration, particularly on the head and

upper back (Fig. 3, B and C). Additionally, cKO mice developed extensive splenomegaly and

lymphadenopathy (Fig. 3D). Histopathological analysis revealed massive lymphocyte and

myeloid cell infiltration in the skin, lungs, liver sinusoids, and colon mucosa (Fig. 3E). Moreover,

cKO mice displayed a significant increase in CD8$^+$ T cell percentages (fig. S3B), along with a

marked rise in cells exhibiting an effector/memory phenotype (CD44hi CD62Llo) (Fig. 3F).

Furthermore, T cells from cKO mice produced elevated levels of IFN-γ, IL-17, and IL-4 in CD4$^+$

T cells, as well as increased IFN-γ production in CD8$^+$ T cells (Fig. 3G and fig. S3, C and D).

These phenotypes closely resembled those observed in *Foxp3*-deficient mice [37] or mice with

depleted T_{reg} cells[38], suggesting a deficiency in immune suppression.

CXXC1 is necessary for the maintenance of T_{reg}-cell suppressive activity

Despite the development of severe autoimmune disease, we observed an increase in both the absolute number and percentage of FOXP3$^+$ T_{reg} cells in the lymph nodes (fig. S4A). The expression level of the FOXP3 protein was only slightly altered in *Cxxc1*-deficient T_{reg} cells (fig. S4B). In an *in vitro* suppression assay, T_{reg} cells from *Foxp3*^{Cre}*Cxxc1*$^{fl/fl}$ and WT mice exhibited similar suppressive effects on naïve T (Tn) cell proliferation (fig. S4C). The expression of the hallmark T_{reg}-cell marker CTLA-4 showed a modest increase in *Cxxc1*-deficient T_{reg} cells compared to WT T_{reg} cells, while the expression of GITR remained unchanged (fig. S4D). To further assess the suppressive capacity of *Cxxc1*-deficient T_{reg} cells *in vivo*, we employed the experimental autoimmune encephalomyelitis (EAE) model. Naïve CD4$^+$ T cells from 2D2 mice were co-transferred with T_{reg} cells from either *Foxp3*^{Cre} or *Foxp3*^{Cre} *Cxxc1*$^{fl/fl}$ mice into *Rag1*$^{-/-}$ recipients, and EAE was induced in these recipient mice. Mice that received only naïve CD4$^+$ T cells from 2D2 mice developed more severe EAE symptoms (Fig. 4A). The addition of WT T_{reg} cells from *Foxp3*^{Cre} mice slightly mitigated EAE progression and reduced Th17 cells in the spinal cord (Fig. 4, A to D). In contrast, *Foxp3*^{Cre} *Cxxc1*$^{fl/fl}$ T_{reg} cells failed to suppress EAE (Fig. 4, A to D), and the cKO mice showed a significant reduction in T_{reg} cells frequency in CNS tissues(Fig. 4E). Finally, we examined the role of CXXC1 in T_{reg} cell-mediated suppression using T cell transfer-induced colitis, in which naïve T cells were transferred to *Rag1*$^{-/-}$ recipients either alone or together with WT or *Foxp3*^{Cre} *Cxxc1*$^{fl/fl}$ T_{reg} cells. The transfer of naïve T cells led to weight loss and intestinal pathology in recipient mice(Fig. 4, F and G). Mice receiving WT

T$_{reg}$ cells continued to gain weight (Fig. 4F) , whereas those that received T$_{reg}$ cells from cKO

mice were unable to prevent colitis and exhibited a reduced percentage of T$_{reg}$ cells (Fig. 4, F to

H). These findings underscore the critical role of CXXC1 in maintaining T$_{reg}$ cell function *in*

vivo.

T$_{reg}$ cell lineage stability and proliferation depend upon CXXC1

T$_{reg}$ cells harbor a diverse T-cell receptor (TCR) repertoire, which likely play a critical role in

their immune suppression function [39-41]. To explore the role of CXXC1 in T$_{reg}$-mediated

suppression, we performed single-cell RNA sequencing (scRNA-seq) combined with TCR

sequencing (TCR-seq) on CD4$^+$YFP$^+$ T$_{reg}$ cells isolated from mouse lymph nodes. After quality

control and removal of doublets, 18,577 cells were retained for further analysis. Through

unsupervised clustering and uniform manifold approximation and projection (UMAP) analysis,

we identified eight distinct T$_{reg}$ cell clusters based on the expression of well-characterized

markers, with a particular focus on two clusters of activated T$_{reg}$ cells that exhibited higher

expression of markers and gene sets relative to naïve T$_{reg}$ cells (Fig. 5A and fig. S5, A to C). A

comparison between *Cxxc1*-deficient and WT T$_{reg}$ cells within each cluster revealed a reduction

in *Cxxc1*-deficient cells in the naïve subsets, while an increase was observed in the Gzmb$^+$ and

H2-Eb1$^+$ subsets (Fig. 5B). To further elucidate the transition of T$_{reg}$ cells along a dynamic

biological timeline, we constructed pseudo-time trajectories using Slingshot [42]. The pseudo-time

gradient depicted a progression from quiescent to activated T$_{reg}$ cells, ultimately encompassing

the Gzmb$^+$ and H2-Eb1$^+$ subsets (Fig. 5C). Given the antigen-specific suppression capabilities of

T_{reg} cells [43,44], we examined their clonal expansion. The analysis revealed that expanded WT

TCR clonotypes (n□≥□2) were predominantly distributed among the Nt5e$^+$ subsets, while

Cxxc1-deficient T_{reg} cells showed expanded clonotypes primarily within the Gzmb$^+$ and H2-Eb1$^+$

subsets(Fig. 5D and Fig. S5D). TCR sharing analysis indicated significant clonotype sharing

among various clusters of WT T_{reg} cells, suggesting a degree of homogeneity. However, the

reduced TCR sharing in *Cxxc1*-deficient cells implies that decreased TCR diversity may impair

the suppressive activity of T_{reg} cells (Fig. 5E) [41]. Furthermore, the *Cxxc1*-deficient group

exhibited lower expression of several T_{reg}-specific genes associated with suppressive functions,

such as*Nt5e*, *Il10*, *Pdcd1*, *Klrg1*, as well as genes that inhibit effector T cells differentiation,

including *Sell* and *Tcf7* (Fig. 5F and fig. S5E). Conversely, *Cxxc1*-deficient T_{reg} cells

demonstrated elevated expression of *Gzmb*, *Il2ra*, and *Cd69* compared to WT, reflecting a profile

indicative of increased activation (Fig. 5, F and G). Additionally, we observed increased

expression of genes linked to Th1-type inflammation, such as *Ifng*, *Tbx21,* and *Hif1a*, in

Cxxc1-deficient T_{reg} cells, likely due to extreme inflammatory conditions (Fig. 5, F to H). The

proportion of FOXP3$^+$Ki67$^+$ T_{reg} cells was lower in cKO mice compared to WT mice (Fig. 5I).

These findings underscore the crucial role of CXXC1 in maintaining T_{reg} cell stability.

***Cxxc1* deletion leads to a reduction in essential T_{reg} cell markers**

To confirm that the deficiency in T_{reg} cell function in T_{reg}-specific *Cxxc1*-deficient animals is due

to intrinsic defects caused by *Cxxc1* deficiency, rather than severe autoimmune inflammation in

Foxp3^{Cre} *Cxxc1*$^{fl/fl}$ mice, we examined *Cxxc1*-sufficient and *Cxxc1*-deficient T_{reg} cell subsets in

heterozygous $Foxp3^{Cre/+}$ $Cxxc1^{fl/fl}$ (designated as "het-KO") and littermate $Foxp3^{Cre/+}$ $Cxxc1^{fl/+}$

(designated as "het-WT") female mice (Fig. 6A). Notably, het-KO female mice did not exhibit

overt signs of autoimmunity, as random X-chromosome inactivation led to the coexistence of

both *Cxxc1*-cKO and *Cxxc1*-WT T_{reg} cells. However, both the frequency and absolute numbers

of $FOXP3^{+}YFP^{+}$ T_{reg} cells within the total T_{reg} population were reduced in het-KO mice

compared to their counterparts in het-WT littermates, indicating that *Cxxc1* deficiency imposes a

competitive disadvantage on T_{reg} cells (Fig. 6B). Additionally, *Cxxc1*-deficient YFP^{+} T_{reg} cells

failed to upregulate the proliferation marker Ki-67 (Fig. 6C). Moreover, YFP^{+} T_{reg} cells in

het-KO female mice showed reduced expression of key genes essential for suppressive function,

including *Icos*, *Il2ra*, *Ctla4*, *and Tnfrsf18*, compared to $YFP^{-}T_{reg}$ cells from the same mice (Fig.

6D). To investigate the molecular program affected by the deletion of *Cxxc1* in T_{reg} cells, we

performed RNA-sequencing (RNA-seq) analysis on $CD4^{+}YFP^{+}$ T_{reg} cells isolated from het-WT

and het-KO mice. We then conducted a differential gene expression (DGE) analysis based on the

RNA-seq data. Among all expressed genes, 865 were upregulated ("Up" genes) and 761 were

downregulated ("Down" genes) by ≥1.5-fold in $CD4^{+}YFP^{+}$ T_{reg} cells from het-KO mice

compared to het-WT mice, with a false discovery rate (FDR)–adjusted *P*-value cutoff of 0.05 (fig.

S6A). Gene Ontology (GO) enrichment analysis revealed that the downregulated genes in

Cxxc1-deficient T_{reg} cells were predominantly enriched in pathways related to the negative

regulation of immune system process and regulation of cell–cell adhesion (fig. S6B). The

Cxxc1-deficient T_{reg} cells also showed reduced expression of several genes associated with T_{reg}

cell suppressive function, including *Il10*, *Tigit*, *Lag3*, *Icos*, *Nt5e*(encoding CD73) and *Itgae*

(encoding CD103) (fig. S6C). Thus, although YFP$^-$ WT T$_{reg}$ cells can prevent autoimmunity in

het-KO mice, the absence of *Cxxc1* in YFP$^+$ T$_{reg}$ cells leads to the dysregulation of key T$_{reg}$ cell

markers under steady-state conditions.

The FOXP3-CXXC1 complex regulates the expression of key factors in T$_{reg}$ cells that are

associated with the breadth of H3K4me3

CXXC1 binds to unmethylated CpG DNA via its N-terminal CXXC finger domain, facilitating

its interaction with DNA methyltransferase 1 (DNMT1). This binding stabilizes the DNMT1

protein, thereby regulating DNA methylation [45,46]. To investigate whether CXXC1 depletion

affects DNA methylation in T$_{reg}$ cells, we performed whole genome bisulfite sequencing (WGBS)

on T$_{reg}$ cells isolated from both WT and cKO mice. On average, *Cxxc1*-deficient T$_{reg}$ cells

exhibited no significant changes in DNA methylation at gene loci or across genome-wide CpG

sites, irrespective of chromosomal region (fig. S7, A to C). Furthermore, *Cxxc1* knockout T$_{reg}$

cells did not show a significant increase in DNA methylation at key T$_{reg}$ signature gene loci (fig.

S7D). Given the pivotal role of MLL4-mediated H3K4me1 in establishing the enhancer

landscape and facilitating long-range chromatin interactions during T$_{reg}$ cell development[15], we

performed CUT&Tag to assess changes in H3K4me1 levels in *Cxxc1*-deficient T$_{reg}$ cells. This

analysis revealed that H3K4me1 levels were similar in both WT and *Cxxc1*-deficient T$_{reg}$ cells

(fig. S7, E to G).

While H3K4me3 modifications typically form sharp 1-to 2-kb peaks around promoters, some

genes exhibit broader H3K4me3 regions, referred to as broad H3K4me3 domains

(H3K4me3-BDs), which can extend to cover part or all of the gene's coding sequences (up to 20

kb) [47,48]. Broad H3K4me3 domains are preferentially associated with genes essential for the

identity or function of specific cell types [47,49] and have been implicated in enhancing

transcriptional elongation and increasing enhancer activity [49]. To further explore the connection

between broad H3K4me3 domains and the expression of immune-regulatory genes, we analyzed

genes containing broad H3K4me3 regions. We classified the H3K4me3 domains surrounding

transcription start sites (TSSs) into three categories: broad (more than 5 kb), medium (between 1

and 5 kb), and narrow (less than 1 kb) (Fig. 7A). Notably, *Cxxc1*-deficient T_{reg} cells exhibited

weaker H3K4me3 signals compared to WT cells within the broad H3K4me3 domains where

CXXC1 binding is prominent (Fig. 7, A and B). Using the criteria established by Benayoun *et*

al.[47], which defines the top 5% of the widest H3K4me3 domains as BDs, we observed similar

enrichment results (fig. S7, H and I). We then compared three groups of genes: BD-associated

genes with reduced H3K4me3 levels following *Cxxc1* deletion, genes with direct CXXC1

binding, and genes with direct FOXP3 binding. The Venn diagram revealed that the majority of

genes (283 out of 294, 96%) with CXXC1 binding and reduced H3K4me3 levels overlap with

FOXP3-bound genes, suggesting that CXXC1 presence enhances FOXP3 binding within the

broad H3K4me3 domain (Fig. 7C). Furthermore, GO term analysis indicated that BD-associated

genes are enriched in biological processes related to the negative regulation of immune system

processes (Fig. 7D). Genome browser views displayed the enrichments of FOXP3, CXXC1, and

H3K4me3 at key signature genes in T_{reg} cells, such as *Ctla4, Il2ra, Icos*, and *Tnfrsf18*, with

lower H3K4me3 densities observed at these loci in *Cxxc1*-deficient T_{reg} cells (Fig. 7E). Similar

patterns were observed at core genes involved in T_{reg} homeostasis and suppressive function (e.g.,

Lag3, *Nt5e*, *Ikzf4*, and *Cd28*) (fig. S7, J)[50,51]. These findings suggest that CXXC1 and FOXP3

collaboratively promote sustained T_{reg} cell homeostasis and function by preserving the H3K4me3

modification at key T_{reg} cell genes.

Discussion

T_{reg} cells specifically express the transcription factor FOXP3, which is essential for maintaining

T_{reg} lineage stability and suppressive function[52]. However, FOXP3 alone is insufficient to fully

regulate the transcriptional signature and functionality of T_{reg} cells; its interaction with protein

partners is crucial for this regulation. In this study, we identify CXXC1 as a key transcription

factor that, by acting as a critical cofactor of FOXP3, orchestrates the T_{reg} transcriptional

program , which in turn plays a vital role in T_{reg} cell homeostasis and function.

FOXP3 is known to promote histone H3 acetylation at the promoters and enhancers of its target

genes, such as *Il2ra*, *Ctla4*, and *Tnfrsf18*, following T_{reg} cell activation, thereby functioning as a

transcriptional activator[53,54]. Conversely, FOXP3 can act as a repressor by silencing target genes

like *Il2* and *Ifng* through the induction of histone H3 deacetylation, mediated by the recruitment

of histone deacetylases and transcriptional co-repressors[10,55]. Additionally, FOXP3 exerts this

repression by recruiting the Ezh2-containing polycomb repressive complex to target genes during

activation, as FOXP3-repressed genes are associated with H3K27me3 deposition and reduced

chromatin accessibility[11,56]. Recent studies have begun to explore the biological significance of

H3K4me3 breadth, revealing a positive correlation between H3K4me3 breadth and gene

expression [49,57,58]. These studies also suggest that H3K4me3 breadth contributes to defining

specific cell identities during development and disease, including systemic autoimmune diseases

like systemic lupus erythematosus and various cancers [47,59-62]. Our study demonstrates that the

loss of *Cxxc1* leads to reduced H3K4me3 levels, predominantly in genes with broader peaks,

such as *Il2ra, Tnfrsf18, Ctla4 and Icos*, directly impairing their suppressive function in T_{reg} cells.

Although our research primarily focused on the role of CXXC1 in T_{reg} cells, it is plausible that

similar mechanisms may be operative in other cell types.

The T_{reg}-specific deletion of *Cxxc1* leads to a rapid and fatal autoimmune disorder, characterized

by systemic inflammation and tissue damage, underscoring the essential role of CXXC1 in

maintaining immune self-tolerance within T_{reg} cells. Interestingly, despite this severe phenotype,

our findings show that H3K4me1 levels were comparable between WT and *Cxxc1*-deficient T_{reg}

cells. In contrast, MLL4 is critical for T_{reg} cell development in the thymus, primarily by

regulating H3K4me1, though it is not required for peripheral T_{reg} cell function[15]. Numerous

studies investigating the effectors of T_{reg} cell-mediated suppression have identified a wide range

of molecules and mechanisms. These include the upregulation of the inhibitory co-stimulatory

receptor CTLA-4, which initiates inhibitory signaling; the sequestration of the T-cell growth

factor IL-2 via CD25; the secretion of inhibitory cytokines, such as interleukin (IL)-10, IL-35

and TGF-β and the activity of ectoenzymes CD39 and CD73 on the T_{reg} cells surface, which

convert extracellular ATP, a potent pro-inflammatory mediator, into its anti-inflammatory

counterpart, adenosine[41,63]. Mechanistically, we demonstrated that CXXC1 interacts with FOXP3

to regulate T_{reg} cell function by trimethylating H3K4 at broad H3K4me3 domains of multiple

genes involved in suppressive functions such as *Il2ra, Nt5e, and Ctla4*. Additionally, MLL1,

another KMT, controls T_{reg} cell activation and function by specifically regulating H3K4

trimethylation at genes encoding key T_{reg}-related molecules such as *Tigit*, *Klrg1*, *Tbx21*, *Cxcr3*,

and serves as a crucial epigenetic regulator in establishing a stable Th1-T_{reg} lineage[14].

Our results offer provide novel insights into the suppressive functions, heterogeneity, and

regulatory mechanisms of T_{reg} cells. Maintaining T_{reg} cell homeostasis and function remains a

significant challenge in harnessing T_{reg} cells for the treatment of autoimmune diseases and the

prevention of graft rejection.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

We thank B. Li (Shanghai Jiao Tong University School of Medicine, Shanghai, China) for

providing *Foxp3*[YFP-Cre] mice, and X. L. Liu (Shanghai Institutes for Biological Sciences, Chinese

Academy of Sciences) for his generous gifts of cell lines. We thank Y. Huang, Y. Li, J. Wan,

C.Guo, and Z. Lin from the Core Facilities, Zhejiang University School of Medicine for their

technical support and S. Hong, Y. Ding, H. Jin, Q. Wang, and X. Zhang from Animal Facilities,

Zhejiang University, for feeding the mice.

Footnotes

This work was supported by grants from the National Natural Science Foundation of China

(32341002, 32030035, 32321002, 32100693 and 32270839), the National Key R & D Program

of China (2023YFA1800202), the Zhejiang Provincial Natural Science Foundation of China

(LZ21C080001, LZ23C070003), Science and Technology Innovation 2030-Major Project

(2021ZD0200405), Key project of the Experimental Technology Program of Zhejiang University

(SZD202203).

Author contributions

L.W. and L.S.: supervision, conceptualization, project administration, and writing—review and

editing. L.W., L.S. and X.M: funding acquisition. X.M: investigation, methodology, project

administration, and writing— original draft. Y.Z.and K.L.: bioinformatics analysis. Y. W., X.L.,

J.C., C.L., Y.Z., S.W., S.T., Q.X., L.D., X.S., and X.G.: investigation and methodology.

Figure legends

Fig. 1. H3K4me3 is required for FOXP3-dependent gene activation in T_{reg} cells.

(**A**)Venn diagram showing overlap of H3K4me3 enriched regions (this study) and FOXP3

binding sites (Konopacki *et al*.2019) in sorted CD4$^+$YFP$^+$ T_{reg} cells (left). Genomic distribution

of overlapped peaks(right). Note that the overlapped peaks are predominantly enriched at

promoters.

(**B**)Venn diagram showing overlap of H3K27me3 enriched regions (Wei *et al.*2009) and FOXP3

binding sites in T_{reg} cells (left). Genomic distribution of overlapped peaks (right). Note that the

overlapped peaks are predominantly enriched at promoters.

(**C**)Heatmap showing enrichment of H3K27me3, H3K4me3, and FOXP3 surrounding the TSS.

Unsupervised K-means clustering was conducted on H3K27me3 and H3K4me3 signals.

(**D**)Heatmap showing gene expression levels in T_{reg} cells (RNA-seq data was obtained from Oh

*et al.*2017) The clusters were consistent as in Fig. 1C.

(**E**)Gene Ontology (GO) pathway analysis of different clusters.

Fig. 2. CXXC1 interacts with FOXP3 in T_{reg} cell.

(**A**)Average CXXC1 CUT&Tag signals around promoters in T_{reg} cells. IgG was used as the

control.

(**B**)Venn diagrams showing the overlap of FOXP3 binding genes, CXXC1 binding genes, and

H3K4me3 enriched genes in T_{reg} cells. Gene promoters covered by FOXP3 binding sites,

CXXC1 binding sites, or exhibited high H3K4me3 levels were defined as FOXP3-bound genes,

CXXC1-bound genes, or H3K4me3-enriched genes.

(**C**)Heatmaps showing FOXP3 ChIP-seq and CXXC1 CUT&Tag signals at indicated regions.

(**D**)Interaction between FOXP3 and CXXC1 was assessed by co-IP (forward and reverse)

using T_{reg} cell lysates.

(**E**)Immunofluorescence for FOXP3 and CXXC1 colocalization in T_{reg} cells. Scale bars, 2 μm.

Fig.3. *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice spontaneously develop a fatal early-onset inflammatory

disorder.

(**A**)Survival curves of *Foxp3*^{cre} (black line) and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ (red line) mice ($n = 20$).

(**B**)Gross body weight of *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice ($n\square=\square6$).

(**C**)A representative image of *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice.

(**D**)Representative images showing the spleen and peripheral lymph nodes from *Foxp3*^{cre} and

Foxp3^{cre}*Cxxc1*$^{fl/fl}$ mice.

(**E**)Hematoxylin and eosin staining of the skin, lung, liver, and colon from *Foxp3*^{cre} and *F*

oxp3^{cre}*Cxxc1*$^{fl/fl}$ mice (scale bar, $100\square$μm).

(**F**)Flow cytometry analysis of CD62L and CD44 expression on peripheral lymph node

CD4$^+$ and CD8$^+$T cells from *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice (left). Right, frequency of

CD44^{low}CD62L^{hi} and CD44^{hi}CD62L^{low} population in CD4$^+$ or CD8$^+$ T cells ($n = 9$).

(**G**)Lymph node cells from *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice were stimulated ex vivo with

PMA + ionomycin for 4h and analyzed for IFN-γ expressing in CD4$^+$ or CD8$^+$ T cells using flow

cytometry (left). Right, percentages of IFN-γ$^+$CD4$^+$ or IFN-γ$^+$CD8$^+$ T cells in the lymph nodes of

Foxp3^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice ($n = 7$).

All mice analyzed were 18-20 days old unless otherwise specified. Error bars show mean ± SD.

The log-rank survival curve was used for survival analysis in A, and multiple unpaired *t*-test or

two-tailed Student's *t*-test were used for statistical analyses in B, F, and G (****$P<0.0001$). The

flow cytometry results are representative of three independent experiments.

Fig. 4. CXXC1 is essential for T$_{reg}$ cells to suppress T cell-mediated EAE and colitis.

(**A**)Mean clinical scores for EAE in *Rag1*$^{-/-}$ recipients of 2D2 CD4$^+$ T cells, either alone or in

combination with *Foxp3*^{cre} or *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice after immunization with MOG35–55,

complete Freund's adjuvant (CFA), and pertussis toxin (n=5).

(**B** and **C**)Representative histology of the spinal cord of $Rag1^{-/-}$ mice after EAE induction.

Hematoxylin and eosin(H&E) staining (upper), Luxol fast blue (F&B) staining (lower). Scale

bars, 50 μm (400×) and 200 μm (100×).

(**D**)Representative flow cytometry plots and quantification of the the percentages of IFNγ$^+$ or

IL-17A$^+$ CD4$^+$Vβ11$^+$ T cells (n=5).

(**E**)Statistical analysis of the percentage CD4$^+$ FOXP3$^+$ T$_{reg}$ cell in CNS tissues 14 days after

EAE induction.

(**F**)Changes in body weight of $Rag1^{-/-}$ mice after colitis induction(n=6).

(**G**)Haematoxylin and eosin (H&E) staining of colons from T cell-induced colitis mice 6 weeks

after T cell transfer. Scale bars, 50 μm (400×) and 200 μm (100×).

(**H**)Statistical analysis of the percentage CD4$^+$ FOXP3$^+$ T$_{reg}$ cell in the spleen, mesenteric lymph

nodes, and colon 6 weeks after colitis induction.

Error bars show mean ± SD. *P* values are determined by a two-tailed Student's *t*-test or two-way

ANOVA and Holm–Sidak post hoc test (A, D, and E). (*P<0.05, **P<0.01, ***P□<□0.001,

****P<0.0001).

Fig. 5. Single-cell transcriptomics reveals distinct T$_{reg}$ cell populations.

(**A**)UMAP plot showing clusters identified based on variable gene expression of sorted

CD4$^+$YFP$^+$ T$_{reg}$ cells. Each dot represents a cell, and each color corresponds to a different

population of cell types. Clustering analysis revealed 8 distinct T$_{reg}$ cell populations.

(**B**)Mean fold changes in cluster abundance between $Foxp3^{cre}$ and $Foxp3^{cre}Cxxc1^{fl/fl}$ mice.

(**C**)Pseudotime trajectories of T_{reg} cells based on Slingshot, color-coded by T_{reg} cell

subpopulations.

(**D**)Visualization of density and clonotype richness across T_{reg} clusters from *Foxp3*[cre] and

Foxp3[cre]*Cxxc1*[fl/fl] mice.

(**E**)TCR sharing of expanded clonotypes across all possible combinations of T_{reg} cells from

Foxp3[cre] and *Foxp3*[cre]*Cxxc1*[fl/fl] mice.

(**F**)Heatmap showing Z scores for the average expression of T_{reg}-specific genes in each cluster

between *Foxp3*[cre] and *Foxp3*[cre]*Cxxc1*[fl/fl] mice.

(**G**-**H**)Representative flow cytometry plots and quantification of (G) expression of CD25, CD69,

ICOS, T-bet, and (H) IFN-γ in CD4$^+$YFP$^+$ T_{reg} cells from *Foxp3*[cre] and *Foxp3*[cre]*Cxxc1*[fl/fl] mice

(n=5 CD25, n=10 CD69, n=5 ICOS, n=6 T-bet, n=5 IFN-γ).

(**I**)Ki-67 expression (left) and frequency (right) in CD4$^+$FOXP3$^+$ T_{reg} cells from *Foxp3*[cre] a

-nd *Foxp3*[cre]*Cxxc1*[fl/fl] mice (n=6).

Error bars show mean ± SD. *P* values are determined by a two-tailed Student's *t*-test (G-H).

(**P<0.01, ***P□<□0.001, ****P<0.0001). The flow cytometry results are representative of

three independent experiments.

Fig. 6. *Cxxc1*-deficient T_{reg} cells in het-KO female mice exhibit reduced suppression

markers under steady-state conditions.

(**A**)Schematic representation of wild-type and Cre-positive T_{reg} cells in female *Foxp3*[Cre/+]mice.

(**B**)Flow cytometry analysis of the YFP$^-$FOXP3$^+$ (WT) and YFP$^+$FOXP3$^+$ (KO) T_{reg} cells in

Foxp3[Cre/+] *Cxxc1*[fl/+] (het-WT) and *Foxp3*[Cre/+] *Cxxc1*[fl/fl] (het-KO) female mice (left), along with

the frequency and absolute numbers of YFP$^+$ cells within the total T$_{reg}$ population (right)(n=5).

(**C**)Flow cytometry analysis of Ki-67expression (left) and MFI (right) in YFP$^-$ and YFP$^+$ cells

within the CD4$^+$FOXP3$^+$ T$_{reg}$ cells from 6- to 8-week-old het-KO female mice (n=4).

(**D**)Representative flow cytometry plots and quantification of ICOS, CD25, CTLA4, and GITR

expression in YFP$^-$ and YFP$^+$ cells within CD4$^+$FOXP3$^+$ T$_{reg}$ cells from het-KO female mice

(n=5).

Error bars show mean ± SD. *P* values are determined by a two-tailed Student's *t*-test (G-H).

(**P<0.01, ***P□<□0.001, ****P<0.0001). The flow cytometry results are representative of

three independent experiments.

Fig. 7. CXXC1 regulates Foxp3-dependent molecule H3K4 trimethylation in T$_{reg}$ cells.

(**A-B**)Heatmaps showing H3K4me3(A)and CXXC1(B) signals centered on narrow, medium, and

broad domains. The top panel shows the average CUT&Tag signals around indicated domains.

(**C**)Venn diagrams showing the overlap of FOXP3 binding genes, CXXC1 binding genes, and

H3K4me3-BD-associated genes with decreased H3K4me3 levels after CXXC1 depletion in T$_{reg}$

cells.

(**D**)Gene Ontology (GO) pathway analysis of the overlap peaks.

(**E**)Representative genome browser view showing the enrichments of FOXP3, CXXC1, and

H3K4me3 in T$_{reg}$ cells.

References

1   Sakaguchi, S., Yamaguchi, T., Nomura, T. & Ono, M. Regulatory T cells and immune

tolerance. Cell 133, 775-787, doi:10.1016/j.cell.2008.05.009 (2008).

2   Josefowicz, S. Z., Lu, L. F. & Rudensky, A. Y. in *Annual Review of Immunology, Vol 30* Vol.

30 *Annual Review of Immunology* (ed W. E. Paul)   531-564 (2012).

3   Singer, B. D., King, L. S. & D'Alessio, F. R. Regulatory T cells as immunotherapy. Frontiers

in Immunology 5, doi:10.3389/fimmu.2014.00046 (2014).

4   Ohkura, N. & Sakaguchi, S. Transcriptional and epigenetic basis of Treg cell development

and function: its genetic anomalies or variations in autoimmune diseases. Cell Research 30,

465-474, doi:10.1038/s41422-020-0324-7 (2020).

5   Littman, D. R. & Rudensky, A. Y. Th17 and Regulatory T Cells in Mediating and

Restraining Inflammation. Cell 140, 845-858, doi:10.1016/j.cell.2010.02.021 (2010).

6   Sakaguchi, S., Vignali, D. A. A., Rudensky, A. Y., Niec, R. E. & Waldmann, H. The

plasticity and stability of regulatory T cells. Nature Reviews Immunology 13, 461-467,

doi:10.1038/nri3464 (2013).

7   Fontenot, J. D., Gavin, M. A. & Rudensky, A. Y. Foxp3 programs the development and

function of CD4+CD25+ regulatory T cells. Nat Immunol 4, 330-336, doi:10.1038/ni904 (2003).

8   Van Gool, F., Nguyen, M. L. T., Mumbach, M. R., Satpathy, A. T., Rosenthal, W. L.,

Giacometti, S., Le, D. T., Liu, W., Brusko, T. M., Anderson, M. S., Rudensky, A. Y., Marson, A.,

Chang, H. Y. & Bluestone, J. A. A Mutation in the Transcription Factor Foxp3 Drives T Helper 2

Effector Function in Regulatory T Cells. Immunity 50, 362-+, doi:10.1016/j.immuni.2018.12.016

(2019).

9   Katoh, H., Qin, Z. S., Liu, R., Wang, L., Li, W., Li, X., Wu, L., Du, Z., Lyons, R., Liu, C.-G.,

Liu, X., Dou, Y., Zheng, P. & Liu, Y. FOXP3 Orchestrates H4K16 Acetylation and H3K4

Trimethylation for Activation of Multiple Genes by Recruiting MOF and Causing Displacement

of PLU-1. Molecular Cell 44, 770-784, doi:10.1016/j.molcel.2011.10.012 (2011).

10   Wang, L., Liu, Y., Han, R., Beier, U. H., Bhatti, T. R., Akimova, T., Greene, M. I., Hiebert, S.

W. & Hancock, W. W. FOXP3(+) regulatory T cell development and function require

histone/protein deacetylase 3. Journal of Clinical Investigation 125, 1111-1123,

doi:10.1172/jci77088 (2015).

11   Arvey, A., van der Veeken, J., Samstein, R. M., Feng, Y., Stamatoyannopoulos, J. A. &

Rudensky, A. Y. Inflammation-induced repression of chromatin bound by the transcription factor

Foxp3 in regulatory T cells. Nature Immunology 15, 580-587, doi:10.1038/ni.2868 (2014).

12   Ruthenburg, A. J., Allis, C. D. & Wysocka, J. Methylation of lysine 4 on histone H3:

Intricacy of writing and reading a single epigenetic mark. Molecular Cell 25, 15-30,

doi:10.1016/j.molcel.2006.12.014 (2007).

13   Takahashi, Y.-h., Westfield, G. H., Oleskie, A. N., Trievel, R. C., Shilatifard, A. & Skiniotis,

G. Structural analysis of the core COMPASS family of histone H3K4 methylases from yeast to

human. Proceedings of the National Academy of Sciences of the United States of America 108,

20526-20531, doi:10.1073/pnas.1109360108 (2011).

14   Wang, T., Guo, J., Li, L., Jin, Q., Zhang, F., Hou, B., Zhang, Y. & Zhou, X. The histone

lysine methyltransferase MLL1 regulates the activation and functional specialization of

regulatory T cells. Cell Reports 43, doi:10.1016/j.celrep.2024.114222 (2024).

15   Placek, K., Hu, G., Cui, K., Zhang, D., Ding, Y., Lee, J.-E., Jang, Y., Wang, C., Konkel, J. E.,

Song, J., Liu, C., Ge, K., Chen, W. & Zhao, K. MLL4 prepares the enhancer landscape for Foxp3

induction via chromatin looping. Nature Immunology 18, 1035-+, doi:10.1038/ni.3812 (2017).

16   Yang, Y., Yang, Y., Chan, K. & Couture, J.-F. Analyzing the impact of CFP1 mutational

landscape on epigenetic signaling. Faseb Journal 35, doi:10.1096/fj.202100427R (2021).

17   Tate, C. M., Lee, J.-H. & Skalnik, D. G. CXXC finger protein 1 restricts the Setd1A histone

H3K4   methyltransferase   complex   to   euchromatin.   Febs   Journal   277,   210-223,

doi:10.1111/j.1742-4658.2009.07475.x (2010).

18   Cao, W., Guo, J., Wen, X., Miao, L., Lin, F., Xu, G., Ma, R., Yin, S., Hui, Z., Chen, T., Guo,

S., Chen, W., Huang, Y., Liu, Y., Wang, J., Wei, L. & Wang, L. CXXC finger protein 1 is critical

for T-cell intrathymic development through regulating H3K4 trimethylation. Nat Commun 7,

11687, doi:10.1038/ncomms11687 (2016).

19 Hui, Z., Zhou, L., Xue, Z., Zhou, L., Luo, Y., Lin, F., Liu, X., Hong, S., Li, W., Wang, D., Lu,

L., Wang, J. & Wang, L. Cxxc Finger Protein 1 Positively Regulates GM-CSF-Derived

Macrophage Phagocytosis Through Csf2rα-Mediated Signaling. Frontiers in Immunology 9,

doi:10.3389/fimmu.2018.01885 (2018).

20   Lin, F., Meng, X., Guo, Y., Cao, W., Liu, W., Xia, Q., Hui, Z., Chen, J., Hong, S., Zhang, X.,

Wu, C., Wang, D., Wang, J., Lu, L., Qian, W., Wei, L. & Wang, L. Epigenetic initiation of the

T(H)17 differentiation program is promoted by Cxxc finger protein 1. Science Advances 5,

doi:10.1126/sciadv.aax1608 (2019).

21 Shen, X., Gao, X., Luo, Y., Xu, Q., Fan, Y., Hong, S., Huang, Z., Liu, X., Wang, Q., Chen, Z.,

Wang, D., Lu, L., Wu, C., Liang, H. & Wang, L. Cxxc finger protein 1 maintains homeostasis

and function of intestinal group 3 innate lymphoid cells with aging. Nature Aging 3, 965-+,

doi:10.1038/s43587-023-00453-7 (2023).

22 Cao, W., Guo, J., Wen, X., Miao, L., Lin, F., Xu, G., Ma, R., Yin, S., Hui, Z., Chen, T., Guo,

S., Chen, W., Huang, Y., Liu, Y., Wang, J., Wei, L. & Wang, L. CXXC finger protein 1 is critical

for T-cell intrathymic development through regulating H3K4 trimethylation. Nature

Communications 7, doi:10.1038/ncomms11687 (2016).

23 Zhou, L., Wang, S., Hu, W., Liu, X., Xu, L., Tong, B., Zhang, T., Xue, Z., Guo, Y., Zhao, J.,

Lu, L., Fan, H., Qian, W., Chen, J., Chen, W. & Wang, L. T cell proliferation requires ribosomal

maturation in nucleolar condensates dependent on DCAF13. Journal of Cell Biology 222,

doi:10.1083/jcb.202201096 (2023).

24 Chou, W.-C., Guo, Z., Guo, H., Chen, L., Zhang, G., Liang, K., Xie, L., Tan, X., Gibson, S.

A., Rampanelli, E., Wang, Y., Montgomery, S. A., Brickey, W. J., Deng, M., Freeman, L., Zhang,

S., Su, M. A., Chen, X., Wan, Y. Y. & Ting, J. P. Y. AIM2 in regulatory T cells restrains

autoimmune diseases. Nature 591, 300-+, doi:10.1038/s41586-021-03231-w (2021).

25 Zeng, H., Yang, K., Cloer, C., Neale, G., Vogel, P. & Chi, H. mTORC1 couples immune

signals and metabolic programming to establish T(reg)-cell function. Nature 499, 485-490,

doi:10.1038/nature12297 (2013).

26 Lu, D., Liu, L., Sun, Y., Song, J., Yin, Q., Zhang, G., Qi, F., Hu, Z., Yang, Z., Zhou, Z., Hu,

Y., Zhang, L., Ji, J., Zhao, X., Jin, Y., McNutt, M. A. & Yin, Y. The phosphatase PAC1 acts as a T

cell suppressor and attenuates host antitumor immunity. Nature Immunology 21, 287-+,

doi:10.1038/s41590-019-0577-9 (2020).

27   Chen, J., Liu, K., Luo, Y., Kang, M., Wang, J., Chen, G., Qi, J., Wu, W., Wang, B., Han, Y.,

Shi, L., Wang, K., Han, X., Ma, X., Liu, W., Ding, Y., Wang, L., Liang, H., Wang, L. & Chen, J.

Single-Cell Profiling of Tumor Immune Microenvironment Reveals Immune Irresponsiveness in

Gastric   Signet-Ring   Cell   Carcinoma.   Gastroenterology   165,   88-103,

doi:10.1053/j.gastro.2023.03.008 (2023).

28   Marson, A., Kretschmer, K., Frampton, G. M., Jacobsen, E. S., Polansky, J. K., MacIsaac, K.

D., Levine, S. S., Fraenkel, E., von Boehmer, H. & Young, R. A. Foxp3 occupancy and

regulation of key target genes during T-cell stimulation. Nature 445, 931-935,

doi:10.1038/nature05478 (2007).

29   Zheng, Y., Josefowicz, S. Z., Kas, A., Chu, T.-T., Gavin, M. A. & Rudensky, A. Y.

Genome-wide analysis of Foxp3 target genes in developing and mature regulatory T cells. Nature

445, 936-940, doi:10.1038/nature05563 (2007).

30   Ono, M., Yaguchi, H., Ohkura, N., Kitabayashi, I., Nagamura, Y., Nomura, T., Miyachi, Y.,

Tsukada, T. & Sakaguchi, S. Foxp3 controls regulatory T-cell function by interacting with

AML1/Runx1. Nature 446, 685-689, doi:10.1038/nature05673 (2007).

31   Fu, W., Ergun, A., Lu, T., Hill, J. A., Haxhinasto, S., Fassett, M. S., Gazit, R., Adoro, S.,

Glimcher, L., Chan, S., Kastner, P., Rossi, D., Collins, J. J., Mathis, D. & Benoist, C. A multiply

redundant genetic switch 'locks in' the transcriptional signature of regulatory T cells. Nature

Immunology 13, 972-980, doi:10.1038/ni.2420 (2012).

32   Ohkura, N., Hamaguchi, M., Morikawa, H., Sugimura, K., Tanaka, A., Ito, Y., Osaki, M.,

Tanaka, Y., Yamashita, R., Nakano, N., Huehn, J., Fehling, H. J., Sparwasser, T., Nakai, K. &

Sakaguchi, S. T Cell Receptor Stimulation-Induced Epigenetic Changes and Foxp3 Expression

Are Independent and Complementary Events Required for Treg Cell Development. Immunity 37,

785-799, doi:10.1016/j.immuni.2012.09.010 (2012).

33   Wei, G., Wei, L., Zhu, J., Zang, C., Hu-Li, J., Yao, Z., Cui, K., Kanno, Y., Roh, T.-Y.,

Watford, W. T., Schones, D. E., Peng, W., Sun, H.-w., Paul, W. E., O'Shea, J. J. & Zhao, K.

Global Mapping of H3K4me3 and H3K27me3 Reveals Specificity and Plasticity in Lineage Fate

Determination   of   Differentiating   CD4(+)   T   Cells.   Immunity   30,   155-167,

doi:10.1016/j.immuni.2008.12.009 (2009).

34   Konopacki, C., Pritykin, Y., Rubtsov, Y., Leslie, C. S. & Rudensky, A. Y. Transcription factor

Foxp1 regulates Foxp3 chromatin binding and coordinates regulatory T cell function. Nature

Immunology 20, 232-+, doi:10.1038/s41590-018-0291-z (2019).

35   Oh, H., Grinberg-Bleyer, Y., Liao, W., Maloney, D., Wang, P., Wu, Z., Wang, J., Bhatt, D. M.,

Heise, N., Schmid, R. M., Hayden, M. S., Klein, U., Rabadan, R. & Ghosh, S. An NF-kappaB

Transcription-Factor-Dependent Lineage-Specific Transcriptional Program Promotes Regulatory

T Cell Identity and Function. Immunity 47, 450-465 e455, doi:10.1016/j.immuni.2017.08.010

(2017).

36   Rubtsov, Y. P., Rasmussen, J. P., Chi, E. Y., Fontenot, J., Castelli, L., Ye, X., Treuting, P.,

Siewe, L., Roers, A., Henderson, W. R., Jr., Muller, W. & Rudensky, A. Y. Regulatory T

cell-derived interleukin-10 limits inflammation at environmental interfaces. Immunity 28,

546-558, doi:10.1016/j.immuni.2008.02.017 (2008).

37   Fontenot, J. D., Gavin, M. A. & Rudensky, A. Y. Foxp3 programs the development and

function of CD4(+)CD25(+) regulatory T cells. Nature Immunology 4, 330-336,

doi:10.1038/ni904 (2003).

38   Kim, J. M., Rasmussen, J. P. & Rudensky, A. Y. Regulatory T cells prevent catastrophic

autoimmunity throughout the lifespan of mice. Nature Immunology 8, 191-197,

doi:10.1038/ni1428 (2007).

39   Shevyrev, D. & Tereshchenko, V. Treg Heterogeneity, Function, and Homeostasis. Frontiers

in Immunology 10, doi:10.3389/fimmu.2019.03100 (2020).

40   Zagorulya, M., Yim, L., Morgan, D. M., Edwards, A., Torres-Mejia, E., Momin, N.,

McCreery, C. V., Zamora, I. L., Horton, B. L., Fox, J. G., Wittrup, K. D., Love, J. C. & Spranger,

S. Tissue-specific abundance of interferon-gamma drives regulatory T cells to restrain

DC1-mediated priming of cytotoxic T cells against lung cancer. Immunity 56, 386-+,

doi:10.1016/j.immuni.2023.01.010 (2023).

41   Dikiy, S. & Rudensky, A. Y. Principles of regulatory T cell function. Immunity 56, 240-255,

doi:10.1016/j.immuni.2023.01.004 (2023).

42   Street, K., Risso, D., Fletcher, R. B., Das, D., Ngai, J., Yosef, N., Purdom, E. & Dudoit, S.

Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. Bmc Genomics

19, doi:10.1186/s12864-018-4772-0 (2018).

43   Tarbell, K. V., Yamazaki, S., Olson, K., Toy, P. & Steinman, R. M. CD25+ CD4+ T cells,

expanded with dendritic cells presenting a single autoantigenic peptide, suppress autoimmune

diabetes. Journal of Experimental Medicine 199, 1467-1477, doi:10.1084/jem.20040180 (2004).

44 Hori, S., Haury, M., Coutinho, A. & Demengeot, J. Specificity requirements for selection

and effector functions of CD25+4+ regulatory T cells in anti-myelin basic protein T cell receptor

transgenic mice. Proceedings of the National Academy of Sciences of the United States of

America 99, 8213-8218, doi:10.1073/pnas.122224799 (2002).

45 Butler, J. S., Lee, J.-H. & Skalnik, D. G. CFP1 Interacts with DNMT1 Independently of

Association with the Setd1 Histone H3K4 Methyltransferase Complexes. DNA and Cell Biology

27, 533-543, doi:10.1089/dna.2007.0714 (2008).

46 Butler, J. S., Palam, L. R., Tate, C. M., Sanford, J. R., Wek, R. C. & Skalnik, D. G. DNA

Methyltransferase Protein Synthesis Is Reduced in CXXC Finger Protein 1-Deficient Embryonic

Stem Cells. DNA and Cell Biology 28, 223-231, doi:10.1089/dna.2009.0854 (2009).

47 Benayoun, B. A., Pollina, E. A., Ucar, D., Mahmoudi, S., Karra, K., Wong, E. D., Devarajan,

K., Daugherty, A. C., Kundaje, A. B., Mancini, E., Hitz, B. C., Gupta, R., Rando, T. A., Baker, J.

C., Snyder, M. P., Cherry, J. M. & Brunet, A. H3K4me3 Breadth Is Linked to Cell Identity and

Transcriptional Consistency. Cell 158, 673-688, doi:10.1016/j.cell.2014.06.027 (2014).

48 Zacarias-Cabeza, J., Belhocine, M., Vanhille, L., Cauchy, P., Koch, F., Pekowska, A.,

Fenouil, R., Bergon, A., Gut, M., Gut, I., Eick, D., Imbert, J., Ferrier, P., Andrau, J.-C. &

Spicuglia, S. Transcription-Dependent Generation of a Specialized Chromatin Structure at the

TCRβ Locus. Journal of Immunology 194, 3432-3443, doi:10.4049/jimmunol.1400789 (2015).

49 Chen, K., Chen, Z., Wu, D., Zhang, L., Lin, X., Su, J., Rodriguez, B., Xi, Y., Xia, Z., Chen,

X., Shi, X., Wang, Q. & Li, W. Broad H3K4me3 is associated with increased transcription

elongation and enhancer activity at tumor-suppressor genes. Nature Genetics 47, 1149-+,

doi:10.1038/ng.3385 (2015).

50 Zhang, R., Huynh, A., Whitcher, G., Chang, J., Maltzman, J. S. & Turka, L. A. An obligate

cell-intrinsic function for CD28 in Tregs. Journal of Clinical Investigation 123, 580-593,

doi:10.1172/jci65013 (2013).

51 Gokhale, A. S., Gangaplara, A., Lopez-Occasio, M., Thornton, A. M. & Shevach, E. M.

Selective deletion of Eos (Ikzf4) in T-regulatory cells leads to loss of suppressive function and

development of systemic autoimmunity. Journal of Autoimmunity 105,

doi:10.1016/j.jaut.2019.06.011 (2019).

52 Wan, Y. Y. & Flavell, R. A. Regulatory T-cell functions are subverted and converted owing

to attenuated Foxp3 expression. Nature 445, 766-770, doi:10.1038/nature05479 (2007).

53 Chen, C., Rowell, E. A., Thomas, R. M., Hancock, W. W. & Wells, A. D. Transcriptional

regulation by Foxp3 is associated with direct promoter occupancy and modulation of histone

acetylation. Journal of Biological Chemistry 281, 36828-36834, doi:10.1074/jbc.M608848200

(2006).

54 Morikawa, H., Ohkura, N., Vandenbon, A., Itoh, M., Nagao-Sato, S., Kawaji, H., Lassmann,

T., Carninci, P., Hayashizaki, Y., Forrest, A. R. R., Standley, D. M., Date, H., Sakaguchi, S. &

Consortium, F. Differential roles of epigenetic changes and Foxp3 expression in regulatory T

cell-specific transcriptional regulation. Proceedings of the National Academy of Sciences of the

United States of America 111, 5289-5294, doi:10.1073/pnas.1312717110 (2014).

55 Pan, F., Yu, H., Dang, E. V., Barbi, J., Pan, X., Grosso, J. F., Jinasena, D., Sharma, S. M.,

McCadden, E. M., Getnet, D., Drake, C. G., Liu, J. O., Ostrowski, M. C. & Pardoll, D. M. Eos

Mediates Foxp3-Dependent Gene Silencing in CD4+ Regulatory T Cells. Science 325,

1142-1146, doi:10.1126/science.1176077 (2009).

56 DuPage, M., Chopra, G., Quiros, J., Rosenthal, W. L., Morar, M. M., Holohan, D., Zhang, R.,

Turka, L., Marson, A. & Bluestone, J. A. The Chromatin-Modifying Enzyme Ezh2 Is Critical for

the Maintenance of Regulatory T Cell Identity after Activation. Immunity 42, 227-238,

doi:10.1016/j.immuni.2015.01.007 (2015).

57 Liu, X., Wang, C., Liu, W., Li, J., Li, C., Kou, X., Chen, J., Zhao, Y., Gao, H., Wang, H.,

Zhang, Y., Gao, Y. & Gao, S. Distinct features of H3K4me3 and H3K27me3 chromatin domains

in pre-implantation embryos. Nature 537, 558-+, doi:10.1038/nature19362 (2016).

58 Sze, C. C., Ozark, P. A., Cao, K., Ugarenko, M., Das, S., Wang, L., Marshall, S. A.,

Rendleman, E. J., Ryan, C. A., Zha, D., Douillet, D., Chen, F. X. & Shilatifard, A. Coordinated

regulation of cellular identity-associated H3K4me3 breadth by the COMPASS family. Science

Advances 6, doi:10.1126/sciadv.aaz4764 (2020).

59 Dahl, J. A., Jung, I., Aanes, H., Greggains, G. D., Manaf, A., Lerdrup, M., Li, G., Kuan, S.,

Li, B., Lee, A. Y., Preissl, S., Jermstad, I., Haugen, M. H., Suganthan, R., Bjoras, M., Hansen, K.,

Dalen, K. T., Fedorcsak, P., Ren, B. & Klungland, A. Broad histone H3K4me3 domains in mouse

oocytes modulate maternal-to-zygotic transition. Nature 537, 548-+, doi:10.1038/nature19360

(2016).

60 Zhang, B., Zheng, H., Huang, B., Li, W., Xiang, Y., Peng, X., Ming, J., Wu, X., Zhang, Y.,

Xu, Q., Liu, W., Kou, X., Zhao, Y., He, W., Li, C., Chen, B., Li, Y., Wang, Q., Ma, J., Yin, Q.,

Kee, K., Meng, A., Gao, S., Xu, F., Na, J. & Xie, W. Allelic reprogramming of the histone

modification H3K4me3 in early mammalian development. Nature 537, 553-+,

doi:10.1038/nature19361 (2016).

61 Zhang, Z., Shi, L., Dawany, N., Kelsen, J., Petri, M. A. & Sullivan, K. E. H3K4

tri-methylation breadth at transcription start sites impacts the transcriptome of systemic lupus

erythematosus. Clinical Epigenetics 8, doi:10.1186/s13148-016-0179-4 (2016).

62 Wong, S. H. K., Goode, D. L., Iwasaki, M., Wei, M. C., Kuo, H.-P., Zhu, L., Schneidawind,

D., Duque-Afonso, J., Weng, Z. & Cleary, M. L. The H3K4-Methyl Epigenome Regulates

Leukemia Stem Cell Oncogenic Potential. Cancer Cell 28, 198-209,

doi:10.1016/j.ccell.2015.06.003 (2015).

63 Deaglio, S., Dwyer, K. M., Gao, W., Friedman, D., Usheva, A., Erat, A., Chen, J.-F., Enjyoji,

K., Linden, J., Oukka, M., Kuchroo, V. K., Strom, T. B. & Robson, S. C. Adenosine generation

catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune suppression.

Journal of Experimental Medicine 204, 1257-1265, doi:10.1084/jem.20062512 (2007).

Fig.1

A H3K4me3 enriched regions / FOXP3 binding sites
17516 | 8520 | 3036
Promoter 56%, Exon 6%, Intron 17%, Intergenic 22%

B H3K27me3 enriched regions / FOXP3 binding sites
40792 | 299 | 13153
Promoter 60%, Exon 5%, Intron 7%, Intergenic 28%

C H3K4me3 | H3K27me3 | FOXP3 — Cluster 1, Cluster 2, Cluster 3, Cluster 4; Enrichment High/Low; -5.0 TSS 5.0Kb

D RNA-seq — Cluster 1, Cluster 2, Cluster 3, Cluster 4; Expression Level 6.0 to -6.0

E Cluster 1, Cluster 2, Cluster 3, Cluster 4 gene ontology dot plots

Fig.2

Fig.3

Fig.4

A

2D2 Only
2D2+Treg(*Foxp3*^cre)
2D2+Treg(*Foxp3*^cre*Cxxc1*^fl/fl)

2D2 Only vs 2D2+Treg(Foxp3^cre), ** *p*=0.0013
2D2 Only vs 2D2+Treg(*Foxp3*^cre*Cxxc1*^fl/fl), *p*=ns
2D2+Treg(*Foxp3*^cre) vs 2D2+Treg(*Foxp3*^cre*Cxxc1*^fl/fl), ** *p*=0.0012

Fig.5

Fig.6

A

Foxp3^WT/YFP-Cre mosaics Treg cells
(Female mice)

X Chr random inactivation

Foxp3^WT
Foxp3^YFP-Cre

B

Gated on LN CD4^+cells

● YFP^-
● YFP^+

Foxp3^cre/+Cxxc1^fl/+   Foxp3^cre/+Cxxc1^fl/fl

FOXP3

YFP

7.83   2.75   9.59   1.02

YFP^+ in FOXP3^+ cells (%)

**

YFP^+ Cell number (x10^5)

**
2.5
2.0
1.5
1.0
0.5
0.0

C

Gated on CD4^+Foxp3^+cells

YFP^-   YFP^+

Ki-67

14.8   4.41

Ki-67^+ cells (%)

****

D

Isotype
--- YFP^-
— YFP^+

Gated on LN CD4^+Foxp3^+cells

Count

ICOS   CD25   CTLA4   GITR

● YFP^-
● YFP^+

ICOS MFI

***

CD25 MFI

***

CTLA4 (MFI)

***

GITR(MFI)

***
30000
20000
10000
0

Fig.7