CXXC-finger protein 1 associates with FOXP3 to stabilize homeostasis and suppressive functions of regulatory T cells

Xiaoyu Meng, [1,2,3†] Yezhang Zhu, [4,5 †] Kuai Liu, [1,2,3 †] Yuxi Wang,[6] Xiaoqian Liu, [1,2,3] Chenxin Liu,[2] Yan Zeng, [1,2,3] Shuai Wang, [1,2,3] Xianzhi Gao, [1,2,3] Xin Shen, [7] Jing Chen, [8] Sijue Tao, [6] Qianying Xu, [2] Linjia Dong, [9] Li Shen, [10,11,]* and Lie Wang[1,2,3,6,12,]**

[1]Institute of Immunology and Bone Marrow Transplantation Center, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.

[2]Zhejiang University School of Medicine, Hangzhou, China.

[3]Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, China.

[4]Department of Hematology, Tongji Hospital, School of Medicine, Tongji University, Shanghai, China

[5]Shanghai Key Laboratory of Signaling and Disease Research, Frontier Science Center of Stem Cell Research, National Stem Cell Translational Resource Center, School of Life Sciences and Technology, Tongji University, Shanghai, China

[6]Laboratory Animal Center, Zhejiang University, Hangzhou, China.

[7]Co-Facility Center, Zhejiang University School of Medicine, Hangzhou, China

[8]Department of Gastrointestinal Surgery, The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China.

[9]School of Basic Medical Sciences and Forensic Medicine, Hangzhou Medical College, Hangzhou, China

[10]MOE Key Laboratory of Biosystems Homeostasis & Protection and Zhejiang Provincial

Key Laboratory for Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang

University, Hangzhou, China.

[11]Department of Orthopedics Surgery, the Second Affiliated Hospital, School of Medicine,

Zhejiang University, Hangzhou, China

[12]Future Health Laboratory, Innovation Center of Yangtze River Delta, Zhejiang University,

Jiaxing, China.

[†] These authors contributed equally to this work.

* Correspondence: wanglie@zju.edu.cn; li_shen@zju.edu.cn.

Abstract

FOXP3-expressing regulatory T (T_{reg}) cells play a pivotal role in maintaining immune

homeostasis and tolerance, with their activation being crucial for preventing various

inflammatory responses. However, the mechanisms governing the epigenetic program in T_{reg}

cells during their dynamic activation remain unclear. In this study, we demonstrate that

CXXC finger protein 1 (CXXC1) interacts with the transcription factor FOXP3 and facilitates

the regulation of target genes by modulating H3K4me3 deposition. *Cxxc1* deletion in T_{reg}

cells leads to severe inflammatory disease and spontaneous T-cell activation, with impaired

immunosuppressive function. As a transcriptional regulator, CXXC1 promotes the expression

of key T_{reg} functional markers under steady-state conditions, which are essential for the

maintenance of T_{reg} cell homeostasis and their suppressive functions. Epigenetically, CXXC1

binds to the genomic regulatory regions of T_{reg} program genes in mouse T_{reg} cells,

overlapping with FOXP3 binding sites. Given its critical role in T_{reg} cell homeostasis,

CXXC1 presents itself as a promising therapeutic target for autoimmune diseases.

Introduction

Regulatory T (T_{reg}) cells are a distinct subset of $CD4^+$ T cells that play a critical role in

maintaining immune homeostasis and self-tolerance by suppressing excessive or aberrant

immune responses to foreign or self-antigens [1-3]. These cells can be further categorized into

thymus-derived regulatory T cells (tT_{reg} cells), periphery-derived T_{reg} cells (pT_{reg} cells), and

induced T_{reg} cells (iT_{reg} cells) [4]. They uniquely express the transcription factor FOXP3, a

member of the forkhead winged-helix family, which is essential for T_{reg} cell lineage

commitment and suppressive function [5,6]. Deletion or mutation of the *Foxp3* gene leads to a

range of immunological disorders, including allergies, immunopathology, and autoimmune

diseases in both mice and humans [7,8].

It is well established that FOXP3 recruits various cofactors to form complexes that either

promote or repress the expression of downstream genes, with histone and DNA modifications

playing pivotal roles in this process. FOXP3 can activate or repress the transcription of key

regulators of T_{reg} cell activation and function by recruiting the histone acetyltransferases

(HATs) or histone deacetylases (HDACs)[9,10]. Notably, FOXP3-bound sites exhibit enrichment

of H3K27me3, a modification essential for FOXP3-mediated repressive chromatin

remodeling under inflammatory conditions[11]. However, the direct role of FOXP3 as a

transcriptional activator through interactions with epigenetic regulators, particularly via

modulation of H3K4 trimethylation, remains poorly documented.

In mammals, six proteins have been identified that catalyze H3K4 methylation. These

proteins contain the SET domain and include MLL1 (KMT2A), MLL2 (KMT2B), MLL3

(KMT2C), MLL4 (KMT2D), SETD1A, and SETD1B [12,13]. For example, MLL1 plays a

critical role as an epigenetic regulator in T_{reg} cell activation and functional specialization[14].

Additionally, Placek *et al.* demonstrated that MLL4 is essential for T_{reg} cell development by

catalyzing H3K4me1 at distant unbound enhancers through chromatin looping [15]. H3K4me3,

which is enriched at the transcription start site (TSS) and the CpG island (CGI), converts

chromatin into active euchromatin by recruiting activating factors [16]. CXXC finger protein

1(CXXC1, also known as CFP1), which contains a SET1 interaction domain (SID), is

required for binding to the histone H3K4 methyltransferases SETD1A and SETD1B [17].

Previous studies have shown that CXXC1 regulates promoter patterns during T cell

maturation, GM-CSF-derived macrophage phagocytosis, TH17 cell differentiation, and the

function of ILC3 cells with aging by modulating H3K4me3 modification[18-21]. Despite the

well-documented role of CXXC1 in various immune effector cells, its role in T_{reg} cells

remains unclear.

Here, we demonstrate that CXXC1 interacts with FOXP3 and enhances the expression of

FOXP3 target genes. T_{reg} cell-specific deletion of *Cxxc1* triggers systemic autoimmunity,

accompanied by multiorgan inflammation, ultimately resulting in early-onset fatal

inflammatory disease in mice. *Cxxc1*-deficient T_{reg} cells exhibit a disadvantage in

proliferation and homeostasis, even in non-inflammatory mice where coexisting wild-type

(WT) T_{reg} cells were present. Moreover, *Cxxc1*-deficient T_{reg} cells display intrinsic defects in

the expression of key suppression molecules, including CTLA-4, CD25, ICOS, and GITR.

Consistent with these findings, mechanistic studies strongly suggest that CXXC1 functions as

an essential cofactor of FOXP3, playing a crucial role in sustaining H3K4me3 modifications

at key T_{reg} cell genes.

Materials and Methods

Mice

All mice used in this study were bred for a minimum of seven generations on a C57BL/6

background. Mouse experiments, or cells from mice of the same genotype, compared

littermates or age-matched control animals. The $Cxxc1^{fl/fl}$ mouse strain has been previously

described [22]. The $Foxp3^{cre}$ mice (JAX,016959) were generously provided by Bin Li

(Shanghai Jiao Tong University School of Medicine, Shanghai, China). CD45.1

(NM-KI-210226) mice were purchased from the Nanjing Biomedical Research Institute of

Nanjing University. $Rag1^{-/-}$ mice (stock#T004753) were purchased from GemPharmatech.

2D2 (MOG35-55-specific TCR transgenic) mice were graciously supplied by Prof. Linrong

Lu (Zhejiang University School of Medicine, Hangzhou, Zhejiang, China). $Foxp3^{cre}$ (WT)

and $Foxp3^{cre} Cxxc1^{fl/fl}$ (cKO) were utilized at 3 weeks old unless otherwise stated. All mice

were housed in the Zhejiang University Laboratory Animal Center under specific

pathogen-free conditions, and all animal experimental procedures were approved by the

Zhejiang University Animal Care and Use Committee (approval no.ZJU20230246).

Cell culture

HEK 293T cells were obtained from ATCC, and Plat E cells were kindly provided by Prof.

Xiaolong Liu (Shanghai Institutes for Biological Sciences). Both cell lines were cultured in

Dulbecco-modified Eagle medium (DMEM) containing 10% (v/v) fetal bovine serum (FBS),

supplemented with 1% penicillin/streptomycin.

Immunofluorescence (IF) Staining

As previously described [23], coverslips were treated with a 0.01% poly-L-lysine solution

(P4707; Sigma) for 10 minutes, air-dried, and then coated with $CD4^+$ YFP^+ T_{reg} cells. The

cells were then fixed in 4% formaldehyde for 15 minutes at room temperature, permeabilized

with 0.2% Triton X-100, and blocked with 1% BSA. Antibodies against CXXC1 (ab198977;

Abcam) and FOXP3(17-5773-82; Invitrogen) were diluted in Image iT FX signal enhancer

(I3693; Invitrogen) and incubated with cells overnight at 4 °C. After washing with

phosphate-buffered saline (PBS), the cells were incubated with a goat anti-rabbit antibody

Alexa Fluor 594 (1:250;10015289; Invitrogen) secondary antibody and stained with DAPI

(200 ng/mL; D523; Dojindo). Slides were washed with PBS and sealed with an antifade

solution(P36934; Invitrogen) before imaging with an Olympus FV3000 fluorescence

microscope. The images were visualized using the FV31-SW software.

Co-immunoprecipitation and western blot

Harvest the appropriate transfected cell lines and primary cells from culture and wash them

with ice-cold PBS. Lyse the cells in NETN300 buffer (300 mM NaCl, 0.5 mM EDTA,

0.5% (v/v) NP-40, 20 mM Tris-HCl pH 8.0) supplemented with a protease inhibitor (1:100,

P8340, Sigma-Aldrich) and PMSF (1 mM) on ice for 10 minutes. Take a small portion of the

whole-cell lysate as input, and incubate the remaining lysate with either Anti-FLAG M2

Beads (M8823; Sigma) or Anti-HA Beads (HY-K0201; MCE) on a rotator at 4°C overnight.

For the endogenous Co-IP assay targeting CXXC1 and FOXP3, incubate the cell lysate with

protein G magnetic beads along with anti-CXXC1 (ab198977; Abcam) or anti-FOXP3

(14-4774-82; Invitrogen) antibodies on a rotator at 4°C overnight. Wash the beads three times

with IP buffer (100 mM NaCl, 0.5 mM EDTA, 0.5% (v/v) NP-40, 20 mM Tris-HCl pH

8.0) to remove non-specific binding. Boil the washed beads with 1× Laemmli sample buffer

(1610747; Bio-Rad) to elute the bound proteins. Separate the denatured proteins by

SDS-PAGE. Transfer the separated proteins onto PVDF membranes (Millipore) for

immunoblotting.Immunoblot the PVDF membranes (IPVH00010) with the following

antibodies: anti-CXXC1 (1:1000; ab198977; Abcam), anti-FOXP3 (1:500; 14-7979-80;

Invitrogen), anti-FlAG (1:1000; 14793; Cell Signaling Technology), anti-HA (1:1000; 3724S;

Cell Signaling Technology).Detect the immunoblotted proteins using a secondary

HRP-conjugated goat anti-rabbit antibody (1:1000; HA1001-100; Huabio) and visualize the

bands using an appropriate detection method.

ELISA

Serum samples from 3-week-old WT and KO mice were analyzed for total IgG and IgE

concentrations using ELISA kits (88-50630-88; eBioscience) according to the manufacturer's

instructions. Half-area ELISA plates were coated with Coating Buffer and incubated

overnight at 4°C. After washing with PBST (PBS, 1 mM EDTA, 0.05% Tween-20), the plates

were blocked with 5% BSA in PBS for 30 minutes at room temperature. Serum was diluted to

the appropriate concentration with blocking buffer and incubated overnight at 4°C. After

washing with PBST, the plates were incubated with HRP-conjugated anti-mouse

IgG(1033-05; SouthernBiotech) and IgE(1110-05; SouthernBiotech) antibodies (1:2000 in

1% BSA/PBST) at 37°C for 1 hour. Following washing, TMB substrate was added, and the

reaction was stopped with 2 M H□SO□ after sufficient color development (1–15 minutes).

Absorbance at 450 nm was measured within 30 minutes.

Lymphocyte Isolation and Flow Cytometry

Cells from lymphoid organs were prepared by mechanical disruption between frosted slides,

while non-lymphoid organs were processed enzymatically. For lung tissue, minced samples

were digested in RPMI containing 100 μg/mL DNase I (9003-98-9;Sigma-Aldrich) and 2

mg/mL Collagenase D (LS004188;Worthington Biochemical) at 37°C for 1.5 hours. Liver

tissue was minced and digested in RPMI supplemented with 100 μg/mL DNase I and 1

mg/mL Collagenase D at 37°C for 30 minutes, with lymphocytes isolated using a 40-70%

Percoll (GE Healthcare) gradient. For intestinal tissue, the samples were first incubated in

DMEM containing 3% FBS, 0.2% HEPES, 0.5 M EDTA, and 0.145 mg/mL dithiothreitol for

10 minutes. This was followed by digestion with 50 mg/mL DNase I and 145 mg/mL

Collagenase II (Worthington Biochemical) in DMEM at 37°C for 5 minutes. Lymphocytes

were then isolated using an 80% and 40% Percoll gradient.

For surface marker analysis, cells were incubated for 15 minutes with purified anti-mouse

CD16/32 antibody (101320; BioLegend) to block Fc receptors. After blocking, cells were

stained with the indicated antibodies for surface markers. To determine cytokine expression,

cells were stimulated for 4 hours at 37□ with phorbol12-myristate13-acetate (50ng/mL;

S1819; Beyotime), ionomycin (1mg/ml; S1672; Beyotime), and brefeldin A

(BFA;00-4506-51; Invitrogen). After stimulation, cells were labeled with a fixable viability

dye and surface markers. Cells were then fixed and permeabilized according to the

manufacturer's instructions (00-8222-49; Invitrogen). For transcription factor staining,

samples were fixed using the FOXP3/Transcription Factor Staining Buffer Set (00-5523-00;

Invitrogen). Flow cytometry was conducted using a BD Fortessa (BD Biosciences) or

LongCyte (Beijing Challen Biotechnology Co., Ltd.) system. Flow cytometry data were

acquired and analyzed using FlowJo software.

The following antibodies were purchased from Invitrogen or BioLegend: Zombie Viole

t f-ixable viability(423113), Zombie NIR fixable(423105), CD25(PC61), CD8α(53-6.7),

CD62L(MEL-14), PD-1(J43), CD44(IM7), IL-17A(TC1118H10), KL-RG1(2F1), CD4

(GK1.5), TCRβ(H57-597), IFN-γ(XMG1.2), FOXP3(FJK-16s), CTLA-4(UC10-4B9), IC

OS(15F9), GITR(DTA-1), LAG3(C9B7W), TCRVβ11(RR3-15), CD45RB(C363-16A) C

D69 (H1.2F3), MHCII(M5/114.15.2), CD80(16-10A1), T-bet(eBio4B10), IKZF4(ESB7C

2), IL-4(11B11), CCR7(4B12), CD73(TY/11.8).

Real-time PCR

Total RNA was extracted from T$_{reg}$ cells using the RNAAiso Plus(9109; Takara) reagent

according to the manufacturer's instructions, and cDNA synthesis was performed using the

Prime Script RT Reagent Kit (Takara). TB Green Premix Ex Taq (RR420A; Takara) was used

for quantitative real-time PCR (qPCR). The expression levels of target mRNA were

normalized to the level of *β-actin* expression. The primers for qPCR are as follows:

Cxxc1 qPCR Forward: ATCCGGGAATGGTACTGTCG

Cxxc1 qPCR Reverse: CTGTGGAGAAGATTTGTGGG

β-actin qPCR Forward: CTGTCCCTGTATGCCTCTG

β-actin qPCR Reverse: ATGTCACGCACGATTTCC

CD4$^+$T and YFP$^+$ T$_{reg}$ cells adoptive transfer in Experimental autoimmune

encephalomyelitis (EAE)

CD4$^+$YFP$^+$ T$_{reg}$ cells from *Foxp3*[cre] and *Foxp3*[cre] *Cxxc1*[fl/fl] mice were enriched using the

Mouse CD4 T Cell Isolation Kit (480005; Biolegend) and then sorted using the BD Aria II

flow cytometer. Naïve CD4$^+$T cells from 2D2 (MOG35-55-specificTCR transgenic) mice

were isolated using the Mouse CD4 Naïve T cell Isolation Kit480039 (480039; Biolegend).

As previously described[24], 2D2 naïve CD4$^+$ T cells alone (5 x 10^5 per mouse), or 2D2 naïve

CD4$^+$ T cells (5x 10^5 per mouse) together with WT or *Foxp3*[cre]*Cxxc1*[fl/fl] T$_{reg}$ cells (2x10^5 per

mouse), were transferred into *Rag1*$^{-/-}$ mice via the tail vein. One day after cell transfer, the

recipient mice were inoculated subcutaneously (s.c.) with 200μg MOG35-55 peptide

(MEVGWYRSPFSRVVHLYRNGK; GenemeSynthesis) emulsified in complete Freund's

adjuvant (CFA) (F5506; Sigma). Intravenous administration of 200 ng of Pertussis toxin (181;

List Biological Laboratories) was performed on days 0 and 2 after peptide inoculation. The

severity of EAE was monitored and blindly graded using a clinical score from 0 to 5: 0, no

clinical signs; 1, limp tail; 2, paraparesis (weakness, incomplete paralysis of one or two hind

limbs); 3, paraplegia (complete paralysis of two hind limbs); 4, paraplegia with forelimb

weakness or paralysis; 5, dying or death.

Isolation lymphocytes from the central nervous system（CNS）

On day 14 after EAE induction, mice were perfused with transcardially administered PBS to

eliminate contaminating blood cells in the central nervous system (CNS). The forebrain and

cerebellum were dissected to expose the spinal cord, which was then carefully removed from

the spinal canal. The fresh spinal cord was harvested and cut into 2 mm pieces. The CNS

tissue pieces were homogenized using a syringe and passed through a 70 µM cell strainer to

obtain a single-cell suspension. The single-cell suspension was digested with collagenase D

(2µg/ml;11088858001; Roche) and deoxyribonuclease I (DNase I; 1 µg/ml; DN25;

Sigma-Aldrich) at 37°C for 20 minutes under rotation. After digestion, the cell suspension

was centrifuged to pellet the cells. The cell pellets were resuspended in 40% Percoll and

layered onto a discontinuous Percoll gradient. Centrifugation at 80% Percoll allowed for the

separation of cells at the 40–80% Percoll interface, which were collected as CNS

mononuclear cells. The collected CNS mononuclear cells were washed with PBS to remove

any remaining Percoll. CNS mononuclear cells were stimulated for 4 hours with PMA and

ionomycin in the presence of Brefeldin A to induce cytokine production. After stimulation,

cells were fixed, rendered permeable, and stained with appropriate antibodies for intracellular

cytokine detection.

Histological analyses

The lungs, skin, liver, and colon were excised from three-week-old mice. Prior to histological

analysis, the samples were fixed in formalin, embedded in paraffin, and stained with

hematoxylin and eosin. For CNS histology, spinal cords were fixed in 4% paraformaldehyde,

paraffin-embedded, sectioned, and stained with Luxol Fast Blue and hematoxylin and eosin

(H&E). To examine colon histology, colons from $Rag1^{-/-}$ hosts were similarly processed and

stained with H&E.

Adoptive transfer colitis model

Colitis was induced following the protocol described[25]. In brief, CD4$^+$ YFP$^+$ T$_{reg}$ cells were

isolated from 3-week-old CD45.2$^+$$Foxp3^{cre}Cxxc1^{fl/fl}$ and CD45.2$^+$ $Foxp3^{cre}$ mice. A total of

2x10^5 T$_{reg}$ cells from each group were mixed with 4x10^5 T$_{eff}$ cells (CD45.1$^+$CD4$^+$CD45RB^{hi})

sorted from CD45.1$^+$ mice and transferred into the $Rag1^{-/-}$ mice via intraperitoneal injection.

T$_{eff}$ cells alone were transferred as a control group. Mouse body weight was measured weekly

post-adoptive transfer. The percentage change in body weight was calculated by comparing

the current weight with the initial weight on day 0. Mice were euthanized when any had

reached 80% of their initial body weight. The large intestines were sectioned into 4μm thick

slices and stained with hematoxylin.

In vitro T$_{reg}$ suppression assay

Naïve CD4$^+$ T cells isolated from WT mice were labeled with CFSE (C34554; Invitrogen).

CD4$^+$YFP$^+$ T$_{reg}$ cells from $Foxp3^{cre}$ and $Foxp3^{cre}Cxxc1^{fl/fl}$ mice were cultured with naïve

CD4$^+$ T cells (1□10^5 cells) at various ratios in the presence of 2μg/mL anti-CD3(16-0031-85;

Invitrogen) and 3μg/mL anti-CD28(16-0281-85; Invitrogen). On day 3, cells were analyzed

by flow cytometry.

CUT&Tag

CUT&Tag assays of CD4$^+$YFP$^+$ T$_{reg}$ cells were conducted as previously described[26]. Briefly,

approximately 1x10^5 single cells were carefully pipetted into wash buffer twice. The pelleted

cells were resuspended in wash buffer, activated concanavalin (BP531; Bangs Laboratories),

and incubated for 15 minutes at room temperature. Cells bound to the beads were

resuspended in Dig-Wash Buffer and incubated with a 1:50 dilution of primary antibodies

(rabbit anti-H3K4me3, Active Motif,39016; rabbit anti-CXXC1, abcam, ab198977; rabbit

anti-FOXP3, abcam, ab 150743;normal IgG, Cell Signaling, 2729) at 4 □ overnight. The

beads were incubated with a secondary antibody (goat anti-rabbit IgG; SAB3700883;

Sigma-Aldrish) diluted 1:100 in Dig-Wash buffer for 60 minutes at room temperature. Cells

were treated with Hyperactive pG-Tn5 Transposase (S602; Vazyme) diluted in Dig-300

Buffer for 1 hour at room temperature. The cells were subsequently resuspended in

Tagmentation buffer (10mM MgCl2 in Dig-300 Buffer) and incubated at 37□ for 1 hour. To

halt tagmentation, 10μl was spiked with 0.5M EDTA, 3μl with 10%SDS, and 3μl with

20mg/ml Proteinase K and incubated at 55□ for 1h. DNA library amplification was

performed according to the manufacturer's instructions and purified using VAHTS DNA

Clean Beads (N411; Vazyme). Libraries were sequenced on the Illumina NovaSeq platform

(Annoroad Gene Technology).

CUT&Tag and ChIP-seq data analysis

FOXP3 ChIP-seq data was obtained from GSE121279. H3K27me3 ChIP-seq data was

obtained from GSE14254. CUT&Tag and ChIP-seq reads were trimmed to 50 bp and aligned

against the mouse genome build mm9 using Bowtie2(v2.3.4.1) with default parameters. All

PCR duplicates and unmapped reads were removed. Peak calling was performed using

MACS2(v2.1.1.20160309) and signal tracks for each sample were generated using the

'wigToBigWig' utility of UCSC. We classified the H3k4me3 peaks around TSSs into three

groups: broad (>5 kb), medium (1-5kb), and narrow (<1 kb). The top 5% of the widest peaks

were considered as broad peaks. The average intensity profiles were generated using

deepTools (v2.5.4). Motif analysis was performed using the 'findmotifsGenome.pl' command

inHomer2 package. Epigenetic factors were identified using the Epigenetic Factor

Database(https://epifactors.autosome.org/) and then screened for those that exclusively

regulate the expression of their target genes by modulating the deposition of H3K4me3.

Genomic distribution was analyzed using the "genomation" R package. GO pathway analysis

was performed using the "clusterProfiler" R package. The sequencing information of

CUT&Tag data used in this study is summarized in Supplementary Table S1.

Clustering analysis

Promoters were defined as ±2 kb regions flanking the annotated TSS. Reads in promoters

were counted using the "coverage" command in bedtools (v2.26.0) and further normalized to

RPKM. The k-means clustering of H3K4me3 and H3K27me3 enrichment at promoters was

conducted using the "kmeans" function in R.

RNA-seq and data analysis

Total RNA was extracted from sorted CD4☐YFP☐ T_{reg} cells using the RNeasy Plus Mini Kit

(Qiagen, #74134), following the manufacturer's protocol. RNA-Seq libraries were

constructed and sequenced by Haplox (Nanchang, China), using an Illumina platform with

paired-end reads of 150 bp. RNA-seq data was obtained from GSE82076. Raw reads were

trimmed to 50 bp and mapped to the mouse genome (mm9) using TopHat (v2.1.1) with

default parameters. Only uniquely mapped reads were kept for downstream analysis. The

RNA abundance of each gene was quantified using Cufflinks (v2.2.1).

Single-Cell RNA-Sequencing

A total of 300,000 sort-purified CD4$^+$YFP$^+$ T_{reg} cells from *Foxp3*^{cre} and *Foxp3*^{cre} *Cxxc1*$^{fl/fl}$

were resuspended in BD Pharmingen Stain Buffer (FBS) (554656; BD). Single cells were

isolated using a chromium controller (BD platform, BD Bioscience) according to the

manufacturer's instructions, as previously described[27]. The single cells were labeled with

sample tags using the BD Mouse Immune Single-Cell Multiplexing Kit (633793; BD).

Following standard protocols, cDNA amplification and library construction were performed

to generate scRNA-seq libraries.

Targeted scRNA-seq data processing

The raw FASTQ files were processed by BD Rhapsody using the Targeted analysis pipeline.

After alignment and filtering, the distribution-based error correction(DBEC)-adjusted

molecules were loaded into R studio(version 4.3.2). All subsequent analyses were performed

using the package Seurat (version 4.4.0) with default parameters. Specifically, the scRNA-seq

data counts were log-normalized. All targeted genes were scaled and then were used for

Principal components analysis (PCA). The batch effects were removed by the

HarmonyMatrix function in the Harmony package (version 1.1.0). The first 20 principal

components were used to calculate nonlinear dimensionality reduction using RunUMAP.

Differential gene expression (DEGs) between clusters was assessed using the FindAllMarkers

function. The clusters were then annotated based on DEGs. Barplots were generated using

ggplot2 (version 3.4.4). Heatmaps were generated using pheatmap (version 1.0.12).

Analysis of the single-cell TCR-seq repertoire

Raw V(D)J fastq reads were processed using BD Rhapsody Pipeline and then were analyzed

using scRepertoire (version 1.12.0). The TCR clonotype was called using the nucleotide

sequence of the CDR3 region for both TCR alpha and beta chains. For cells with multiple

chains, the top two clonotypes with the highest expression were selected for downstream

analysis. Clonal overlap between different cell types was calculated using the clonalOverlap

function of scRepertoire. A clonotype was defined as expansion if it could be detected in at

least two cells.

scRNA-seq trajectory analysis

UMAP embeddings obtained from the Seurat package were projected into the

Slingshot(version 2.10.0) package to construct pseudotime Trajectories for T_{reg} cells. Naïve

subsets were set as the root state.

Whole genome bisulfite sequencing (WGBS) and data analysis

Sorted CD4$^+$YFP$^+$ T$_{reg}$ cells(3$\square$10^6) were lysed in cell lysis buffer to release DNA. The

bisulfite-treated DNA was used to prepare the sequencing library. DNA libraries were

transferred to the Illumina Platform for sequencing using 150 bp paired-end reads. Raw reads

were trimmed using TrimGalore (v0.4.4) with default parameters. Subsequently, the reads

were mapped against the mm9 reference genome using Bismark v0.19.0 with parameters

'--bowtie2'. PCR duplicates were removed and the methylation levels were calculated using

'bismark_methylation_extractor'. We calculated the mean CpG methylation levels of various

genome elements: promoter, 5$'$-UTR, exon, intron, 3$'$-UTR, genebody, intergenic, CGIs, and

repeats using in-house scripts. The sequencing information of WGBS data used in this study

is summarized in Supplementary Table S2.

Statistical analysis

The statistical significance analysis was performed using Prism 8.0 (GraphPad). Error bars

are presented as mean ± SD. *P* values of < 0.05 were deemed statistically significant (**P <*

0.05, **P < 0.01, ***P < 0.001 and *****P < 0.000*). Statistical analyses were performed with

unpaired Student's *t*-test or two-way ANOVA and Holm–Sidak post hoc test.

Results

FOXP3 binds regulatory loci primed for activation and repression in T$_{reg}$ cells

FOXP3-mediated gene expression is well-recognized, with several studies highlighting its

dual role as both a transcriptional activator and repressor [28-32]. However, the mechanisms by

which FOXP3 regulates T_{reg}-specific gene transcription via epigenetic modifications remain

incompletely understood. To investigate these mechanisms, we employed CUT&Tag to

generate genome-wide H3K4me3 maps in T_{reg} cells. To complement this, we compared our

data with an H3K27me3 ChIP-Seq dataset from Wei *et al.*[33] focusing on previously identified

FOXP3-bound loci(Konopacki *et al.* 2019)[34]. This integrative analysis allowed us to identify

FOXP3-dependent genes associated with either H3K4me3 (indicative of transcriptional

activation) or H3K27me3 (indicative of repression) deposition. These findings provide

insight into how FOXP3 modulates the epigenetic landscape to regulate T_{reg} cell function.

As expected, H3K4me3 was enriched at gene promoters (fig. S1, A and B). A Venn diagram

revealed overlap between FOXP3 binding sites and H3K4me3 peaks, with minimal overlap

with H3K27me3 peaks (Fig. 1, A and B). The overlapping regions between FOXP3 binding

sites and H3K4me3 or H3K27me3 peaks were predominantly located at promoters (Fig. 1, A

and B). To further investigate the role of FOXP3 in promoting H3K4me3 deposition, we

compared H3K4me3 levels between FOXP3-positive T_{reg} cells and FOXP3-negative

conventional T cells (Tconv). Consistent with our hypothesis, H3K4me3 abundance was

higher at T_{reg}-specific gene loci (e.g., *Tnfrsf18*, *Ctla4*, *Il2ra* and *Nt5e*) in T_{reg} cells compared

to Tconv cells (Fig. 1,C and fig. S1, C). Notably, the selection of these loci was guided by

prior studies identifying genes specifically associated with T_{reg} cell function[35]. This finding

highlights the role of FOXP3 in shaping the epigenetic landscape of T_{reg} cells through

enhanced H3K4me3 deposition at these critical loci.

To further characterize these modifications,we clustered promoters into four groups based on

the enrichment of H3K4me3 and H3K27me3. Clusters 1 and 3 showed strong enrichment of

H3K4me3; cluster 2 was enriched with H3K27me3; and cluster 4 showed weak enrichment

of both modifications. FOXP3 preferentially bound to clusters 1 and 3 prom-oters, which

displayed high H3K4me3 levels (Fig. 1C). Correspondingly, genes in these clusters exhibited

high transcription levels, as shown by reanalysis of previously published RNA-seq data (Oh

et al. 2017) [36] （Figure 1D, right）. In contrast, genes with H3K27me3 enrichment at their

promoters were transcribed at low levels.

Gene Ontology (GO) analysis of these four clusters revealed distinct functional roles. Cluster

1 was enriched in genes involved in mRNA processing, covalent chromatin modification, and

histone modification, while cluster 3 was enriched in genes related to DNA repair and

mitochondrion organization (Fig. 1E). Cluster 2, enriched with H3K27me3, was associated

with pattern specification process, while cluster 4 showed no correlation with T_{reg} cells.

Notably, signature T_{reg} cell genes such as *Tnfrsf18*, *Nrp1*, *Stat5a*, *Lag3*, *Icos*, and *Pdcd1* were

enriched in clusters 1 and 3, showing strong H3K4me3 marks (fig. S1C). Conversely, genes

like *Hic1*, *Trp73,* and *Rnf157*, associated with inflammatory responses, were enriched for

H3K27me3 in clusters 2 (fig. S1C). These findings collectively support the conclusion that

FOXP3 contributes to transcriptional activation in T_{reg} cells by promoting H3K4me3

deposition at target loci, while also regulating gene expression directly or indirectly through

other epigenetic modifications.

CXXC1 interacts with FOXP3 and binds H3K4me3-enriched sites in T_{reg} cells

We conducted an enrichment analysis of known motifs at the overlapping peaks of FOXP3

ChIP-seq and H3K4me3 CUT&Tag in T_{reg} cells to identify epigenetic factors that directly

interact with FOXP3 to mediate chromatin remodeling and transcriptional reprogramming.

Motif analysis of the overlapping peaks between FOXP3 binding sites and regions enriched

in H3K4me3 revealed that, in addition to transcription factors, the most abundant motif

associated with H3K4me3 was the epigenetic factor CXXC1(fig. S2A). To investigate this

further, we performed CUT&Tag for endogenous CXXC1 in T_{reg} cells to examine the

genome-wide co-occupancy of CXXC1 and FOXP3. Over half of these CXXC1 binding sites

were located at promoter regions (fig. S2B). Additionally, CXXC1 exhibited strong binding

at transcription start site (TSS) and CpG islands (CGIs)(Fig. 2A and fig. S2C). As illustrated

by the Venn diagram (Fig. 2B), more than half of the FOXP3-bound genes and

H3K4me3-enriched genes were also bound by CXXC1. Similarly, more than half of

CXXC1 peaks were overlapped with FOXP3 peaks (fig. S2D). Furthermore, the

CXXC1-specific and FOXP3-specific binding sites also demonstrated modest binding of

FOXP3 and CXXC1, respectively (Fig. 2C). These findings indicate that FOXP3 and

CXXC1 share a substantial number of target genes in T_{reg} cells. To confirm this interaction,

we further validated the reciprocal immunoprecipitation of both endogenous and exogenous

CXXC1 and FOXP3 (Fig. 2D and fig. S2E). An immunofluorescence assay revealed

predominant colocalization of CXXC1 with FOXP3 in the nucleus (Fig. 2E). Overall, these

results suggest that CXXC1 primarily functions as a coactivator of FOXP3-driven

transcription in T_{reg} cells.

Complete ablation of *Cxxc1* in T$_{reg}$ cells leads to a fatal autoimmune disease

To investigate the role of CXXC1 in T$_{reg}$-cell homeostasis and function, we generated *Foxp3*^{Cre}*Cxxc1*$^{fl/fl}$ mice (conditional knockout [cKO] mice) by crossing *Cxxc1*$^{fl/fl}$ with *Foxp3*$^{YFP-Cre}$ [37] mice, thereby specifically deleting *Cxxc1* in T$_{reg}$ cells. The effective depletion of *Cxxc1* in T$_{reg}$ cells was confirmed through quantitative PCR (qPCR) and Western blotting (fig. S3A). Notably, cKO mice appeared normal at birth but later exhibited spontaneous mortality starting around three weeks of age (Fig. 3A). Deletion of *Cxxc1* in T$_{reg}$ cells led to the development of severe inflammatory disease, characterized by reduced body size, stooped posture, crusting of the eyelids, ears, and tail, and skin ulceration, particularly on the head and upper back (Fig. 3, B and C). Additionally, cKO mice developed extensive splenomegaly and lymphadenopathy (Fig. 3D). *Foxp3*^{Cre}*Cxxc1*$^{fl/fl}$ mice exhibited elevated serum levels of anti-dsDNA autoantibodies and IgG , along with a modest increase in IgE concentration (Figure 3E and fig. S3B). Histopathological analysis revealed massive lymphocyte and myeloid cell infiltration in the skin, lungs, liver sinusoids, and colon mucosa (Fig. 3F). In full agreement with the aforementioned severe autoimmune diseases, *Foxp3*^{Cre}*Cxxc1*$^{fl/fl}$ mice had decreased percentages and numbers of CD4$^+$ Foxp3$^+$ T$_{reg}$ cells in small intestine lamina propria (LPL), liver and lung(Fig. 3G and fig. S3C). Moreover, cKO mice displayed an increase in CD8$^+$ T cell percentages (fig. S3D), along with a marked rise in cells exhibiting an effector/memory phenotype (CD44hi CD62Llo) (Fig. 3H). Furthermore, T cells from cKO mice produced elevated levels of IFN-γ, IL-17, and IL-4 in CD4$^+$ YFP$^-$ T cells, as well as increased IFN-γ production in CD8$^+$ T cells (Fig. 3I and fig. S3, E and F). These phenotypes closely resembled those observed in *Foxp3*-deficient mice [38] or mice with depleted T$_{reg}$ cells[39],

suggesting a deficiency in immune suppression.

CXXC1 is necessary for the maintenance of T$_{reg}$-cell suppressive activity

Despite the development of severe autoimmune disease, we observed an increase in both the

absolute number and percentage of FOXP3$^+$ T$_{reg}$ cells in the lymph nodes (fig. S4A). The

expression level of the FOXP3 protein was only slightly altered in *Cxxc1*-deficient T$_{reg}$ cells

(fig. S4B). In an *in vitro* suppression assay, T$_{reg}$ cells from *Foxp3*^{Cre}*Cxxc1*$^{fl/fl}$ and WT mice

exhibited similar suppressive effects on naïve T (Tn) cell proliferation (fig. S4C). The

expression of the hallmark T$_{reg}$-cell marker CTLA-4 showed a modest increase in

Cxxc1-deficient T$_{reg}$ cells compared to WT T$_{reg}$ cells, while the expression of GITR remained

unchanged (fig. S4D). To further assess the suppressive capacity of *Cxxc1*-deficient T$_{reg}$ cells

in vivo, we employed the experimental autoimmune encephalomyelitis (EAE) model. Naïve

CD4$^+$ T cells from 2D2 mice were co-transferred with T$_{reg}$ cells from either *Foxp3*^{Cre} or

Foxp3^{Cre} *Cxxc1*$^{fl/fl}$ mice into *Rag1*$^{-/-}$ recipients, and EAE was induced in these recipient mice.

Mice that received only naïve CD4$^+$ T cells from 2D2 mice developed more severe EAE

symptoms (Fig. 4A). The addition of WT T$_{reg}$ cells from *Foxp3*^{Cre} mice slightly mitigated

EAE progression and reduced Th17 cells in the spinal cord (Fig. 4, A to D). In contrast,

Foxp3^{Cre} *Cxxc1*$^{fl/fl}$ T$_{reg}$ cells failed to suppress EAE (Fig. 4, A to D), and the cKO mice

showed a reduction in T$_{reg}$ cells frequency in CNS tissues(Fig. 4E). Finally, we examined the

role of CXXC1 in T$_{reg}$ cell-mediated suppression using T cell transfer-induced colitis, in

which naïve T cells were transferred to *Rag1*$^{-/-}$ recipients either alone or together with WT or

Foxp3^{Cre} *Cxxc1*$^{fl/fl}$ T$_{reg}$ cells. The transfer of naïve T cells led to weight loss and intestinal

pathology in recipient mice(Fig. 4, F and G). Mice receiving WT T$_{reg}$ cells continued to gain

weight (Fig. 4F) , whereas those that received T$_{reg}$ cells from cKO mice were unable to

prevent colitis and exhibited a reduced percentage of T$_{reg}$ cells (Fig. 4, F to H). These findings

underscore the critical role of CXXC1 in maintaining T$_{reg}$ cell function *in vivo*.

T$_{reg}$ cell lineage homeostasis and proliferation depend upon CXXC1

T$_{reg}$ cells harbor a diverse T-cell receptor (TCR) repertoire, which likely play a critical role in

their immune suppression function [40-42]. To explore the role of CXXC1 in T$_{reg}$-mediated

suppression, we performed single-cell RNA sequencing (scRNA-seq) combined with TCR

sequencing (TCR-seq) on CD4$^+$YFP$^+$ T$_{reg}$ cells isolated from mouse lymph nodes. After

quality control and removal of doublets, 18,577 cells were retained for further analysis.

Through unsupervised clustering and uniform manifold approximation and projection

(UMAP) analysis, we identified eight distinct T$_{reg}$ cell clusters based on the expression of

well-characterized markers, with a particular focus on two clusters of activated T$_{reg}$ cells that

exhibited higher expression of markers and gene sets relative to naïve T$_{reg}$ cells (Fig. 5A and

fig. S5, A to C). A comparison between *Cxxc1*-deficient and WT T$_{reg}$ cells within each cluster

revealed a reduction in *Cxxc1*-deficient cells in the naïve subsets, while an increase was

observed in the Gzmb$^+$ and H2-Eb1$^+$ subsets (Fig. 5B). To further elucidate the transition of

T$_{reg}$ cells along a dynamic biological timeline, we constructed pseudo-time trajectories using

Slingshot [43]. The pseudo-time gradient depicted a progression from quiescent to activated T$_{reg}$

cells, ultimately encompassing the Gzmb$^+$ and H2-Eb1$^+$ subsets (Fig. 5C). Given the

antigen-specific suppression capabilities of T$_{reg}$ cells [44,45], we examined their clonal

expansion. The analysis revealed that expanded WT TCR clonotypes (n$\geq$2) were

predominantly distributed among the Nt5e$^+$ subsets, while *Cxxc1*-deficient T$_{reg}$ cells showed

expanded clonotypes primarily within the Gzmb$^+$ and H2-Eb1$^+$ subsets(Fig. 5D and Fig.

S5D). TCR sharing analysis indicated clonotype sharing among various clusters of WT T$_{reg}$

cells, suggesting a degree of homogeneity. However, the reduced TCR sharing in

Cxxc1-deficient T$_{reg}$ cells implies that decreased TCR diversity may impair the suppressive

activity of T$_{reg}$ cells (Fig. 5E) [42]. Furthermore, the *Cxxc1*-deficient group exhibited lower

expression of several T$_{reg}$-specific genes associated with suppressive functions, such as*Nt5e*,

Il10, *Pdcd1*, *Klrg1*, as well as genes that inhibit effector T cells differentiation, including *Sell*

and *Tcf7* (Fig. 5F and fig. S5E). Conversely, *Cxxc1*-deficient T$_{reg}$ cells demonstrated elevated

expression of *Gzmb*, *Il2ra*, and *Cd69* compared to WT, reflecting a profile indicative of

increased activation (Fig. 5, F and G). Additionally, we observed increased expression of

genes linked to Th1-type inflammation, such as *Ifng*, *Tbx21,* and *Hif1a*, in *Cxxc1*-deficient

T$_{reg}$ cells, likely due to extreme inflammatory conditions (Fig. 5, F to H). The proportion of

FOXP3$^+$Ki67$^+$ T$_{reg}$ cells was lower in cKO mice compared to WT mice (Fig. 5I). These

findings underscore the crucial role of CXXC1 in in maintaining T$_{reg}$ cell function and

homeostasis.

Intrinsic *Cxxc1* deficiency impairs T$_{reg}$ cell suppression, proliferation, and molecular

programs.

To confirm that the deficiency in T$_{reg}$ cell function in T$_{reg}$-specific *Cxxc1*-deficient animals is

due to intrinsic defects caused by *Cxxc1* deficiency, rather than severe autoimmune

inflammation in *Foxp3*^{Cre} *Cxxc1*^{fl/fl} mice, we examined *Cxxc1*-sufficient and *Cxxc1*-deficient

T_{reg} cell subsets in heterozygous *Foxp3*^{Cre/+} *Cxxc1*^{fl/fl} (designated as "het-KO") and littermate

Foxp3^{Cre/+} *Cxxc1*^{fl/+} (designated as "het-WT") female mice (Fig. 6A). Notably, het-KO female

mice did not exhibit overt signs of autoimmunity, as random X-chromosome inactivation led

to the coexistence of both *Cxxc1*-cKO and *Cxxc1*-WT T_{reg} cells. However, both the frequency

and absolute numbers of FOXP3$^+$YFP$^+$ T_{reg} cells within the total T_{reg} population were reduced

in het-KO mice compared to their counterparts in het-WT littermates, indicating that *Cxxc1*

deficiency imposes a competitive disadvantage on T_{reg} cells (Fig. 6B). Additionally,

Cxxc1-deficient YFP$^+$ T_{reg} cells failed to upregulate the proliferation marker Ki-67 (Fig. 6C).

Moreover, YFP$^+$ T_{reg} cells in het-KO female mice showed reduced expression of key genes

essential for suppressive function, including ICOS, CD25,CTLA4, and GITR, compared to

YFP$^-$$T_{reg}$ cells from the same mice (Fig. 6D). Consistently, we confirmed the impaired

suppressive function of T_{reg} cells from heterozygous *Foxp3*^{Cre/+} *Cxxc1*^{fl/fl} mice in *vitro* and in

vivo (Fig. 6E and fig. S6A-E). To investigate the molecular program affected by the deletion

of *Cxxc1* in T_{reg} cells, we performed RNA-sequencing (RNA-seq) analysis on CD4$^+$YFP$^+$ T_{reg}

cells isolated from het-WT and het-KO mice. We then conducted a differential gene

expression (DGE) analysis based on the RNA-seq data. Among all expressed genes, 865 were

upregulated ("Up" genes) and 761 were downregulated ("Down" genes) by ≥1.5-fold in

CD4$^+$YFP$^+$ T_{reg} cells from het-KO mice compared to het-WT mice, with a false discovery

rate (FDR)–adjusted *P*-value cutoff of 0.05 (fig. S6F). Gene Ontology (GO) enrichment

analysis revealed that the downregulated genes in *Cxxc1*-deficient T_{reg} cells were

predominantly enriched in pathways related to the negative regulation of immune system

process and regulation of cell–cell adhesion (fig. S6G). The *Cxxc1*-deficient T$_{reg}$ cells also

showed reduced expression of several genes associated with T$_{reg}$ cell suppressive function,

including *Il10*, *Tigit*, *Lag3*, *Icos*, *Nt5e*(encoding CD73) and *Itgae* (encoding CD103) (fig.

S6H). Thus, while YFP☐ WT T$_{reg}$ cells effectively prevent autoimmunity in het-KO mice, the

absence of *Cxxc1* in YFP☐ T$_{reg}$ cells disrupts key T$_{reg}$ cell marker expression and impairs

their suppressive function under steady-state conditions.

The FOXP3-CXXC1 complex regulates the expression of key factors in T$_{reg}$ cells that

are associated with the breadth of H3K4me3

CXXC1 binds to unmethylated CpG DNA via its N-terminal CXXC finger domain,

facilitating its interaction with DNA methyltransferase 1 (DNMT1). This binding stabilizes

the DNMT1 protein, thereby regulating DNA methylation [46,47]. To investigate whether

CXXC1 depletion affects DNA methylation in T$_{reg}$ cells, we performed whole genome

bisulfite sequencing (WGBS) on T$_{reg}$ cells isolated from both WT and cKO mice. On average,

Cxxc1-deficient T$_{reg}$ cells exhibited no changes in DNA methylation at gene loci or across

genome-wide CpG sites, irrespective of chromosomal region (fig. S7, A to C). Furthermore,

Cxxc1 knockout T$_{reg}$ cells did not show a increase in DNA methylation at key T$_{reg}$ signature

gene loci (fig. S7D). Given the pivotal role of MLL4-mediated H3K4me1 in establishing the

enhancer landscape and facilitating long-range chromatin interactions during T$_{reg}$ cell

development[15], we performed CUT&Tag to assess changes in H3K4me1 levels in

Cxxc1-deficient T$_{reg}$ cells. This analysis revealed that H3K4me1 levels were similar in both

WT and *Cxxc1*-deficient T$_{reg}$ cells (fig. S7, E to G).

While H3K4me3 modifications typically form sharp 1-to 2-kb peaks around promoters, some

genes exhibit broader H3K4me3 regions, referred to as broad H3K4me3 domains

(H3K4me3-BDs), which can extend to cover part or all of the gene's coding sequences (up to

20 kb) [48,49]. Broad H3K4me3 domains are preferentially associated with genes essential for

the identity or function of specific cell types [48,50] and have been implicated in enhancing

transcriptional elongation and increasing enhancer activity [50]. To further explore the

connection between broad H3K4me3 domains and the expression of immune-regulatory

genes, we analyzed genes containing broad H3K4me3 regions. We classified the H3K4me3

domains surrounding transcription start sites (TSSs) into three categories: broad (more than 5

kb), medium (between 1 and 5 kb), and narrow (less than 1 kb) (Fig. 7A). Notably,

$Cxxc1$-deficient T_{reg} cells exhibited weaker H3K4me3 signals compared to WT cells within

the broad H3K4me3 domains where CXXC1 binding is prominent (Fig. 7, A and B). Using

the criteria established by Benayoun et $al.$[48], which defines the top 5% of the widest

H3K4me3 domains as BDs, we observed similar enrichment results (fig. S7, H and I). We

then compared three groups of genes: BD-associated genes with reduced H3K4me3 levels

following $Cxxc1$ deletion, genes with direct CXXC1 binding, and genes with direct FOXP3

binding. The Venn diagram revealed that the majority of genes (283 out of 294, 96%) with

CXXC1 binding and reduced H3K4me3 levels overlap with FOXP3-bound genes, suggesting

that CXXC1 presence enhances FOXP3 binding within the broad H3K4me3 domain (Fig.

7C). Furthermore, GO term analysis indicated that BD-associated genes are enriched in

biological processes related to the negative regulation of immune system processes (Fig. 7D).

Genome browser views displayed the enrichments of FOXP3, CXXC1, and H3K4me3 at key

signature genes in T_{reg} cells, such as *Ctla4, Il2ra, Icos*, and *Tnfrsf18*, with lower H3K4me3

densities observed at these loci in *Cxxc1*-deficient T_{reg} cells (Fig. 7E). Similar patterns were

observed at core genes involved in T_{reg} homeostasis and suppressive function (e.g., *Lag3*,

Nt5e, *Ikzf4*, and *Cd28*) (fig. S7, J)[51,52]. These findings suggest that CXXC1 and FOXP3

collaboratively promote sustained T_{reg} cell homeostasis and function by preserving the

H3K4me3 modification at key T_{reg} cell genes.

We previously demonstrated that the FOXP3-CXXC1 complex plays a key role in

modulating H3K4me3 deposition at T_{reg}-specific gene loci. To further clarify whether *Cxxc1*

deletion affects FOXP3 binding to its target genes, we performed CUT&Tag experiments to

compare FOXP3 binding profiles between WT and *Cxxc1* KO T_{reg} cells. The results revealed

that most FOXP3-bound regions in WT T_{reg} cells were similarly enriched in KO T_{reg} cells,

indicating that *Cxxc1* deletion does not impair FOXP3's DNA-binding ability (fig. S8A and

B). Together, these findings suggest that the regulatory role of CXXC1 in T_{reg} cells is

mediated through its effect on H3K4me3 deposition rather than altering FOXP3's binding to

DNA.

Discussion

T_{reg} cells specifically express the transcription factor FOXP3, which is essential for

maintaining T_{reg} lineage stability and suppressive function[53]. However, FOXP3 alone is

insufficient to fully regulate the transcriptional signature and functionality of T_{reg} cells; its

interaction with protein partners is crucial for this regulation. In this study, we identify

CXXC1 as a critical epigenetic regulator and functional cofactor of FOXP3. Although

CXXC1 is not required for FOXP3's DNA-binding activity, as evidenced by similar FOXP3

binding patterns in WT and *Cxxc1*-deficient T$_{reg}$ cells (CUT&Tag analysis), it plays an

essential role in maintaining H3K4me3 modifications at FOXP3 target loci. By acting as both

an epigenetic regulator and a FOXP3 cofactor, CXXC1 ensures the stability of the T$_{reg}$

transcriptional program, highlighting its pivotal role in preserving T$_{reg}$ cell functionality and

immune homeostasis.

FOXP3 is known to promote histone H3 acetylation at the promoters and enhancers of its

target genes, such as *Il2ra*, *Ctla4*, and *Tnfrsf18*, following T$_{reg}$ cell activation, thereby

functioning as a transcriptional activator[54,55]. Conversely, FOXP3 can act as a repressor by

silencing target genes like *Il2* and *Ifng* through the induction of histone H3 deacetylation,

mediated by the recruitment of histone deacetylases and transcriptional co-repressors[10,56].

Additionally, FOXP3 exerts this repression by recruiting the Ezh2-containing polycomb

repressive complex to target genes during activation, as FOXP3-repressed genes are

associated with H3K27me3 deposition and reduced chromatin accessibility[11,57]. Recent

studies have begun to explore the biological significance of H3K4me3 breadth, revealing a

positive correlation between H3K4me3 breadth and gene expression [50,58,59]. These studies

also suggest that H3K4me3 breadth contributes to defining specific cell identities during

development and disease, including systemic autoimmune diseases like systemic lupus

erythematosus and various cancers [48,60-63]. Our study demonstrates that the loss of *Cxxc1*

leads to reduced H3K4me3 levels, predominantly in genes with broader peaks, such as *Il2ra,*

Tnfrsf18, Ctla4 and Icos, directly impairing their suppressive function in T$_{reg}$ cells. Although

our research primarily focused on the role of CXXC1 in T$_{reg}$ cells, it is plausible that similar

mechanisms may be operative in other cell types.

The T_{reg}-specific deletion of *Cxxc1* leads to a rapid and fatal autoimmune disorder,

characterized by systemic inflammation and tissue damage, underscoring the essential role of

CXXC1 in maintaining immune self-tolerance within T_{reg} cells. Interestingly, despite this

severe phenotype, our findings show that H3K4me1 levels were comparable between WT and

Cxxc1-deficient T_{reg} cells. In contrast, MLL4 is critical for T_{reg} cell development in the

thymus, primarily by regulating H3K4me1, though it is not required for peripheral T_{reg} cell

function[15]. Numerous studies investigating the effectors of T_{reg} cell-mediated suppression

have identified a wide range of molecules and mechanisms. These include the upregulation of

the inhibitory co-stimulatory receptor CTLA-4, which initiates inhibitory signaling; the

sequestration of the T-cell growth factor IL-2 via CD25; the secretion of inhibitory cytokines,

such as interleukin (IL)-10, IL-35 and TGF-β and the activity of ectoenzymes CD39 and

CD73 on the T_{reg} cells surface, which convert extracellular ATP, a potent pro-inflammatory

mediator, into its anti-inflammatory counterpart, adenosine[42,64]. Mechanistically, we

demonstrated that CXXC1 interacts with FOXP3 to regulate T_{reg} cell function by

trimethylating H3K4 at broad H3K4me3 domains of multiple genes involved in suppressive

functions such as *Il2ra, Nt5e, and Ctla4*. Additionally, MLL1, another KMT, controls T_{reg} cell

activation and function by specifically regulating H3K4 trimethylation at genes encoding key

T_{reg}-related molecules such as *Tigit*, *Klrg1*, *Tbx21*, *Cxcr3*, and serves as a crucial epigenetic

regulator in establishing a stable Th1-T_{reg} lineage[14].

Our findings demonstrate that *Cxxc1*-deficient T_{reg} cells exhibit reduced H3K4me3 levels at

T_{reg}-specific loci, indicating that CXXC1 plays a critical role in regulating this epigenetic

modification. This is consistent with previous studies showing that CXXC1 acts as a

non-catalytic component of the Set1/COMPASS complex [65-68], which includes the H3K4

methyltransferases SETD1A and SETD1B, the primary enzymes responsible for H3K4

trimethylation. Given this, we propose that CXXC1 supports H3K4me3 deposition in T_{reg}

cells by interacting with and stabilizing the activity of the Set1/COMPASS complex. Further

studies are required to directly investigate the interactions between CXXC1 and these

methyltransferases in the T_{reg} cell context.

Our results offer provide novel insights into the suppressive functions, heterogeneity, and

regulatory mechanisms of T_{reg} cells. Maintaining T_{reg} cell homeostasis and function remains a

challenge in harnessing T_{reg} cells for the treatment of autoimmune diseases and the

prevention of graft rejection.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

We thank B. Li (Shanghai Jiao Tong University School of Medicine, Shanghai, China) for

providing *Foxp3*$^{YFP-Cre}$ mice, X.Y. Zhou(Chinese Academy of Science (CAS), Beijing, China)

for providing the FOXP3 CUT&Tag antibody, and X. L. Liu (Shanghai Institutes for

Biological Sciences, Chinese Academy of Sciences) for his generous gifts of cell lines. We

thank Y. Huang, Y. Li, J. Wan, C.Guo, and Z. Lin from the Core Facilities, Zhejiang

University School of Medicine for their technical support and S. Hong, Y. Ding, H. Jin, Q.

Wang, and X. Zhang from Animal Facilities, Zhejiang University, for feeding the mice.

Footnotes

This work was supported by grants from the National Natural Science Foundation of China

(32341002, 32030035, 32321002, 32100693, and 32270839), the National Key R & D

Program of China (2023YFA1800202, 2024YFF0728703, 2022YFA1103702,

2022YFA1103200), the Zhejiang Provincial Natural Science Foundation of China

(LZ21C080001, LZ23C070003), Science and Technology Innovation 2030-Major Project

(2021ZD0200405), Key project of the Experimental Technology Program of Zhejiang

University (SZD202203), and the Fundamental Research Funds for the Central Universities

(226-2024-00161).

Author contributions

657 L.W. and L.S.: supervision, conceptualization, project administration, and writing—review

658 and editing. L.W., L.S. and X.M: funding acquisition. X.M: investigation, methodology,

659 project administration, and writing— original draft. Y.Z.and K.L.: bioinformatics analysis. Y.

660 W., X.L., J.C., C.L., Y.Z., S.W., S.T., Q.X., L.D., X.S., and X.G.: investigation and

661 methodology.

Figure legends

Fig. 1. H3K4me3 is required for FOXP3-dependent gene activation in T_{reg} cells.

(**A**)Venn diagram showing overlap of H3K4me3 enriched regions (this study) and FOXP3

binding sites (Konopacki *et al.*2019) in sorted CD4$^+$YFP$^+$ T$_{reg}$ cells (left). Genomic

distribution of overlapped peaks(right). Note that the overlapped peaks are predominantly

enriched at promoters.

(**B**)Venn diagram showing overlap of H3K27me3 enriched regions (Wei *et al.*2009) and

FOXP3 binding sites in T$_{reg}$ cells (left). Genomic distribution of overlapped peaks (right).

Note that the overlapped peaks are predominantly enriched at promoters.

(**C**)Representative genome browser view showing the enrichments of H3K4me3 and FOXP3

inTconv or T$_{reg}$ cells.

(**D**)Heatmap showing enrichment of H3K27me3, H3K4me3, and FOXP3 surrounding the

TSS. Unsupervised K-means clustering was conducted on H3K27me3 and H3K4me3 signals.

(**E**)Heatmap showing gene expression levels in T$_{reg}$ cells (RNA-seq data was obtained from

Oh *et al.*2017) The clusters were consistent as in Fig. 1C.

(**E**)Gene Ontology (GO) pathway analysis of different clusters.

Fig. 2. CXXC1 interacts with FOXP3 in T$_{reg}$ cell.

(**A**)Average CXXC1 CUT&Tag signals around promoters in T$_{reg}$ cells. IgG was used as the

control.

(**B**)Venn diagrams showing the overlap of FOXP3 binding genes, CXXC1 binding genes, and

H3K4me3 enriched genes in T$_{reg}$ cells. Gene promoters covered by FOXP3 binding sites,

CXXC1 binding sites, or exhibited high H3K4me3 levels were defined as FOXP3-bound

genes, CXXC1-bound genes, or H3K4me3-enriched genes.

(**C**)Heatmaps showing FOXP3 ChIP-seq and CXXC1 CUT&Tag signals at indicated regions.

(**D**)Interaction between FOXP3 and CXXC1 was assessed by co-IP (forward and reverse)

using T_{reg} cell lysates.

(**E**)Immunofluorescence for FOXP3 and CXXC1 colocalization in T_{reg} cells. Scale bars, 2

μm.

Fig.3. *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice spontaneously develop a fatal early-onset inflammatory

disorder.

(**A**)Survival curves of *Foxp3*^{cre} (black line) and *Foxp3*^{cre}*Cxxc1*^{fl/fl} (red line) mice (n = 20).

(**B**)Gross body weight of *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice ($n = 6$).

(**C**)A representative image of *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice.

(**D**)Representative images showing the spleen and peripheral lymph nodes from *Foxp3*^{cre} and

Foxp3^{cre}*Cxxc1*^{fl/fl} mice.

(**E**)ELISA quantification of anti-dsDNA IgG in the serum of *Foxp3*^{cre}

and *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice (n = 4).

(**F**)Hematoxylin and eosin staining of the skin, lung, liver, and colon from *Foxp3*^{cre} an

d *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice (scale bar, 100 μm).

(**G**)Representative flow cytometry plots of CD4 Foxp3 T_{reg} cells isolated from the

small intestinal lamina propria (LPL), liver, and lung of *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*^{fl/f}

^l mice.

(**H**)Flow cytometry analysis of CD62L and CD44 expression on peripheral lymph node

CD4⁺YFP⁻ and CD8⁺T cells from *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice (left). Right, frequency

of CD44^{low}CD62L^{hi} and CD44^{hi}CD62L^{low} population in CD4⁺ YFP⁻ or CD8⁺ T cells (n = 6).

(**I**)Lymph node cells from *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice were stimulated ex vivo with

PMA + ionomycin for 4h and analyzed for IFN-γ expressing in $CD4^+$ YFP^- or $CD8^+$ T cells

using flow cytometry (left). Right, percentages of IFN-γ^+CD4^+ YFP^- or IFN-γ^+CD8^+ T cells

in the lymph nodes of *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice (*n* = 6).

All mice analyzed were 18-20 days old unless otherwise specified. Error bars show mean ±

SD. The log-rank survival curve was used for survival analysis in A, and multiple unpaired

t-test or two-tailed Student's *t*-test were used for statistical analyses in B, E, Gand H

(**P<0.01, ***P□<□0.001, ****P<0.0001). The flow cytometry results are representative

of three independent experiments.

Fig. 4. CXXC1 is essential for T_{reg} cells to suppress T cell-mediated EAE and colitis.

(**A**)Mean clinical scores for EAE in $Rag1^{-/-}$ recipients of 2D2 $CD4^+$ T cells, either alone or in

combination with *Foxp3*^{cre} or *Foxp3*^{cre}*Cxxc1*^{fl/fl} mice after immunization with MOG35–55,

complete Freund's adjuvant (CFA), and pertussis toxin (n=5).

(**B** and **C**)Representative histology of the spinal cord of $Rag1^{-/-}$ mice after EAE induction.

Hematoxylin and eosin(H&E) staining (upper), Luxol fast blue (F&B) staining (lower). Scale

bars, 50 μm (400×) and 200 μm (100×).

(**D**)Representative flow cytometry plots and quantification of the the percentages of IFNγ$^+$ or

IL-17A$^+$ $CD4^+$Vβ11$^+$ T cells (n=5).

(**E**)Statistical analysis of the percentage $CD4^+$ FOXP3$^+$ T_{reg} cell in CNS tissues 14 days after

EAE induction.

(**F**)Changes in body weight of $Rag1^{-/-}$ mice after colitis induction(n=6).

(**G**)Haematoxylin and eosin (H&E) staining of colons from T cell-induced colitis mice 6

weeks after T cell transfer. Scale bars, 50 μm (400×) and 200 μm (100×).

(**H**)Statistical analysis of the percentage CD4$^+$ FOXP3$^+$ T$_{reg}$ cell in the spleen, mesenteric

lymph nodes, and colon 6 weeks after colitis induction.

Error bars show mean ± SD. *P* values are determined by a two-tailed Student's *t*-test or

two-way ANOVA and Holm–Sidak post hoc test (A, D, E and H). (*P<0.05, **P<0.01,

***P□<□0.001, ****P<0.0001).

Fig. 5. Single-cell transcriptomics reveals distinct T$_{reg}$ cell populations.

(**A**)UMAP plot showing clusters identified based on variable gene expression of sorted

CD4$^+$YFP$^+$ T$_{reg}$ cells. Each dot represents a cell, and each color corresponds to a different

population of cell types. Clustering analysis revealed 8 distinct T$_{reg}$ cell populations.

(**B**)Mean fold changes in cluster abundance between *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice.

(**C**)Pseudotime trajectories of T$_{reg}$ cells based on Slingshot, color-coded by T$_{reg}$ cell

subpopulations.

(**D**)Visualization of density and clonotype richness across T$_{reg}$ clusters from *Foxp3*^{cre} and

Foxp3^{cre}*Cxxc1*$^{fl/fl}$ mice.

(**E**)TCR sharing of expanded clonotypes across all possible combinations of T$_{reg}$ cells from

Foxp3^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice.

(**F**)Heatmap showing Z scores for the average expression of T$_{reg}$-specific genes in each

cluster between *Foxp3*^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice.

(**G**-**H**)Representative flow cytometry plots and quantification of (G) expression of CD25,

CD69, ICOS, T-bet, and (H) IFN-γ in CD4$^+$YFP$^+$ T$_{reg}$ cells from

Foxp3^{cre} and *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice (n=5 CD25, n=10 CD69, n=5 ICOS, n=6 T-bet, n=5

IFN-γ).

(**I**)Ki-67 expression (left) and frequency (right) in CD4$^+$FOXP3$^+$ T$_{reg}$ cells from *Foxp3*

^{cre} a-nd *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$ mice (n=6).

Error bars show mean ± SD. *P* values are determined by a two-tailed Student's *t*-test (G-I).

(****P<0.01, ***P□<□0.001, ****P<0.0001**). The flow cytometry results are representative

of three independent experiments.

Fig. 6. *Cxxc1*-deficient T$_{reg}$ cells exhibit functional impairment and disrupted

homeostasis in steady-state conditions.

(**A**)Schematic representation of wild-type and Cre-positive T$_{reg}$ cells in female

Foxp3$^{Cre/+}$mice.

(**B**)Flow cytometry analysis of the YFP$^-$FOXP3$^+$ (WT) and YFP$^+$FOXP3$^+$ (KO) T$_{reg}$ cells in

Foxp3$^{Cre/+}$ *Cxxc1*$^{fl/+}$ (het-WT) and *Foxp3*$^{Cre/+}$ *Cxxc1*$^{fl/fl}$ (het-KO) female mice (left), along

with the frequency and absolute numbers of YFP$^+$ cells within the total T$_{reg}$ population

(right)(n=5).

(**C**)Flow cytometry analysis of Ki-67expression (left) and MFI (right) in YFP$^-$ and YFP$^+$ cells

within the CD4$^+$FOXP3$^+$ T$_{reg}$ cells from 6- to 8-week-old het-KO female mice (n=4).

(**D**)Representative flow cytometry plots and quantification of ICOS, CD25, CTLA4, and

GITR expression in YFP$^-$ and YFP$^+$ cells within CD4$^+$FOXP3$^+$ T$_{reg}$ cells from het-KO female

mice (n=5).

(**E**)Suppression of CFSE-labelled Tn cell proliferation by different ratios of CD4$^+$YFP$^+$ T$_{reg}$

cells from *Foxp3*$^{Cre/+}$ *Cxxc1*$^{fl/+}$ and *Foxp3*$^{Cre/+}$ *Cxxc1*$^{fl/fl}$ female mice. On the right, the

percentage of proliferated responding T cells is presented (n=5).

Error bars show mean ± SD. *P* values are determined by a two-tailed Student's *t*-test (B-E).

(ns, not significant. *$P<0.05$,**$P<0.01$, ***$P<0.001$, ****$P<0.0001$). The flow

cytometry results are representative of three independent experiments.

Fig. 7. CXXC1 regulates Foxp3-dependent molecule H3K4 trimethylation in T$_{reg}$ cells.

(**A**-**B**)Heatmaps showing H3K4me3(A)and CXXC1(B) signals centered on narrow, medium,

and broad domains. The top panel shows the average CUT&Tag signals around indicated

domains.

(**C**)Venn diagrams showing the overlap of FOXP3 binding genes, CXXC1 binding genes, and

H3K4me3-BD-associated genes with decreased H3K4me3 levels after CXXC1 depletion in

T$_{reg}$ cells.

(**D**)Gene Ontology (GO) pathway analysis of the overlap peaks.

(**E**)Representative genome browser view showing the enrichments of FOXP3, CXXC1, and

H3K4me3 in T$_{reg}$ cells.

References

1   Sakaguchi, S., Yamaguchi, T., Nomura, T. & Ono, M. Regulatory T cells and immune

tolerance. Cell 133, 775-787, doi:10.1016/j.cell.2008.05.009 (2008).

2   Josefowicz, S. Z., Lu, L. F. & Rudensky, A. Y. in *Annual Review of Immunology, Vol 30*

Vol. 30 *Annual Review of Immunology* (ed W. E. Paul)   531-564 (2012).

3   Singer, B. D., King, L. S. & D'Alessio, F. R. Regulatory T cells as immunotherapy.

Frontiers in Immunology 5, doi:10.3389/fimmu.2014.00046 (2014).

4   Ohkura, N. & Sakaguchi, S. Transcriptional and epigenetic basis of Treg cell

development and function: its genetic anomalies or variations in autoimmune diseases. Cell

Research 30, 465-474, doi:10.1038/s41422-020-0324-7 (2020).

5   Littman, D. R. & Rudensky, A. Y. Th17 and Regulatory T Cells in Mediating and

Restraining Inflammation. Cell 140, 845-858, doi:10.1016/j.cell.2010.02.021 (2010).

6   Sakaguchi, S., Vignali, D. A. A., Rudensky, A. Y., Niec, R. E. & Waldmann, H. The

plasticity and stability of regulatory T cells. Nature Reviews Immunology 13, 461-467,

doi:10.1038/nri3464 (2013).

7   Fontenot, J. D., Gavin, M. A. & Rudensky, A. Y. Foxp3 programs the development and

function of CD4+CD25+ regulatory T cells. Nat Immunol 4, 330-336, doi:10.1038/ni904

(2003).

8   Van Gool, F., Nguyen, M. L. T., Mumbach, M. R., Satpathy, A. T., Rosenthal, W. L.,

Giacometti, S., Le, D. T., Liu, W., Brusko, T. M., Anderson, M. S., Rudensky, A. Y., Marson,

A., Chang, H. Y. & Bluestone, J. A. A Mutation in the Transcription Factor Foxp3 Drives T

Helper 2 Effector Function in Regulatory T Cells. Immunity 50, 362-+,

doi:10.1016/j.immuni.2018.12.016 (2019).

9   Katoh, H., Qin, Z. S., Liu, R., Wang, L., Li, W., Li, X., Wu, L., Du, Z., Lyons, R., Liu,

C.-G., Liu, X., Dou, Y., Zheng, P. & Liu, Y. FOXP3 Orchestrates H4K16 Acetylation and

H3K4 Trimethylation for Activation of Multiple Genes by Recruiting MOF and Causing

Displacement of PLU-1. Molecular Cell 44, 770-784, doi:10.1016/j.molcel.2011.10.012

(2011).

10   Wang, L., Liu, Y., Han, R., Beier, U. H., Bhatti, T. R., Akimova, T., Greene, M. I.,

Hiebert, S. W. & Hancock, W. W. FOXP3(+) regulatory T cell development and function

require histone/protein deacetylase 3. Journal of Clinical Investigation 125, 1111-1123,

doi:10.1172/jci77088 (2015).

11   Arvey, A., van der Veeken, J., Samstein, R. M., Feng, Y., Stamatoyannopoulos, J. A. &

Rudensky, A. Y. Inflammation-induced repression of chromatin bound by the transcription

factor Foxp3 in regulatory T cells. Nature Immunology 15, 580-587, doi:10.1038/ni.2868

(2014).

12   Ruthenburg, A. J., Allis, C. D. & Wysocka, J. Methylation of lysine 4 on histone H3:

Intricacy of writing and reading a single epigenetic mark. Molecular Cell 25, 15-30,

doi:10.1016/j.molcel.2006.12.014 (2007).

13   Takahashi, Y.-h., Westfield, G. H., Oleskie, A. N., Trievel, R. C., Shilatifard, A. &

Skiniotis, G. Structural analysis of the core COMPASS family of histone H3K4 methylases

from yeast to human. Proceedings of the National Academy of Sciences of the United States

of America 108, 20526-20531, doi:10.1073/pnas.1109360108 (2011).

14   Wang, T., Guo, J., Li, L., Jin, Q., Zhang, F., Hou, B., Zhang, Y. & Zhou, X. The histone

lysine methyltransferase MLL1 regulates the activation and functional specialization of

regulatory T cells. Cell Reports 43, doi:10.1016/j.celrep.2024.114222 (2024).

15   Placek, K., Hu, G., Cui, K., Zhang, D., Ding, Y., Lee, J.-E., Jang, Y., Wang, C., Konkel, J.

E., Song, J., Liu, C., Ge, K., Chen, W. & Zhao, K. MLL4 prepares the enhancer landscape for

Foxp3 induction via chromatin looping. Nature Immunology 18, 1035-+, doi:10.1038/ni.3812

(2017).

16   Yang, Y., Yang, Y., Chan, K. & Couture, J.-F. Analyzing the impact of CFP1 mutational

landscape on epigenetic signaling. Faseb Journal 35, doi:10.1096/fj.202100427R (2021).

17   Tate, C. M., Lee, J.-H. & Skalnik, D. G. CXXC finger protein 1 restricts the Setd1A

histone H3K4 methyltransferase complex to euchromatin. Febs Journal 277, 210-223,

doi:10.1111/j.1742-4658.2009.07475.x (2010).

18   Cao, W., Guo, J., Wen, X., Miao, L., Lin, F., Xu, G., Ma, R., Yin, S., Hui, Z., Chen, T.,

Guo, S., Chen, W., Huang, Y., Liu, Y., Wang, J., Wei, L. & Wang, L. CXXC finger protein 1 is

critical for T-cell intrathymic development through regulating H3K4 trimethylation. Nat

Commun 7, 11687, doi:10.1038/ncomms11687 (2016).

19   Hui, Z., Zhou, L., Xue, Z., Zhou, L., Luo, Y., Lin, F., Liu, X., Hong, S., Li, W., Wang, D.,

Lu, L., Wang, J. & Wang, L. Cxxc Finger Protein 1 Positively Regulates GM-CSF-Derived

Macrophage Phagocytosis Through Csf2rα-Mediated Signaling. Frontiers in Immunology 9,

doi:10.3389/fimmu.2018.01885 (2018).

20   Lin, F., Meng, X., Guo, Y., Cao, W., Liu, W., Xia, Q., Hui, Z., Chen, J., Hong, S., Zhang,

X., Wu, C., Wang, D., Wang, J., Lu, L., Qian, W., Wei, L. & Wang, L. Epigenetic initiation of

the T(H)17 differentiation program is promoted by Cxxc finger protein 1. Science Advances

5, doi:10.1126/sciadv.aax1608 (2019).

21 Shen, X., Gao, X., Luo, Y., Xu, Q., Fan, Y., Hong, S., Huang, Z., Liu, X., Wang, Q., Chen,

Z., Wang, D., Lu, L., Wu, C., Liang, H. & Wang, L. Cxxc finger protein 1 maintains

homeostasis and function of intestinal group 3 innate lymphoid cells with aging. Nature

Aging 3, 965-+, doi:10.1038/s43587-023-00453-7 (2023).

22 Cao, W., Guo, J., Wen, X., Miao, L., Lin, F., Xu, G., Ma, R., Yin, S., Hui, Z., Chen, T.,

Guo, S., Chen, W., Huang, Y., Liu, Y., Wang, J., Wei, L. & Wang, L. CXXC finger protein 1 is

critical for T-cell intrathymic development through regulating H3K4 trimethylation. Nature

Communications 7, doi:10.1038/ncomms11687 (2016).

23 Zhou, L., Wang, S., Hu, W., Liu, X., Xu, L., Tong, B., Zhang, T., Xue, Z., Guo, Y., Zhao,

J., Lu, L., Fan, H., Qian, W., Chen, J., Chen, W. & Wang, L. T cell proliferation requires

ribosomal maturation in nucleolar condensates dependent on DCAF13. Journal of Cell

Biology 222, doi:10.1083/jcb.202201096 (2023).

24 Chou, W.-C., Guo, Z., Guo, H., Chen, L., Zhang, G., Liang, K., Xie, L., Tan, X., Gibson,

S. A., Rampanelli, E., Wang, Y., Montgomery, S. A., Brickey, W. J., Deng, M., Freeman, L.,

Zhang, S., Su, M. A., Chen, X., Wan, Y. Y. & Ting, J. P. Y. AIM2 in regulatory T cells

restrains autoimmune diseases. Nature 591, 300-+, doi:10.1038/s41586-021-03231-w (2021).

25 Zeng, H., Yang, K., Cloer, C., Neale, G., Vogel, P. & Chi, H. mTORC1 couples immune

signals and metabolic programming to establish T(reg)-cell function. Nature 499, 485-490,

doi:10.1038/nature12297 (2013).

26 Lu, D., Liu, L., Sun, Y., Song, J., Yin, Q., Zhang, G., Qi, F., Hu, Z., Yang, Z., Zhou, Z.,

Hu, Y., Zhang, L., Ji, J., Zhao, X., Jin, Y., McNutt, M. A. & Yin, Y. The phosphatase PAC1

acts as a T cell suppressor and attenuates host antitumor immunity. Nature Immunology 21,

287-+, doi:10.1038/s41590-019-0577-9 (2020).

27 Chen, J., Liu, K., Luo, Y., Kang, M., Wang, J., Chen, G., Qi, J., Wu, W., Wang, B., Han,

Y., Shi, L., Wang, K., Han, X., Ma, X., Liu, W., Ding, Y., Wang, L., Liang, H., Wang, L. &

Chen, J. Single-Cell Profiling of Tumor Immune Microenvironment Reveals Immune

Irresponsiveness in Gastric Signet-Ring Cell Carcinoma. Gastroenterology 165, 88-103,

doi:10.1053/j.gastro.2023.03.008 (2023).

28 Marson, A., Kretschmer, K., Frampton, G. M., Jacobsen, E. S., Polansky, J. K., MacIsaac,

K. D., Levine, S. S., Fraenkel, E., von Boehmer, H. & Young, R. A. Foxp3 occupancy and

regulation of key target genes during T-cell stimulation. Nature 445, 931-935,

doi:10.1038/nature05478 (2007).

29 Zheng, Y., Josefowicz, S. Z., Kas, A., Chu, T.-T., Gavin, M. A. & Rudensky, A. Y.

Genome-wide analysis of Foxp3 target genes in developing and mature regulatory T cells.

Nature 445, 936-940, doi:10.1038/nature05563 (2007).

30 Ono, M., Yaguchi, H., Ohkura, N., Kitabayashi, I., Nagamura, Y., Nomura, T., Miyachi,

Y., Tsukada, T. & Sakaguchi, S. Foxp3 controls regulatory T-cell function by interacting with

AML1/Runx1. Nature 446, 685-689, doi:10.1038/nature05673 (2007).

31 Fu, W., Ergun, A., Lu, T., Hill, J. A., Haxhinasto, S., Fassett, M. S., Gazit, R., Adoro, S.,

Glimcher, L., Chan, S., Kastner, P., Rossi, D., Collins, J. J., Mathis, D. & Benoist, C. A

multiply redundant genetic switch 'locks in' the transcriptional signature of regulatory T cells.

Nature Immunology 13, 972-980, doi:10.1038/ni.2420 (2012).

32   Ohkura, N., Hamaguchi, M., Morikawa, H., Sugimura, K., Tanaka, A., Ito, Y., Osaki, M.,

Tanaka, Y., Yamashita, R., Nakano, N., Huehn, J., Fehling, H. J., Sparwasser, T., Nakai, K. &

Sakaguchi, S. T Cell Receptor Stimulation-Induced Epigenetic Changes and Foxp3

Expression Are Independent and Complementary Events Required for Treg Cell

Development. Immunity 37, 785-799, doi:10.1016/j.immuni.2012.09.010 (2012).

33   Wei, G., Wei, L., Zhu, J., Zang, C., Hu-Li, J., Yao, Z., Cui, K., Kanno, Y., Roh, T.-Y.,

Watford, W. T., Schones, D. E., Peng, W., Sun, H.-w., Paul, W. E., O'Shea, J. J. & Zhao, K.

Global Mapping of H3K4me3 and H3K27me3 Reveals Specificity and Plasticity in Lineage

Fate   Determination   of   Differentiating   CD4(+)   T   Cells.   Immunity   30,   155-167,

doi:10.1016/j.immuni.2008.12.009 (2009).

34   Konopacki, C., Pritykin, Y., Rubtsov, Y., Leslie, C. S. & Rudensky, A. Y. Transcription

factor Foxp1 regulates Foxp3 chromatin binding and coordinates regulatory T cell function.

Nature Immunology 20, 232-+, doi:10.1038/s41590-018-0291-z (2019).

35   Hill, J. A., Feuerer, M., Tash, K., Haxhinasto, S., Perez, J., Melamed, R., Mathis, D. &

Benoist, C. Foxp3 transcription-factor-dependent and -independent regulation of the

regulatory   T   cell   transcriptional   signature.   Immunity   27,   786-800,

doi:10.1016/j.immuni.2007.09.010 (2007).

36   Oh, H., Grinberg-Bleyer, Y., Liao, W., Maloney, D., Wang, P., Wu, Z., Wang, J., Bhatt, D.

M., Heise, N., Schmid, R. M., Hayden, M. S., Klein, U., Rabadan, R. & Ghosh, S. An

NF-kappaB   Transcription-Factor-Dependent   Lineage-Specific   Transcriptional   Program

Promotes   Regulatory   T   Cell   Identity   and   Function.   Immunity   47,   450-465   e455,

doi:10.1016/j.immuni.2017.08.010 (2017).

37   Rubtsov, Y. P., Rasmussen, J. P., Chi, E. Y., Fontenot, J., Castelli, L., Ye, X., Treuting, P.,

Siewe, L., Roers, A., Henderson, W. R., Jr., Muller, W. & Rudensky, A. Y. Regulatory T

cell-derived interleukin-10 limits inflammation at environmental interfaces. Immunity 28,

546-558, doi:10.1016/j.immuni.2008.02.017 (2008).

38   Fontenot, J. D., Gavin, M. A. & Rudensky, A. Y. Foxp3 programs the development and

function of CD4(+)CD25(+) regulatory T cells. Nature Immunology 4, 330-336,

doi:10.1038/ni904 (2003).

39   Kim, J. M., Rasmussen, J. P. & Rudensky, A. Y. Regulatory T cells prevent catastrophic

autoimmunity throughout the lifespan of mice. Nature Immunology 8, 191-197,

doi:10.1038/ni1428 (2007).

40   Shevyrev, D. & Tereshchenko, V. Treg Heterogeneity, Function, and Homeostasis.

Frontiers in Immunology 10, doi:10.3389/fimmu.2019.03100 (2020).

41   Zagorulya, M., Yim, L., Morgan, D. M., Edwards, A., Torres-Mejia, E., Momin, N.,

McCreery, C. V., Zamora, I. L., Horton, B. L., Fox, J. G., Wittrup, K. D., Love, J. C. &

Spranger, S. Tissue-specific abundance of interferon-gamma drives regulatory T cells to

restrain DC1-mediated priming of cytotoxic T cells against lung cancer. Immunity 56, 386-+,

doi:10.1016/j.immuni.2023.01.010 (2023).

42   Dikiy, S. & Rudensky, A. Y. Principles of regulatory T cell function. Immunity 56,

240-255, doi:10.1016/j.immuni.2023.01.004 (2023).

43   Street, K., Risso, D., Fletcher, R. B., Das, D., Ngai, J., Yosef, N., Purdom, E. & Dudoit, S.

Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. Bmc

Genomics 19, doi:10.1186/s12864-018-4772-0 (2018).

44   Tarbell, K. V., Yamazaki, S., Olson, K., Toy, P. & Steinman, R. M. CD25+CD4+T cells,

expanded with dendritic cells presenting a single autoantigenic peptide, suppress autoimmune

diabetes. Journal of Experimental Medicine 199, 1467-1477, doi:10.1084/jem.20040180

(2004).

45   Hori, S., Haury, M., Coutinho, A. & Demengeot, J. Specificity requirements for selection

and effector functions of CD25+4+ regulatory T cells in anti-myelin basic protein T cell

receptor transgenic mice. Proceedings of the National Academy of Sciences of the United

States of America 99, 8213-8218, doi:10.1073/pnas.122224799 (2002).

46   Butler, J. S., Lee, J.-H. & Skalnik, D. G. CFP1 Interacts with DNMT1 Independently of

Association with the Setd1 Histone H3K4 Methyltransferase Complexes. DNA and Cell

Biology 27, 533-543, doi:10.1089/dna.2007.0714 (2008).

47   Butler, J. S., Palam, L. R., Tate, C. M., Sanford, J. R., Wek, R. C. & Skalnik, D. G. DNA

Methyltransferase Protein Synthesis Is Reduced in CXXC Finger Protein 1-Deficient

Embryonic Stem Cells. DNA and Cell Biology 28, 223-231, doi:10.1089/dna.2009.0854

(2009).

48   Benayoun, B. A., Pollina, E. A., Ucar, D., Mahmoudi, S., Karra, K., Wong, E. D.,

Devarajan, K., Daugherty, A. C., Kundaje, A. B., Mancini, E., Hitz, B. C., Gupta, R., Rando,

T. A., Baker, J. C., Snyder, M. P., Cherry, J. M. & Brunet, A. H3K4me3 Breadth Is Linked to

Cell   Identity   and   Transcriptional   Consistency.   Cell   158,   673-688,

doi:10.1016/j.cell.2014.06.027 (2014).

49   Zacarias-Cabeza, J., Belhocine, M., Vanhille, L., Cauchy, P., Koch, F., Pekowska, A.,

Fenouil, R., Bergon, A., Gut, M., Gut, I., Eick, D., Imbert, J., Ferrier, P., Andrau, J.-C. &

Spicuglia, S. Transcription-Dependent Generation of a Specialized Chromatin Structure at the

TCRβ Locus. Journal of Immunology 194, 3432-3443, doi:10.4049/jimmunol.1400789

(2015).

50   Chen, K., Chen, Z., Wu, D., Zhang, L., Lin, X., Su, J., Rodriguez, B., Xi, Y., Xia, Z.,

Chen, X., Shi, X., Wang, Q. & Li, W. Broad H3K4me3 is associated with increased

transcription elongation and enhancer activity at tumor-suppressor genes. Nature Genetics 47,

1149-+, doi:10.1038/ng.3385 (2015).

51   Zhang, R., Huynh, A., Whitcher, G., Chang, J., Maltzman, J. S. & Turka, L. A. An

obligate cell-intrinsic function for CD28 in Tregs. Journal of Clinical Investigation 123,

580-593, doi:10.1172/jci65013 (2013).

52   Gokhale, A. S., Gangaplara, A., Lopez-Occasio, M., Thornton, A. M. & Shevach, E. M.

Selective deletion of Eos (Ikzf4) in T-regulatory cells leads to loss of suppressive function

and   development   of   systemic   autoimmunity.   Journal   of   Autoimmunity   105,

doi:10.1016/j.jaut.2019.06.011 (2019).

53   Wan, Y. Y. & Flavell, R. A. Regulatory T-cell functions are subverted and converted

owing to attenuated Foxp3 expression. Nature 445, 766-770, doi:10.1038/nature05479

(2007).

54   Chen, C., Rowell, E. A., Thomas, R. M., Hancock, W. W. & Wells, A. D. Transcriptional

regulation by Foxp3 is associated with direct promoter occupancy and modulation of histone

acetylation.   Journal   of   Biological   Chemistry   281,   36828-36834,

doi:10.1074/jbc.M608848200 (2006).

55   Morikawa, H., Ohkura, N., Vandenbon, A., Itoh, M., Nagao-Sato, S., Kawaji, H.,

Lassmann, T., Carninci, P., Hayashizaki, Y., Forrest, A. R. R., Standley, D. M., Date, H., Sakaguchi, S. & Consortium, F. Differential roles of epigenetic changes and Foxp3 expression in regulatory T cell-specific transcriptional regulation. Proceedings of the National Academy of Sciences of the United States of America 111, 5289-5294, doi:10.1073/pnas.1312717110 (2014).

56   Pan, F., Yu, H., Dang, E. V., Barbi, J., Pan, X., Grosso, J. F., Jinasena, D., Sharma, S. M., McCadden, E. M., Getnet, D., Drake, C. G., Liu, J. O., Ostrowski, M. C. & Pardoll, D. M. Eos Mediates Foxp3-Dependent Gene Silencing in CD4+ Regulatory T Cells. Science 325, 1142-1146, doi:10.1126/science.1176077 (2009).

57 DuPage, M., Chopra, G., Quiros, J., Rosenthal, W. L., Morar, M. M., Holohan, D., Zhang, R., Turka, L., Marson, A. & Bluestone, J. A. The Chromatin-Modifying Enzyme Ezh2 Is Critical for the Maintenance of Regulatory T Cell Identity after Activation. Immunity 42, 227-238, doi:10.1016/j.immuni.2015.01.007 (2015).

58   Liu, X., Wang, C., Liu, W., Li, J., Li, C., Kou, X., Chen, J., Zhao, Y., Gao, H., Wang, H., Zhang, Y., Gao, Y. & Gao, S. Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation embryos. Nature 537, 558-+, doi:10.1038/nature19362 (2016).

59 Sze, C. C., Ozark, P. A., Cao, K., Ugarenko, M., Das, S., Wang, L., Marshall, S. A., Rendleman, E. J., Ryan, C. A., Zha, D., Douillet, D., Chen, F. X. & Shilatifard, A. Coordinated regulation of cellular identity-associated H3K4me3 breadth by the COMPASS family. Science Advances 6, doi:10.1126/sciadv.aaz4764 (2020).

60   Dahl, J. A., Jung, I., Aanes, H., Greggains, G. D., Manaf, A., Lerdrup, M., Li, G., Kuan, S., Li, B., Lee, A. Y., Preissl, S., Jermstad, I., Haugen, M. H., Suganthan, R., Bjoras, M.,

Hansen, K., Dalen, K. T., Fedorcsak, P., Ren, B. & Klungland, A. Broad histone H3K4me3

domains in mouse oocytes modulate maternal-to-zygotic transition. Nature 537, 548-+,

doi:10.1038/nature19360 (2016).

61   Zhang, B., Zheng, H., Huang, B., Li, W., Xiang, Y., Peng, X., Ming, J., Wu, X., Zhang, Y.,

Xu, Q., Liu, W., Kou, X., Zhao, Y., He, W., Li, C., Chen, B., Li, Y., Wang, Q., Ma, J., Yin, Q.,

Kee, K., Meng, A., Gao, S., Xu, F., Na, J. & Xie, W. Allelic reprogramming of the histone

modification   H3K4me3   in   early   mammalian   development.   Nature   537,   553-+,

doi:10.1038/nature19361 (2016).

62   Zhang, Z., Shi, L., Dawany, N., Kelsen, J., Petri, M. A. & Sullivan, K. E. H3K4

tri-methylation breadth at transcription start sites impacts the transcriptome of systemic lupus

erythematosus. Clinical Epigenetics 8, doi:10.1186/s13148-016-0179-4 (2016).

63   Wong, S. H. K., Goode, D. L., Iwasaki, M., Wei, M. C., Kuo, H.-P., Zhu, L.,

Schneidawind, D., Duque-Afonso, J., Weng, Z. & Cleary, M. L. The H3K4-Methyl

Epigenome Regulates Leukemia Stem Cell Oncogenic Potential. Cancer Cell 28, 198-209,

doi:10.1016/j.ccell.2015.06.003 (2015).

64   Deaglio, S., Dwyer, K. M., Gao, W., Friedman, D., Usheva, A., Erat, A., Chen, J.-F.,

Enjyoji, K., Linden, J., Oukka, M., Kuchroo, V. K., Strom, T. B. & Robson, S. C. Adenosine

generation catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune

suppression. Journal of Experimental Medicine 204, 1257-1265, doi:10.1084/jem.20062512

(2007).

65   Lee, J. H. & Skalnik, D. G. CpG-binding protein (CXXC finger protein 1) is a component

of the mammalian set1 histone H3-Lys4 methyltransferase complex, the analogue of the yeast

Set1/COMPASS complex. Journal of Biological Chemistry 280, 41725-41731,

doi:10.1074/jbc.M508312200 (2005).

66 Thomson, J. P., Skene, P. J., Selfridge, J., Clouaire, T., Guy, J., Webb, S., Kerr, A. R. W.,

Deaton, A., Andrews, R., James, K. D., Turner, D. J., Illingworth, R. & Bird, A. CpG islands

influence chromatin structure via the CpG-binding protein Cfp1. Nature 464, 1082-U1162,

doi:10.1038/nature08924 (2010).

67 Shilatifard, A. in *Annual Review of Biochemistry, Vol 81* Vol. 81 *Annual Review of*

Biochemistry (ed R. D. Kornberg)   65-95 (2012).

68 Brown, D. A., Di Cerbo, V., Feldmann, A., Ahn, J., Ito, S., Blackledge, N. P., Nakayama,

M., McClellan, M., Dimitrova, E., Turberfield, A. H., Long, H. K., King, H. W., Kriaucionis,

S., Schermelleh, L., Kutateladze, T. G., Koseki, H. & Klose, R. J. The SET1 Complex Selects

Actively Transcribed Target Genes via Multivalent Interaction with CpG Island Chromatin.

Cell Reports 20, 2313-2327, doi:10.1016/j.celrep.2017.08.030 (2017).

1049

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6

Fig.7

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 4

A

Lymph node

*Foxp3*cre   *Foxp3*cre*Cxxc1*fl/fl

CD4

FOXP3

9.95   19.40

● *Foxp3*cre
● *Foxp3*cre*Cxxc1*fl/fl

CD4+ FOXP3+ (%)

**

Cell number (x10⁴)

****

B

--- *Foxp3*cre
—— *Foxp3*cre*Cxxc1*fl/fl

Count

FOXP3

● *Foxp3*cre
● *Foxp3*cre*Cxxc1*fl/fl

Foxp3 MFI

ns

C

*Foxp3*cre

79.1   13.4   29.2   60.5

0 : 1

*Foxp3*cre*Cxxc1*fl/fl

21.1   30.1   55.1

Count

CFSE   1 : 2   1 : 4   1 : 8

Regulatory T cell to effector T cell ratio

● *Foxp3*cre
● *Foxp3*cre*Cxxc1*fl/fl

Percentage(%)

ns   ns   ns

1:2   1:4   1:8

D

▩ Isotype --- *Foxp3*cre —— *Foxp3*cre*Cxxc1*fl/fl

CD4+YFP+

Count

CTLA-4   GITR

● *Foxp3*cre
● *Foxp3*cre*Cxxc1*fl/fl

CTLA4(MFI)

**

GITR (MFI)

ns

Supplementary Figure 5

A

Ccr7
Tcf7
Lef1
Sell
Myc
Bcl2
Cxcr3
Nrp1
Klrg1
Il1rl1
Igals3
Cxcr3
Rora
Gzmb
Il10
Gzma
Tigit
Lag3
Pdcd1
Tnfsf14
Nt5e
Lap3
Tnfrsf9
Tmem97
Irf8
Cd160
H2-Eb1
H2-Ab1
H2-Aa
H2-DMb2
H2-DMa
Mki67
Ube2c
Top2a
Aurkb

Naive subsets
pre activated subsets
Klrg1$^+$ activated subsets
Gzmb$^+$ activated subsets
Nt5e$^+$ activated subsets
Tnfsf9$^+$ activated subsets
H2-Eb1$^+$ activated subsets
Mki67$^+$ subsets

B

Foxp3^{cre} *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$

UMAP_2

UMAP_1

- Naive subsets
- pre actived subsets
- Klrg1$^+$ actived subsets
- Gzmb$^+$ actived subsets
- Nt5e$^+$ actived subsets
- Tnfrsf9$^+$ actived subsets
- H2−Eb1$^+$ actived subsets
- Mki67$^+$ subsets

C

Frequency

Foxp3^{cre} *Foxp3*^{cre} *Cxxc1*$^{fl/fl}$

- Naive subsets
- pre activated subsets
- Klrg1$^+$ activated subsets
- Gzmb$^+$ actived subsets
- Nt5e$^+$ activated subsets
- Tnfrsf9$^+$ activated subsets
- H2−Eb1$^+$ activated subsets
- Mki67$^+$ subsets

D

- Single (0<X<=1)
- Small (1<X<=5)
- Medium (5<X<=20)
- NA

UMAP 2

UMAP 1

E

Isotype --- *Foxp3*^{cre} — *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$

CD4$^+$YFP$^+$

Count

KLRG1   CD62L   CD73   PD-1

● *Foxp3*^{cre} ● *Foxp3*^{cre}*Cxxc1*$^{fl/fl}$

KLRG1(MFI)   CD62L(MFI)   CD73(MFI)   PD-1 (MFI)

****   ****   ****   ****

Supplementary Figure 6

A

2D2 Only vs 2D2+Treg(*Foxp3*^cre/+^*Cxxc1*^fl/+^), ****$p<0.0001$
2D2 Only vs 2D2+Treg(*Foxp3*^cre/+^*Cxxc1*^fl/fl^), p=ns
2D2+Treg(*Foxp3*^cre/+^*Cxxc1*^fl/+^) vs 2D2+Treg(*Foxp3*^cre/+^*Cxxc1*^fl/fl^), *** p=0.0007

Supplementary Figure 7

Supplementary Figure 8

A

*Foxp3*cre   *Foxp3*cre *Cxxc1*fl/fl

Foxp3 peaks

B

chr1:60,953,251-60,985,265

chr4:155,395,406-155,408,039

*Foxp3*cre

*Foxp3*cre *Cxxc1*fl/fl

Ctla4   *Tnfrsf18*

chr2:11,470,358-11,706,417

chr9:88,164,041-88,325,194

*Foxp3*cre

*Foxp3*cre *Cxxc1*fl/fl

Il2ra   *Nt5e*