1

2 **_Clostridioides difficile_ in Equidae necropsied in north-western France, in 2019-2021:**

3 **common carriage of toxigenic strains and virulence of emerging ribotype 017**

Sandrine Petry[1,a], Jackie Tapprest[1,b], Karine Maillard[2], Frédéric Barbut[3], Fabien Duquesne[1,c], Sofia

Kozak[1,d], Nathalie Foucher[1,e], Maud Bernez-Romand[1,f], Ludovic Bridoux[4,¤], Isabelle Poquet[4, *]

[1] ANSES, Normandy Laboratory for Animal Health, Physiopathology and Epidemiology of Equine

Diseases Unit (PhEED), Goustranville, France

[a] Sandrine.PETRY@anses.fr

[b] present address: jackie.tapprest@agriculture.gouv.fr; Direction Générale de l'Enseignement et de

la Recherche (DGER), Ministère de l'Agriculture, de la Souveraineté Alimentaire et de la Forêt,

Paris, France

[c] Fabien.DUQUESNE@anses.fr

[d] present address: sofiakozak@hotmail.fr

[e] Nathalie.FOUCHER@anses.fr

[f] maud.bernez-romand@anses.fr

[2] LABÉO Frank Duncombe, Caen, France

present address: karine.maillard@calvados.fr; Pôle Système d'Information des Ressources

Humaines, Département du Calvados, France

[3] National Reference Laboratory for _C. difficile_, Hôpital Saint Antoine, Assistance Publique Hôpitaux

de Paris (AP-HP) Paris, France and INSERM S-1139, Université Paris-Cité, Paris, France

frederic.barbut@aphp.fr

[4] Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350, Jouy-en-Josas, France

[¤] ludovic.bridoux@inrae.fr

* Corresponding author: isabelle.poquet@inrae.fr, ORCID https://orcid.org/0000-0003-3514-6520

Abstract

Clostridioides difficile is an anaerobic, spore-forming entero-pathogen, causing diarrhea in humans and animals. We studied French Equidae as a model host to evaluate *C. difficile* occurrence, diversity, virulence and circulation. We took advantage of necropsies of Equidae in northwestern France (Normandy), from 2019 to 2021 (n=100), to systematically recover the caecal content and, if watery and possibly diarrheal, any intestinal content. We searched for *C. difficile* and characterized isolates for their toxin gene profile and PCR-ribotype according to the WEBRIBO database. We performed free toxin detection to reach a high-standard diagnosis of infection.

Twenty-seven Equidae were positive for *C. difficile* and twenty had a toxigenic strain, including one animal also colonized by a non-toxigenic strain. Toxigenic isolates belonged to eight common ribotypes: 078, 126 (three-toxin profile), 005, 012, 020, 181, AI-53 (two-toxin profile), 017 (one-toxin profile), and non-toxigenic isolates to ribotypes 009, 035 and 439. Ribotypes 017 and 009 were predominant (n=5). In two premises, some Equidae shared the same ribotype, 020 or 009, suggesting potential transmission clusters. In four animals with signs of diarrhea and toxigenic isolates, the detection of free toxin demonstrated *C. difficile* infection. These isolates were of ribotype 126 (n=1) or 017 (n=3) and the least susceptible to vancomycin. Two animals died from their infection by ribotype 017, which is emerging in Asia but infrequent in Europe.

French Equidae represented an underestimated reservoir of pathogenic *C. difficile* ribotypes and one of them was highly virulent, both preoccupying findings in a One Health perspective*.*

Importance

C. difficile, a major entero-pathogen widely disseminated in the environment, is a One Health issue. Animals are raising concern as human contamination sources. Equidae are in close contact with humans and exposed to the same risk factors. The systematic survey of Equidae necropsied from 2019 to 2021 in the first breeding region in France, revealed that 20% harboured pathogenic strains. These belonged to prevalent, clinically important ribotypes, raising the possibility of zoonotic transmission. A high-standard diagnosis of *C. difficile* infection (CDI), recommended in European hospitals, but rare in animals, revealed CDI in 4% of the population, and prior antibiotic treatment was a major risk factor. A ribotype known as emerging in Asia was highly virulent and caused a local infection cluster.

French Equidae as underestimated reservoirs of *C. difficile* and notably of ribotype 017 are of concern in a One Health perspective and should be carefully surveilled*.*

Keywords

Introduction

C. difficile is a spore-forming anaerobe responsible for diarrhea and colitis (1). It is a major cause of health-care associated diarrhea and a rising cause of community-acquired infections. *C. difficile* infections (CDIs), which are transmitted by spores by the fecal-oral route, can be severe, with complications and recurrences, and life threatening. TcdA and TcdB toxins are responsible for symptoms, even though TcdB alone is sufficient for disease, and CDT, the binary toxin, has an accessory role (2). Non-toxigenic strains are unable to cause disease, while toxigenic strains can cause disease or be asymptomatically carried (3). Antibiotics, hospitalization and age are the main risk factors for CDI (1). Antibiotics notably induce microbiota dysbiosis, which favors spore germination and vegetative multiplication. CDIs are treated by fidaxomicin and vancomycin (4), which contribute to dysbiosis and therefore relapses. Spores and biofilms could contribute to persistence between relapses (5, 6). The last resort treatment is fecal microbiota transplant (FMT) (4).

C. difficile is a One Health issue (7-10). It causes diarrhea in most, wild, farm and companion animals (8, 9, 11-13). *C. difficile* strains can jump between hosts, leading to inter-species transmission (10, 14-16). Equidae, which are closely interacting with humans, are relevant to zoonotic and anthropo-zoonotic transmission. Importantly, the healthcare of these monogastric animals of high economic value (https://www.ifce.fr) is careful, and they have the same main risk factors for CDI as humans, antibiotics and hospitalization. In both hosts*, C. difficile* is responsible for outbreaks and sporadic cases (17-21). Both have long been treated by the same antibiotics, metronidazole and vancomycin, until fidaxomicin substituted for metronidazole in the hospital, and can be treated by FMT (22). As infected populations, their main difference is age. While <3-years-old children are mainly asymptomatic carriers (3), CDI can affect foals (17, 18).

Data about *C. difficile* importance in horses (7, 9, 11, 19) remain scarce in France. *C. difficile* testing is not systematic and the detection of a toxigenic strain is not sufficient to diagnose CDI (3). Here, we studied *C. difficile* in Equidae necropsied in Normandy, the first breeding region in France (12,600 foals and 145,000 Equidae in 2021, https://www.ifce.fr). Necropsy (23) was the opportunity both to recover digestive contents from a population with diverse clinical pictures, and to get access to animal data as detailed as patient data, together with the cause of their death. Our objectives were to evaluate i) *C. difficile* occurrence, diversity and putative circulation among French Equidae, and ii) the prevalence of CDI or carriage after a *post-mortem* diagnosis including free toxin detection.

Materials and methods

Equine necropsy

Equine necropsies (ANSES, PhEED) are performed to determine the cause of animal death, using a standardized procedure (23). It includes examination of the cadaver, evisceration and organ examination. Animal identification data, previous treatments, necropsy observations, pathogen identification and death cause are registered (https://sitesv2.anses.fr/fr/minisite/resumeq/pr%C3%A9sentation-de-resumeq) (23).

In 2018, the intestinal contents of five necropsied Equidae with *post-mortem* signs of endo-enterotoxaemia and diarrhea were collected for a pilot study. Then, for all animals necropsied from May 2019 to August 2021 (except fetus and stillborn foals), digestive contents were systematically recovered in the caecum, and in case of a watery consistency, in any other intestine segment, notably the colon. All samples except one caecal content (129) were preserved as aliquots at 4°C, -20°C and -70°C.

This study followed institutional guidelines and was approved by the scientific committee of IFCE (https://www.ifce.fr) that funded the project. All data are anonymous.

Standard search for entero-pathogens

Escherichia coli colonies were isolated on Eosin Methylene Blue plates and streaked on blood plates to see hemolysis, and *Salmonella* spp. on XLD plates (Thermo Scientific). *C. difficile* was detected on ChromID (BioMérieux) and *Clostridium perfringens* and *Paeniclostridium sordellii* on Columbia agar containing 5% sheep blood (BioMérieux), under anaerobiosis. Matrix-Assisted Laser Desorption/Ionization Time-Of-Flight (MALDI-TOF) Mass Spectrometry (MS) (MBT Smart mass spectrometer, Compass and FlexControl softwares, Bruker Daltonics) confirmed identification. Rotavirus was identified by PCR.

Systematic *C. difficile* search

Any digestive sample was directly streaked onto ChromID or CLO plates (BioMérieux) and incubated for 48h at 37°C under anaerobiosis. In case of a negative result, incubation for 3-7 days in BHI broth supplemented by cysteine (0.1%), yeast extract (5 g/L), taurocholate (1 g/L), D-cycloserine (250 µg/mL) and cefoxitin (8 µg/mL) (OXOID), allowed enrichment before ChromID plating. Suspicious, irregular and/or black colonies were screened by PCRs targeting *tpi*, *tcdA* and *tcdB* genes (24, 25). Identification was confirmed by MALDI-TOF MS. A single colony of each confirmed strain per positive sample was grown in BHI broth at 37°C under anaerobiosis and this culture preserved in glycerol 17% at -70°C except in two cases: two initially detected but not immediately preserved strains (non-toxigenic in animal #48 and toxigenic in #54) could not be recovered again.

C. difficile characterization

All preserved isolates (N=34 from n=25 Equidae) were characterized by multiplex PCR targeting: i) *tpi* gene, ii) wild-type or truncated toxin genes and iii) *lok* fragment characterizing non-

149 toxigenic strains, followed by capillary gel electrophoresis (26). For ribotyping, PCR profiles after
150 capillary gel-based electrophoresis (27) were analyzed with GeneMapper software (Thermo Fischer
151 Scientific, Villebon-sur-Yvette, France) and ribotypes assigned according to database WEBRIBO
152 (https://webribo.ages.at/). The first isolates were characterized by Multi Locus Sequence Typing
153 (28).
154
155 **Vancomycin or metronidazole susceptibility**
156   The Minimum Inhibitory Concentration (MIC) was determined using Etest (bioMérieux) on
157 Brucella blood agar (Becton Dickinson) supplemented with vitamin K1 (10 mg/L), hemin (5 mg/L)
158 and defibrinated horse blood (5%) (OXOID), after 24 h at 37°C.
159
160 **Toxin detection**
161   *C. difficile* toxins (TcdA and TcdB) and glutamate dehydrogenase (GDH) were detected in
162 digestive contents using an Enzymatic Immuno-Assay (Quick Check Complete, Alere), according to
163 manufacturer's instructions. Toxin detection was performed to diagnose CDI, as recommended in
164 Europe (3).
165
166 **Statistical analysis**
167   Pearson's Chi-squared test or Fisher's exact test when appropriate were used. Unknown
168 values were not imputed. Statistical significance: p-value <0.05.

Results

Necropsied Equidae

We studied all one hundred Equidae (except fetuses and stillborn foals) necropsied at ANSES in Normandy from 2019 to 2021, whatever clinical picture or death cause (Table S1). All animals except three originated from Normandy (Table S1). Most were females (67%) and <1-year-old animals (59%), notably foals (<6-months, 48%) (Table 1). They represented nine breeds, mostly Thoroughbred (42%), Trotters (31%) and Saddlebred horses (11%) (Table 1). Animal death causes included infections, non-infectious diseases and trauma. The two most prevalent causes were endo-enterotoxaemia (12%), an infection characterized by toxin-mediated inflammation (20) and infection by *Rhodococcus equi* (9%), common in foals (29).

C. difficile prevalence

Digestive contents were collected from i) the caecum (CAE) in all cases (n=100 animals) and ii) any other intestinal segment in case of watery consistency (CDL), possibly indicating diarrhea (n=30) (Table S1). All preserved digestive contents were tested for *C. difficile* presence, directly or after enrichment. Candidate colonies were screened by standard PCRs (24, 25) and confirmed by MALDI-TOF. Twenty-seven animals were positive for *C. difficile*, in their two samples in nine cases (Table S1). *C. difficile* presence in Equidae was independent of animal sex, age, breed, antibiotic treatment or hospitalization, but significantly correlated to signs of diarrhea or endo-enterotoxaemia (Table 1).

Pathogenicity

C. difficile isolates were preserved from twenty-five Equidae and named '*Cd*E' (Table 2). They form the first French library of *C. difficile* strains originating from Equidae, CloDifEqui (Table 2). Their pathogenicity profile was characterized by multiplex PCR (26). Non-toxigenic isolates (N=9) originated from seven animals, including six Thoroughbred, while toxigenic isolates (N=25), with one- to three-toxin profiles, originated from nineteen animals of six breeds (Table 2, Table S1). One animal (#29) was co-colonized by a non-toxigenic strain (CAE) and a toxigenic one (CDL). In all other animals with two isolates, both were either non-toxigenic or toxigenic with the same toxin profile (Table 2).

Ribotype diversity

CloDifEqui isolates were assigned to eleven ribotypes and four clades (1, 2, 4 and 5) (Table 2, Table S2) (27). Toxigenic isolates were of eight ribotypes: 126 and 078 (three-toxin profile: TcdA, TcdB, CDT and same *tcdC* truncation), 020, 005, 012, 181 and AI-53 (TcdA, TcdB), and 017 (TcdB). Non-toxigenic isolates belonged to ribotypes 009, 035 and 439 (Table 2, Table S2). The two most prevalent ribotypes (n=5 animals) were 017 (N=8 isolates) and 009 (N=7).

206　　　　　The two isolates of animal #29 belonged to ribotypes 439 and 005. For all other pairs
207　　originating from the same animal, both isolates shared the same ribotype (Table 2), suggesting their
208　　phylogenetic proximity and raising the possibility of a single clonal lineage colonizing animal gut.

Metronidazole or vancomycin susceptibility

All strains were susceptible to metronidazole and vancomycin (Table 2). The median MIC was 0.13 mg/L for metronidazole and 0.5 mg/L for vancomycin. The strains least susceptible to metronidazole (MIC=0.38 mg/L) or vancomycin (MIC=1.0 mg/L) were of ribotype 017 and 126 respectively. Most ribotype 017 strains were poorly susceptible to each antibiotic (Table 2).

Putative circulation

We cross-analyzed *C. difficile* ribotype and the corresponding Equidae location (Table S1, Figure 1, Table 2). In two premises between Caen and Pont-Audemer, few animals shared the same ribotype: i) two co-located Equidae necropsied within a few days in 2019 (#42 and #43) displayed toxigenic ribotype 020, and ii) four co-located Equidae, necropsied in 2019 (#5, #18 and #21) and 2020 (#61), carried ribotype 009. This suggests that a single clonal lineage could have colonized the co-located animals and that each premises could represent a transmission cluster.

CDI or asymptomatic carriage

We tested the presence of free TcdA and TcdB toxins in animal samples using a standard enzymatic immuno-assay. To check its specificity when applied to equine digestive contents rather than human stools, we included samples expected to be toxin-negative: i) all nine samples displaying a non-toxigenic *C. difficile* (Table 3) and ii) twenty-seven *C. difficile*-negative samples, including fifteen CDL and all samples from animals having endo-enterotoxaemia signs (Table S1, Table S3). No toxins could be detected in any of them, while GDH was positive in most of the former, as expected (Table S3, Table 3).

CDI was demonstrated by the detection of free toxins, together with GDH, in the two samples of four animals (#3, #32, #36 and #37), which displayed signs of diarrhea (CDL in the colon) and toxigenic *C. difficile* (ribotype 126 or 017) as the only detected entero-pathogen (Table 3, Figure 2). In the remaining sixteen Equidae with a toxigenic *C. difficile*, no free toxins could be detected in any sample (Table 3). Half of them consistently showed no diarrhea signs (#1, #2, #9, #12, #19, #85, #90 and #92). For all remaining animals except one (#57), diarrhea signs could have an independent cause: another entero-pathogen (*C. perfringens* in #43, with *P. sordellii* in #29 and #42 or with both *P. sordellii* and rotavirus in #53, and a parasite in #78) or the intestinal disease that caused death (acute folded colon in #54 and intestinal lymphoma in #81) (Table S1). Finally, sixteen Equidae were asymptomatic carriers of toxigenic *C. difficile,* while possibly having an independent diarrhea.

In the subpopulation displaying toxigenic *C. difficile*, animals with CDI and asymptomatic carriers were similar according to sex, age and non-pathognomonic signs of endo-enterotoxaemia or diarrhea, but they significantly differed according to breed and antibiotic treatment (Table 1). In contrast to a minority of asymptomatic carriers, all animals with CDI were Trotters and most had received antibiotics (Table S1, Table 1). This showed antibiotics to be a risk factor for CDI in the equine population.

Cause of animal death

We examined the cause of death of the four animals with CDI (Table S1). The fulminant death of animal #3, a 15-days-old foal, was unrelated to CDI and due to small intestine intussusception (Table S1). Animal #37 died from multiple (respiratory, musculoskeletal and digestive) infections (including endo-enterotoxaemia), probably due to *Klebsiella pneumoniae*, identified in lungs, lung abscess and hock muscle, and/or *C. difficile* (Table S1).

CDI was the only identified cause of the endo-enterotoxaemia and death of two animals, #32 and #36 (Table S1). Ribotype 017 was implicated in both cases and probably also in another death in 2018 (animal #0, which displayed signs of diarrhea, had received antibiotics and died from endo-enterotoxaemia) (Table 2, Table S1).

Discussion

C. difficile **prevalence and diversity**

C. difficile prevalence in French necropsied Equidae (27%) was unanticipated by many professionals, probably because its carriage was previously overlooked. Similar prevalence has been reported in European live horses: 4-33% in general and 5-63% for ill animals (9). The dead condition of animals apparently did not result in any major bias, probably because death causes were very diverse (Table S1).

C. difficile strains from Equidae belonged to diverse, already described ribotypes (9, 30). Of note, the historical epidemic ribotype 027 (1), described in horses (19), was not identified here, probably because of the changing *C. difficile* epidemiology and 027 prevalence decrease (31), notably in France (https://www.pasteur.fr/fr/file/44921/download) (30, 32). Most toxigenic ribotypes found in Equidae are very common in humans. Ribotypes 020, 078, 126 and 005 on the one hand and 017 and 020 on the other hand were among the Top10 and Top20 ribotypes found in patients in Europe in 2013-2014 (30). French Equidae therefore contributed to the dissemination of clinically important ribotypes, a preoccupying finding.

Most toxigenic ribotypes evidenced here are also common in animals (9, 11). Ribotypes 126, 020 and 012 had been found in farm animals (9). Ribotype 078, prevalent in animals, notably pigs (9), was rather uncommon here (N=1 isolate), but the phylogenetically close ribotype 126 (33) was represented (N=3) (Table 2). Of note, most toxigenic ribotypes identified here have been detected in diverse sources in Europe, molluscs (005, 012, 017, 020, 078, 126 and AI-53) (34), potatoes (35) and vegetables (012, 020, 078 and 126) (11). The noteworthy diversity and ubiquity of ribotypes identified here highlighted Equidae as important players in the dissemination, circulation and transmission of pathogenic *C. difficile*, a One Health issue.

The co-localization of Equidae sharing the same ribotype (Figure 1) suggested transmission between them, even though the full understanding of *C. difficile* transmission and phylogeny would require further genome-wide investigations. Regarding the co-located carriers of prevalent ribotype 009, described in Czech horses (36), one was a foal born in 2020 (#61), after the death of the others in 2019 (#5, #18, #21) (Table S1), indicating indirect transmission and spore persistence in the stud farm environment for months.

CDI and carriage

To reach a high standard diagnosis of CDI in necropsied Equidae with *post-mortem* signs of diarrhea, we combined the amplification of toxin genes, a method of high sensitivity and negative prediction value, to toxin detection, a method of high specificity and positive prediction value, as recommended in European hospitals (3). Toxin detection is uncommon in animals, notably Equidae (37-39), after *post-mortem* examination (40). Here, toxin detection demonstrated CDI in 4% of the population.

C. difficile carriage concerned 23% of animals. Equine signs of diarrhea or of endo-enterotoxaemia were significantly correlated to *C. difficile* presence (Table S1, Table 1) and

specifically carriage (p-value <0.005). Consistently, more *C. difficile* carriers (n=7/23) than *C. difficile*-negative animals (n=9/73) displayed another entero-pathogen (bacterium, virus, parasite), with a strong trend towards significance (p < 0.06). The most prevalent was *C. perfringens* in four *C. difficile* carriers (#29, #42, #43 and #53) and it was the most probable cause of the endo-enterotoxaemia and death of at least one of them (#43) (Table S1). The co-colonization or co-infection by *C. difficile* and *C. perfringens* have been described in horses (41, 42), dogs (43), pigs (44) and humans (45).

Our most complicated case was animal #29, which displayed three pathogen species and two *C. difficile* strains, toxigenic and non-toxigenic (Table S1, Table 2). Animal co-colonization by a toxigenic strain and a non-toxigenic one of ribotype 009 or 010 has been reported in companion animals (46). Non-toxigenic strains have been shown to prevent CDI or reduce its severity in animal models (47) and, for a ribotype 009 strain, also in commercial piglets (44).

All Equidae with CDI, in contrast to a few asymptomatic carriers of a toxigenic *C. difficile*, were Trotters (Table 1, Table S1). The condition of their microbiota could have promoted CDI. Indeed, most Trotters with CDI had been treated by antibiotics, favoring microbiota dysbiosis (Table 1, Table S1). On the contrary, none of the six Thoroughbred asymptomatically carrying a toxinogenic strain had received antibiotics. Moreover, the four Thoroughbred having received antibiotics carried non-toxinogenic strains of ribotype 009 (Table 1, Table S1), possible able to protect against a toxigenic strain (44, 47).

CDIs in Equidae had been caused by isolates of ribotypes 126 or 017, which were the CloDifEqui isolates least susceptible to vancomycin (MIC=1.0mg/L and 0.75mg/L respectively), and the latter were also poorly susceptible to metronidazole (MIC=0.19-0.38mg/L). This suggested that Equidae treatments against CDIs could select strains more difficult to eradicate and able to better persist.

The ribotypes virulent in Equidae differ by their reported prevalence. Ribotype 126 was the sixth most prevalent ribotype in European patients in 2013-2014 (30) and is also, together with its close relative ribotype 078, prevalent in production animals, notably pigs (9, 48). In Brittany, a French region close to Normandy, biogas plants using porcine, bovine and poultry manure almost exclusively provided ribotypes 078 and 126 (26). Ribotype 017, known as an emerging ribotype in Asia (49), is less frequent in Europe: it was the twelfth most prevalent ribotype in 2013-2014 (30). In France, its prevalence was low in 2011-2014 (30) and 2020 (https://www.pasteur.fr/fr/sante-publique/CNR/les-cnr/bacteries-anaerobies-botulisme/rapports-d-activite), and it was undetected in ten hospitals in 2022 (32). As ribotype 017 is also uncommon in animals (9, 49) and has not been detected in Brittany (26), its prevalence in French Equidae (5%) was noticeable. It was responsible for three CDI cases (#32, #36 and #37) in a small area (Figure 1) within six weeks (Table S1), raising the possibility of a local infection cluster. Ribotype 017, whose virulence was of concern, could be emerging in Equidae and/or in Normandy. Finally, French Equidae, as a reservoir for *C. difficile* clinically important ribotypes, would deserve careful monitoring in a One Health perspective.

References

1. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. 2016. *Clostridium difficile* infection. Nat Rev Dis Primers 2:16020.

2. Kordus SL, Thomas AK, Lacy DB. 2022. *Clostridioides difficile* toxins: mechanisms of action and antitoxin therapeutics. Nat Rev Microbiol 20:285-298.

3. Crobach MJ, Planche T, Eckert C, Barbut F, Terveer EM, Dekkers OM, Wilcox MH, Kuijper EJ. 2016. European Society of Clinical Microbiology and Infectious Diseases: update of the diagnostic guidance document for *Clostridium difficile* infection. Clin Microbiol Infect 22 Suppl 4:S63-81.

4. van Prehn J, Reigadas E, Vogelzang EH, Bouza E, Hristea A, Guery B, Krutova M, Noren T, Allerberger F, Coia JE, Goorhuis A, van Rossen TM, Ooijevaar RE, Burns K, Scharvik Olesen BR, Tschudin-Sutter S, Wilcox MH, Vehreschild M, Fitzpatrick F, Kuijper EJ, Guideline Committee of the European Study Group on Clostridioides d. 2021. European Society of Clinical Microbiology and Infectious Diseases: 2021 update on the treatment guidance document for *Clostridioides difficile* infection in adults. Clin Microbiol Infect 27 Suppl 2:S1-S21.

5. Frost LR, Cheng JKJ, Unnikrishnan M. 2021. *Clostridioides difficile* biofilms: A mechanism of persistence in the gut? PLoS Pathog 17:e1009348.

6. Poquet I, Saujet L, Canette A, Monot M, Mihajlovic J, Ghigo JM, Soutourina O, Briandet R, Martin-Verstraete I, Dupuy B. 2018. *Clostridium difficile* Biofilm: Remodeling Metabolism and Cell Surface to Build a Sparse and Heterogeneously Aggregated Architecture. Front Microbiol 9:2084.

7. Lim SC, Knight DR, Riley TV. 2020. *Clostridium difficile* and One Health. Clin Microbiol Infect 26:857-863.

8. Rodriguez C, Taminiau B, Van Broeck J, Delmee M, Daube G. 2016. *Clostridium difficile* in Food and Animals: A Comprehensive Review. Adv Exp Med Biol 932:65-92.

9. Rodriguez Diaz C, Seyboldt C, Rupnik M. 2018. Non-human *C. difficile* Reservoirs and Sources: Animals, Food, Environment. Advances in Experimental Medicine and Biology 1050:227-243.

10. Warriner K, Xu C, Habash M, Sultan S, Weese SJ. 2017. Dissemination of *Clostridium difficile* in food and the environment: Significant sources of *C. difficile* community-acquired infection? J Appl Microbiol 122:542-553.

11. Bolton D, Marcos P. 2023. The Environment, Farm Animals and Foods as Sources of *Clostridioides difficile* Infection in Humans. Foods 12:1094.

12. Weese JS. 2020. *Clostridium* (*Clostridioides*) *difficile* in animals. Journal of Veterinary Diagnostic Investigation: Official Publication of the American Association of Veterinary Laboratory Diagnosticians, Inc 32:213-221.

13. Janezic S, Zidaric V, Pardon B, Indra A, Kokotovic B, Blanco JL, Seyboldt C, Diaz CR, Poxton IR, Perreten V, Drigo I, Jiraskova A, Ocepek M, Weese JS, Songer JG, Wilcox MH, Rupnik M. 2014. International *Clostridium difficile* animal strain collection and large diversity of animal associated strains. BMC Microbiol 14:173.

14. Hensgens MPM, Keessen EC, Squire MM, Riley TV, Koene MGJ, de Boer E, Lipman LJA, Kuijper EJ, (ESGCD) ESoCMaIDSGfCd. 2012. *Clostridium difficile* infection in the community: a zoonotic disease? Clinical Microbiology and Infection: The Official Publication of the European Society of Clinical Microbiology and Infectious Diseases 18:635-645.

15. Knetsch CW, Connor TR, Mutreja A, van Dorp SM, Sanders IM, Browne HP, Harris D, Lipman L, Keessen EC, Corver J, Kuijper EJ, Lawley TD. 2014. Whole genome sequencing reveals potential spread of *Clostridium difficile* between humans and farm animals in the Netherlands, 2002 to 2011. Euro Surveill 19:20954.

16. Eyre DW, Cule ML, Wilson DJ, Griffiths D, Vaughan A, O'Connor L, Ip CLC, Golubchik T, Batty EM, Finney JM, Wyllie DH, Didelot X, Piazza P, Bowden R, Dingle KE, Harding RM, Crook DW, Wilcox MH, Peto TEA, Walker AS. 2013. Diverse sources of *C. difficile* infection identified on whole-genome sequencing. N Engl J Med 369:1195-205.

17. Baverud V. 2004. *Clostridium difficile* diarrhea: infection control in horses. Vet Clin North Am Equine Pract 20:615-30.

18. Diab SS, Songer G, Uzal FA. 2013. *Clostridium difficile* infection in horses: a review. Veterinary Microbiology 167:42-49.

19. Hain-Saunders NMR, Knight DR, Bruce M, Riley TV. 2022. *Clostridioides difficile* infection and One
Health: an equine perspective. Environmental Microbiology 24:985-997.

20. Uzal FA, Navarro MA, Asin J, Henderson EE. 2022. Clostridial Diseases of Horses: A Review. Vaccines
10:318.

21. Zhang K, Martinod S. 2014. *Clostridium difficile* Infection in Horses. Austin J Vet Sci & Anim Hus 1.

22. Mullen KR, Yasuda K, Divers TJ, Weese JS. 2018. Equine faecal microbiota transplant: Current
knowledge, proposed guidelines and future directions. Equine Vet Educ 30:151-160.

23. Tapprest J, Foucher N, Linster M, Laloy E, Cordonnier N, Amat JP, Hendrikx P. 2019. Resumeq: A
Novel Way of Monitoring Equine Diseases Through the Centralization of Necropsy Data. Front Vet Sci
6:135.

24. Lemee L, Dhalluin A, Testelin S, Mattrat MA, Maillard K, Lemeland JF, Pons JL. 2004. Multiplex PCR
targeting *tpi* (triose phosphate isomerase), *tcdA* (Toxin A), and *tcdB* (Toxin B) genes for toxigenic
culture of *Clostridium difficile*. J Clin Microbiol 42:5710-4.

25. Persson S, Jensen JN, Olsen KE. 2011. Multiplex PCR method for detection of *Clostridium difficile*
tcdA, *tcdB*, *cdtA*, and *cdtB* and internal in-frame deletion of *tcdC*. J Clin Microbiol 49:4299-300.

26. Le Maréchal C, Gateau C, Poezevara T, Couturier J, Rouxel S, Syed Zaidi R, Houard E, Pourcher A-M,
Denis M, Barbut F. 2020. Characterization of *Clostridioides difficile* strains isolated from manure and
digestate in five agricultural biogas plants. Anaerobe 62:102180.

27. Ferraris L, Couturier J, Eckert C, Delannoy J, Barbut F, Butel MJ, Aires J. 2019. Carriage and
colonization of *C. difficile* in preterm neonates: A longitudinal prospective study. PLoS One
14:e0212568.

28. Griffiths D, Fawley W, Kachrimanidou M, Bowden R, Crook DW, Fung R, Golubchik T, Harding RM,
Jeffery KJ, Jolley KA, Kirton R, Peto TE, Rees G, Stoesser N, Vaughan A, Walker AS, Young BC, Wilcox
M, Dingle KE. 2010. Multilocus sequence typing of *Clostridium difficile*. J Clin Microbiol 48:770-8.

29. Duquesne F, Houssin E, Sevin C, Duytschaever L, Tapprest J, Fretin D, Hebert L, Laugier C, Petry S.
2017. Development of a multilocus sequence typing scheme for *Rhodococcus equi*. Vet Microbiol
210:64-70.

30. Freeman J, Vernon J, Pilling S, Morris K, Nicholson S, Shearman S, Longshaw C, Wilcox MH, Group P-
ELSoARaPCdRS. 2018. The ClosER study: results from a three-year pan-European longitudinal
surveillance of antibiotic resistance among prevalent *Clostridium difficile* ribotypes, 2011-2014.
Clinical Microbiology and Infection: The Official Publication of the European Society of Clinical
Microbiology and Infectious Diseases 24:724-731.

31. Turner NA, Smith BA, Lewis SS. 2019. Novel and emerging sources of *Clostridioides difficile* infection.
PLoS Pathog 15:e1008125.

32. Le Monnier A, Candela T, Mizrahi A, Bille E, Bourgeois-Nicolaos N, Cattoir V, Farfour E, Grall I,
Lecointe D, Limelette A, Marcade G, Poilane I, Poupy P, Kansau I, Zahar J-R, Pilmis B, Hartmann C,
Kazhalawi A, Lambert-Bordes S, Bleunven S, Bedos J-P, Greder-Belan A, Rigaudea S, Lecuyer H,
Jousset A, Lebeaux D, Levy B, Rabate C, Collignon A, Batah J, Francois V, Sebbane G, Woerther P-L,
Loggia G, Michon J, Verdon R, Samba D, Méar J-B, Guillard T, Nguyen Y, Banisadr F, Delmer A,
Himberlin C, Diallo S, Furet I, Achouri B, Reksa A, Jouveshomme S, Menage E, Philippart F, et al. 2022.
One-day prevalence of asymptomatic carriage of toxigenic and non-toxigenic *Clostridioides difficile* in
10 French hospitals. Journal of Hospital Infection 129:65-74.

33. Knight DR, Kullin B, Androga GO, Barbut F, Eckert C, Johnson S, Spigaglia P, Tateda K, Tsai PJ, Riley TV.
2019. Evolutionary and Genomic Insights into *Clostridioides difficile* Sequence Type 11: a Diverse
Zoonotic and Antimicrobial-Resistant Lineage of Global One Health Importance. mBio 10.

34. Agnoletti F, Arcangeli G, Barbanti F, Barco L, Brunetta R, Cocchi M, Conedera G, D'Este L, Drigo I,
Spigaglia P, Mazzolini E. 2019. Survey, characterization and antimicrobial susceptibility of *Clostridium*
difficile from marine bivalve shellfish of North Adriatic Sea. International Journal of Food
Microbiology 298:74-80.

35. Tkalec V, Viprey V, Davis G, Janezic S, Sente B, Devos N, Wilcox M, Davies K, Rupnik M, consortium C-
C. 2022. *Clostridioides difficile* positivity rate and PCR ribotype distribution on retail potatoes in 12
European countries, January to June 2018. Euro Surveillance: Bulletin Europeen Sur Les Maladies
Transmissibles = European Communicable Disease Bulletin 27:2100417.

36. Kecerova Z, Cizek A, Nyc O, Krutova M. 2019. *Clostridium difficile* isolates derived from Czech horses
are resistant to enrofloxacin; cluster to clades 1 and 5 and ribotype 033 predominates. Anaerobe
56:17-21.
37. Arroyo LG, Staempfli H, Weese JS. 2007. Molecular analysis of *Clostridium difficile* isolates recovered
from horses with diarrhea. Veterinary Microbiology 120:179-183.
38. Magdesian KG, Leutenegger CM. 2011. Real-time PCR and typing of *Clostridium difficile* isolates
colonizing mare-foal pairs. Veterinary Journal (London, England: 1997) 190:119-123.
39. Ossiprandi MC, Buttrini M, Bottarelli E, Zerbini L. 2010. Preliminary molecular analysis of *Clostridium
difficile* isolates from healthy horses in northern Italy. Comparative Immunology, Microbiology and
Infectious Diseases 33:e25-29.
40. Zakia LS, Arroyo LG, Gomez DE, Boerlin P, Surette MG, Lillie BN. 2023. Postmortem examination of
equids >1-year-old with enterotyphlocolitis in Ontario: a retrospective descriptive study. Journal of
Veterinary Diagnostic Investigation 35:350-354.
41. Schoster A, Arroyo LG, Staempfli HR, Shewen PE, Weese JS. 2012. Presence and molecular
characterization of *Clostridium difficile* and *Clostridium perfringens* in intestinal compartments of
healthy horses. BMC veterinary research 8:94.
42. Uzal FA, Diab SS, Blanchard P, Moore J, Anthenill L, Shahriar F, Garcia JP, Songer JG. 2012. *Clostridium
perfringens* type C and *Clostridium difficile* co-infection in foals. Veterinary Microbiology 156:395-
402.
43. Diniz AN, Coura FM, Rupnik M, Adams V, Stent TL, Rood JI, de Oliveira CA, Jr., Lobato FCF, Silva ROS.
2018. The incidence of *Clostridioides difficile* and *Clostridium perfringens netF*-positive strains in
diarrheic dogs. Anaerobe 49:58-62.
44. Oliveira Junior CA, Silva ROS, Lage AP, Coura FM, Ramos CP, Alfieri AA, Guedes RMC, Lobato FCF.
2019. Non-toxigenic strain of *Clostridioides difficile* Z31 reduces the occurrence of *C. difficile* infection
(CDI) in one-day-old piglets on a commercial pig farm. Vet Microbiol 231:1-6.
45. Forero AJ, Muñoz M, Camargo M, Soto-De León SC, Ríos-Chaparro DI, Birchenall C, Pinilla D, Pardo
JM, Josa DF, Patarroyo MA, Ramírez JD. 2019. High frequency of toxigenic *Clostridium difficile* and
Clostridium perfringens coinfection among diarrheic patients at health care facility-onset (HCFO) and
community-onset (CO) centers in Bogotá, Colombia. Gut Pathogens 11:27.
46. Alves F, Castro R, Pinto M, Nunes A, Pomba C, Oliveira M, Silveira L, Gomes JP, Oleastro M. 2023.
Molecular epidemiology of *Clostridioides difficile* in companion animals: Genetic overlap with human
strains and public health concerns. Frontiers in Public Health 10:1070258.
47. Gerding DN, Sambol SP, Johnson S. 2018. Non-toxigenic *Clostridioides* (Formerly *Clostridium*) *difficile*
for Prevention of *C. difficile* Infection: From Bench to Bedside Back to Bench and Back to Bedside.
Front Microbiol 9:1700.
48. Schneeberg A, Neubauer H, Schmoock G, Baier S, Harlizius J, Nienhoff H, Brase K, Zimmermann S,
Seyboldt C. 2013. *Clostridium difficile* Genotypes in Piglet Populations in Germany. Journal of Clinical
Microbiology 51:3796-3803.
49. Imwattana K, Knight DR, Kullin B, Collins DA, Putsathit P, Kiratisin P, Riley TV. 2019. *Clostridium
difficile* ribotype 017 - characterization, evolution and epidemiology of the dominant strain in Asia.
Emerg Microbes Infect 8:796-807.

Figure Legends

Figure 1. Geographical location of *C. difficile*-positive Equidae in Normandy and PCR-ribotypes of their strains

Each *C. difficile*-positive Equidae is located on a map of Normandy according to the latitude and longitude coordinates of its premises (Table S1). Different symbols are used depending on *C. difficile* toxin profile: squares for a three-toxin profile (TcdA TcdB and CDT, for ribotypes 078 and 126), circles for two-toxin profile (TcdA and TcdB, for ribotypes 005, 012, 020, 181 and AI-53), triangles for a TcdB profile (for ribotype 017) and finally diamonds for non-toxigenic strains (ribotypes: 009, 035 and 439) (Table 2). *C. difficile* ribotype (Table 2) is indicated in a box, and if several animals of a single premises harboured the same ribotype (Table S1, Table 2), their number is shown between brackets. *C. difficile* pathogenicity is also indicated by a colour code: non-toxinogenic in green, and toxigenic in either orange or red depending on whether *C. difficile* was asymptomatically carried or responsible for a CDI.

Figure 2. Diagnosis of CDI or carriage in *C. difficile*-positive Equidae

All *C. difficile*-positive animals are shown as a wedge of a pie chart. For each positive animal, the pathogenicity and ribotype of *C. difficile*, and the result of diagnosis, are indicated in four concentric rings.

First (core): *C. difficile* pathogenicity and toxin profile (see Table 2). Non-toxigenic ('Non-Tox') and toxigenic strains ('Tox') are indicated on a green or orange background, respectively. 'Both' is indicated when a non-toxigenic strain and a toxigenic strain were co-colonizing the same animal (#29, bi-coloured wedge). The toxin profile of toxigenic strains is specified using the following abbreviations: A for toxin A (TcdA), B for toxin B (TcdB), CDT for binary toxin and '?' when the presence of *cdt* genes has not been tested.

Second: *C. difficile* ribotype (according to WEBRIBO database) (Table 2)

Third: Animal identification number

Fourth (outer ring): Diagnosis (Table 3). Infection (CDI), asymptomatic carriage of a toxigenic strain or carriage of a non-toxigenic strain are on a red, orange or green background respectively.

Table 1. Equine population: data summary

	C. difficile presence			C. difficile pathogenicity				Diagnosis (animals with a toxigenic C. difficile)		
	No	Yes	P-value	Non-toxigenic	Toxigenic	Both	P-value	Carriage	Infection	P-value
Year										
2019	32	15	0.60	3	11	1	0.91	8	4	0.30
2020	28	8		3	5	0		5	0	
2021	13	4		1	3	0		3	0	
Animal data										
Sex										
Female	46	21	0.44	7	13	1	0.34	11	3	1.00
Male, gelding	26	6		0	6	0		5	1	
Unknown	1	0								
Age										
Foal	33	15	0.55	6	9	0	0.42	6	3	0.79
Yearling	10	1		0	1	0		1	0	
Young	5	2		0	2	0		2	0	
Adult	25	9		1	7	1		7	1	
Race										
Thoroughbred	30	12	0.87	6	6	0	0.15	6	0	**0.04**
Trotter	23	8		1	6	1		3	4	
Saddlebred horse	9	2		0	2	0		2	0	
Other (6 breeds)	9	5		0	5	0		5	0	
Unknown	2	0								
• Thoroughbred versus Other	30 / 43	12 / 15	0.82	6 / 1	6 / 13	0 / 1	**0.02**	6 / 10	0 / 4	0.27
• Trotter versus Other	23 / 50	8 / 19	1.0	1 / 6	6 / 13	1 / 0	0.29	3 / 13	4 / 0	**0.007**
***Post-mortem* signs of**										
Diarrhea										
One sample per animal: CAE	59	11	**0.0001**	3	8	0	1.0	8	0	0.12
Two samples per animal: CAE + CDL	14	16		4	11	1		8	4	
Endo/enterotoxⴰmia										
Yes	7	13	**0.0001**	3	9	1	1.0	7	3	0.58
No	65	14		4	10	0		9	1	
Unknown	1	0								
Risk factors										
Previous antibiotic treatment										
Yes	24	11	0.83	5	6	0	0.31	3	3	**0.014**
No	43	14		2	11	1		12	0	
Unknown	6	2		0	2	0		1	1	
Previous hospitalization										
Yes	22	11	0.57	4	7	0	0.65	7	0	0.25
No	49	16		3	12	1		9	4	
Unknown	2	0		3	12	1		9	4	

Table 1 Legend

The equine population composition is shown. The number of necropsied animals is indicated according to year or main animal data: i) individual

data (sex, age, breed), ii) *post-mortem* observations (signs of diarrhea, endo-enterotoxaemia), iii) *ante-mortem* data (previous antibiotic

treatment or hospitalization) (Table S1). The equine population composition was statistically analyzed according to: i) *C. difficile* presence or not

(all animals), ii) *C. difficile* pathogenicity or not (*C. difficile*-positive animals) and finally iii) the diagnosis of CDI or asymptomatic carriage (for

positive animals with a toxigenic *C. difficile*).

Table 2. *C. difficile* strains isolated from necropsied Equidae: molecular characterization

Strain Name	Equidae n°	Intestinal content	Method Enrichment	tpi[a]	tcdA[b]	tcdB[a]	lok[c]	tcdC[d] deletion	cdt[e]	Toxin profile[f]	Ribotype[g] (WEBRIBO)	MLST[h]	MIC Vancomycin (mg/L)[i]	MIC Metronidazole (mg/L)[i]
2019-2021														
Cd E-3	3	CAE	-							TcdA TcdB CDT	126	ST11	**1.00**	0.09
Cd E-3-CDL		CDL	-	+	+	+	-	-39	+		126	ST11	**1.00**	0.06
Cd E-19	19	CAE	-							TcdA TcdB CDT			0.38	0.05
Cd E-53	53	CAE§	+	+	+	+	-	-39	+		078	nd	0.50	0.09
Cd E-2	2	CAE	+								005	nd	**0.75**	0.13
Cd E-90	90	CAE	+	+	+	+	-	0	T		005	nd	0.25	0.05
Cd E-29-CDL	29*	CDL	-									nd	**0.75**	0.13
Cd E-9	9	CAE	-	+	+	+	-	0	T		012	ST54	0.50	0.05
Cd E-85	85	CAE	-								012	nd	0.50	0.05
Cd E-42	42	CAE	+							TcdA TcdB	020	nd	**0.75**	0.19
Cd E-42-CDL		CDL	+									nd	0.50	0.13
Cd E-43	43	CAE	-	+	+	+	-	0	T		020	ST2	0.50	0.19
Cd E-43-CDL		CDL	-									ST2	0.50	**0.25**
Cd E-57	57	CAE§	-									nd	0.50	0.13
Cd E-78	78	CAE§	+									nd	0.50	0.19
Cd E-81-CDL	81	CDL§	+	+	+	+	-	0	T		181	nd	0.38	0.19
Cd E-1	1	CAE	+	+	+	+	-	0	T		AI-53	nd	0.25	0.13
Cd E-12	12	CAE	-								017	ST37	0.50	0.09
Cd E-32	32	CAE	-									ST37	**0.75**	**0.38**
Cd E-32-CDL		CDL	-									ST37	**0.75**	0.19
Cd E-36	36	CAE	-	+	del	+	-	0	-	TcdB	017	ST37	**0.75**	**0.25**
Cd E-36-CDL		CDL	-									ST37	**0.75**	0.19
Cd E-37	37	CAE	-									nd	**0.75**	**0.25**
Cd E-37-CDL		CDL	-									ST37	**0.75**	**0.38**
Cd E-92	92	CAE	-									nd	0.50	**0.25**
Cd E-5	5	CAE	-								009	ST3	0.50	0.09
Cd E-18	18	CAE	-									ST3	**0.75**	0.13
Cd E-18-CDL		CDL	-									ST3	0.50	0.09
Cd E-21	21	CAE	-	+	-	-	+	-	-	Non-toxigenic	009	nd	0.13	0.09
Cd E-60	60	CAE	-									nd	**0.75**	0.09
Cd E-60-CDL		CDL	-									nd	0.50	0.06
Cd E-61-CDL	61	CDL§	+									nd	**0.75**	0.06
Cd E-91	91	CAE	-	+	-	-	+	-	-		035	nd	0.13	0.13
Cd E-29	29*	CAE	+	+	-	-	+	-	–		439	nd	0.50	0.05
2018														
CdE-0-CDL	0#	CDL	+	+	del	+	-	0	-	TcdB	017	nd	0.38	0.032

Table 2 Legend.

All preserved strains are listed and their origin (animal and sample) indicated. All data about their

molecular characterization: multiplex PCR results, toxin profile, PCR-ribotype and Multi Locus

Sequence Type (MLST) are shown. Their level of susceptibility to vancomycin or metronidazole

(MICs) is also indicated.

[a] *tpi, tcdB*: PCR-fragment of wild-type size ('+') or no amplification ('-')

[b] *tcdA*: PCR-fragment of wild-type size ('+') or of shorter size indicative of a deletion ('del') or no
amplification ('-')

[c] lok: PCR-fragment of 117 bp ('+', i. e. PaLoc deletion, non-toxigenic strain) or no amplification ('-')

[d] *tcdC*: deletion of 39 bp ('-39') or no deletion ('0', i. e. PCR-fragment of wild-type size) or no
amplification ('-')

[e] *cdt*: PCR-fragment of wild-type size ('+') or truncated PCR-fragment ('T', i. e. of a pseudogene) or
no amplification ('-')

[f] Toxin profile: the combination of toxins encoded on the strain genome (according to the PCR-
established genotype)

[g] PCR-ribotype assigned according to the WEBRIBO database

[h] MLST: Multi-Locus Sequence Type; nd: not determined

[i] MIC: Minimum Inhibitory concentration; the two highest MIC values are in bold.

[*] animal co-colonized by two different strains (one recovered from CAE and one from CDL)

[¤] animal necropsied at the end of 2018 (pilot study)

[§] the single sample that provided a *C. difficile* strain (out of the two recovered ones)

Table 3. Diagnosis of CDI or carriage

Equidae n°	Intestinal sample	C. difficile Strain[a]	Pathogenicity	Toxin profile[b]	Ribotype	GDH[d]	Free Toxins[d] (TcdA TcdB)	Diagnosis[d]
3	CAE	Cd E-3	Toxigenic	TcdA TcdB CDT	126	+	+	CDI
	CDL	Cd E-3-CDL	Toxigenic	TcdA TcdB CDT	126	+	+	
32	CAE	Cd E-32	Toxigenic	TcdB	017	+	+ (very weak)	
	CDL	Cd E-32-CDL	Toxigenic	TcdB	017	+	+ (very weak)	
36	CAE	Cd E-36	Toxigenic	TcdB	017	+	+	
	CDL	Cd E-36-CDL	Toxigenic	TcdB	017	+	+	
37	CAE	Cd E-37	Toxigenic	TcdB	017	+	+	
	CDL	Cd E-37-CDL	Toxigenic	TcdB	017	+	+	
19	CAE	Cd E-19	Toxigenic	TcdA TcdB CDT	126	-	-	Asymptomatic carriage
53	CAE	Cd E-53	Toxigenic	TcdA TcdB CDT	078	-	-	
	CDL	none	NA		NA	-	-	
54	CAE	none	NA		NA	-	-	
	CDL	not preserved	Toxigenic	TcdA TcdB ?	ND	-	-	
2	CAE	Cd E-2	Toxigenic	TcdA TcdB	005	+ (very weak)	-	
29*	CAE	Cd E-29	Non-Toxigenic	None	439	+ (very weak)	-	
	CDL	Cd E-29-CDL	Toxigenic	TcdA TcdB	005	+ (very weak)	-	
90	CAE	Cd E-90	Toxigenic	TcdA TcdB	005	-	-	
9	CAE	Cd E-9	Toxigenic	TcdA TcdB	012	+	-	
85	CAE	Cd E-85	Toxigenic	TcdA TcdB	012	+	-	
42	CAE	Cd E-42	Toxigenic	TcdA TcdB	020	-	-	
	CDL	Cd E-42-CDL	Toxigenic	TcdA TcdB	020	-	-	
43	CAE	Cd E-43	Toxigenic	TcdA TcdB	020	+	-	
	CDL	Cd E-43-CDL	Toxigenic	TcdA TcdB	020	+	-	
57	CAE	Cd E-57	Toxigenic	TcdA TcdB	020	-	-	
	CDL	none	NA		NA	-	-	
78	CAE	Cd E-78	Toxigenic	TcdA TcdB	020	-	-	
	CDL	none	NA		NA	-	-	
81	CAE	none	NA		NA	-	-	
	CDL	Cd E-81-CDL	Toxigenic	TcdA TcdB	181	-	-	
1	CAE	Cd E-1	Toxigenic	TcdA TcdB	AI-53	-	-	
12	CAE	Cd E-12	Toxigenic	TcdB	017	-	-	
92	CAE	Cd E-92	Toxigenic	TcdB	017	-	- +	
5	CAE	Cd E-5	Non-toxigenic		009	+	-	Carriage
18	CAE	Cd E-18	Non-toxigenic		009	+	-	
	CDL	Cd E-18-CDL	Non-toxigenic		009	+	-	
21	CAE	Cd E-21	Non-toxigenic		009	-	-	
60	CAE	Cd E-60	Non-toxigenic		009	+	-	
	CDL	Cd E-60-CDL	Non-toxigenic		009	+	-	
61	CAE	none	NA		NA	-	-	
	CDL	Cd E-61-CDL	Non-toxigenic		009	+	-	
91	CAE	Cd E-91	Non-toxigenic		035	+	-	
48	CAE	none	NA		NA	-	-	
	CDL	not preserved	Non-toxigenic		ND	-	-	

Table 3 Legend.

All data about *C. difficile*-positive Equidae and results are indicated: i) signs of putative diarrhea as

inferred from the *post-mortem* observed watery consistency of intestinal sample (CDL, in bold), ii)

presence of a *C. difficile* strain and its pathogenicity (toxigenic [in bold] or non-toxigenic strain) and iii)

free toxin detection. Animals are grouped according to diagnosis: CDI, asymptomatic carriage of a

toxigenic strain or carriage of a non-toxigenic strain.

ª Strain: preserved strain name or 'not preserved' (a strain that was initially detected but could not be

recovered) or 'none' (no detectable *C. difficile*)

ᵇ Toxin profile, according to the PCR-established genotype of the strain. In one case, '?' is indicated as

no PCR targeting *cdt* genes has been performed.

* co-colonized animal

NA: not applicable

ND: not determined

584 **Supporting information**
585
586

Table S1. Animal data

The registered data of all necropsied Equidae are shown.

Table S2. Sequence Type (MLST) of CloDifEqui strains

The Sequence Type (MLST) of the first isolated strains was characterized.

Table S3. Toxin detection in *C. difficile*-negative samples

Toxin detection was performed in samples expected to give a negative result as a control for assay
specificity (also see Table 3). The samples from *C. difficile*-negative animals that have been assayed
are listed here.

Statements

Funding

This work was funded by IFCE (Institut Français du Cheval et de l'Equitation, France, https://www.ifce.fr, Research project 2019-009-Rech CloDifEqui), ANSES and INRAE.

For the purpose of Open Access, a CC-BY public copyright licence has been applied by the authors to the present document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising from this submission.

Authorship contribution statement

SP and IP designed the whole project, submitted it to IFCE for funding. JT, NF and MBR performed animal necropsies and intestinal content sampling, registered and analyzed *post-mortem* data. JT established the cause of animal death. KM performed bacteriological and molecular analyses to identify pathogens. SP, KM, FD, SK, LB and IP searched and isolated *C. difficile* strains and performed the primary molecular characterization of strains. FB performed final molecular characterization and free toxin detection. SP and IP analyzed all data. IP wrote the manuscript and SP, JT, FB, FD, SK and IP edited it. All authors read the manuscript and agreed to submit it.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Data availability

All data are available in the manuscript.

Acknowledgements

We deeply thank Lydia Baudet (ANSES, Normandy Laboratory for Animal Health, Physiopathology and Epidemiology of Equine Diseases Unit, Goustranville, France), Rabab Syed Zaidi and Lina Ma (CNR *C. difficile*, Hôpital Saint Antoine, Paris, France) for excellent technical assistance.

Figure 1

Figure 2