A comparison of microglial morphological complexity analysis in adult mouse brain samples using 2-dimensional and 3-dimensional image analysis tools

Murray, C.J.[1], Tunderman, E.D.[1,2], Vecchiarelli, H.A.[1], González Ibáñez, F.[1,3,4], Tremblay, M-È[1,3,4,5,6,7].

1. Division of Medical Sciences, University of Victoria, Victoria, BC., Canada.
2. Department of Translational Neuroscience – Brain Center, University Medical Center Utrecht, Utrecht University
3. Axe Neurosciences, Centre de Recherche du CHU de Québec-Université Laval, City, QC, Canada.
4. Départment de Médecine Moléculaire, Université Laval, Québec City, QC, Canada
5. Neurology and Neurosurgery Department, McGill University, Montréal, QC, Canada.
6. Department of Biochemistry and Molecular Biology, University of British Columbia, Vancouver, BC, Canada.
7. Centre for Advanced Materials and Related Technology (CAMTEC), University of Victoria, BC, Canada.

Main Points:

- The 3D microglial morphology analysis method indicates maintained and novel significant differences between groups when compared to the 2D analysis method.

- The data suggest that the 3D analysis method may be better able to detect certain aspects of microglial morphology (i.e., soma size) compared to the 2D analysis method.

- Although somewhat distinct, overall conclusions with respect to microglial adaptations to stimuli are similar using the 2D and 3D analysis methods.

- Using a nested statistical design alters statistical outcomes in a way that more accurately takes into account individual animal variability compared to non-nested approaches.

Abstract

Characterizing cell morphology has been an essential aspect of neuroscience for over a century to provide essential insights into cellular function and dysfunction. Microglia, the resident innate immune cells of the central nervous system, undergo drastic changes in morphology in response to various stimuli, with many classifications proposed in recent years. Increased availability of advanced analysis software to study microglial morphology represents a step forward in the field. However, whether the use of advanced analysis tools provides equivalent or varied outcomes remains undetermined. This study re-analyzed raw data, previously processed using a standard 2D microglial morphology analysis method, using 3D analysis methods. The published article observed significant changes in microglial morphology in the mouse ventral hippocampus after administration of a ketogenic diet and exposure to repeated social defeat stress in young adult male

mice. Overall, we observed different statistical outcomes in the 3D dataset compared to the previously published 2D results, with both maintained and new findings. This may indicate that the 3D analysis method is better able to capture minute changes in morphology. However, overall conclusions on microglial morphology changes remain consistent between methods. Lastly, we highlight the difference between a more mainstream statistical design and a nested design, which takes into account individual cell and animal variability. Overall, we highlight and discuss the difference between 2D and 3D microglia morphology analysis and explore the contribution of individual cell and animal variability to statistical outcomes.

Keywords: Microglial morphology, Image analysis tools, 2D, 3D, Comparative methodology, Mouse, Hippocampus, Stress, Ketogenic diet.

1 **Introduction**

Microglia, the resident innate immune cells of the central nervous system (CNS), are critical to
maintain CNS homeostasis and perform immune surveillance (Kreutzberg, 1996; Šimončičová et
al., 2022). In healthy conditions, microglia constantly communicate with neurons and other cells
of the CNS, promoting neuronal and glial cell health by clearing debris, pruning synapses,
modulating inflammation, facilitating neuronal connectivity and plasticity, and neuronal/glial
survival (Colonna & Butovsky, 2017; Gao et al., 2023). These essential functions are generally
accompanied by changes in microglial morphology, with healthy adult grey matter microglia often
exhibiting small somas with highly motile, ramified processes (i.e., surveillant morphotype) that
enable them to interact dynamically with their microenvironment, as observed in adult mouse
neocortex (Nimmerjahn et al., 2005; Tremblay et al., 2010). However, microglia in white matter
or early life stages can display different morphologies in rodent models and humans, reflecting
their distinct environmentally-induced functional states (Ambrose et al., 2020; Paolicelli et al.,
2022; Tremblay, 2020; Vidal-Itriago et al., 2022). Upon detecting pathogens or signs of damage
and debris, microglia can shift toward a reactive pro-inflammatory state, triggered by pathogen-
associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs) via
pattern recognition receptors (PRRs), like Toll-like receptors (TLRs) (Colonna & Butovsky, 2017).
In response to such adverse signals, microglia undergo a variety of morphological changes, often
retracting and thickening their processes, presenting somatic swelling, and sometimes becoming

hyper-ramified. Indeed, the existence of mixed morphological states that are distinct from the

surveillant morphotype of homeostatic microglia in health and disease and the diversity within

these populations is often overlooked. In line with this, microglia reactivity and morphological

changes are not binary, but exist as a spectrum of morphologies, each signifying different

microglial states that often reflect and shape the local environment they inhabit (Fig. 1).

Understanding this diversity is important. Even though morphology is not a direct measure of

function, there is substantial evidence linking specific microglial morphological states with distinct

transcriptomic profiles, for instance, suggesting that these morphological changes have significant

functional implications (Gyoneva et al., 2019; Paolicelli et al., 2022).

For example, upon detecting PAMPs and/or DAMPs caused by infection, injury, apoptotic

cells and/or cellular debris, microglia can retract their branches, undergo somatic swelling, and

adopt a less complex structure, which is often associated with an increased phagocytic activity

(Colonna & Butovsky, 2017; B. M. Davis et al., 2017, 2017; Doorn, Moors, et al., 2014; Ginhoux

et al., 2010; Morrison et al., 2017; Savage et al., 2019). In more severe cases, such as traumatic

brain injury, neurodegenerative disease, or chronic inflammation, microglia can further retract

their branches and adopt an amoeboid morphology, which is associated with enhanced migration

efficiency to sites of damage and phagocytic activity in the affected areas (Doorn, Goudriaan, et

al., 2014; Wicks et al., 2022). This ameboid microglia morphology is frequently observed in cases

of severe or chronic inflammation, but also during healthy states, such as during development

when microglial migration and phagocytosis are prevalent (Badanjak et al., 2021; Doorn, Moors,

et al., 2014; Perez-Pouchoulen et al., 2015; VanRyzin et al., 2019). Another distinct morphology

that inhabits the far-end of the spectrum is the hyper-ramified microglia, which has been observed

to be more prominent in stress resilient mice, indicating a potential link between this morphology

and an adaptive response to stress (Fujikawa & Jinno, 2022). In addition, there is diversity within

these contexts as well, with many intermediate or more polarized forms also being present, such

as bushy microglia—characterized by a large soma and surrounded by short, stubby branches—

within the same populations (Wicks et al., 2022). For a more extensive list of microglia

morphologies, please refer to Table 1.

In order to quantify these distinct microglial morphotypes, a suite of parameters has been

developing since Pío del Río Hortega first discovered microglia in 1919. Some of the most

commonly used parameters are soma area/volume, convex hull, circularity, branch length, Sholl

analysis, and more (Leyh et al., 2021). Although each descriptor can give useful information as to

how microglia are responding to any given stimuli, combinations of parameters in relation to one

another can give increased insight into individual morphotypes. For example, the morphology

index provides a readout of overall cell shape based on the ratio between soma area and

arborization area (i.e., the larger the value, the greater the soma size in relation to branching area),

whereas the Schoenen Ramification Index provides a readout of the overall branching complexity

of the cell, represented by the ratio between the maximum number of Sholl intersections and the

number of primary branches of the cell (Schoenen, 1982; Tremblay et al., 2012).

Given the association between microglial morphology and environmental pressures and/or

stimuli, microglial morphology analysis is a crucial tool for investigating their functional state in

CNS pathology. Techniques commonly used to assess microglial morphology often involve either

two-dimensional (2D) maximum intensity projection analysis (Karperien et al., 2013; Kozlowski

& Weimer, 2012; Torres-Platas et al., 2014; Verdonk et al., 2016) or more advanced three-

dimensional (3D) reconstruction analysis methods (e.g., Neurolucida or Imaris software) (Erny et

al., 2015; Perego et al., 2013). However, it remains unclear whether these differing approaches

yield comparable results and whether the conclusions drawn from each method are equally reliable

and accurate. The potential discrepancies between these methods highlight a broader, increasing

issue within the field of neuroscience: the lack of standardization in data generation, collection,

and analysis, which significantly contributes to reproducibility challenges and translational

failures (Green et al., 2022).

To this end, using Imaris 10.0.2, this study re-analyzed raw data in 3D from a previously

published study from our group that revealed significant changes in 2D microglia morphology in

the mouse ventral hippocampus after administration of a ketogenic diet (KD) or control diet (CD)

and exposure to repeated social defeat stress in young adult male mice. The original analysis

focused on comparing key morphological parameters, including microglial soma area, arborization

area, and the morphology index, across different experimental groups. These groups comprised

stress-susceptible (SUS) and stress-resistant (RES) mice, further categorized based on their diet

(KD *vs* CD). The 2D analysis highlighted distinct changes in microglial morphology, such as

differences in soma size and branching complexity. By extending this analysis into 3D, using the

same dataset, we aim to provide a comprehensive understanding of the differences between 2D

and 3D morphology analysis methods.

Methods

The raw data utilized in this methods manuscript were obtained from those published previously (González Ibáñez et al., 2023). Therefore, we only outline aspects of the method section necessary to contextualize these results.

Animals

All animal experiments were approved by the Université Laval's animal ethics committee (Comité de protection des animaux de L'Université Laval), in accordance with the guidelines set forth by the Canadian Council on Animal Care. For in depth details concerning animal experiments and the protocols used for the ketogenic diet, social defeat paradigm, social interaction test to determine resistant *vs* susceptible mice to stress, euthanasia, and tissue preparation, please refer to González Ibáñez et al., (2023). Briefly, 7–8-week-old male C57BL/6J mice (Jackson Laboratories) were perfused with phosphate-buffered saline (PBS; 50 mM pH 7.4) followed by a mixture of 4% paraformaldehyde and 0.2% glutaraldehyde in phosphate buffer (PB; 100 mM, pH 7.4) (González Ibáñez et al., 2023).

Fluorescence Staining and Confocal Imaging

For a detailed description of the methods used for immunofluorescence staining and confocal imaging, please refer to González Ibáñez et al., (2023). Briefly, 50 μm coronal sections were incubated with sodium citrate at 70 ºC, 0.1% $NaBH_4$, and blocking buffer before overnight incubation at 4 ºC with mouse anti-ionized calcium-binding adapter molecule 1 (IBA1; 1:150) and rabbit anti-transmembrane protein 119 (TMEM119; 1:300) primary antibodies. The following day, the sections were incubated with corresponding fluorescent secondary antibodies for 90 min and then with DAPI (1:20k) for 5 min. The sections were then mounted and coverslipped. Microglia (IBA1[+]/TMEM119[+]) (n=3–5 mice/group; 32–94 microglia/group) were then imaged at 40x with a Quorum WaveFX spinning disc confocal microscope (Quorum Technologies, Guelph, ON, Canada) with an ORCA-R2 camera (512 x 512 pixels; Hamamatsu Photonics, Hamamatsu, Japan). Z-stack images were taken with a Z-interval of 0.33 μm.

Microglial Morphology

This study utilized confocal images from a previously published paper from our research group— González Ibáñez et al., (2023)—to reanalyze previous 2D data with the 3D morphological analysis

software, Imaris (v10.0.2 Oxford Instruments). All microglial analysis was conducted blind to experimental conditions.

The following protocol was utilized to reconstruct microglia within the Imaris workspace for morphological characterization:

1. Raw czi. image files were imported into the Imaris 'Arena' and converted to the ims. file format using the built in Imaris file converter feature.

2. Once the image was selected, a new surface was created and the 'Skip Automatic Creation, Edit Manually' button was selected for manual reconstruction of the soma.

   a. Under the 'Contour' tab, the 'Time' drawing mode with an Insert Vertex every 300 ms was used to trace the soma (DAPI$^+$/IBA1$^+$). The perimeter of the soma was traced every 2–5 slices throughout the Z distance of the soma, which proved to accurately represent the volume of the soma qualitatively.

   b. The 'Create Surface' button was then selected, which allows Imaris to interpolate the slices not traced based off of the slices that were traced, completing the soma reconstruction. (Fig. 2B)

3. Next, we added another object using the 'Add New Filaments' selection for reconstruction of the microglial branches.

4. We then selected the 'Skip Automatic Creation, Edit Manually' button, opened the 'Edit' tab and selected 'Import Soma…' under the 'Process Filaments' section.

   a. Our manual soma reconstruction object was selected.

5. We then opened the 'Creation' tab, checked the 'Keep Data' box and proceeded to recompute the filaments.

   a. The 'Tree Autopath Algorithm' was used with default creation parameters.

   b. The source channel was selected and the 'Use Existing Segment' option was selected under the 'Starting Points' heading.

   c. Soma model was calculated and subsequently used as the starting point for filament tracing.

   d. The thinnest filament diameter was set at 1 μm.

   e. The seed point threshold was manually adjusted so that all visible branches extending from the soma contained at least one seed point at the base and terminal end. The 'Classify Seed Points' option was selected.

f. For seed points classification, a portion of seed points were manually selected or discarded before selecting 'Train and Predict'—an option within Imaris that uses machine learning to select or discard seed points based off of your previous selections.

g. No 'Terminal Segment Postfilter' was selected.

h. Filaments were then inspected and manually edited if needed. (Fig. 2C)

6. We then utilized the 'Filament Sholl Analysis24' plugin, acquired from Ironhorse1618 (2023): https://github.com/Ironhorse1618/Python3.7-Imaris-XTensions.

a. Sholl Sphere Radius was set to 5 µm.

b. 'Create Convex Hull surface' was selected. (Fig. 2D)

7. Finally, under the statistics tab, the volume of the convex hull, volume of the soma, total Sholl intersections and critical radius were extracted.

Although many pre-processing options exist in Imaris, we chose not to use these in order to keep the images as similar as possible to those used for the 2D morphology analysis with the aim to identify if the 3D aspect of this analysis alters results.

For morphology, we analyzed soma volume (μm^3), convex hull volume (μm^3), and morphology index (soma volume/convex hull volume) for the 3D analysis method, which respectively compares with soma area (μm^2), arborization area (μm^2), and morphology index (soma area/arborization area) quantified using the 2D analysis method performed by González Ibáñez et al., (2023). In addition, we quantified the total number of Sholl intersections at 5 µm radius intervals and the critical radius (radius with the most Sholl intersections) for microglia using the 3D analysis method.

Statistics

Statistical testing was performed in an identical manner as in González Ibáñez et al., (2023) to allow for comparison. However, due to the nature of the 3D analysis, some images that contained smaller Z-stacks and therefore had cut-off branches in the Z-plane previously used for the 2D analysis, could not be used. In addition, some images had multiple microglia analyzed where 2 or more fully intact microglia were present. Therefore, the sample size for the current study is slightly different (n=3–5 mice/group; 32–94 microglia/group) compared to González Ibáñez et al., (2023) (n=3–5 mice/group; 51–88 microglia/group).

Data shown are reported as means ± standard error of the mean (SEM). Normality of the

data was assessed using the Shapiro-Wilk test. The data did not pass the normality test, but due to

the fact that no non-parametric test yet exists for our study design, parametric tests were used.

We first sought to perform statistical tests identical to those performed by González Ibáñez

et al., (2023) to be able to compare our 3D analysis with their 2D data. This was accomplished by

performing a 2-way analysis of variance (ANOVA) test with diet and stress as our independent

factors. Our *a priori* hypothesis was that we would find similar significant results to González

Ibáñez et al., (2023). To test this hypothesis, we also performed Tukey's *post-hoc* test to assess

these comparisons, as did González Ibáñez et al., (2023). This analysis was performed using

GraphPad Prism (v10.0.3 Graphstats Technologies).

To more accurately showcase the individual animal variability within each group, we also

performed a nested 2-way ANOVA, with the animals nested within stress and stress within diet.

This analysis was not performed by González Ibáñez et al., (2023). Due to this, we have re-

analyzed the raw 2D microglia morphology data using a nested design. To accomplish this, we

used a mixed-effects model, where diet and stress are fixed effects and animal is a random effect.

We then computed the estimated marginal means for the combinations between diet and stress

before performing pairwise comparisons between these means. The Tukey's *post-hoc* test was

applied to perform the pairwise comparisons, to allow for comparison with the non-nested data.

The Kenward-Roger degrees-of-freedom method was used to estimate the degrees of freedom.

This analysis was performed in RStudio v2024.04.2+764.

For all analyses, statistical significance was reached when *p* value < 0.05 and potential

changes were indicated at *p* < 0.1. No statistical outliers were removed. All graphs were created in

GraphPad Prism v10.0.3.

Results

Direct 2D *vs* 3D Comparison

We first sought to perform morphological analyses that were most similar to those performed in

González Ibánez et al., (2023) for direct comparison between the 2D and 3D morphology analysis

methods. These analyses were performed for all cellular values within each group. For ease of

reference, we will break up this section per morphological characteristic. Please refer to Table 2

for a comparison between different analysis methods and statistical designs in terms of statistical
differences.

Soma Area vs *Soma Volume*

In terms of interactions, González Ibáñez et al., (2023) reported a significant interaction of stress
x diet ($F_{(2,374)}$=5.63, p=0.0039) and a main effect of stress ($F_{(2,374)}$=9.63, p=<0.0001) for soma area.
Our 3D analysis supports this, with a significant interaction observed for stress x diet ($F_{(2,339)}$=8.32,
p=0.0003) and a main effect of stress ($F_{(2,339)}$=17.39, p=<0.0001) with a similar confidence
compared to the 2D analysis.

Further investigation between groups revealed a significant difference (p=0.0169) between
the CD CTRL and KD CTRL groups, indicating larger soma area in the KD CTRL group in the
2D analysis. However, in our 3D analysis, we did not observe a significant difference between
these groups (p=0.1196) (Fig. 2B,E). In addition, González Ibáñez et al., (2023) reported a
significant difference (p=0.0002) between the CD CTRL and CD SUS group, with microglia from
the CD SUS having a larger soma area. Our 3D analysis supports this finding, with an observed
significant difference (p=0.0004) between these groups, with a similar confidence to that reported
in the 2D analysis (Fig. 2B,E). In addition, our 3D analysis also found significant differences
between KD CTRL and KD SUS (p=0.0189) (larger soma for KD SUS), KD CTRL and KD RES
(p=0.0133) (larger soma for KD CTRL), KD SUS and KD RES (p=<0.0001) (larger soma for KD
SUS), and CD RES and KD RES (p=0.0480) (larger soma for CD RES) groups (Fig. 2B,E) that
was not observed in the 2D analysis (p=0.9974, p=0.5111, p=0.3346 and p=0.9770, respectively).

Arborization Area vs *Convex Hull Volume*

For arborization area, González Ibánez et al., (2023) reported a significant interaction of stress x
diet ($F_{(2,374)}$=5.42, p=0.0048) and a main effect of stress ($F_{(2,374)}$=4.61, p=0.0105). Similarly, our
3D analysis revealed a significant interaction between stress x diet ($F_{(2,339)}$=15.00, p=<0.0001), but
we did not observe a main effect of stress ($F_{(2,339)}$=2.17, p=0.1153). However, we did observe a
significant main effect of diet ($F_{(1,339)}$=10.61, p=0.0012).

Similar to the soma area, the 2D analysis resulted in a significant difference between CD
CTRL and KD CTRL (p=0.0391), with a larger arborization area in the KD CTRL group. This
finding was supported by our 3D analysis (p=<0.0001) (Fig. 2D,F). The 2D analysis also resulted
in a significant difference between CD CTRL and CD RES (p=0.0017) groups, with CD RES
having a higher arborization area, but this finding was not found by our 3D analysis (p=0.9051)

(Fig. 2D,F). In addition to these, our 3D analysis revealed significant differences between CD CTRL and CD SUS ($p=<0.0001$) (larger convex hull for CD SUS), CD SUS and CD RES ($p=0.0066$) (larger convex hull for CD SUS), and CD RES and KD RES ($p=0.0119$) (larger convex hull for KD RES) groups (Fig. 2D,F). These differences were not observed in the 2D analysis ($p=0.3333$, $p=0.4992$ and $p=0.7312$, respectively).

Morphology Index

González Ibáñez et al. (2023) reported a significant main effect of stress ($F_{(2,374)}=9.63$, $p=<0.0001$) for the morphology index. Our 3D analysis supports this finding ($F_{(2,339)}=6.94$, $p=0.0011$), and further revealed a significant interaction of stress x diet ($F_{(2,339)}=8.27$, $p=0.0003$).

Further analysis revealed a significant difference between KD SUS and KD RES groups in the 2D ($p=0.0066$) and 3D ($p=<0.0001$) analysis, with the KD SUS group having a higher value (Fig. 2G). Our 3D analysis further revealed significant differences between KD CTRL and KD SUS ($p=0.0001$) (higher value for KD SUS), as well as CD SUS and KD SUS ($p=0.0047$) (higher value for KD SUS) groups (Fig. 2G). These results were not observed in the 2D analysis ($p=0.1016$ and $p=0.9974$, respectively).

Overall, these results suggest that although similar, the 2D and 3D analysis methods do result in distinct differences between the analyzed groups, with often more differences observed in the 3D analysis method. These differences observed with the 3D method are also accompanied by a higher confidence.

Further 3D Morphological Characterization

In addition to soma volume, convex hull volume and morphology index, we quantified the number of Sholl intersections at 5 μm radius intervals, and the critical radius (i.e., the radius with the highest number of intersections). Although these measurements are possible to quantify in 2D, they were not performed in González Ibáñez et al., (2023).

The critical radius analysis did not indicate any significant differences between groups (Fig. 2I). However, the Sholl analysis revealed a significant interaction between stress x diet ($F_{(2,339)}=6.39$, $p=0.0019$), and main effects of stress ($F_{(2,339)}=10.29$, $p=<0.0001$) and diet ($F_{(2,339)}=6.61$, $p=0.0106$). Our *post-hoc* analysis indicated that there is a significant difference between CD CTRL and KD CTRL ($p=0.0006$), KD CTRL and KD SUS ($p=<0.0001$), as well as

KD SUS and KD RES (p=0.0051) groups (Fig. 2C,H). Among these differences, KD CTRL and

KD RES both have a greater number of total Sholl intersections compared to their respective

comparisons.

Exploration of a Nested Statistical Design

We next sought to investigate how the 2D and 3D microglia morphology analysis methods could

differ with the utilization of a different statistical method that also takes into account individual

animal variability. As a note, the data reported in González Ibáñez et al., (2023) used individual

cell values as a sample size to examine the variability of microglial morphology between cells,

and did not use animal means which do not take into account intercellular variability. However,

the nested design takes into account both cellular and animal variability. Therefore, we performed

a nested 2-way ANOVA in RStudio with the raw 2D microglia morphology data from González

Ibáñez et al., (2023) and the re-analyzed data from the 3D microglia morphology method.

Soma Area vs. *Soma Volume*

The 2D data analyzed with a nested 2-way ANOVA revealed a significant interaction of stress x

diet ($F_{(2,374)}$=5.76, p=0.0035) and a main effect of stress ($F_{(2,374)}$=6.97, p=0.0011), which is

supported by the 3D data (stress x diet ($F_{(2,339)}$=9.66, p=<0.0001); stress ($F_{(2,339)}$=22.28,

p=<0.0001)).

Tukey's *post-hoc* indicated that there was a significant difference between CD CTRL and

CD SUS (p=0.0028) groups in the 2D analysis (Fig. 3Aii), with CD SUS having larger soma area.

We did not observe any significant differences between groups in the 3D analysis (Fig. 3Ai).

However, we observed a potential increase in soma volume in the KD SUS group, compared to

the KD RES (p=0.0659) group in the 3D analysis (Fig. 3Ai).

Arborization Area vs. *Convex Hull Volume*

For the 2D arborization area, a significant interaction was observed for stress x diet ($F_{(2,374)}$=6.09,

p=0.0025), as well as a main effect for stress ($F_{(2,374)}$=3.76, p=0.0243). The 3D analysis for convex

hull volume resulted in a significant interaction of stress x diet ($F_{(2,339)}$=16.94, p=<0.0001), as well

as main effects of stress ($F_{(339)}$=5.03, p=0.0071) and diet ($F_{(2,339)}$=9.21, p=0.0026). However, we

observed no significant differences after the *post-hoc* test for either analysis method (Fig. 3Bi,ii).

Morphology Index

The nested analysis revealed significant main effects of stress for both the 2D ($F_{(2,374)}$=10.75,
p=<0.0001) and 3D ($F_{(2,339)}$=9.06, p=0.0001) analysis methods. Additionally, a significant
interaction was observed in the 3D analysis between stress x diet ($F_{(2,339)}$=11.13, p=<0.0001). We
observed no significant differences after the *post-hoc* test for either analysis method (Fig. 3Ci,ii).
3D Sholl Analysis
We observed a significant interaction between stress and diet ($F_{(2,339)}$=8.69, p=0.0002), as well as
main effects of stress ($F_{(2,339)}$=11.28, p=<0.0001) and diet ($F_{(2,339)}$=20.91, p=<0.0001) for the 3D
Sholl analysis. However, there were no significant differences between groups after our *post-hoc*
analysis (Fig. 3D). No statistically significant effects were found for critical radius in the 3D
dataset (Fig. 3E).
305
306 Overall, when statistical tests more accurately account for individual animal variability within the
307 groups, many of the statistically significant comparisons between individual groups using whole
308 group data are not observed, especially for the 3D dataset.
309

Discussion

This methods research paper set out to compare 2D and 3D morphology analysis tools of
IBA1[+]/TMEM119[+] microglia from the ventral hippocampus of adult male mice imaged using a
confocal microscope (40x) from a previously published research paper from our group, González
Ibáñez et al., (2023). The 2D microglia morphology method used in González Ibáñez et al., (2023)
is a well-established methodology that has been utilized in multiple publications over the past
decade (Basilico et al., 2022; Bordeleau et al., 2020, 2021; Chen et al., 2024; Enlow et al., 2021;
Fernández De Cossío et al., 2021; González Ibáñez et al., 2023; Hui et al., 2019; Hui, St-Pierre,
Detuncq, et al., 2018; Hui, St-Pierre, El Hajj, et al., 2018; Lecours et al., 2020; Paasila et al., 2019;
Picard et al., 2021; Savage et al., 2020; Tremblay et al., 2012). The 3D microglial morphology
method used in this study was performed using Imaris v10.0.2, a well-known and powerful
analysis software for 3D microscopy data (Haass-Koffler et al., 2012; Nguyen & Thompson-Peer,
2021; Rose et al., 2024; Singaravelu et al., 2017; Testen et al., 2024; VanRyzin et al., 2019).
Although microglial/glial cell morphology analysis has not been extensively performed using
Imaris in comparison to 2D methods, its potential—based on the available options within the
Imaris workspace—is notable.

Overall, we found measurable differences between the two methods, with the 3D analysis

method producing overlapping, but also distinct significant differences between groups to a greater

extent than with the 2D analysis. In addition, a nested statistical design—which more accurately

takes into account individual animal variability—also produced distinct results compared to total

cell/group data, which perhaps highlights more robust findings.

2D vs *3D Microglia Morphology Analysis*

The most striking differences between the 2D and 3D methods were related to the soma. Our

previous 2D analysis by González Ibáñez et al., (2023) indicated two statistically different specific

comparisons between groups, whereas the 3D analysis indicated five. However, the differences

observed between CD CTRL and KD CTRL in the 2D analysis were not found in the 3D analysis.

It is not surprising that soma should produce the most varied responses between these methods.

The 2D analysis involves a Z-stack image that is converted to a maximum intensity projection

image, resulting in a soma area that is equal to the widest portion of the soma, whereas the 3D

analysis takes into account soma directionality in all three planes, as well as indentations and

inconsistencies in soma shape. These can be common in microglia, and many other cells, especially

when in apposition to other cells or structures, such as blood vessels or axons (Stratoulias et al.,

2019). Therefore, it is possible that the 3D analysis is better able to capture slight variations in

soma shape, revealing minute changes to cell populations within groups that the 2D analysis

method cannot.

A similar argument can also be made for arborization area. Our 3D analysis indicated

significant differences in arborization area (measured by convex hull volume) between CD CTRL

and KD CTRL groups, a finding which has been observed by our group and others (González

Ibáñez et al., 2023; Gzielo et al., 2019). However, there were also distinct interactions found in 2D

that the 3D analysis did not capture, and *vice versa*. The most likely cause of this is the fact that

although microglia maintain distinct territories, microglial branches do somewhat overlap in 3D

space, and microglial clusters can form during an insult (e.g., lesions in multiple sclerosis), as well

as in health (Hume et al., 1983; Van Horssen et al., 2012). The maximum projection images make

it extremely challenging to differentiate between overlapping branches, resulting in slightly less

accurate depictions of arborization area. This being said, the 2D analysis method is by no means

considered incorrect, but is instead another method with strengths and limitations that should be

contemplated before performing analysis. This is also the case with 3D analysis. Although
differentiating between overlapping branches is relatively simple, 3D analysis software (*e.g.*,
Imaris) does not always accurately depict branch length. For example, a recent study comparing
different morphology analysis tools for neurons of the *Drosophila* larvae reported that although
automatic detection using 3D analysis software (*e.g.*, Imaris) was the most efficient and convenient
method for dendrite reconstruction of the methods tested, automatic detection of dendrites in
Imaris resulted in significantly shorter dendritic lengths compared to manually traced dendrites
(Nguyen & Thompson-Peer, 2021). However, they also reported that this is somewhat mitigated
by the ability to easily edit reconstructed objects manually, a feature that we also relied upon in
our analysis. It is also important to note that the 2D microglial morphology analysis has a similar
number of possible quantifiable measurements compared to the 3D analysis, further bolstering its
capabilities (Leyh et al., 2021).

Although the 3D method may be able to pick up on more subtle changes to morphology,
such as soma indentations, and, by extension, between groups, robust morphological changes are
still captured by the 2D analysis. For example, the morphology index—a more descriptive measure
of changing microglial morphology as it takes into account both soma size and arborization area—
was observed to be significantly higher in the KD SUS group compared to the KD RES group in
the 2D and 3D analysis methods (Fig. 2G).

Nonetheless, despite the noticeable differences between these two methods, the overall
findings of González Ibáñez et al., (2023) for microglial morphology using 2D analysis methods
are supported by the 3D analysis method. Ketogenic diet and stress do have an effect on microglial
morphology at baseline, and based on the 3D morphological data, the findings reported by
González Ibáñez et al., (2023)—that ketogenic diet can help prevent some of the morphological
adaptations microglia have in response to repeated social defeat stress—are supported. Similar
findings between 2D and 3D methods with respect to microglial morphology have also been found
by others. For example, microglia process length was found to be significantly shorter in the
inferior temporal cortex from post-mortem human brain tissue with Alzheimer's disease pathology
compared to age-matched controls using ImageJ (2D) and Imaris software (3D) (Davies et al.,
2017).

Important Considerations for Method Choice

Although the most important consideration for choosing an analysis method is arguably the
research question at hand, varied financial and temporal constraints always accompany each
method and influence choice. The 2D analysis performed by González Ibáñez et al., (2023) was
performed using ImageJ, a free open-source software funded by the National Institutes of Health
(NIH). This software has a wide-array of easily installed plugins that can perform an impressive
number of analyses for cellular morphology and much more. Another free open-sourced software
that can be used for cellular morphology is QuPath. QuPath is extremely user-friendly and has
some built-in analytical features, such as cell detection. These programs can also be combined,
where more easily made annotations performed in QuPath can be sent to ImageJ through QuPath,
and subsequently analyzed using the very convenient ImageJ plugins. Additionally, users can enter
commands and write scripts in both QuPath and ImageJ, allowing for customizable workflows that
can drastically enhance analysis speed and contribute to reproducibility.

Although Z-stacks can be imported into both QuPath and ImageJ—as well as the fact that
ImageJ has some 3D analysis plugins—these programs are quite limited in 3D at this time. This
being said, recent years have seen a surge in interest in 3D morphology characterization, and open-
source plugins for ImageJ (e.g., GIANI), as well as other programs such as Microscopy Image
Browser, are becoming more powerful and convenient for use in 3D (Barry et al., 2022; Belevich
et al., 2016). However, in terms of time, these options still take decidedly longer to analyze cells
in 3D compared to software built for 3D analysis, like Imaris, especially considering batch analysis
capabilities, where analysis parameters can be applied to multiple images at a time. For example,
manually tracing neurons from the *Drosophila* larvae took an average of 21 minutes per image,
whereas Imaris had an average tracing time of 7 minutes per image (Nguyen & Thompson-Peer,
2021).

Imaris is an incredibly powerful tool for visualization and analysis of 3D microscopy data.
Built-in plugins are available in the software, but custom plugins can also be mapped and
performed in Imaris. For example, this project utilized a plugin built by Ironhorse1618 (2023):
https://github.com/Ironhorse1618/Python3.7-Imaris-XTensions, which extracts and sends
reconstructed objects from Imaris to Python 3.7. Subsequent calculations (i.e., soma volume, Sholl
analysis, convex hull, etc.) are performed in Python before the data is sent back to Imaris for ease
of visualization and use. Overall, software like Imaris is extremely powerful, versatile and
efficient, but also financially restrictive. The upfront costs for software such as these are

significant, and are often accompanied by yearly subscription fees. However, more affordable options through core imaging facilities, that in some cases can be accessed remotely, may exist.

Overall, the decision to use either 2D or 3D analysis methods for cell morphology depends heavily on the research question, but also on financial constraints, time and experience of the researcher, as was also highlighted by Green et al., (2022). In Green et al., (2022), inconsistencies between ImageJ-based methods for quantifying microglial morphology in 2D were highlighted. It was suggested that combining methods would provide the most accurate depiction of microglial morphology, as was previously discussed in this paper (Green et al., 2022). However, the authors also emphasized that single-cell analyses are preferable to aggregate data calculated from the means of many microglia within the same photomicrograph (Green et al., 2022). We agree with this suggestion, as means compiled from multiple biological replicates may mask natural variability within the microglial population in terms of morphology. However, single-cell analyses may also result in certain biological replicates driving the dataset—since animal variability is common—and somewhat arbitrarily increase the sample size (Voelkl et al., 2020). To account for this, a nested or Bayesian statistical design can be utilized (Coventry & Bartlett, 2024; Galbraith et al., 2010).

Our study revealed significant differences between data analyzed with a nested 2-way ANOVA with 'animal' as our random factor *vs* a 2-way ANOVA. Both the 2D and 3D datasets analyzed with the nested statistical design resulted in much fewer significant differences between groups. The 3D data in particular was drastically different, as no statistically significant comparisons between individual groups were maintained using the nested design. A likely reason for this is that the pooled individual data points for each group used in the 2-way ANOVA assumes they are all independent from one another, and does not account for similarities within each individual animal, ultimately resulting in an increase in Type 1 errors (Galbraith et al., 2010). Therefore, the nested approach potentially yields more accurate and robust statistically significant effects when compared to pooled group data used in the 2-way ANOVA. However, it is important to note that the Tukey *post-hoc* test is a particularly stringent test, as it makes all possible comparisons despite their biological/experimental relevance, and adjusts *p*-values to account for this. A *post-hoc* test that computes multiple comparisons based off of selected comparisons (e.g., Šídák or Bonferonni multiple comparisons tests) that are biologically and experimentally relevant may be more appropriate. Nonetheless, our goal was to compare the non-nested 2-way ANOVA

with a nested 2-way ANOVA, meaning that the *post-hoc* test, as long as it is consistent between the two designs, is somewhat irrelevant.

Overall, we believe that this methods paper will help shed new light on the differences between 2D and 3D microglial morphology analysis. We also aim to stress that although outcomes are different, general tendencies and overall findings are consistent between the two, and therefore no one method is considered correct or incorrect. Instead, the right method of choice should depend on the research question and whether or not it can be sufficiently and accurately answered using 2D or 3D image analysis tools.

Conclusion

We re-analyzed raw microglia confocal microscopy images in 3D from a previously published paper from our group that analyzed microglia morphology in 2D in the ventral hippocampus of adult mice. Overall, the 3D analysis method supports many of the observations made using the 2D analysis method, but also indicates many significant differences between groups that were not observed using the 2D method. This suggests that the 3D analysis method may be better able to capture minute changes in microglial morphology, possibly due to superior visualization of the cell compared to collapsed 2D images. However, overall conclusions about microglial morphological adaptation to ketogenic diet and stress from both methods are similar, highlighting the fact that both methods are appropriate for this type of analysis. In addition, we compare data analyzed using a nested and non-nested design, showing how individual animal variability can influence datasets.

Acknowledgments

We acknowledge with respect the ləkʷəŋən peoples on whose traditional territory the University of Victoria stands and the Songhees, Esquimalt and WSÁNEĆ peoples whose historical relationships with the land continue to this day. We also acknowledge the animal work done by Drs. Kaushik Sharma, Nathalie Vernoux and Kanchan Bisht, who are authors on the original manuscript. In addition, we acknowledge that image analysis for this work was performed in the University of British Columbia Life Sciences Institute Imaging Core Facility, RRID: SCR_023783. We also thank the Imaris application support team for their guidance for the use of Imaris software.

M-ÈT is a Tier II Canada Research Chair in *Neurobiology of Aging and Cognition* and
holds funding through the Natural Sciences and Engineering Research Council of Canada (RGPIN:
2024-06043), Canadian Institutes of Health Research (CIHR) (PJT461831) and the Canadian
Foundation for Innovation (John R. Evans Leaders Fund: 39965). HAV was a CIHR Fellow,
Michael Smith Health Research BC Research Trainee and was supported by a Brain Canada
Training Fellowship and BC Women's Health Research Institute Fellowship. FGI is a Michael
Smith Health Research BC research trainee and was supported by a full doctoral scholarship by
the Mexican Council of Science and Technology (CONACYT, now CONAHCYT). CJM was
supported by a CIHR CGS-M scholarship, and a graduate grant from the Branch Out Neurological
Foundation. EDT was funded by the KF Hein grant, Jo Kolk grant, Hendrik Muller grant, and the
UMC Utrecht Strategic Network Development grant.

References

Ambrose, N., Rodriguez, M., Waters, K. A., & Machaalani, R. (2020). Microglia in the human

infant brain and factors that affect expression. *Brain, Behavior, & Immunity - Health*, *7*,

100117. https://doi.org/10.1016/j.bbih.2020.100117

Bachstetter, A. D., Ighodaro, E. T., Hassoun, Y., Aldeiri, D., Neltner, J. H., Patel, E., Abner, E. L., &

Nelson, P. T. (2017). Rod-shaped microglia morphology is associated with aging in two

human autopsy series. *Neurobiology of Aging*, *52*, 98–105.

https://doi.org/10.1016/j.neurobiolaging.2016.12.028

Badanjak, K., Fixemer, S., Smajić, S., Skupin, A., & Grünewald, A. (2021). The Contribution of

Microglia to Neuroinflammation in Parkinson's Disease. *International Journal of*

Molecular Sciences, *22*(9), 4676. https://doi.org/10.3390/ijms22094676

Barry, D. J., Gerri, C., Bell, D. M., D'Antuono, R., & Niakan, K. K. (2022). GIANI – open-source

software for automated analysis of 3D microscopy images. *Journal of Cell Science*,

135(10), jcs259511. https://doi.org/10.1242/jcs.259511

Basilico, B., Ferrucci, L., Ratano, P., Golia, M. T., Grimaldi, A., Rosito, M., Ferretti, V., Reverte, I.,

Sanchini, C., Marrone, M. C., Giubettini, M., De Turris, V., Salerno, D., Garofalo, S., St-

Pierre, M., Carrier, M., Renzi, M., Pagani, F., Modi, B., … Ragozzino, D. (2022). Microglia

control glutamatergic synapses in the adult mouse hippocampus. *Glia*, *70*(1), 173–195.

https://doi.org/10.1002/glia.24101

Belevich, I., Joensuu, M., Kumar, D., Vihinen, H., & Jokitalo, E. (2016). Microscopy Image

Browser: A Platform for Segmentation and Analysis of Multidimensional Datasets. *PLOS*

Biology, *14*(1), e1002340. https://doi.org/10.1371/journal.pbio.1002340

Bordeleau, M., Fernández De Cossío, L., Lacabanne, C., Savage, J. C., Vernoux, N., Chakravarty,

M., & Tremblay, M.-È. (2021). Maternal high-fat diet modifies myelin organization,

microglial interactions, and results in social memory and sensorimotor gating deficits in

adolescent mouse offspring. *Brain, Behavior, & Immunity - Health*, *15*, 100281.

https://doi.org/10.1016/j.bbih.2021.100281

Bordeleau, M., Lacabanne, C., Fernández De Cossío, L., Vernoux, N., Savage, J. C., González-

Ibáñez, F., & Tremblay, M.-È. (2020). Microglial and peripheral immune priming is

partially sexually dimorphic in adolescent mouse offspring exposed to maternal high-fat

diet. *Journal of Neuroinflammation*, *17*(1), 264. https://doi.org/10.1186/s12974-020-

01914-1

Chen, K., Qi, X., Zhu, L., Li, M., Cong, B., & Li, Y. (2024). Quantitative analysis of microglia

morphological changes in the hypothalamus of chronically stressed rats. *Brain Research*

Bulletin, *206*, 110861. https://doi.org/10.1016/j.brainresbull.2023.110861

Colonna, M., & Butovsky, O. (2017). Microglia Function in the Central Nervous System During

Health and Neurodegeneration. *Annual Review of Immunology*, *35*(1), 441–468.

https://doi.org/10.1146/annurev-immunol-051116-052358

Coventry, B. S., & Bartlett, E. L. (2024). Practical Bayesian Inference in Neuroscience: Or How I

Learned to Stop Worrying and Embrace the Distribution. *Eneuro*, *11*(7), ENEURO.0484-

23.2024. https://doi.org/10.1523/ENEURO.0484-23.2024

Davalos, D., Grutzendler, J., Yang, G., Kim, J. V., Zuo, Y., Jung, S., Littman, D. R., Dustin, M. L., &

Gan, W.-B. (2005). ATP mediates rapid microglial response to local brain injury in vivo.

Nature Neuroscience, *8*(6), 752–758. https://doi.org/10.1038/nn1472

Davies, D. S., Ma, J., Jegathees, T., & Goldsbury, C. (2017). Microglia show altered morphology

and reduced arborization in human brain during aging and A lzheimer's disease. *Brain*

Pathology, *27*(6), 795–808. https://doi.org/10.1111/bpa.12456

Davis, B. M., Salinas-Navarro, M., Cordeiro, M. F., Moons, L., & De Groef, L. (2017).

Characterizing microglia activation: A spatial statistics approach to maximize information

extraction. *Scientific Reports*, *7*(1), 1576. https://doi.org/10.1038/s41598-017-01747-8

Davis, E. J., Foster, T. D., & Thomas, W. E. (1994). Cellular forms and functions of brain microglia.

Brain Research Bulletin, *34*(1), 73–78. https://doi.org/10.1016/0361-9230(94)90189-9

Doorn, K. J., Goudriaan, A., Blits-Huizinga, C., Bol, J. G. J. M., Rozemuller, A. J., Hoogland, P. V. J.

M., Lucassen, P. J., Drukarch, B., Van De Berg, W. D. J., & Van Dam, A. (2014). Increased

Amoeboid Microglial Density in the Olfactory Bulb of P arkinson's and A lzheimer's

Patients. *Brain Pathology*, *24*(2), 152–165. https://doi.org/10.1111/bpa.12088

Doorn, K. J., Moors, T., Drukarch, B., Van De Berg, W. D., Lucassen, P. J., & Van Dam, A.-M.

(2014). Microglial phenotypes and toll-like receptor 2 in the substantia nigra and

hippocampus of incidental Lewy body disease cases and Parkinson's disease patients.

Acta Neuropathologica Communications, *2*(1), 90. https://doi.org/10.1186/s40478-014-

0090-1

Enlow, W., Bordeleau, M., Piret, J., Ibáñez, F. G., Uyar, O., Venable, M.-C., Goyette, N.,

Carbonneau, J., Tremblay, M.-E., & Boivin, G. (2021). Microglia are involved in

phagocytosis and extracellular digestion during Zika virus encephalitis in young adult

immunodeficient mice. *Journal of Neuroinflammation*, *18*(1), 178.

https://doi.org/10.1186/s12974-021-02221-z

Erny, D., de Angelis, A. L. H., Jaitin, D., Wieghofer, P., Staszewski, O., David, E., Keren-Shaul, H.,

Mahlakoiv, T., Jakobshagen, K., Buch, T., Schwierzeck, V., Utermöhlen, O., Chun, E.,

Garrett, W. S., McCoy, K. D., Diefenbach, A., Staeheli, P., Stecher, B., Amit, I., & Prinz, M.

(2015). Host microbiota constantly control maturation and function of microglia in the

CNS. *Nature Neuroscience*, *18*(7), 965–977. https://doi.org/10.1038/nn.4030

Fernández De Cossío, L., Lacabanne, C., Bordeleau, M., Castino, G., Kyriakakis, P., & Tremblay,

M.-È. (2021). Lipopolysaccharide-induced maternal immune activation modulates

microglial CX3CR1 protein expression and morphological phenotype in the hippocampus

and dentate gyrus, resulting in cognitive inflexibility during late adolescence. *Brain,*

Behavior, and Immunity, *97*, 440–454. https://doi.org/10.1016/j.bbi.2021.07.025

Fujikawa, R., & Jinno, S. (2022). Identification of hyper-ramified microglia in the CA1 region of

the mouse hippocampus potentially associated with stress resilience. *European Journal*

of Neuroscience, *56*(8), 5137–5153. https://doi.org/10.1111/ejn.15812

Galbraith, S., Daniel, J. A., & Vissel, B. (2010). A Study of Clustered Data and Approaches to Its

Analysis. *The Journal of Neuroscience*, *30*(32), 10601–10608.

https://doi.org/10.1523/JNEUROSCI.0362-10.2010

Gao, C., Jiang, J., Tan, Y., & Chen, S. (2023). Microglia in neurodegenerative diseases: Mechanism

and potential therapeutic targets. *Signal Transduction and Targeted Therapy*, *8*(1), 359.

https://doi.org/10.1038/s41392-023-01588-0

Ginhoux, F., Greter, M., Leboeuf, M., Nandi, S., See, P., Gokhan, S., Mehler, M. F., Conway, S. J.,

Ng, L. G., Stanley, E. R., Samokhvalov, I. M., & Merad, M. (2010). Fate Mapping Analysis

Reveals That Adult Microglia Derive from Primitive Macrophages. *Science*, *330*(6005),

841–845. https://doi.org/10.1126/science.1194637

Giordano, K. R., Denman, C. R., Dubisch, P. S., Akhter, M., & Lifshitz, J. (2021). An update on the

rod microglia variant in experimental and clinical brain injury and disease. *Brain*

Communications, *3*(1), fcaa227. https://doi.org/10.1093/braincomms/fcaa227

González Ibáñez, F., Halvorson, T., Sharma, K., McKee, C. G., Carrier, M., Picard, K., Vernoux, N.,

Bisht, K., Deslauriers, J., Lalowski, M., & Tremblay, M.-È. (2023). Ketogenic diet changes

microglial morphology and the hippocampal lipidomic profile differently in stress

susceptible versus resistant male mice upon repeated social defeat. *Brain, Behavior, and*

Immunity, *114*, 383–406. https://doi.org/10.1016/j.bbi.2023.09.006

Green, T. R. F., Murphy, S. M., & Rowe, R. K. (2022). Comparisons of quantitative approaches for

assessing microglial morphology reveal inconsistencies, ecological fallacy, and a need for

standardization. *Scientific Reports*, *12*(1), 18196. https://doi.org/10.1038/s41598-022-

23091-2

Gyoneva, S., Hosur, R., Gosselin, D., Zhang, B., Ouyang, Z., Cotleur, A. C., Peterson, M., Allaire,

N., Challa, R., Cullen, P., Roberts, C., Miao, K., Reynolds, T. L., Glass, C. K., Burkly, L., &

Ransohoff, R. M. (2019). *Cx3cr1-* deficient microglia exhibit a premature aging

transcriptome. *Life Science Alliance*, *2*(6), e201900453.

https://doi.org/10.26508/lsa.201900453

Gzielo, K., Soltys, Z., Rajfur, Z., & Setkowicz, Z. K. (2019). The Impact of the Ketogenic Diet on

Glial Cells Morphology. A Quantitative Morphological Analysis. *Neuroscience*, *413*, 239–

251. https://doi.org/10.1016/j.neuroscience.2019.06.009

Haass-Koffler, C. L., Naeemuddin, M., & Bartlett, S. E. (2012). An Analytical Tool that Quantifies

Cellular Morphology Changes from Three-dimensional Fluorescence Images. *Journal of*

Visualized Experiments, *66*, 4233. https://doi.org/10.3791/4233

Hellwig, S., Brioschi, S., Dieni, S., Frings, L., Masuch, A., Blank, T., & Biber, K. (2016). Altered

microglia morphology and higher resilience to stress-induced depression-like behavior in

CX3CR1-deficient mice. *Brain, Behavior, and Immunity*, *55*, 126–137.

https://doi.org/10.1016/j.bbi.2015.11.008

Hinwood, M., Morandini, J., Day, T. A., & Walker, F. R. (2012). Evidence that microglia mediate

the neurobiological effects of chronic psychological stress on the medial prefrontal

610   cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, *22*(6), 1442–1454.

611   https://doi.org/10.1093/cercor/bhr229

612   Hinwood, M., Tynan, R. J., Charnley, J. L., Beynon, S. B., Day, T. A., & Walker, F. R. (2013). Chronic

613   stress induced remodeling of the prefrontal cortex: Structural re-organization of

614   microglia and the inhibitory effect of minocycline. *Cerebral Cortex (New York, N.Y.: 1991)*,

615   *23*(8), 1784–1797. https://doi.org/10.1093/cercor/bhs151

616   Holloway, O. G., Canty, A. J., King, A. E., & Ziebell, J. M. (2019). Rod microglia and their role in

617   neurological diseases. *Seminars in Cell & Developmental Biology*, *94*, 96–103.

618   https://doi.org/10.1016/j.semcdb.2019.02.005

619   Hui, C. W., Bhardwaj, S. K., Sharma, K., Joseph, A. T., Bisht, K., Picard, K., Tremblay, M.-È., &

620   Srivastava, L. K. (2019). Microglia in the developing prefrontal cortex of rats show

621   dynamic changes following neonatal disconnection of the ventral hippocampus.

622   *Neuropharmacology*, *146*, 264–275. https://doi.org/10.1016/j.neuropharm.2018.12.007

623   Hui, C. W., St-Pierre, A., El Hajj, H., Remy, Y., Hébert, S. S., Luheshi, G. N., Srivastava, L. K., &

624   Tremblay, M.-È. (2018). Prenatal Immune Challenge in Mice Leads to Partly Sex-

625   Dependent Behavioral, Microglial, and Molecular Abnormalities Associated with

626   Schizophrenia. *Frontiers in Molecular Neuroscience*, *11*, 13.

627   https://doi.org/10.3389/fnmol.2018.00013

628   Hui, C. W., St-Pierre, M.-K., Detuncq, J., Aumailley, L., Dubois, M.-J., Couture, V., Skuk, D.,

629   Marette, A., Tremblay, J. P., Lebel, M., & Tremblay, M.-È. (2018). Nonfunctional mutant

630   Wrn protein leads to neurological deficits, neuronal stress, microglial alteration, and

immune imbalance in a mouse model of Werner syndrome. *Brain, Behavior, and*

Immunity, *73*, 450–469. https://doi.org/10.1016/j.bbi.2018.06.007

Hume, D. A., Perry, V. H., & Gordon, S. (1983). Immunohistochemical localization of a

macrophage-specific antigen in developing mouse retina: Phagocytosis of dying neurons

and differentiation of microglial cells to form a regular array in the plexiform layers. *The*

Journal of Cell Biology, *97*(1), 253–257. https://doi.org/10.1083/jcb.97.1.253

Jensen, M. B., Gonzalez, B., Castellano, B., & Zimmer, J. (1994). Microglial and astroglial

reactions to anterograde axonal degeneration: A histochemical and

immunocytochemical study of the adult rat fascia dentata after entorhinal perforant

path lesions. *Experimental Brain Research*, *98*(2). https://doi.org/10.1007/BF00228413

Karperien, A., Ahammer, H., & Jelinek, H. F. (2013). Quantitating the subtleties of microglial

morphology with fractal analysis. *Frontiers in Cellular Neuroscience*, *7*, 3.

https://doi.org/10.3389/fncel.2013.00003

Kettenmann, H., Hanisch, U.-K., Noda, M., & Verkhratsky, A. (2011). Physiology of microglia.

Physiological Reviews, *91*(2), 461–553. https://doi.org/10.1152/physrev.00011.2010

Kettenmann, H., Kirchhoff, F., & Verkhratsky, A. (2013). Microglia: New roles for the synaptic

stripper. *Neuron*, *77*(1), 10–18. https://doi.org/10.1016/j.neuron.2012.12.023

Kozlowski, C., & Weimer, R. M. (2012). An Automated Method to Quantify Microglia

Morphology and Application to Monitor Activation State Longitudinally In Vivo. *PLoS*

ONE, *7*(2), e31814. https://doi.org/10.1371/journal.pone.0031814

Kreutzberg, G. W. (1996). Microglia: A sensor for pathological events in the CNS. *Trends in*

Neurosciences, *19*(8), 312–318. https://doi.org/10.1016/0166-2236(96)10049-7

Ladeby, R., Wirenfeldt, M., Garcia-Ovejero, D., Fenger, C., Dissing-Olesen, L., Dalmau, I., &

Finsen, B. (2005). Microglial cell population dynamics in the injured adult central

nervous system. *Brain Research Reviews*, *48*(2), 196–206.

https://doi.org/10.1016/j.brainresrev.2004.12.009

Lawson, L. J., Perry, V. H., Dri, P., & Gordon, S. (1990). Heterogeneity in the distribution and

morphology of microglia in the normal adult mouse brain. *Neuroscience*, *39*(1), 151–170.

https://doi.org/10.1016/0306-4522(90)90229-w

Lecours, C., St-Pierre, M.-K., Picard, K., Bordeleau, M., Bourque, M., Awogbindin, I. O., Benadjal,

A., Ibanez, F. G., Gagnon, D., Cantin, L., Parent, M., Di Paolo, T., & Tremblay, M.-E. (2020).

Levodopa partially rescues microglial numerical, morphological, and phagolysosomal

alterations in a monkey model of Parkinson's disease. *Brain, Behavior, and Immunity*, *90*,

81–96. https://doi.org/10.1016/j.bbi.2020.07.044

Leyh, J., Paeschke, S., Mages, B., Michalski, D., Nowicki, M., Bechmann, I., & Winter, K. (2021).

Classification of Microglial Morphological Phenotypes Using Machine Learning. *Frontiers*

in Cellular Neuroscience, *15*, 701673. https://doi.org/10.3389/fncel.2021.701673

Morrison, H., Young, K., Qureshi, M., Rowe, R. K., & Lifshitz, J. (2017). Quantitative microglia

analyses reveal diverse morphologic responses in the rat cortex after diffuse brain injury.

Scientific Reports, *7*(1), 13211. https://doi.org/10.1038/s41598-017-13581-z

Morsch, M., Radford, R., Lee, A., Don, E. K., Badrock, A. P., Hall, T. E., Cole, N. J., & Chung, R.

(2015). In vivo characterization of microglial engulfment of dying neurons in the

zebrafish spinal cord. *Frontiers in Cellular Neuroscience*, *9*, 321.

https://doi.org/10.3389/fncel.2015.00321

Nguyen, C., & Thompson-Peer, K. L. (2021). Comparing Automated Morphology Quantification

Software on Dendrites of Uninjured and Injured Drosophila Neurons. *Neuroinformatics*,

19(4), 703–717. https://doi.org/10.1007/s12021-021-09532-9

Nimmerjahn, A., Kirchhoff, F., & Helmchen, F. (2005). Resting Microglial Cells Are Highly Dynamic

Surveillants of Brain Parenchyma in Vivo. *Science*, *308*(5726), 1314–1318.

https://doi.org/10.1126/science.1110647

Paasila, P. J., Davies, D. S., Kril, J. J., Goldsbury, C., & Sutherland, G. T. (2019). The relationship

between the morphological subtypes of microglia and Alzheimer's disease

neuropathology. *Brain Pathology*, *29*(6), 726–740. https://doi.org/10.1111/bpa.12717

Paolicelli, R. C., Sierra, A., Stevens, B., Tremblay, M.-E., Aguzzi, A., Ajami, B., Amit, I., Audinat, E.,

Bechmann, I., Bennett, M., Bennett, F., Bessis, A., Biber, K., Bilbo, S., Blurton-Jones, M.,

Boddeke, E., Brites, D., Brône, B., Brown, G. C., … Wyss-Coray, T. (2022). Microglia states

and nomenclature: A field at its crossroads. *Neuron*, *110*(21), 3458–3483.

https://doi.org/10.1016/j.neuron.2022.10.020

Perego, C., Fumagalli, S., & De Simoni, M.-G. (2013). Three-dimensional Confocal Analysis of

Microglia/macrophage Markers of Polarization in Experimental Brain Injury. *Journal of*

Visualized Experiments : JoVE, *79*, 50605. https://doi.org/10.3791/50605

Perez-Pouchoulen, M., VanRyzin, J. W., & McCarthy, M. M. (2015). Morphological and

Phagocytic Profile of Microglia in the Developing Rat Cerebellum. *Eneuro*, *2*(4),

ENEURO.0036-15.2015. https://doi.org/10.1523/ENEURO.0036-15.2015

Picard, K., Bisht, K., Poggini, S., Garofalo, S., Golia, M. T., Basilico, B., Abdallah, F., Ciano

Albanese, N., Amrein, I., Vernoux, N., Sharma, K., Hui, C. W., Savage, J. C., Limatola, C.,

Ragozzino, D., Maggi, L., Branchi, I., & Tremblay, M.-È. (2021). *Microglial-glucocorticoid*

receptor depletion alters the response of hippocampal microglia and neurons in a chronic

unpredictable mild stress paradigm in female mice. https://doi.org/10.5167/UZH-

208855

Ritzel, R. M., Doran, S. J., Glaser, E. P., Meadows, V. E., Faden, A. I., Stoica, B. A., & Loane, D. J.

(2019). Old age increases microglial senescence, exacerbates secondary

neuroinflammation, and worsens neurological outcomes after acute traumatic brain

injury in mice. *Neurobiology of Aging*, *77*, 194–206.

https://doi.org/10.1016/j.neurobiolaging.2019.02.010

Rose, S. E., Williams, C. A., Hailey, D. W., Mishra, S., Kirkland, A., Keene, C. D., Garden, G. A.,

Jayadev, S., & Young, J. E. (2024). Advancements in high-resolution 3D microscopy

analysis of endosomal morphology in postmortem Alzheimer's disease brains. *Frontiers*

in Neuroscience, *17*, 1321680. https://doi.org/10.3389/fnins.2023.1321680

Savage, J. C., Carrier, M., & Tremblay, M.-È. (2019). Morphology of Microglia Across Contexts of

Health and Disease. In O. Garaschuk & A. Verkhratsky (Eds.), *Microglia* (Vol. 2034, pp.

13–26). Springer New York. https://doi.org/10.1007/978-1-4939-9658-2_2

Savage, J. C., St-Pierre, M.-K., Carrier, M., El Hajj, H., Novak, S. W., Sanchez, M. G., Cicchetti, F., &

Tremblay, M.-È. (2020). Microglial physiological properties and interactions with

synapses are altered at presymptomatic stages in a mouse model of Huntington's

disease pathology. *Journal of Neuroinflammation*, *17*(1), 98.

https://doi.org/10.1186/s12974-020-01782-9

Schoenen, J. (1982). The dendritic organization of the human spinal cord: The dorsal horn.

Neuroscience, *7*(9), 2057–2087. https://doi.org/10.1016/0306-4522(82)90120-8

Schwarz, J. M., Sholar, P. W., & Bilbo, S. D. (2012). Sex differences in microglial colonization of

the developing rat brain. *Journal of Neurochemistry*, *120*(6), 948–963.

https://doi.org/10.1111/j.1471-4159.2011.07630.x

Shahidehpour, R. K., Higdon, R. E., Crawford, N. G., Neltner, J. H., Ighodaro, E. T., Patel, E., Price,

D., Nelson, P. T., & Bachstetter, A. D. (2021). Dystrophic microglia are associated with

neurodegenerative disease and not healthy aging in the human brain. *Neurobiology of*

Aging, *99*, 19–27. https://doi.org/10.1016/j.neurobiolaging.2020.12.003

Šimončičová, E., Gonçalves de Andrade, E., Vecchiarelli, H. A., Awogbindin, I. O., Delage, C. I., &

Tremblay, M.-È. (2022). Present and future of microglial pharmacology. *Trends in*

Pharmacological Sciences, *43*(8), 669–685. https://doi.org/10.1016/j.tips.2021.11.006

Singaravelu, J., Zhao, L., Fariss, R. N., Nork, T. M., & Wong, W. T. (2017). Microglia in the primate

macula: Specializations in microglial distribution and morphology with retinal position

and with aging. *Brain Structure and Function*, *222*(6), 2759–2771.

https://doi.org/10.1007/s00429-017-1370-x

Smith, K. L., Kassem, M. S., Clarke, D. J., Kuligowski, M. P., Bedoya-Pérez, M. A., Todd, S. M.,

Lagopoulos, J., Bennett, M. R., & Arnold, J. C. (2019). Microglial cell hyper-ramification

and neuronal dendritic spine loss in the hippocampus and medial prefrontal cortex in a

mouse model of PTSD. *Brain, Behavior, and Immunity*, *80*, 889–899.

https://doi.org/10.1016/j.bbi.2019.05.042

Stence, N., Waite, M., & Dailey, M. E. (2001). Dynamics of microglial activation: A confocal time-

lapse analysis in hippocampal slices. *Glia*, *33*(3), 256–266.

Stratoulias, V., Venero, J. L., Tremblay, M., & Joseph, B. (2019). Microglial subtypes: Diversity

within the microglial community. *The EMBO Journal*, *38*(17).

https://doi.org/10.15252/embj.2019101997

Svahn, A. J., Graeber, M. B., Ellett, F., Lieschke, G. J., Rinkwitz, S., Bennett, M. R., & Becker, T. S.

(2013). Development of ramified microglia from early macrophages in the zebrafish optic

tectum. *Developmental Neurobiology*, *73*(1), 60–71.

https://doi.org/10.1002/dneu.22039

Taylor, S. E., Morganti-Kossmann, C., Lifshitz, J., & Ziebell, J. M. (2014). Rod Microglia: A

Morphological Definition. *PLOS ONE*, *9*(5), e97096.

https://doi.org/10.1371/journal.pone.0097096

Testen, A., VanRyzin, J. W., Bellinger, T. J., Kim, R., Wang, H., Gastinger, M. J., Witt, E. A., Franklin,

J. P., Vecchiarelli, H. A., Picard, K., Tremblay, M.-È., & Reissner, K. J. (2024). *Abstinence*

from cocaine self-administration promotes microglia pruning of astrocytes which drives

cocaine-seeking behavior. https://doi.org/10.1101/2024.09.20.614128

Torres-Platas, S. G., Comeau, S., Rachalski, A., Bo, G. D., Cruceanu, C., Turecki, G., Giros, B., &

Mechawar, N. (2014). Morphometric characterization of microglial phenotypes in human

cerebral cortex. *Journal of Neuroinflammation*, *11*, 12. https://doi.org/10.1186/1742-

2094-11-12

Tremblay, M.-È. (2020). A Diversity of Cell Types, Subtypes and Phenotypes in the Central

Nervous System: The Importance of Studying Their Complex Relationships. *Frontiers in*

Cellular Neuroscience, *14*, 628347. https://doi.org/10.3389/fncel.2020.628347

Tremblay, M.-È., Lowery, R. L., & Majewska, A. K. (2010). Microglial Interactions with Synapses

Are Modulated by Visual Experience. *PLoS Biology*, *8*(11), e1000527.

https://doi.org/10.1371/journal.pbio.1000527

Tremblay, M.-È., Zettel, M. L., Ison, J. R., Allen, P. D., & Majewska, A. K. (2012). Effects of aging

and sensory loss on glial cells in mouse visual and auditory cortices. *Glia*, *60*(4), 541–

558. https://doi.org/10.1002/glia.22287

Van Horssen, J., Singh, S., Van Der Pol, S., Kipp, M., Lim, J. L., Peferoen, L., Gerritsen, W., Kooi, E.-

J., Witte, M. E., Geurts, J. J., De Vries, H. E., Peferoen-Baert, R., Van Den Elsen, P. J., Van

Der Valk, P., & Amor, S. (2012). Clusters of activated microglia in normal-appearing white

matter show signs of innate immune activation. *Journal of Neuroinflammation*, *9*(1),

602. https://doi.org/10.1186/1742-2094-9-156

VanRyzin, J. W., Marquardt, A. E., Argue, K. J., Vecchiarelli, H. A., Ashton, S. E., Arambula, S. E.,

Hill, M. N., & McCarthy, M. M. (2019). Microglial Phagocytosis of Newborn Cells Is

Induced by Endocannabinoids and Sculpts Sex Differences in Juvenile Rat Social Play.

Neuron, *102*(2), 435-449.e6. https://doi.org/10.1016/j.neuron.2019.02.006

Verdonk, F., Roux, P., Flamant, P., Fiette, L., Bozza, F. A., Simard, S., Lemaire, M., Plaud, B.,

Shorte, S. L., Sharshar, T., Chrétien, F., & Danckaert, A. (2016). Phenotypic clustering: A

novel method for microglial morphology analysis. *Journal of Neuroinflammation*, *13*(1),

153. https://doi.org/10.1186/s12974-016-0614-7

Vidal-Itriago, A., Radford, R. A. W., Aramideh, J. A., Maurel, C., Scherer, N. M., Don, E. K., Lee, A.,

Chung, R. S., Graeber, M. B., & Morsch, M. (2022). Microglia morphophysiological

diversity and its implications for the CNS. *Frontiers in Immunology*, *13*, 997786.

https://doi.org/10.3389/fimmu.2022.997786

Voelkl, B., Altman, N. S., Forsman, A., Forstmeier, W., Gurevitch, J., Jaric, I., Karp, N. A., Kas, M. J.,

Schielzeth, H., Van De Casteele, T., & Würbel, H. (2020). Reproducibility of animal

research in light of biological variation. *Nature Reviews Neuroscience*, *21*(7), 384–393.

https://doi.org/10.1038/s41583-020-0313-3

Wake, H., Moorhouse, A. J., Jinno, S., Kohsaka, S., & Nabekura, J. (2009). Resting Microglia

Directly Monitor the Functional State of Synapses In Vivo and Determine the Fate of

Ischemic Terminals. *The Journal of Neuroscience*, *29*(13), 3974–3980.

https://doi.org/10.1523/JNEUROSCI.4363-08.2009

Wicks, E. E., Ran, K. R., Kim, J. E., Xu, R., Lee, R. P., & Jackson, C. M. (2022). The Translational

Potential of Microglia and Monocyte-Derived Macrophages in Ischemic Stroke. *Frontiers*

in Immunology, *13*, 897022. https://doi.org/10.3389/fimmu.2022.897022

Yuan, T.-F., Liang, Y.-X., Peng, B., Lin, B., & So, K.-F. (2015). Local proliferation is the main source

of rod microglia after optic nerve transection. *Scientific Reports*, *5*, 10788.

https://doi.org/10.1038/srep10788

Ziebell, J. M., Taylor, S. E., Cao, T., Harrison, J. L., & Lifshitz, J. (2012). Rod microglia: Elongation,

alignment, and coupling to form trains across the somatosensory cortex after

experimental diffuse brain injury. *Journal of Neuroinflammation*, *9*, 247.

https://doi.org/10.1186/1742-2094-9-247

Figures

Figure 1 | Microglia exist within a continuum in which their transcriptional and morphological characteristics often mirror the context of their surrounding microenvironment. These states, including ramified, bushy, and amoeboid morphologies, are dynamic and can change throughout the lifespan or in response to various stimuli. While individual cells may show variability within a single state, an overall shift in microglial morphology can occur depending on specific environmental or pathological signals.

Figure 2 | Representative images of microglia (Ai) stained with immunofluorescent antibodies against IBA1 (red) (Aii), TMEM119 (green) (Aiv) and DAPI (blue) (Aiii) in the Imaris workspace. Soma reconstruction (B) in Imaris is shown alongside filament tracing and sholl analysis (C) and convex hull (D). Microglia morphological parameters from 3D analysis methods are shown (E–G, I and J). Microglia show significant adaptations in all morphological parameters, except for critical radius, in response to diet and stress (E–G, I and J). In addition, the 3D analysis method consistently results in a lower morphology index compared to the 2D analysis method performed in González Ibáñez et al., (2023). n=3–5 mice/group; n=32–94 microglia/group. Statistical significance was assessed using a 2-way ANOVA with Tukey *post hoc* test and data are expressed as mean ± standard error of the mean, where *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. In addition #<0.05, ##<0.01, ###<0.001 depicts a main effect of stress. CD: control diet; KD: ketogenic diet; CTRL: non-stressed controls; SUS: susceptible to stress; RES: resistant to stress; a.u: arbitrary units.

Figure 3 | Comparison between the 2D and 3D microglia morphology analysis methods using a nested 2-way ANOVA. Significant differences were found for soma volume where the KD SUS group has a larger soma volume compared to the KD RES group, whereas soma area revealed significant differences between CD CTRL and KD CTRL, and CD CTRL and CD SUS groups (Ai,ii). Significant differences were found between CD CTRL and KD CTRL, as well as CD SUS for convex hull volume, whereas no significant effects were found for arborization area (Bi,ii). The 3D method revealed significant differences between KD SUS and KD RES groups for morphology index, with KD RES having a smaller ratio, compared to no significant effects observed in the 2D method (Ci,ii). Total sholl intersections analyzed in the 3D method only did not show any significant effects (D), and neither did the critical radius (E). n=3–5 mice/group; n=32–94 microglia/group. Statistical significance was assessed using a nested 2-way ANOVA with Tukey's *post hoc* test and data are expressed as mean ± standard error of the mean, where *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$. In addition #<0.05, ##<0.01, ###<0.001 depicts a main effect of stress. CD: control diet; KD: ketogenic diet; CTRL: non-stressed controls; SUS: susceptible to stress; RES: resistant to stress; a.u: arbitrary units.

Table 1 | Prominent examples of microglia morphologies and their associated significance.

Morphology	Characteristics	Functions	Associated Conditons	Model	References
Hyper-ramified	Highly ramified, with complex and branched branches; appear elongated and thin	Adaptive response to acute and chronic stress; surveillance; potentially neuroprotective	Observed in stress-resilient mice	Mouse; rat	(Fujikawa & Jinno, 2022; Hellwig et al., 2016; Hinwood et al., 2012, 2013; Smith et al., 2019)
Ramified (Homeostatic)	Small soma with long, highly branched branches actively surveilling the brain environment	Surveillance of CNS environment by extending and retracting thin ramified branches; detection of damage or pathogen associated patterns; maintenance of homeostasis	Healthy brain; baseline condition	Human; mouse; rat; zebrafish	(Colonna & Butovsky, 2017; Davalos et al., 2005, 2005; Ginhoux et al., 2010; Nimmerjahn et al., 2005; Svahn et al., 2013; Torres-Platas et al., 2014; Tremblay et al., 2010; Wake et al., 2009)
Hypertrophic (Bushy)	Enlarged soma with increased ramification; shorter and thicker branches with increased complexity	Early response to injury; may facilitate the transition to more immune activated states	Initial injury or inflammation response	Human; mouse; rat	(Ambrose et al., 2020; Jensen et al., 1994; Ladeby et al., 2005; Wicks et al., 2022)
Rod	Straight, rod-shaped appearance; aligned branches; often observed near damaged neurons	Involved in response to axonal damage; potential guidance and repair functions; may interact directly with neurons	Acute injury; CNS infections (e.g. encephalitis); aging; neurodegeneration	Human; mouse; rat	(Bachstetter et al., 2017; Giordano et al., 2021; Holloway et al., 2019; Taylor et al., 2014; Yuan et al., 2015; Ziebell et al., 2012)
Amoeboid	Enlarged soma with very few or no branches; appears rounded and amoeboid-like	Highly phagocytic; involved in clearing debris and dead cells	Developing brain; clearing debris and apoptotic cells; acute injury response; demyelination conditions	Human; mouse; rat; zebrafish	(E. J. Davis et al., 1994; Kettenmann et al., 2011, 2013; Kreutzberg, 1996; Lawson et al., 1990; Morsch et al., 2015; Schwarz et al., 2012;

					Stence et al., 2001; Svahn et al., 2013)
Dystrophic	Fragmented swollen branches; reduced branching and phagocytic activity; senescence marker-positive	Impaired phagocytic function, may contribute to neurodegeneration and impaired tissue repair; potentially a form of microglia senescence	Neurodegenerative disorders (e.g., Alzheimer's, Parkinson's); aging; traumatic brain injury		(Ritzel et al., 2019; Shahidehpour et al., 2021)

Table 2 | Comparison of statistical outcomes for statistically significant differences across different analysis methods and statistical designs. n=3–5 mice/group; n=32–94 microglia/group. Statistical significance was assessed using a 2-way or nested 2-way ANOVA with Tukey *post hoc* test and data are expressed as mean ± standard error of the mean, where $p < 0.05$ is considered significant.

Characteristic	Interaction	3D	Nested 3D	2D	Nested 2D
Soma	CD CTRL *vs.* KD CTRL	ns. 0.1196	ns. 0.8137	**0.0169**	ns. 0.1687
	CD CTRL *vs.* CD SUS	**0.0004**	ns. 0.4421	**0.0002**	**0.0283**
	CD RES *vs.* KD RES	**0.048**	ns. 0.4818	ns. 0.9770	ns. 0.9918
	KD CTRL *vs.* KD SUS	**0.0189**	ns. 0.7274	ns. 0.9974	ns. 0.9983
	KD CTRL *vs.* KD RES	**0.0133**	ns. 0.4940	ns. 0.5111	ns. 0.7481
	KD SUS *vs.* KD RES	**<0.0001**	ns. 0.0659	ns. 0.3346	ns. 0.5859
Branch Spread	CD CTRL *vs.* KD CTRL	**<0.0001**	ns. 0.1567	**0.0391**	ns. 0.6611
	CD CTRL *vs.* CD SUS	**<0.0001**	ns. 0.1597	ns. 0.3333	ns. 0.9075
	CD CTRL *vs.* CD RES	ns. 0.9051	ns. 0.9902	**0.0017**	ns. 0.3920
	CD SUS *vs.* CD RES	**0.0066**	ns. 0.4246	ns. 0.4992	ns. 0.9217
	CD RES *vs.* KD RES	**0.0119**	ns. 0.5199	ns. 0.7312	ns. 0.9879
Morphology Index	CD SUS *vs.* KD SUS	**0.0047**	ns. 0.9272	ns. 0.9974	ns. 0.9996
	KD CTRL *vs.* KD SUS	**0.0001**	ns. 0.7065	ns. 0.1016	ns. 0.6801
	KD SUS *vs.* KD RES	**<0.0001**	ns. 0.2301	**0.0066**	ns. 0.3561

Total Sholl Intersections	CD CTRL *vs.* KD CTRL	**0.0006**	ns. 0.5854	NA	NA
	KD CTRL *vs.* KD SUS	**<0.0001**	ns. 0.3989	NA	NA
	KD SUS *vs.* KD RES	**0.0051**	ns. 0.7791	NA	NA
Critical Radius	ns.	ns.	ns.	NA	NA

CD=Control diet, KD=ketogenic diet, CTRL=control stress, SUS=susceptible phenotype to stress, RES=resistant phenotype to stress, ns.=not significant, NA=not applicable. Numerical values are *p*-values.