Actomyosin Contractility is a Potent Suppressor of Mesoderm Induction by

Human Pluripotent Stem Cells

Loic Fort[1,3], Wenjun Wang[2], and Ian Macara[1,3]

[1] Dept of Cell and Developmental Biology, Vanderbilt University School of Medicine Nashville TN 37240 USA

[2] Dept of Biomedical Engineering, Vanderbilt University Nashville TN 37240 USA

[3] Authors for correspondence

Abstract

The activation of WNT signaling in human pluripotent stem cells drives efficient conversion to

lateral mesoderm, which can be further differentiated into cardiomyocytes. Stabilization of the

WNT effector β-catenin promotes expression of mesoderm-specifying genes such as Brachyury

(Bra/T, encoded by *TBXT*) and Eomesodermin and drives an epithelial-mesenchymal transition

(EMT) that occurs at 49-52 hrs post WNT activation. Interestingly, WNT activation also triggers

an early contraction of stem cell colonies, with increased myosin light chain (MLC)

phosphorylation, resulting to cellular morphological changes. Small molecule inhibitors and

genetic approaches were applied to determine if contractility is related to mesoderm

differentiation. Contrary to expectation, suppressing contractility accelerated differentiation;

Bra/T induction was enhanced, and EMT initiated as early as 24 hrs post WNT activation.

Conversely, expression of constitutively active Rho kinase or myosin light chain kinase, to

increase contractility, efficiently blocked any induction of mesoderm, including EMT.

Mechanistically, reduced contractility promoted active β-catenin levels in the cytoplasm and

nucleus. We propose that the early contraction of human pluripotent stem cell colonies in

response to WNT activation initially delays mesoderm induction by restraining accumulation of

active β-catenin, suggesting a role for contractility as a temporal gate.

Introduction

Mechanical forces play major roles at many stages of animal development: Starting as early as embryo compaction at the 4-cell stage[1], forces are crucial through gastrulation and later during tissue morphogenesis. Internal forces are largely generated by acto-myosin contractility, and in epithelia are transmitted across cell boundaries by intercellular junctions, enabling collective organization. Tension is generated by these junctions and by adhesion to the extracellular matrix (ECM). Early embryogenesis is marked by waves of cell fate specification to create the three major lineages, followed by further differentiation, and these events are likely coordinated with changes in physical forces created by the cells, to which they then respond. Mechanistically, however, the connections between cell fate and mechanical force remain unclear.

Human embryonic stem cell (hESC) and induced pluripotent stem cell (hiPSC) colonies can be induced to select the mesoderm lineage and eventually to differentiate into functional cardiomyocytes. Pluripotent stem cells are strongly epithelial and lineage specification relies on an epithelial-to-mesenchymal transition (EMT) that is reminiscent of some aspects of gastrulation. When treated with the morphogen BMP4, hESCs can self-organize into 2D "gastruloids" [2], and if plated on compliant hydrogel surfaces they will form "gastrulation-like" nodes that express Brachyury, a mesoderm marker. Micropatterning and traction stress mapping demonstrated that regions of high tension promote Brachyury expression, through release of β-catenin from the cell junctions and consequent activation of WNT signaling, which is known to be required for mesoderm induction [3].

Mesoderm can also be generated from human pluripotent stem cells by direct WNT pathway activation using the small molecule GSK3β inhibitor Chiron 99021 (hereafter CHIR), which protects β-catenin from degradation. Whether mechanical forces play any role in this pathway to mesoderm induction is unknown. We observed that hiPSCs induced by CHIR to differentiate into mesoderm show colony contraction, increased phospho-MLC (myosin light chain) and increased optical retardance, which is a non-invasive measure of actomyosin contractility. From this correlation between contractility and mesoderm induction, plus prior work on BMP4-driven mesoderm in hESCs, we expected that inhibiting actomyosin activity would block induction. Surprisingly, however, we obtained the opposite result: suppressing MLC phosphorylation using small molecule inhibitors or genetic tools substantially promoted mesoderm induction, while

further increasing actomyosin contractility was sufficient to block differentiation and EMT

completely. Mechanistically, we found that promoting cell relaxation led to an increase in active

β-catenin in the cytoplasm and nucleus.

Results

<u>WNT-mediated mesoderm specification increases actomyosin contractility</u>

To explore how cells sense, integrate and respond to physical cues, we first investigated

whether differentiating cells experience differential forces along their conversion from hiPSC to

mesoderm. hiPSCs were treated with CHIR to stabilize β-catenin and thereby activate the WNT

pathway, triggering pluripotency exit and commitment to primitive streak and mesoderm

lineages. We noticed that hiPSC colonies showed rapid retraction of their edgea following CHIR

addition **(Supp. Fig 1A)**, consistent with observations from other groups [4]. Having previously

reported an essential and permissive role for apoptosis during mesoderm specification [5], we

tested whether colony retraction was related to the wave of cell death that occurs following

CHIR addition. However, co-treatment with a pan-Caspase inhibitor (Q-VD-OPH) did not affect

retraction of colony edges **(Supp. Fig 1A)**, ruling out an effect of cell density and cell death.

Commitment to the mesoderm lineage also resulted in thicker epithelial layers,

containing taller cells **(Supp. Fig 1B-D)**, suggesting profound morphological changes that are

highly dependent on the cellular contractile state. Consistent with the role of nuclei as

mechanosensor [6,7], differentiating cells exhibit significantly smaller nuclear area after 48h of

CHIR treatment **(Supp. Fig 1E-F)**. To address the variation of cell density, we correlated each of

our nuclear area measurements with the actual cell number **(Supp. Fig 1G)**. As expected,

nuclear area negatively correlates with cell density before differentiation. However, we did not

find any correlation following CHIR treatment, together suggesting that the smaller nuclear size

at 48h of differentiation is unlikely to be related to cell density. While these observations were

indirect, they point towards a differential contractile status of early in mesoderm commitment,

switching from low contractile undifferentiated cells to high contractility. To directly test for

higher contractility, differentiating cells were stained for phospho-T18/S19 Myosin Light Chain 2

(ppMLC), a key regulator of contractility. Immunofluorescences reveal a gradual increase of

ppMLC2 intensity over time, with the formation of supracellular fibers extending across several

cells **(Supp. Fig 1H-I)**. Finally, we took advantage of the birefringence property of actin fibers to

measure light retardance using quantitative polarization microscopy. Birefringence of a material

changes proportionally to the applied strain, providing an orthogonal way to probe for

contractile status during mesoderm commitment. Confirming our previous data, retardance

values increased during hiPSC-to-mesoderm conversion **(Supp. Fig 1J)**. Specificity of the

retardance value as a readout for contractility was tested by incubating undifferentiated hiPSCs

with a ROCK inhibitor (Y-27632) to suppress MLC phosphorylation. Together, these data suggest

that hiPSC differentiating along the cardiac mesoderm trajectory experience increasing

contractile forces mediated by MLC2 activity.

Inhibition of actomyosin contractility promotes hiPSc and hESC conversion to the mesoderm

lineage

We hypothesized that increased contractility might be required for pluripotent cell to

acquire their mesoderm identity. To test this hypothesis, hiPSCs were differentiated in the

presence of ROCK inhibitor, Y-27632. We first confirmed that MLC2 phosphorylation was

reduced by ROCK inhibition **(Supp. Fig 2A-B)**. Contrary to expectations, blocking contractility

strongly promoted Brachyury expression, a mesoderm marker (encoded by the *TBXT* gene, also

known as *T* gene) **(Fig 1A-B)**. Expression of other key mesoderm (*MESP1* and *TBX6*) and EMT

(*SNAI1* and *SNAI2*) genes was also upregulated following Y-27632 treatment **(Fig 1C-F)**. To rule

out an effect for ROCK inhibition on global transcription, we examined expression levels of

genes unrelated to mesodermal fate and did not detect any response to decreased contractility,

suggesting a specific role of actomyosin contractility in cell fate specification **(Supp. Fig 2C)**.

Normal hiPSCs synchronously undergo EMT at 49-52hrs post-CHIR addition. Brachyury

expression is initially restricted to the colony edges at 24hrs, before broader expression across

the colony at later timepoints **(Fig. 1G-H)**. However, Y-27632 treatment resulted in markedly

accelerated EMT, as shown by loss of the epithelial marker ZO-1 from tight junctions **(Fig. 1G, I)**

and increased expression of the mesenchymal marker Slug **(Supp. Fig 2D)**, plus more

widespread Brachyury expression **(Fig. 1G-H)**. To validate these data, we used the more specific

ROCK inhibitor, H-1152, which led to similar decreases in ppMLC2 **(Supp. Fig 2E-F)** and

recapitulated the early EMT and higher induction of mesoderm markers **(Supp. Fig 2G-M)**.

Importantly, these effects of ROCK inhibition were conserved in hESCs expressing a knock-in T-

mNeonGreen fusion [3] **(Fig. 1J-K)** and in wild-type H9 hESCs treated with CHIR. Importantly,

there was no effect of reduced contractility on exit from the pluripotency state **(Fig 1L and**

Supp. Fig 2N, P).

Because apoptosis-mediated ATP release is crucial to enable mesoderm fate

specification [5], we probed for PARP cleavage as an apoptosis marker but did not detect any

difference following addition of Y-27632 or H-1152 **(Supp. Fig 2L and 2N-O)**. MLC2

phosphorylation was previously shown to be spatially regulated by ROCK and Myosin Light

Chain Kinase (MLCK) in polarized and non polarized cells [8,9]. Therefore, we tested the

involvement of MLCK by treating cells with ML-7, a potent MLCK inhibitor [10]. ML-7 led to

efficient reduction in ppMLC2 **(Fig. 1N-O)** and strongly enhanced Brachyury expression at 48h

(Fig. 1P-Q), without affecting cell death. Together these data unexpectedly shows that

pharmacological blocking of MLC2-mediated force generation enhances mesoderm

commitment quantitatively and accelerates the EMT required for conversion to the cardiac

mesoderm.

<u>Genetic suppression of actomyosin contractility promotes stem cell conversion to the</u>

<u>mesoderm lineage</u>

To rule out off target effects of the small molecule inhibitors, we designed a genetic

approach to suppress contractility in human cells. Dephosphorylation of ppMLC is mediated by

Myosin Light Chain Phosphatase, an heterotrimer composed of a catalytic subunit PP1cβ, a

Myosin phosphatase-targeting regulatory subunit (MYPT1) and a protein of unknown function

M20 [11,12] **(Fig 2A-B)**. Based on previous work describing an optogenetic construct to control

actomyosin contractility [13], we created a truncated MYPT1, fused with a nuclear export

sequence and a mNG reporter (**Fig 2B**), that is expected to act as a constitutively active mutant

and promote MLC2 dephosphorylation. To validate the activity of this new tool, we cloned this

construct (referred to as MYPT1^{CA}-NES-mNG) under a doxycycline-inducible promoter and

established a stable hiPSC line. Global reduction of ppMLC2 was observed by immunoblot and

immunofluorescence after doxycycline addition (**Fig 2C-E**). Strikingly, activating MYPT1^{CA}

recapitulated the effects of pharmaceutical inhibition of contractility: differentiating MYPT1^{CA}

cells displayed enhanced Brachyury expression (**Fig 2F-J**) and earlier induction of the EMT genes

Snail and Slug (**Fig 2F**) following Doxycycline induction, compared to controls. Apoptosis

induction was not affected in this system (**Fig 2H**). We conclude that reducing MLC2-driven

contractility using both pharmaceutic and genetic strategies, accelerates mesoderm identity.

<u>Contractility is a potent inhibitor of mesoderm identity</u>

Next, we sought to test if constitutively increased contractility impacts hiPSC-to-mesoderm

conversion. We stably expressed constitutively active ROCK2 or MLCK (ROCK2^{CA} and MLCK^{CA})

under a doxycycline inducible promoter with ubiquitous expression of a mVenus reporter. Green

transduced cells were mixed with unlabeled WT hiPSCs, pre-treated or not with doxycycline to

turn on ROCK2^{CA} or MLCK^{CA} before starting the differentiation protocol for 72h (**Fig 3A**). We first

confirmed that mVenus-positive cells treated with doxycycline showed signs of higher

contractility. In the co-culture set up with ROCK2^{CA}/WT or MLCK^{CA}/WT cells treated with

Doxycycline, mVenus-positive clusters showed strong cortical F-actin and higher ppMLC2

staining organized as fibers compared to WT mVenus-negative cells which show more diffused

staining (**Supp. Fig 3A, C**). These data support the idea that mVenus-positive cells experience

higher contractility, allowing us to test whether cell fate was affected. ROCK2^{CA}/WT co-cultures

were treated with water (+ Veh) or Doxycycline (+Dox) and stained for mesoderm (Bra/T) (**Fig.**

3B), primitive streak (EOMES) (**Fig 3D**) and EMT markers (Slug and ZO-1) (**Fig 3F**). mVenus-

positive and negative area were drawn based on the mVenus coverage across each cluster

(**Supp. Fig 3B-D**) and percentages of cells expressing each marker were reported. Following

doxycycline induction, mVenus-positive clusters showed a dramatic decrease in mesoderm,

primitive streak and mesenchymal markers compared to mVenus-negative cells or Vehicle-

treated cells (**Fig. 3C-E-G**). Because we previously showed that MLCK inhibition affects

mesoderm commitment the same way as ROCK inhibition (**Fig. 1P-Q**), we sought to address the

effect of MLCK-driven contractility on cell identity. Importantly, MLCK^{CA}-positive clusters failed

to commit to the primitive streak and mesoderm lineage and to trigger the EMT program

required for mesoderm identity (**Supp. Fig 3E-G**). These data support the hypothesis the

surprising conclusion that constant and pre-mature activation of ROCK/MLCK-mediated

contractility prevents mesoderm conversion and EMT and is sufficient to counteract the effect

of CHIR.

<u>Reduced cell contractility promotes nuclear accumulation of stable β-Catenin</u>

Mechanistically, we first hypothesized that Hippo signaling, one of the main mechanosensitive

pathways in epithelial tissues, drives mesoderm identity enhancement upon cell relaxation. YAP,

a downstream effector of the Hippo pathway, has been previously associated with cell fate

patterning in 2D gastruloids [14] Activation of the Hippo pathway leads to phosphorylation of YAP

at Ser127, restricting its localization to the cytosol and promoting degradation. In contrast, non-

phosphorylated YAP translocates to the nucleus and forms a complex with co-transcriptional

activator TEAD1-4, together regulating gene expression. Surprisingly, however, phosphorylated

YAP is not affected during differentiation or by cell relaxation (**Supp. Fig 4 A-B**). We next directly

probed for *CTGF* and *CYR61* expression, two well-described YAP target genes. Gene expression

strongly decreased during differentiation but was not affected by cellular contractile status

(**Supp. Fig 4C**). Together, these data rule out a significant role for the Hippo pathway in the

acceleration of mesodermal differentiation following cell relaxation.

Next, we investigated the positive feedback loop reported during BMP4-driven

mesoderm commitment, which relies on mechanical tension-dependent secretion of canonical

WNT ligands [3]. We treated our MYPT1^{CA}-NES-mNG cells with doxycycline to trigger cell

relaxation and probed for canonical and non-canonical WNT ligands. As a positive control,

expression of the mesoderm marker *TBX6* was increased in low contractile cells compared to

Vehicle-treated control cells. Interestingly, canonical WNT ligand expression (*WNT3A* and

WNT8A) was strongly upregulated by suppression of contractility, while non-canonical *WNT4*

expression was unaffected by CHIR treatment nor by the reduction of contractility (**Supp. Fig

4E**). To test whether increased WNT3A/8A expression was responsible for increased mesoderm

commitment in low-contractility cells, we blocked WNT ligand processing and secretion using

IWP-2 (Porcupine inhibitor). Addition of IWP-2 did not rescue mesoderm gene expression in Y-

27632-treated cells, suggesting that this pathway is unlikely to drive mesoderm specification

enhancement (**Supp. Fig 4F**).

We then turned our attention to β-catenin, which exists in epithelial cells both as a

cytosolic molecule and bound to E-Cadherin at adherens junctions and plays key

mechanosensitive functions during development and tissue homeostasis [3,15]. Whether each

pool shares common regulatory mechanisms is still debated [16]. In the absence of WNT signaling,

cytosolic β-catenin is associated with a destruction complex, where it is phosphorylated at

Ser45 by CK1, which primes the protein for subsequent phosphorylation by GSK3β, driving its

ubiquitination and degradation. WNT activation disrupts the destruction complex, permitting

the accumulation of stable, non-phosphorylated β-catenin. Therefore, active non-phospho β-

catenin (hereafter active β-catenin) signaling can be assessed from the level of non-

phosphorylated Ser45. We evaluated active β-catenin shuttling to the nucleus by isolating

cytosolic and nuclear fractions from MYPT1^{CA} cells (**Fig 4A**). Purity of each fraction was assessed

from enrichment of a nuclear and cytosolic marker, Lamin A/C and GAPDH respectively (**Supp.**

Fig 4D). Strikingly, active β-catenin was strongly upregulated in both the cytosolic and the

nuclear fractions in low contractile differentiating cells (24h + Dox) (**Fig. 4B, C**). Immunostaining

confirmed the enrichment of active β-catenin in the nucleus 24h post differentiation (**Fig. 4D-F**).

We validated this observation by imaging a mEGFP-β-Catenin knock-in line treated with CHIR in

normal or low contractility (Y-27632) states. First, we noticed cytosolic accumulation of mEGFP-

β-catenin within 5h of CHIR treatment, regardless of the presence of Y-27632. However,

consistent with the immunostaining data, mEGFP-β-catenin later became enriched in the nuclei

of cells under low contractile stress (**Supp. Fig 4G-H**). Collectively, these findings suggest that a

cell relaxation-dependent nuclear accumulation of β-catenin potentiates the WNT signaling-

dependent mesoderm program. However, this model can only hold true if the concentration of

CHIR used in our system (7.5 µM) does not saturate the WNT signaling response. To test this

premise, we doubled the concentration of CHIR and probed for expression of WNT target genes

as a read-out for WNT signaling (**Fig. 4G**). *LEF1* and *AXIN2* expression responded in a dose-

dependent manner to CHIR. Interestingly, however, at any given CHIR concentration, inhibition

of cell contractility by addition of Y-27632 further enhanced WNT-target gene responses.

Together, these results demonstrate that increased active β-Catenin translocation to the nucleus

following cell relaxation amplifies WNT response and enhances mesoderm commitment, in a

synergistic way with CHIR.

Discussion

Synchronization of developmental processes is key to development. Germ cell lineage

commitment relies on tight spatio-temporal control of genetic programs, which is concomitant

to profound morphological changes during gastrulation. Here, we aimed to investigate how

differentiating cells sense and integrate physical cues. Our data reveal an unexpected and

paradoxical role for acto-myosin contractility as a potent inhibitor of the mesoderm lineage. We

report that low contractility promotes nuclear localization of β-catenin, which correlates with

higher mesoderm marker expression. β-catenin exists as two pools, one bound to the adherens

junctions through E-Cadherin and one cystosolic pool, part of the WNT pathway. The regulation

and exchange of β-catenin molecules between these 2 pools remains poorly understood. E-

Cadherin binding to β-catenin prevents its nuclear translocation and co-activator transcriptional

activity [17] SRC-mediated phosphorylation of β-catenin at residue Y654 reduces by 6-fold the

binding interaction [18] so one possibility might be that SRC activity is dependent on contractility

in pluripotent stem cells. Alternatively, Gayrard et al [19] reported a mechanism by which release

of cell-cell tension is required for β-catenin release through SRC-dependent phosphorylation of

FAK. Therefore, release of tension leads to β-catenin accumulation in the cystosolic

compartment and its translocation into the nucleus as observed by cell fractionation and

confocal imaging (Fig. 4B, D and Supp. Fig 4G).

In several in vivo and in vitro models, including ours, initiation of mesoderm commitment

correlates with tissue deformation and cell contraction [3,20–23]. In *Drosophilia,* contraction of the

posterior end of the embryo is mediated by supercellular active myosin II cables reminiscent

with our observation in 2D [24]. Higher contractility also leads to deformation of nuclear

morphology, as reported by others [21,25]. Therefore, how can we reconciliate this response with

the fact that reducing contractility promotes mesoderm identity? We propose a model in which

increasing contractility following the first 24h of differentiation dampens CHIR response to

temporally synchronize mesoderm identity. Indeed, in the developing embryo, specification of

lineages required deep morphological changes and cell compaction occur, mostly mediated by

actomyosin contractility [26–28]. It is widely accepted that high density cells, similar to ingressing

cells during gastrulation, have low contractility compared to cells at low density, both in vitro

and in vivo [29]. Our work suggest that this mechanical gate prevents mesoderm identity until

mechanical tension is released. Inducing low contractility prior to CHIR treatment, disrupts this

mechanical gate, resulting to accelerated mesoderm identity and EMT.

Our data demonstrate that contractile forces are strongly antagonist for mesoderm induction

They confirm and expand on a previous study using Y-27632 to enhance mesendodermal

differentiation [30], while ruling out potential off target effect of the drug. Our study also suggests

that cell fate can be modulated by actomyosin contractility. In *Xenopus,* decreased ROCK-

dependent actomyosin contractility promotes ectoderm-to-mesoderm transition, with

ectodermal cells adopting mesoderm-like properties [31], together emphasizing the plasticity of

cell fate during development. While we have not tested for alternative cell fate following CHIR

addition, our differentiation protocol was adapted from Lian et al [32,33] and was not reported to

lead to ectodermal identity. Additionally, we note that differentiating cells lose their

pluripotency marks, regardless of their contractile status, suggesting that the rate of

pluripotency exit is not affected (Supp. Fig 2N, P).

We further report that low contractile cells show premature EMT and induction of mesoderm

specification, and link it to higher nuclear β-catenin signaling. These findings balance previous

reports suggesting that mechanical tension promotes Brachyury expression in Bilateria [34] and in

gastrulation-like models, through mechanosensitive regulation of β-catenin at high cell adhesion

tension [3]. Muncie et al proposed a dual function for mechanical tension: First, BMP4-driven

Brachyury expression relies on mechanical stretch of cell junction that triggers a β-catenin-

mechanosensitive response by exposing the conserved Y654 residue. Phosphorylation by SRC-

family kinase destabilizes junctional β-catenin and initiates its nuclear translocation. A positive

feedback loop reinforces mesoderm commitment through increased canonical WNT ligand

secretion. These discrepancies may be explained by differences in the differentiation protocol

(BMP4 versus CHIR) and possibly by the substrate/micropattern used [35], providing differential

biochemical and mechanical inputs. Indeed, our protocol directly activates WNT pathway by

inhibition of the destruction complex, which is in contrast with BMP4 activation. Moreover,

while we noticed increased canonical WNT ligand expression following cell relaxation, we ruled

out a feed-forward loop by blocking WNT ligand secretion (Supp. Fig 4D-E).

Alternative mechanisms might also complement this pathway. For instance, a previous study

shows that hiPSC epithelial cohesiveness is a crucial barrier for BMP4-signaling activation, by

blocking BMP4 diffusion to the basolaterally-localized BMP receptors and restricting Brachyury

expression to the colony edge [36]. However, in our system, the use of the small molecule GSK3β

inhibitor CHIR as a differentiating cue rules out a similar initial mechanism. While WNT target

genes have now been extremely well characterized in multiple systems, it remains unclear how,

β-catenin affects temporally its effectors. The dynamics nature of β-catenin, and the plasticity of

WNT activation in undifferentiated cells [37], may explain the different outcomes resulting from

specific differentiation protocols.

Recent reports highlight the extreme sensitivity of hiPSC to cell density. For example, the key

mechanical transducer ETV4 was linked to cell crowding and fate determination, with an

ON/OFF response dependent on cell density [38]. Similarly, radial patterning of 2D gastruloid was

strongly affected by a 1.5 fold change in cell density, while WNT dynamics was affected by only a

1.15 fold increase of cell density [39]. In our system, we showed that while nuclear area tends to

correlate with cell density in basal condition, this correlation was lost during differentiation.

Additionally, cell density is highly linked with Hippo pathway activity and we showed that Hippo

pathway is not differentially regulated during mesoderm induction, confirming previous data

from our lab at a later timepoint [5]. Unlike other studies on mammary cell lines [40] we did not

notice any effect of ROCK inhibition on YAP phosphorylation state. One explanation might be

that PSCs are strongly epithelial and, as previously reported, are insensitive to Hippo activation-

dependent cell-cell interaction[41].

Limitations of the study

In this study, we aimed to understand if and how mechanical tension is important for mesoderm

lineage specification. To this end, we used a simple 2D system, allowing us to control most

parameters while only perturbing actomyosin contractility. However, this 2D system is limited by

its physiological implication. Translation of these data into 2.5D and 3D models of gastrulation

(2D and 3D gastruloid respectively), and alternatively into early mouse development will allow

the generalization these findings. Moreover, while we consistently observed higher β-catenin in

the nucleus upon cell relaxation, further work will be required to address whether it reflects an

increase of import or a decrease of export from the nucleus.

Finally, it remains to address whether increased nuclear β-catenin results to an increase binding

to its target gene promoters, such as Brachyury promoter, which contains two TCF/LEF-1/β-

catenin binding sites within the first 500pb [42,43]. Alternatively, further work on chromatin

rearrangement will be required to better understand whether enhancer/repressor get spatially

modulated by mechanical tension.

Acknowledgements

We thank Cynthia Reinhart-King for giving us access to the QPOL facility. We thank the Weaver

lab (UCSF) and the Gardel lab (University of Chicago) for sharing the T-mNG knock-in hESC line.

We also thank all members of the Macara lab, especially Vaishna Vamadevan and Christian de

Caestecker for valuable suggestions and advice. This work was funded by an American Heart

Association postdoctoral fellowship (Award ID 23POST1018347 to L.F.), a Vanderbilt

Mechanobiology fellowship (to L.F.) and a grant from the National Institutes of Health

(GM070902 to I.G.M.).

Author contributions

L.F. conceived the study, performed and quantified all experiments except the QPOL, and wrote

the manuscript. W.W. imaged and quantified the QPOL data. I.G.M. supervised the project, and

edited the manuscript

Declaration of interest

The authors declare no competing interests.

Material & Methods

Research compliance and ethical regulation

All experiments using human ESCs were performed using the WA09 (H9) cell line under the

supervision of the Vanderbilt Institutional Human Pluripotent Cell Research Oversight

Committee (protocol IRB no. 160146). No cells were sourced directly from human embryos.

Reagents

Common laboratory reagents used in this study are listed in Supplementary Table 1

Cell culture and maintenance

Human GM25256 iPSCs were obtained from the Coriell Institute. The mEGFP–CTNNB1 knock-in

GM25256 iPSC cell line was obtained from the Allen Cell Collection, Coriell Institute (cell ID

AICS-0058-067). The human ESC line H9 (WA09) was obtained from the WiCell Research

Institute. The T-mNG knock-in hESC H9 was made by Valerie Weaver lab (UCSF) and kindly

provided by Margaret Gardel (University of Chicago). The iPSC and ESC H9 lines were cultured

on Matrigel-coated six-well plates (Matrigel diluted at 42 µg ml−1 in DMEM/F12 medium) and

cultured in mTeSR1 medium. The medium was changed daily until the cells reached 70%

confluency. Cells were passaged using Gentle cell dissociation reagent for 4 min, resuspended in

mTseR1 medium as small clusters and replated at 1:7.

HEK293T cells were obtained from the ATCC. Cells were grown in 10% FBS DMEM medium and

passaged at 1:10 every 2-3 days.

All cell lines used in this study were maintained at 37 °C under 5% CO_2.

<u>Cell freezing and thawing</u>

Human iPSCs and ESCs were harvested from culture dishes using Gentle cell dissociation

reagent and centrifuged at 120g for 3 min. The pellets were resuspended in mTeSR1 medium

supplemented with 10% DMSO and aliquoted into cryovials. The cells were first transferred to

−80 °C for 24 h before long-term storage in liquid nitrogen. iPSCs and ESCs were slowly thawed

using mTeSR1, centrifuged and resuspended in mTeSR1 supplemented with 10 µM Y-27632 for

24 h.

<u>Mesoderm differentiation protocol</u>

Differentiation was initiated when cells reach 70-80% confluency. Cells were treated with 7.5

µM CHIR99021 (hereafter CHIR) diluted in RPMI media supplemented with 1X B27 minus

insulin and 100U/mL Penicillin and 100 µg/mL streptomycin (1X) for up to 72 hours.

For pharmaceutical inhibition of contractility, cells were pre-treated overnight with respective

drug or vehicle (water) diluted in mTeSR1. Differentiation was initiated by switching to CHIR

media as described above, supplemented with respective drug or vehicle (water). Similarly, cells

were pre-induced overnight with Doxycycline or vehicle (water) diluted in mTeSR1 before

initiating the differentiation protocol in the presence of Doxycycline or water. For concentration

of each drug, please refer to Supplementary Table 1.

<u>Lentiviral preparation</u>

HEK193T were grown at 40% confluency in 10 cm plate and transfected using Calcium-

Phosphate. Briefly, 20 µg of lentiviral plasmid, 15 µg of pSPAX2 (Addgene #8454) and 6 µg

pMD2G (Addgene #12260) were mixed with 450 μL sterile water and 50 μL of 2.5M CaCl$_2$

solution, previously filter sterilized. DNA mixture was added dropwise into 500 μL of 2X HeBS

(50 mM HEPES, 10 mM KCl, 12 mM dextrose, 280 mM NaCl and 1.5 mM Na2PO4, pH 7.04) with

constant vortexing. Solution was added dropwise on HEK 293T cells for 6h. Medium was

replaced with 8mL of fresh 10% FBS DMEM medium for 48h. Supernatant was filtered through a

0.22 μm membrane and concentrated using a 100kDa cut-off Amicon centrigal unit. Aliquots

were frozen at -80 °C.

SDS-PAGE and Western blotting

Cells were washed with ice cold 1X PBS and lysed with 70 μL RIPA buffer (150 mM NaCl, 10 mM

Tris–HCl pH 7.5, 1 mM EDTA, 1% Triton X-100, 0.1% SDS) supplemented with and 1X protease

and phosphatase inhibitors. Mechanical disruption was performed by scrapping the cells off the

plate, transferring the lysate into a 1.5mL Eppendorff tube and vortexing for 5 seconds followed

by 5 min incubation on ice. Soluble proteins were collected by centrifugation at 16000 g (13200

rpm) for 10min at 4°C. Protein concentration was measured using Precision Red following the

manufacturer's instructions. Lysates in sample buffer were boiled for 5min.

For ppMLC blotting, cells were washed with 1X PBS and directly lysed using 250 μL of boiling 1X

Sample buffer. Lysates were sonicated 3 times for 10 seconds with 60 seconds incubation on ice

between each blast.

30 μg of proteins were resolved on 1.5mm thick bis-tris acrylamide gels and transferred onto

0.2 μm nitrocellulose membrane using BioRad wet transfer system (50 V for 2hours).

Membranes were blocked for > 30min at room temperature (RT) with 5% BSA in 1X TBS-T

(10 mM Tris pH 8.0, 150 mM NaCl and 0.5% Tween 20) and incubated overnight at 4°C and

gentle rocking with primary antibodies diluted in blocking buffer (Supplementary Table 2).

Membranes were washed 3 times with 1X TBS-T and incubated for 1h at RT with AlexaFluor-

conjugated secondary antibodies (Supplementary Table 2). After 3 additional washes in 1X TBS-

T, membranes were scanned using Li-Cor Odyssey CLx system, analyzed and processed using

Image StudioLite v. 5.2.

Immunofluorescence

Cells were cultured on Matrigel-coated Ø 12mm coverglass, fixed with 4% paraformaldehyde for

10min, permeabilized for 5min at RT (20mM Glycine, 0.05% Triton X-100 in 1X PBS), and

incubated with blocking buffer for > 30min (5% BSA in 1X PBS). Coverslips were transferred into

a dark humidified chamber and incubated with primary and AlexaFluor-conjugated secondary

antibodies (Supplementary Table 2) diluted in blocking buffer for 2 hours. Coverslips were

washed in 1X PBS and mounted on glass slide using Fluoromount-G mounting solution.

For ppMLC staining, samples were incubated overnight with primary antibody.

<u>Image acquisition, analysis and processing</u>

Micrographs were obtained using an inverted Nikon A1-R confocal microscope equipped with a

x20 objective (Numerical aperture NA 0.75), a x40 oil objective (NA 1.20) and a x100 oil

objective (NA 1.40). Z-stacks covering the entire cell height were acquired. Maximum Intensity

Projections were obtained and images were analyzed and processed using Fiji (version

2.1.0/1.54f).

<u>MYPT1^{CA}-NES-mNeonGreen cloning strategy</u>

gBlock corresponding to truncated human MYPT1 (aa 1-300) was built in fusion with a PKI super

NES and mNeonGreen sequence. gBlock was amplified using Fusion polymerase and the

following primers: Fwd 5'-CTGCTGACCGGTACCATGGCGGACGCGAAGC-3' and Rev 5'-

CATCATACGCGTCTACGATCCGCCACCGC-3'. PCR product and recipient backbone (pInducer10b-

HA-KRas G12V Addgene #164928) were digested with AgeI-HF and MluI-HF for 1h at 37°C, gel

purified and ligated overnight at 16°C using T4 ligase. Ligation reactions were transformed into

chemically competent E.Coli Stbl3 strain by heatshock (45 sec at 42°C) and selected on

Ampicillin plate. Clones were isolated and set up for MiniPrep (QIAprep Spin MiniPrep kit). The

presence of the insert was confirmed by restriction digest check. Positive clones were sent for

full plasmid sequencing provided by Genewiz Plasmid-EZ service before MidiPrep using Takara

Bio NucleoBond XtraMidi kit.

<u>Flow cytometry analysis</u>

T-mNeonGreen knock-in hESC were transduced with a lentivector expressing Histone 2B-

mScarlet (pWPI-H2B-mScarlet). Cells were pre-treated overnight with Vehicle (water) or 10 µM

Y-27632. Differentiation was initiated for 24 hours as described above and cells were harvested

as single cells. Briefly, cells were washed and gentle cell dissociation buffer was added for 8 min

at 37°C. Cell suspension was centrifuged and resuspended in 500 μL PBS. Cells were passed

through a Ø 40 μm strainer directly into 500 μL of 8% PFA (final concentration 4%) and fixed for

10 min on a wheel before sending them for cytometry using a 5-laser Fortessa analyzer (70 μm

nozzle). Singlets were selected based of forward and side scatter. mScarlet (PE-Texas-RedA) and

mNeonGreen (GFP-A) population were selected by drawing a gate on double negative WT hESC

shifted by one order of magnitude. Percentage of mNeonGreen/mScarlet double positive was

reported.

<u>RNA isolation and RT-qPCR</u>

Cells were washed and RNAs were isolated using RNeasy mini kit following manufacturer's

instructions. 1 μg of RNA was reverse transcribed to cDNA using the SuperScript III first-strand

synthesis kit. cDNA was diluted 1:5 in water and mixed with Maxima SYBR Green/1mM of each

primer (Supplementary Table 3). Quantitative PCR was performed on a BioRad CFX96

thermocycler. C_t values from technical triplicates were averaged and normalized to GAPDH using

the $\Delta\Delta C_t$ formula.

<u>Cell Fractionation</u>

Following appropriate treatment, cells were washed with ice cold 1X PBS and collected as single

cell suspension using gentle cell dissociation as previously described. Cell suspension was

washed with cold 1X PBS and resuspended in 500 μL of cold PBS. 100 μL of the cell suspension

was set aside and mixed with 30 μL of 4X sample buffer. Sample was sonicated for 15 sec at 20%

power 3 times, boiled 5min and stored at -20 °C (Whole Cell Lysate fraction or WCL).

Cell fractionation was performed using Cell signaling kit (#9038) and their protocol. Briefly, the

remaining 400 μL of cell suspension was centrifuged at 500 x g at 4°C for 5min. Supernatant was

removed and pellet was resuspended in 250 μL of CIB buffer, vortexed for 5 sec and incubated

on ice for 5 min. Samples were centrifuged at 500 x g at 4°C for 5min and supernatant was

stored in pre-chilled tube (= Cytoplasmic fraction). Pellet was resuspended in 250 μL of MIB

buffer, vortexed for 15 sec and incubated on ice for 5 min. Samples were centrifuged at 8000 x g

at 4°C for 5min and supernatant was discarded (= Membrane and organelle fraction). Pellet was

resupended in 125 µL of CyNIB buffer, sonicated for 5 sec at 20% power 3 times (cytoskeletal

and nuclear fraction). Each fraction was mixed with the appropriate volume of 4X sample buffer

to obtain a final concentration of 1X. Samples were boiled for 5min and centrifuged before

loading 15 µL.

<u>Reproducibility and statistics</u>

All datasets were analyzed using GraphPad Prism10 (version 10.0.3). Individual measurements

(n) were averaged for each biological repeat (N). Datasets were tested for normality (Shapiro-

Wilk test) before applying the appropriate statistical test on the biological repeat (N), except if

mentioned differently in the figure legend. For dataset displayed as superplots, individual

measurements and biological repeats are represented by small and large dots respectively. Each

dataset for a biological repeat is color coded. Error bar represent S.D. Significance levels are

given as follows and exact P values are indicated in each figure: NS, not significant ($P \geq 0.05$);

$*P < 0.05$; $**P < 0.01$; $***P < 0.001$; and $****P < 0.0001$.

All experiments, including representative MIPs and western blots, were performed at least three

times independently as biological repeats, unless stated otherwise in the legends. No statistical

method was used to pre-determine the sample size, the experiments were not randomized and

the investigators were not blinded to allocation during experiments and outcome assessment.

Figure legends

<u>Figure 1: Pharmaceutical inhibition of actomyosin contractility promotes hiPSc and hESC</u>

<u>conversion to the mesoderm lineage</u>

A, B – Immunoblot of Brachyury during hiPSC-to-mesoderm commitment (0-48h) in the

presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Molecular weights (M.W.) are

displayed on the right side (A). Brachyury expression was quantified by densitometry and

normalized to α-Tubulin as loading control across N=3 independent biological repeats (B). Mean

and S.D. are displayed. Two-tailed unpaired t test was performed.

C-F – Relative expression of mesoderm markers (*MESP1, TBX6*) (C, D) and EMT markers (*SNAI1,*

SNAI2) (E, F) 48h post CHIR treatment in the presence (+Y-27) or absence (+Veh) of ROCK

inhibitor. N=3 independent biological repeats except for *MESP1* N=4 independent biological

repeats. Mean and S.D. are displayed. Two-tailed unpaired t test was performed.

G-I – Representative MaxIP immunofluorescences during hiPSC-to-mesoderm commitment (0-

48h) in the presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Cells were stained with a

nuclear marker (DNA), EMT marker (ZO-1 – Inverted LUT) and Mesoderm marker (T/Bra–

Inverted LUT). Scale bar = 50 µm (G). Quantification of Brachyury expression (H) and ZO-1

intactness (I) is represented as violin plots with individual measurements (small dots) averaged

for each biological repeat (large dots). Median (Plain red line) and quartiles (Dotted black lines).

n=10-15 technical repeats across N=3 independent biological repeats. One-way ANOVA with

Tukey's multiple comparisons post-test was performed on biological repeats.

J, K – Analysis of Brachyury expression by flow cytometry following 24h CHIR treatment in the

presence (+Y-27632) or absence (+Veh) of ROCK inhibitor using Bra-mNeonGreen knock-in H9

hESC line (J). Quantification of the percentage of cells mNeonGreen-positive from N=4

independent biological repeats. Mean and S.D. are displayed. Two-tailed unpaired t test was

performed (K).

L, M – Immunoblot of Brachyury (mesoderm) and Nanog (Pluripotency marker) using hESC H9 in

the presence (+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Molecular weights (M.W.) are

displayed on the right side (L). Brachyury expression was quantified by densitometry and

normalized to α-Tubulin as loading control across N=3 independent biological repeats (M).

Mean and S.D. are displayed. Two-tailed unpaired t test was performed.

N, O – Representative MaxIP immunofluorescence of hiPSCs treated with a MLCK inhibitor (ML-

7) or Vehicle (DMSO) and stained for nuclei (DNA) and phospho T18/S19 Myosin Light Chain 2

(ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views of the yellow dotted ROI are

shown for ppMLC2. Scale bar = 20 µm (N). Percentage of cellular area positive for ppMLC2

following Vehicle or ML-7 treatment is reported as violin pots. Median (Plain red line) and

quartiles (Dotted black lines). n=15 technical repeats across N=3 independent biological repeats.

Two-tailed unpaired t test was performed on the biological repeats.

P, Q - Immunoblot of Brachyury (mesoderm) and PARP (Cell death marker) using hiPSCs in the

presence (+ ML-7) or absence (+ Veh) of MLCK inhibitor at basal state (0h) and during CHIR

treatment (24h, 48h). Molecular weights (M.W.) are displayed on the right side (P). Brachyury

expression was quantified by densitometry and normalized to α-Tubulin as loading control

across N=3 independent biological repeats. Mean and S.D. are displayed. Mann-Whitney test

was performed for the 24h timepoints and two-tailed unpaired t test was performed for the 48h

timepoints (Q).

<u>Figure 2: Genetic inhibition of actomyosin contractility promotes hiPSc conversion to the</u>

<u>mesoderm lineage</u>

A – Summary of the main regulators of MLC phosphorylation.

B – Schematic representation of full length MYPT1 (1030 aa), interacting with PP1c phosphatase

and protein of unknown function M20. Truncation of MYPT1 Cterm Δ301-1030 leads to a

constitutive recruitment and activation of PP1C. Truncated MYPT1 was fused with NES-

mNeonGreen and is referred to as MYPT1^{CA}-NES-mNG.

C – Immunoblot of phospho T18/S19 MLC2 (ppMLC2) and total MLC2 in WT hiPSCs treated with

Doxycycline or hiPSC expressing MYPT1^{CA}-NES-mNG in the presence or absence of Doxycycline.

α-Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side

D, E – Representative MaxIP immunofluorescence of hiPSC expressing MYPT1^{CA}-NES-mNG and

treated or not with Doxycycline. Cells were fixed and stained for phospho T18/S19 MLC2

(ppML2 - Magenta), DNA (blue) and tight junction marker ZO-1 (Orange). Scale bar = 50 μm (D).

Fraction of the cellular area positive for ppMLC2 following Doxycycline induction of MYPT1^{CA}-

NES-mNG hiPSC and reported as violin plots. Median (Plain red line) and quartiles (Dotted black

lines). n=32 technical repeats across N=3 independent biological repeats. Two-tailed unpaired t

test was performed on the biological repeats (E).

F, G – Representative immunoblot for Brachyury (mesoderm marker), Snail and Slug (EMT

markers) using WT and MYPT1^{CA}-NES-mNG hiPSCs, following CHIR treatment (0h-48h) in the

presence or absence of Doxycycline. Molecular weights (M.W.) are displayed on the right side

(F). Brachyury expression was quantified by densitometry and normalized to α-Tubulin as

loading control across N=3-4 independent biological repeats. Mean and S.D. are displayed. A

one-way ANOVA with Tukey's multiple comparisons post-test was performed (G).

H – Immunoblot of PARP (cell death marker) and Brachyury (Mesoderm marker) during

mesoderm commitment (0h-48h CHIR) using MYPT1^{CA}-NES-mNG hiPSCs induced with

Doxycycline. Molecular weights (M.W.) are displayed on the right side.

I, J – Representative MaxIP immunofluorescences of WT and MYPT1^{CA}-NES-mNG hiPSCs treated

with CHIR for 24h +/- Doxycycline. Cells were stained for nuclei (DNA), EMT marker (ZO-1 –

Inverted LUT) and Mesoderm marker (T/Bra – Inverted LUT). Scale bar = 100 µm (I).

Quantification of Brachyury expression is presented as violin plots. Median (Plain red line) and

quartiles (Dotted black lines). n=15-16 technical repeats across N=3 independent biological

repeats. One-way ANOVA with Šidák's multiple comparisons post-test was performed on

biological repeats (J).

Figure 3: Increased actomyosin contractility is sufficient to block mesoderm commitment and

EMT

A – Experimental design for co-culture experiment. hiPSC were transduced with lentivector

expressing ROCK2^{CA} or MLCK^{CA}. Transduced cells are mVenus positive and are mixed with WT

mVenus-negative hiPSCs. ROCK2^{CA}/MLCK^{CA} expression is induced by addition of Doxycycline 24h

prior to initiating mesoderm differentiation. Doxycycline induction was maintained during the 3

days of differentiation using CHIR before fixing and staining.

B-H – Representative MaxIP immunofluorescence for Brachyury (B), EOMES (D), Slug (F) and ZO-

1 (H) in Vehicle (H2O) and Doxycycline-induced ROCK2^{CA} co-culture. mVenus-positive cell

clusters are highlighted by an orange dotted line. Individual channels are inverted LUT. Scale bar

= 50 µm. For (H), magnified mVenus-positive and mVenus-negative areas are shown as insets.

Scale bar = 20 µm. Quantification of Brachyury (C), EOMES (E) and Slug-positive cells (G) is

reported as violin plot, comparing mVenus-positive vs mVenus-negative cluster in the presence

or absence of Doxycycline. For Brachyury, n = 7 (+ Veh) and n=15 (+ Dox) across N=3

independent biological repeats. For EOMES, n=9 across N=3 independent biological repeats. For

Slug, n=9 (+ Veh) and n=10 (+ Dox) across N=3 independent biological repeats. One-way ANOVA

with Tukey's multiple comparisons post-test was performed (C, E, G).

Figure 4: Reduced cell contractility promotes nuclear accumulation of active β-Catenin

A – Experiment design for treatments and cell fractionation.

B-C – Representative immunoblot as explained in (A). MYPT1^{CA}-NES-mNG hiPSCs were induced

with CHIR for 24h in the presence or absence of Doxycycline. Whole cell lysate (WCL) was

collected and loaded as input while cytosolic and nuclear fractions were extracted. Fraction

purity was probed using cytosolic (GAPDH) and nuclear (Lamin A/C) markers. Active β-Catenin

(non-phospho S45) enrichment was probed for each fraction and α-Tubulin was used as loading

control. Molecular weights (M.W.) are displayed on the right side (B). Quantification of active β-

Catenin enrichment in the cytosolic (left) and nuclear (right) fractions. N=3 independent

biological repeats. Mean and S.D. are displayed. One-way ANOVA with Šidák's multiple

comparisons post-test was performed (C).

D-F – Representative confocal slice and MaxIP for active non-phospho S45 β-Catenin (Inverted

LUT) and mNeonGreen respectively. MYPT1^{CA}-NES-mNG hiPSCs were fixed at 0h and 24h post

CHIR treatment complemented or not with Doxycycline. Scale bar = 50 μm. Insets represent a

magnified view of the active non-phospho S45 β-Catenin with an example of the nuclei binary

mask overlaid (pink). Scale bar = 10 μm (D). Schematic representation of the quantification

pipeline. (E). Quantification of active non-phospho S45 β-Catenin mean intensity as depicted in

(E). n = 10-12 technical repeats across N=3 independent biological repeats. Mean and S.D. are

displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed (F).

G – Relative expression of WNT target genes (*LEF1* and *AXIN1*) with increasing concentration of

CHIR in the presence (Magenta) or absence (Cyan) of ROCK inhibitor Y-27632. N=7 independent

biological repeats. Mean and S.E.M. are displayed. One-way ANOVA with Šidák's multiple

comparisons post-test was performed.

<u>Supplementary Figure 1: hiPSC differentiation along the cardiac mesoderm increases</u>

<u>actomyosin-driven contractility (Related to Figure 1)</u>

A – Bright field imaging of WT hiPSCs treated with CHIR in the presence or absence of a pan-

Caspase inhibitor Q-VD-OPH. Retraction area is highlighted in yellow. Time is presented as

hh:mm. Scale bar = 20 μm.

B, C – Confocal XZ orthogonal views (B) and Maximum Intensity Projections (MaxIP) (C) of

hiPSCs at basal state (0h CHIR) or post-CHIR treatment (24h and 48h CHIR), stained for DNA

(Cyan) and F-Actin (Grey). Scale bar = 10 μm.

D – Quantification of the monolayer height at 0h, 24h and 48h post CHIR treatment from (B).

Violin plots represent individual measurements (small dots) averaged for each biological repeat

(large dots). Median (Plain red line) and quartiles (Dotted black lines). n=32 technical repeats

across N=7 independent biological repeats. A Kruskal-Wallis test with Dunn's multiple

comparison post-test was performed on biological repeats.

E, F – Representative MaxIP immunofluorescence of WT hiPSC during CHIR treatment. Fixed

cells were stained for DNA marker (Hoechst). Scale bar = 20 µm (E). Quantification of average

nuclear size from n=15 fields of view across N=3 independent biological repeats. Median (Plain

red line) and quartiles (Dotted black lines) are displayed. Two-tailed unpaired t test was

performed (F).

G – Correlation of cell area over cell density for each field of view during CHIR treatment.

Biological repeats are color-coded. Pearson correlation coefficient is displayed.

H – Representative MaxIP immunofluorescence of hiPSCs at basal state (0h) or post-CHIR

treatment (24 and 48h) stained for phospho T18/S19 MLC2 (ppMLC2). Pixels are color-coded by

intensity using the fire LUT.

I – Percentage of total imaged area that is ppMLC2-positive at 0h, 24h, 48h post CHIR is

reported as violin plots. Median (Plain red line) and quartiles (Dotted black lines). n=15

technical repeats across N=3 independent biological repeats. A Kruskal-Wallis test with Dunn's

multiple comparison post-test was performed on the technical repeats.

J – Retardance measurements obtained from hiPSCs at basal state (0h), post-CHIR treatment

(24h, 48h, 72h) or treated with Y-27632 at 0h (Y-27) using Quantitative Polarization Microscopy

imaging. Median (Plain red line) and quartiles (Dotted black lines). n=240 technical repeats for

0h, 24h, 48h, 72h and n=120 technical repeats for Y-27 across N=3 independent biological

repeats. A one-way ANOVA with Šidák's multiple comparisons post-test was performed on the

biological repeats.

<u>Supplementary Figure 2: Pharmaceutical inhibition of actomyosin contractility promotes hiPSc</u>

<u>and hESC conversion to the mesoderm lineage (Related to Figure 2)</u>

A, B - Representative MaxIP immunofluorescences of hiPSCs treated with a ROCK inhibitor (Y-

27632) or Vehicle (H_2O) and stained for nuclei marker (DNA) and phospho T18/S19 Myosin Light

Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 µm. Magnified views of the yellow dotted ROI

are shown for ppMLC2. Scale bar = 20 µm (A). Fraction of the cellular area positive for ppMLC2

following Y-27632 treatment and reported as violin pots. Median (Plain red line) and quartiles

(Dotted black lines). n=30 technical repeats across N=3 independent biological repeats. Two-

tailed unpaired t test was performed on the biological repeats (B).

C – Relative expression of unrelated genes 48h post CHIR treatment in the presence (+Y-27) or

absence (+Veh) of ROCK inhibitor. N=4 independent biological repeats. Mean and S.D. are

displayed. Two-tailed unpaired t test was performed.

D - Representative MaxIP immunofluorescences of hiPSC treated 48h with CHIR in the presence

(+ Y-27632) or absence (+ Veh) of ROCK inhibitor. Cells were stained for nuclei marker (DNA),

EMT markers (ZO-1 and Slug – Inverted LUT). Scale bar = 50 μm.

E, F - Representative MaxIP immunofluorescences of hiPSCs treated with a ROCK inhibitor (H-

1152) or Vehicle (H_2O) and stained for nuclei marker (DNA) and phospho T18/S19 Myosin Light

Chain 2 (ppMLC2 – Inverted LUT). Scale bar = 50 μm. Magnified views of the yellow dotted ROI

are shown for ppMLC2. Scale bar = 20 μm (E). Fraction of the cellular area positive for ppMLC2

following H-1152 treatment and reported as violin pots. Median (Plain red line) and quartiles

(Dotted black lines). n=15 technical repeats across N=3 independent biological repeats. Two-

tailed unpaired t test was performed on the biological repeats (F).

G-K - Relative expression of mesoderm markers (*TBXT*, *MESP1, TBX6*) (G-I) and EMT markers

(*SNAI1, SNAI2*) (J, K) 48h post CHIR treatment in the presence (+H-1152) or absence (+Veh) of

ROCK inhibitor. N=3 independent biological repeats. Mean and S.D. are displayed. Two-tailed

unpaired t test was performed.

L – Immunoblot probing for mesoderm marker expression (T/Bra) and cell death marker (PARP

cleavage) during hiPSC-to-mesoderm commitment (0h-48h CHIR) in the presence of ROCK

inhibitor (+H-1152). Addition of the pan-caspases inhibitor Q-VD-OPH (+ Q-VD) to the

differentiation medium totally blocks mesoderm commitment and was used as negative control.

α-Tubulin was used as loading control. Molecular weights (M.W.) are displayed on the right side.

M – Representative MaxIP immunofluorescences for mesoderm marker Brachyury (T/Bra –

Inverted LUT) and nuclei. Cells were fixed 24h post-CHIR treatment in the presence or absence

of ROCK inhibitor H-1152.

N-P – Representative immunoblot of PARP (cell death), T/Bra (mesoderm), Oct4 and Nanog

(Pluripotency markers) using hESC H9 in the presence (+ Y-27632) or absence (+ Veh) of ROCK

inhibitor. Molecular weights (M.W.) are displayed on the right side (N). PARP cleavage (O) and

Nanog expression (P) was quantified by densitometry and normalized to α-Tubulin as loading

control across N=4 (PARP) and N=3 (Nanog) independent biological repeats. Mean and S.D. are

displayed. Two-tailed unpaired t test was performed.

<u>Supplementary Figure 3: Increased actomyosin contractility is sufficient to block mesoderm</u>

<u>commitment and EMT (Related to Figure 3)</u>

A – Representative MaxIP immunofluorescence of Doxycycline-induced ROCK2^{CA} hiPSCs

(mVenus-positive) mixed with WT hiPSCs (mVenus-negative). Cells were stained for DNA

(Nuclei), F-actin (top – Inverted LUT) and phospho T18/S19 Myosin Light Chain 2 (Bottom -

ppMLC2 – Inverted LUT). Scale bar = 50 μm. Magnified views comparing mVenus-positive

(Green dotted ROI) and mVenus-negative (Black dotted ROI) areas are shown for F-Actin and

ppMLC2. Scale bar = 10 μm.

B – Percentage of the ROI area positive for mVenus based on cluster considered as mVenus-

Negative and Positive using the ROCK2^{CA} line. On average, mVenus negative clusters have <15%

of their area positive for mVenus. Mean and S.E.M are displayed. n=7 (no Dox) and n=15 fields

of view (+Dox) across N=3 independent biological experiments.

C – Representative MaxIP immunofluorescence of Doxycycline-induced MLCK^{CA} hiPSCs (mVenus-

positive) mixed with WT hiPSCs (mVenus-negative). Cells were stained for DNA (Nuclei), F-actin

(top – Inverted LUT) and phospho T18/S19 Myosin Light Chain 2 (Bottom - ppMLC2 – Inverted

LUT). Scale bar = 50 μm. Magnified views comparing mVenus-positive (Green dotted ROI) and

mVenus-negative (Black dotted ROI) areas are shown for F-Actin and ppMLC2. Scale bar = 10

μm.

D – Percentage of the ROI area positive for mVenus based on cluster considered as mVenus-

Negative and Positive using the MLCK^{CA} line. On average, mVenus negative clusters have <15%

of their area positive for mVenus. Mean and S.E.M are displayed. n=7 (no Dox) and n=18 fields

of view (+Dox) across N=3 independent biological experiments.

E, F - Representative MaxIP immunofluorescence for Brachyury in Vehicle (+Veh) and

Doxycycline-induced MLCK^{CA} co-culture. mVenus-positive cell clusters are highlighted by an

orange dotted line. Individual channels are inverted LUT. Scale bar = 50 μm (E). Percentage of

Brachyury-positive nuclei is reported as violin plot, comparing mVenus-positive vs mVenus-

negative cluster in the presence or absence of Doxycycline. Median (Plain red line) and quartiles

(Dotted black lines). n=7 (+Veh) and n=18 (+ Dox) fields of view (small dots) across N=3

independent biological repeats (large dots). One-way ANOVA with Tukey's multiple comparisons

post-test was performed (F).

G - Representative MaxIP immunofluorescence for EOMES, Slug and ZO-1 in Doxycycline-

induced MLCK^{CA} co-culture. mVenus-positive cell clusters are highlighted by an orange dotted

line. Individual channels are inverted LUT. Scale bar = 50 μm.

<u>Supplementary Figure 4: Reduced cell contractility promotes nuclear accumulation of active β-</u>

<u>Catenin (Related to Fig. 4)</u>

A, B – Immunoblot for phospho S127 YAP (pYAP)and total YAP following differentiation (0-48h

CHIR) in the presence or absence of ROCK inhibitor Y-27632. Brachyury expression was also

probed as an internal positive control and α-Tubulin was used as loading control. Molecular

weights (M.W.) are displayed on the right side (A). Ratio of phospho/total YAP was quantified

across N=4 independent biological repeats. Mean and S.D. are displayed. One-way ANOVA with

Šidák's multiple comparisons post-test was performed.

C – Relative expression of YAP-target genes *CTGF* and *CYR61* during differentiation in the

presence or absence of ROCK inhibitor Y-27632. N=4 independent biological repeats. Mean and

S.D. are displayed. One-way ANOVA with Šidák's multiple comparisons post-test was performed.

D - Relative expression of canonical WNT (*WNT3A, WNT8A*), non-canonical WNT (*WNT4*) and

mesoderm marker (*TBX6*) in MYPT1^{CA}-NES-mNG cells treated with CHIR (0-48h) in the presence

of Vehicle or Doxycycline. N=3 independent biological repeats. Mean and S.D. are displayed.

One-way ANOVA with Šidák's multiple comparisons post-test was performed.

E – Relative expression of *TBX6* and *MESP1* (mesoderm markers) following CHIR treatment in

the presence of Y-27632 complemented or not with 7.5 μM Porcupine inhibitor (IWP-2 or DMSO

respectively). N=3 independent biological repeats. Mean and S.D. are displayed. One-way

ANOVA with Tukey's multiple comparisons post-test was performed.

F – Compartment purity following fractionation was assessed by quantifying Lamin A/C and

GAPDH enrichment in both, cytosolic and nuclear fractions across N=4 independent biological

repeats.

G, H – Still pictures from confocal timelapse imaging of mEGFP-β-Catenin knock-in hiPSCs

treated with CHIR +/- Y-26732. Time is shown in hours. Scale bar = 10 μm (G). mEGFP-β-Catenin

intensity is reported as a scan line plot across a cell at 0h and 20h (H).

References

1.   Firmin, J. *et al.* Mechanics of human embryo compaction. *Nature* 629, 646–651
(2024).
2.   Deglincerti, A. *et al.* Self-organization of human embryonic stem cells on
micropatterns. *Nat Protoc* 11, 2223–2232 (2016).
3.   Muncie, J. M. *et al.* Mechanical Tension Promotes Formation of Gastrulation-like
Nodes and Patterns Mesoderm Specification in Human Embryonic Stem Cells. *Dev*
Cell 55, 679-694.e11 (2020).
4.   Hookway, C. *et al.* A human induced pluripotent stem (hiPS) cell model for the
holistic study of epithelial to mesenchymal transitions (EMTs). *BioRxiv* (2024).
5.   Fort, L., Gama, V. & Macara, I. G. Stem cell conversion to the cardiac lineage requires
nucleotide signalling from apoptosing cells. *Nat Cell Biol* 24, 434–447 (2022).
6.   Lomakin, A. J. *et al.* The nucleus acts as a ruler tailoring cell responses to spatial
constraints. *Science (1979)* 370, (2020).
7.   de Leeuw, N. F., Budhathoki, R., Russell, L. J., Loerke, D. & Blankenship, J. T. Nuclei as
mechanical bumpers during epithelial remodeling. *Journal of Cell Biology* 223,
(2024).
8.   Totsukawa, G. *et al.* Distinct roles of ROCK (Rho-kinase) and MLCK in spatial
regulation of MLC phosphorylation for assembly of stress fibers and focal adhesions
in 3T3 fibroblasts. *J Cell Biol* 150, 797–806 (2000).
9.   Shi, Y., Ohler, L. S., de Jesus, B. N. & Dickinson, D. J. Opposing actomyosin pools
generate cortical flows that establish epithelial polarity. *Biorxiv* (2024).
10. Makishima, M. *et al.* Induction of differentiation of human leukemia cells by
inhibitors of myosin light chain kinase. *FEBS Lett* 287, 175–177 (1991).
11. Matsumura, F., Yamakita, Y. & Yamashiro, S. Myosin Phosphatase-targeting Subunit 1
Controls Chromatid Segregation. *Journal of Biological Chemistry* 286, 10825–10833
(2011).
12. Kiss, A., Erdődi, F. & Lontay, B. Myosin phosphatase: Unexpected functions of a long-
known enzyme. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1866,
2–15 (2019).
13. Yamamoto, K. *et al.* Optogenetic relaxation of actomyosin contractility uncovers
mechanistic roles of cortical tension during cytokinesis. *Nat Commun* 12, 7145
(2021).
14. Stronati, E. *et al.* YAP1 regulates the self-organized fate patterning of hESC-derived
gastruloids. *Stem Cell Reports* 17, 211–220 (2022).
15. Röper, J.-C. *et al.* The major β-catenin/E-cadherin junctional binding site is a primary
molecular mechano-transductor of differentiation in vivo. *Elife* 7, (2018).
16. van der Wal, T. & van Amerongen, R. Walking the tight wire between cell adhesion
and WNT signalling: a balancing act for β-catenin. *Open Biol* 10, (2020).
17. Orsulic, S., Huber, O., Aberle, H., Arnold, S. & Kemler, R. E-cadherin binding prevents
β-catenin nuclear localization and β-catenin/LEF-1-mediated transactivation. *J Cell*
Sci 112, 1237–1245 (1999).
18. Roura, S., Miravet, S., Piedra, J., de Herreros, A. G. & Duñach, M. Regulation of E-
cadherin/Catenin Association by Tyrosine Phosphorylation. *Journal of Biological*
Chemistry 274, 36734–36740 (1999).

19. Gayrard, C., Bernaudin, C., Déjardin, T., Seiler, C. & Borghi, N. Src- and confinement-
dependent FAK activation causes E-cadherin relaxation and β-catenin activity.
Journal of Cell Biology 217, 1063–1077 (2018).

20. Leptin, M. & Grunewald, B. Cell shape changes during gastrulation in *Drosophila*.
Development 110, 73–84 (1990).

21. Skory, R. M. *et al.* The nuclear lamina couples mechanical forces to cell fate in the
preimplantation embryo via actin organization. *Nat Commun* 14, 3101 (2023).

22. Caldarelli, P. *et al.* Self-organized tissue mechanics underlie embryonic regulation.
Nature 633, 887–894 (2024).

23. Rozbicki, E. *et al.* Myosin-II-mediated cell shape changes and cell intercalation
contribute to primitive streak formation. *Nat Cell Biol* 17, 397–408 (2015).

24. Rozbicki, E. *et al.* Myosin-II-mediated cell shape changes and cell intercalation
contribute to primitive streak formation. *Nat Cell Biol* 17, 397–408 (2015).

25. Dahl, K. N., Kahn, S. M., Wilson, K. L. & Discher, D. E. The nuclear envelope lamina
network has elasticity and a compressibility limit suggestive of a molecular shock
absorber. *J Cell Sci* 117, 4779–4786 (2004).

26. Maître, J.-L. *et al.* Adhesion Functions in Cell Sorting by Mechanically Coupling the
Cortices of Adhering Cells. *Science (1979)* 338, 253–256 (2012).

27. Krieg, M. *et al.* Tensile forces govern germ-layer organization in zebrafish. *Nat Cell
Biol* 10, 429–436 (2008).

28. Turlier, H. & Maître, J.-L. Mechanics of tissue compaction. *Semin Cell Dev Biol* 47–48,
110–117 (2015).

29. Loza, A. J. *et al.* Cell density and actomyosin contractility control the organization of
migrating collectives within an epithelium. *Mol Biol Cell* 27, 3459–3470 (2016).

30. Maldonado, M., Luu, R. J., Ramos, M. E. P. & Nam, J. ROCK inhibitor primes human
induced pluripotent stem cells to selectively differentiate towards mesendodermal
lineage via epithelial-mesenchymal transition-like modulation. *Stem Cell Res* 17,
222–227 (2016).

31. Kashkooli, L., Rozema, D., Espejo-Ramirez, L., Lasko, P. & Fagotto, F. Ectoderm to
mesoderm transition by down-regulation of actomyosin contractility. *PLoS Biol* 19,
e3001060 (2021).

32. Lian, X. *et al.* Robust cardiomyocyte differentiation from human pluripotent stem
cells via temporal modulation of canonical Wnt signaling. *Proceedings of the
National Academy of Sciences* 109, (2012).

33. Lian, X. *et al.* Directed cardiomyocyte differentiation from human pluripotent stem
cells by modulating Wnt/β-catenin signaling under fully defined conditions. *Nat
Protoc* 8, 162–175 (2013).

34. Brunet, T. *et al.* Evolutionary conservation of early mesoderm specification by
mechanotransduction in Bilateria. *Nat Commun* 4, 2821 (2013).

35. Llewellyn, J., Charrier, A., Cuciniello, R., Helfer, E. & Dono, R. Substrate stiffness
alters layer architecture and biophysics of human induced pluripotent stem cells to
modulate their differentiation potential. *iScience* 27, 110557 (2024).

36. Vasic, I. *et al.* Loss of TJP1 disrupts gastrulation patterning and increases
differentiation toward the germ cell lineage in human pluripotent stem cells. *Dev Cell*
58, 1477-1488.e5 (2023).

37. Pagella, P. *et al.* The time-resolved genomic impact of Wnt/β-catenin signaling. *Cell
Syst* 14, 563-581.e7 (2023).

38.   Yang, S. *et al*. ETV4 is a mechanical transducer linking cell crowding dynamics to
lineage specification. *Nat Cell Biol* 26, 903–916 (2024).
39.   Rufo, J. *et al*. An explainable map of human gastruloid morphospace reveals
gastrulation failure modes and predicts teratogens. *BioRvix* (2024).
40.   Dupont, S. *et al*. Role of YAP/TAZ in mechanotransduction. *Nature* 474, 179–183
(2011).
41.   Pagliari, S. *et al*. YAP–TEAD1 control of cytoskeleton dynamics and intracellular
tension guides human pluripotent stem cell mesoderm specification. *Cell Death*
Differ 28, 1193–1207 (2021).
42.   Yamaguchi, T. P., Takada, S., Yoshikawa, Y., Wu, N. & McMahon, A. P. T (Brachyury) is
a direct target of Wnt3a during paraxial mesoderm specification. *Genes Dev* 13,
3185–3190 (1999).
43.   Arnold, S. J. *et al*. Brachyury is a target gene of the Wnt/β-catenin signaling pathway.
Mech Dev 91, 249–258 (2000).
870

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

Supplementary Figure 1 (Related to Figure 2)

Supplementary Figure 3 (Related to Figure 3)

Supplementary Figure 4 (Related to Figure 4)